Ketogenesis is dispensable for the metabolic adaptations to caloric restriction

Chung-Yang Yeh[1,2], Lexie Borgelt[1,2], Brynn J. Vogt[1,2], Alyssa A. Clark[1,2], Ted T. Wong[1,2], Isaac Grunow[1,2],

Michelle M. Sonsalla[1,2,3], Reji Babygirija[1,2,4], Yang Liu[1,2,6], Michaela E. Trautman[1,2], Mariah F. Calubag[1,2,4], Bailey

A. Knopf[1,2,4], Fan Xiao[1,2,5], and Dudley W. Lamming[1,2,3,4,5,6,7,8*]

[1]Department of Medicine, University of Wisconsin-Madison, Madison, WI 53705, USA
[2]William S. Middleton Memorial Veterans Hospital, Madison, WI 53705, USA
[3]Comparative Biomedical Sciences Graduate Training Program, University of Wisconsin-Madison, Madison, WI 53706, USA
[4]Graduate Program in Cellular and Molecular Biology, University of Wisconsin-Madison, Madison, WI 53706, USA
[5]Nutrition and Metabolism Graduate Program, University of Wisconsin-Madison, Madison, WI 53706, USA
[6]Endocrinology and Reproductive Physiology Graduate Program, University of Wisconsin-Madison, Madison, WI 53706, USA
[7]University of Wisconsin Comprehensive Diabetes Center, University of Wisconsin-Madison, Madison, WI, 53705, USA
[8]University of Wisconsin Carbone Comprehensive Cancer Center, University of Wisconsin-Madison, Madison, WI 53705, USA

[*]**Correspondence**

Dudley W. Lamming, PhD
Associate Professor of Medicine
University of Wisconsin-Madison
1685 Highland Ave
MFCB Rm 4147
Madison, WI 53705, USA
dlamming@medicine.wisc.edu
Tel: 608-262-7341

Abstract

Caloric restriction (CR) robustly extends the health and lifespan of diverse species. When fed once daily, CR-treated mice rapidly consume their food and endure a prolonged fast between meals. As fasting is associated with a rise in circulating ketones, we decided to investigate the role of ketogenesis in CR using mice with whole-body ablation of *Hmgcs2*, the rate-limiting enzyme producing the main ketone body β-hydroxybutyrate (βHB). Here, we report that *Hmgcs2* is largely dispensable for many metabolic benefits of CR, including CR-driven changes in adiposity, glycemic control, liver autophagy, and energy balance. Although we observed sex-specific effects of *Hmgcs2* on insulin sensitivity, fuel selection, and adipocyte gene expression, the overall physiological response to CR remains robust in mice lacking *Hmgcs2*. To gain insight into why deletion of *Hmgcs2* does not disrupt CR, we measured fasting βHB levels as mice began a CR diet. Surprisingly, as CR-fed mice adapt to CR, they no longer engage high levels of ketogenesis during the daily fast. Our work suggests that the benefits of long-term CR in mice are not mediated by ketogenesis.

Keywords: caloric restriction, dietary restriction, metabolism, ketogenesis, ketones, BHB, metabolic health.

Introduction

Caloric restriction (CR) extends healthspan and longevity in a variety of species ranging from yeast, worms, and flies to mice and non-human primates, with benefits for healthy aging seen as well in short-term human studies [1]. CR-fed mice are typically fed one meal per day, containing 20-40% fewer calories than animals in a matched *ad libitum* (AL) control group with 24-hour access to food. This style of CR leads to the mice binge eating their daily food allotment in 2-3 hours, self-imposing a prolonged fast between meals [2-4]. This daily fasting period contributes to the benefits of CR in both wild-type mice [5] and Alzheimer's disease mice [6].

Fasting elicits a plethora of strong physiological adaptations [7]. One of the key metabolic responses, as blood glucose levels are depleted during fasting, is to initiate ketogenesis and produce ketone bodies from fatty acids [8]. Of the three ketone bodies – acetoacetate, acetone, and 3-β-hydroxybutyrate (βHB) – βHB is the most abundant and well-studied. βHB acts as both an energy currency and as a signaling molecule, for instance, through direct post-translational modifications and the modulation of histone deacetylases [9]. Fasting, ketogenic diets, and dietary supplementation with exogenous ketones significantly elevate circulating βHB levels and can elicit benefits in Alzheimer's disease animal models [10-12]. A critical unanswered question is whether fasting-induced βHB contributes to the beneficial effects of CR [7].

To determine how ketogenesis contributes to the metabolic response to CR, we generated a mouse model lacking *Hmgcs2*, the rate-limiting enzyme in the formation of ketones from fatty acid-derived acetyl-CoA. We placed both male and female *Hmgcs2$^{-/-}$* (KO) mice on either an *ad libitum* (AL) diet or a 30% reduced CR diet (CR), with CR-fed animals receiving food once per day at the beginning of the light cycle. We extensively characterized the metabolic phenotypes of these animals, finding that KO mice broadly received the same benefits from CR as wild-type (WT) mice, with few genotype-diet interactions. To understand why the loss of *Hmgcs2* led to unexpectedly limited physiological consequences, we measured the βHB level over 24 hrs. Surprisingly, the WT mice treated with prolonged CR do not produce high levels of circulating βHB as would the matching AL mice when fasting. In a separate cohort of mice, we demonstrated that this downregulation in ketogenesis occurs during the first few weeks of CR initiation. Together, our results suggest that ketogenesis is dispensable for the metabolic benefits of CR and that ketones are not elevated during long-term CR due to adaptations to recurrent daily fasting.

Results

Hmgcs2 deletion does not alter the effects of CR on weight and body composition

To investigate the role of ketogenesis in caloric restriction (CR), we generated mice lacking whole body *Hmgcs2* expression in the C57BL/6J genetic background. At 9 weeks of age, male and female Hmgcs2$^{+/+}$ (WT) and Hmgcs2$^{-/-}$ (KO) mice were singly housed and either maintained on an *ad libitum* diet (AL) or "stepped down" to a 30% reduced CR diet, fed once per day within 1 hr from the start of the light cycle (**Fig. 1A**). The CR-fed mice were pair-fed according to the AL group, with meal size adjusted weekly. Food consumption and body weight (BW) were tracked weekly, and metabolic testing began after 8 weeks on diet; following the completion of the experiments we confirmed that HMGCS2 protein was undetectable in KO mice across multiple tissues, including the liver, small intestine (jejunum and ileum), large intestine (colon), and kidney (**Figs. 1B-1E**). Notably, in WT mice of both sexes, CR strongly induced HMGCS2 expression in the kidney but not in other tissues (**Fig. 1E**).

As anticipated, CR suppressed weight gain over a 7-month period in both WT and KO male mice (**Fig. 1F**). Throughout the experiment, the AL group did not exhibit notable changes in food consumption (**Supplemental Fig. 1A**). Knockout of *Hmgcs2* did not affect the ability of CR to suppress the gain of fat (**Supplemental Fig. 1B**) and lean mass (**Supplemental Fig. 1C**), and we observed the corresponding changes in body composition (**Fig. 1G; Supplemental Fig. 1D**). Nearly identical observations were made in the female mice (**Figs. 1I-1J; Supplemental Figs. 1E-1H**). Analysis of final changes in BW, fat, and lean mass revealed only robust diet effects and no significant effect of genotype (**Figs. 1H & 1K**).

Hmgcs2 deletion improves insulin sensitivity in female mice

To determine the effect of *Hmgcs2* deletion on glucose metabolism, we performed glucose, insulin, and pyruvate tolerance tests. As our previous work has shown that time since feeding dictates insulin sensitivity in CR [13], tests were conducted after either 21 hrs or 7 hrs of fasting, which corresponds to the CR feeding time of approximately 6 am, or 7 hrs after CR feeding at approximately 4 pm, respectively.

We observed an overall positive effect of CR on glucose metabolism. CR improved glucose tolerance in both male and female mice regardless of the length of fasting; however, there was no effect of genotype or a genotype x diet interaction in either sex (**Figs. 2A & 2E; Supplemental Figs. 2A & 3A**). In agreement with our

previous observations, CR improves insulin sensitivity after 21 hrs of fasting in both sexes (**Figs. 2B & 2F**), yet the shorter 7 hr fasting period revealed post-prandial insulin resistance in CR-fed animals (**Supplemental Figs. 2B & 3B**). Interestingly, the deletion of *Hmgcs2* had an overall beneficial effect in females regardless of the length of the fast, significantly increasing insulin sensitivity in female AL mice when insulin sensitivity was assessed after 21 hrs of fasting (**Figs. 2F; Supplemental Fig. 3B**).

A pyruvate tolerance test examines the ability of the liver to suppress gluconeogenesis. We found that CR improves pyruvate tolerance in all groups (**Figs. 2C & 2G; Supplemental Figs. 2C & 3C**). In females, although there was no overall effect of genotype, we observed a significant genotype-diet interaction, suggesting that *Hmgcs2* contributes to CR-mediated improvements in pyruvate tolerance (**Fig. 2G**).

In a glucose-stimulated insulin secretion test after 21 hrs fasting, there was a trend in the suppression of insulin secretion in the males by CR (p=0.0643) (**Supplemental Figs. 2D and 3D**). There was no overall effect of diet, genotypes, or their interaction on insulin sensitivity as assessed by HOMA-IR in either sex (**Figs. 2D & 2H**). However, there was a trend (p=0.1470) towards improved insulin sensitivity in KO females (**Fig. 2H**). Overall, while we identified some interesting sex-specific genotype effects and genotype-diet interactions, we do not find the ablation of *Hmgcs2* to significantly impede the main metabolic characteristics of CR.

Hmgcs2 deletion selectively alters energy balance

We utilized indirect calorimetry to determine the effects of *Hmgcs2* knockout on CR metabolism. In the males, CR significantly increased energy expenditure of the KO mice in the light cycle and there is a genotype-diet interaction (**Fig. 3A**). In both males and females, CR significantly reduced energy expenditure in the dark cycle, reflecting a time when the animals would be fasting (**Figs. 3A & 3C**). The measurement of respiratory exchange ratio is an indicator of fuel selection. Due to the distinct feeding pattern of the CR animals, there is a significant, but expected, increase in RER during the light cycle and decrease in RER during the dark cycle of both male and female mice (**Figs. 3B & 3D**). In males, there is a genotype-diet effect in the light cycle and the dark cycle, the latter of which shows significantly increased RER in the KO mice treated with CR during the fasting period (**Fig. 3B**). In females, no significant genotype effect was observed in the binned data, although the post CR-feeding RER increase in the light cycle appears to be dampened by the loss of *Hmgcs2* (**Fig. 3D**).

CR-feeding led to an increase in activity in the light cycle for both males and females; the average activity level is not different in any of the groups in the dark phase (**Supplemental Figs. 4A & 4B**).

CR effects on iWAT adipocyte physiology and liver autophagy do not require *Hmgcs2*

We have previously shown that CR induces a significant reprogramming of adipocyte metabolism [14]. We characterized the inguinal white adipose tissues (iWAT) in WT and KO mice after 7 months of CR feeding. At the time of sacrifice, the total weight of iWAT is less in the CR animals, and CR robustly reduced adipocyte size (**Figs. 4A-4C & 4F-4H**). qPCR analysis of the iWAT of male mice treated with CR found robust upregulation of the thermogenic gene *Ucp1*, and genes that facilitate lipid metabolism, including *Dgat1*, *Elovl3*, *Cidea*, *Fasn*, and *Acc1*. Interestingly, the loss of *Hmgcs2* in males upregulates *Ucp1* even further and blunts the effect of CR on the expression of *Cidea* (**Figs. 4D-4E**). In the iWAT of the females, the same set of genes were likewise significantly upregulated by CR, except for *Ucp1* and *Acc1,* with no effect of *Hmgcs2* in the expression of *Ucp1* or *Cidea* (**Figs. 4I-4J**). In addition, the knockout of *Hmgcs2* suppressed *Lipe* and *Atgl* in females, but not in males (**Supplemental Figs. 5A-5B**).

Autophagy is induced by CR and can also be induced by either a ketogenic diet [15] or βHB signaling [16-18]. In male liver, CR robustly upregulated the total expression of the autophagosomal marker LC3 and the autophagy receptor p62 regardless of genotype (**Fig. 5A**). On the other hand, LC3 B/A ratio, and mTORC1-associated protein phosphorylation on pS757-ULK, pS240/244-S6, and pT37/46-4E-BP1 were unchanged (**Supplemental Fig. 5C**). In females, we did not detect any changes in the expression of autophagy proteins (**Fig. 5B**), but there was a slight decrease in the phosphorylation of pT37/46-4E-BP1 induced by CR (**Supplemental Fig. 5D**). None of the protein targets we investigated have a response to the KO genotype.

Ketogenesis engagement is minimized after prolonged CR

To further understand why the loss of *Hmgcs2* had no overt effects on CR physiology, we sought to determine the ketone levels of these animals throughout the course of a day. We measured βHB over a period of 24 hrs, with each group in their typical experimental condition (**Fig. 6A; Supplemental Figs. 6A-6B**). Although WT mice on the CR diet exhibited slightly higher βHB levels than those on the AL diet, the differences are negligible at most timepoints, with peak βHB concentrations remaining below 1.0 mmol/L. This contradicts the

long-held assumption that CR should significantly increase ketone production [8]. To further investigate this phenomenon, we measured the circulating βHB level of the AL mice while fasting, starting at the beginning of the light cycle (**Fig. 6B; Supplemental Figs. 6C-6D**). Predictably, ketone levels rise in the WT group only, with ketogenesis particularly engaged after the dark cycle begins. After 12-16 hrs of fasting, circulating βHB levels peaked at 2.5+ mmol/L, much higher than that achieved by the mice treated with CR. For comparison, a dotted line is placed at the 1 mmol/L level in both graphs. This rise in βHB level was not observed in the fasting AL KO group, which still exhibited fluctuations in βHB readings, but at a much lower level.

Finally, we looked to further characterize the development of this resistance to ketogenesis by CR-fed mice by measuring fasting ketone levels during the first 8 weeks of CR initiation. In a separate cohort of mice, we measured the fasting ketone levels of all groups once per week at the conclusion of a 21 hr fast. As part of this experiment, AL-fed mice were fasted only once per week, whereas CR-fed mice experience self-administered daily fasting as a result of their feeding behavior (**Fig. 6C**). We found that in both males and females, the circulating βHB level achieved by the WT CR-fed mice begin to drop below that of the WT AL-fed mice after 2 weeks of CR conditioning (**Figs. 6D-6E**). These results are strong evidence that the adaptation to daily fasting in CR-fed mice results in drastically reduced circulating ketones levels in response to fasting.

Caloric restriction enhances ketone metabolism

This significant reduction in circulating ketone levels could result from either a lack of ketone production or an increase in ketone consumption. To determine whether the ability of these animals for ketogenesis and ketolysis is altered, we treated them with either exogenous medium-chained triglycerides (MCT) or βHB, respectively. Consumption of βHB-precursors, such as medium triglyceride (MCT) oil, acutely increases circulating βHB in humans [19,20] and in mice [21]. After oral administration of MCT oil, we monitored the circulating level of βHB for 4 hrs. We found that this treatment reliably engaged ketogenesis, particularly in females, with WT female mice fed the CR diet showing significantly higher levels of circulating βHB than AL-fed WT females (**Figs. 7A-7B**).

In the ketone tolerance test, we injected exogenous sodium βHB and monitored circulating βHB levels for 3 hrs. This approach has been shown to readily raise circulating βHB levels in mice [22-24]. We confirmed a rapid absorption of the exogenous βHB into circulation, where the peak level is observed at 15-30 min after

injection and steadily decreases thereafter. In both males and females, CR feeding clearly enhanced the metabolism of this excess ketone challenge (**Figs. 7C-7D**). In females only, KO of *Hmgcs2* further enhanced ketone catabolism, significantly decreasing the βHB area under the curve in AL-fed mice. Together, these results suggest that CR mice have enhanced capacity for both ketogenesis and ketolysis if the appropriate substrates are available.

To summarize our findings, a visualization of all discrete data is rank-sorted by the difference between the WT and KO mice under the CR feeding condition and provided in heatmap form (**Figs. 7E-7F**). As expected, the top traits involve ketone metabolism, while the arrangement of the remaining data is less predictable.

Discussion

The precise mechanisms by which geroprotective dietary interventions promote healthy aging continue to be elusive. Here, we provide clear evidence that ketogenesis is dispensable for the many metabolic benefits of a CR diet, and that adaptation to daily fasting simply discourages βHB production. Nevertheless, the current literature has conflicting reports on whether the CR diet actually increases circulating βHB. In a fed-once-daily 40% CR diet, fasting blood βHB levels appears to decrease in mice, with a counterbalanced enhancement in liver glycogen [25]. To this aim, the contribution of enhanced glycogen metabolism on physiological health has yet to be elucidated. In aged rats, a fed-once-daily 40% CR diet increases circulating ketone levels [26]. In a mouse model of prostate cancer, a two-days-a-week (5:2) intermittent fasting protocol caused urinary ketone to be initially highly elevated, but returns to normal levels after 8 fasting episodes [27].

Our ability to interpret human data to determine if this response is conserved is limited, as most published studies have either a low number of participants or lack a control group. Several studies show that fasting or CR in overweight or obese subjects do induce ketogenesis. In a patient case report, a 5:2 fasting protocol with 36-hrs fasting windows observed that circulating βHB remained highly elevated even after 8 months of practice [28]. Obese (BMI >30) asthma patients treated with alternate day 20% CR reliably have increased ketone production for at least 8 weeks [29]. However, some very long-term studies of CR suggest that our results also apply to people. These include a study of young overweight women (BMI 24-40) treated with 25% daily energy restriction; while there was an initial increase in serum ketone level at month 1 and 3, this difference was reduced with time and disappears by month 6; meanwhile in the same study, 25% CR in the form of severe restriction limited to 2 day/week appears to maintain elevated ketone levels over 6 months [30]. The phase II CALERIE clinical trial was one of the most comprehensive CR patient studies, and unique in that it focused on non-obese adults. After 24 months, patients practicing ~11.9% CR (prescribed as 25%) exhibited a modest increase in fasting plasma ketones compared to the AL group (p=0.055; [31]), with a very low incidence of urinary ketosis (4.4% in CR vs 1.9% in AL; [32]). On balance, these studies suggest that while CR and fasting initially induces ketogenesis in the overweight and obese, over the long-term CR does not drastically elevate ketone levels in most non-obese individuals.

In addition to its role as an energy currency, βHB is known to act as an effective signaling molecule that elicits multiple benefits [9]. βHB is an inhibitor of histone deacetylases (HDACs) [33] and contributes to a recently

identified post-translational protein modification, lysine β-hydroxybutyrylation [34]. While the concentration of βHB that needs to be reached for its signaling properties has not been established, our results suggest that the βHB levels reached during CR may be a poor surrogate to investigate this, as βHB concentration in a CR-treated mouse is not much higher than that of a non-fasting AL animal. This implies that many CR benefits are achieved either through another mechanism distinct from the actions of βHB, or that the effect of βHB only requires a small local increase that is undetectable in circulation.

Acetoacetate is the canonical precursor to βHB and while the conversion between the two molecules is reversible, deletion of hepatic *Hmgcs2* has been shown to deplete both plasma acetoacetate and βHB [35]. However, despite the deletion of *Hmgcs2* in our study, we observed non-zero amounts of circulating βHB. This has been observed previously by others [35], and is most likely explained by two routes of non-canonical ketogenesis. First, endogenous gut microbiota can produce butyrate, which can directly contribute to plasma βHB levels through a molecular pathway independent of HMGCS2 [36]. The branched chain amino acid leucine can also be catabolized to produce HMG-CoA, the precursor to acetoacetate, without HMGCS2, accounting for roughly 4.4% of the ketone bodies in starved rats [37]. Nevertheless, it is clear from our data that these alternate sources of ketone are dwarfed by canonical *Hmgcs2*-dependent ketogenesis, which is completely abolished by the deletion of *Hmgcs2* and not compensated for by the non-canonical pathways.

The main limitation in this study is that we cannot test the role of HMGCS2 in every CR response, owing to the wide range of known biological effects. While we focused on metabolism, CR can elicit other benefits that require their own investigation, including improved brain function [38], mitochondria health [39], and disease mitigation [1]. Similarly, we only evaluated these effects in one strain of mice, and did not evaluate the role of HMGCS2 in CR-induced lifespan extension. The role of ketones in different genetic background, and its roles on lifespan is an important future study, as there have been mouse strains identified for which CR do not extend or even shorten lifespan [40-42].

It is a long-held assumption that the fasting component of CR induces significant ketogenesis, contributing to the benefits of the diet [8]. Here, we provide proof that this assumption – like many others about the mechanisms by which CR functions [43,44] – may be incorrect. Mice completely lacking the capacity for fasting-induced ketogenesis still receive most of the metabolic benefits of CR, and indeed the long-term adaptation to

CR downregulate ketogenesis. Further work should be done to understand the mechanisms of this physiological "memory" for fasting. Altogether, our result demonstrates that it may be wise for us to consider the effects of dietary interventions not solely as acute responses, but more as part of a biological framework consisting of long-term adaptations.

Materials and Methods

Statistical analysis. Where an analysis for a 2-factorial experimental design is possible, we performed 2-way ANOVA with Sidak's multiple comparison post hoc test. Exact p-value of the post hoc tests are provided in the source data file. All significance test calculations were done with GraphPad Prism ver10.4.0.

Animals. All procedures were performed in accordance to institutional guidelines and were approved by the Institutional Animal Care and Use Committee of William S. Middleton Memorial Veterans Hospital (Madison, WI, USA) and the University of Wisconsin-Madison. *Hmgcs2* KO mice were generated by the Advanced Genome Editing Lab of the University of Wisconsin-Madison Biotechnology Center, which performed pronuclear microinjection of a CRISPR/Cas9 construct that resulted in the insertion of a fragmented gene duplication downstream of exon 2, rendering the loss of functional gene transcript expression. This has been confirmed with long-read (PacBio) genome sequencing. The loss of detectable HMGCS2 protein and the ability to engage canonical ketogenesis has been confirmed as presented.

CR-fed mice are fed at the beginning of the light phase, typically between 6:00 to 7:00 am. All mice were fed the Teklad Global 18% Protein Rodent Diet (Envigo 2018). CR is implemented in a step-down protocol of increasing 10% per week until 30% restriction is reached. Animals are adapted to the feeding protocol for 8 weeks prior to physiological testing. All mice were maintained at a temperature of approximately 22ºC, and health checks were performed daily by the facility staff. Relative humidity of the animal room ranges between 30-70%, with an average of 42%. Mice were housed in a SPF facility in static microisolator cages under 12:12 light cycle conditions with *ad libitum* access to water.

Animals were temporarily housed in the Oxymax/CLAMS-HC metabolic chamber system (Columbus Instruments) during indirect calorimetry experiments. Mice were housed in this system for ~48 continuous hours. The first ~24 hours of data was discarded as acclimation period with a subsequent continuous 24 hours period utilized for data analysis.

Food consumption of the AL group is monitored weekly. All animals in this experiment were singly housed. Body composition was determined using an EchoMRI Body Composition Analyzer.

RT-qPCR. CR mice were euthanized 4-5 hrs after feeding and AL mice were fasted overnight and refed at the same time as CR feeding. Tissues were flash frozen using L_2N. mRNA extraction was performed on iWAT using TRI reagent (AM9738; Invitrogen) and cDNA was generated using Superscript III reverse transcriptase (56575; Invitrogen) according to manufacturer instructions. RT-qPCR was carried out using SYBR green (*Power* SYBR Green PCR Master Mix; 4367659; Applied Biosystems). The following primers were used: *Dgat1* F 5'TGG TGT GTG GTG ATG CTG ATC'3 R 5'GCC AGG CGC TTC TCA A3'; *Elovl3* F 5'ATG CAA CCC TAT GAC TTC GAG 3' R 5'ACG ATG AGC AAC AGA TAG ACG 3'; *Cidea* F 5'GAA TAG CCA GAG TCA CCT TCG 3', R 5'AGC AGA TTC CTT AAC ACG GC3'; *Fasn* F 5'CCCCTCTGTTAATTGGCTCC 3', R 5'TTGTGGAAGTGCAGGTTAGG 3'; *Acc1* F 5'AAG GCT ATG TGA AGG ATG 3', R 5'CTG TCT GAA GAG GTT AGG 3'; *Ucp1* F 5' GCA TTC AGA GGC AAA TCA GC 3', R 5' GCC ACA CCT CCA GTC ATT AAG 3'; *Lipe* F 5' CTG AGA TTG AGG TGC TGT CG 3', R 5' CAA GGG AGG TGA GAT GGT AAC 3'; *Atgl* F 5' ATA TCC CAC TTT AGC TCC AAG G 3', R 5' CAA GTT GTC TGA AAT GCC GC 3'; *β-Act* F 5' GAT GTA TGA AGG CTT TGG TC 3', R 5' TGT GCA CTT TTA TTG GTC TC 3'. qPCR outlier analysis was performed using Prism 10.4.0 built-in function ROUT (Q=1%).

Western Blot. Liver tissues for Western Blotting were flash frozen using L_2N and homogenized in RIPA buffer containing EDTA-free Protease and Phosphotase Inhibitor Mini Tablet (Thermo Scientific, A32961). Quantification of all proteins are normalized to the loading control and expressed as a percentage of the WT-AL group. Total protein quantifications are normalized to HSP90 and phosphoresidues were normalized to the total protein. Antibodies were purchased from Cell Signaling Technology: Hmgcs2 (20940), LC3A/B (12741), HSP90 (4877), ULK1 (8054), pS757-ULK (14202), 4E-BP1 (9644), pT37/46-p4E-BP1 (2855), S6 (2217), pS240/244-S6 (2215), HRP-linked anti-rabbity secondary (7074). The P62 antibody was purchased from Progene Biotechnik (GP62-C). HRP-linked anti-guinea pig secondary antibody was purchased from American Research Products (90001).

In vivo **procedures.** Metabolic tolerance tests were carried out after either 21 hrs or 7 hrs fasting as indicated. The 21 hrs fast tests were carried out at the beginning of the light cycle while the 7 hrs fast tests were carried out ~3 hrs prior to the dark cycle. In the glucose and insulin tolerance test, mice were injected with glucose (1 g/kg; Sigma-Adrich, G7021) or insulin (0.50 and 0.75 U/kg; Novolin) intraperitoneally (I.P.). For glucose-stimulated insulin secretion, 2 g/kg of glucose was used. Blood glucose was monitored via a Bayer Contour glucometer. Lipid tolerance test (LTT) and ketone tolerance test (KTT) were performed on animals after ~4 hrs fast post the CR feeding period. In the KTT, βHB in the form of a sodium salt (CAS-No: 150-83-4; Sigma-Aldrich) is administered at 2 g/kg via I.P. injections. In the lipid tolerance test, MCT oil, containing mostly capric acid and caprylic acid (M4200; CAS 73398-61-5; Spectrum Chemical), was delivered via oral gavage (5 mL/kg). Ketone level was measured using KetoBM handheld ketone meter.

iWAT Histology. After sacrifice, a portion of the iWAT was fixed in formaldehyde. Paraffin sections were stained with H&E and quantified using imageJ for adipocyte size. For each mouse, 2-4 fields of view were scored by three blinded researchers and averaged together for analysis.

DATA AVAILABILITY STATEMENT

The data that support the plots within this article and other findings of this study, including full scans of western blot images, are provided as Source Data files.

DECLARATION OF INTERESTS

DWL has received funding from, and is a scientific advisory board member of, Aeovian Pharmaceuticals, which seeks to develop novel, selective mTOR inhibitors for the treatment of various diseases.

ACKNOWLEDGEMENTS

We would like to thank all members of the Lamming lab for their assistance and input. The Lamming laboratory is supported in part by the NIH/NIA (AG056771, AG062328, AG081482, and AG084156 to D.W.L.), NIH/NIDDK (DK125859 to D.W.L.) and startup funds from the University of Wisconsin-Madison School of Medicine and Public Health and Department of Medicine to D.W.L. C-Y.Y. was supported in part by a NIA F32 postdoctoral fellowship (F32AG077916) and a NIA K99 award (K99AG084921). M.F.C. is supported by F31AG082504. R.B. is supported by F31AG081115. M.E.T. is supported by a F99/K00 award AG083290. The authors used the University of Wisconsin-Madison Genome Editing and Animal Models (GEAM) Core for generating the animals and the UW Carbone Cancer Center Experimental Animal Pathology Laboratory (P30CA014520) for the iWAT staining. The Lamming lab was supported in part by the U.S. Department of Veterans Affairs (IS1-BX005524), and this work was supported using facilities and resources from the William S. Middleton Memorial Veterans Hospital. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. This work does not represent the views of the Department of Veterans Affairs or the United States Government.

References

1. Green, C.L., Lamming, D.W., and Fontana, L. (2022). Molecular mechanisms of dietary restriction promoting health and longevity. Nat Rev Mol Cell Biol 23, 56-73. 10.1038/s41580-021-00411-4.
2. Duregon, E., Pomatto-Watson, L., Bernier, M., Price, N.L., and de Cabo, R. (2021). Intermittent fasting: from calories to time restriction. Geroscience 43, 1083-1092. 10.1007/s11357-021-00335-z.
3. Acosta-Rodríguez, V.A., de Groot, M.H.M., Rijo-Ferreira, F., Green, C.B., and Takahashi, J.S. (2017). Mice under Caloric Restriction Self-Impose a Temporal Restriction of Food Intake as Revealed by an Automated Feeder System. Cell Metab 26, 267-277.e262. 10.1016/j.cmet.2017.06.007.
4. Bruss, M.D., Khambatta, C.F., Ruby, M.A., Aggarwal, I., and Hellerstein, M.K. (2010). Calorie restriction increases fatty acid synthesis and whole body fat oxidation rates. Am J Physiol Endocrinol Metab 298, E108-116. 10.1152/ajpendo.00524.2009.
5. Pak, H.H., Haws, S.A., Green, C.L., Koller, M., Lavarias, M.T., Richardson, N.E., Yang, S.E., Dumas, S.N., Sonsalla, M., Bray, L., et al. (2021). Fasting drives the metabolic, molecular and geroprotective effects of a calorie-restricted diet in mice. Nat Metab 3, 1327-1341. 10.1038/s42255-021-00466-9.
6. Babygirija, R., Han, J.H., Sonsalla, M.M., Matoska, R., Calubag, M.F., Green, C.L., Tobon, A., Yeh, C.Y., Vertein, D., Schlorf, S., et al. (2024). Fasting is required for many of the benefits of calorie restriction in the 3xTg mouse model of Alzheimer's disease. bioRxiv. 10.1101/2024.09.19.613904.
7. Mihaylova, M.M., Chaix, A., Delibegovic, M., Ramsey, J.J., Bass, J., Melkani, G., Singh, R., Chen, Z., Ja, W.W., Shirasu-Hiza, M., et al. (2023). When a calorie is not just a calorie: Diet quality and timing as mediators of metabolism and healthy aging. Cell Metabolism 35, 1114-1131. https://doi.org/10.1016/j.cmet.2023.06.008.
8. de Cabo, R., and Mattson, M.P. (2019). Effects of Intermittent Fasting on Health, Aging, and Disease. N Engl J Med 381, 2541-2551. 10.1056/NEJMra1905136.
9. Newman, J.C., and Verdin, E. (2017). β-Hydroxybutyrate: A Signaling Metabolite. Annu Rev Nutr 37, 51-76. 10.1146/annurev-nutr-071816-064916.
10. Madhavan, S.S., Roa Diaz, S., Peralta, S., Nomura, M., King, C.D., Ceyhan, K.E., Lin, A., Bhaumik, D., Foulger, A.C., Shah, S., et al. (2025). β-hydroxybutyrate is a metabolic regulator of proteostasis in the aged and Alzheimer disease brain. Cell Chem Biol 32, 174-191.e178. 10.1016/j.chembiol.2024.11.001.
11. Ye, Y., Fu, C., Li, Y., Sun, J., Li, X., Chai, S., Li, S., Hou, M., Cai, H., Wang, Z., and Wu, M. (2024). Alternate-day fasting improves cognitive and brain energy deficits by promoting ketone metabolism in the 3xTg mouse model of Alzheimer's disease. Exp Neurol 381, 114920. 10.1016/j.expneurol.2024.114920.
12. Di Lucente, J., Persico, G., Zhou, Z., Jin, L.-W., Ramsey, J.J., Rutkowsky, J.M., Montgomery, C.M., Tomilov, A., Kim, K., Giorgio, M., et al. (2024). Ketogenic diet and BHB rescue the fall of long-term potentiation in an Alzheimer's mouse model and stimulates synaptic plasticity pathway enzymes. Communications Biology 7, 195. 10.1038/s42003-024-05860-z.
13. Pak, H.H., Grossberg, A.N., Sanderfoot, R.R., Babygirija, R., Green, C.L., Koller, M., Dzieciatkowska, M., Paredes, D.A., and Lamming, D.W. (2024). Non-canonical metabolic and molecular effects of calorie restriction are revealed by varying temporal conditions. Cell Reports 43. 10.1016/j.celrep.2024.114663.
14. Calubag, M., Ademi, I., Yeh, C.-Y., Babygirija, R., Bhoopat, A., Kasza, I., Sonsalla, M., Green, C., Pak, H., and Lamming, D. (2022). FGF21 has a sex-specific role in calorie-restriction-induced beiging of white adipose tissue in mice. Aging Biology.
15. Liśkiewicz, D., Liśkiewicz, A., Grabowski, M., Nowacka-Chmielewska, M.M., Jabłońska, K., Wojakowska, A., Marczak, Ł., Barski, J.J., and Małecki, A. (2021). Upregulation of hepatic autophagy under nutritional ketosis. J Nutr Biochem 93, 108620. 10.1016/j.jnutbio.2021.108620.
16. Gómora-García, J.C., Montiel, T., Hüttenrauch, M., Salcido-Gómez, A., García-Velázquez, L., Ramiro-Cortés, Y., Gomora, J.C., Castro-Obregón, S., and Massieu, L. (2023). Effect of the Ketone Body, D-β-Hydroxybutyrate, on Sirtuin2-Mediated Regulation of Mitochondrial Quality Control and the Autophagy-Lysosomal Pathway. Cells 12. 10.3390/cells12030486.
17. Abdelhady, R., Saber, S., Ahmed Abdel-Reheim, M., Mohammad, S.A.M., Alfaifi, J., M, I.E.A., L, A.S., A, I.F., E, A.E., H, S.E.-W., et al. (2023). Unveiling the therapeutic potential of exogenous β-hydroxybutyrate for chronic colitis in rats: novel insights on autophagy, apoptosis, and pyroptosis. Front Pharmacol 14, 1239025. 10.3389/fphar.2023.1239025.
18. Roohy, F., Siri, M., Kohansal, K., Ghalandari, A., Rezaei, R., Maleki, M.H., Shams, M., Monsef, A., and Dastghaib, S. (2024). Targeting apoptosis and unfolded protein response: the impact of β-

hydroxybutyrate in clear cell renal cell carcinoma under glucose-deprived conditions. Mol Biol Rep *51*, 168. 10.1007/s11033-023-08977-2.

19. Heidt, C., Fobker, M., Newport, M., Feldmann, R., Fischer, T., and Marquardt, T. (2023). Beta-Hydroxybutyrate (BHB), Glucose, Insulin, Octanoate (C8), and Decanoate (C10) Responses to a Medium-Chain Triglyceride (MCT) Oil with and without Glucose: A Single-Center Study in Healthy Adults. Nutrients *15*. 10.3390/nu15051148.

20. Norgren, J., Sindi, S., Sandebring-Matton, A., Kåreholt, I., Daniilidou, M., Akenine, U., Nordin, K., Rosenborg, S., Ngandu, T., and Kivipelto, M. (2020). Ketosis After Intake of Coconut Oil and Caprylic Acid-With and Without Glucose: A Cross-Over Study in Healthy Older Adults. Front Nutr *7*, 40. 10.3389/fnut.2020.00040.

21. Shoji, H., Kunugi, H., and Miyakawa, T. (2022). Acute and chronic effects of oral administration of a medium-chain fatty acid, capric acid, on locomotor activity and anxiety-like and depression-related behaviors in adult male C57BL/6J mice. Neuropsychopharmacol Rep *42*, 59-69. 10.1002/npr2.12226.

22. Kraeuter, A.K., Mashavave, T., Suvarna, A., van den Buuse, M., and Sarnyai, Z. (2020). Effects of beta-hydroxybutyrate administration on MK-801-induced schizophrenia-like behaviour in mice. Psychopharmacology (Berl) *237*, 1397-1405. 10.1007/s00213-020-05467-2.

23. Yum, M.S., Ko, T.S., and Kim, D.W. (2012). Anticonvulsant Effects of β-Hydroxybutyrate in Mice. J Epilepsy Res *2*, 29-32. 10.14581/jer.12008.

24. Sun, W., Wen, M., Liu, M., Wang, Q., Liu, Q., Li, L., Siebert, H.C., Loers, G., Zhang, R., and Zhang, N. (2022). Effect of β-hydroxybutyrate on behavioral alterations, molecular and morphological changes in CNS of multiple sclerosis mouse model. Front Aging Neurosci *14*, 1075161. 10.3389/fnagi.2022.1075161.

25. Hu, L., Xia, X., Zong, Y., Gu, Y., Wei, L., and Yin, J. (2022). Calorie Restriction Enhanced Glycogen Metabolism to Compensate for Lipid Insufficiency. Molecular Nutrition & Food Research *66*, 2200182. https://doi.org/10.1002/mnfr.202200182.

26. Lin, A.L., Zhang, W., Gao, X., and Watts, L. (2015). Caloric restriction increases ketone bodies metabolism and preserves blood flow in aging brain. Neurobiol Aging *36*, 2296-2303. 10.1016/j.neurobiolaging.2015.03.012.

27. Thomas, J.A., 2nd, Antonelli, J.A., Lloyd, J.C., Masko, E.M., Poulton, S.H., Phillips, T.E., Pollak, M., and Freedland, S.J. (2010). Effect of intermittent fasting on prostate cancer tumor growth in a mouse model. Prostate Cancer Prostatic Dis *13*, 350-355. 10.1038/pcan.2010.24.

28. Borer, K.T. (2024). Twice-Weekly 36-Hour Intermittent Fasting Practice Attenuates Hunger, Quadruples ß-Hydroxybutyrate, and Maintains Weight Loss: A Case Report. Cureus *16*, e57979. 10.7759/cureus.57979.

29. Johnson, J.B., Summer, W., Cutler, R.G., Martin, B., Hyun, D.H., Dixit, V.D., Pearson, M., Nassar, M., Telljohann, R., Maudsley, S., et al. (2007). Alternate day calorie restriction improves clinical findings and reduces markers of oxidative stress and inflammation in overweight adults with moderate asthma. Free Radic Biol Med *42*, 665-674. 10.1016/j.freeradbiomed.2006.12.005.

30. Harvie, M.N., Pegington, M., Mattson, M.P., Frystyk, J., Dillon, B., Evans, G., Cuzick, J., Jebb, S.A., Martin, B., Cutler, R.G., et al. (2011). The effects of intermittent or continuous energy restriction on weight loss and metabolic disease risk markers: a randomized trial in young overweight women. Int J Obes (Lond) *35*, 714-727. 10.1038/ijo.2010.171.

31. Huffman, K.M., Parker, D.C., Bhapkar, M., Racette, S.B., Martin, C.K., Redman, L.M., Das, S.K., Connelly, M.A., Pieper, C.F., Orenduff, M., et al. (2022). Calorie restriction improves lipid-related emerging cardiometabolic risk factors in healthy adults without obesity: Distinct influences of BMI and sex from CALERIE™ a multicentre, phase 2, randomised controlled trial. EClinicalMedicine *43*, 101261. 10.1016/j.eclinm.2021.101261.

32. Ferguson, B.S., Sahoo, P., McGrail, E., Francois, A., and Stratton, M.S. (2022). Modestly Increased Incidence of Ketosis in Caloric Restriction Does not Significantly Alter the Effects of Caloric Restriction. The Journal of nutrition, health and aging *26*, 657-662. https://doi.org/10.1007/s12603-022-1815-7.

33. Shimazu, T., Hirschey, M.D., Newman, J., He, W., Shirakawa, K., Le Moan, N., Grueter, C.A., Lim, H., Saunders, L.R., Stevens, R.D., et al. (2013). Suppression of oxidative stress by β-hydroxybutyrate, an endogenous histone deacetylase inhibitor. Science *339*, 211-214. 10.1126/science.1227166.

34. Huang, H., Zhang, D., Weng, Y., Delaney, K., Tang, Z., Yan, C., Qi, S., Peng, C., Cole, P.A., Roeder, R.G., and Zhao, Y. (2021). The regulatory enzymes and protein substrates for the lysine β-hydroxybutyrylation pathway. Sci Adv *7*. 10.1126/sciadv.abe2771.

35. Feola, K., Venable, A.H., Broomfield, T., Villegas, M., Fu, X., Burgess, S., and Huen, S.C. (2024). Hepatic ketogenesis is not required for starvation adaptation in mice. Molecular Metabolism *86*, 101967. https://doi.org/10.1016/j.molmet.2024.101967.

36. Chen, H.C., Liu, Y.W., Chang, K.C., Wu, Y.W., Chen, Y.M., Chao, Y.K., You, M.Y., Lundy, D.J., Lin, C.J., Hsieh, M.L., et al. (2023). Gut butyrate-producers confer post-infarction cardiac protection. Nat Commun *14*, 7249. 10.1038/s41467-023-43167-5.

37. Thomas, L.K., Ittmann, M., and Cooper, C. (1982). The role of leucine in ketogenesis in starved rats. Biochem J *204*, 399-403. 10.1042/bj2040399.

38. Hadem, I.K.H., Majaw, T., Kharbuli, B., and Sharma, R. (2019). Beneficial effects of dietary restriction in aging brain. J Chem Neuroanat *95*, 123-133. 10.1016/j.jchemneu.2017.10.001.

39. Lanza, I.R., Zabielski, P., Klaus, K.A., Morse, D.M., Heppelmann, C.J., Bergen, H.R., 3rd, Dasari, S., Walrand, S., Short, K.R., Johnson, M.L., et al. (2012). Chronic caloric restriction preserves mitochondrial function in senescence without increasing mitochondrial biogenesis. Cell Metab *16*, 777-788. 10.1016/j.cmet.2012.11.003.

40. Mulvey, L., Wilkie, S.E., Borland, G., Griffiths, K., Sinclair, A., McGuinness, D., Watson, D.G., and Selman, C. (2021). Strain-specific metabolic responses to long-term caloric restriction in female ILSXISS recombinant inbred mice. Mol Cell Endocrinol *535*, 111376. 10.1016/j.mce.2021.111376.

41. Mitchell, S.J., Madrigal-Matute, J., Scheibye-Knudsen, M., Fang, E., Aon, M., Gonzalez-Reyes, J.A., Cortassa, S., Kaushik, S., Gonzalez-Freire, M., Patel, B., et al. (2016). Effects of Sex, Strain, and Energy Intake on Hallmarks of Aging in Mice. Cell Metab *23*, 1093-1112. 10.1016/j.cmet.2016.05.027.

42. Liao, C.Y., Rikke, B.A., Johnson, T.E., Diaz, V., and Nelson, J.F. (2010). Genetic variation in the murine lifespan response to dietary restriction: from life extension to life shortening. Aging Cell *9*, 92-95. 10.1111/j.1474-9726.2009.00533.x.

43. Pearson, K.J., Lewis, K.N., Price, N.L., Chang, J.W., Perez, E., Cascajo, M.V., Tamashiro, K.L., Poosala, S., Csiszar, A., Ungvari, Z., et al. (2008). Nrf2 mediates cancer protection but not prolongevity induced by caloric restriction. Proc Natl Acad Sci U S A *105*, 2325-2330. 10.1073/pnas.0712162105.

44. Yu, D., Tomasiewicz, J.L., Yang, S.E., Miller, B.R., Wakai, M.H., Sherman, D.S., Cummings, N.E., Baar, E.L., Brinkman, J.A., Syed, F.A., and Lamming, D.W. (2019). Calorie-Restriction-Induced Insulin Sensitivity Is Mediated by Adipose mTORC2 and Not Required for Lifespan Extension. Cell reports *29*, 236-248 e233. 10.1016/j.celrep.2019.08.084.

Figure 1.

Figure Legends

Figure 1. The ablation of ketogenesis does not alter the effect of caloric restriction on body weight and body composition.

(A) Main experimental setup of the caloric restriction study. **(B-E)** Western blot confirmation of the ablation of Hmgcs2 in the KO animals of both AL and CR groups in the liver (B), the small intestine jejunum and ileum (C), the large intestine colon (D), and the kidney (E). **(F-H)** Male mice body weight (F), body composition % (G), and final changes in body composition (H) over 7 months. For (F) and (G), at the beginning of the experiment, WT-AL, KO-AL, WT-CR, KO-CR, n=7,6,8,7 respectively; for (H) n=5,5,8,6. **(I-K)** Female mice body weight (I), body composition % (J), and final changes in body composition (K) over 7 months. For (I) and (J), at the beginning of the experiment, WT-AL, KO-AL, WT-CR, KO-CR, n=6,7,7,9 respectively; for (K) n=6,6,7,8. (H & K) *p<0.05, ***p<0.001, ****p<0.0001, Sidak's test post 2-way ANOVA. Data presented as mean ± SEM.

Figure 2.

Figure 2. The ablation of ketogenesis selectively improves insulin sensitivity in females.

(A-D) After 21 hrs fasting in male mice, the results of glucose tolerance test (A; 1 g/kg; I. P.), insulin tolerance test (B; 0.5 U/kg; I. P.), pyruvate tolerance test (C; 2 g/kg; I. P.), and HOMA-IR measurements (D). For WT-AL, KO-AL, WT-CR, and KO-CR in each experiment, GTT n=7,6,8,7; ITT n=7,6,8,6; PTT n=7,6,8,6; HOMA-IR n=7,5,8,7. **(E-H)** After 21 hrs fasting in female mice, the results of glucose tolerance test (E; 1 g/kg; I. P.), insulin tolerance test (F; 0.5 U/kg; I. P.), pyruvate tolerance test (G; 2 g/kg; I. P.), and HOMA-IR measurements (H). For WT-AL, KO-AL, WT-CR, and KO-CR in each experiment, GTT n=6,7,7,8; ITT n=6,7,6,8; PTT n=6,7,7,8; HOMA-IR n=9,7,8,7. (A-H) *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, Sidak's test post 2-way ANOVA. Data presented as mean ± SEM.

Figure 3.

Figure 3. The ablation of ketogenesis induce sex-specific changes in RER.

(A-B) Male mice energy expenditure (A) and respiratory exchange ratio (B) expressed over a 24 hrs period and binned by light/dark cycle. For WT-AL, KO-AL, WT-CR, and KO-CR, n=7,6,8,7. **(C-D)** Female mice energy expenditure (C) and respiratory exchange ratio (D) expressed over a 24 hrs period and binned by light/dark cycle. For WT-AL, KO-AL, WT-CR, and KO-CR, n=6,7,7,8.

(A-D) *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, Sidak's test post 2-way ANOVA, completed separately for each cycle. Data presented as mean ± SEM.

Figure 4.

Figure 4. Loss of ketogenesis does not alter the effects of CR on iWAT physiology.

(A-C) In male mice, representative H&E staining of the iWAT (A), iWAT weight (B) and the average size of adipocytes (C). For WT-AL, KO-AL, WT-CR, and KO-CR, iWAT weight n=5,3,5,4; adipocyte size n=4,3,4,4. **(D-E)** Male iWAT gene expression of *Ucp1* (D), n=6 each group; iWAT gene expression level of lipid processing genes (E) *Dgat1*, *Elovl3*, *Cidea*, *Fasn*, and *Acc1*. n=6 each group for *Dgat*1, *Fasn*, and *Acc1*; for WT-AL, KO-AL, WT-CR, and KO-CR, Elovl3 n=5,6,6,5; *Cidea* n=6,5,6,5. **(F-H)** In female mice, representative H&E staining of the iWAT (F), iWAT weight (G) and the average size of adipocytes (H). For WT-AL, KO-AL, WT-CR, and KO-CR, iWAT weight n=5,4,5,3; adipocyte size n=4 each group. **(I-J)** Female iWAT gene expression of *Ucp1* (I). For WT-AL, KO-AL, WT-CR, and KO-CR, n=6,5,6,6. Female iWAT gene expression level of lipid processing genes (J) *Dgat1*, *Elovl3*, *Cidea*, *Fasn*, and *Acc1*. n=6 each group for *Dgat1* and *Acc1*; *Elovl3* n=4,6,4,6; *Cidea* n=6,6,5,6; Fasn n=6,6,4,6. (B-E & G-J) *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, Sidak's test post 2-way ANOVA (E & J) conducted separately for each gene. Data presented as mean ± SEM.

Figure 5.

Figure 5. Caloric restriction upregulates autophagy-related proteins in the liver regardless of ketogenesis.

(A-B) Liver western blots in male (A) and female (B) mice for p62 and LC3. n=9 each group. *$p < 0.05$, Sidak's test post 2-way ANOVA conducted separately for each protein. Data presented as mean ± SEM.

Figure 6.

Figure 6. Acclimation to a recurrent fasting caloric restriction protocol suppresses ketogenesis.

(A) Circulating βHB level of male + female mice measured every 4 hrs under typical experimental conditions (food present for AL groups/CR groups fed at 6 am). For WT-AL, KO-AL, WT-CR, and KO-CR, n=8,8,14,8. **(B)** Circulating βHB level of typically *ad libitum*-fed male + female mice fasting starting at 6 am. n=8 each group.

(C) Diagram of the CR acclimation experiment using a separate cohort of mice. **(D-E)** Fasting βHB level of male (D) and female (E) mice during the acclimation phase of the CR protocol. For WT-AL, KO-AL, WT-CR, and KO-CR, male n=6,7,4,5; female n=5,4,5,5. (A & D-E) *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$, Sidak's test post 2-way ANOVA. (B) *$p<0.05$, t-test. Data presented as mean ± SEM.

Figure 7.

Figure 7. Caloric restriction enhances ketogenesis and ketolysis capacity.

(A-B) Lipid tolerance test using MCT oil (5 ml/kg; P. O.) in male (A) and female (B) mice. Circulating βHB is unchanged in the KO mice of both sexes while CR significantly enhances ketogenic ability in the lipid tolerance test. For WT-AL, KO-AL, WT-CR, and KO-CR, male n=5,3,5,5; female n=5 each group. **(C-D)** Ketone tolerance test using sodium 3-β-hydroxybutyrate in saline (2 g/kg; I. P.) 4 hrs after CR-feeding and AL fasting. Ketone consumption is improved in male (C) and female (D) mice treated with CR. Male n=5 each group; for WT-AL, KO-AL, WT-CR, and KO-CR, female n= 5,5,4,5. **(E-F)** Heatmap representation of the discrete dataset in this study for male (E) and female (F), expressed as Z-score comparisons within individual variables, sorted by the difference of WT-CR vs KO-CR. (A-D) *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, Sidak's test post 2-way ANOVA. Data presented as mean ± SEM.