Microtubule anchoring and coupling of CD20 to the RhoA/Rock1 pathway

Kathrin Kläsener[1,2,3,a,✉], Cindy Eunhee Lee[4,9,a], Julian Bender[5], Angela Naumann[2,6], Lena
Reimann[1,7], Geoffroy Andrieux[8], Nadja Herrmann[2], Roland Nitschke[2,6], Klaus Warnatz[3,9],
Bettina Warscheid[5], Reinhard E. Voll[3], Michael Reth[1,2,3,✉]

[1] Biology III, Faculty of Biology, University of Freiburg, Freiburg, Germany
[2] Signaling Research Centers BIOSS and CIBSS, University of Freiburg, Freiburg, Germany
[3] Department of Rheumatology and Clinical Immunology, Medical Center –Faculty of Medicine,
University of Freiburg, Freiburg, Germany
[4] Faculty of Science and Technology, University of Canberra, Canberra, Australia
[5] Biochemistry II, Theodor-Boveri-Institute, Biocenter, Faculty of Chemistry and Pharmacy, University of
Würzburg, Würzburg, Germany
[6] Life Imaging Center and Signalling Research Centres CIBSS and BIOSS, University of Freiburg,
Freiburg, Germany.
[7] Celonic AG, Basel, Switzerland
[8] Institute of Medical Bioinformatics and Systems Medicine, Medical Center - Faculty of Medicine,
University of Freiburg, Freiburg, Germany
[9] Center for Chronic Immunodeficiency (CCI), Medical center - Faculty of medicine, University of
Freiburg, Freiburg, Germany
[a] equal contributors

* Correspondence:

✉Dr. Michael Reth. Department of Rheumatology and Clinical Immunology, Medical Center University
of Freiburg, Germany and BIOSS, Signalhaus Freiburg, Schänzlestr.18 D-79104 Freiburg,
Tel.: +49 761 2032718, E-mail: michael.reth@bioss.uni-freiburg.de
24
✉Dr. Kathrin Kläsener. Department of Rheumatology and Clinical Immunology, Medical Center
University of Freiburg, Germany and BIOSS, Signalhaus Freiburg, Schänzlestr.18 D-79104 Freiburg,
Tel.: +49 761 2032730, E-mail: kathrin.klaesener@bioss.uni-freiburg.de

Abstract

CD20 is a B cell-specific four-helix transmembrane protein and a prominent target of
therapeutic anti-CD20 antibodies. We have recently described that CD20 is localized within a
membrane nanocluster harboring the IgD class B cell antigen receptor (IgD-BCR) where it
functions as a gatekeeper for the resting state of naïve human B lymphocytes. How CD20
exerts its gatekeeper function is not yet known. Using Ramos and human peripheral blood B
cells, we show here that the serine/threonine kinase PKCδ, constitutively phosphorylates
serine residues at the cytosolic tails of CD20. The phosphorylated CD20 becomes a target for
14-3-3 adaptor proteins that link CD20 to the RhoA GDP/GTP exchange factor GEF-H1 and
the microtubule (MT) network controlling the stability of the IgD-BCR nanocluster. Binding of
anti-CD20 antibodies results in MT disassembly and the replacement of the GEF-H1/CD20
complex by a RhoA-GTP/ROCK1/CD20 complex which drives actomyosin contractility. Our
study suggests that CD20 not only maintains the resting state, but also orchestrates the
MT/actin switch in active B lymphocytes. This could have implications for treatment with anti-
CD20 antibodies and may help to optimize therapeutic protocols.

Keywords: anti-CD20 therapeutic antibody / B lymphocyte / MT network / PKCδ / RhoA-
Rock1 pathway.

Introduction

Upon a successful production of an immunoglobulin heavy and light chain pair, pre-B cells
become immature B cells expressing an IgM-class B cell antigen receptor (IgM-BCR). The
IgM-BCR plays an important role for the central tolerance process where immature B cells
undergo negative selection to eliminate self-reactive clones. After surviving selection,
immature B cells further develop into transitional B cells that migrate to secondary lymphoid
organs and then into mature naive B cells co-expressing an IgM and IgD-BCR[1]. According to
the clonal selection theory, mature B cells only become fully activated upon binding to their
cognate antigen. Thus, most mature naive B cells remain in a dormant state and the IgD-BCR
plays an important role for the maintenance of this resting state. We have previously shown
that the IgM-BCR and IgD-BCR are localized in different membrane nanoclusters[2]. The IgD-
BCR nanocluster has a raft-type lipid composition and harbors important B cell coreceptors
such as CD19, CD20, CD40 and the BAFF receptor[3–6]. Interestingly, most proteins residing
within the IgD nanocluster including the IgD-BCR reach their full expression level only at the
mature naive B cell stage. This is also the case for CD20, a 37 kD non-glycosylated
phosphoprotein with four transmembrane domains that form dimers or higher oligomers on the
B cell surface[7–9]. The CD20 protein is encoded by the MS4A1 gene and a member of the MS4A
gene family. Due to the lack of a known ligand the biological function of CD20 has been
regarded as an enigma of B cell biology[10,11]. However, newer data show that CD20 is a highly
interactive protein that helps in the regulation and functional organization of other receptors on
the B cell surface[12]. For example, CD70 has recently been shown to require the presence of
CD20 for immune synapse formation and efficient T: B collaboration[13].

CD20 is best known as a prominent target of therapeutic monoclonal antibodies (mAb) such
as rituximab (RTX), obinutuzumab (GA101), or ocrelizumab used for the treatment of B cell
chronic lymphocytic leukemia (B-CLL), malignant B cell lymphomas or human autoimmune
diseases such as rheumatoid arthritis (RA), systemic lupus erythematosus (SLE) and, more
recently, multiple sclerosis (MS)[14–18]. The anti-CD20 mAbs are thought to eliminate targeted B
cells by antibody-dependent cellular cytotoxicity (ADCC) mechanisms, phagocytosis[19], or
induction of cell cycle arrest and initiation of an apoptosis program in some B cell lines[20]. The
success of the anti-CD20 immunotherapy could be further improved by a more profound insight
into the biological function of CD20. An important advance for this goal is the finding that CD20
is part of the IgD-BCR nanocluster where it functions as a gatekeeper for the resting state of
naïve human B lymphocytes[21]. Here, we show that constitutively phosphorylated serine
residues in the N- and C-terminal cytosolic tails of CD20 play an important role for this

gatekeeper function by coupling CD20 to the microtubule (MT) network and the RhoA/ROCK1
signaling pathway.
81
82 **Results**
83
CD20 is a substrate of PKCδ acting as a gatekeeper for the resting state of B cells
The resting state of naïve human B lymphocytes is maintained by several negative regulators
including PKCdelta (PKCδ), a member of the novel serine/threonine protein kinase C (PKC)
family[22,23]. PKCδ has been found to be constitutively associated with the BCR signaling subunit
Igα (CD79a) and negatively regulates the BCR signaling[24]. To study the gatekeeper function
of PKCδ we employed the CRISPR/Cas9 method to generate two PKCδ-deficient (PKCδ KO)
clones of the human Burkitt lymphoma cell line Ramos (Fig. S1A and B). The PKCδ KO Ramos
cells were viable and could be maintained in culture. We first compared the expression of B
cell marker proteins on the surface of the PKCδ KO line to the wild-type (wt) Ramos cells by
flow cytometric analysis (Fig. 1A). Normalization of the protein expression levels on PKCδ KO
to Ramos wt cells shows that the loss of PKCδ is associated with a 25-fold increase of the B
cell activation marker CD69 on Ramos B cells. The PKCδ deficiency also results in the
upregulation of membrane proteins that are associated with the IgD-class nanocluster on
resting B cells, namely CD20, CD19, CD40 and the IgD-BCR, whereas proteins that are
associated with the IgM-BCR, such as CD45 are downregulated. In addition, two membrane
proteins that function as negative B cell signaling regulators, namely CD22 and the IgG-binding
Fc receptor FcγRIIb are upregulated in the PKCδ KO Ramos line (Fig. 1A). The 20-fold
upregulation of the inhibitory FcγRIIb (CD32b) is interesting because mutations of CD32b are
associated with human autoimmune diseases such as systemic lupus erythematosus (SLE)[25].
In summary, this study shows that PKCδ KO Ramos B cells are activated and counteract this
activation by upregulating the IgD-class nanocluster and negative signaling regulators. These
findings support the notion that, similar to CD20, PKCδ acts as a gatekeeper for the "resting
state" of mature human B cells[22].
Next, we conducted a phosphoproteome analysis to monitor the difference in the level of
constitutive serine/threonine phosphorylation in Ramos wt and the PKCδ KO cell line (Fig. 1B).
In this analysis we identified several serine residues in the N-terminal and C-terminal cytosolic
tail of CD20 that are constitutively phosphorylated by PKCδ. Thus, PKCδ and CD20, two
gatekeepers of the resting state of B lymphocytes, are connected by an enzyme/substrate
relationship. Global serine/threonine phospho-proteome analysis of Ramos wt and PKCδ KO
cells also reveal specific alterations in the phosphorylation of components of the RhoA/Rock-
1 pathway and the MT cytoskeletal network described below (Fig. S1C). A comparison of the
RNA-seq data obtained from the two cell lines indicated that, compared to Ramos wt, the PKCδ
KO cells displayed an increased expression of genes encoding for proteins involved in MT
depolymerization and a loss of cortical cytoskeletal stability (Fig. S2A and S2B).

CD32b is associated with the IgD-BCR nanocluster and stabilizes CD20 expression

The 20-fold upregulation of CD32b on the surface of PKCδ KO Ramos B cells suggests that this inhibitory Fc receptor also has a constitutive gatekeeper function for the resting B cell stage[26]. To test for this, we generated Ramos clones with a CD32b deficiency (CD32b KO) or CD32b overexpression (CD32b-tr). Ramos wt cells express very low amounts of CD32b on their surface, but the CD32b KO line clearly showed that this level can be further reduced to background levels (Fig. 2A). Transfection of Ramos B cells with a human CD32b lentiviral expression vector resulted in a 500-fold increase of CD32b on the surface of CD32b-tr Ramos cells (Fig. 2B). To test for alterations of B cell marker proteins on CD32b-deficient or - overexpressing Ramos cells (CD32-tr), we directly compared these cells with Ramos wt cells in a flow cytometric analysis. Loss of CD32b resulted in a reduced expression of proteins associated with the IgD-class nanocluster such as CD20, CD40, CXCR4, and the IgD-BCR, and a higher expression of CD19 and the IgM-BCR, as well as the B cell activation marker CD69 (Fig. 2C). Notably, CD19 and the IgM-BCR interact with each other on activated B lymphocytes and the expression pattern on CD32b KO Ramos cells suggests that the loss of CD32b results in a partial B cell activation[27]. The comparison of CD32b-tr to Ramos wt cells shows an opposite phenotype, namely the increased expression of proteins associated with the IgD-class nanocluster (Fig. 2D). The effect of the CD32b expression on the stability of the IgD-class nanocluster suggests that CD32b is part of this membrane nanocluster. To test for this, we employed the Fab-based proximity ligation assay (Fab-PLA), which monitors the 10-20 nanometer proximity between the IgD-BCR and CD32b on the cell surface (Fig. 2E and 2F)[28]. The IgD:CD32b PLA signal monitored from Ramos wt is drastically increased upon CD32b overexpression in CD32-tr, reaching a level that is similar to the PLA signal monitored by human healthy donor (HD) B cells. As a negative control for this analysis, we tested Ramos CD32b KO cells, which show no PLA signal. The comparable IgD-BCR/CD32b PLA signal obtained from the analysis of CD32b-tr and HD B cells suggests a similar expression of CD32b on these two cell types. This was verified by a comparative flow cytometry analysis of Ramos CD32b-tr and HD B cells that also confirmed the higher CD20 expression on the Ramos CD32b-tr B cells (Fig. S3). Because of these features and the more stable IgD-class nanocluster we chose CD32b-tr Ramos B cells for our further studies of the cellular function of CD20.

Function of CD20 serine residues and their association with 14-3-3 adaptor proteins

The serine/threonine phosphoproteome analysis of Ramos and its PKCδ-deficient subclones suggests that 4 serine residues in the N-terminal cytosolic tail (S7, S35/36, S49) and 3 serine residues in the C-terminal cytosolic tail (S225, S253/254) of CD20 are constitutively phosphorylated by PKCδ. Interestingly, the three most significant CD20 serine phosphorylation sites (S35/36, S225) are found in the context of an RxxS motif that, once phosphorylated,

could become a binding target of the 14-3-3 adaptor family members, which function by binding
to serine/threonine-phosphorylated intracellular proteins and altering the conformation,
activity, and subcellular localization of their binding partners (Fig. 3A)[29–31]. Additionally, another
serine residue (S221) is located near the plasma membrane. Neither the phosphorylated nor
the unphosphorylated form of S221 was detected in the phosphoproteome, perhaps because
of an adjacent palmitoylation at C220[32]. S221 lies within a classical RxxSxP binding motif for
14-3-3. To eliminate all possibilities of 14-3-3 binding, S221 was also mutated to alanine. To
test for the function of the identified serine residues, we generated expression vectors for WT
or mutant CD20 proteins carrying a serine to alanine (S/A) exchange of either the 4 N-terminal
(Nmut), the 4 C-terminal (Cmut), or all 8 serine residues (NCmut). For purification purposes,
the expressed CD20 proteins also carry a C-terminal flag-tag (Fig. 3B). Next, we employed the
CRISPR/Cas-9 method to generate CD20-deficient Ramos B cells that were transfected with
retroviral expression vectors encoding either CD20 WT or tail mutants (Nmut, Cmut, or NCmut)
of CD20. The obtained CD20 transfectants were sorted and further transfected with the human
CD32b lentiviral expression vector. Flow cytometry analysis showed that all four double-
transfectants expressed CD20 and CD32b at levels similar to the CD32b-tr Ramos line (Fig.
3C). A similar expression was also found for CD19 as well as the IgM- and IgD-BCR. However,
in comparison to the CD32b-tr Ramos line and the CD20 WT transfectants, Ramos cells
expressing the Nmut, Cmut, or NCmut CD20 mutant displayed a higher expression of the
activation marker CD83 suggesting that the S/A exchange partially compromises the
gatekeeper function of CD20 (Fig. 3C, far right)[33].
The finding that 4 of the 8 serine residues under study are part of an RxxS motif sequence
suggests that CD20 is not only constitutively phosphorylated by PKCδ but also constitutively
associated with a 14-3-3 adaptor protein. To test for this, we used the 1-PLA assay to monitor
the colocalization of CD20 with a 14-3-3 adaptor protein in CD32b-tr Ramos cells and
CD32b/CD20 double-transfectants expressing in addition to CD32b, either CD20 WT or one
of three different tail mutants of CD20 (Fig. 3D and 3E). The 1-PLA assay employs CD20 and
14-3-3 specific DNA-coupled antibodies that detect a colocalization of the two proteins in the
10-40 nanometer range[28]. A colocalization of CD20 with 14-3-3 is detected in the CD20 WT,
but to a lesser extent in the CD20 Nmut, Cmut or NCmut Ramos cells. Western blot analysis
of anti-flag immunoprecipitates of Ramos cell lysates confirms that, compared to CD20 WT,
the S/A mutants of CD20 are less well associated with the 29 kD 14-3-3 protein (Fig. 3F).
189
The GEF-H1 activity is regulated by the cytoplasmic tails of CD20
14-3-3 is a dimeric adaptor protein that can stabilize a CD20 homodimer and/or the
heterodimeric association of CD20 with other RxxS motif-containing proteins. In a preliminary
study, we found that exposure of B cells to anti-CD20 antibodies activates the RhoA/Rock1
signaling pathway (see below). A key activator of this pathway is the GDP/GTP exchange
protein GEF-H1 (also known as ARHGEF2), which is constitutively associated with the MT

network in resting B cells[34–39]. The intrinsically disordered regions in the C-terminal tail of GEF-
H1[35] carries an RRxxR and an RxxSxP motif (Fig. 4A). The RRxxR motif is a known binding
target for light chains of the dynein motor complex (such as LC8 or Tctex-1) and is thought to
be one of the interfaces linking GEF-H1 to MTs[37]. The serine residue S886 within the RxxSxP
motif of GEF-H1 is phosphorylated by the p21-activated kinase-2 (PAK2), a modification that
is known to inhibit GEF-H1 activity[40]. Binding of a 14-3-3 adaptor protein to the phosphorylated
RxxSxP motif may connect GEF-H1 to the constitutively phosphorylated CD20 tails. To test
for this, we used the 1-PLA assay and found a colocalization of CD20 and GEF-H1 in the CD20
WT transfectant and to a lesser extent in the CD20 Nmut, Cmut or NCmut transfected Ramos
cells (Fig. 4B and 4C). Western blot analysis of anti-flag immunoprecipitates of CD20 proteins
in the Ramos cell lysates confirmed that the tail mutants of CD20 are less well associated with
GEF-H1 than CD20 WT (Fig. 4D). In particular, the NC double-mutant of CD20 is hardly
associated with GEF-H1, compared to CD20 WT.
The amount of the GDP/GTP exchange active GEF-H1 protein in the cytosol of Ramos B cells
can be tested by a pulldown assay employing a G17A trapping mutant of RhoA, which
stabilizes the intermediate RhoA/GEF-H1 interaction[41]. For this, lysates of the different Ramos
cell lines were incubated with GST-RhoA-G17A-coupled GST beads, and the bound GEF-H1
was monitored by western blotting. Compared to CD20 WT, the cytosol of Ramos cells
expressing the Nmut, Cmut or NCmut contains a higher amount of active GEF-H1/RhoA
complexes (Fig. 4E and 4F). Taken together, these data suggest that the constitutively
phosphorylated serine residues within the cytosolic tails of CD20 regulate the RhoA/Rock-1
pathway via the binding and sequestering of GEF-H1.
218
Exposure to anti-CD20 antibody activates the RhoA/Rock-1 pathway
We next asked whether or not the exposure of Ramos B cells to a therapeutic anti-CD20
antibody such as rituximab (RTX) alters the interaction of CD20 with components of the
RhoA/Rock-1 pathway. For this we employed the 1-PLA assays to monitor changes in the
proximity of CD20 to 14-3-3, GEF-H1, RhoA and Rock-1 in Ramos CD32b-tr as well as CD20
WT or CD20 NCmut transfectants before or after a 5 min exposure to RTX (Fig. 5). The RTX
treatment reduced the CD20/14-3-3 proximity in CD32b-tr and CD20 WT transfected Ramos
cells to the level of unstimulated NCmut Ramos cells (Fig. 5A and 5B) and the same was found
for the CD20/GEF-H1 proximity (Fig. 5C and 5D). Interestingly, the 1-PLA assays show an
increased proximity of CD20 and RhoA (Fig. 5E and 5F) as well as CD20 and Rock-1 in RTX-
treated Ramos cells expressing CD20 WT but not those expressing CD20 NCmut. This study
suggests that in RTX-treated B cells, CD20 stabilizes a RhoA/Rock-1 complex at the inner
leaflet of the plasma membrane in a 14-3-3-dependent manner[42,43]. Importantly, the C-
terminus of Rock-1 also carries an evolutionary conserved RxxS motif. Thus, it is feasible that
CD20 functions not only as a gatekeeper of the resting state, but also as a switch that promotes
activation of the RhoA/Rock-1 pathway in antigen- or RTX-exposed B cells. As a control, we

determined whether active GEF-H1 is formed after RTX activation following GDP/GTP
exchange. The pull-down assay with the G17A trapping mutant of RhoA shows an increase in
active GEF-H1/RhoA complexes, especially in Ramos, CD32b-tr and WT cells (Fig. S4A and
4B).
We then used the phosphoflow assays to monitor alteration in the phosphorylation of members
of the RhoA/Rock-1 pathway after the RTX treatment of CD32b-tr, CD20 WT or NCmut Ramos
cells (Fig. S5). Exposure of CD32b-tr or CD20 WT Ramos cells to RTX results in a reduction
of the S886 phosphorylation of GEF-H1 and the loss of the inhibitory S188 phosphorylation of
RhoA (Fig. S5A and 5B). The same treatment resulted in an increased phosphorylation of
tyrosine Y914 of Rock-1, a modification that is associated with Rock-1 activation (Fig.
S5C)[44,45]. Importantly, all these amino acids are less well phosphorylated in the NCmut Ramos
cells and their phosphorylation state is not significantly altered after exposure of the cells to
RTX. Together these data demonstrate that the cytosolic serine/threonine residues studied
connect CD20 to the RhoA/Rock-1 signaling pathway.
Exposure to the therapeutic antibody RTX activates the RhoA/Rock-1 signaling pathway not
only in Ramos but also naïve human B cells. This is indicated by the time-dependent alteration
of the phosphorylation status of GEF-H1, RhoA and Rock-1 upon the treatment of HD B cells
with 10 ug/mL of RTX (Fig. S5D-5F). The 1-PLA assays of HD B cells before or after a 5 min
exposure to RTX show a reduction of the 14-3-3 (Fig. 6A and 6B) and GEF-H1 (Fig. 6C and
6D) proximity to CD20 and at the same time a strong recruitment of RhoA (Fig. 6E and 6F)
and Rock-1 (Fig. 6G and 6H) in the vicinity of CD20 in the RTX-treated cells. Thus, the CD20
engagement not only liberates and activates GEF-H1 but also results in an increased coupling
of the engaged CD20 to the RhoA/Rock-1 signaling module. Interestingly not only the RTX
treatment but also exposure of HD B cells to a MT polymerization inhibitor such as nocodazole
(Noc) changes the CD20 association with components of the RhoA/Rock-1 signaling pathway
(Fig. 6E-6H).
261

Destabilization of MTs and the IgD-BCR nanocluster by anti-CD20 antibodies
It is well known that GEF-H1 is associated with the MT network and that MT destabilization
results of GEF-H1 liberation and increased activity[34,36–38,46,47]. The result of our 1-PLA assay
(see above) suggests that the activation of RhoA/Rock-1 signaling pathway by RTX is
mediated by the destabilization of the MT cytoskeleton. To test for this, we pretreated HD B
cells for 1 hour with either Noc or with the MT stabilizer docetaxel (Tax) and then measured
RTX-induced changes by flow cytometry (Fig. 7A and 7B). The RTX-induced reduction of
CD20 abundance and the increased Rock1 phosphorylation can be prevented by pretreating
the HD B cells for 1 hour with Tax, whereas the Noc-pretreated HD B cells show already an
Rock1 activation with no further increase by RTX. Thus, the immediate effect of the binding of
RTX to CD20 seems to be the destabilization of the MT network resulting in GEF-H1 liberation
and the activation of the RhoA/Rock-1 signaling pathway.

MT dynamics are tightly regulated. To test whether binding of RTX to CD20 can influence or
induce the destabilization of MTs, Ramos CD32b-tr cells were treated with RTX and labeled
with mouse anti-alpha-tubulin to visualize tubulin structures by immunofluorescence
microscopy. In the Airyscan Super-Resolution (SR) imaging results (Fig. S6) we could visualize
the peeling of MT polymers, the outwardly curved spiral protofilaments at the end of shortening
MT upon RTX exposure (Fig. S6A and B). In a second approach with HD naïve B cells (Fig.
S7A-C), Airyscan (SR) confocal images of the labeled α-tubulin showed a spatial distribution
of small "ram's-horn" MT intermediates upon RTX treatment (Fig. S7B)[48,49]. Interestingly, a
pretreatment with Tax before RTX exposure could stabilize the B cell integrity and reduce MT
depolymerization (Fig. S7C).

On resting B cells CD20 and CD19 reside in the same protein/lipid nanocluster in close
proximity to the IgD-BCR, and the exposure to RTX leads to IgD-BCR/CD19 dissociation and
the formation of an IgM-BCR/CD19 complex[21]. Taking advantage of the Fab-PLA technique,
we next asked whether or not the MT network is involved in the nanoscale receptor
organization and reformation. Indeed, the exposure of the HD B cells to either RTX or Noc
results in a dissolution of the IgD-BCR nanocluster and IgM-BCR/CD19 conjugation. The RTX-
induced nanoscale receptor rearrangements are partially prevented by the pretreatment of the
HD B cells with Tax (Fig. 7C-7F). Taken together, these data show that in resting B cells the
nanocluster of the IgD-BCR is connected to the MT network and that the cytosolic tails of CD20
play an important role in this connection and its formation.

Discussion

PKCδ and CD20: a kinase/substrate pair with a gatekeeper function
We previously have shown that CD20 is stabilizing the membrane nanocluster of the IgD-BCR
and acts as a gatekeeper for the resting state of B lymphocytes[21]. Here we describe that serine
residues within the cytoplasmic tails of CD20 play an important role for this gatekeeper
function. Specifically, we show that 6 serine residues of CD20 are constitutively
phosphorylated by the serine/threonine kinase PKCδ and that this modification couples CD20
to the MT network in resting and the RhoA/ROCK1 signaling pathway in activated B cells. The
exchange of the serine to alanine residues did not alter the amount of CD20 on the B cell
surface, but it has an impact on the B cell activation state as the CD20 mutated Ramos cells
displayed a higher expression of the CD83 activation marker[33]. These data suggest that the
phosphorylated serine residues are connected to the gatekeeper function of CD20. The notion
that PKCδ also acts as a gatekeeper for the resting state of B lymphocytes is supported by
several genetic data[22,24,50,51]. The PKCδ KO mice develop autoimmunity[52] and loss-of-function
mutations of PKCδ in humans are associated with autoimmune diseases such as systemic
lupus erythematosus (SLE) or a B cell lymphoproliferative syndrome[53]. Our finding that CD20

is a substrate of PKCδ provides a novel mechanistic insight in the development of these

autoimmune diseases.

PKCδ has multiple substrates at the membrane, in the cytosol and inside the nucleus[50]. Thus,

the targeting of PKCδ to a specific subcellular location may determine its biological function in

different settings. Like many other kinases, PKCδ is regulated by an autoinhibitory mechanism

where the N-terminal regulatory domain is folded over the C-terminal catalytic domain, thus

inhibiting the kinase activity. The specific targeting of PKCδ is connected with an opening of

the autoinhibited conformation and kinase activation. For example, the binding of the N-

terminal C1A/B domain to diacylglycerol (DAG) recruits the open, active form of PKCδ to the

plasma membrane[54]. The location and activity of PKCδ can also be regulated by

posttranslational modifications. Interestingly, the SRC-family kinase Lyn can phosphorylate

and bind to tyrosine 52 of PKCδ[55,56]. The formation of the Lyn/PKCδ complex activates both

kinases and places them at the inner leaflet of the plasma membrane. We have previously

found that, after an Igα phosphorylation, the IgD-BCR is associated with PKCδ[24]. It is thus

feasible that an active Lyn/PKCδ complex is formed within the IgD-BCR nanocluster, thereby

explaining the constitutive serine phosphorylation of CD20 at this location.

CD20 binds GEF-H1 and inhibits the GDP/GTP exchange activity in resting B cells

Four of the 8 mutated serine residues of CD20 are part of conserved RxxS or RxxSxP motifs

that are well-known binding sites for 14-3-3 adaptor proteins[57]. A constitutive CD20/14-3-3

association in resting human B cells was confirmed by a 1-PLA and immune precipitation. The

RxxS and RxxSxP sequence is localized at the CD20 N-terminus and C-terminus, respectively.

The binding of the dimeric 14-3-3 adaptor to these sequences can either alter the

intramolecular conformation of the two cytosolic tails of CD20 or stabilize the formation of a

CD20 homodimer. Alternatively, the 14-3-3 dimer could also connect CD20 to other signaling

molecules[58,59]. Using a 1-PLA and immune precipitation assays we confirmed that the

GDP/GTP exchange factor GEF-H1, which carries a conserved RxxSxP motif at its

unstructured C-terminus, is one of the binding partners of the constitutively phosphorylated

CD20. The serine residue S886 within the conserved RxxSxP motif of GEF-H1 is constitutively

phosphorylated by Pak2, a modification that is associated with the inhibition of the GDP/GTP

exchange activity[60]. Thus, two serine/threonine kinases, PKCδ and Pak2, control the formation

and silencing of the CD20/GEF-H1 complex.

The activity of GEF-H1 is regulated by the cytoskeleton[47]. Specifically, GEF-H1 is intimately

associated with the MT network and this association inhibits its GDP/GTP exchange activity.

There are two (non-mutually exclusive) mechanisms how MT binding inhibits GEF-H1. In the

first one, polymerized MT stabilize the autoinhibited conformation in which the N-terminal C1

domain is folded over the PH domain of GEF-H1[61]. The second mechanism involves the

binding of the light chain protein Tctex-1 of the dynein motor complex to the RRxxR binding

motif at the unstructured C-terminal tail of GEF-H1[62,63]. In this way the dimeric dynein motor

complex couples GEF-H1 to polymerized MT. The RxxSxP motif that connects GEF-H1 to the

phosphorylated CD20 tails lies 60 amino acids downstream of the Tctex-1 binding site. Thus,

in resting B cells the autoinhibited GDP/GTP exchange factor functions as an adaptor that

connects CD20 to the MT network. Based on the finding that anti-CD20 antibodies can

modulate the calcium response in B lymphocytes, it has previously been suggested that CD20

acts as a calcium channel[64,65]. This suggestion is not supported by the CD20 structure which

did not reveal a calcium permeable pore[7]. However, we noticed that the N-terminus of the

store-operated calcium channel ORAI 1 carries an RRxxR and an RxxS motif[66]. Thus, Tctex-

1 and the 14-3-3 adaptor could couple CD20 to the ORAI1 channel and this connection could

explain the observed effect of anti-CD20 antibodies on the calcium response in B lymphocytes.

CD20 can orchestrate the MT-to-actin switch during B cell activation

In migrating, polarized B lymphocytes, the MT organization center (MTOC) is found in the

vicinity of the trailing edge and the MT network is involved in the organization of the uropod at

this location[67]. We have found that in polarized B cells the IgD-BCR nanocluster is localized at

the uropod whereas at the leading-edge the IgM-BCR is associated with microvilli structures

that are formed by core bundles of actin filaments[68]. The interaction of CD20 to the MT network

may help to stabilize the IgD-BCR nanocluster[69]. Interestingly, apart from the CD20/GEF-H1

complex the IgD-BCR nanocluster harbors several receptors that carry in their cytosolic

juxtamembrane region a RRxxR (BAFF-R) or a related RRxxK motif (CD19) for Tctex-1

binding. Thus, multiple contact points to the MT network may promote the stability of the IgD-

BCR nanocluster[70].

In polarized lymphocytes RhoA and ROCK1 are also found at the uropod and a RhoA

deficiency or an inhibition of ROCK1 kinase activity prevents the formation of this structure[67,71].

RhoA undergoes geranylgeranylation at its C-terminal CAAX motif and thus is constitutively

bound via its lipid anchor to the inner leaflet of the plasma membrane[72]. In spite of its

permanent membrane association, RhoA in resting B cells is not found in the vicinity of CD20

and the IgD-BCR nanocluster as the PLA has shown. Only upon MT destabilization and the

following dissociation of the GEF-H1/CD20 complex RhoA and ROCK1 are found in the vicinity

of CD20. Interestingly, ROCK1 carries a RxxS motif close to its C-terminus and it is feasible

that the GEF-H1/CD20 complex in resting B cells is replaced by a RhoA-GTP/ROCK1/CD20

complex in activated B cells. The GEF-H1 activation and liberation from the MT network could

have a dual impact on ROCK1 activity and membrane location. The increased GDP/GTP

exchange activity of GEF-H1 generates RhoA-GTP that attaches to the RhoA binding domain

of ROCK1 and the GEF-H1-deprived CD20/14-3-3 complex is free to bind to ROCK1. Both

binding events counteract ROCK1 autoinhibition and localizes the active ROCK1 at the IgD-

BCR nanocluster where ROCK1 phosphorylates the myosin light chain (among other

substrates), resulting in the formation of myosin-II filaments and actomyosin stress fibers with

increased contractility[73]. The stress fiber formation in polarized, antigen-exposed activated B
cells[58,74] may be associated with uropod retraction, the dissociation of the IgD-BCR
nanocluster and the observed movement of CD20 and CD19 to the IgM-BCR. According to
this scenario CD20 is not only a gatekeeper of the resting state but also an actomyosin bundles
associated transporter that promotes the immunoreceptor coupling and organization motif
(ICOM)-dependent conjugation of the IgM-BCR with the CD19 coreceptors in activated B
lymphocytes[75].
396

The effect of anti-CD20 antibodies on cytoskeleton reorganization

Previously it was thought that binding of anti-CD20 antibodies did not have a direct effect on
the physiology of targeted B cells, but simply provided a flag for their destruction via
phagocytosis or ADCC mechanisms[76]. However, several studies indicated that anti-CD20
antibodies induce an apoptosis program in the targeted B cells[77]. We found that exposing B
cells to anti-CD20 antibodies results in the dissolution of MTs and MT instability, which is
associated with apoptosis [38,44,78]. Furthermore, caspase 3, an important effector of the
apoptosis program, can cleave and activate ROCK1[44]. Interestingly, it is the cleaved, active
ROCK1 fragment, which carries the C-terminal RxxS motif and therefore could still be
associated with CD20. In this way the active kinase domain of ROCK1 remains bound to the
plasma membrane in a RhoA-GTP independent manner. A recent 3-dimensional super-
resolution study has localized the RTX-engaged CD20 on actin bundles within microvilli
structures[79]. This is in line with our findings that the IgM-BCR is associated with an actin-based
network of ridges of microvilli structures and that CD20 is found in the vicinity of the IgM BCR
after exposure to either cognate antigen or anti-CD20 antibodies[68]. Together these data
support the assignment of CD20 as a switch factor, flexible to interact with the MT and
alternatively the actomyosin cytoskeleton in resting and activated B lymphocytes, respectively.
The finding that the exposure to anti-CD20 antibodies results in dissolution of MT bundles and
protofilaments and that MT-destabilizing drugs can mimic the effect of an anti-CD20 treatment
of B lymphocytes not only gives a new insight in the therapeutic effect of the anti-CD20 therapy
but also could have an impact on the improvement of such a therapy. It is known that anti-
CD20 treatment results in the reduction of the amount of CD20 on the B cell surface and this
could reduce the targeting efficiency[80,81]. The exposure of healthy donor B cells to MT
stabilizing drugs such a Taxol derivates could prevent the downregulation of antibody-bound
CD20 and thus increase the B cell targeting function of the anti-CD20 therapy. However, there
also exist data showing that the ADCC-dependent killing of anti-CD20 labelled B cells by NK
cells requires dynamic alterations of the MT network. If this is the case, a combination of anti-
CD20 therapy with MT-destabilizing drugs such a vincristine could enhance the NK-mediated
cytotoxicity of anti-CD20-targeted B cells[82]. While these potential modification of the anti-CD20
therapy requires more studies, the cellular function of CD20 as an adaptor between antigen

receptors and the cytoskeleton is now established and CD20 is no longer an enigma of B cell
biology.
429

430   **Materials & Methods**
431

Human naïve B cells
Primary naïve B cells were obtained from fresh buffy coats, provided by the Institute for
Transfusion Medicine and Gene Therapy, ITG, Freiburg. Peripheral blood mononuclear cells
(PBMCs) were separated by Ficoll gradient centrifugation of whole blood and negatively
selected using EasySep Human Naïve B Cell Isolation Kit (Stemcell). Prior to experiments,
primary B cells were controlled for purity and rested overnight. The study was approved by the
Institutional Review Board of the University Freiburg (Ethical votes No 507/16 and 336/16).
439

Cell culture
The human Burkitt lymphoma B cell line Ramos was obtained from American Type Culture
Collection (ATCC, Ramos cat. CRL-1923, RRID: CVCL 1646) and stably transfected with
ecotropic receptor (EcoR) to allow murine retroviral transfection with MMLV particles. Ramos
B cells were cultured in RPMI 1640-Glutamaxx medium, 10% FCS (PAN, Biotech), 10 units/mL
penicillin/streptomycin (Gibco), 20 mM hepes (Gibco), and 50 mM β-mercapto-ethanol (Sigma)
in a humidified saturated atmosphere at 37 °C with 5% CO_2.
For the quantitative phosphoproteomic study, Ramos B cells were labelled using stable isotype
labelling by amino acids in cell culture (SILAC). Light labelling was performed using 12C6L-
arginine/12C6-L-lysine. For medium-heavy and heavy labelling 13C6L-arginine/D4-L-lysine
and 13C615N4-L-arginine/13C615N2-L-lysine were used, respectively.
451

Generation of the Ramos CD32b KO and CD20 KO cell lines
The human CD32b and CD20 gene were render defective in the Ramos B cell system by the
CRISPR/Cas9 technique using the SgRNAs agcacgttgatccactgggg for CD32b KO (ex4) and
gccctattgctatgcaatctgg for the CD20 KO (ex3), respectively. The Neon Transfection System
(Invitrogen) was used to introduced a ribonucleoprotein (RNP) complex with the pre-
assembled Cas9 protein and guide RNA into the Ramos B cells according to the genome
editing method of Integrated DNA Technology (IDT). In brief: Ramos cell medium was changed
the day before transfection. 1×10^6 Ramos cells were pelleted by centrifugation and
resuspended in 9 µL buffer R (Neon), 1µL of RNP complex, and 2µL of electroporation
enhancer (IDT). RNP complex contained diluted sgRNA (annealed crRNA and tracrRNA in a
1:1 ratio in IDTE buffer) and Cas9 endonuclease in buffer R (NEON). Electroporation was
performed in 10 µL NEON tips at 1350 V, 30 ms, single pulse. The transfected cells were first
recovered for 72hs at 37°C and 5% CO2 without antibiotics and then subjected to complete
RPMI medium. Successful transfected Ramos cells were batch sorted using BioRad cell sorter

in three different rounds and monitored throughout experiments. Deletion of the target gene
was verified by flow cytometry.
468
Generation of the Ramos PKCδ (PRKCD) gene KO
The sequence AACCCAATCATAGCAGAGC or GCTGAGTTCAGTGAGTGC encoding for the
specific gRNA was inserted into a lentivirus CRISPR vector (Addgene). The constructs were
co-transfected with an mCherry expression vector into Ramos cells. The cells were incubated
at 37°C for 24 hours in a humidified CO2 incubator. The following day, single cells were sorted
for mCherry reporter gene expression with the BD FACSAria™ III Cell Sorter. The sorted single
cells were cultured for 3-4 weeks in complete RPMI medium supplied with 10% FBS in a
humidified CO_2 incubator at 37°C. The expected PRKCD gene deficiency was verified by
Sanger sequencing and the loss of PKCδ protein expression by Western blotting.
478
Lentiviral transfection for the generation of CD32b-tr Ramos B cells
The human embryonic kidney HEK 293T cells were cultured in complete DMEM GlutaMAX
medium supplemented with 10% FCS, 10 mM HEPES, 10 µM sodium pyruvate, 50 units/mL
penicillin and 50 µg/mL streptomycin in a humidified saturated atmosphere at 37 °C with 7.5%
CO_2. Lentivirus particles were obtained by co-transfecting HEK 293T cells with pLenti-
hCD32b-IRES-GFP, pCMVΔR8.74 and pMD2vsvG plasmids using Polyethyleneimine (PEI,
Polysciences). Viral supernatant was harvested, sterile filtered and combined 24h and 48h
post-transfection. Lentiviral particles were enriched by centrifugation (4 hs, 10.000 g, 8 °C)
after placing them in a 1:5 ratio on a 10% sucrose layer. The lentiviral pellet was resuspended
in DMEM Glutamaxx without supplements and stored at –80 °C. The viral titers were assessed
by determining the multiplicity of infection (MOI). Briefly, $5x10^4$ HEK 293T cells per well were
seeded in a p24 well plate in 1 mL medium. An aliquot of the concentrated virus was diluted
1:100 in medium. Various volumes (0, 1, 5, 10, 25, and 50 µL) of lentivirus dilution were added
to the cells. After 48h of incubation the GFP expression was analyzed by flow cytometry and
lentivirus titer was calculated: Transduction units/mL = (number of cells x % of GFP⁺ cells x
dilution factor) / (mL of lentivirus dilution).
495
Construction of WT or S/A mutated CD20-flag expression vectors
The CD20-flag vector was modified by geneblocks (IDT) encoding the N-terminal or C-terminal
tails with S/A exchanges at the chosen 4 positions, respectively. Geneblocks were solved in
IDTE buffer according manufacturer's protocol and subsequently inserted in the pMIG CD20-
flag-IRES-GFP vector using infusion cloning method. Plasmid were sequenced (GATC,
eurofins) and used to generate the CD20WT, CD20Nmut, CD20Cmut and CD20NCmut
Ramos cell lines.
503
Retroviral transfection of CD20 mutants:

Murine retrovirus-containing supernatants were obtained by transfecting Phoenix-eco cells
with LipoJet (SignaGen) according to manufacturer's protocol and different pMIG-CD20-IRES-
GFP constructs, or empty GFP-control plasmid. After 48h viral supernatant was collected,
sterile filtered and used directly or stored in -80°C. For transduction 4×10^5 newly generated
CD20 KO Ramos cells were resuspended in 1 mL of viral supernatant containing Polybrene
(1 µg/mL) and spin infected by centrifugation (180 min, 300g, 37°C). The viral supernatant was
removed and cells were cultured and sorted for GFP expression (BioRad cell sorter).
512

RNA sequencing
Wildtype Ramos or PRKCD Knockout Ramos were cultured with or without IgM (10ug/ml)
stimulation for 4 hours and then RNAs were extracted by RNeasy plus kit (Qiagen cat# 74143).
RNA libraries for RNAseq were prepared using the NEBNext® Ultra™ II Directional RNA
Library Prep Kit for Illumina (NEB cat # E7760L) according to manufacturer's instructions and
samples were sequenced 84-bases long on the NextSeq® 550 with high output flowcell
(NextSeq® 500/550 High Output Kit v2).
520

GEO data
Downstream analyses were conducted using R (version 4.4.0). Raw read counts were
normalized for library size and composition using the trimmed mean of M-values (TMM)
method. Differential gene expression analysis comparing the unstimulated Ramos and PKCδ
KO cell conditions was performed with the edgeR package (version 4.2.2)[83]. Gene set
enrichment analysis was carried out using the clusterProfiler package (version 4.12.6)[84]with
gene sets from the Molecular Signatures Database (MSigDB, version 7)[85]. For all statistical
analyses, significance was defined as an adjusted p-value < 0.05.
529

Flow cytometry analysis and phospho-flow detection
For surface staining, $1–20 \times 10^5$ cells were Fc-blocked (Human TruStain, Biolegend), stained
with fluorescent antibodies on ice, washed twice, and then their fluorescence measured with
a FACS Attune NxT (Life technologies). For intracellular staining and phospho-flow analysis,
the cells were activated for the requested time points, immediately fixed with 4% PFA, washed
twice and permeabilized with 0.5% saponin (Quillaja bark, Sigma) containing protease and
phosphatase inhibitors (HALT, Thermo Fisher) for 30 min. Samples were washed and stained
overnight at 4°C. If necessary secondary antibody staining was performed, the background
was controlled using only the secondary antibody. Data were exported in FCS-3.0 format and
analyzed with FlowJo software (TreeStar).
540

Western Blotting
Unstimulated or activated Ramos B cells were collected and immediately lysed in 2x Laemmli
or RIPA buffer for 30 min on ice. Equal amounts of adjusted and cleared lysates (Qubit 3.0;

life technologies) were subjected to SDS–PAGE on 10% mini precast gels (7bioscience). After

transfer, the PVDF membrane (GE Healthcare Amersham) was blocked with 5% BSA in PBS

and 0.1% Tween20. The membrane was first exposed to primary antibodies specific for the

proteins under study (CD20, 14-3-3 or GEF-H1) and then washed. A horseradish (HRP)

peroxidase-conjugated goat anti-rabbit or goat anti-mouse antibody was used as secondary

antibody before detection with ECL chemiluminescent substrate (Bio-Rad).

Affinity purification of CD20-flag and GEF-H1

The anti-flag immunoprecipitation was performed according to the manufacturer's protocol

(chromotek). In brief, 10^7 Ramos cells were lysed for 30 min in ice-cold lysis buffer

supplemented with a protease/phosphatase inhibitor cocktail (HALT, Thermo Fisher). Lysates

were centrifuged for 10 min at 17000 g and diluted in 300µL dilution buffer. 50 µL of total lysate

were kept for the input analysis. Diluted lysates were rotated overnight with anti-flag agarose

beads. Beads were then sedimented and washed extensively. Bound proteins were eluted with

2x SDS-PAGE Laemmli buffer and samples were boiled at 95°C for 5 min and subjected to

SDS-PAGE and Western blotting.

RhoA(G17A), the nucleotide-binding deficient mutant of RhoA binds strongly to GEF-H1 and

can be used for GEF-H1 purification and quantification. For this a glutathione S-transferase

(GST) coupled RhoA(G17A) fusion protein was produced in E. coli bacteria and the GST-

RhoA(G17A) loaded on Glutathione Sepharose beads. The GST-RhoA(G17A)-beads were

mixed with Ramos B cell lysate and rotated overnight at 4°C. Beads were washed extensively

with lysis buffer, boiled for 5 min and the cleared supernatant subjected to SDS-PAGE and

western blotting.

Proximity ligation assay (PLA)

The Fab-PLA was performed as described earlier[28]. In brief: for Fab-PLA, the F(ab)- fragments

were prepared from the corresponding antibodies using the Pierce Fab Micro Preparation Kit

(Thermo Fisher). The produced F(ab)-fragments or primary antibodies (1-PLA) coupled with

PLA probemaker (Sigma-Aldrich) after buffer exchange (Zeba™ spin desalting columns,

Thermo Fisher). Ramos or HD naïve B cells were settled on polytetrafluoroethylene (PTFE)

slides (Thermo Fisher Scientific) for 30 min at 37°C and kept unstimulated or activated with

RTX [10µg / mL] for 5 min and then fixed for 20 min with 4% paraformaldehyde.

After extensive washings the slides were blocked for 30 min with blocking buffer (25 µg/mL

sonicated salmon sperm DNA, 250 µg/mL BSA, 1 M glycine). PLA was performed with the

Duolink In Situ Orange (Sigma-Aldrich). Slides were directly mounted with DAPI Fluoromount-

G (Southern Biotech) to visualize the PLA signals in relation to the nuclei. For each experiment,

a minimum of 1000 HD B cells or Ramos cells were analyzed with CellProfiler-3.0.0. In brief:

metadata of images were exported and the intensity of the input frames were rescaled to the

full intensity range in the RGB scheme. PLA signals (size of PLAdots 4-10 pixels) in a distance

of 10 pixels to the Nuclei (size 15-80 pixel) with a threshold of 0.0075 of at least 1000 cells
were measured and subjected to analysis.
585

586   **Imaging, Image analysis, and Data processing**
587   All microscope images for PLA were acquired using Leica DMi8 microscope equipped with an
588   HC PL APO 63×/1.40-0.60 oil immersion objective lens and analyzed with CellProfiler 3.0.0.
589   and Prism software (GraphPad, La Jolla, CA).
590

Microtubule staining
MT were stained as described[86]. In brief: Ramos B cells were settled on uncoated glass bottom
chambers (IBIDI) for 30 min at 37°C, kept unstimulated or activated with RTX for 20 min and
then fixed for 20 min with 4% paraformaldehyde and 0.1% glutaraldehyde. After blocking and
permeabilization with 250 µg/mL BSA, 1 M glycine 0.5% saponine for 30 min cells were stained
with anti-α tubulin (Sigma Aldrich) over night and with secondary anti mouse Alexa fluor-555
antibody (Invitrogen). Fixed Ramos cells in IBIDI chambers were kept in PBS until microscope
imaging.
HD B cells were settled on PTFE slides (Thermo Fisher) for 30 min at 37°C, kept unstimulated,
or activated with RTX [10µg/mL] for 5 min, with or without pretreatment with Taxfor 1h and
then fixed for 20 min with 4% paraformaldehyde and 0.1% glutaraldehyde. After blocking and
permeabilization with 250 µg/mL BSA, 1 M glycine 0.5% saponine for 30 min cells were stained
with anti-α tubulin (Sigma Aldrich) over night and with secondary anti mouse Alexa fluor-555
antibody (Invitrogen). Slides were directly mounted with Fluoromount-G mounting-solution
(Invitrogen) and subjected to microscope analysis.
606

Microscopy
The high-resolution visualization of MTs was performed using a LSM 880 Airyscan laser
scanning confocal microscope attached to an inverted microscope Axio Observer Z1 (Carl
Zeiss Microscopy). The Alexa Fluor 555 labelled α-tubulin was excited with 561nm, the
emission was detected with the Airyscan detector in SuperResolution (SR) mode. For all cells
z-stacks were acquired with a Plan-Apochromat 63x/1.40 Oil DIC M27 (Fig. S7) or a LD LCI
Plan-Apochromat 40x/1.20 autocorr (Fig. S6) objective. The immersion medium was oil (63x)
or water (40x).
Zeiss Software ZEN Black 2.3 SP1 FP3 (release 14.0.22.201) was used for acquisition.
(acquisition settings for Fig. S7: EM filter SP615, pixel size xy:0.049µm, z-step size: 0.199µm,
pixel dwell time 1µs, bi-directional scanning, for Fig. S6: no EM filter, pixel size xy:0.049µm, z-
step size: 0.243µm or 1µm, pixel dwell time 1,92µs, bi-directional scanning). All images were
processed with the above ZEN Black software using the Airyscan algorithm and with Zen Blue
(Version 3.9.3) for subset and maximum intensity projection. Airyscan processed images
shown-in Fig. S6 are single z-planes from an image z-stack and in Fig. S7 maximum intensity

projections of the full image z-stack. The display gain is set to 0.75 and the intensity value to

Best fit.

Statistical analysis

Students t-test was used for the experiments if not otherwise described to determine statistical

significance provided by Prism10 software (GraphPad, La Jolla, CA).

MT network stabilization or destabilization

For the stabilization of MT HD naïve B cells or Ramos B cells were treated for 1 h with 100 nM

docetaxel trihydrate (MedChemExpress) solved in DMSO and diluted in RPMI culture medium.

For the destabilization of MT cells were treated for 1 h with 10 µM nocodazole (Selleck-

Chemicals) solved in DMSO and diluted in RPMI culture medium.

Rituximab (RTX) treatment

RTX was kindly provided by F. Hoffmann-La Roche-AG. For naïve HD B cells or Ramos cell

treatment RTX was diluted to a final concentration of 10 µg/mL and experiments were

performed at least three times.

Phosphopeptide enrichment

SILAC-labeled Ramos B-cells (wild-type and knockout) were starved by serum depletion (FCS)

for 30 min. Following starvation, cells were harvested by centrifugation at 1,200 rpm for 5 min,

washed twice with starvation medium. As a control, mock treatment was performed using PBS.

After activation, cells were rapidly snap-frozen in liquid nitrogen to preserve their signaling

status. Cell lysis was performed using GdmHCl buffer (6 M GdmHCl, 100 mM Tris-HCl pH 8.5,

10 mM TCEP, and 40 mM chloroacetamide). The cell pellets were resuspended in the lysis

buffer, followed by sonication (two cycles of 30 seconds). Protein denaturation was achieved

by incubating the lysates at 95°C for 5 min. To remove cellular debris, the lysates were

centrifuged at 3,500 x g for 30 min at 4°C. Protein concentration was determined using a

Bradford assay. To precipitate proteins, four volumes of ice-cold acetone were added to the

lysate. The protein precipitates were collected by centrifugation and washed with ice-cold

acetone to remove residual contaminants. Proteins were digested using a combined Lys-C /

trypsin digestion. Phosphopeptides were enriched using the Easyphos workflow. The

phosphopeptide-enriched eluates were desalted using C18 StageTips (3M Company), pooled,

and stored at -80°C until further analysis.

LC-MS/MS analysis

Peptide mixtures, reconstituted in 0.1% TFA, were analyzed by nano HPLC-ESI-MS/MS using

an UltiMate 3000 RSLCnano HPLC system (Thermo Fisher Scientific, Dreieich, Germany)

online coupled to an QExactive Plus mass spectrometer (Thermo Fisher Scientific, Bremen,

Germany). The RSLC system was equipped with C18 trap columns (nanoEase M/Z Symmetry
C18 Trap; 20 mm length, 180 µm inner diameter, 5 µm particle size, 100 Å pore size, Waters
Corporation, Milford, MA) and an analytical C18 reversed-phase nano LC column (nanoEase
M/Z HSS C18 T3; 250 mm length, 75 mm inner diameter, 1.8 µm particle size, 100 Å pore
size, Waters Corporation, Milford, MA). A binary solvent system consisting of 0.1% (v/v) formic
acid (FA) as solvent A and 80% (v/v) acetonitrile (ACN)/0.1% (v/v) FA as solvent B was
employed for peptide separation. Peptide mixtures were loaded, washed and preconcentrated
on the pre-column for 5 min using solvent A and a flow rate of 10 µL/min. A gradient was then
applied at a flow rate of 300 nL/min ranging from 4 to 39% B in 140 min, 39–54% B in 15 min,
54%-95% in 5 min and 3 min at 95% B. Eluted peptides were transferred to a fused silica
emitter for electrospray ionization enabled by a Nanospray Flex ion source with DirectJunction
adaptor (Thermo Fisher Scientific), applying a spray voltage of 1.6 kV and a capillary
temperature of 250°C. MS/MS data were acquired in data-dependent mode using the following
parameters: MS precursor scans at m/z 370–1700 with a resolution of 70,000 (at m/z 400);
automatic gain control (AGC) of 3×10^6 ions; a maximum injection time (IT) of 60 ms; a top 12
method for higher-energy collisional dissociation of multiply charged precursor ions with a
normalized collision energy of 28%. MS/MS scans from 200 to 2000 m/z were recorded at a
resolution of 35,000. The AGC for MS/MS scans was set to 1×10^5 with a maximum IT of 120
ms and a dynamic exclusion time of 45 s. Data are available via ProteomeXchange with
identifier PXD063667.
681
MS data analysis
MaxQuant (version 2.6.7.0) with its integrated Andromeda search engine[87,88] was employed.
MS/MS data were searched against the human proteome set with isoforms downloaded from
UniProt (105300 entries). Protein identification was performed using MaxQuant default
settings (including carbamidomethylation of cysteine residues as fixed modification and N-
terminal acetylation and methionine oxidation as variable modifications), with the following
exceptions: 'Arg6;Lys4' and 'Arg10;Lys8' were specified as medium and heavy modification
labels, respectively; a maximum of three missed cleavages was allowed; 'Phospho (STY)',
'Oxidation (M)' and 'Acetyl (Protein N-term)' were selected as variable modifications; and the
options 'match between runs' and 'requantify' were activated.
Ratios between labels (normalized by MaxQuant) were extracted from the
Phospho(STY)Sites.txt file, reverse entries and potential contaminants were removed, and the
list was filtered for at least two unique peptides per phosphosite. Phosphosite ratios were
normalized to the respective ratio at the protein group level using the median ratio of all protein
group ratios referenced in the 'Protein group IDs' column of the Phospho(STY)Sites.txt file.
Resulting phospho site ratios were filtered so that only sites with valid values in >=3 of 4
replicates were retained. Significance of differential regulation was tested using linear models
implemented in the LIMMA package and the resulting fold changes and significance level were

plotted. Data analysis was performed in Python 3.10 using the autoprot wrapper [89] for data analysis and plotting. All original code for MS data analysis has been deposited at Zenodo (https://doi.org/10.5281/zenodo.15350913).

Acknowledgments

We thank Dr. Christian Klein and the Roche Innovation Center Zurich for financial support and anti-CD20 antibodies and Falk Nimmerjahn, University of Erlangen for the CD32B plasmid. The generation of the CRISPR-Cas9 PKCδ plasmid and RNA sequencing was made possible by the help of Dr. Claudio Mussolino and Dr. Baerbel Keller. We thank them for their assistance. We thank Dr. Lise Leclercq for correcting the manuscript. We also thank the Lighthouse Core Facility, which is funded in part by the Medical Faculty, the University of Freiburg (Project Numbers 2023/A2-Fol; 2021/B3-Fol), the DKTK, and the DFG (Project Number 450392965).

Funding: This work was supported by the Roche research contract ZVK-2023101202 and an RO1 grant of the National Institutes of Health (NIH) under the award number A031503 (SMA, MR). The Zeiss LSM 880 microscope was funded by the Deutsche Forschungsgemeinschaft (DFG), Project No. 31778431, Work in the Warscheid lab was supported by the DFG SPP2453 (project number 541758684), FOR2743 (project 9) and TRR130 (project C01).

Disclosure and competing interest statement

The authors declare no conflict of interests.

Supplements

Main figures 1-7

Supplemental figures S1-S7

Structured Methods - Reagents and Tools Table

Extended table 1 Geneblocks

Graphical abstract

Scheme of the resting B cell IgD-BCR cluster connected to the microtubule network

The activated B cell IgM-BCR connected by Rock1 to the actin network

Figure 1

A

B

Figure 1:

Phenotype of PKCδ KO Ramos B cells and identification of PKCδ substrate proteins.

Comparison of the expression of B cell marker protein on the surface of PKCδ KO (brown) and WT Ramos B cells (blue) by flow cytometry. The expression data of PKCδ KO are normalized to that of WT Ramos cells set to 100%. **B**. Volcano plot showing quantitative changes of the constitutive serine/threonine phosphorylation of substrate proteins of WT in comparison to PKCδ KO Ramos B cells. On the X-axis the plot depicts log2 fold change of phosphorylated serine or threonine residues in the whole proteome of the two cell lines. The Y-axis shows the log10 FDR adjusted p-value <0.05. The significant changes of phosphorylated CD20 serine residues are indicated and shown in orange.

Figure 2

E

Fab-PLA IgD:CD32b

F

Figure 2:

Functional association of the Fc-receptor CD32b with the IgD-BCR nanocluster.

A, B. Flow cytometry analysis of CD32b expression on the surface of CD32b KO (beige) or

CD32b-tr (orange) in comparison to WT Ramos B cells (blue). The unstained control (US) is

shown in light grey. **C, D.** Flow cytometry analysis of the expression of B cell marker protein

on the surface of CD32b KO (beige) or CD32b-tr (orange) in comparison to WT Ramos B cells

(blue). The expression data of CD32b KO or CD32b-tr cells are normalized to those of WT

Ramos set to 100%. **E.** Fab-PLA study of the IgD-BCR:CD32b proximity on the surface of WT

Ramos (upper left), CD32b-tr, (upper right), CD32b KO (lower left), and isolated healthy donor

(HD) B cells (lower right). PLA dots in red and the nuclei stained blue by DAPI, Scale bar 5

µm, **F.** Quantification of Fab-PLA data with the CellProfiler and calculation of the significance

between the groups with PRISM 10, one-way ANOVA, the Students t-test shows no significant

difference between CD32-tr and HD B cells. Mean value are show as red bars.

Figure 3

A

B

C

804

805

D

1-PLA CD20:14-3-3

WT   Nmut

Cmut   NCmut

E

1-PLA CD20:14-3-3

F

anti flag   total lysate

14-3-3
29kD

CD20
30kD

806

807

808   **Figure 3:**

809   **Mutational analysis of CD20 serine residues and their association with 14-3-3**

810   **A.** Protein sequence of the N-terminal (top) and part of the C-terminal cytoplasmic tail of human

811   CD20 (bottom). Below the WT sequence the sequence with the introduced S/A mutations (in

812   bold letters) at the N-terminal (Nmut) and C-terminal (Cmut) cytoplasmic tail of CD20 are

813   shown. The location of the RxxS motifs implicated in 14-3-3 binding is shown below the

814   sequence **B.** Membrane topology of human CD20 with a C-terminal flag-tag (CD20-flag). The

815   location of the two RxxS motifs is indicated with red boxes **C**. Flow cytometry analysis of

816   CD32b-tr Ramos cells (orange) and reconstituted Ramos cells expressing CD20 WT (black),

817   Nmut (green), Cmut (dark blue) or NCmut (red). The Ramos cells are tested for the expression

818   of CD20, CD32b, IgM, IgD, and CD83. The unstained (US) CD32-tr cells are shown in grey**.**

D. Fab-PLA study of the CD20:14-3-3 proximity in WT (upper left), Nmut (upper right), Cmut (lower left), and NCmut (lower right) Ramos B cells. PLA dots in red and the nuclei stained blue by DAPI, Scale bar 5 μm. **E**. Quantification of the PLA data by CellProfiler. Significant differences between groups were calculated with PRISM 10, one-way ANOVA. Students t-test shows no significant difference between CD32-tr and WT Ramos cells. **F.** Western blot analysis of the CD20-flag/14-3-3 association in WT, Nmut, Cmut and NCmut Ramos B cells. The amount of 14-3-3 protein in the anti-flag immunoprecipitated and total lysates is show on the left and right, respectively. Lysates were taken from at least 3 independently generated Ramos cell lines.

Figure 4

A

B

1-PLA CD20:GEF-H1

WT Nmut
Cmut NCmut

C

1-PLA CD20:GEF-H1

D

E

F

Figure 4:

Constitutive association of CD20 with the GDP/GTP exchange factor GEF-H1.

A. Schematic drawing of the domain structure of the RhoA guanine nucleotide exchange factor

GEF-H1 (ARHGEF2). The functional domains are drawn as blue boxes. The location of the

RRxxR motif implicated in dynein light chain binding, and the Pak2 phosphorylated S886 within

the 14-3-3 binding **R**xx**S**x**P** motif is indicated. **B**. 1-PLA study of the CD20: GEF-H1 proximity

in WT (upper left), Nmut (upper right), Cmut (lower left), and NCmut (lower right) Ramos B

cells. PLA dots in red and the nuclei stained blue by DAPI, Scale bar 5 µm. **C**. Quantification

of PLA of B. with CellProfiler. Significant differences between groups were calculated with

PRISM 10, one-way ANOVA. Students t-test shows no significant difference between CD32-

tr and WT Ramos cells. **D.** Western blot analysis of the constitutive CD20-flag/GEF-H1

association in WT, Nmut, Cmut and NCmut Ramos B cells. The amount of GEF-H1 protein in

the anti-flag immunoprecipitated and total lysates is show on the left and right, respectively. **E.**

Western blot analysis of free GEF-H1 in the lysates of Ramos and CD32b-tr cells as well as

reconstituted Ramos cells expressing CD20 WT, Nmut, Cmut or NCmut. The amount of GEF-

H1 purified with the GST-RhoAG17A beads and in the total lysate is shown at the top and

bottom, respectively. **F.** Quantification of the Western blot data with Image Studio Lite (Licor)

showing the difference of signal intensity in the immunoprecipitate (IP) versus that of the total

lysates (TL).

Figure 5

917
918

919 **Figure 5:**

920 **Loss of GEF-H1 and gain of the RhoA/Rock1 association of CD20 upon RTX treatment**

1-PLA study of **A.** the CD20:14-3-3, **C.** the CD20:GEF-H1, **E.** the CD20:RhoA and **F.** the CD20:ROCK1 proximity in CD32b-tr Ramos cells (left) and reconstituted Ramos cells expressing WT (middle) or NCmut CD20 (right). The studied Ramos cells were either untreated (top) or exposed for 5 min to RTX (bottom). PLA dots in red and the nuclei stained blue by DAPI, Scale bar 5 µm. **B**, **D, G, H.** Quantification of the 1-PLA data with CellProfiler. Significant differences between two groups were calculated with PRISM 10 Student's t-test.

Figure 6

Figure 6:

Altered CD20 association in RTX or nocodazole treated HD B cells

1-PLA study of **A.** the CD20:14-3-3, **C.** the CD20:GEF-H1, **E.** the CD20:RhoA and **F.** the

CD20:ROCK1 proximity in unstimulated (left) and HD B cells exposed for 5 min to RTX (middle)

or treated for 1 h with Noc (right). PLA dots in red and the nuclei stained blue by DAPI, Scale

bar 5 μm. **B**, **D, G, H.** Quantification of the 1-PLA data with CellProfiler. Significant differences

between two groups were calculated with PRISM 10 Student's t-test.

Figure 7

A

CD20 surface abundance

B

Rock1 pY914

C   **Fab-PLA IgD:CD19**

D

E   **Fab-PLA IgM:CD19**

F

Figure 7:

Stabilization of the IgD-BCR nanocluster and CD20 by the MT network

A. CD20 expression on the surface of untreated (UT) HD B cells or B cells stimulated for 5 min

with RTX. As indicated, some of these HD B cells were pretreated for 1 h with either

nocodazole (Noc) or docetaxel (Tax). Shown is a summary of the MFI data of >3 flow cytometry

experiments. PRISM10 and the students-tests were used to calculated the significant

differences between the experimental groups. **B**. Y914 phosphorylation and activation of Rock1 in untreated (UT) HD B cells or B cells stimulated for 5 min with RTX. As indicated, HD B cells were kept untreated, or pretreated for 1 h with nocodazole (Noc), or docetaxel (Tax). A summary of the MFI data of >3 flow cytometry experiments is presented. Students-tests (Prism10) were used to calculated the significant differences between the groups. Fab-PLA study of **C.** the IgD-BCR:CD19 or **E.** the IgM-BCR:CD19 proximity of HD B cells, kept untreated or stimulated for 5 min with RTX without or with pretreatment of the B cells with either Noc or Tax for 1h as indicated. PLA dots in red and the nuclei stained blue by DAPI, Scale bar 5 µm. **D, F.** Quantification of the Fab-PLA data with CellProfiler. Significant differences between groups were calculated with PRISM 10 Student's t-test.

Figure S1

A

clone 6'-1 6'-5 6'-8 6'-9 6'-11 6'-15 6'-17 6'-22

PKCδ
80kD

β-actin
42kD

B

clone 6'-1 6'-5 6'-8 6'-9 6'-11 6'-15 6'-17 6'-22

PKCδ pT505
80kD

β-actin
42kD

established *PRKCD* KO B cell-line

C

Volcano plot:
- PLEK: S117
- PLEC: S4155/S4165
- KLC1: S600
- MYO1E: S980
- MAP4: S375
- DYNC1LI1: T513
- MAP4: S1970
- STMN1: S16/S25
- STMN1: S63
- RFTN1: S199
- DYNC1LI1: S510
- MAP1S: T612/S614
- PAK2: S2
- ARHGEF2: S175/S178
- ARHGEF2: S154/S164
- MARK3: S390
- MAP1S: S629/S631
- MARK3: S461/S464
- ARHGEF2: S933
- ARHGEF2: S152
- PAK2: S141

Y-axis: -Log₁₀ Adj. P Value (0.0, 0.2, 0.4, 0.6, 0.8, 1.0, 1.2)
X-axis: Fold-Change WT vs. PKCδ KO (-2, -1, 0, 1)

1014

1015

Supplementary figure 1:

Western blot analysis of PKCδ after CRISPR-Cas9 mediated PKCδ KO. Lysates were stained

for PKCδ **A**. or PKCδ phospho-threonine 505 (PKCδ pT505) **B**. Clones 6'-9 and 6'-22 were

selected for further PKCδ KO studies. **C**. Volcano blot indicating the changes of the constitutive

serine or threonine protein phosphosites of PKCδ KO compared to Ramos cells. On the X-axis

the plot depicts log2 fold change of phosphorylated serine or threonine residues in the whole

proteome of the two cells lines. The Y-axis shows the log10 FDR adjusted p-value. Decreased

phosphorylation of serine residues of the MT pathway are shown in dark green, increase in

light green. The phosphorylated serine residues of MT proteins are indicated.

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

Figure S2

A

GOBP_REGULATION_OF_MICROTUBULE_DEPOLYMERIZATION_Mut_Rest-WT_Rest_plot

B

Supplementary figure 2:

Gene set enrichment analysis (GSEA) of genes associated with destabilized MT and cytoskeleton. **A.** The plot shows enrichment of genes associated with positive MT disassembly in PKCδ KO compared to Ramos WT cells. Vertical black lines indicate where genes overexpressed by MT depolymerization fall in the ranked list, (p-value 0.011 and FDR 0.81). **B.** As indication for the disconnection from the actin-cytoskeleton to the plasma membrane the plot shows the decrease of genes essential for the localization of the portion of the cytoskeleton that lies just beneath to the plasma membrane in PKCδ KO compared to Ramos WT cells, (p-value 0.00018 and FDR 0.091). The Normalized Enrichment Score (blue line) considers the ranked list of expression differences between two independent unstimulated PKCδ KO Ramos cell clones and Ramos WT cells.

Figure S3

CD32b CD20 IgD CD19 IgM

US
CD32b-tr
HD

Supplementary figure 3:
Due to the expression of the FcγRIIb receptor on Ramos, the surface abundance of critical B cell proteins more closely resembles that of naive B cells.

Flow cytometry analysis shows the comparison of the expression of B cell surface protein abundance on healthy donor (HD) negative selected naïve B cells (green), Ramos (black) and FcγRIIb receptor transfected (CD32b-tr, orange) Ramos cells. Unstained (US) control in grey).

Figure S4

A

B

1096
1097

Supplementary figure 4:

Western blot analysis of free GEF-H1 in the lysate of Ramos and CD32b-tr cells as well as
reconstituted Ramos cells expressing CD20 WT, Nmut, Cmut or NCmut after activation with
RTX for 5 min. The amount of GEF-H1 purified with the GST-RhoAG17A beads and in the
total lysate is shown at the top and bottom, respectively. **F.** Quantification of the Western blot
data with Image Studio Lite (Licor) showing the difference of signal intensity in the
immunoprecipitate (IP) versus that of the total lysates (TL).

Figure S5

1124
1125

Supplementary figure 5:

Intracellular phospho-flow cytometry analysis of intracellular signal proteins upon RTX

treatment for 5 min compared to unstimulated CD20 WT and NCmut S>A transfectant show:

A Loss of inhibitory GEF-H1 phosphorylation Serine[886], which is a prerequisite for the activation

of RhoA. **B** Loss of inhibitory RhoA Serine[188] phosphorylation, which renders RhoA active.

RhoA-GTP allows the increase of **C** active Rock1, phosphorylated at Y[914].

Students t-tests calculated with PRISM 10, show significant differences between the resting

and stimulated state of CD32b-tr and WT cells, but not between S>A mutants. Representative

NCmut CD20 mutant is presented. **D- F** Intracellular phospho-flow of the same protein residues

as in **A-C** but on HD naive B cells and as a time course after RTX stimulation for 0, 1, 5, 20

min. **D** inhibitory GEF-H1 S[886] phosphorylation, **E** inhibitory RhoA S[188], and **F** activatory Rock1

Y914 phosphorylation. Significant differences between the groups were calculated with PRISM

10, one-way ANOVA. Activatory phosphorylations are depicted in red, inhibitory are shown in

blue.

1140
1141
1142

Figure S6

A

Unstimulated

B

RTX for 20 min

Supplementary figure 6:

Airyscan confocal microscopy images of Ramos CD32b-tr cells **A.** unstimulated or **B.** incubated with RTX for 20 min. Cells were fixed and stained for α-tubulin with Alexa Fluor 555 to visualize MTs. Images present the mid-section of an Airyscan processed z-stack. White arrows in **B.** indicate the peeling of the curved protofilaments away from MT in the treated cells. Scale bar: 5µm.

Figure S7

A
Unstimulated

B
RTX for 5 min

C
Tax, 1h, RTX for 5 min

Supplementary figure 7:

Maximum intensity projection of Airyscan (SR) confocal z-stacks of HD naïve B cells **A.** untreated, **B.** upon exposure to RTX for 5 min or **C**. with Taxol pretreatment for 1h before exposure to RTX for 5 min. Cells were fixed and stained for α-tubulin with Alexa Fluor 555. In **B.** background is brightened due to higher presence of dead or destroyed cells. Scale bar: 5µm.

References:

1. Getahun A. Role of inhibitory signaling in peripheral B cell tolerance*. *Immunol Rev*. 2022;307(1):27-42. doi:10.1111/imr.13070

2. Kläsener K, Maity PC, Hobeika E, Yang J, Reth M. B cell activation involves nanoscale receptor reorganizations and inside-out signaling by Syk. *Elife*. 2014;3:e02069. doi:10.7554/eLife.02069

3. Horikawa K, Nishizumi H, Umemori H, Aizawa S, Takatsu K, Yamamoto T. *Distinctive Roles of Fyn and Lyn in IgD-and IgM-Mediated Signaling*. Vol 11.; 1999.

4. Becker M, Hobeika E, Jumaa H, Reth M, Maity PC. CXCR4 signaling and function require the expression of the IgD-class B-cell antigen receptor. *Proceedings of the National Academy of Sciences*. Published online May 1, 2017. doi:10.1073/PNAS.1621512114

5. Geisberger R, Lamers M, Achatz G. The riddle of the dual expression of IgM and IgD. *Immunology*. 2006;118(4):429-437. doi:10.1111/j.1365-2567.2006.02386.x

6. Maity PC, Blount A, Jumaa H, Ronneberger O, Lillemeier BF, Reth M. B cell antigen receptors of the IgM and IgD classes are clustered in different protein islands that are altered during B cell activation. *Sci Signal*. 2015;8(394):ra93. doi:10.1126/scisignal.2005887

7. Rougé L, Chiang N, Steffek M, et al. Structure of CD20 in complex with the therapeutic monoclonal antibody rituximab. *Science (1979)*. 2020;367(6483):1224-1230. doi:10.1126/science.aba3526

8. Deans JP, Li H, Polyak MJ. CD20-mediated apoptosis: signalling through lipid rafts. *Immunology*. 2002;107(2):176-182. Accessed July 16, 2015. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1782791&tool=pmcentrez&rendertype=abstract

9. Reinhardt SCM, Masullo LA, Baudrexel I, et al. Ångström-resolution fluorescence microscopy. *Nature*. 2023;617(7962):711-716. doi:10.1038/s41586-023-05925-9

10. Pavlasova G, Mraz M. The regulation and function of CD20: An "enigma" of B-cell biology and targeted therapy. *Haematologica*. 2020;105(6):1494-1506. doi:10.3324/haematol.2019.243543

11. Riley JK, Sliwkowski MX. CD20: a gene in search of a function. *Semin Oncol*. 2000;27(6 Suppl 12):17-24.

12. Müller M, Gräbnitz F, Barandun N, et al. Light-mediated discovery of surfaceome nanoscale organization and intercellular receptor interaction networks. *Nat Commun*. 2021;12(1). doi:10.1038/s41467-021-27280-x

13. Arp AB, Abel Gutierrez A, ter Beest M, et al. CD70 recruitment to the immunological synapse is dependent on CD20 in B cells. *Proceedings of the National Academy of Sciences*. 2025;122(16). doi:10.1073/pnas.2414002122

14. Kosmas C, Stamatopoulos K, Stavroyianni N, Tsavaris N, Papadaki T. Anti-CD20-based therapy of B cell lymphoma: State of the art. *Leukemia*. 2002;16(10):2004-2015. doi:10.1038/sj.leu.2402639

15. Smith MR. Rituximab (monoclonal anti-CD20 antibody): Mechanisms of action and resistance. *Oncogene*. 2003;22(47):7359-7368. doi:10.1038/sj.onc.1206939

16. Margoni M, Preziosa P, Filippi M, Rocca MA. Anti-CD20 therapies for multiple sclerosis: current status and future perspectives. *J Neurol*. 2022;269(3):1316-1334. doi:10.1007/s00415-021-10744-x

17. Jazirehi AR, Bonavida B. Cellular and molecular signal transduction pathways modulated by rituximab (rituxan, anti-CD20 mAb) in non-Hodgkin's lymphoma: Implications in chemosensitization and therapeutic intervention. *Oncogene*. 2005;24(13):2121-2143. doi:10.1038/sj.onc.1208349

18. Klein C, Jamois C, Nielsen T. Anti-CD20 treatment for B-cell malignancies: current status and future directions. *Expert Opin Biol Ther*. 2021;21(2):161-181. doi:10.1080/14712598.2020.1822318

19.   Payandeh Z, Bahrami AA, Hoseinpoor R, et al. The applications of anti-CD20
antibodies to treat various B cells disorders. *Biomedicine and Pharmacotherapy*.
2019;109:2415-2426. doi:10.1016/j.biopha.2018.11.121

20.   Boross P, Leusen JHW. Mechanisms of action of CD20 antibodies. *Am J Cancer Res*.
2012;2(6):676-690.

21.   Kläsener K, Jellusova J, Andrieux G, et al. CD20 as a gatekeeper of the resting state of
human B cells. *Proc Natl Acad Sci U S A*. 2021;118(7). doi:10.1073/pnas.2021342118

22.   Salzer E, Santos-Valente E, Keller B, Warnatz K, Boztug K. Protein Kinase C δ: a
Gatekeeper of Immune Homeostasis. *J Clin Immunol*. 2016;36(7):631-640.
doi:10.1007/s10875-016-0323-0

23.   Lim PS, Sutton CR, Rao S. Protein kinase C in the immune system: From signalling to
chromatin regulation. *Immunology*. 2015;146(4):508-522. doi:10.1111/imm.12510

24.   Pracht C, Minguet S, Leitges M, Reth M, Huber M. Association of protein kinase C-δ
with the B cell antigen receptor complex. *Cell Signal*. 2007;19(4):715-722.
doi:10.1016/j.cellsig.2006.07.023

25.   Sjef Verbeek J, Hirose S, Nishimura H. The complex association of fcγriib with
autoimmune susceptibility. *Front Immunol*. 2019;10(OCT).
doi:10.3389/fimmu.2019.02061

26.   Karnell JL, Dimasi N, Karnell FG, et al. CD19 and CD32b Differentially Regulate
Human B Cell Responsiveness. *The Journal of Immunology*. 2014;192(4):1480-1490.
doi:10.4049/jimmunol.1301361

27.   McCaleb MR, Miranda AM, Ratliff KC, Torres RM, Pelanda R. CD19 Is Internalized
Together with IgM in Proportion to B Cell Receptor Stimulation and Is Modulated by
Phosphatidylinositol 3-Kinase in Bone Marrow Immature B Cells. *Immunohorizons*.
2023;7(1):49-63. doi:10.4049/immunohorizons.2200092

28.   Kläsener K, Yang J, Reth M. *Study B Cell Antigen Receptor Nano-Scale Organization*
by in Situ Fab Proximity Ligation Assay. Vol 1707.; 2018. doi:10.1007/978-1-4939-
7474-0_12

29.   Tinti M, Johnson C, Toth R, Ferrier DEK, Mackintosh C. Evolution of signal
multiplexing by 14-3-3-binding 2R-ohnologue protein families in the vertebrates. *Open*
Biol. 2012;2(7):120103. doi:10.1098/rsob.120103

30.   Pitasse-Santos P, Hewitt-Richards I, Abeywickrama Wijewardana Sooriyaarachchi
MD, Doveston RG. Harnessing the 14-3-3 protein–protein interaction network. *Curr*
Opin Struct Biol. 2024;86. doi:10.1016/j.sbi.2024.102822

31.   Omasits U, Ahrens CH, Müller S, Wollscheid B. Protter: interactive protein feature
visualization and integration with experimental proteomic data. *Bioinformatics*.
2014;30(6):884-886. doi:10.1093/bioinformatics/btt607

32.   Ivaldi C, Martin BR, Kieffer-Jaquinod S, et al. Proteomic analysis of S-Acylated
proteins in human B cells reveals palmitoylation of the immune regulators CD20 and
CD23. *PLoS One*. 2012;7(5). doi:10.1371/journal.pone.0037187

33.   Li Z, Ju X, Silveira PA, Abadir E, Hsu W hsun. CD83 : Activation Marker for Antigen
Presenting Cells and Its Therapeutic Potential. 2019;10(June):1-9.
doi:10.3389/fimmu.2019.01312

34.   Ren Y, Li R, Zheng Y, Busch H. *Cloning and Characterization of GEF-H1, a*
Microtubule-Associated Guanine Nucleotide Exchange Factor for Rac and Rho
GTPases*. Vol 273.; 1998. http://www.jbc.org

35.   Zenke FT, Krendel M, DerMardirossian C, King CC, Bohl BP, Bokoch GM. p21-
activated Kinase 1 Phosphorylates and Regulates 14-3-3 Binding to GEF-H1, a
Microtubule-localized Rho Exchange Factor. *Journal of Biological Chemistry*.
2004;279(18):18392-18400. doi:10.1074/jbc.M400084200

36.   Krendel M, Zenke FT, Bokoch GM. Nucleotide exchange factor GEF-H1 mediates
cross-talk between microtubules and the actin cytoskeleton. *Nat Cell Biol*.
2002;4(4):294-301. doi:10.1038/ncb773

37. Meiri D, Marshall CB, Mokady D, et al. Mechanistic insight into GPCR-mediated
activation of the microtubule-associated RhoA exchange factor GEF-H1. *Nat Commun*.
2014;5. doi:10.1038/ncomms5857
38. Kashyap AS, Fernandez-Rodriguez L, Zhao Y, et al. GEF-H1 Signaling upon
Microtubule Destabilization Is Required for Dendritic Cell Activation and Specific
Anti-tumor Responses. *Cell Rep*. 2019;28(13):3367-3380.e8.
doi:10.1016/j.celrep.2019.08.057
39. Kopf A, Kiermaier E. Dynamic Microtubule Arrays in Leukocytes and Their Role in
Cell Migration and Immune Synapse Formation. *Front Cell Dev Biol*. 2021;9.
doi:10.3389/fcell.2021.635511
40. Kosoff R, Chows HY, Radus M, Chernoffs J. Pak2 kinase restrains mast cell FcεRI
receptor signaling through modulation of rho protein guanine nucleotide exchange
factor (GEF) activity. *Journal of Biological Chemistry*. 2013;288(2):974-983.
doi:10.1074/jbc.M112.422295
41. Waheed F, Speight P, Dan Q, Garcia-Mata R, Szaszi K. Affinity precipitation of active
Rho-GEFs using a GST-tagged mutant Rho protein (GST-RhoA(G17A)) from
epithelial cell lysates. *J Vis Exp*. 2012;(61). doi:10.3791/3932
42. Sluchanko NN, Gusev NB. 14-3-3 Proteins and regulation of cytoskeleton.
Biochemistry (Moscow). 2010;75(13):1528-1546. doi:10.1134/S0006297910130031
43. Brandwein D, Wang Z. Interaction between Rho GTPases and 14-3-3 proteins. *Int J*
Mol Sci. 2017;18(10). doi:10.3390/ijms18102148
44. Julian L. *Investigating the Role of Caspase Cleavage of ROCK1 in Tissue Homeostasis*
and Tumour Development.; 2015.
45. Olson MF. Rho-associated coiled-coil containing kinases (ROCK): structure,
regulation, and functions. *Small GTPases*. 2014;5:e29846. doi:10.4161/sgtp.29846
46. Birkenfeld J, Nalbant P, Yoon SH, Bokoch GM. Cellular functions of GEF-H1, a
microtubule-regulated Rho-GEF: is altered GEF-H1 activity a crucial determinant of
disease pathogenesis? *Trends Cell Biol*. 2008;18(5):210-219.
doi:10.1016/j.tcb.2008.02.006
47. Pineau J, Pinon L, Mesdjian O, Fattaccioli J, Duménil AML, Pierobon P. Microtubules
restrict F-actin polymerization to the immune synapse via GEF-H1 to maintain polarity
in lymphocytes. Published online January 3, 2022. doi:10.1101/2022.01.03.473915
48. Mandelkow EM, Mandelkow E, Milligan RA. *Microtubule Dynamics and Microtubule*
Caps: A Time-Resolved Cryo-Electron Microscopy Study.
49. Downing KH, Nogales E. Cryoelectron Microscopy Applications in the Study of
Tubulin Structure, Microtubule Architecture, Dynamics and Assemblies, and
Interaction of Microtubules with Motors. In: *Methods in Enzymology*. Vol 483.
Academic Press Inc.; 2010:121-142. doi:10.1016/S0076-6879(10)83006-X
50. Basu A, Worth F, States U. Two Faces of Protein Kinase C δ: The Contrasting Roles
of PKC δ in Cell Survival and Cell Death. 2015;(August). doi:10.1100/tsw.2010.214
51. Gada KD, Logothetis DE. PKC regulation of ion channels: The involvement of PIP2.
Journal of Biological Chemistry. 2022;298(6). doi:10.1016/j.jbc.2022.102035
52. Miyamoto. Increased proliferation of B cells and auto-immunity in mice lacking
protein kinase Cd. *Nature*. 2002;(volume 416):865.
53. Sun Kuehn H, Niemela JE, Rangel-Santos A, et al. Loss-of-function of the protein
kinase C d (PKCd) causes a B-cell lymphoproliferative syndrome in humans. *Blood*.
2013;121(16):3117-3125. doi:10.1182/blood
54. Merino-Cortés S V, Gardeta SR, Roman-Garcia S, et al. *Diacylglycerol Kinase ζ*
Promotes Actin Cytoskeleton Remodeling and Mechanical Forces at the B Cell
Immune Synapse. Vol 13.; 2020. https://www.science.org
55. Chari R, Kim S, Murugappan S, Sanjay A, Daniel JL, Kunapuli SP. Lyn, PKC-δ,
SHIP-1 interactions regulate GPVI-mediated platelet-dense granule secretion. *Blood*.
2009;114(14):3056-3063. doi:10.1182/blood-2008-11-188516

56. Kajimoto T, Sawamura S, Tohyama Y, Mori Y, Newton AC. Protein kinase C δ-specific activity reporter reveals agonist-evoked nuclear activity controlled by Src family of kinases. *Journal of Biological Chemistry*. 2010;285(53):41896-41910. doi:10.1074/jbc.M110.184028

57. Johnson C, Crowther S, Stafford MJ, Campbell DG, Toth R, MacKintosh C. Bioinformatic and experimental survey of 14-3-3-binding sites. *Biochemical Journal*. 2010;427(1):69-78. doi:10.1042/BJ20091834

58. Pennington K, Chan T, Torres M, Andersen J. The dynamic and stress-adaptive signaling hub of 14-3-3: emerging mechanisms of regulation and context-dependent protein–protein interactions. *Oncogene*. 2018;37(42):5587-5604. doi:10.1038/s41388-018-0348-3

59. Paul AL, Denison FC, Schultz ER, Zupanska AK, Ferl RJ. 14-3-3 Phosphoprotein interaction networks - Does isoform diversity present functional interaction specification? *Front Plant Sci*. 2012;3(AUG). doi:10.3389/fpls.2012.00190

60. Joo E, Olson MF. Regulation and functions of the RhoA regulatory guanine nucleotide exchange factor GEF-H1. *Small GTPases*. 2021;12(5-6):358-371. doi:10.1080/21541248.2020.1840889

61. Azoitei ML, Noh J, Marston DJ, et al. Spatiotemporal dynamics of GEF-H1 activation controlled by microtubule- And Src-mediated pathways. *Journal of Cell Biology*. 2019;218(9):3077-3097. doi:10.1083/JCB.201812073

62. Mok YK, Lo KWH, Zhang M. Structure of Tctex-1 and Its Interaction with Cytoplasmic Dynein Intermediate Chain. *Journal of Biological Chemistry*. 2001;276(17):14067-14074. doi:10.1074/jbc.M011358200

63. Meiri D, Marshall CB, Greeve MA, et al. Mechanistic Insight into the Microtubule and Actin Cytoskeleton Coupling through Dynein-Dependent RhoGEF Inhibition. *Mol Cell*. 2012;45(5):642-655. doi:10.1016/j.molcel.2012.01.027

64. van de Ven AAJM, Compeer EB, Bloem AC, et al. Defective calcium signaling and disrupted CD20-B-cell receptor dissociation in patients with common variable immunodeficiency disorders. *J Allergy Clin Immunol*. 2012;129(3):755-761.e7. doi:10.1016/j.jaci.2011.10.020

65. Li H, Ayer LM, Polyak MJ, et al. The CD20 calcium channel is localized to microvilli and constitutively associated with membrane rafts: Antibody binding increases the affinity of the association through an epitope-dependent cross-linking-independent mechanism. *Journal of Biological Chemistry*. 2004;279(19):19893-19901. doi:10.1074/jbc.M400525200

66. Schober R, Waldherr L, Schmidt T, et al. STIM1 and Orai1 regulate Ca 2+ microdomains for activation of transcription. *Biochim Biophys Acta Mol Cell Res*. 2019;1866(7):1079-1091. doi:10.1016/j.bbamcr.2018.11.001

67. Fais S, Malorni W. Leukocyte uropod formation and membrane/cytoskeleton linkage in immune interactions. *J Leukoc Biol*. 2003;73(5):556-563. doi:10.1189/jlb.1102568

68. Saltukoglu D, Özdemir B, Holtmannspötter M, et al. Plasma membrane topography governs the 3D dynamic localization of IgM B cell antigen receptor clusters. *EMBO J*. 2023;42(4):e112030. doi:10.15252/embj.2022112030

69. Mattila PK, Batista FD, Treanor B. Dynamics of the actin cytoskeleton mediates receptor cross talk: An emerging concept in tuning receptor signaling. *J Cell Biol*. 2016;212(3).

70. Wang J, Lin F, Wan Z, et al. *I M M U N O L O G Y Profiling the Origin, Dynamics, and Function of Traction Force in B Cell Activation*.; 2018. https://www.science.org

71. Sanchez-Madrid F. SJ. Bringing up the rear- defining the roles of the uropod. *Nat Rev Mol Cell Biol*. 2009;Volume 10:353-359.

72. Roberts PJ, Mitin N, Keller PJ, et al. Rho family GTPase modification and dependence on CAAX motif-signaled posttranslational modification. *Journal of Biological Chemistry*. 2008;283(37):25150-25163. doi:10.1074/jbc.M800882200

73. Lehtimäki JI, Rajakylä EK, Tojkander S, Lappalainen P. Generation of stress fibers through myosin-driven reorganization of the actin cortex. *Elife*. 2021;10:1-43. doi:10.7554/eLife.60710

74. Rey M, Sanchez-Madrid F, Valenzuela-Fernandez A. *The Role of Actomyosin and the Microtubular Network in Both the Immunological Synapse and T Cell Activation*. Vol 12.; 2007.

75. Reth M. Discovering immunoreceptor coupling and organization motifs. *Front Immunol*. 2023;14. doi:10.3389/fimmu.2023.1253412

76. Shanehbandi D, Majidi J, Kazemi T, Baradaran B, Aghebati-Maleki L. CD20-based Immunotherapy of B-cell Derived Hematologic Malignancies. *Curr Cancer Drug Targets*. 2017;17(5):423-444. doi:10.2174/1568009617666170109151128

77. Deans JP, Li H, Polyak MJ. *CD20-Mediated Apoptosis: Signalling through Lipid Rafts*.; 2002.

78. Chang YC, Nalbant P, Birkenfeld J, Chang ZF, Bokoch GM. 3GEF-H1 couples nocodazole-induced microtubule disassembly to cell contractility via RhoA. *Mol Biol Cell*. 2008;19(5):2147-2153. doi:10.1091/mbc.E07-12-1269

79. Kozlova V, Ledererova A, Ladungova A, et al. CD20 is dispensable for B-cell receptor signaling but is required for proper actin polymerization, adhesion and migration of malignant B cells. *PLoS One*. 2020;15(3). doi:10.1371/journal.pone.0229170

80. Hiraga J, Tomita A, Sugimoto T, et al. Down-regulation of CD20 expression in B-cell lymphoma cells after treatment with rituximab-containing combination chemotherapies: its prevalence and clinical significance. Published online 2009. doi:10.1182/blood-2008-08

81. Marshalek JP, Dragan M, Tomassetti S, Labarbera K, Amaya A, Har ; *E19537 Publication Only CD20 and CD19 Expression Loss in Relapsed or Refractory B-Cell Non-Hodgkin Lymphoma: A Retrospective Cohort*.; 2022.

82. Däbritz JHM, Yu Y, Milanovic M, et al. CD20-targeting immunotherapy promotes cellular senescence in B-cell lymphoma. *Mol Cancer Ther*. 2016;15(5):1074-1081. doi:10.1158/1535-7163.MCT-15-0627

83. Chen Y, Chen L, Lun ATL, Baldoni PL, Smyth GK. edgeR v4: powerful differential analysis of sequencing data with expanded functionality and improved support for small counts and larger datasets. *Nucleic Acids Res*. 2025;53(2). doi:10.1093/nar/gkaf018

84. Wu T, Hu E, Xu S, et al. clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *Innovation*. 2021;2(3). doi:10.1016/j.xinn.2021.100141

85. Subramanian A, Tamayo P, Mootha VK, et al. *Gene Set Enrichment Analysis: A Knowledge-Based Approach for Interpreting Genome-Wide Expression Profiles*.; 2005. www.pnas.orgcgidoi10.1073pnas.0506580102

86. Wang JC, Bolger-Munro M, Gold MR. Visualizing the actin and microtubule cytoskeletons at the b-cell immune synapse using stimulated emission depletion (STED) microscopy. *Journal of Visualized Experiments*. 2018;2018(134). doi:10.3791/57028

87. Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat Biotechnol*. 2008;26(12):1367-1372. doi:10.1038/nbt.1511

88. Cox J, Neuhauser N, Michalski A, Scheltema RA, Olsen J V., Mann M. Andromeda: A peptide search engine integrated into the MaxQuant environment. *J Proteome Res*. 2011;10(4):1794-1805. doi:10.1021/pr101065j

89. Bender J, Mühlhäuser WWD, Zimmerman JP, Drepper F, Warscheid B. Autoprot: Processing, Analysis and Visualization of Proteomics Data in Python. Published online January 23, 2024. doi:10.1101/2024.01.18.571429