1 MANUSCRIPT TITLE

2 Tracing the vertebrate selenoproteome reveals expansions in ray-finned fishes and

3 convergent depletions in tetrapods

4 AUTHORS

5 Max Ticó[1], Marco Mariotti[1*]

6 [1] Department of Genetics, Microbiology and Statistics, Universitat de Barcelona,
7 Barcelona, Catalonia, 08024, Spain

8 * To whom correspondence should be addressed. Email: marco.mariotti@ub.edu

9 ABSTRACT

Selenoproteins incorporate the rare selenium-containing amino acid selenocysteine

(Sec) and play crucial roles for redox homeostasis, stress response, and hormone

regulation. Sec is inserted by co-translational recoding of the UGA codon, normally a

stop. As consequence, selenoproteins are often misannotated in public databases and

require specialized bioinformatic methods and resources.

Here, we present a refined characterization of the composition and evolution of the

vertebrate selenoproteome. Based on analyses of 19 gene families across hundreds of

genomes, we show that extant selenoproteomes were shaped by extensive gene

duplications (53), losses (41), and Sec-to-cysteine (Cys) conversions (19). Tetrapods

including mammals encode 24-25 selenoproteins, with variations in 6 families. Notably,

the same genes underwent convergent evolutionary events in multiple tetrapods,

namely Sec-to-Cys substitutions (SELENOU1, GPX6) and gene losses (SELENOV). In

contrast, ray-finned fish exhibit larger and more dynamic selenoproteomes, reinforcing

the hypothesis that the selective advantage of Sec is stronger in aquatic environments.

We detected selenoprotein duplications spread across the actinopterygian clade

involving 13 families, mainly involved in antioxidant defense. The richest

selenoproteomes were found in Salmoninae and Cyprinoidei fish with 55 and 45
selenoproteins, respectively, owing to whole genome duplications. Among our findings,
the SELENOP family stands out in lampreys, carrying up to an unprecedented 162 UGAs
putatively recoded to Sec.

Our study presents the most comprehensive evolutionary map of vertebrate
selenoproteins to date and delineates the specific selenoproteome of each lineage,
establishing a foundational framework for selenium biology research in the era of
biodiversity genomics.

INTRODUCTION

Selenium is an essential trace element for human health. The biological effects of
selenium are mostly mediated by its incorporation in the non-canonical amino acid
selenocysteine (Sec) (Fairweather-Tait et al. 2011; Labunskyy et al. 2014; Mariotti and
Gladyshev 2022). Sec is the essential component of selenoproteins, a unique class of
proteins with crucial functions in both normal physiology and cancer. Sec is often
referred to as the 21st amino acid due to its distinctive biosynthetic pathway and
incorporation mechanism, which differ fundamentally from those of the 20 canonical
amino acids. Importantly, Sec is inserted in correspondence to the UGA codon, which is
normally a stop codon. The UGA is co-translationally recoded to Sec in selenoprotein
mRNAs in response to local sequence elements, the major of which is called SECIS
element mechanism (Howard and Copeland 2019; Copeland and Howard 2021).
Moreover, unlike standard amino acids which are synthesized then charged onto their
cognate transfer RNAs (tRNAs), Sec is synthesized directly on its own dedicated tRNA,
tRNASec, through a multi-step enzymatic process. The selenium required for this
reaction is provided in the form of selenophosphate, synthesized by protein SEPHS2, an

essential enzyme in Sec metabolism which, intriguingly, is itself a selenoprotein (Manta

et al. 2022).

Since their discovery in the 1970s, Sec and selenoproteins have been extensively studied

across diverse fields of biology (Flohé 2009; Davy and Castellano 2018). Indeed, despite

representing a small fraction of protein-coding genes, selenoproteins perform many

critical functions in redox homeostasis, stress response, hormone maturation, and

protein quality control, thus playing key roles in both physiology and cancer biology. This

has prompted extensive research into their enzymatic functions, selenium metabolism,

and the unique mechanism of Sec biosynthesis. Selenium itself, a trace element with a

narrow margin between deficiency and toxicity, has also drawn interest in nutritional

and physiological contexts (Fairweather-Tait et al. 2011). As a result, selenoproteins are

investigated in a broad range of species, from standard model organisms (e.g., mouse,

zebrafish) to agriculturally relevant animals (e.g., cattle, chicken, salmon). Naturally,

understanding the full set of selenoproteins encoded by an organism is essential for the

meaningful design and interpretation of experiments to elucidate selenium biology.

The human and mouse selenoproteomes were first described in 2003 as comprising 25

and 24 Sec-encoding genes, respectively (Kryukov et al. 2003). Subsequent studies

expanded the known selenoproteomes of selected vertebrates and metazoans

(Castellano et al. 2004; Castellano et al. 2005; Novoselov et al. 2006; Shchedrina et al.

2007; Jiang et al. 2010; Jiang et al. 2012). In our seminal paper in 2012, we provided the

first description of selenoproteome composition and evolution across diverse

vertebrates (Mariotti et al. 2012). Although highly influential, that study was constrained

by the limited availability of sequenced genomes at the time (44).

In the last 10 years, major advances in sequencing and the rise of biodiversity genomics

provided an unprecedented wealth of genomic data (Lewin et al. 2018; Mc Cartney et al.

2024). However, due to their deviation from the standard genetic code, selenoproteins

have not kept pace with the expansion of automated genomics resources. At present,

90% and 50% of selenoprotein genes have substantial annotation errors in Ensembl and

NCBI genomes, respectively (Ticó et al. 2024). This also led to unreliable selenoprotein

data in evolutionary resources dedicated to orthology (e.g. Ensembl Compara (Vilella et

al. 2009), EggNOG (Powell et al. 2014)).

In this manuscript, we present a refined comprehensive characterization of the

selenoproteome of vertebrates, obtained by meticulously tracing the evolution of 19

protein families across hundreds of genomes through sequence analysis and

phylogenetic reconstruction. Our work provides an essential evolutionary framework to

understand the selenoprotein content of any vertebrate genome, expanding on previous

efforts by an order of magnitude.

87 RESULTS

We downloaded 315 genome assemblies from Ensembl representative of all major

lineages of vertebrates, complemented by 25 NCBI genomes from early vertebrates and

outgroups (Methods). Next, we used the software Selenoprofiles to identify homologs

belonging to any of the known metazoan selenoprotein families in those genomes,

resulting in 18,579 gene predictions. Phylogenetic reconstruction was then performed for

each family, and gene and species trees were manually inspected to infer gene

evolutionary events. These comprise gene duplications, gene losses, and the

selenoprotein-specific event of the conversion of a Sec residue to Cys. In selenoprotein

evolution, Cys to Sec events are also possible, but have been observed only along

evolutionary distances much larger than encompassed by vertebrates (Mariotti et al.

2015; Zhang et al. 2022). Each branch of gene trees was manually annotated as a separate

orthologous group, assigning gene names as consistent as possible with existing

resources and literature. Each putative event was critically assessed, following criteria of

maximum parsimony, and seeking consistency with the species tree and support by

independent data (e.g. RNA sequences, other genome assemblies from NCBI) (Methods).

Thanks to the vast diversity of organisms in our dataset, the great majority of events were

shared among multiple species, leading to phylogenetically coherent patterns of gene

occurrence that allowed reliable evolutionary inferences. Yet, two lineages stood out for

their divergence from this general trend. First, bird genomes exhibited many apparent

gene losses scattered across lineages, in a pattern that would imply dozens of

independent events (Supplementary Figure S1). This included the apparent absence of

SEPHS2, a gene that nevertheless is required for biosynthesis of all other selenoproteins

(Mariotti et al. 2015). We ascribed this pattern to a known bias of bird assemblies: they

systematically lack certain GC-rich regions that present technical challenges for

sequencing (Botero-Castro et al. 2017). Indeed, we found RNA evidence of expression

for these putatively lost genes, indicating that they are indeed present in the real

biological genomes of birds. Ultimately, we dismissed all selenoprotein losses in Sauria

(which includes birds) as artefactual.

Second, the fish lineage of Cyprinoidei (e.g. carps, zebrafish *Danio rerio*) exhibited a

massive amount of apparent gene duplications (Supplementary Figure S1), which we

could not precisely disentangle by phylogenetic analysis. We realized that the sublineage

of Cyprininae (e.g. carps) reportedly underwent extensive recent whole genome

duplications and polyploidy, which greatly complicated this task. Therefore, we decided

to limit our analysis to the evolutionary events that occurred at the root of Cyprinoidei,

i.e. shared between zebrafish and Cyprininae, and dismiss the rest.

Overview of the vertebrate selenoproteome

We identified selenoproteins belonging to 19 protein families. The most represented

families were glutathione peroxidases with 1,529 Sec-encoding genes, iodothyronine

deiodinases with 999, and thioredoxin reductases with 721. In general, ray-finned fishes
(Actinopterygii) exhibited richer selenoproteomes (33-55 selenoprotein genes) than
tetrapods (24-25 genes).

While our analyses replicated well earlier reports performed at a smaller scale, we
detected many novel events, particularly gene duplications. The majority affected
Actinopterygii and its sublineages, whose selenoproteome appeared not only larger, but
also more dynamic. In total, we detected 52 gene duplications (40 in Actinopterygii), and
41 gene losses (25 in Actinopterygii). The most selenoprotein-rich organisms were
Salmoninae (e.g. trouts, salmons) with 55 selenoprotein genes, 17 of which emerged by
duplication specifically in this lineage, and Cyprinoidei with 45 selenoproteins, 7 of which
lineage-specific. On the other hand, Sec-to-Cys conversions were more prevalent in
Tetrapoda: we detected 19 of these events overall, 16 of which were in tetrapods.
Hereafter, we describe the selenoproteome evolution for each family separately.

Figure 1. Evolution of the vertebrate selenoproteome. *The evolutionary events shaping*

selenoprotein genes are shown along the species tree. Leaf nodes represent either

individual species (italicized) or collapsed lineages; in the latter case, the number in
brackets indicates how many species from that lineage are included in our main
dataset. Asterisks ('') denote lineages with apparent gene losses that we interpret as*
artefacts caused by known issues in genome assembly quality. Bold numbers in the
rightmost column indicate the number of selenoprotein genes detected in each species
or lineage. Grey rectangles highlight taxonomic clades of interest; "Protacant" refers to
Protacanthopterygii. Animal silhouettes were sourced from https://www.phylopic.org.

<u>Glutathione peroxidases (GPX)</u>

Glutathione peroxidases (GPX) are key antioxidant enzymes that protect cells from
oxidative damage by reducing hydrogen peroxide and lipid peroxides using glutathione
as an electron donor (Toppo et al. 2008; Mariotti et al. 2012). Mammals possess eight
GPX homologs with specialized tissue distribution and functions, including five
selenoproteins: GPX1 (ubiquitous), GPX2 (gastrointestinal), GPX3 (extracellular, kidney),
GPX4 (membrane-associated, ubiquitous), GPX6 (olfactory system); and three Cys
paralogs: GPX5, GPX7, and GPX8. Our phylogenetic analysis (Figure 2, Supplementary
Figure S2) confirms a well-supported evolutionary classification of GPX proteins,
identifying three major groups: GPX1/GPX2, GPX3/GPX5/GPX6, and GPX4/GPX7/GPX8.
These findings align with previous studies on GPX evolution (Toppo et al. 2008; Mariotti
et al. 2012; Xue et al. 2022).

GPX1, GPX2, GPX3, GPX4 are present in all vertebrates, with the only exception of
Cyclostomata that lost GPX1. On the other hand, GPX6 originated at the root of placentals
and exhibited a very dynamic evolution. Multiple independent GPX6 Sec-to-Cys
conversions had been reported (Mariotti et al. 2012), which we confirmed and expanded
in our analyses (Figures 1, 2b, 2d, 3; Supplementary Figure S2). We observed Cys
conversions in Platyrrhini primates (New World monkeys, e.g. marmoset, titi), Muroidea
rodents (e.g. mouse, rat, hamster), Lagomorpha (e.g. rabbit, hare, pika), Feliformia (e.g.

cats, hyenas), Bathyergidae rodents (e.g. mole-rats), and *Marmota marmota marmota*

(Figure 3). Furthermore, GPX6 was lost altogether in Chiroptera (bats).

We identified several lineage-specific GPX gene duplications across vertebrates. In

Cyclostomata (jawless fish), we observed two lineage-specific gene duplications

resulting that yielded GPX2b and GPX3b2. In Chondrichthyes (cartilaginous fish), a

duplication of GPX1 gave rise to a distinct Sec-paralog, GPX1c. We also confirmed three

two previously reported GPX gene duplications at the root of Actinopterygii, leading to

the emergence of GPX1b, and GPX4b. These genes were found in all investigated species

within this lineage, except for Oryziinae (e.g. medaka), which lost GPX1b. Curiously,

Oryziinae also exhibited a Cys-containing paralog emerged by duplication of GPX4b,

named GPX4b2. A third gene, GPX3b, was also reported in Actinopterygii; our analysis

indicates that it originated at the root of Vertebrata, and was subsequently lost in

Tetrapoda.

We also observed additional selenoprotein gene duplications in specific branches of

Actinopterygii: three in Salmoninae yielding GPX1b2, GPX3a2, and GPX4a2; and one in

Cyprinoidei yielding GPX1a2. These duplications were previously reported in an analysis

of 8 fish genomes (Xue et al. 2022). Our study demonstrates that they are lineage-

specific events, so that we have assigned new names to the duplicated genes to reflect

their precise evolutionary trajectories. The correspondence with (Xue et al. 2022) are

provided in Supplementary Table T1.

Figure 2. Phylogeny of the GPX and TXNRD families. *a) Topology of the GPX family, split*

*into three clades shown in panels **c–e**. **b)** Zoomed section of the GPX6 gene tree, used*

here to illustrate this form of representation, abundantly used throughout this paper.

Each leaf correspond to a putative gene whose protein sequence was included inthe

tree. Leaves are color-coded by their label, assigned by Selenoprofiles by examining

which codon aligns to the Sec position in the profile (e.g. green for Sec-UGA, red for

Cys, orange for unaligned). Moving outward, you will see an equally colored

representation of the corresponding aligned protein, which shows its non-gap positions

and includes black lines to map where introns are located in the coding sequence. Most

intronless genes are pseudogenes, which constitute the prevalent false positives of

Selenoprofiles. The outmost circle provides a manual annotation of gene clades, with

colors that indicate the taxonomy of the source of those genes (e.g. blue for

Actinopterygii, green for mammals). Internal nodes of the gene tree represent either

speciation events (red) or duplication events (blue), determined via the species overlap

algorithm. The latter are displayed with size proportional to the number of species

supporting the duplication. Duplication events deemed genuine via manual analysis are

*marked with a star. The GPX gene tree visualization is split into groups: **c)** GPX1/GPX2,*

***d)** GPX3/GPX5/GPX6, **e)** GPX4/GPX7/GPX8. **f)** Gene tree visualization of the TXNRD*

family. Note that selected clades of the gene tree that did not exhibit evolutionary

events were collapsed (e.g. see GPX7 and GPX8 in panel e). Expanded gene trees are

available as Supplementary Figures S2 (GPX) and S5 (TXNRD).

Figure 3. GPX6 multiple sequence alignment. *Selenocysteine residues are shown in green while Cysteine residues are shown in red. Species containing a GPX6–Cysteine homolog are highlighted by lineage.*

Iodothyronine deiodinases (DIO)

Iodothyronine deiodinases (DIO) regulate thyroid hormone metabolism by catalyzing the activation or inactivation of thyroxine (T4) and triiodothyronine (T3) (Darras and Van Herck 2012). The vertebrate DIO family includes three main genes: DIO1, which is involved in peripheral thyroid hormone conversion; DIO2, which plays a critical role in local T3 activation; and DIO3, which inactivates excess thyroid hormones, preventing hyperthyroidism. Through our analyses (Figures 1, 4, Supplementary Figures S1, S3), we detected these three genes in all vertebrate lineages, with the exception of Cyclostomata that lacked DIO1. This gene is present in all investigated Chordates, indicating that DIO1 predate the vertebrate radiation and was then lost in Cyclostomata. All Actinopterygii additionally carried a DIO3 duplicate (DIO3b). We also confirmed two more duplications, one of the ancestral DIO3 gene (also called DIO3a) (Lorgen et al. 2015), to

generate DIO3a2, and the other of DIO2 to yield DIO2b (Figures 1, 4a, 4b, Supplementary Figures S3, S4). Both were previously reported in Salmonids (Lorgen et al. 2015), and we traced them further back to the root of Salmoninae. In Cyclostomata, we identified two lineage-specific duplications resulting in DIO2c and DIO3c.

Figure 4. Phylogeny of the DIO family. *a) Gene tree visualization of the DIO family. See full feature description in Figure 2b. **b)** Protacanthopterygii-specific DIO evolution. Selenoprotein genes are shown as circles. Genes are color-labelled according to panel A. Expanded gene and species trees are available as Supplementary Figures S3 and S4.*

Thioredoxin reductases (TXNRD)

Thioredoxin reductases (TXNRDs) play a central role in maintaining cellular redox balance by catalyzing the reduction of thioredoxins, which are essential for regulating oxidative stress responses, cell proliferation, and redox-regulated signaling cascades (Arnér 2009). Tetrapods have three TXNRD genes: TXNRD1 (cytosolic), TXNRD2 (mitochondrial), and TXNRD3/TGR (testis-specific, containing a peculiar N-terminal Grx domain), each of which participates in distinct physiological processes.

Reportedly, fishes lack the TXNRD3/TGR gene, but their TXNRD1 gene encodes for a Grx-

containing protein as its major isoform, which duplicated to give rise to TXNRD3/TGR in

tetrapods (Dou et al., 2022; Mariotti et al., 2012). Yet, our present analysis (Figures 2f, 5,

Supplementary Figure S5) did not recapitulate TXNRD evolution as previously proposed.

Because we detected TXNRD3 in Chondrichthyes, a lineage predating Actinopterygii, we

argue that this gene emerged earlier than tetrapods, and was subsequently lost in ray-

finned fishes. We did not find TXNRD3 in Cyclostomata, and thus we map the

TXNRD1/TXNRD3 duplication to the root of Gnathostomata. Moreover, because the

Cyclostomata gene clearly clusters with TXNRD1, we identify this clade as the ancestral

one and TXNRD3 as the gene derived from its duplication.

It is reported that the human TXNRD1 gene encodes for a minor transcript isoform that

includes this domain ("TxnRd1_v3"). This is absent in mouse, while in fish it represents

the predominant TXNRD1 variant (Arnér 2009). Therefore, we systemically investigated

the occurrence of Grx domains in the genomic regions of TXNRD1 and TXNRD3 across

all vertebrates (Methods). Our results show that Grx domains are a widespread feature

of TXNRD1 genes (Figure 5). Yet, they have been lost by several tetrapod lineages

independently (Glires, Chiroptera, Sauria, Equus, and Pecora), possibly functionally

replaced by TXNRD3. We argue that the Grx-domain is ancestral feature of TXNRD1.

We further investigated the ancestral state of vertebrates by examining the position of

metazoan outgroups to vertebrates in the TXNRD gene tree (Figure 2f), as well as an

extended gene tree featuring additional outgroups available in PhylomeDB (QFO2020

seed, *Caenorhabditis elegans* phylome ID: 791). Our phylogenetic analyses indicate that

the divergence between TXNRD1/TXNRD3 and TXNRD2 predates the origin of

metazoans, as representatives from both clades are found in diverse animal lineages,

including the nematode *C. elegans*, the annelid *Helobdella robusta*, and the cnidarian

Nematostella vectensis. Secondary events are evident in several lineages: the tunicate

Ciona intestinalis lost TXNRD2, while arthropods such as *Ixodes scapularis* and *D.*

melanogaster lost TXNRD1. Notably, *D. melanogaster* retains two TXNRD2-derived

genes, presumably the result of a secondary duplication. Fitting with our analysis, non-

vertebrate animals carried Grx-domains solely in TXNRD1 genes. Due to this domain,

these invertebrate genes were referred as TGR in literature, despite weak or absent

glutathione reductase activity (Fata et al. 2022). Altogether, we reconstruct that ancestral

vertebrates carried a cytosolic Grx-containing TXNRD1 gene and a mitochondrial

TXNRD2 gene, the former later giving rise to TXNRD3 in jawed vertebrates.

We further detected two TXNRD1 duplications. One occurred in Salmoninae and yielded

TXNRD1b, which retained the Grx domain. Another one gave rise to TXNRD1c in

Acanthomorphata, which did not conserve such domain, and also features an atypical

GNCUG C-terminus (rather than GCUC found across TXNRDs). TXNRD1c was

subsequently convergently lost in several acanthomorph lineages (Figure 1), including

Tetraodontidae, Atherinomorphae, Anabantaria, and Carangaria. Lastly, all Cyclostomata

(except *Eptatretus burgeri*) lost TXNRD2 and thus encode TXNRD1 as their sole gene in

this family.

Figure 5. Phylogenetic distribution of the Grx domain in TXNRD1 and TXNRD3 proteins across vertebrate lineages. *Colors are assigned according to the TXNRD gene tree. A non-collapsed version of this figure is available as Supplementary Figures S6.*

SELENOJ

Selenoprotein J (SELENOJ) shows significant similarity to jellyfish J1-crystallins and, together with them, forms a distinct subfamily within the large family of ADP-ribosylation enzymes. While the homology may suggest a structural role which would be unique among selenoproteins, SELENOJ remains functionally uncharacterized. SELENOJ is reportedly restricted to actinopterygian fishes and sea urchins, with Cys homologs reported in cnidarians (Castellano et al. 2005). Our analyses replicated earlier reports, and additionally highlighted the loss of SELENOJ in Cyclostomata. Moreover, we had previously reported a SELENOJ duplication in medaka and stickleback, yielding SELENOJ2. Here, we traced back this duplication to the root of Euteleosteomorpha

(euteleosts), which included more than half of fishes in our dataset (Figures 1, 6a, 6b,

Supplementary Figures S7, S8). Later, SELENOJ2 was lost independently in

Tetraodontidae (e.g. pufferfish). On the other hand, SELENOJ2 underwent a Sec-to-Cys

conversion in Salmoninae.

<u>SELENOO</u>

Selenoprotein O (SELENOO) is the largest known selenoprotein by molecular weight,

and belongs to a widespread protein family recently functionally characterized as a

mitochondrial pseudokinase that transfers AMP to various protein substrates, implicated

in oxidative stress response and metastasis (Sreelatha et al. 2018; Melo et al. 2025).

Previous studies identified a SELENOO paralog named SELENOO2 in zebrafish, which

was absent in the genomes of fugu, stickleback, medaka, and pufferfish (Mariotti et al.

2012). In our analysis, we replicated these results, and yet we observed the presence of

SELENOO2 in many other fish genomes, including among others all of Cyprinodontoidei

(toothcarps) and Carangaria (e.g. jacks, flatfishes), which implies a complex pattern of

gene evolution (Figures 1, 6c, 6d, Supplementary Figures S9, S10). Ultimately, we

hypothesize that SELENOO2 emerged at the root of Clupeocephala but was

subsequently lost in specific lineages, namely Tetraodontidae, Oryziinae, Anabantaria

(e.g. gourami), and Perciformes (e.g. stickleback).

<u>Methionine sulfoxide reductase B (MSRB)</u>

Methionine sulfoxide reductase B (MSRB, previously known as SELENOR), are enzymes

that reduce oxidized methionine residues, with roles for antioxidant defense and redox-

based modulation of various pathways including actin assembly (Tarrago et al. 2022).

Human possesses one Sec-containing gene, MSRB1, and two Cys paralogs, MSRB2 and

MSRB3. Our analyses indicate that the three genes predate the origin of vertebrates.

Within fish lineages, MSRB1 underwent a complex history of duplications and losses

(Figures 1, 6e, 6f, Supplementary Figures S11, S12). We previously identified a Sec-
containing gene, MSRB1b, in five ray-finned fishes (Mariotti et al. 2012). Our present
analysis refined its origin to the subclade of Osteoglossocephalai, which comprises all
Actinopterygii in our dataset except the MSRB1b-lacking species *Erpetoichthys*
calabaricus and *Lepisosteus oculatus*. This gene was then duplicated in Cyprinoidei
yielding MSRB1b1b, and in Euteleosteomorpha yielding MSRB1b2. The latter gene
duplicated again in Salmoninae, yielding MSRB1b2b, while the Northern pike Esox lucius
(its sister species within Euteleosteomorpha) lost MSRB1b2. On the other hand, MSRB1
has been lost in multiple lineages: Protoacanthopterygii (i.e. the group that includes
Salmoninae as well as *Esox lucius*), Tetraodontidae, Perciformes, Anabantaria,
Cyprinodontoidei, and the Chinese tongue sole *Cynoglossus semilaevis*. Notably, this
reconstruction highlights at least four convergent events of MSRB1 loss in
Percomorphacae, as we found MSRB1 only in two subclades, Carangaria (five genomes)
and non-Perciformes Eupercaria (five genomes). Notably, other selenoprotein families
show similar patterns of convergent losses in this lineage (Figure 1), as discussed later.

Figure 6. Phylogeny of the SELENOJ, SELENOO, and MSRB families. a, c, e) *Gene tree*

visualizations of the SELENOJ, SELENOO, and MSRB families, respectively. For tree

annotation and legend details, see Figure 2b. **b, d, f)** *Actinopterygii species tree*

representations showing the presence of SELENOJ, SELENOO and MSRB paralogs,

respectively. Expanded gene and species trees are available: Supplementary Figure S7,

S8 (SELENOJ), S9, S10 (SELENOO), S11, S12 (MSRB).

<u>SELENOT</u>

Selenoprotein T (SELENOT) is a thioredoxin-like enzyme anchored to the endoplasmic

reticulum (ER) membrane. It is highly conserved throughout evolution, and it has an

essential role in cellular homeostasis and redox regulation (Anouar et al. 2018). Previous

studies reported three SELENOT genes in actinopterygian fishes (Mariotti et al. 2012),

here refined and expanded. We found that most actinopterygian groups possessed two

SELENOT genes, namely SELENOT (aka SELENOT1) and SELENOT2; yet, Salmoninae,

Otophysi and Cyprinoidei exhibited further lineage-specific duplications. Ultimately, we

mapped the origin of SELENOT2 to a duplication of SELENOT1 at the root of

Actinopterygii. SELENOT1 duplicated again to yield SELENOT1b in Otophysi, i.e. the group

including Cyprinoidei (e.g. zebrafish and carps), and Characiphysae (e.g. tetra, piranha).

In Cyprinoidei, further duplications of SELENOT2 and SELENOT1b originated the genes

SELENOT2b and SELENOT1b2, respectively (Figures 1, 7a, Supplementary Figure S13).

Finally, we observed yet another SELENOT1 duplication in Salmoninae, yielding the gene

SELENOT1a2. Interestingly, Cyclostomata also independently expanded this family

through one gene duplication, which yielded a divergent paralog that we designated

SELENOT3.

<u>SELENOU</u>

The Selenoprotein U (SELENOU) family includes selenoprotein genes in birds and fishes,

the latter reported to carry up to three Sec-containing paralogs (Castellano et al. 2004;

Mariotti et al. 2012). Mammals and amphibians are reported to encode only Cys-

homologs, with human containing three paralogs (SELENOU1/PRLX2A,

SELENOU2/PRLX2C, SELENOU3/PRLX2B), involved in diverse functions such as antioxidant defense, prostaglandin biosynthesis, and glycolysis regulation. SELENOU exhibits a scattered pattern of gene occurrence, implying a complex evolutionary history involving gene duplications, losses, and Sec-to-Cys conversions. Here, we provide our best attempt to explain this with current data (Figures 1, 7b, 7c, Supplementary Figures S1, S14).

The split of SELENOU1/2/3 predates metazoa, and all Sec-encoding genes detected in this family belonged to the SELENOU1 branch. Within fish, we mapped a first duplication of the SELENOU1 gene in Osteoglossocephalai, yielding SELENOU1a2. This gene was then lost at the root of the large group of Acanthomorphata, while it converted to Cys homolog in *Esox lucius*. Prior to both of these events, we traced two additional SELENOU1 duplications in Clupeocephala and Acanthomorphata, yielding the SELENOU1c and SELENOU1b genes, respectively. SELENOU1b was then lost at the root of Ovalentaria. Finally, we detected another two independent SELENOU1 duplications in Cyprinoidei and Salmoninae, yielding the gene SELENOU1a1b and SELENOU1a1c, respectively. Within tetrapods, SELENOU1 underwent at least four convergent Sec-to-Cys conversions (Figure 7c), specifically in Theria (i.e. placentals and marsupials), Anura (frogs), Testudines (turtles), and Psittaciformes birds (parrots).

SELENOW and SELENOV

Selenoprotein W (SELENOW) is a widespread glutathione-dependent antioxidant oxidoreductase implicated in resolution of inflammation, with reported roles in muscle growth and differentiation, as well as in protecting neurons from oxidative stress during neuronal development (Jeong et al. 2002; Misra et al. 2023). Selenoprotein V (SELENOV) is a placental-specific putative oxidoreductase expressed in testis and potentially involved in antioxidant defense and ER stress, whose C-terminus shares substantial sequence homology to SELENOW (Zhang et al. 2020).

It was previously reported that SELENOV was lost in *Gorilla gorilla* (Mariotti et al. 2012).

Our present analysis identified an additional case of SELENOV loss, specifically in

Odontoceti (toothed whales, e.g. dolphins, killer whales) (Figures 1, 7d, Supplementary

Figure S15). Regarding the SELENOW branch of this family, mammals have a single gene:

SELENOW, also referred to as SELENOW1. Earlier work, replicated and expanded here,

pinpointed multiple SELENOW homologs in non-mammalian vertebrates, with a clearly

distinct gene clade named SELENOW2. The split of SELENOW1 and SELENOW2

predates the vertebrate radiation. Among tetrapods, we detected SELENOW2 only in

amphibians (Anura), so that we infer that it was lost at the root of Amniota. In fish, the

phylogenetic history of the SELENOW is remarkably complex (FIGURE 7d,

Supplementary Figure S15): it constitutes the most difficult vertebrate selenoprotein

family to solve conclusively, since no simple scenario can explain the observed pattern

of gene presence. In our reconstruction (Figure 1), a duplication of SELENOW2 in

Euteleosteomorpha led to the emergence of SELENOW2c. Further along this lineage,

specifically in Acanthomorphata, SELENOW1 and SELENOW2a were then lost, so that

these genes are absent from the majority of extant fish genomes. On the other hand, we

observed several lineage-specific duplications. Two SELENOW2 duplications occurred

in Cyprinoidei, yielding genes that we named SELENOW2a2 and SELENOW2b.

Salmoninae exhibited another two duplications in the SELENOW2 clade that we

attributed to the SELENOW2 and SELENOW2c genes, yielding SELENOW2a3 and

SELENOW2c2, respectively. Then, SELENOW2c duplicated in Acanthomorphata

yielding SELENOW2c3, and the resulting gene duplication later within this lineage, in

Cyprinodontoidei, giving rise to SELENOW2c3b. A series of scattered losses in the

SELENOW2c3 clade were also detected but challenging to trace with precision. Our

maximum-parsimony reconstruction implies five independent losses of SELENOW2c3

within various Acanthomorphata branches (Figure 1).

Figure 7. Phylogeny of the SELENOT, SELENOU and SELENOW families. a, b, d) Gene

tree visualizations of the SELENOT, SELENOOU and SELENOW families, respectively.

For tree annotation and legend details, see Figure 2b. *c)* SELENOU multiple sequence

alignment. Species are color-labelled according to lineage. Selenocysteine residues are

marked in green and Cysteine residues in red. Expanded gene trees are available:

Supplementary Figures S13 (SELENOT), S14 (SELENOU), S15 (SELENOW).

SELENOP

Selenoprotein P (SELENOP) is the only mammalian selenoprotein with multiple Sec

residues, e.g. 10 in human and mouse, due to its Sec-rich C-terminal tail. SELENOP is a

key player in selenium transport and distribution, serving as the primary selenium carrier

in the bloodstream, delivering selenium to peripheral tissues via receptor-mediated

uptake. We recently characterized the evolution of SELENOP, showing that its Sec

content can vary by two orders of magnitude across metazoa (Baclaocos et al. 2019). Its

C-terminal tail is challenging to identify using genomic data due to its fast evolution, so

that here we mostly focused on gene-level detection. As expected, we identified two

phylogenetic clades of vertebrate genes, multi-Sec SELENOP/SELENOP1, and single-Sec

SELENOP2/SELENOPB. While the former is reportedly ubiquitous in vertebrate

genomes, the latter was lost in Eutheria. Moreover, SELENOP2 was converted to a Cys

homolog in Anura. Our present analysis (Figure 8a, Supplementary Figure S19) confirmed

these findings, and offered additional insights into early vertebrate evolution.

Invertebrate SELENOP homologs are consistently placed at the base of the gene tree,

supporting the hypothesis that the SELENOP1/SELENOP2 duplication arose within the

vertebrate lineage. Notably, Cyclostomata retain the two paralogs, yet both highly

divergent in sequence. One gene occupies a basal position in the tree even compared to

vertebrate outgroups, likely due to long-branch attraction artifacts; this is presumably

the SELENOP1 ortholog. The other Cyclostomata gene clusters within SELENOP2, and

displaying a unique CXXU motif (rather than the more typical UXXC) in its N-terminal

thioredoxin-like domain. Though SELENOP2 of other vertebrates carries a single Sec, in

Cyclostomata it exhibits an extraordinary expansion of Sec content: it harbors 162 and

139 in-frame UGA codons in *Lethenteron reissneri* (Far Eastern brook lamprey) and

Lampetra fluviatilis (European river lamprey), respectively (Supplementary Document

D1). Analogously to other metazoan multi-Sec SELENOP genes (Baclaocos et al. 2019),

the vast majority of these putative Sec residues are encoded within an expanded

modular C-terminal region, and two SECIS elements are detected downstream. Between

the SECISes, we noted the presence of a distinctive region rich in CA dinucleotides

(~1500-1800nt). Highly similar matches in TSA data of other Cyclostomata were

detected all across the putative transcript (Supplementary Document D1). With this

record, lampreys surpasses all other metazoans in the number of Sec-UGAs in SELENOP,

including the previous high of 132 reported in bivalves (Baclaocos et al. 2019).

<u>Other Salmoninae-specific duplications: SELENOE, SELENOF, SELENOH</u>

Our analysis revealed a substantial selenoproteome expansion specific to the lineage of

Salmoninae, which may suggest an increased functional reliance of this clade on

selenium. Besides the aforementioned expansions in the GPX, DIO, TXNRD, MSRB,

SELENOT, SELENOU and SELENOW families (Figure 1), we detected gene duplications

of SELENOE and SELENOF, yielding SELENOEb and SELENOFb, respectively, and another

two for SELENOH, yielding SELENOHb and SELENOHc (Figure 8, Supplementary Figures

S16, S17, S18). Briefly, SELENOF is a ER-resident selenoprotein involved in redox quality

control (Yim et al. 2018), recently pinpointed as tumor suppressor (Zigrossi et al. 2022;

Flowers et al. 2023). Our results show that SELENOF is ubiquitous to vertebrates. On the

other hand, SELENOE (formerly Fep15), a functionally uncharacterized selenoprotein

distantly related to SELENOF (Novoselov et al. 2006), was found as selenoprotein only

in fishes: Anura carry a Cys-homolog, while the rest of tetrapods (Amniota) lost the gene

altogether. Notably, we mapped the origin of SELENOE to the root of Gnathostomata,

since we did not find it in Cyclostomata nor in vertebrate outgroups. Finally, SELENOH

is a widespread nucleolar oxidoreductase that regulates redox homeostasis and

suppresses DNA damage through the p53 pathway, and it is implicated in cancer

proliferation (Wu et al. 2014; Cox et al. 2016; Bertz et al. 2018). Apart from the

Salmoninae-specific duplications, no other gene evolution event was detected for

SELENOE, SELENOF, and SELENOH, with the only exception of the loss of SELENOH in

Cyclostomata.

Notably, by inspecting the positions of introns relative to the alignment of protein

sequences (Figures 2, 4, 7), we noticed that they were conserved in all these Salmoninae-

specific gene duplications. This indicates a mechanism of DNA-based gene duplication

(as opposed to RNA-based transposition). The previously reported whole genome

duplication (WGD) at the root of salmonids (Lien et al. 2016) is therefore a most plausible

explanation for the origin of these genes. Importantly, this implies that we expect other

(non-Salmoninae) Salmonids to be also affected.

Figure 8. Phylogeny of SELENOP, SELENOE, SELENOOF, and SELENOH . a, b, c, d) *Gene tree visualizations of the* **SELENOP, SELENOE, SELENOOF, and SELENOH** *families, respectively. For tree annotation and legend details, see Figure 2b. Expanded gene trees are available: Supplementary Figure S19 (SELENOP), S16 (SELENOE), S17 (SELENOF), S18 (SELENOH).*

Other families: SEPHS2, SELENOL, SELENOK, SELENOM, SELENON, SELENOS

Hereafter, we briefly summarize the remaining vertebrate selenoproteome families, wherein our present analysis recapitulated earlier results without novel findings. As

already introduced, SEPHS2 (aka SPS2) is a selenoprotein whose function is critical for

Sec biosynthesis. As expected (Mariotti et al. 2012; Mariotti et al. 2015), we identified

SEPHS2 in all vertebrate lineages (barring its artefactual absence in many bird genome

assemblies, mentioned above). The SEPHS2 gene is intronless only in genomes of

Eutheria (placentals). As previously shown (Mariotti et al. 2012), the ancestral intron-

containing SEPHS2 gene (referred to as SEPHS2a) duplicated by retrotransposition at the

root of Mammalia, yielding an intron-less copy referred to as SEPHS2b. Both copies are

present in extant genomes of Marsupialia, while SEPHS2a was then lost in Eutheria. Our

results confirmed these findings (Supplementary Figures S1, S20). Thus, the sole

placental SEPHS2 gene is actually SEPHS2b. Of note, SEPHS2 duplicated at the root of

vertebrates, and independently also in various other metazoan lineages, to yield a non-

Sec/non-Cys paralog known as SEPHS1/SPS1 (Mariotti et al. 2015). The vertebrate

SEPHS1 gene carries threonine in place of Sec. This kind of conversion is highly unusual

for selenoproteins, since Sec is typically catalytic and it is highly dissimilar from canonical

amino acids other than Cys. Indeed, SEPHS1 appears to lack the enzymatic activity of

SEPHS2 (selenophosphate synthesis from selenide), and instead has a function in gene

expression-mediated control of redox homeostasis (Manta et al. 2022).

SELENOL is a peculiar selenoprotein featuring two nearby Sec residues that can form a

diselenide bond. It features a peroxiredoxin-like motive suggestive of a redox-related

function (Shchedrina et al. 2007). As expected (Mariotti et al. 2012), we detected

SELENOL in nearly all vertebrates except tetrapods, confirming a gene loss at the root of

this lineage (Supplementary Figure S1).

The remaining selenoprotein families were detected in a single Sec-containing copy in

all investigated genomes, barring pseudogenes and spurious absence attributed to

imperfect assembly quality. Curiously, all these single-copy selenoproteins localize to

the ER. They include: SELENOK, involved in calcium homeostasis and resolution of ER

stress (Odunsi et al. 2025); SELENOM and SELENOS, involved in unfolded protein

responses (Shi et al. 2024); and SELENON, involved in calcium and redox homeostasis

(Chernorudskiy et al. 2020) (Supplementary Figure S1).

DISCUSSION

The recent acceleration of genome sequencing and the rise of global biodiversity

initiatives has generated an unprecedented abundance of genomic data across

vertebrate lineages. This surge presents a unique opportunity to revisit fundamental

questions about gene evolution, functional diversity, and lineage-specific adaptations.

Here, we leveraged this genomic wealth to provide an updated view of the composition

and evolution of the vertebrate selenoproteome, obtained through analyses of hundreds

of genomes and meticulous phylogenetic reconstruction which greatly expanded upon

previous endeavors. We reinforced and refined existing findings, and uncovered new

genes and evolutionary events.

Our reconstruction of the ancestral vertebrate selenoproteome consists of 28

selenoprotein genes representing all the 19 known vertebrate families (Figure 1). Our

conclusions mostly overlap with our earlier analysis (Mariotti et al. 2012), with the

exceptions of SELENOE (now mapped to Gnathostomata rather than Vertebrata), and

GPX3b (Vertebrata rather than Actinopterygii). We also concluded that these

selenoproteins emerged specifically at the root of Vertebrata: GPX1, GPX3b, DIO2, DIO3,

SELENOI, SELENOP2. Among these, SELENOI stands out in that the whole family

appeared at the root of vertebrates: there is no homolog that includes the Sec position,

regardless of the aligned amino acid, outside of vertebrates.

On the other hand, our present work substantially expands the selenoproteome in fish

lineages. The conservation of intron structures and the phylogenetic correspondence

with documented whole-genome duplications suggests that most of these gene

duplications resulted from such events (Figure 1). This raises the question of whether

such selenoproteome expansions are adaptive, or are neutral consequences of having a

larger proteome. To gain insights in this matter, we compared the size of our predicted

selenoproteome with that of the annotated proteome for each species (Figure 9). As

expected, the selenoproteome-to-proteome ratio is indeed lower in mammals than fish.

Aquatic environments have long been hypothesized to favor selenoprotein usage, a

trend first suggested by Lobanov et al. from limited genomic data (Lobanov et al. 2007).

Subsequent analyses further corroborated this model by showing relaxed purifying

selection on selenoprotein genes in terrestrial vertebrates (Sarangi et al. 2018), alongside

reduced Sec content in SELENOP, an established proxy for selenium demand (Sarangi et

al. 2018; Baclaocos et al. 2019). Our present analysis with a larger dataset mostly support

this pattern. However, we note that the difference in selenoproteome-to-proteome

ratios between aquatic and terrestrial species is primarily driven by mammalian loss and

actinopterygian expansion. Some lineages diverge from this trend: aquatic Salmoninae

and Cyprinoidei show low ratios, despite having the biggest selenoproteomes across

vertebrates; and land-dwelling Sauropsida exhibit relatively high ratios (Figure 9).

Figure 9. Genomic correlation between the number of predicted selenoproteins and

annotated proteome size. *Each point represents a species and is color-labelled*

according to lineage.

The underlying cause of the selective advantage of Sec in aquatic environments is

arguably yet unclear. It has been proposed that land-dwelling organisms must face (i)

reduced selenium bioavailability and (ii) increased oxygen exposure (Lobanov et al.

2007; Sarangi et al. 2018). While the latter seems intuitively relevant, it contradicts one

of the prevalent proposed biochemical rationale for usage of selenium over cysteine

(Reich and Hondal 2016; Maroney and Hondal 2018): its resistance to (over)oxidation – a

trait that would suggest selenoproteins should be more beneficial in terrestrial

environments. Moreover, our data shows that selenoproteins belonging to antioxidant

families are more prevalent in fish compared to tetrapods. While further research is

required to fully resolve this question, we argue that the current evidence better

supports selenium bioavailability as the most likely evolutionary driver in aquatic vs land

environments.

These observations may depict our own selenoproteins as an unstable remnant of an

aquatic past. Yet, it is worth noting that (i) land-living tetrapods exhibit substantial

purifying selection, preventing Sec to Cys conversions in most genes (Castellano et al.

2009); and (ii) the aquatic selenoproteome is far from static, having expanded under

putative adaptive forces at the root of Actinopterygii and in multiple of its clades. We

found that these duplications occurred primarily in families linked to antioxidant defense,

either with well-established roles (GPX, MSRB) or with putative/indirect functions

(SELENOT, SELENOU, SELENOW). In contrast, no variation was observed in ER-resident

proteins with functions in calcium homeostasis (SELENOK, SELENON) or protein quality

control / unfolded protein response (SELENOM, SELENOS). While this trend was not

absolute – e.g. ER-resident SELENOE and SELENOF duplicated in Salmoninae, and the

frequently duplicated SELENOT is also ER-resident – it suggests that selenoproteome

expansions in Actinopterygii may have been driven primarily by selection for enhanced

antioxidant capacity.

Among tetrapods, our most striking finding is the recurrence of convergent evolutionary

events, including Sec-to-Cys conversions in SELENOU1 and GPX6, as well as

independent losses of SELENOV, all of which are supported by robust phylogenetic

evidence. It is important to note that, because our reconstruction is based on maximum

parsimony, our estimate of such events represents a conservative lower bound.

Interestingly, both SELENOV and GPX6 are the "youngest" mammalian selenoproteins

and they are specifically expressed in the testis, a tissue recognized for its role in the

evolution of new genes (Kaessmann 2010).

Our study has some limitations. First, we restricted our search to genes homologous to

known selenoprotein families, because existing methods for identifying novel

selenoproteins suffer from low specificity (Santesmasses et al. 2020). As a result, we

may have overlooked lineage-specific selenoproteins, particularly in fish lineages where

selenoproteome expansions are evident. Second, we could only detect events within

lineages represented in our dataset. Given the dynamic nature of the Actinopterygii

selenoproteome, additional genome sequences from this group will likely reveal further

duplications. Third, our analysis assumes the species tree as ground truth, which may

not be entirely accurate. For instance, the convergent loss of the same genes in

Tetraodontidae, Perciformes, Anabantaria, and *Cynoglossus semilaevis* (Figure 1) could

be explained as a single ancestral loss if non-Perciform Eupercaria and Carangaria were

instead outgroups of these clades – yet, we found no supporting evidence for this

alternative phylogeny in the literature (Hughes et al. 2018). Fourth, further data or

alternative methodological approaches could refine the evolutionary history of

selenoprotein families with the most complex gene dynamics, such as SELENOW and

SELENOU, and provide deeper insight into lineages with highly complex genome
evolution, such as Cyprininae.

Despite these limitations, we anticipate that our characterization of vertebrate
selenoproteins will largely hold up over time and provide a meaningful framework for
future evolutionary and functional research.

MATERIAL AND METHODS

Genome sequences and species tree

We downloaded all genome assemblies and annotations available in Ensembl version
v.108 (Cunningham et al. 2022) comprising 315 genomes, each corresponding to a
different species or strain. The phylogenetic tree of species was downloaded from
Ensembl Compara (Vilella et al. 2009). To refine our reconstruction of the ancestral
vertebrate selenoproteome, we also analyzed NCBI assemblies from species with
informative taxonomy. Specifically, we included various early-branching vertebrates:
nine cartilaginous fish (Chondrichthyes) and eight jawless vertebrates (Cyclostomata);
several chordates, i.e. close outgroups of vertebrates: two ascidians (Tunicata), four
amphioxiformes (Branchiostoma); and two remote outgroups: sea urchin
Strongylocentrotus purpuratus (Echinodermata, deuterostome) and centipede
Strigamia maritima (Arthropoda, protostome). These 25 genomes, together with the 315
Ensembl assemblies, constituted our main dataset. For follow up analyses to confirm or
dismiss gene evolution events, we then inspected gene predictions on an extended set
of 1,010 genome assemblies downloaded from NCBI, using the NCBI taxonomy tree as
the rough backbone for evolutionary interpretation.

Selenoprotein gene prediction

Selenoprotein gene prediction was performed using Selenoprofiles (Mariotti and Guigó 2010; Santesmasses et al. 2018; Ticó et al. 2024) version v4.5.0, available at https://github.com/marco-mariotti/selenoprofiles. Selenoprofiles is a computational pipeline to identify members of the known selenoprotein families and related proteins. It uses several homology-based gene prediction programs, and employs a set of manually curated multiple sequence alignments of selenoprotein families to scan genomes for homologues. The profiles we used are available at https://github.com/marco-mariotti/selenoprotein_profiles (v1.3.0). We scanned genome assemblies for all known selenoprotein families in metazoa. For most analyses, we focused on the predictions labelled as "selenocysteine", i.e., the genes encoding for selenoproteins, with no apparent pseudogene features (e.g. frameshifts).

Phylogenetic reconstruction of selenoprotein families

The protein sequences of all Selenoprofiles predictions across Ensembl genomes were collected using the Selenoprofiles *join* utility. These were used as input to phylogenetic reconstruction workflows, available in ETE3 v3.1.3 build framework (Huerta-Cepas et al. 2016). Phylogenetic trees were obtained with the "eggnog41" workflow (Powell et al. 2014), which internally uses PhyML v20160115.patched for phylogenetic reconstruction (Guindon et al. 2010). We visualized gene trees with the show_tree program in the biotree_tools package v0.0.6 (https://github.com/marco-mariotti/biotree_tools), based on the ETE3 python module (Huerta-Cepas et al. 2016), specifically adapted for custom visualization in this manuscript. The resulting vectorial images were further annotated manually using the Inkscape software. Sequence and tree manipulations throughout the project were carried out using the programs *alignment_tools* and *tree_tools* from the biotree_tools package. Protein sequence alignments were computed using ClustalO v.1.2.4 and processed with Pyaln v0.1.4 (https://github.com/marco-mariotti/pyaln).

Analysis of gene evolution events

We manually inspected gene and species trees to determine gene evolution events: losses, duplications, and Sec-to-Cys conversions. To detect potential duplications, gene trees were processed using the species overlap algorithm: the number of common species between the left and right branch of every ancestral node was used as a quantitative indicator of a gene duplication in the last common ancestor of those species. Putative duplication nodes were colored in the gene tree visualization, and their size was made proportional to this indicator, facilitating our manual review.

A visualization of the species tree together with all gene predictions in each genome was produced using the *tree drawer* utility available in Selenoprofiles, customized as follows. First, each gene prediction corresponding to a multi-member family were assigned to a mammalian-centric homology group, hereafter referred to as "subfamily" (e.g. GPX1, GPX2, GPX3, etc), using the *orthology* utility of Selenoprofiles. Then, we encoded the expected selenoproteome for every lineage based on literature; e.g. bats, since they are mammals with no further lineage-specific events reported, were expected to carry one Sec-encoding gene for each of GPX1, GPX2, GPX3, GPX4 and GPX6. In our custom *tree drawer* visualization (Supplementary Figure S1), we marked each gene prediction based on the deviation from such expectations; e.g., bats were labelled as missing a Sec-GPX6. This highlighted the presence of extra genes (putative duplications) and missing genes (putative losses). Ultimately, the joint analysis of gene and species trees led us to map each event into the species tree, following criteria of maximum parsimony and consistency with the known phylogenetic relationship among species.

Lastly, each event was critically evaluated to exclude potentially artifacts. We required support by at least two sister species for most events. Single-species events were deemed reliable only if supporting independent data (e.g. RNA sequences) existed. Due

to imperfect quality of genome assemblies, we sought confirmation of each individual

gene loss using independent sequence data from the lineage of interest, as follows. First,

we inspected Selenoprofiles predictions in an extended set of NCBI genomes. If the loss

was not dismissed, we then selected the protein sequence of the putatively lost gene

from a closely related organism, and searched it using TBLASTN at the NCBI web server

(Camacho et al. 2009) in all available RNA data, i.e. transcriptome shotgun assemblies

(TSAs), Expressed Sequence Tags (ESTs), and Reference RNA sequences (refseq_rna).

Grx-domain analysis in TxnRd genes

To investigate the distribution of glutaredoxin (Grx) domains in thioredoxin reductase

(TXNRD) genes, we first collected a set of representative TXNRD protein sequences and

scanned them for conserved domains using InterProScan (Jones et al. 2014). We

extracted the sequences containing the Grx domain and thus manually curated a high-

confidence alignment of Grx domains of TXNRDs. This was used as sequence profile to

search for homologous matches in Ensembl vertebrate genomes using Selenoprofiles.

Next, we cross-analyzed this data with those TXNRD gene predictions in the same

genomes classified as either TXNRD1 or TXNRD3 by the Selenoprofiles lineage utility. We

used a custom script based on pyranges v1 to assign Grx domain predictions to TXNRD1

and TXNRD3 genes, if located within their extended genomic coordinates (±5,000

nucleotides). Unassigned Grx domains were discarded. Finally, we visualized domain

architecture and evolutionary patterns using the ETE3 toolkit (Huerta-Cepas et al. 2016).

Gene nomenclature

We strived to assign gene names as consistent as possible with existing resources and

literature. To do so, we consulted the relevant publications on vertebrate selenoprotein

evolution as well as nomenclature (Regina et al. 2016), as well as current annotations at

Uniprot, Ensembl, and NCBI. The reference recommendation for gene names is to derive

them from the corresponding mammalian orthologs (Regina et al. 2016). The genes emerged from non-mammalian duplications —many of them previously unreported— were named by appending alphabetical or numerical suffixes to the source gene, e.g. GPX4b, SELENOW2. To enhance readability, suffixes are written in lowercase throughout this work. To avoid ambiguity, when creating new gene names we systematically skipped the suffixes "a" and "1" (in favor of "b" or "2"), because these are inconsistently used in the literature to refer to non-mammalian orthologs of single-copy mammalian genes; e.g. SELENOW is sometimes referred to as SELENOW1 in species that also possess the paralog SELENOW2.

DATA AVAILABILITY

The latest Selenoprofiles code is available at https://github.com/marco-mariotti/selenoprofiles4 and documented at https://selenoprofiles4.readthedocs.io/. Selenoprotein sequences used in this study split by family can be found in Supplementary Data D2.

SUPPLEMENTARY DATA

Figure S1. Selenoproteome evolution across vertebrate species. Each row represents a species and each column corresponds to a selenoprotein family. Rectangles represent individual "selenocysteine" Selenoprofiles predictions, color-labeled according to their gene annotation in Ensembl: pink for correctly annotated genes, blue for misannotated genes, grey for genes absent from the Ensembl annotation. Two graphical features are used to indicate the departure from the expected selenoproteome described in 2012 by Mariotti et al (coded in the Expectation_table.csv file at https://github.com/marco-mariotti/selenoprotein_profiles v.1.2.0): yellow represent genes expected in that lineage but undetected by Selenoprofiles; and red crosses mark extra genes compared to expectations, therefore filtered by the Selenoprofiles lineage tool.

Figure S2. Gene tree of the Glutathione peroxidase (GPX) family. *Subfamilies are color-*

labeled as follows: GPX4 in orange, GPX4b in yellow, GPX3b in purple, GPX3 in green,

GPX6 in pink, GPX5 in light purple, GPX2 in beige, GPX1 in brown, and GPX1b in blue.

Nodes highlighted in light blue indicate sequences from Salmoninae species. For a

detailed explanation of the tree layout and annotation features, refer to Figure 2.

Figure S3. Gene tree of the Iodothyronine deiodinase (DIO) family. *Subfamilies are*

color-labeled as follows: DIO1 in orange, DIO2 in yellow, DIO3 in green and DIO3b in

purple. See also Figures 2b (layout explanation) and 4 (compressed tree).

Figure S4. Iodothyronine deiodinase (DIO) evolution across Salmoninae species. *Each*

row represents a species and each column corresponds to a selenoprotein family.

Nodes represent individual Selenoprofiles predictions and are color-coded according

to their annotation in Ensembl: pink for well annotated Sec residue, blue for a

misannotated Sec residue and gray for missing annotations.

Figure S5. Gene tree of the Thioredoxin reductase (TXNRD) family. *Subfamilies are*

color-labeled as follows: TXNRD1 in green, TXNRD2 in orange, TXNRD3 in yellow.

Nodes highlighted in light blue indicate sequences from Salmoninae species. See also

Figures 2b (layout explanation) and 2f (compressed tree).

Figure S6. Vertebrate species tree representing the presence/absence of Grx domain in

TxnRd sequences. *Colors are assigned according to the TXNRD gene tree.*

Figure S7. Gene tree of the Selenoprotein J (SELENOJ) family. *Subfamilies are color-*

labeled as follows: SELENOJ in green and SELENOJ2 in yellow. Nodes highlighted in light

blue indicate sequences from Salmoninae species. See also Figures 2b (layout

explanation) and 6 (compressed tree).

Figure S8. Selenoprotein J (SELENOJ) evolution across Actinopterygii species. *Each row*

represents a species and each column corresponds to a selenoprotein family. Nodes

represent individual Selenoprofiles predictions and are color-coded according to their

amino acid content at the UGA position: green for selenocysteine-containing

sequences, red for cysteine-containing sequences, and grey for other classifications

inferred by Selenoprofiles.

Figure S9. Gene tree of the Selenoprotein O (SELENOO) family. *Subfamilies are color-*

labeled as follows: SELENOO in green and SELENOO2 in orange. Nodes highlighted in

brown and lightpurple indicate sequences from Tetraodontidae and Oryziinae species,

respectively. See also Figures 2b (layout explanation) and 6 (compressed tree).

Figure S10. Selenoprotein O (SELENOO) evolution across Actinopterygii species. *Each*

row represents a species and each column corresponds to a selenoprotein family.

Nodes represent individual Selenoprofiles predictions and are color-coded according

to their amino acid content at the UGA position: green for selenocysteine-containing

sequences, red for cysteine-containing sequences, and grey for other classifications

inferred by Selenoprofiles.

Figure S11. Gene tree of the Methionine sulfoxide reductase B (MSRB) family. *See also*

Figures 2b (layout explanation) and 6 (compressed tree).

Figure S12. Methionine sulfoxide reductase B (MSRB) evolution across Actinopterygii

species. *Each row represents a species and each column corresponds to a*

selenoprotein family. Nodes represent individual Selenoprofiles predictions and are

color-coded according to their amino acid content at the UGA position: green for

selenocysteine-containing sequences, red for cysteine-containing sequences, and grey

for other classifications inferred by Selenoprofiles.

Figure S13. Gene tree of the Selenoprotein T (SELENOT) family. *Subfamilies are color-*

labeled as follows: SELENOT1 in green and SELENOT2 in yellow. Nodes highlighted in

light blue indicate sequences from Salmoninae species, while lightpink and grey

indicate sequences from Cyprinoidei and Otophysi species, respectively. See also

Figures 2b (layout explanation) and 7 (compressed tree).

Figure S14. Gene tree of the Selenoprotein U (SELENOU) family. *Subfamilies are color-*

labeled as follows: SELENOU1a in yellow, SELENOU1a2 in orange, SELENOU1b in green,

SELENO1c in purple and Cysteine-homolog PRLX2A in beige. Nodes highlighted in light

blue indicate sequences from Salmoninae species, while lightpink indicate sequences

from Cyprinoidei. See also Figures 2b (layout explanation) and 7 (compressed tree).

Figure S15. Gene tree of the Selenoprotein V/W (SELENOV/W) family. *Subfamilies are*

color-labeled as follows: SELENOW1 from Actinopterygii species in green, SELENOW1

from tetrapoda species in beige, SELENOV in purple, SELENOW2a in yellow and

SELENOW2c in orange. Nodes highlighted in light blue indicate sequences from

Salmoninae species, while lightpink and lightgrey indicate sequences from Cyprinoidei

and Cyprinodontoidei species, respectively. See also Figures 2b (layout explanation)

and 7 (compressed tree).

Figure S16. Gene tree of the Selenoprotein E (SELENOE) family. *See also Figures 2b*

(layout explanation) and 8 (compressed tree).

Figure S17. Gene tree of the Selenoprotein F (SELENOF) family. *Nodes highlighted in light*

blue indicate sequences from Salmoninae species See also Figures 2b (layout

explanation) and 8 (compressed tree).

Figure S18. Gene tree of the Selenoprotein H (SELENOH) family. *Nodes highlighted in*

light blue indicate sequences from Salmoninae species. See also Figures 2b (layout

explanation) and 8 (compressed tree).

Figure S19. Gene tree of the Selenoprotein P (SELENOP) family. *Nodes highlighted in*

light blue indicate sequences from Salmoninae species. See also Figures 2b (layout

explanation) and 8 (compressed tree).

Figure S20. Gene tree of the SEPHS family. See also Figure 2b (layout explanation).

Table T1. Comparison of Glutathione Peroxidase (GPX) gene nomenclature between Xue et al. and this study. The table lists the gene names assigned to GPX family members in Xue et al. and the corresponding nomenclature used in this study (Ticó et al.). The third column provides relevant notes on differences in naming, such as updated classification, subfamily assignment, or gene duplication interpretations.

Document D1. Sequence analysis of SELENOP2 in lampreys. The document contains the sequences of SELENOP2 genes in two Cyclostomata, obtained via Selenoprofiles in genome data, then corroborated by RNA sequences. SECIS elements were predicted using SECISearch3.

Document D2. Protein sequences of all selenoproteins predicted and analyzed in this study. For details on file structure and content, please refer to the accompanying README file.

FUNDING

MM is funded by grants RYC2019-027746-I, PID2020-115122GA-I00, PID2023-147164NB-I00 funded by MICIU/AEI /10.13039/501100011033 and by "ESF Investing in your future", FEDER, UE.

BIBLIOGRAPHY

Anouar Y, Lihrmann I, Falluel-Morel A, Boukhzar L. 2018. Selenoprotein T is a key player in ER proteostasis, endocrine homeostasis and neuroprotection. *Free Radic Biol Med* 127:145–152.

Arnér ESJ. 2009. Focus on mammalian thioredoxin reductases--important selenoproteins with versatile functions. *Biochim Biophys Acta* [Internet] 1790:495–526. Available from: https://pubmed.ncbi.nlm.nih.gov/19364476/

Baclaocos J, Santesmasses D, Mariotti M, Bierła K, Vetick MB, Lynch S, McAllen R, Mackrill

JJ, Loughran G, Guigó R, et al. 2019. Processive Recoding and Metazoan Evolution of

Selenoprotein P: Up to 132 UGAs in Molluscs. *J Mol Biol* [Internet] 431:4381–4407.

Available from: https://pubmed.ncbi.nlm.nih.gov/31442478/

Bertz M, Kühn K, Koeberle SC, Müller MF, Hoelzer D, Thies K, Deubel S, Thierbach R, Kipp

AP. 2018. Selenoprotein H controls cell cycle progression and proliferation of human

colorectal cancer cells. *Free Radic Biol Med* [Internet] 127:98–107. Available from:

https://pubmed.ncbi.nlm.nih.gov/29330096/

Botero-Castro F, Figuet E, Tilak MK, Nabholz B, Galtier N. 2017. Avian Genomes Revisited:

Hidden Genes Uncovered and the Rates versus Traits Paradox in Birds. *Mol Biol Evol*

[Internet]   34:3123–3131.   Available   from:

https://pubmed.ncbi.nlm.nih.gov/28962031/

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.

BLAST+: architecture and applications. *BMC Bioinformatics* [Internet] 10. Available

from: https://pubmed.ncbi.nlm.nih.gov/20003500/

Castellano S, Andrés AM, Bosch E, Bayes M, Guigó R, Clark AG. 2009. Low

exchangeability of selenocysteine, the 21st amino acid, in vertebrate proteins. *Mol*

Biol   *Evol*   [Internet]   26:2031–2040.   Available   from:

https://pubmed.ncbi.nlm.nih.gov/19487332/

Castellano S, Lobanov A V., Chapple C, Novoselov S V., Albrecht M, Hua D, Lescure A,

Lengauer T, Krol A, Gladyshev VH, et al. 2005. Diversity and functional plasticity of

eukaryotic selenoproteins: Identification and characterization of the SelJ family. *Proc*

Natl   *Acad*   *Sci*   *U*   *S*   *A*   [Internet]   102:16188–16193.   Available   from:

https://www.pnas.org/doi/abs/10.1073/pnas.0505146102

Castellano S, Novoselov S V., Kryukov G V., Lescure A, Blanco E, Krol A, Gladyshev VN,

Guigó R. 2004. Reconsidering the evolution of eukaryotic selenoproteins: a novel

nonmammalian family with scattered phylogenetic distribution. *EMBO Rep*

[Internet] 5:71–77. Available from: https://pubmed.ncbi.nlm.nih.gov/14710190/

Chernorudskiy A, Varone E, Colombo SF, Fumagalli S, Cagnotto A, Cattaneo A, Briens M,

Baltzinger M, Kuhn L, Bachi A, et al. 2020. Selenoprotein N is an endoplasmic

reticulum calcium sensor that links luminal calcium levels to a redox activity. *Proc*

Natl Acad Sci U S A [Internet] 117:21288–21298. Available from:

https://pubmed.ncbi.nlm.nih.gov/32817544/

Copeland PR, Howard MT. 2021. Ribosome Fate during Decoding of UGA-Sec Codons.

Int J Mol Sci [Internet] 22. Available from:

https://pubmed.ncbi.nlm.nih.gov/34948001/

Cox AG, Tsomides A, Kim AJ, Saunders D, Hwang KL, Evason KJ, Heidel J, Brown KK, Yuan

M, Lien EC, et al. 2016. Selenoprotein H is an essential regulator of redox

homeostasis that cooperates with p53 in development and tumorigenesis. *Proc Natl*

Acad Sci U S A [Internet] 113:E5562–E5571. Available from:

https://pubmed.ncbi.nlm.nih.gov/27588899/

Cunningham F, Allen JE, Allen J, Alvarez-Jarreta J, Amode MR, Armean IM, Austine-

Orimoloye O, Azov AG, Barnes I, Bennett R, et al. 2022. Ensembl 2022. *Nucleic Acids*

Res [Internet] 50:D988–D995. Available from:

https://pubmed.ncbi.nlm.nih.gov/34791404/

Darras VM, Van Herck SLJ. 2012. Iodothyronine deiodinase structure and function: from

ascidians to humans. *J Endocrinol* [Internet] 215:189–206. Available from:

https://pubmed.ncbi.nlm.nih.gov/22825922/

Davy T, Castellano S. 2018. The genomics of selenium: Its past, present and future.
Biochim Biophys Acta Gen Subj [Internet] 1862:2427–2432. Available from:
https://pubmed.ncbi.nlm.nih.gov/29859258/

Fairweather-Tait SJ, Bao Y, Broadley MR, Collings R, Ford D, Hesketh JE, Hurst R. 2011.
Selenium in human health and disease. *https://home.liebertpub.com/ars* [Internet]
14:1337–1383.   Available   from:
https://www.liebertpub.com/doi/10.1089/ars.2010.3275

Fata F, Gencheva R, Cheng Q, Lullo R, Ardini M, Silvestri I, Gabriele F, Ippoliti R, Bulman
CA, Sakanari JA, et al. 2022. Biochemical and structural characterizations of
thioredoxin reductase selenoproteins of the parasitic filarial nematodes Brugia
malayi and Onchocerca volvulus. *Redox Biol* [Internet] 51. Available from:
https://pubmed.ncbi.nlm.nih.gov/35276442/

Flohé L. 2009. The labour pains of biochemical selenology: the history of selenoprotein
biosynthesis. *Biochim Biophys Acta* 1790:1389–1403.

Flowers B, Bochnacka O, Poles A, Diamond AM, Kastrati I. 2023. Distinct Roles of
SELENOF in Different Human Cancers. *Biomolecules* [Internet] 13. Available from:
https://pubmed.ncbi.nlm.nih.gov/36979420/

Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New
Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing
the Performance of PhyML 3.0. *Syst Biol* [Internet] 59:307–321. Available from:
https://dx.doi.org/10.1093/sysbio/syq010

Howard MT, Copeland PR. 2019. New Directions for Understanding the Codon
Redefinition Required for Selenocysteine Incorporation. *Biol Trace Elem Res*
[Internet] 192:18–25. Available from: https://pubmed.ncbi.nlm.nih.gov/31342342/

Huerta-Cepas J, Serra F, Bork P. 2016. ETE 3: Reconstruction, Analysis, and Visualization

of Phylogenomic Data. *Mol Biol Evol* [Internet] 33:1635. Available from:

https://pmc.ncbi.nlm.nih.gov/articles/PMC4868116/

Hughes LC, Ortí G, Huang Y, Sun Y, Baldwin CC, Thompson AW, Arcila D, Betancur R, Li

C, Becker L, et al. 2018. Comprehensive phylogeny of ray-finned fishes

(Actinopterygii) based on transcriptomic and genomic data. *Proc Natl Acad Sci U S*

A [Internet] 115:6249–6254. Available from:

https://www.pnas.org/doi/abs/10.1073/pnas.1719358115

Jeong D won, Kim TS, Chung YW, Lee BJ, Kim IY. 2002. Selenoprotein W is a glutathione-

dependent antioxidant in vivo. *FEBS Lett* [Internet] 517:225–228. Available from:

https://pubmed.ncbi.nlm.nih.gov/12062442/

Jiang L, Liu Q, Ni J. 2010. In silico identification of the sea squirt selenoproteome. *BMC*

Genomics [Internet] 11:289. Available from: http://www.biomedcentral.com/1471-

2164/11/289

Jiang L, Ni J, Liu Q. 2012. Evolution of selenoproteins in the metazoan. *BMC Genomics*

[Internet] 13:446. Available from:

http://bmcgenomics.biomedcentral.com/articles/10.1186/1471-2164-13-446

Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell

A, Nuka G, et al. 2014. InterProScan 5: genome-scale protein function classification.

Bioinformatics [Internet] 30:1236–1240. Available from:

https://dx.doi.org/10.1093/bioinformatics/btu031

Kaessmann H. 2010. Origins, evolution, and phenotypic impact of new genes. *Genome*

Res [Internet] 20:1313–1326. Available from:

https://pubmed.ncbi.nlm.nih.gov/20651121/

Kryukov G V., Castellano S, Novoselov S V., Lobanov A V., Zehtab O, Guigó R, Gladyshev
VN. 2003. Characterization of mammalian selenoproteomes. *Science* [Internet]
300:1439–1443. Available from: https://pubmed.ncbi.nlm.nih.gov/12775843/

Labunskyy VM, Hatfield DL, Gladyshev VN. 2014. Selenoproteins: Molecular Pathways
and Physiological Roles. *Physiol Rev* [Internet] 94:739. Available from:
https://pmc.ncbi.nlm.nih.gov/articles/PMC4101630/

Lewin HA, Robinson GE, Kress WJ, Baker WJ, Coddington J, Crandall KA, Durbin R,
Edwards S V., Forest F, Gilbert MTP, et al. 2018. Earth BioGenome Project:
Sequencing life for the future of life. *Proc Natl Acad Sci U S A* [Internet] 115:4325–
4333. Available from: https://pubmed.ncbi.nlm.nih.gov/29686065/

Lien S, Koop BF, Sandve SR, Miller JR, Kent MP, Nome T, Hvidsten TR, Leong JS, Minkley
DR, Zimin A, et al. 2016. The Atlantic salmon genome provides insights into
rediploidization. *Nature 2016 533:7602* [Internet] 533:200–205. Available from:
https://www.nature.com/articles/nature17164

Lobanov A V., Fomenko DE, Zhang Y, Sengupta A, Hatfield DL, Gladyshev VN. 2007.
Evolutionary dynamics of eukaryotic selenoproteomes: Large selenoproteomes
may associate with aquatic life and small with terrestrial life. *Genome Biol* [Internet]
8:1–16.   Available   from:
https://genomebiology.biomedcentral.com/articles/10.1186/gb-2007-8-9-r198

Lorgen M, Casadei E, Król E, Douglas A, Birnie MJ, Ebbesson LOE, Nilsen TO, Jordan WC,
Jørgensen EH, Dardente H, et al. 2015. Functional divergence of type 2 deiodinase
paralogs in the Atlantic salmon. *Curr Biol* [Internet] 25:936–941. Available from:
https://pubmed.ncbi.nlm.nih.gov/25802152/

Manta B, Makarova NE, Mariotti M. 2022. The selenophosphate synthetase family: A
review. *Free Radic Biol Med* 192:63–76.

Mariotti M, Gladyshev VN. 2022. Selenocysteine-containing proteins. *Redox Chemistry*

and Biology of Thiols:405–421.

Mariotti M, Guigó R. 2010. Selenoprofiles: profile-based scanning of eukaryotic genome

sequences for selenoprotein genes. *Bioinformatics* [Internet] 26:2656–2663.

Available from: https://dx.doi.org/10.1093/bioinformatics/btq516

Mariotti M, Ridge PG, Zhang Y, Lobanov A V., Pringle TH, Guigo R, Hatfield DL, Gladyshev

VN. 2012. Composition and evolution of the vertebrate and mammalian

selenoproteomes. *PLoS One* [Internet] 7. Available from:

https://pubmed.ncbi.nlm.nih.gov/22479358/

Mariotti M, Santesmasses D, Capella-Gutierrez S, Mateo A, Arnan C, Johnson R, D'Aniello

S, Yim SH, Gladyshev VN, Serras F, et al. 2015. Evolution of selenophosphate

synthetases: emergence and relocation of function through independent

duplications and recurrent subfunctionalization. *Genome Res* [Internet] 25:1256–

1267. Available from: http://genome.cshlp.org/content/25/9/1256.full

Maroney MJ, Hondal RJ. 2018. Selenium versus sulfur: Reversibility of chemical reactions

and resistance to permanent oxidation in proteins and nucleic acids. *Free Radic Biol*

Med [Internet] 127:228–237. Available from:

https://pubmed.ncbi.nlm.nih.gov/29588180/

Mc Cartney AM, Formenti G, Mouton A, De Panis D, Marins LS, Leitão HG, Diedericks G,

Kirangwa J, Morselli M, Salces-Ortiz J, et al. 2024. The European Reference Genome

Atlas: piloting a decentralised approach to equitable biodiversity genomics. *npj*

Biodiversity 2024 3:1 [Internet] 3:1–17. Available from:

https://www.nature.com/articles/s44185-024-00054-6

Melo LMN, Sabatier M, Ramesh V, Szylo KJ, Fraser CS, Pon A, Mitchell EC, Servage KA,

Allies G, Westedt I V., et al. 2025. Selenoprotein O Promotes Melanoma Metastasis

and Regulates Mitochondrial Complex II Activity. *Cancer Res* [Internet] 85:942–955.

Available from: https://pubmed.ncbi.nlm.nih.gov/39700395/

Misra S, Lee TJ, Sebastian A, McGuigan J, Liao C, Koo I, Patterson AD, Rossi RM, Hall MA,

Albert I, et al. 2023. Loss of selenoprotein W in murine macrophages alters the

hierarchy of selenoprotein expression, redox tone, and mitochondrial functions

during inflammation. *Redox Biol* [Internet] 59. Available from:

https://pubmed.ncbi.nlm.nih.gov/36516721/

Novoselov S V., Hua D, Lobanov A V., Gladyshev VN. 2006. Identification and

characterization of Fep15, a new selenocysteine-containing member of the Sep15

protein family. *Biochem J* [Internet] 394:575–579. Available from:

https://pubmed.ncbi.nlm.nih.gov/16236027/

Odunsi A, Kapitonova MA, Woodward G, Rahmani E, Ghelichkhani F, Liu J, Rozovsky S.

2025. Selenoprotein K at the intersection of cellular pathways. *Arch Biochem*

Biophys [Internet] 764. Available from: https://pubmed.ncbi.nlm.nih.gov/39571956/

Powell S, Forslund K, Szklarczyk D, Trachana K, Roth A, Huerta-Cepas J, Gabaldón T,

Rattei T, Creevey C, Kuhn M, et al. 2014. eggNOG v4.0: nested orthology inference

across 3686 organisms. *Nucleic Acids Res* [Internet] 42:D231–D239. Available from:

https://dx.doi.org/10.1093/nar/gkt1253

Regina BF, Gladyshev VN, Arnér ES, Berry MJ, Bruford EA, Burk RF, Carlson BA, Castellano

S, Chavatte L, Conrad M, et al. 2016. Selenoprotein gene nomenclature. *Journal of*

Biological Chemistry [Internet] 291:24036–24040. Available from:

https://www.jbc.org/action/showFullText?pii=S0021925820346731

Reich HJ, Hondal RJ. 2016. Why Nature Chose Selenium. *ACS Chem Biol* [Internet] 11:821–

841. Available from: https://pubmed.ncbi.nlm.nih.gov/26949981/

Santesmasses D, Mariotti M, Gladyshev VN. 2020. Bioinformatics of Selenoproteins.

Antioxid Redox Signal [Internet] 33:525–536. Available from:

https://pubmed.ncbi.nlm.nih.gov/32031018/

Santesmasses D, Mariotti M, Guigó R. 2018. Selenoprofiles: A Computational Pipeline for

Annotation of Selenoproteins. *Methods in Molecular Biology* [Internet] 1661:17–28.

Available from: https://link.springer.com/protocol/10.1007/978-1-4939-7258-6_2

Sarangi GK, Romagne F, Castellano S. 2018. Distinct Patterns of Selection in Selenium-

Dependent Genes between Land and Aquatic Vertebrates. *Mol Biol Evol* [Internet]

35:1744–1756. Available from: https://dx.doi.org/10.1093/molbev/msy070

Shchedrina VA, Novoselov S V., Malinouski MY, Gladyshev VN. 2007. Identification and

characterization of a selenoprotein family containing a diselenide bond in a redox

motif. *Proc Natl Acad Sci U S A* [Internet] 104:13919–13924. Available from:

https://www.pnas.org/doi/abs/10.1073/pnas.0703448104

Shi Z, Han Z, Chen J, Zhou JC. 2024. Endoplasmic reticulum-resident selenoproteins and

their roles in glucose and lipid metabolic disorders. *Biochim Biophys Acta Mol Basis*

Dis [Internet] 1870. Available from: https://pubmed.ncbi.nlm.nih.gov/38763408/

Sreelatha A, Yee SS, Lopez VA, Park BC, Kinch LN, Pilch S, Servage KA, Zhang J, Jiou J,

Karasiewicz-Urbańska M, et al. 2018. Protein AMPylation by an Evolutionarily

Conserved Pseudokinase. *Cell* [Internet] 175:809-821.e19. Available from:

https://pubmed.ncbi.nlm.nih.gov/30270044/

Tarrago L, Kaya A, Kim HY, Manta B, Lee BC, Gladyshev VN. 2022. The selenoprotein

methionine sulfoxide reductase B1 (MSRB1). *Free Radic Biol Med* [Internet] 191:228–

240. Available from: https://pubmed.ncbi.nlm.nih.gov/36084791/

Ticó M, Sullivan E, Guigó R, Mariotti M. 2024. Overcoming the widespread flaws in the

annotation of vertebrate selenoprotein genes in public databases. *bioRxiv*

[Internet]:2024.10.30.620813.   Available   from:

https://doi.org/10.1101/2024.10.30.620813

Toppo S, Vanin S, Bosello V, Tosatto SCE. 2008. Evolutionary and structural insights into

the multifaceted glutathione peroxidase (Gpx) superfamily. *Antioxid Redox Signal*

[Internet]   10:1501–1513.   Available   from:

https://pubmed.ncbi.nlm.nih.gov/18498225/

Vilella AJ, Severin J, Ureta-Vidal A, Heng L, Durbin R, Birney E. 2009. EnsemblCompara

GeneTrees: Complete, duplication-aware phylogenetic trees in vertebrates.

Genome   Res   [Internet]   19:327–335.   Available   from:

http://genome.cshlp.org/content/19/2/327.full

Wu RTY, Cao L, Chen BPC, Cheng WH. 2014. Selenoprotein H suppresses cellular

senescence through genome maintenance and redox regulation. *J Biol Chem*

[Internet]   289:34378–34388.   Available   from:

https://pubmed.ncbi.nlm.nih.gov/25336634/

Xue Y, Chen L, Li B, Xiao J, Wang H, Dong C, Li X, Xu P. 2022. Genome-wide mining of

gpx gene family provides new insights into cadmium stress responses in common

carp (Cyprinus carpio). *Gene* 821:146291.

Yim SH, Everley RA, Schildberg FA, Lee SG, Orsi A, Barbati ZR, Karatepe K, Fomenko DE,

Tsuji PA, Luo HR, et al. 2018. Role of Selenof as a Gatekeeper of Secreted Disulfide-

Rich Glycoproteins. *Cell Rep* [Internet] 23:1387–1398. Available from:

https://pubmed.ncbi.nlm.nih.gov/29719252/

Zhang X, Xiong W, Chen LL, Huang JQ, Lei XG. 2020. Selenoprotein V protects against

endoplasmic reticulum stress and oxidative injury induced by pro-oxidants. *Free*

Radic   Biol   Med [Internet] 160:670–679. Available from:

https://pubmed.ncbi.nlm.nih.gov/32846216/

Zhang Y, Jin J, Huang B, Ying H, He J, Jiang L. 2022. Selenium Metabolism and

Selenoproteins in Prokaryotes: A Bioinformatics Perspective. *Biomolecules 2022,*

Vol. 12, Page 917 [Internet] 12:917. Available from: https://www.mdpi.com/2218-

273X/12/7/917/htm

Zigrossi A, Hong LK, Ekyalongo RC, Cruz-Alvarez C, Gornick E, Diamond AM, Kastrati I.

2022. SELENOF is a new tumor suppressor in breast cancer. *Oncogene 2021 41:9*

[Internet] 41:1263–1268. Available from: https://www.nature.com/articles/s41388-

021-02158-w

1066

1067