1 Title: Genome sequencing confirms cryptic diversity and potentially adaptive changes in gene
2 content of Lepidoptera-infecting Apicomplexa
3

4 Andrew J. Mongue[1], Peter Kuperus[2], Astrid T. Groot[2]
5
6 [1]Department of Entomology and Nematology, University of Florida, USA
7 [2]Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Netherlands

8
**Abstract**

Apicomplexan parasites of insects represent a largely unexplored branch of the tree of life. Their
small size and lack of informative phenotypic traits can obscure their diversity and many
lineage-specific adaptations to different host species. To better understand how these parasites
have evolved, we have sequenced and annotated the genome of a neogregarine parasite of
moths including *Helicoverpa armigera*. This parasite, which we term *Helicoverpa*-infecting
*Ophryocystis*, has a tiny 7 Megabase genome with only 2,200 protein-coding genes. Using
phylogenomic methods, we place it as the closest currently sequenced relative to *Ophryocystis*
*elektroscirrha* and its sister lineage, both of which infect milkweed butterflies. We then explore
evolution of the ATPase gene family, which is known to be reduced in other *Ophryocystis*
lineages. *Helicoverpa*-infecting *Ophryocystis* has more ATPase genes than other *Ophryocystis*,
which may relate to the interactions of parasites with phytochemistry encountered in their insect
hosts. These results demonstrate both the amount of uncertainty remaining in the insect-
infecting Apicomplexa as well as the utility of applying modern genomic approaches to their
study.

24
**Introduction**
Apicomplexa are ancient and diverse unicellular eukaryotic parasites (Levine, 2018). Members
of the phylum infect most multicellular animals, including both vertebrates like humans,
(Abrahamsen et al., 2004; Gardner et al., 2002; Levine, 2018) and invertebrates (Gao et al.,
2020; McLaughlin & Myers, 1970; Templeton et al., 2010). This evolutionary longevity and host
breadth makes them intriguing targets for evolutionary genetic study: what molecular and
genomic features are required to exploit so many hosts across the tree of life?
32
Some species, such as the causal parasite of malaria, *Plasmodium*, have understandably
received intensive study (Bushman et al., 2016; Gardner et al., 2002; Krishna et al., 2001).
However, much of the diversity remains largely unexplored, especially that of invertebrate-
infecting Apicomplexa (Boisard & Florent, 2020). Better understanding of these parasites is
important both to make sense of the evolution and diversity of life at a basic level (Boisard &
Florent, 2020; DeBarry & Kissinger, 2011) and to develop potential biological controls of pest
arthropods (e.g., as control agents of mosquitos, Tseng, 2007). The first step towards these two
broader goals is to gain a better molecular understanding of invertebrate-infecting Apicomplexa
that are already being used as model systems in the laboratory.
42
Recent sequencing of two *Ophryocystis* lineages (Mongue et al., 2023) that parasitize milkweed
butterflies (Nymphalidae Danainae) has already helped to shed light on the host-parasite

interactions of this system. In particular, larvae of butterflies in the genus *Danaus* are specialists
on milkweed plants in the genus *Asclepias*, which contain toxic cardiac glycoside
phytochemicals (Agrawal et al., 2012; Brower & Fink, 1985). These compounds block the
function of sodium/potassium pump ATPases in most organisms (Matsui & Schwartz, 1968), but
monarchs possess amino acid adaptations that make them resistant to cardiac glycosides
(Aardema & Andolfatto, 2016). Given the toxic effects of milkweed phytochemicals on ATPase
function generally (Kelly et al., 1985; Matsui & Schwartz, 1968), and the demonstrated inhibitory
effect of milkweed chemistry on apicomplexan parasite growth (de Roode et al., 2008; De
Roode et al., 2011), an intuitive hypothesis was that the milkweed phytochemicals would inhibit
ATPase. However, in sequencing the genome of *Ophryocystis*, the authors found that these
parasites lack sodium pump ATPases entirely (Mongue et al., 2023). This unexpected finding
led to the hypothesis that exposure to milkweed phytochemicals could lead to the loss of
ATPases, although it was difficult to draw conclusions about parasite adaptation with only a pair
of sequences from milkweed butterfly parasites and no relatives to act as comparisons.
59
Here we report the genome sequence of another *Ophryocystis* parasite of moths in the genus
*Helicoverpa* (Lepidoptera: Noctuidae). This parasite has been described as *Ophryocystis*
*elektroscirrha*-like (Gao et al., 2020) and is visually nearly identical to *O. elektroscirrha* in the
most visible oocyst stage. However, unlike *O. elektroscirrha's* host, *Helicoverpa* moths are
generalist crop pests whose diet does not include cardiac glycoside producing plants (Nasreen
& Mustafa, 2000; Zalucki et al., 1986). As such, their parasites should not be exposed to the
selective pressures of ATPase inhibiting phytochemicals. The *Ophryocystis* parasite of
*Helicoverpa* thus makes a useful outgroup to understand the genetics of ATPases and how they
compare to *O. elektroscirrha*. To explore the evolutionary genetics of ATPases, as well as to
generate resources for another insect-infecting apicomplexan model, we present the genome of
a previously unsequenced *Ophryocystis* species and explore the molecular evolution of
ATPases.
72
73  **Methods**
74
*Sample collection and DNA extraction*
Oocyst purification was performed on *Chloridea virescens* moths that had been infected with
*Ophryocystis elektroscirrha*-like parasite, collected originally from *Helicoverpa armigera* (see
Gao et al. 2020). To remove debris and isolate oocyst, wings were removed from dead, infected
moths. Groups of 10 moths were pooled into 50 mL tubes (eight sets, totalling 80 moths). Each
tube was supplemented with 40 mL sterile water and 1 mL Dreft detergent solution and
subjected to two rounds of 10-minute sonication. Samples were then manually shaken for 10
minutes, minimizing foam formation. Large insect fragments were removed before the tube was
centrifuged at 1,000 rpm for 2 minutes in a fixed-angle rotor, and oocysts were collected from
the black sediment layer. The oocyst-rich black layer (Figure 1, left) was collected into a 2 mL
tube, washed twice with 1 mL water/Dreft solution, and vortexed before centrifuged at 10,000
rpm for 4 minutes. The black oocyst pellet was retained, while the brown layer of moth scales
was removed by pipetting. Oocysts were air-dried and resuspended in 80 µL TE buffer. Oocyst
integrity and purity were assessed microscopically (~2 µL per sample, Figure 1, middle).

DNA extraction was performed using a modified CTAB protocol (Doyle and Doyle, 1987). First,

a total of 10 to 20 zirconium beads (1 mm) were added to the spore suspension, followed by

homogenization using a Precellys system (6,800 rpm). Oocyst disruption was confirmed

microscopically (Figure 1, right).

**Figure 1. Oocyst purification for DNA extraction. Left:** Oocyst-rich suspension collected from *C. virescens* adults. **Middle:** Cleaned oocyst suspension. **Right:** Disrupted oocysts for DNA extraction.

For DNA extraction, 600 µL of 2% CTAB solution

and 6 µL Proteinase K (Thermo Scientific, 20 mg/mL) were added, followed by incubation at

56°C for 1 hour. DNA was purified by adding chloroform:isoamyl alcohol (24:1), vortexing, and

centrifuging at 14,000 rpm for 8 minutes. The upper aqueous phase transferred to a new tube,

and the extraction was repeated. DNA was precipitated with 96% ethanol and incubated at -

20°C for 30 minutes, followed by centrifugation at 14,000 rpm for 10 minutes at 10°C. The DNA

pellet was washed with 80% ethanol, centrifuged under the same conditions, and resuspended

in 50 µL TE buffer containing 2 µL 5M NaCl and 2 µL RNase (1 mg/mL). Samples were

incubated at 37°C for 30 minutes, after which DNA was re-precipitated using 150 µL 96%

ethanol, incubated at -20°C for 30 minutes, and centrifuged again. The pellet was washed with

200 µL 80% ethanol, air-dried, and resuspended in 25 µL sterile water. DNA concentration and

purity were assessed using a Implen nanophotometer (N60 Touch), yielding 8.9 ng/µL (260/280

= 1.9; 260/230 = 1.6).

*Sequencing and genome assembly*

Library preparation started with 204 ng of DNA and followed the Oxford Nanopore genomic-dna-

by-ligation-sqk-lsk110 protocol. After the final adapter ligation was performed, DNA

concentration was measured on a nanophotometer (2.6 ng/µL; 260/280 = 1.3; 260/230 = 0.4).

Sequencing was performed on an Oxford Nanopore Flongle flow cell (FLO-FLG001) loaded with

13 ng library prepped DNA and run for 24 hours.

A minimum quality filter was implemented to remove low quality (<Q10) reads from the

generated data. Genome assembly was performed using Geneious Prime (version 2023.0.4)

(https://www.geneious.com/prime/) with the Flye and Minimap2 plugins. The flye assembler was

run in –nano-raw mode with an estimated genome size parameter of 10m, based on results

from existing gregarine genomes.

*Phylogenomics*

To assess evolutionary relationships, a phylogenomics pipeline that was previously successfully implemented in insects by Mongue et al. (2024) was employed for Apicomplexa. In brief, the genome assemblies listed in **Table 1** were obtained from online databases and BUSCO v4.14 (Manni et al., 2021) was run in genome mode using the apicomplexa_odb10 dataset to identify single copy orthologs in each species. Next, scripts from the BUSCO_phylogenomics git repo (https://github.com/jamiemcg/BUSCO_phylogenomics) were used to parse complete single copy orthologs found in at least 70% of the sampled species. Subsequently, FastTree v2.1.11 (Price et al., 2009) was used to construct gene trees for each orthologous protein with the JTT model of molecular evolution (Jones et al., 1992) and the CAT approximation for variable rates at each site (Stamatakis, 2006). Finally, ASTRAL (Zhang et al., 2018) was used to generate a single consensus species tree with local quartet posterior probabilities for support (Sayyari & Mirarab, 2016).The tree was manually rooted with a dinoflagellate, *Symbiodinium microadriaticum*, as the outgroup.

| Species | Accession | Reference |
|---|---|---|
| *Ascogregarina taiwanensis* | PRJNA27765 | (Templeton et al., 2010) |
| *Cryptosporidium meleagridis* | PRJNA1022047 | (Penumarthi et al., 2024) |
| *Cryptosporidium parvum* | PRJNA144 | (Abrahamsen et al., 2004) |
| *Eimeria praecox* | PRJEB71489 | (Blake et al., 2024) |
| *Gregarina niphandrodes* | PRJNA259233 | Unpublished |
| *Ophryocystis elektroscirrha* | PRJNA906508 | (Mongue et al., 2023) |
| *Ophryocystis elektroscirrha*-like | https://doi.org/10.5281/zenodo.7817702 | (Mongue et al., 2023) |
| *Helicoverpa*-infecting *Ophryocystis* | [TBD] | This publication |
| *Plasmodium falciparum* | PRJNA13173 | (Gardner et al., 2002) |
| *Plasmodium yoelii* | PRJEB5706 | (Otto et al., 2014) |
| *Porospora gigantea* A | PRJNA734792 | (Boisard & Florent, 2020) |
| *Porospora gigantea* B | PRJNA734792 | (Boisard & Florent, 2020) |
| *Sarcocystis neurona* | PRJNA227351 | (Dangoudoubiyam et al., 2024) |
| *Symbodinium microadriaticum* | PRJNA292355 | (Aranda et al., 2016) |
| *Theileria orientalis* | PRJDA49463 | (Hayashida et al., 2012) |

**Table 1. Genomic resources for phylogenomic analyses.** For each species we list the data accession and publication, where available.

*Gene annotation*

A gene annotation method was chosen that has proven successful for other *Ophryocystis*
genomes (Mongue et al., 2023). Specifically, the GenSAS pipeline (Humann et al., 2019) was
used to generate gene models as follows. To start, repetitive sequences were identified with
RepeatMasker v4.1.1 (Smit et al., 2019) and masked with RepeatModeler v2.0.1 (Flynn et al.,
2020) and in GenSAS (Humann et al., 2019). Based on previous success, GeneMarkES v4.48
(Ter-Hovhannisyan et al., 2008) was used to annotate genes in the repeat-masked assembly,
then BUSCO v5.2.2 (Manni et al., 2021) was used to query the annotation with the
apicomplexa_odb10 dataset of putatively conserved orthologs.
163
*General orthology analysis*
With the sequencing and annotation of this parasite, there are now three genomes and
annotations from lineages in Ophyrocystidae, enough information to begin exploring overall
patterns of gene conservation in this family. To do so, ProteinOrtho v5.11 (Lechner et al., 2011)
was used to search for orthologs between *Helicoverpa*-infecting *Ophryocystis* and the two
existing *Ophryocystis* lineages from milkweed butterflies. Basic summary statistics of gene
conservation are presented in the results.
171
*ATPase gene family analysis*
Because the *Helicoverpa* parasite represents a closer relative to other *Ophryocystis* lineages
than other sequenced Apicomplexa and its host does not associate with cardiac glycosides, the
ATPase genes from this genome serve as interesting negative control contrasts to those
described from *O. elektroscirrha* and its sister lineage (Mongue et al., 2023). To investigate the
ATPase genes, we followed our previous procedure (Mongue et al. (2023) by querying the new
gene annotation with the amino acid sequence of a known ATPase gene (GenBank accession:
CCJ05450.1) using BLAST+ v2.9.0 (Camacho et al., 2009). ATPase function was confirmed via
reciprocal BLAST on NCBI. Once confirmed, these new sequences were added to a database
of primarily apicomplexan ATPases that was initially compiled by Lehane et al. (2019) and
further developed by Mongue et al. (2023). These sequences were aligned with CLUSTAL
Omega's web-hosted tool (Sievers & Higgins, 2014) and manually trimmed the output to ~260
conserved amino acids. To generate a phylogenetic tree, this trimmed alignment was passed to
the *phangorn* package in R (Schliep, 2011) using the LG+G(4)+I model of substitution and 100
bootstrap replicates to give confidence. With this final phylogenetic tree, the function of newly
identified ATPases was inferred based on sequence similarity to those with existing functional
annotations.
189
190  **Results**
191
192  *Genome assembly*
193  We generated 249,580 Oxford Nanopore reads with an N50 of ~2,100 Kb, totaling 304.9 Mb of
194  sequence data. Of these, 172,880 passed base-calling quality filter (>Q10) and were used for
195  downstream assembly, totaling 172.9 Mb. Primary assembly stats are found in **Table 2**. Of note,
196  only two contigs had lengths <1,000 bp.
197

| Assembly Statistics | Assembly (flye) |
|---|---|
| Total length | 7,447,837 bp |
| Sequence count | 191 |
| N50 | 63,090 bp |
| L50 | 35 |
| GC content | 31.2% |
| BUSCO completeness (apicomplexa_odb10) | C:33.8%[S:33.6%, D:0.2%],F:9.9%,M: 56.3% |

**Table 2. Genome assembly statistics.**

*Phylogenomics*

**Figure 1. Phylogenetic placement of *Helicoverpa* infecting *Ophryocystis*.** *Ophryocystis*
parasites of *H. armigera* are sister to the lineages of *Ophryocystis* found in milkweed butterflies.
Node labels show posterior quartet support values.

As expected, but now confirmed via genomic analysis, the *Ophryocystis* parasite of *H. armigera*
is genetically distinct from *O. elektroscirrha* that infects *D. plexippus,* and in fact represents the
outgroup to the two lineages of milkweed butterfly infecting *Ophryocystis* (**Figure 1**).

211 *Ascogregrina taiwnensis*, a parasite of mosquitos (Templeton et al., 2010), is the closest
212 sequenced relative of *Ophryocystis* as currently defined. More generally, all of these genomes
213 fall into a monophyletic clade with other species in the subclass Gregarinasina, all of which are
214 parasites of arthropods.
215
*Repeat and gene content*
As expected of a very small genome, we recovered a very low repeat content (9.74%), which
lower even that the ~12% seen in *O. elektroscirrha* (Mongue et al., 2023). All of these were
simple repeat motifs; RepeatModeler found no complex families (e.g. retrotransposons).
Likewise, we recovered a mere 2,225 genes encoding 2,177 proteins. This gene set contained
61.2% (n=273) of expected complete and single copy orthologs, 0.7% (n=3) duplicated
complete BUSCOs, 3.6% (16) fragmented and 34.5% (n=154) missing BUSCOs. Despite the
low absolute numbers, we note that this annotation marks an improvement over the BUSCOs
identified directly from the genome, strongly suggesting this annotation is not failing to capture
genes in the underlying genome.
226
*General patterns of orthology*
We recovered 2,094 genes conserved in at least two of the three species in some capacity, *i.e.*
either 1-to-1 or 1-to-many. Starting with pairwise orthology, we found 556 genes conserved in
*O. elektroscirrha* and *O. elektroscirrha*-like but absent in *Helicoverpa*-infecting *Ophryocystis*; all
of these were 1-to-1 strictly conserved orthologs. Another 489 were found in *Helicoverpa*-
infecting *Ophryocystis* and *O. elektroscirrha* but not *O. elektroscirrha*-like, and finally a mere 56
proteins were conserved in *Helicoverpa*-infecting *Ophryocystis* and *O. elektroscirrha*-like but not
*O. elektroscirrha*. After accounting for these pairwise orthologs, we identified 993 proteins
conserved across all three lineages. Of these, only 15 were 1-to-many orthologs, and in each of
those cases, the relationship was a 1-to-1-to-2 orthology, in other words most parsimoniously
single lineage-specific gene duplication events. Specifically, we recovered 5 duplications in
*Helicoverpa*-infecting *Ophryocystis*, 9 duplicate orthologs in *O. elektroscirrha*, and 1 single
duplicate ortholog in *O. elektroscirrha*-like. Simple subtraction of these counts from the total
annotated gene sets leaves 633 lineage specific genes in *Helicoverpa*-infecting *Ophryocystis*,
648 unique to *O. elektroscirrha*, and 674 unique to *O. elektroscirrha*-like. Results are
summarized in **Table 3.**
243

| Lineage | Vs. *Helicoverpa*-infecting *Ophryocystis* | Vs. *O. elektroscirrha* | Vs. *O. elektroscirrha*-like | Vs. all | Lineage specific |
|---|---|---|---|---|---|
| *Helicoverpa*-infecting *Ophryocystis* | 2,177 | 489 | 56 | 993 (+5) | 634 |
| *O. elektroscirrha* | 489 | 2,695 | 556 | 993 (+9) | 648 |
| *O. elektroscirrha*-like | 56 | 556 | 2,280 | 993 (+1) | 674 |

**Table 3.** Summary of gene conservation and novelty across three lineages of Ophryocystidae.
Greyed out diagonal cells represent self-vs-self comparisons and list the total gene content. For
the Vs. all comparison, 993 is the number of 1-to-1-to-1 orthologs, with the parenthetical giving
the number of lineage-specific duplicate genes.

*ATPase gene family evolution*

**Figure 2. Maximum likelihood tree of ATPase family genes across Apicomplexa.** ATPase
genes from *Helicoverpa* infecting *Ophryocystis* are denoted with filled red triangles, those of
other *Ophryocystis* with open black triangles. Functions are inferred from relation to other
apicomplexan ATPases with known function.

In total, we identified four ATPases from *Helicoverpa* infecting *Ophryocystis* (**Figure 2**, closed
red triangles). Three of these are orthologous to ATPases found in the genomes of *O.*
*elektroscirrha* and the related *O. elektroscirrha*-like (**Figure 2**, open black triangles). One of
these (HeliOph.00g018340.m01) is inferred to be a sarcoplasmic/endoplasmic reticulum
Calcium ATPase (SERCA), though with a notably longer branch and lower bootstrap support
compared to the other two *Ophryocystis* SERCAs. The other two are orthologous to a pair of
*Ophryocystis* ATPases (HeliOph.00g000870.m01and HeliOph.00g012140.m01) that were
previously identified as the Plasma Membrane-bound Calcium (PMCA) ATPases. In our current
analyses however, these ATPases are confidently placed apart from other PMCAs and showed
reciprocal BLAST hits to phospholipid transporting ATPases. For each of the three ATPases
above, the gene phylogeny recapitulates the species phylogeny, with the newly sequenced

*Ophryocystis* being the outgroup to the other two *Ophryocystis* sequences. Finally, *Helicoverpa*
infecting *Ophryocystis* has a fourth ATPase (HeliOph.00g001320.m01) not found in the other
*Ophryocystis.* This ATPase is confidently placed within the Plasma Membrane-bound Calcium
(PMCA) ATPases of *Cryptosporidium* and other Apicomplexa (**Figure 2**).
272
**Discussion**
We offer the first sequence of *Helicoverpa*-infecting *Ophryocystis*, an apicomplexan parasite of
moths in the genus *Helicoverpa*. This parasite has been called "*Ophryocystis elektroscirrha*-
like", although previous phylogenetic analysis of 18S rRNA placed it in a separate clade from
other similar parasites of *O. elektroscirrha sensu stricto* (Gao et al., 2020). Our present analyses
confirm the distinction between these visually similar parasites at the genomic and genetic level.
279
*Comparative genomics of* Ophryocystis
To start with, the genome of *Helicoverpa*-infecting *Ophryocystis* spans only 7.1 Mb, roughly 1.7
Mb smaller than the two other published *Ophryocystis* genomes and with fewer protein coding
genes (Mongue et al., 2023). Although we lack independent estimates for genome sizes in this
clade (e.g. flow cytometry), the consistency of reports of small genomes in other *Ophryocystis*
(Mongue et al., 2023) and their closest sequenced outgroup *Ascogregarina taiwanensis* (a mere
6.1 Mb, Templeton et al., 2010) gives us increasing confidence that these gregarine
Apicomplexa are well-assembled with truly small genomes. Likewise, the consistently small set
of protein-coding genes (~2,200 – 2,600) raise questions about the core set of genes necessary
for biological function in Apicomplexa. Reductions in genome size and gene content of parasitic
species are well-known phenomena (Keeling, 2004; Keeling & Slamovits, 2005), but these three
parasites of Lepidoptera apparently have half the gene content of even other arthropod-infecting
gregarines in the genus *Porospora* (Boisard et al., 2022). More research is merited on this
large-scale reduction in protein coding genes overall, but as a start, we have parsed basic
patterns of gene conservation.
295
Across the three lineages of *Ophryocystis,* less than half of the gene content (993 genes) is
strictly conserved. When accounting for the pairwise orthologous relationships, each lineage is
left with <700 lineage-specific genes based on current analysis. Although these numbers are
still too large to fully manually explore, additional data from related, yet un-sequenced parasites
(Müller□Theissen et al., 2025) can only shrink these estimates. It may soon be tractable to
explore the genetic basis of host-specificity of these parasites based on a relatively small
number of candidate genes. For now, we explore already identified set of genes-of-interest.
303
*ATPase evolution in* Ophryocystis
ATPases are class of proteins with diverse functions, but generally act to actively transport
molecules either within or into and out of cells (Chène, 2002). For single-celled Apicomplexa
important functions like creating ion gradients for cell-cell signaling (Kelly et al., 1985) are not
relevant, but even single cells need to maintain proper salt balance and transport molecules
against the concentration gradient. Some Apicomplexa possess ATP4 type sodium pumps
(Lehane et al., 2019) but these have never been reported in *Ophryocystis* (Mongue et al., 2023);
the *Helicoverpa*-infecting *Ophryocystis* apparently lacks this class of ATPases as well. Other

gregarine Apicomplexa do possess calcium ATPases, both intracellular and membrane-bound
(Lehane et al., 2019).

Mongue et al. previously reported a pair of PMCAs in milkweed butterfly infecting *Ophryocystis*
albeit with especially long branches (2023). However, in our current re-analysis, we identify
these ATPases as Type V phospholipid transporters, with a pair of orthologs in *Helicoverpa*-
infecting *Ophryocystis*. This parasite has a more confidently assigned PMCA, closely related to
PMCAs of other Apicomplexa and lacking in the two *Danaus Ophryocystis.* Although still not
conclusive, the pattern produced by these new data and analyses is intriguing. Parasites of
*Danaus* butterflies, which feed on plants rich in cardiac glycosides (Ackery & Vane-Wright,
1984), apparently lack any membrane-bound ion pumping ATPases. *Helicoverpa*, the host
genus of this parasite, has a diet that does not include plants containing cardiac glycosides
(Nasreen & Mustafa, 2000; Zalucki et al., 1986) and does retain a PMCA, like other gregarines
(Mongue et al., 2023). Even though cardiac glycosides are not thought to pass through cell
membranes, their inhibitory effects on cell membrane ion pumps has documented knock-on
changes to intracellular calcium concentrations (Lee et al., 1970). Given that cardiac glycosides
have some inhibitory properties for calcium pump ATPases as well (Kelly et al., 1985), it is
possible that the loss of PMCAs in milkweed butterfly parasites is adaptive (Albalat & Cañestro,
2016), or at least neutral in a host environment where such ATPases would be routinely
inhibited.

Finally, all three *Ophryocystis* have intracellular SERCA ATPases, though the branch length
between *Helicoverpa* infecting *Ophryocystis* and the other two is longer than most others in this
class of ATPases. The long branch length between SERCA ATPases of *Helicoverpa*-infecting
*Ophryocystis* and the milkweed butterfly parasites may reflect an evolutionary history of
changes to SERCA sequence in *Ophryocystis* prior to the loss of PMCA ATPases. This
evolutionary history is still speculative and even if true does not explain the continued inhibitory
effects of cardiac glycosides on *O. elektroscirrha* (de Roode et al., 2008). Nevertheless, the
sequencing of another *Ophryocystis* genome has helped refine hypotheses on molecular
mechanisms of host-parasite co-evolution.

**Conclusions**
We have sequenced and annotated the genome of *Helicoverpa*-infecting *Ophryocystis.* Our
results add a valuable resource to studying host-parasite dynamics. Comparisons with other
sequenced *Ophryocystis* offer some additional clarity on adaptation to host biochemistry as well.
More generally, these analyses begin to shed light on a poorly sampled portion of the
apicomplexan tree of life and demonstrate the crucial role that genomics can play in resolving
between organisms with few to no informative phenotypic differences.

**Data availability**
Because this parasite species has yet to be formally described, it does not fit easily into
frameworks like NCBI that require existing species metadata. Instead, raw sequencing reads,
assembly, annotation, and orthology datasets are all available at
https://doi.org/10.5281/zenodo.15856773.

**References**

356
357
**References**
Aardema, M. L., & Andolfatto, P. (2016). Phylogenetic incongruence and the evolutionary
origins of cardenolide-resistant forms of Na+, K+-ATPase in Danaus butterflies.
*Evolution*, *70*(8), 1913–1921.
Abrahamsen, M. S., Templeton, T. J., Enomoto, S., Abrahante, J. E., Zhu, G., Lancto, C. A.,
Deng, M., Liu, C., Widmer, G., & Tzipori, S. (2004). Complete genome sequence of the
apicomplexan, Cryptosporidium parvum. *Science*, *304*(5669), 441–445.
Ackery, P. R., & Vane-Wright, R. I. (1984). *Milkweed butterflies, their cladistics and biology,*
*being an account of the natural history of the Danainae, a subfamily of the Lepidoptera,*
*Nymphalidae.*
Agrawal, A. A., Petschenka, G., Bingham, R. A., Weber, M. G., & Rasmann, S. (2012). Toxic
cardenolides: Chemical ecology and coevolution of specialized plant–herbivore
interactions. *New Phytologist*, *194*(1), 28–45.
Albalat, R., & Cañestro, C. (2016). Evolution by gene loss. *Nature Reviews Genetics*, *17*(7),
379–391.
Aranda, M., Li, Y., Liew, Y. J., Baumgarten, S., Simakov, O., Wilson, M. C., Piel, J., Ashoor, H.,
Bougouffa, S., & Bajic, V. B. (2016). Genomes of coral dinoflagellate symbionts highlight
evolutionary adaptations conducive to a symbiotic lifestyle. *Scientific Reports*, *6*(1),
39734.
Blake, D. P., of Life, W. S. I. T., & Darwin Tree of Life Consortium. (2024). The genome
sequence of the Coccidian parasite, Eimeria praecox (Apicomplexa: Eucoccidiorida).
*Wellcome Open Research*, *9*, 331.
Boisard, J., Duvernois-Berthet, E., Duval, L., Schrével, J., Guillou, L., Labat, A., Le Panse, S.,
Prensier, G., Ponger, L., & Florent, I. (2022). Marine gregarine genomes reveal the

breadth of apicomplexan diversity with a partially conserved glideosome machinery.

*BMC Genomics*, *23*(1), 1–22.

Boisard, J., & Florent, I. (2020). Why the–omic future of Apicomplexa should include gregarines.

*Biology of the Cell*, *112*(6), 173–185.

Brower, L. P., & Fink, L. S. (1985). A Natural Toxic Defense System: Cardenolides in Butterflies

versus Birds. *Annals of the New York Academy of Sciences*, *443*(1), 171–188.

https://doi.org/10.1111/j.1749-6632.1985.tb27072.x

Bushman, M., Morton, L., Duah, N., Quashie, N., Abuaku, B., Koram, K. A., Dimbu, P. R.,

Plucinski, M., Gutman, J., & Lyaruu, P. (2016). Within-host competition and drug

resistance in the human malaria parasite Plasmodium falciparum. *Proceedings of the*

*Royal Society B: Biological Sciences*, *283*(1826), 20153038.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T.

L. (2009). BLAST+: Architecture and applications. *BMC Bioinformatics*, *10*(1), 1–9.

Chène, P. (2002). ATPases as drug targets: Learning from their structure. *Nature Reviews Drug*

*Discovery*, *1*(9), 665–673.

Dangoudoubiyam, S., Norris, J. K., Namasivayam, S., de Paula Baptista, R., Cannes do

Nascimento, N., Camp, J., Schardl, C. L., Kissinger, J. C., & Howe, D. K. (2024).

Temporal gene expression during asexual development of the apicomplexan Sarcocystis

neurona. *Msphere*, e00111-24.

de Roode, J. C., Pedersen, A. B., Hunter, M. D., & Altizer, S. (2008). Host plant species affects

virulence in monarch butterfly parasites. *Journal of Animal Ecology*, *77*(1), 120–126.

https://doi.org/10.1111/j.1365-2656.2007.01305.x

De Roode, J. C., Rarick, R. M., Mongue, A. J., Gerardo, N. M., & Hunter, M. D. (2011). Aphids

indirectly increase virulence and transmission potential of a monarch butterfly parasite

by reducing defensive chemistry of a shared food plant. *Ecology Letters*, *14*(5), 453–

461.

DeBarry, J. D., & Kissinger, J. C. (2011). Jumbled genomes: Missing Apicomplexan synteny.

*Molecular Biology and Evolution*, *28*(10), 2855–2871.

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F.

(2020). RepeatModeler2 for automated genomic discovery of transposable element

families. *Proceedings of the National Academy of Sciences*, *117*(17), 9451–9457.

Gao, K., Muijderman, D., Nichols, S., Heckel, D. G., Wang, P., Zalucki, M. P., & Groot, A. T.

(2020). Parasite-host specificity: A cross-infection study of the parasite Ophryocystis

elektroscirrha. *Journal of Invertebrate Pathology*, *170*, 107328.

Gardner, M. J., Hall, N., Fung, E., White, O., Berriman, M., Hyman, R. W., Carlton, J. M., Pain,

A., Nelson, K. E., & Bowman, S. (2002). Genome sequence of the human malaria

parasite Plasmodium falciparum. *Nature*, *419*(6906), 498–511.

Hayashida, K., Hara, Y., Abe, T., Yamasaki, C., Toyoda, A., Kosuge, T., Suzuki, Y., Sato, Y.,

Kawashima, S., & Katayama, T. (2012). Comparative genome analysis of three

eukaryotic parasites with differing abilities to transform leukocytes reveals key mediators

of Theileria-induced leukocyte transformation. *MBio*, *3*(5), e00204-12.

Humann, J. L., Lee, T., Ficklin, S., & Main, D. (2019). Structural and functional annotation of

eukaryotic genomes with GenSAS. In *Gene prediction* (pp. 29–51). Springer.

Jones, D. T., Taylor, W. R., & Thornton, J. M. (1992). The rapid generation of mutation data

matrices from protein sequences. *Bioinformatics*, *8*(3), 275–282.

Keeling, P. J. (2004). Reduction and compaction in the genome of the apicomplexan parasite

Cryptosporidium parvum. *Developmental Cell*, *6*(5), 614–616.

Keeling, P. J., & Slamovits, C. H. (2005). Causes and effects of nuclear genome reduction.

*Current Opinion in Genetics & Development*, *15*(6), 601–608.

Kelly, R. A., O'Hara, D. S., Canessa, M. L., Mitch, W. E., & Smith, T. W. (1985).

Characterization of digitalis-like factors in human plasma. Interactions with NaK-ATPase

and cross-reactivity with cardiac glycoside-specific antibodies. *Journal of Biological Chemistry*, *260*(21), 11396–11405.

Krishna, S., Woodrow, C., Webb, R., Penny, J., Takeyasu, K., Kimura, M., & East, J. M. (2001). Expression and functional characterization of a Plasmodium falciparum Ca2+-ATPase (PfATP4) belonging to a subclass unique to apicomplexan organisms. *Journal of Biological Chemistry, 276*(14), 10782–10787.

Lechner, M., Findeiss, S., Steiner, L., Marz, M., Stadler, P. F., Prohaska, S. J., Findeiß, S., Steiner, L., Marz, M., Stadler, P. F., & Prohaska, S. J. (2011). Proteinortho: Detection of (Co-)orthologs in large-scale analysis. *BMC Bioinformatics*, *12*, 124. https://doi.org/10.1186/1471-2105-12-124

Lee, K. S., Shin, M. R., Kang, D. H., & Chen, K. (1970). Studies on the mechanism of cardiac glycoside action. *Biochemical Pharmacology*, *19*, 1055–1069.

Lehane, A. M., Dennis, A. S., Bray, K. O., Li, D., Rajendran, E., McCoy, J. M., McArthur, H. M., Winterberg, M., Rahimi, F., & Tonkin, C. J. (2019). Characterization of the ATP4 ion pump in Toxoplasma gondii. *Journal of Biological Chemistry*, *294*(14), 5720–5734.

Levine, N. D. (2018). *The Protozoan Phylum Apicomplexa: Volume 2*. CRC Press.

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO update: Novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Molecular Biology and Evolution*, *38*(10), 4647–4654.

Matsui, H., & Schwartz, A. (1968). Mechanism of cardiac glycoside inhibition of the (Na+-K+)-dependent ATPase from cardiac tissue. *Biochimica et Biophysica Acta (BBA)-Enzymology*, *151*(3), 655–663.

McLaughlin, R. E., & Myers, J. (1970). Ophryocystis elektroscirrha sp. N., a Neogregarine Pathogen of the Monarch Butterfly Danaus plexippus (L.) and the Florida Queen

Butterfly D. gilippus berenice Cramer1. *The Journal of Protozoology*, *17*(2), 300–305.

https://doi.org/10.1111/j.1550-7408.1970.tb02375.x

Mongue, A. J., Markee, A., Grebler, E., Liesenfelt, T., & Powell, E. C. (2024). Genome report:

Genome sequence of tuliptree scale, Toumeyella liriodendri (Gmelin), an ornamental

pest insect. *G3: Genes, Genomes, Genetics*, jkae231.

Mongue, A. J., Martin, S. H., Manweiler, R. E., Scullion, H., Koehn, J. L., De Roode, J. C., &

Walters, J. R. (2023). Genome sequence of Ophryocystis elektroscirrha, an

apicomplexan parasite of monarch butterflies: Cryptic diversity and response to host-

sequestered plant chemicals. *BMC Genomics*, *24*(1), 278.

Müller-Theissen, M. L., Barriga, P., Yan, K., Gottdenker, N. L., & Altizer, S. M. (2025). Museum

collections reveal a global range of Ophryocystis parasites in Danaus butterflies.

*Ecological Entomology*.

Nasreen, A., & Mustafa, G. (2000). Biology of Helicoverpa armigera (Hübner) reared in

laboratory on natural diet. *Pakistan Journal of Biological Sciences*, *3*(10), 1668–1669.

Otto, T. D., Böhme, U., Jackson, A. P., Hunt, M., Franke-Fayard, B., Hoeijmakers, W. A.,

Religa, A. A., Robertson, L., Sanders, M., & Ogun, S. A. (2014). A comprehensive

evaluation of rodent malaria parasite genomes and gene expression. *BMC Biology*, *12*,

1–18.

Penumarthi, L. R., Baptista, R. P., Beaudry, M. S., Glenn, T. C., & Kissinger, J. C. (2024). A

new chromosome-level genome assembly and annotation of Cryptosporidium

meleagridis. *bioRxiv*.

Price, M. N., Dehal, P. S., & Arkin, A. P. (2009). FastTree: Computing large minimum evolution

trees with profiles instead of a distance matrix. *Molecular Biology and Evolution*, *26*(7),

1641–1650.

Sayyari, E., & Mirarab, S. (2016). Fast Coalescent-Based Computation of Local Branch Support

from Quartet Frequencies. *Molecular Biology and Evolution*, *33*(7), 1654–1668.

https://doi.org/10.1093/molbev/msw079

Schliep, K. P. (2011). phangorn: Phylogenetic analysis in R. *Bioinformatics*, *27*(4), 592–593.

Sievers, F., & Higgins, D. G. (2014). Clustal omega. *Current Protocols in Bioinformatics*, *48*(1),

3–13.

Smit, A., Hubley, R., & Green, P. (2019). *2013–2015. RepeatMasker Open-4.0.*

Stamatakis, A. (2006). *Phylogenetic models of rate heterogeneity: A high performance*

*computing perspective*. 8-pp.

Templeton, T. J., Enomoto, S., Chen, W.-J., Huang, C.-G., Lancto, C. A., Abrahamsen, M. S., &

Zhu, G. (2010). A genome-sequence survey for Ascogregarina taiwanensis supports

evolutionary affiliation but metabolic diversity between a Gregarine and Cryptosporidium.

*Molecular Biology and Evolution*, *27*(2), 235–248.

Ter-Hovhannisyan, V., Lomsadze, A., Chernoff, Y. O., & Borodovsky, M. (2008). Gene

prediction in novel fungal genomes using an ab initio algorithm with unsupervised

training. *Genome Research*, *18*(12), 1979–1990.

Tseng, M. (2007). Ascogregarine parasites as possible biocontrol agents of mosquitoes. *Journal*

*of the American Mosquito Control Association*, *23*(sp2), 30–34.

Zalucki, M., Daglish, G., Firempong, S., & Twine, P. (1986). The biology and ecology of

Heliothis-armigera (Hubner) and Heliothis-punctigera Wallengren (Lepidoptera,

Noctuidae) in Australia-What do we know. *Australian Journal of Zoology*, *34*(6), 779–

814.

Zhang, C., Rabiee, M., Sayyari, E., & Mirarab, S. (2018). ASTRAL-III: polynomial time species

tree reconstruction from partially resolved gene trees. *BMC Bioinformatics*, *19*(6), 15–30.

506