A Plasma Membrane Vesicle Imaging-Based Platform for Studying Membrane Fusion

Inbar Yosibash[1,2*], Suman Khan[3*], Alexandra Lichtenstein[4], Raviv Dharan[1,2], Alisa Vaknin[1], Susan Daniel[5], Ori Avinoam[3#], Raya Sorkin[1,2#]

[1]School of Chemistry, Faculty of Exact Sciences, Tel Aviv University, Israel.

[2]Center for Physics and Chemistry of Living Systems, Tel Aviv University, 6997801, Tel Aviv, Israel

[3]Department of Biomolecular Sciences, Weizmann Institute of Science, Rehovot, Israel.

[4]Cellular and Molecular Imaging Center, Faculty of Medicine, Tel- Aviv University, Israel.

[5]Robert Frederick Smith School of Chemical and Biomolecular Engineering, Cornell University, Ithaca, NY, 14853, USA

* These authors contributed equally to this work

#corresponding authors: rsorkin@tauex.tau.ac.il, ori.avinoam@weizmann.ac.il

Abstract:

Membrane fusion is fundamental to biological processes such as fertilization and viral infection. However, studying fusion mechanisms in live cells is challenging due to their complex composition and topography. To address this, we developed a cell-free system based on giant plasma membrane vesicles (GPMVs), enabling us to study the binding and fusion of the SARS-CoV-1 Spike protein with the human ACE2 receptor in a simplified yet physiologically relevant membrane environment. We generated two distinct GPMV populations- one mimicking the viral membrane and the other mimicking the host cell membrane, with the former expressing Spike and cytoplasmic mCherry, and the latter expressing ACE2-GFP. Mixing of these two GPMV populations led to docking of Spike GPMVs onto ACE2 GPMVs. Using live confocal microscopy, we followed and quantified the docking kinetics, confirming receptor accumulation at the docking interface prior to fusion. Fusion was triggered using trypsin, a protease known to activate Spike, and was monitored and quantified using confocal fluorescence microscopy. To quantify fusion efficiency at high throughput, we used Imaging Flow Cytometry. We conclude that the GPMV-based platform offers a versatile tool for dissecting fusion mechanisms beyond viral interactions. By enabling control over experimental conditions, this system provides new opportunities for studying fusion processes with applications in drug delivery, therapeutic design, and fundamental membrane biology research.

Introduction:

Membrane fusion is the process by which two distinct lipid bilayers combine to form a single, continuous membrane. This is a fundamental cellular process essential for various functions such as viral entry and release, intracellular transport, and cell-cell fusion during fertilization and muscle formation (Jena, 2000, Martens and McMahon, 2008, Arnold, 1995). Many enveloped viruses, such as coronaviruses, exploit membrane fusion to infiltrate host cells by merging their lipid envelope with host cell membranes (Tang et al., 2020, Millet and Whittaker, 2018). This generally necessitates fusion proteins that mediate the fusion reaction and lower the energy barrier for fusion.

In the case of coronaviruses such as SARS-CoV-1 (CoV-1), the fusion process is mediated by a type I membrane glycoprotein called Spike, which protrudes from the virus envelope. This protein has two structural domains: a receptor-binding domain, S1, and a fusion mediating domain, S2 (Bosch et al., 2003). The S2 domain contains a fusion peptide that is highly conserved across all CoVs (Madu et al., 2009, Tang et al., 2020). CoV-1 enters host cells by first binding to the ACE2 receptor on the host cell surface. This binding is followed by conformational changes in the Spike protein, triggered by proteolytic cleavage, which promote fusion and enable the delivery of the viral genome into the host cell cytosol (Harrison, 2008, Belouzard et al., 2012, Tang et al., 2020). SARS-CoV-1 can utilize both plasma membrane fusion and endosomal entry pathways, with the preferred route depending on cell type and the availability of specific host proteases.

Here we present a new high- throughput and simple method to study protein interactions and membrane fusion using Giant Plasma Membrane Vesicles (GPMVs). These are cell-derived vesicles that largely retain the compositional lipid and protein complexity of the parent cell plasma membrane but lack cortical actin cytoskeleton and intracellular processes (Sezgin, 2022). By combining GPMVs with high-throughput analysis using Imaging Flow Cytometry, we can rapidly obtain quantitative data on fusion efficiency. We demonstrate the applicability of our approach by examining the fusion process between the SARS-CoV-1 Spike protein and its human receptor, ACE2. We generated two distinct GPMV populations: one expressing ACE2 conjugated to GFP and the other expressing Spike with cytoplasmic mCherry to examine receptor binding and membrane fusion under controlled conditions. We monitored the kinetics of receptor binding, visualized docking interactions, and quantified fusion events following proteolytic activation. We compared Imaging Flow Cytometry and confocal microscopy based quantitative image analysis and obtained a good agreement, demonstrating the reliability of our new method. We further depleted cholesterol using methyl-β-cyclodextrin and observed a corresponding reduction in fusion efficiency, consistent with previously reported findings (Glende et al., 2008, Sanders et al., 2021).

Compared to traditional approaches such as liposome-based fusion assays, supported bilayer systems, or live-cell assays, our GPMV-based platform offers several unique advantages. Unlike synthetic liposomes, GPMVs retain a near-native lipid and protein environment, providing a more realistic membrane model while effectively removing active cellular processes that could affect fusion (Sezgin, 2022). In contrast to supported membranes, GPMVs are free, so their interactions with each other are not influenced by the support (Dimova, 2019). To reveal mechanistic details of fusion processes, it is advantageous to use single vesicles with natively embedded membrane proteins and to capture large numbers of interactions. Our method is broadly applicable beyond SARS-CoV-1 fusion, enabling detailed investigation of membrane fusion mechanisms relevant to drug delivery, antiviral strategies, and fundamental questions in membrane biology.

Results:

Forming GPMVs with labeled content or proteins

We aimed to study protein-protein interactions and fusion using a simplified and controlled system of GPMVs that express distinct proteins of interest. To mimic virus and host membranes, we generated two distinct populations of GPMVs derived from mammalian cells: ACE2- GFP and Spike with cytoplasmic mCherry. (Fig. 1 *A*; see Materials for details, Supplementary movie 1). Fluorescence microscopy of the cell cultures confirmed expression of the fluorescent proteins, and imaging of the GPMVs confirmed the successful incorporation of ACE2-GFP into the respective GPMV membranes,

and mCherry into the cytoplasm of the Spike GPMVs (Fig. 1 *B*). The distinct localization of these proteins in separate vesicles and the added fluorescent labeling provide a controlled system to test whether receptor binding interactions and fusion occur at given conditions.

Figure 1*: The formation of GPMVs: (A) Schematic illustration of GPMV formation. (B) Microscopy images of Flp-In TREx 293 cells expressing ACE2-GFP/ HEK293 cells expressing Spike and cytoplasmic mCherry (left Images), scale bar 500µm. The isolated GPMVs contained Spike in their membrane and cytoplasmic mCherry/ ACE2-GFP in their membrane (right Images), scale bar 5 µm.*

Characterizing protein-protein interactions using GPMVs.

The interaction between the human ACE2 receptor and the SARS-CoV-1 Spike protein was characterized by examining the docking behavior of the two GPMV populations upon their mixing in the presence of Ca^{2+} (Fig. 2 *A*). Live confocal microscopy revealed that upon mixing, GPMVs expressing ACE2-GFP docked onto the Spike expressing GPMVs, with ACE2-GFP fluorescence accumulating at the docking interface over time (Fig. 2 *B*). To characterize this interaction, we measured the kinetics of ACE2-GFP accumulation at the docking site. In these experiments, vesicle pairs were brought into contact in a controlled manner: one vesicle was held with a micropipette and brought into contact with another vesicle (Fig. 2 *C*). This setup enabled precise control over vesicle–vesicle interactions. Additionally, real-time tracking of docking events showed that ACE2 accumulation at the interface reached equilibrium at ~90 seconds (n = 12, Fig. 2 D and *E*, Supplementary movie 2). To assess the enrichment, we compared the photon counts per pixel of ACE2-GFP at the interface to its baseline membrane levels in freely moving GPMVs, we observed a significant enrichment (P < 0.0001, n = 53, Fig. 2 *F*). We also performed parallel experiments with freely moving GPMVs in solution using live confocal microscopy. We observed interface formation between non-immobilized vesicles exhibited a similar equilibration time of 85.4 seconds (n = 23, Supplementary figures 1 and 2, Supplementary movie 3), suggesting that the binding kinetics of ACE2 are consistent regardless of vesicle mobility state. Together, these results indicate that GPMVs provide a robust approach for studying receptor-binding interactions in a controlled system.

Figure 2: *Visualization of receptor binding between Spike and ACE2 GPMVs: (A) Schematic illustration of the experimental procedure of GPMV docking. (B) Time-lapse live confocal microscopy imaging of GPMV docking. Both types of GPMVs mixed with 2mM Ca²⁺ and their docking dynamics were observed. The white arrowheads indicate a representative docking event. Time stamps (in seconds) denote the progression of docking, scale bar 10μm. (C) Schematic illustration of the experimental setup of (D). (D) Time-lapse live confocal microscopy imaging of GPMV docking using the C-trap confocal microscope and micropipette aspiration. A GPMV expressing ACE2- GFP (yellow) was trapped using a micropipette and brought close to a Spike-expressing GPMV (magenta) to observe docking dynamics. The time stamps (in seconds) indicate the progression from initial contact to docking, scale bar 2μm. (E) Kinetics of reaching docking steady state. Exponential fitting of GPMV docking kinetics measured using live confocal microscopy and micropipette aspiration. The plot shows normalized photon counts per pixel over time, representing ACE2-GFP accumulation at the docking interface. The black curve represents an exponential fit (n = 12 vesicles). The docking process reached a steady state at ~90 seconds (indicated by the red dashed line), where the grey dashed line determines time zero. (F) Violin plot comparing the photon counts per pixel of the interface and the ACE2-GFP membrane of the docked GPMVs (n = 53 vesicles). The black dots represent experiments conducted, and the red line represents the median (P < 0.0001, Mann- Whitney).*

Triggering and Quantifying fusion using GPMVs.

To test whether membrane fusion can be triggered in our GPMV system, we examined if proteolytic activation by trypsin (Yamazaki et al., 2023, Qiu et al., 2006, Kim et al., 2022) induces full fusion between docked Spike and ACE2-expressing GPMVs (Fig. 3 *A*). Live confocal microscopy revealed that upon trypsin addition, full fusion events occurred, as indicated by content mixing between the GPMVs (Fig. 3 *B*). Time-lapse microscopy demonstrated that fusion can be observed in real time (Supplementary movie 4 and 5). We observed that the majority of the fusion events occur within 3 minutes of triggering (Supplementary movie 4). Next, we quantified the fusion index 2 hours after triggering. Fusion efficiency was quantified by analyzing several fields of view two hours post-trypsin treatment, capturing unfused, docked, and fully fused GPMV (Fig. 3 *B*). We have measured 4 fields of view each experiment from which we extracted ~159 vesicles per field. The fusion index, calculated as the percentage of fused vesicles relative to the less-abundant vesicle population, was ~30±5 (N = 3). Note that vesicles that undergo fusion can continue fusing with additional vesicles, as can be seen in Figure 3 *B*, where docking of vesicles that have both fluorescent labels is observed.

Figure 3: *Visualization of trypsin-triggered Spike-ACE2 GPMV full fusion in real time. (A) Schematic illustration of the experimental procedure of GPMVs fusion. (B) Endpoint states. Representative field including unfused Spike with mCherry GPMV (magenta arrowhead), unfused ACE2-GFP GPMV (orange arrowhead), fused GPMV (white arrowhead), and GPMV docking (green arrowhead), scale bar 10µm.*

High Throughput fusion analysis using Imaging Flow Cytometry

Confocal microscopy is a valuable tool for quantifying fusion, but it has limitations. There are many out-of-plane vesicles that complicate the data analysis. Furthermore, the experiments are time-consuming and low-throughput. For higher-throughput quantification, we performed ImageStream analysis to distinguish between unfused single vesicles and fused GPMVs at two hours post-trypsin treatment (Fig. 4 *A*, supplementary Fig. 3), and obtained a fusion index of ~33±9 (N = 7). Fusion efficiency measured by ImageStream analysis was consistent with results obtained using confocal microscopy (Fig. 4B). These results confirm that proteolytic activation is sufficient to induce fusion of

docked GPMVs, providing a quantitative framework for studying viral membrane fusion in a controlled, cell-free system.

Next, we sought to determine whether the ImageStream approach could distinguish differences in fusion efficiency. To this end, we tested the effect of plasma membrane cholesterol depletion in both GPMV populations- Spike and ACE2 expressing- using 1mM methyl-β-cyclodextrin, a manipulation previously shown to impair SARS-CoV-1 fusion (Glende et al., 2008). Imaging Flow Cytometry revealed a significant ~2.5-fold reduction in fusion efficiency compared to control (P < 0.001, Fig. 4 C), demonstrating both the requirement of cholesterol for effective fusion and the ability of GPMVs to faithfully recapitulate this aspect of the process. Importantly, these results highlight the suitability of ImageStream for the quantitative analysis of membrane fusion.

Figure 4: *ImageStream can assess fusion efficiency and identify differences under cholesterol depletion. (A) ImageStream Imaging Flow Cytometry GPMVs fusion endpoint (i.e., 2 hours after triggering). Representative images show brightfield, mCherry (yellow), GFP (magenta), and composite channels for individual GPMVs and fused vesicles. Top row displays GPMV expressing ACE2- GFP (yellow), middle row displays GPMV expressing Spike (labeled with mCherry) and bottom row displays a fused GPMV showing the colocalization of GFP and mCherry signals in the composite image, scale bar 5µm. (B) Comparison of fusion index (%) between ImageStream and confocal microscopy. The fusion index was calculated as the number of GPMVs showing colocalization of mCherry and GFP divided by the less abundant GPMV population and multiplied by 100. Data are shown for ImageStream fusion index (~33±9, N = 7) and confocal microscopy (~30±5 , N = 3). The black dots represent experiments conducted, and the red line represents the median. (C) Cells expressing ACE2-GFP and cells expressing Spike-mCherry were treated with 1mM methyl-β - cyclodextrin for 30 minutes prior to GPMV generation. Fusion index was quantified using ImageStream, and normalized to untreated controls. The black line represents the overall mean, and each outlined shape indicates the mean of a single experiment. Cholesterol depletion resulted in a significant reduction in fusion (P < 0.001, Mann- Whitney test). Colors correspond to independent experiments.*

Discussion:

Membrane fusion is essential in many physiological and pathological processes, including viral infection and fertilization (Jena, 2000, Martens and McMahon, 2008, Arnold, 1995). Due to its importance, various experimental methods have been developed over the years to study fusion.

Here, we introduce a new method to investigate membrane fusion and protein- protein interactions in a controlled, cell-free, yet biologically relevant environment using GPMVs.

To demonstrate that GPMVs can be effectively used to test fusion, we used the well-established membrane fusion reaction mediated by the viral Spike protein and the human ACE2 receptor as a model. To characterize the ACE2- Spike interactions, we tracked in real time the accumulation of ACE2- GFP at the interface between GPMVs using confocal fluorescence microscopy within our optical tweezers instrument. The accumulation kinetics reached a steady state within approximately 90 seconds (Fig. 2 *E*), suggesting a rapid and specific interaction between Spike and ACE2. To assess binding specificity, we used GPMVs expressing the membrane protein TSPAN4 conjugated to GFP, and observed no docking events under identical conditions (Supplementary Fig. 4). The kinetics was similar with micropipette-immobilized vesicles and the freely diffusing vesicles (Supplementary Fig. 2), confirming that surface interactions did not impact the binding between ACE2 and Spike at these conditions. This result demonstrates that the GPMV system is a faithful model for studying early events in viral entry.

To quantitatively study fusion, we imaged GPMVs following their mixing and fusion-triggering using confocal fluorescence microscopy. Confocal microscopy provides high-resolution, real-time insights into the fusion process (Fig. 3 *B*). Additionally, it enables various real-time manipulations, such as photoactivation and photobleaching experiments. However, it presented certain limitations: the presence of out-of-plane vesicles complicated accurate quantification as it was often difficult to determine whether vesicles observed in the focal plane were part of a fusion event or simply overlapping, and the low-throughput nature of confocal imaging limited the number of vesicles that could be analyzed in a reasonable time frame, reducing statistical power. In each experiment, we captured four fields of view, with an average of 159 vesicles per field. To measure a larger population of free-standing, non- supported vesicles, we expanded the use of Imaging Flow Cytometry (ImageStream), a technique commonly used for extracellular vesicles (EVs) and synthetic vesicles. ImageStream enables the simultaneous collection of high-resolution images and flow cytometric data from thousands of individual vesicles allowing sensitive detection of single vesicles, morphological characterization, and the ability to distinguish true vesicle fusion events from vesicle overlap or background noise (Clark, 2015, Gorgens et al., 2019). This dual capability is especially valuable when working with heterogeneous vesicle populations, where accurate classification is critical (Botha et al., 2021, Lannigan and Erdbruegger, 2017). ImageStream has also been shown to detect differences in vesicle size, shape, and fluorescence intensity (Godino et al., 2023), and to detect different EVs in complex biological fluids such as blood plasma (Woud et al., 2022).

We mixed ACE2 harboring GPMVs with a population of GPMVs with activated spike protein, and screened the vesicles using flow cytometry (Fig. 4 *A,* Supplementary Fig 3). ImageStream flow cytometry is highly suitable for our assay due to its high-throughput capability, enabling us to analyze approximately 5,000 vesicles within 10 minutes. The fusion index obtained using confocal microscopy was ~30±5 and ~33±9 in ImageStream (Fig. 4 *B*). The similarity between ImageStream and confocal microscopy results further emphasized the applicability of ImageStream for quantitative fusion studies. In our hands, ImageStream effectively distinguished fused GPMVs from single vesicles two hours post-trypsin treatment, complementing the lower-throughput, high-resolution data obtained by confocal microscopy.

To test whether differences in fusion efficiency can be detected with ImageStream, we depleted plasma membrane cholesterol using 1mM methyl-β-cyclodextrin (MβCD), a water-soluble oligosaccharide composed of β(1−4)- linked glucopyranose units arranged in a closed ring structure. Its external surface is hydrophilic, while its inner cavity is hydrophobic. This structure enables MβCD

to enhance the solubility of hydrophobic compounds, such as cholesterol, by encapsulating them within its inner cavity (Castro et al., 2013). Cholesterol depletion disrupts membrane microdomains enriched in cholesterol (Glende et al., 2008, Lu et al., 2008) which are critical for viral entry processes. Consistent with previous studies (Glende et al., 2008, Sanders et al., 2021), We observed an approximately 2.5- fold reduction in fusion efficiency following cholesterol depletion (Fig. 4 *C*).

Several experimental approaches have been commonly used to study membrane fusion. Liposome ensemble assays rely on bulk fluorescence measurements, typically using dequenching to monitor lipid and content mixing between vesicles (Gui and Lee, 2018). These assays provide a quantitative ensemble measure of fusion, however, they average over large populations of vesicles, cannot resolve individual fusion events or intermediates, and are prone to artifacts from leakage and non-linear quenching (Dimova, 2019). Single-vesicle fusion assays allow real-time observation, such as kinetics of single vesicle fusion events (Villamil Giraldo et al., 2022, Cervantes et al., 2025), and are often based on large or small unilamellar vesicles (LUVs and SUVs). This assay includes vesicle fusion to supported planar membranes or to other immobilized vesicles. Fluorescent labeling enables the detection of fusion intermediates using fluorescence microscopy. However, interactions with the support can hinder membrane dynamics, and the small size of the vesicles makes it difficult to control membrane tension (Dimova, 2019). On larger scales, giant unilamellar vesicles (GUVs) fusion assays enable direct visualization of fusion events (Witkowska et al., 2021) and morphological changes using optical and electron microscopy (Grunst et al., 2024, Mangala Prasad et al., 2022). Although fusion between GUVs permits observation of lipid and content mixing as well as mechanical manipulation of membrane tension, reconstituting fusion proteins in GUVs is technically challenging (Dimova, 2019). Another common approach is to use whole cells (Reuter et al., 2023). While this preserves native membrane complexity, studying fusion mechanisms in live cells is challenging due to their complex composition and topography (Sezgin, 2022). However, our system has limitations: GPMVs lack cytoskeletal structure and active machinery, which can be necessary for certain fusion events in live cells. Importantly, both interacting membranes are of the same size and exhibit negligible curvature, unlike viral membranes, which have high curvature that can influence fusion (Golani and Schwarz, 2023). Additionally, vesicle size heterogeneity and variations in protein expression levels may introduce some variability in efficiency and kinetics. Future improvements, such as using size-sorted vesicles and methods to achieve a more uniform protein expression, could further improve the reproducibility and precision of this assay.

Our GPMV-based platform offers a powerful tool for studying membrane fusion in a biologically relevant yet simplified environment. Beyond viral entry, this approach could be applied to investigate fundamental mechanisms of membrane fusion and screen potential inhibitors. Future studies could expand this system to other fusion proteins and environmental conditions, further broadening its applicability in membrane biology and therapeutic development.

Materials and Methods:

Cell Culture- HEK293T (ATCC CRL-3216) or Flp-In T-REx 293 stable ACE2- GFP cells were cultured in DMEM medium (Sartorius 01-055-1A) containing 10% Fetal bovine serum (Gibco A5256701), 2mM glutamate (Sartorius 03-020-1A), 1mM sodium pyruvate (Sartorius 03-042-1B) and 1% penicillin–streptomycin (Gibco 15140122) in 100mm cell culture plates (Corning) at 37°C and 5% CO_2. Dissociation of the cells was achieved with 1mL of trypsin-EDTA (Bio-Lab 004715233100). Cells were collected after centrifugation at 100g for 3 minutes, diluted 1:20, plated on a 100mm plate and cultured until the confluency reached 50–60% and no more than 24 hours before transfection.

Plasmid Preparation and Stable Cell Lines Generation- The plasmid encoding SARS-CoV-1 Spike, including its map is provided in the Supplementary (Supplementary Fig. 5). The plasmid encoding mCherry was obtained from Clontech (catalog no. 632523). Plasmids scale-up was performed using Nucleobond midi-plasmid purification kit (Macherey-Nagel no. 740410). The sequence was verified by Sanger sequencing (ZABAM Instrumentation and Service, Tel Aviv University).

The hACE2-eGFP gene (Addgene plasmid # 154962) was subcloned into pcDNA™5/FRT/TO (Thermo Fisher Scientific) using NEBuilder® HiFi DNA Assembly (NEB) following the manufacturer's protocol. The primers (IDT) used for the PCR amplification were:

Insert - 5' AACTTAAGCTTGGTACCGAGGCTAGCGTTTAAACGGGC 3' and

5' ACTGGACTAGTGGATCCGAGTTACTTGTACAGCTCGTCC 3',

Vector - 5' CTCGGATCCACTAGTCCAG 3' and 5' CTCGGTACCAAGCTTAAG 3'.

The eGFP-pcDNA™5/FRT/TO plasmid was a gift from Dr. Ariel Ben-Bassat (Tel Aviv University). The cloned gene sequences were verified by Sanger sequencing (ZABAM Instrumentation and Service, Tel Aviv University). Stable cell lines were generated by Lipofectamine™ 2000-mediated transfection of ~1 million Flp-In™ T-REx™ 293 cells (Thermo Fischer Scientific) plated in a 6-well plate (Corning) with 2.7µg of pOG44 (Thermo Fisher Scientific) and 0.3µg of gene of interest containing pcDNA™5/FRT/TO constructs. Cells were maintained at 37°C, 5% CO_2 in Dulbecco's Modified Eagle Medium (Sartorius) supplemented with 10% FBS, 2mM L-Glutamine (Thermo Fischer Scientific), and 1mM Sodium Pyruvate (Sartorius). For stable selection, the media was supplemented with 50µg/mL hygromycin and 15µg/mL blasticidin (Invivogen) three days post-transfection, with hygromycin increased to 100µg/mL after seven days. Additionally, 100µg/mL Penicillin and 100µg/mL Streptomycin were added to maintain sterility. Media was replaced every 2–3 days until sufficient cell islands formed, after which cells were expanded for cryopreservation.

HEK293T Cell Transfection and Flp-In T-REx 293 Cell Induction - HEK293T or Flp-In T-REx 293 stable ACE2- GFP cells were plated at 20% confluency on a 100mm plate coated with 10µg/mL Poly-L-lysine (Sigma-Aldrich) to keep the cells attached and to minimize cell debris in solution. At 50% confluency, HEK293 cells were transiently cotransfected with 12µg mCherry plasmid and 12µg Spike plasmid using 60µl Lipofectamine 2000 (Invitrogen, Thermo Fisher scientific) according to the manufacture's protocols. Flp-In T-REx 293 stable ACE2- GFP cells were induced by adding 1µg/mL tetracycline (Sigma-Aldrich) to the medium. The cells were grown at 37°C and 5% CO_2 for 24 to 36 hours to obtain an optimal protein expression.

Depletion of Cholesterol in Cells Prior to GPMV Production- To deplete cholesterol, 100 mm plates containing cells expressing either Spike and mCherry or ACE2-GFP were washed twice with 10 mL PBS. Then, 1 mM Methyl-β-cyclodextrin (Sigma-Aldrich, C4555) was added to the plates, followed by incubation for 30 minutes at 37°C and 5% CO_2. In parallel, control cells (expressing either Spike-mCherry or ACE2-GFP) were incubated in DMEM without MβCD for the same duration. Both cell populations—Spike-mCherry–expressing cells and ACE2-GFP–expressing cells—underwent cholesterol depletion prior to GPMV production for the fusion experiments. For control experiments, GPMVs were prepared from cells treated with DMEM only.

GPMVs Formation- Following protein expression, 100mm plates with cells were washed with 10mL GPMV buffer (20mM HEPES and 150mM NaCl at pH 7.4). GPMV vesiculation was induced using a published methods (Gerstle et al., 2018). Briefly, the cells were washed and incubated with 4mL active vesiculation buffer (GPMV buffer supplemented with 2mM $CaCl_2$, 1.9mM dithiothreitol (DTT)

and 27.6mM formaldehyde) at room temperature for at least 4 hours in the dark, during which GPMVs are spontaneously formed and released into the supernatant. The GPMVs were collected using a cut pipette tip. A centrifugation step of 100g for 3 minutes was employed to remove any cell debris, then, the supernatant was collected. The GPMVs were used on the same day.

Sample Preparation for ImageStream and Confocal Fluorescence experiments- For all experimental setups, GPMVs expressing ACE2-GFP and GPMVs expressing Spike-mCherry were mixed in the presence of 2 mM $CaCl_2$ and incubated at 37°C and 5% CO_2 for at least 2 hours, allowing membrane docking to occur.

Docking Assays: After incubation, samples were directly used for docking analysis.

ImageStream Assays: Following the 2-hour docking incubation, samples were further incubated for 2 hours with 0.83 g/L trypsin (Sigma-Aldrich, T1426) at 37°C and 5% CO_2 prior to imaging.

Docking Kinetics (C-trap Confocal): The mixed GPMVs were immediately imaged following the addition of $CaCl_2$, and docking kinetics was monitored in real time.

Control experiments with TSPAN4-GFP GPMVs - HEK293T cells were seeded at 20% confluency on 100 mm plates pre-coated with 10 μg/mL poly-L-lysine (Sigma-Aldrich, P6282) to ensure cell adhesion and reduce the presence of cell debris in solution. Upon reaching ~50% confluency, cells were transiently transfected with 10 μg of a TSPAN4-GFP expression plasmid (Dharan et al., 2022) using 20 μL of Lipofectamine 2000 (Invitrogen, Thermo Fisher Scientific), and incubated at 37°C in a 5% CO_2 atmosphere for 24–36 hours to allow for protein expression. GPMVs were subsequently prepared as described above. For docking experiments, TSPAN4-GFP-expressing GPMVs were mixed with GPMVs expressing Spike fused to cytoplasmic mCherry in the presence of 2 mM $CaCl_2$, and incubated at 37°C with 5% CO_2 for at least 2 hours.

Confocal and Micropipette Aspiration Setup- The docking kinetics experiment was performed using a C-trap confocal fluorescence optical tweezers setup (LUMICKS) made of an inverted microscope based on a 60X water-immersion objective (NA 1.2). A micropipette aspiration setup including a micromanipulator (Sensapex) holding a micropipette with a diameter of 5μm (Biological industries) was connected to a Fluigent EZ-25 pump and integrated into the optical tweezer instrument. Before and after each experiment, the zero-suction pressure was found by aspirating a polystyrene bead (3.43μm, Spherotech) into the pipette and reducing the suction pressure until the bead stopped moving. Using micropipette aspiration, one ACE2- GFP GPMV was trapped and brought into proximity with Spike mCherry GPMV while tracking the time until reaching the steady state of the docking. The micropipette pressure used was less than 0.1mbar and therefore the GPMV tension was not affected. Experiments were performed at room temperature.

Spinning Disk Confocal Microscopy- Live receptor binding and fusion experiments were performed using a Yokogawa automatic Spinning Disk confocal scanning unit (CSU-W1-T2) mounted on an inverted Olympus IX83 microscope (37°C, with 5% CO_2). 60× 1.4 NA oil immersion objectives were used for data acquisition. Images were captured by back-illuminated Prime 95B sCMOS cameras (Photometrics) controlled by VisiView software (Visitron Systems GmbH). Confocal fluorescence excitation was done with solid-state laser diodes (488nm for eGFP and 561nm for mCherry). The following fluorescence emission filters were used: 525/50nm for eGFP and 609/54nm for mCherry.

Intensity Analysis at Membrane Interfaces of Immobilized GPMVs- A custom graphical user interface (GUI) was developed to assist manual analysis of fluorescence intensity distributions at membrane interfaces from time-series microscopy images. Raw TIFF sequences were first pre-

processed using percentile-based contrast normalization for visualization while preserving the original bit depth for quantitative analysis.

To distinguish between interface and non-interface membrane regions with precise spatial control, a dual region of interest (ROI) approach was implemented where two rotatable rectangular regions were manually positioned on each frame: (1) Interface ROI, placed directly at the membrane contact site to quantify local enrichment; and (2) Reference ROI, positioned at a non-interface membrane region to establish baseline intensity values from the same image. Each ROI could be freely rotated and repositioned with sub-pixel accuracy to accommodate membrane geometries ensuring accurate tracking of membrane interfaces throughout the time series.

Mean fluorescence intensities were calculated for both interface and reference regions at each time point by averaging all pixel values within each mask. For time-series analysis, ROI configurations could be propagated across frames with position correction capabilities to maintain consistent region definitions throughout the sequence. Results were automatically exported as CSV files containing time points and corresponding mean intensity values for both regions.

Intensity Analysis at Membrane Interfaces of Moving GPMVs- To quantify intensity distributions at membrane interfaces from time-series fluorescence microscopy data of free-floating GPMVs, we integrated deep learning-based segmentation with quantitative intensity profiling.

Two-channel TIFF stacks were first preprocessed with percentile-based contrast normalization for segmentation and tracking while preserving the original bit depth for quantitative analysis. For vesicle tracking, reference points on both vesicles of interest within the same frame were manually defined. These points served as input to a SAM2 (Segment Anything Model 2) neural network which propagated segmentation masks across all frames, enabling robust tracking and segmentation of freely moving and deformable GPMVs throughout the time series.

To analyze the membrane interface, we applied computational geometry to detect intersection regions between vesicles, generating perpendicular profile lines that extended from the interface into both vesicles. This automated approach eliminates the both bais and labor-intensive nature of manual ROI placement in our previous method, while providing more precise localization of interface boundaries even as GPMVs pair move and rotate.

Membrane-specific "donut" masks were created through adaptive boundary detection based on intensity gradients at membrane peripheries. The algorithm automatically identified membrane boundaries by analyzing radial intensity profiles, where values exceeding 75% of peak intensity defined the inner membrane boundary, while the outer boundary was established at 80% of peak intensity along the outward gradient. This adaptive approach ensures consistent isolation of membrane regions despite variations in membrane thickness or fluorescence intensity.

Intensity measurements were calculated for both interface and non-interface membrane regions by extracting pixel values along the profiling lines within the donut masks. The analysis was exported as CSV files containing time points and corresponding intensity values for both regions.

IMultispectral Imaging Flow Cytometry (ImageStreamX)- GPMVs were prepared and mixed as described above and imaged using multispectral imaging flow cytometry (ImageStreamX markII flow cytometer; Cytek Biosciences). Single and focused population of GPMVs was gated using brightfield channel: Gradient RMS (feature measures image contrast or focus quality) and morphological features (area and aspect ratio of GPMVs). Between 4 and 8 thousand events of single focused GPMVs were gathered for each sample and visualized. eGFP fluorescence was measured using

488nm wavelength laser,200mW for excitation and channel 2 band emission spectrum of 480-560nm. mCherry fluorescence was measured using laser 561nm wavelength, 200mW for excitation and channel 4 band emission spectrum of 595-642nm. Experiments were performed at room temperature.

ImageStream Fusion Quantification- To quantify the number of fused GPMVs obtained from ImageStream, all image channels (brightfield, GFP, and mCherry) were first normalized to 300 pixels along the largest dimension while preserving the aspect ratio with Hermite interpolation (Supplementary Fig. 6). Following this pre-processing, all image channels were segmented using the Cellpose neural network to generate instance masks of the GPMVs in all three channels.

To identify spherical GPMVs from cellular membrane aggregates and debris, a weighted sphericity assessment was performed on each GPMV instance mask obtained from the BF channel as reference. The following circularity metrics and weights were used: (1) Perimeter-Area (Bryant, 2008) Circularity (35% weight)— implementing the isoperimetric quotient ($4\pi \cdot area/perimeter^2$) to identify shapes approaching perfect circularity; (2) Distance Transform Circularity (35% weight) — quantifying radial uniformity by calculating the variation coefficient of centroid-to-boundary distances and transforming it via exponential function (e^($-2 \cdot variation$)) to penalize irregularities; (3) Convexity-Based Circularity (15% weight) — measuring the area-to-convex-hull-area ratio to identify concavities that deviate from ideal vesicle morphology; and (4) Eccentricity-Based Circularity (15% weight) — calculated as (1-eccentricity) to account for elongation. Values greater than 0.71 were used to select round GPMVs.

To determine whether each reference circular GPMV was eGFP-positive, mCherry-positive, or both, cross-channel colocalization analysis was performed pairwise (i.e., BF-GFP and BF-mCherry masks), requiring mask area overlap (≥70%) with respect to the reference channel. Based on this cross-channel colocalization, GPMVs were classified as either eGFP-positive, mCherry-positive, or both, while GPMVs lacking eGFP or mCherry signal were discarded. The fusion index was calculated as $\frac{GPMVs\ positive\ for\ both\ GFP\ and\ mCherry}{min(GFP-positive,\ mCherry-positive)} \times 100$, using the minimum channel positivity count as the denominator to compensate for potential variations in fluorophore labeling efficiency between markers.

Fusion Index Normalization- Fusion index values were normalized to the mean of the control group. For each experiment, the mean of the control replicates was calculated, and both control and treated (1mM methyl-β-cyclodextrin) values were divided by this average. This normalization set the control group to a mean of 1, allowing the treated group to be expressed as fold change relative to control. Normalized values were used for analysis and visualization.

Docking Normalization and Quantification- To quantify the ACE2-eGFP accumulation at the docking interface, fluorescence values were normalized where 0% represented the intensity before the start of interaction and 100% corresponded to the intensity when steady state was reached, following the formula: normalized value = [(original value - minimum value) / (maximum value - minimum value)] × 100. For each interacting pair, the start of accumulation was manually identified and set as the zero time point for analysis. The normalized data were then fitted to characterize the accumulation kinetics and determine the time to equilibrium using the exponential plateau model: Y = YM - (YM - Y0) × exp(-K × X), where Y0 is the starting fluorescence intensity at the manually aligned starting point, YM is the maximum fluorescence intensity at steady state, K is the rate constant (inverse of time units), and X is time. Both normalization and curve fitting were performed using GraphPad Prism built-in functions.

Methyl- β- cyclodextrin Versus Control Normalization- Raw fusion index values were normalized to the average of the control group within each experiment. Specifically, the mean fusion index of the control samples was calculated, and all corresponding control and 1mM treatment values were divided by this mean.

Code Availability-

Code used for the data analysis is available at:

https://gitlab.com/sorkin-lab/gpmv-fusion-project

Acknowledgements:

I.Y. acknowledges Financial support from the ADAMA Center. We thank Dr. Ziv Porat, Head of the Flow Cytometry Unit at the Weizmann Institute of Science, for valuable discussions regarding the ImageStream experiments. R.S. acknowledges support by the Israel Science Foundation (Grant No. 1289/20), and the NSF-BSF (Grant No. 2021793). Co-Funded by the European Union (ERC ReMembrane 101077502). Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them. O.A. also acknowledges funding from the Minerva Foundation with funding from the Federal German Ministry for Education and Research and the Minna James Heineman Foundation, the Henry Chanoch Krenter Institute for Biomedical Imaging and Genomics, the Schwartz Reisman Collaborative Science Program, and the Yeda-Sela Center for Basic Research. O.A. is an incumbent of the Miriam Berman presidential development chair.

References:

ARNOLD, K. 1995. Chapter 19 - Cation-Induced Vesicle Fusion Modulated by Polymers and Proteins,. *In:* LIPOWSKY, R. A. S., E. (ed.) *Handbook of Biological Physics.* North-Holland.

BELOUZARD, S., MILLET, J. K., LICITRA, B. N. & WHITTAKER, G. R. 2012. Mechanisms of coronavirus cell entry mediated by the viral spike protein. *Viruses,* 4**,** 1011-33.

BOSCH, B. J., VAN DER ZEE, R., DE HAAN, C. A. & ROTTIER, P. J. 2003. The coronavirus spike protein is a class I virus fusion protein: structural and functional characterization of the fusion core complex. *J Virol,* 77**,** 8801-11.

BOTHA, J., PUGSLEY, H. R. & HANDBERG, A. 2021. Conventional, High-Resolution and Imaging Flow Cytometry: Benchmarking Performance in Characterisation of Extracellular Vesicles. *Biomedicines,* 9.

BRYANT, J., SANGWIN, C. 2008. *How Round Is Your Circle? Where Engineering and Mathematics Meet,* -------Princeton, New Jersey, Princeton University Press.

CASTRO, B. M., TORRENO-PINA, J. A., VAN ZANTEN, T. S. & GRACIA-PARAJO, M. F. 2013. Biochemical and imaging methods to study receptor membrane organization and association with lipid rafts. *Methods Cell Biol,* 117**,** 105-22.

CERVANTES, M., MANNSVERK, S., HESS, T., FILIPE, D., VILLAMIL GIRALDO, A. & KASSON, P. M. 2025. Single-Virus Microscopy of Biochemical Events in Viral Entry. *JACS Au,* 5**,** 399-407.

CLARK, R. 2015. Imaging flow cytometry enhances particle detection sensitivity for extracellular vesicle analysis. *Nat Methods,* 12**,** i−ii.

DHARAN, R., GOREN, S., CHEPPALI, S. K., SHENDRIK, P., BRAND, G., VAKNIN, A., YU, L., KOZLOV, M. M., & SORKIN, R. 2022. Transmembrane proteins tetraspanin 4 and CD9 sense membrane curvature. Proc. Natl. Acad. Sci. U. S. A., 119, e2208993119

DIMOVA, R. B. L. A. R. 2019. Chapter Six - Fusion assays for model membranes: a critical review. *Advances in Biomembranes and Lipid Self-Assembly,* 30**,** 229-270.

GERSTLE, Z., DESAI, R. & VEATCH, S. L. 2018. Giant Plasma Membrane Vesicles: An Experimental Tool for Probing the Effects of Drugs and Other Conditions on Membrane Domain Stability. *Methods Enzymol,* 603**,** 129-150.

GLENDE, J., SCHWEGMANN-WESSELS, C., AL-FALAH, M., PFEFFERLE, S., QU, X., DENG, H., DROSTEN, C., NAIM, H. Y. & HERRLER, G. 2008. Importance of cholesterol-rich membrane microdomains in the interaction of the S protein of SARS-coronavirus with the cellular receptor angiotensin-converting enzyme 2. *Virology,* 381**,** 215-21.

GODINO, E., RESTREPO SIERRA, A. M. & DANELON, C. 2023. Imaging Flow Cytometry for High-Throughput Phenotyping of Synthetic Cells. *ACS Synth Biol,* 12**,** 2015-2028.

GOLANI, G. & SCHWARZ, U. S. 2023. High curvature promotes fusion of lipid membranes: Predictions from continuum elastic theory. *Biophys J,* 122**,** 1868-1882.

GORGENS, A., BREMER, M., FERRER-TUR, R., MURKE, F., TERTEL, T., HORN, P. A., THALMANN, S., WELSH, J. A., PROBST, C., GUERIN, C., BOULANGER, C. M., JONES, J. C., HANENBERG, H., ERDBRUGGER, U., LANNIGAN, J., RICKLEFS, F. L., EL-ANDALOUSSI, S. & GIEBEL, B. 2019. Optimisation of imaging flow cytometry for the analysis of single extracellular vesicles by using fluorescence-tagged vesicles as biological reference material. *J Extracell Vesicles,* 8**,** 1587567.

GRUNST, M. W., QIN, Z., DODERO-ROJAS, E., DING, S., PREVOST, J., CHEN, Y., HU, Y., PAZGIER, M., WU, S., XIE, X., FINZI, A., ONUCHIC, J. N., WHITFORD, P. C., MOTHES, W. & LI, W. 2024. Structure and inhibition of SARS-CoV-2 spike refolding in membranes. *Science,* 385**,** 757-765.

GUI, L. & LEE, K. K. 2018. Influenza Virus-Liposome Fusion Studies Using Fluorescence Dequenching and Cryo-electron Tomography. *Methods Mol Biol,* 1836**,** 261-279.

HARRISON, S. C. 2008. Viral membrane fusion. *Nat Struct Mol Biol,* 15**,** 690-8.

JENA, B. P. 2000. Insights on membrane fusion. *Cell Biol Int,* 24**,** 769-71.

KIM, Y., JANG, G., LEE, D., KIM, N., SEON, J. W., KIM, Y. H. & LEE, C. 2022. Trypsin enhances SARS-CoV-2 infection by facilitating viral entry. *Arch Virol,* 167**,** 441-458.

LANNIGAN, J. & ERDBRUEGGER, U. 2017. Imaging flow cytometry for the characterization of extracellular vesicles. *Methods,* 112**,** 55-67.

LU, Y., LIU, D. X. & TAM, J. P. 2008. Lipid rafts are involved in SARS-CoV entry into Vero E6 cells. *Biochem Biophys Res Commun,* 369**,** 344-9.

MADU, I. G., ROTH, S. L., BELOUZARD, S. & WHITTAKER, G. R. 2009. Characterization of a highly conserved domain within the severe acute respiratory syndrome coronavirus spike protein S2 domain with characteristics of a viral fusion peptide. *J Virol,* 83**,** 7411-21.

MANGALA PRASAD, V., BLIJLEVEN, J. S., SMIT, J. M. & LEE, K. K. 2022. Visualization of conformational changes and membrane remodeling leading to genome delivery by viral class-II fusion machinery. *Nat Commun,* 13**,** 4772.

MARTENS, S. & MCMAHON, H. T. 2008. Mechanisms of membrane fusion: disparate players and common principles. *Nat Rev Mol Cell Biol,* 9**,** 543-56.

MILLET, J. K. & WHITTAKER, G. R. 2018. Physiological and molecular triggers for SARS-CoV membrane fusion and entry into host cells. *Virology,* 517**,** 3-8.

QIU, Z., HINGLEY, S. T., SIMMONS, G., YU, C., DAS SARMA, J., BATES, P. & WEISS, S. R. 2006. Endosomal proteolysis by cathepsins is necessary for murine coronavirus mouse hepatitis virus type 2 spike-mediated entry. *J Virol,* 80**,** 5768-76.

REUTER, N., CHEN, X., KROPFF, B., PETER, A. S., BRITT, W. J., MACH, M., UBERLA, K. & THOMAS, M. 2023. SARS-CoV-2 Spike Protein Is Capable of Inducing Cell-Cell Fusions Independent from Its Receptor ACE2 and This Activity Can Be Impaired by Furin Inhibitors or a Subset of Monoclonal Antibodies. *Viruses,* 15.

SANDERS, D. W., JUMPER, C. C., ACKERMAN, P. J., BRACHA, D., DONLIC, A., KIM, H., KENNEY, D., CASTELLO-SERRANO, I., SUZUKI, S., TAMURA, T., TAVARES, A. H., SAEED, M., HOLEHOUSE, A.

S., PLOSS, A., LEVENTAL, I., DOUAM, F., PADERA, R. F., LEVY, B. D. & BRANGWYNNE, C. P. 2021. SARS-CoV-2 requires cholesterol for viral entry and pathological syncytia formation. *Elife,* 10.

SEZGIN, E. 2022. Giant plasma membrane vesicles to study plasma membrane structure and dynamics. *Biochim Biophys Acta Biomembr,* 1864**,** 183857.

TANG, T., BIDON, M., JAIMES, J. A., WHITTAKER, G. R. & DANIEL, S. 2020. Coronavirus membrane fusion mechanism offers a potential target for antiviral development. *Antiviral Res,* 178**,** 104792.

VILLAMIL GIRALDO, A. M., MANNSVERK, S. & KASSON, P. M. 2022. Measuring single-virus fusion kinetics using an assay for nucleic acid exposure. *Biophys J,* 121**,** 4467-4475.

WITKOWSKA, A., HEINZ, L. P., GRUBMULLER, H. & JAHN, R. 2021. Tight docking of membranes before fusion represents a metastable state with unique properties. *Nat Commun,* 12, 3606.

WOUD, W. W., VAN DER POL, E., MUL, E., HOOGDUIJN, M. J., BAAN, C. C., BOER, K. & MERINO, A. 2022. An imaging flow cytometry-based methodology for the analysis of single extracellular vesicles in unprocessed human plasma. *Commun Biol,* 5**,** 633.

YAMAZAKI, E., YAZAWA, S., SHIMADA, T., TAMURA, K., SAGA, Y., ITAMOCHI, M., INASAKI, N., HASEGAWA, S., MORINAGA, Y., OISHI, K. & TANI, H. 2023. Activation of SARS-CoV-2 by trypsin-like proteases in the clinical specimens of patients with COVID-19. *Sci Rep,* 13**,** 11632.