1   Gene dosage effects of 22q11.2 copy number variants on in-vivo measures of white matter
2   axonal density and dispersion
3

Rune Boen[1], Julio E. Villalón-Reina[2], Leila Kushan[1], Kathleen P. O'Hora[1], Hoki Fung[1], Nadine

Parker[3], Ibrahim A. Akkouh[4], Dag Alnæs[3], Ruth O'Hara[5], Matthew John Marzelli[5], Lara Foland-

Ross[5], Christina French Chick[5], Isabelle Cotto[5], Allan L. Reiss[5], Joachim Hallmayer[5], Paul M.

Thompson[2], Ole A. Andreassen[3], Ida E. Sønderby[3,4,6], Carrie E. Bearden[1,7]

[1] Semel Institute for Neuroscience and Human Behavior, University of California Los Angeles,

Los Angeles, CA, USA.

[2] Imaging Genetics Center, Mark and Mary Stevens Neuroimaging and Informatics Institute,

University of Southern California, Los Angeles, CA, USA

[3] Centre for Precision Psychiatry, Division of Mental Health and Addiction, Oslo University

Hospital & University of Oslo, Oslo, Norway

[4] Department of Medical Genetics, Oslo University Hospital, Oslo, Norway

[5] Stanford University School of Medicine, Stanford, CA, USA

[6] KG Jebsen Centre for Neurodevelopmental Disorders, University of Oslo, Oslo, Norway.

[7] Department of Psychology, University of California Los Angeles, Los Angeles, CA, USA.

Correspondence: Carrie E. Bearden
A7-460 Semel Institute, Los Angeles, CA 90095, USA
cbearden@mednet.ucla.edu

Keywords: 22q11.2 copy number variant, NODDI, white matter, axonal density, axonal
dispersion

27   Abstract

22q11.2 deletion (22qDel) and duplication (22qDup) carriers have an increased risk of

neurodevelopmental disorders and exhibit altered brain structure, including white matter

microstructure. However, the underlying cellular architecture and age-related changes

contributing to these white matter alterations remain poorly understood. Neurite orientation

dispersion and density imaging (NODDI) was used on mixed cross-sectional and longitudinal

data to examine group differences and age-related trajectories in measures of axonal density (i.e.,

intracellular volume fraction; ICVF), axonal orientation (orientation dispersion index; ODI) and

free water diffusion (isotropic volume fraction; ISO) in 50 22qDel (n scans = 69, mean age =

20.7, age range = 7.4-51.1, 64.0% female) and 24 22qDup (n scans = 34, mean age = 21.6, age

range = 8.3-49.4, 54.2% female) carriers, and 890 controls (n scans = 901, mean age = 21.9, age

range = 7.8-51.1, 54.5% female). The results showed widespread gene dosage effects, with

higher ICVF in 22qDel and lower ICVF in 22qDup compared to controls, and region-specific

effects of the 22qDel and 22qDup on ODI and ISO measures. However, 22qDel and 22qDup

carriers did not exhibit altered age-related trajectories relative to controls. Observed differences

in ICVF suggest higher white matter axonal density in 22qDel and lower axonal density in

22qDup compared to controls. Conversely, differences in ODI are highly localized, indicating

region-specific effects on axonal dispersion in white matter. We do not find evidence for altered

developmental trajectories of axonal density or dispersion among 22q11.2 CNV carriers,

suggesting stable disruptions to neurodevelopmental events before childhood.

Introduction

22q11.2 copy number variants (CNVs) are rare, recurrent structural genetic variants that have

profound impacts on neurodevelopment and confer increased risk for several developmental

neuropsychiatric disorders, including autism spectrum disorder (ASD) and attention

deficit/hyperactivity disorder (ADHD)(1–4). The 22q11.2 deletion (22qDel) is estimated to occur

in one in ~3700 live births(1), where 90% of cases are estimated to be due to de novo events(5).

The 22qDel is also associated with a range of medical and neurobehavioral outcomes, including

congenital heart defects, immune deficiencies, and intellectual disability(1,6,7), and confers a

markedly increased risk of psychosis(1–4). The 22q11.2 duplication (22qDup) is estimated to

occur in one in ~1600 live births(1) and, conversely, confers a putative reduced risk of

psychosis(4,8,9). Moreover, the 22qDup is more likely to be inherited rather than de novo, and

may have a more variable clinical phenotype presentation(10,11) compared to the 22qDel.

Despite their divergent risk for psychotic illness, both 22qDel and 22qDup are associated with

elevated rates of ASD, ADHD, and intellectual disability compared to controls(1,3,12), and

disrupt dosage-sensitive genes that play fundamental roles during neurodevelopment. Thus, in

contrast to research on individuals with behaviorally defined psychiatric disorders, studying a

molecularly confirmed CNV like the 22q11.2 locus provides a genetically traceable and

hypothesis-driven probe of brain alterations in individuals with a high risk of developing brain

disorders. Moreover, reciprocal CNVs (i.e., gain and loss of genomic material at the same locus)

allows for investigation of converging and diverging neurodevelopmental pathways of brain

alterations by examining gene dosage effects on brain phenotypes.

22q11.2 CNVs have been associated with medium to large differences in several measures of

brain structure, including grey and white matter(13–16), suggesting altered cellular architecture.

However, the complex neuronal composition and dispersion of the cortical mantle of the brain

makes it challenging to study the cellular architecture in MRI-derived measures of grey matter,

such as cortical thickness. In contrast, the white matter consists of predominantly axons, which

can be probed using diffusion MRI (dMRI) in vivo. As water diffusion is restricted or hindered

by biological tissue, dMRI can be used to generate measures of white matter microstructure that

is influenced by the cellular architecture along white matter tracts(17–19). For instance, water

molecules in an environment of dense and coherent white matter axons will diffuse along the

primary direction of the axons, which has been commonly measured by diffusion tensor imaging

(DTI)-derived metrics such as fractional anisotropy (FA, i.e., standardized value of anisotropic

diffusion) and mean diffusivity (i.e., measure of diffusion in all directions).

A previous consortium-based DTI-study of 22qDel carriers reported an overall pattern of higher

FA and lower diffusivity across most white matter tracts in 22qDel carriers compared to

controls(13), which could reflect an increase in the cumulative cellular membrane circumference,

increase in smaller tortuous axons(13), and/or higher density of axons with disproportionately

small diameters(7, 14) compared to controls. Indeed, other studies have suggested other lines of

evidence for increased axonal density in 22qDel carriers, such as excessive prenatal overgrowth

of thalamocortical axons in a cellular model of the 22qDel(21) and a higher frequency of deep

layer cortical neurons in a post-mortem examination of a 3-month-old infant with 22qDel(22). In

contrast to 22qDel, only one prior small dMRI study has included 22qDup carriers(17). Despite

the modest sample size, the results indicated widespread lower FA and higher diffusivity in

22qDup carriers compared to controls(23). Although the biological underpinnings of these

differences are unknown, the opposing pattern in 22qDel and 22qDup carriers suggests a gene

dosage effect, which may reflect divergent effects of the 22q11.2 CNV on the cellular

architecture of white matter tracts. Indeed, the 22q11.2 locus is a hotspot for genes critical to

white matter development, including axonal morphology and function in white matter such as

regulation of axonal diameter, growth and branching (e.g., *ZDHHC8*(24,25), *TBX1*(26),

RTN4R/Nogo-66(27), *TXNRD2*(28)), as well as dendritic morphology and neuronal excitability

(e.g., *DGCR8*(29,30)). In addition, both 22qDel and 22qDup affect expression of genes outside

the 22q11.2 locus(21,29,31,32), including genes that influence axonal morphology(21).

However, to our knowledge, no previous studies have directly compared axonal density in white

matter tracts between 22qDel and 22qDup carriers to determine if a gene dosage effect exists.

Moreover, such effects may be region-specific due to the unique developmental trajectories of

white matter fibers. For example, limbic, commissural and projection fibers develop and mature

earlier than association fibers(33–36). Indeed, it has been postulated that the nature of the white

matter alterations in 22qDel carriers may differ between commissural and association fibers(13),

potentially influenced by the maturational patterns of these fiber tracts. Specifically, the presence

of commissural fibers are observed in the first trimester to second trimester(37–42), whereas the

emergence of association fibers occurs later during the second trimester(38,39,42,43). The

number of axons peak around birth for commissural fibers(44,45), followed by axonal

elimination(45,46), axon growth and myelination from birth through adolescence (42,47,48). The

association fibers are the least mature fibers during infancy(49) and are the last to mature, as

indicated by developmental changes in DTI measures well into adulthood(36). Thus, it is

plausible that the 22q11.2 CNV may have maturation-specific effects on commissural and

association tracts.

Conventional DTI measures are inherently nonspecific to the underlying white matter cellular

architecture(18,50,51) and can be influenced by alterations to axonal density and/or dispersion

that can be driven by early (e.g., axonal proliferation or axonal pruning) or later (e.g., axonal

diameter expansion) neurodevelopmental events. In contrast, neurite orientation dispersion and

density imaging (NODDI) is a dMRI method that is better suited to test the hypothesis of altered

axonal density in 22q11.2 CNV carriers, as it provides measures of the biophysical properties of

white matter tracts, including axonal density (i.e., intracellular volume fraction, ICVF), axonal

dispersion (i.e., orientation dispersion index, ODI), and free water diffusion/cerebrospinal fluid

(i.e., isotropic water diffusion, ISO). It is important to note that NODDI generates metrics that

provide indirect proxies of underlying microstructural properties and cannot directly disentangle

the specific biological events (e.g., along-axon diameter variations and/or glial changes) that

drive these signal differences(52). Nevertheless, as a novel technique that disentangles the

composite signal of FA derived from DTI, NODDI metrics allow for more precise and

biologically meaningful inferences about the underlying white matter architecture. Moreover,

NODDI-derived measures have been found to correlate with their histological counterparts(53–

55) and to be altered in several neuropsychiatric disorders, including psychosis and ASD(56).

Thus, examining NODDI measures in 22q11.2 CNV carriers may be informative for detecting

converging neurobiological patterns with idiopathic developmental neuropsychiatric disorders.

In addition, ICVF has been found to be more sensitive to age-related differences in white matter

microstructure compared to DTI-derived metrics during development(57). As such, this metric

can yield novel insight into age-specific neurobiological events that underlie changes in white

matter microstructure(58–60), such as prenatal proliferation of axons(44,45) and prolonged

periods of axonal diameter expansion across childhood and adolescence(58). Thus, by mapping

the neurodevelopmental trajectories of the axonal architecture across white matter regions, we

can obtain important insight into the maturational mechanisms that underlie altered white matter

observed in 22qDel and 22qDup carriers.

Here, we aim to characterize the effects of 22q11.2 CNVs on axonal density and axonal

dispersion in white matter regions. We hypothesize gene-dosage effects on axonal density in

white matter regions such that 22qDel carriers will exhibit higher and 22qDup carriers will

exhibit lower axonal density, respectively. We also aim to investigate spatiotemporal effects of

the 22q11.2 CNV on axonal architecture by examining age-related trajectories using mixed

cross-sectional and longitudinal data, and by leveraging a large dataset of typically developing

controls from the Human Connectome Project (HCP)(61,62). We reasoned that alterations to

prenatal neurodevelopmental processes will manifest as robust group differences in NODDI

measures, whereas alterations to later neurodevelopmental processes will be reflected by altered

age-related trajectories in NODDI measures across childhood and adolescence.

156   Methods

Participants

A total of 964 participants with 1,004 scans were included in the analysis, including 50 22qDel

carriers, 24 22qDup carriers and 890 age and sex matched typically developing controls (Table

1). The 22qDel carriers were recruited from either the University of California, Los Angeles

(UCLA) or Stanford University, whereas the 22qDup carriers were recruited from UCLA only.

The control group included unrelated healthy control participants recruited from UCLA and

Stanford University, and healthy control participants derived from the HCP, including HCP-

Development(61) and HCP-Aging(62) to establish robust developmental trajectories from

childhood to adulthood (see sTable 1-2 for baseline characteristics across scanner sites, sTable 3

for binned age groups, sFigures 1-2).

Diffusion MRI acquisition, preprocessing and quality control

Multi-shell dMRI was acquired using a 3T Siemens Prisma scanner with the following

parameters: TE = 89.2ms, TR = 3230ms, flip angle = 78°, slice thickness = 1.5mm, voxel size =

1.5mm isotropic obtained with 7 b = 0s/mm^2 images, and 3 b = 200s/mm^2, 6 b = 500s/mm^2, 46 b

= 1500s/mm^2 and 46 b = 3000s/mm^2 diffusion weighted volumes. The scans were acquired with

both anterior to posterior and posterior to anterior phase encoding directions, resulting in a total

of 216 volumes, following the HCP acquisition protocol(63). Manual inspections and statistical

analyses were conducted to exclude participants with excessive head movement (see

supplementary note 1 for details). We also included dMRI data derived from the HCP-D

(https://www.humanconnectome.org/study/hcp-lifespan-development) and HCP-A

(https://www.humanconnectome.org/study/hcp-lifespan-aging) as described elsewhere(61,62).

All the raw dMRIs were denoised with LPCA and corrected for Gibbs ringing artifacts using

DiPy (64,65); susceptibility induced distortions, head movement, and eddy currents using

TOPUP and EDDY in FSL (66,67). Afterwards, we ran the diffusion preprocessing pipeline

from HCP (https://github.com/Washington-University/HCPpipelines). NODDI was fitted using

the dMIPY package (https://github.com/AthenaEPI/dmipy). The NODDI derived scalar maps

were registered to the ENIGMA-DTI template(68) and average skeletonized ROI-based

measures were calculated based on the Johns Hopkins University White Matter atlas(69)

following the ENIGMA-DTI protocol (https://enigma.ini.usc.edu/protocols/dti-protocols/)(68),

including anterior corona radiata (ACR), anterior limb of internal capsule (ALIC), body of

corpus callosum (BCC), corpus callosum (CC), cingulum cingulate gyrus (CGC), cingulum

hippocampus (CGH), corona radiata (CR), corticospinal tract (CST), extreme/external capsule

(EC), fornix (FX) , fornix (crus)/stria terminalis (FXST), genu of corpus callosum (GCC),

internal capsule (IC), posterior corona radiata (PCR), posterior limb of the internal capsule

(PLIC), posterior thalamic radiation (PTR), retrolenticular part of the internal capsule (RLIC),

splenium of corpus callosum (SCC), superior corona radiata (SCR), superior fronto-occipital

fasciculus (SFO), superior longitudinal fasciculus (SLF), sagittal stratum (SS), tapetum of the

corpus callosum (TAP), uncinate fasciculus (UNC) (see sFigure 3 for a brain map). Due to

repeated measures across scanner sites, we harmonized all of the NODDI measures using

longCombat(70). Due to the focus on white matter regions, which mostly consist of myelinated

axons(71), we define ICVF as a measure of axonal density, ODI as a measure of axonal

dispersion and ISO as a measure of free water diffusivity(51).

Statistical analyses

Group comparisons

We used generalized additive mixed models (GAMM) in R(72) to test for group differences in ICVF, ISO and ODI across 27 variables (24 regions of interest (ROIs) and 3 average measures). Here, we compared 22qDel carriers to controls, 22qDup carriers to controls, and 22qDel carriers to 22qDup carriers. In addition, we ran gene dosage analyses on NODDI measures by treating copy number as a continuous variable (22qDel = 1, control = 2, and 22qDup = 3). All group analyses were adjusted for the smoothed age effect using cubic regression splines, sex, and repeated measures through the inclusion of participant ID as random intercept. All GAMM analyses were fitted using default arguments, where k = -1 is set for the smoothed age terms and the use of restricted maximum likelihood when fitting all models. The standardized beta is reported and used as a measure of effect size. To account for multiple testing, we applied FDR correction across 75 (i.e., dependent variables) * 4 (i.e., three group analyses and one gene dosage analysis) = 300 comparisons. Finally, we conducted several sensitivity analyses to test the robustness of the results by including 1) participants from the same scanner site only, and by adjusting for 2) cerebral white matter volume, 3) intracranial volume or 4) cerebrospinal fluid volume (see supplementary note 2 for details).

Developmental trajectories

To examine the developmental trajectories of the NODDI measures, we examine the smoothed effect of age, with cubic regression splines using GAMM, in controls, 22qDel carriers and 22qDup carriers. All age-related analyses were conducted on site-harmonized data, and adjusted for the fixed effect of sex and the random effect of subject ID. First, we present the smoothed effect of age for the control group, along with the corresponding FDR-corrected p-values specific to the control group analysis, to establish typical developmental trajectories across white matter regions. The effective degrees of freedom (edf) represent the complexity of the smoothed age term, where values close to 1 indicate a linear relationship and higher values indicate non-linear relationships between chronological age and the response variable. We then compared the developmental trajectories of 22qDel and 22qDup carriers to the control group. A significant interaction term between the smoothed-age term and 22qDel or 22qDup indicates a difference in

233 developmental trajectory compared to controls. To account for multiple testing, we applied FDR-

234 correction across all the interaction terms (i.e., age*22qDel and age*22qDup).

236   Results

Gene dosage effects of the 22q11.2 CNV on intracellular volume fraction

Mean ICVF was significantly greater in 22qDel carriers compared to controls, and significantly

greater across all regions except for FX and CGH. In contrast, 22qDup carriers showed

significantly lower mean ICVF, and lower ICVF across all white matter regions compared to

controls. Reflecting a similar pattern, there was a negative gene dosage effect of the 22q11.2

CNV (i.e., decreasing ICVF with increasing copy number) on mean ICVF and regional ICVF

across all white matter regions (Figure 1, left panel, sTable 3). Sensitivity analyses revealed that

scanner site, cerebral white matter volume, intracranial volume, and cerebrospinal fluid volume

did not influence these results (sFigure 4, sTable 5-8).

Higher mean, but regionally variable effects, of 22q11.2 CNV on orientation dispersion

index

Both 22qDel and 22qDup carriers showed greater mean ODI compared to controls; however,

there were region-specific differences. 22qDel carriers had higher ODI in limbic (i.e., FXST,

CGH), commissural (i.e., GCC) and association (i.e., SS, SLF, SFO, EC) fiber tracts, but also

directionally variable effects on projection tracts (i.e., higher ODI in ACR and PTR; lower ODI

in IC, PLIC, ALIC and CST) compared to controls. 22qDup carriers showed higher ODI in

projection (i.e., IC, PLIC, ALIC, CST), commissural (i.e., BCC) and association (i.e., CGC,

UNC, EC) fiber tracts, but also lower ODI in PCR, where projections fibers predominate,

compared to controls (Figure 1, middle panel, sTable 3). The sensitivity analyses revealed the

same pattern, albeit with fewer significant results; opposing effects of the 22qDel and 22qDup on

ODI in projection (i.e., PTR, PLIC, IC) and association regions (i.e., SS and SLF) yielded the

most robust effects (sFigure 5, sTable 5-8).

Reduced free water diffusion in 22qDup carriers

There was no significant difference in mean ISO between 22qDel and controls. Regionally,

however, 22qDel carriers exhibited higher ISO in a few projection tracts (i.e., PCR, ACR, PTR),

and directionally variable effects on limbic (i.e., higher in FX; lower in FXST) commissural

fibers (i.e., higher in TAP; lower in SCC) compared to controls (Figure 1, right panel).

Sensitivity analyses revealed only higher ISO in the FX and PTR in 22qDel compared to controls

after adjusting for intracranial volume, white matter volume (significant for FX and PTR), and

cerebrospinal fluid volume (significant for FX; sFigure 3, sTable 3). 22qDup carriers showed

lower mean ISO compared to controls, primarily driven by lower ISO in white matter regions

predominated by projection fibers (RLIC, CR, PCR, SCR, ACR, PTR) and association fibers

(UNC, SS, SLF, SFO, Figure 1, right panel). However, similar to the 22qDel carriers, 22qDup

carriers also exhibited higher ISO in FX and lower ISO in FXST. Overall, sensitivity analyses

yielded the same pattern of significant group-level differences between 22qDup and controls,

except for the FX, UNC, RLIC and ACR that were no longer significant (sFigure 6, sTable 5-8).

22q11.2 CNV carriers show typical developmental trajectories in axonal density and

dispersion

In controls, there were significant age effects on ICVF, ODI and ISO across all white matter

regions from childhood to adulthood, except for ODI in the TAP (sTable 9). Most of the

measures showed a non-linear age effect across white matter regions, except for ODI in ALIC,

CST, EC, SFO, SLF and SS, which showed a linear age effect indicated by their effective

degrees of freedom (i.e., edf = 1). The age-related trajectories of 22qDel and 22qDup carriers did

not statistically differ from the age-related trajectories of the control group for the global and

regional measures of ICVF (Figure 2), ODI (Figure 3) or ISO (Figure 4) after adjusting for

multiple comparisons (sTable 10). Follow-up analyses also revealed that several group

differences, most prominently in ICVF, are already evident in childhood and persist through

adolescence and into adulthood (sFigure 7, sTable 11).

291   Discussion

Hitherto, this is the largest study to examine biophysical dMRI-derived measures in brain white

matter among 22q11.2 CNV carriers. The results showed widespread higher ICVF in 22qDel and

lower ICVF in 22qDup compared to controls, where gene dosage at the 22q11.2 locus was

associated with lower ICVF across all white matter regions, suggesting a strong effect of the

22q11.2 CNV on axonal density. Further, we found higher mean axonal dispersion (ODI for both

22qDel and 22qDup compared to controls, but also directionally variable effects of the 22q11.2

CNV on ODI across white matter regions; overall, having more copies of 22q11.2 genes was

associated with lower ODI in regions where association, projection and limbic fibers

predominate. However, we also observed that more copies at the 22q11.2 genomic locus were

associated with higher ODI in IC and CST regions. In addition, we observed lower ISO in

22qDup carriers compared to controls in several white matter tracts where projection and

association fibers predominate. Finally, we find that 22qDel and 22qDup carriers showed similar

developmental trajectories to controls in all NODDI measures across childhood and adolescence.

The current study extends on the previous findings of higher FA in 22qDel(13) and lower FA in

22qDup(23) relative to typically developing controls, by focusing on measures of two of the

main contributors to FA: axonal density and axonal dispersion(51). Across regions, we found

robust evidence for higher axonal density in 22qDel carriers and lower axonal density in 22qDup

carriers compared to controls, suggesting that axonal density is the main contributor to

differences in FA alterations in white matter microstructure among 22q11.2 CNV carriers. It

should be noted, however, that some white matter regions have previously been found to show

lower (i.e., FXST, SLF, EC) or no difference in FA (i.e., RLIC, UNC, SS, PTR, SFO) in 22qDel

carriers compared to controls(13). Here, we find that all these regions, except for the RLIC and

UNC, were characterized by both higher axonal density and higher axonal dispersion in 22qDel

carriers compared to controls. Thus, it seems that higher axonal dispersion counteracts the

influence of higher ICVF, yielding lower or similar FA relative to controls in 22qDel carriers.

The greatest dosage effects of the 22q11.2 genomic region on ICVF were found in commissural

tracts and the corona radiata, which consist of fibers in the corpus callosum and projection fibers

from motor cortex, respectively. Such strong opposing effects on white matter architecture may

be driven by mechanisms that are directly related to the deletion and duplication of genes within

the 22q11.2 locus. However, the exact mechanisms underlying the dosage effect on these tracts

are unknown, but it should be noted that the corpus callosum has been found to be larger in

children with the 22qDel, which contrasts with the overall pattern of lower brain volume in

22qDel carriers(14,15). As corpus callosum is made up by subregions where the size of the

axons typically differ in humans, e.g., the relative density of thick and thin axons (i.e., the axon

diameter distribution) differs across the corpus callosum(73–75), it seems plausible that axonal

overgrowth or axonal diameter expansion may underlie alterations to axonal density of

commissural fibers and corpus callosum volume. Indeed, deletion of genes within the 22q11.2

locus (e.g., *ZDHHC8* and *RTN4R/Nogo-66)* has been linked to axonal growth and branching in

mice models of the 22qDel(24,27). Others have shown that heterozygous deletion of the *TBX1*

among mice can result in relatively more axons with small diameters and no large diameters in

the fimbria compared to wild-type mice(26), which is in line with the predictions from previous

diffusion MRI studies on 22q11.2 deletion carriers(13,20).

Our results also revealed higher ODI in 22qDel carriers for a few regions that are predominated

by association fibers. Lower FA has previously been found in these white matter regions in

22qDel compared to controls(13). While lower FA can indicate both lower axonal density and/or

higher axonal dispersion/crossing fibers(51), our result of increased ODI indicates that this is

primarily driven by higher axonal dispersion in association tracts in 22qDel carriers relative to

controls. Moreover, long association fibers typically originate from upper cortical layers and

show protracted development compared to limbic and projection fibers(33–36), which may

indicate that the white matter fibers originating from upper cortical layers show different

morphology compared to fibers originating from deeper cortical layers. This interpretation aligns

with several lines of previous research, including a mouse model of 22qDel showing fewer

neural progenitors(76), shorter axons, and less dendritic branching in neurons from the upper

cortical layers relative to wild type mice(28). The latter finding was observed after deletion of

the *TXNRD2* gene*,* which also resulted in fewer long distance cortical connections and more

local connections, which may yield higher axonal dispersion for long distance/association fibers

in 22qDel carriers (e.g., lower alignment and/or rerouting through compensation of alternative or

less direct paths). In addition, a post-mortem study reported lower frequency of neurons in the

upper cortical layers in a 22qDel carrier(22), and a neuroimaging study reported altered axonal

morphology (i.e., utilizing ultra-strong dMRI gradients sensitive to axonal morphology) and

lower white matter volume in association tracts in 22qDel carriers compared to controls(20),

supporting this interpretation.

Overall, the results point towards a divergent pattern of white matter microstructure between

22qDel and 22qDup carriers; however, there were a few points of convergence. Such diverging

and converging effects are of particular interest in research on genomic structural variants, where

gene dosage effects are often observed for neuroimaging derived features(16,23,77) but where

the clinical phenotypic profile may converge (e.g., increased risk of ASD for both 22qDel and

22qDup carriers(12)). In the current study, we observed that a few regions exhibit similar

directions of effect for both 22qDel and 22qDup, including higher ODI in EC, higher ISO in FX

and lower ISO in FXST. These results indicate higher axonal dispersion in EC in both 22qDel

and 22qDup, whereas the differences in FX and FXST may be driven by CSF partial volume in

the voxels at the fornix and adjacent areas. White matter alterations in the external capsule have

also been associated with idiopathic ASD (i.e., lower FA in twins with autism compared to

control twins)(78) and symptom severity, and it is one of the white matter regions most

susceptible to environmental factors in individuals with ASD(78). Moreover, a recent pilot study

found within-subject white matter changes in the external capsule among toddlers with ASD

after a behavioral intervention that improved verbal and communication skills(79), possibly

implicating a role of the external capsule in individuals with or at risk for speech and

communication problems. Our results suggest altered axonal circuitry in the external capsule

may be implicated in the behavioral phenotypes of ASD commonly observed in 22q11.2 CNV

carriers, although future research on the relationship between external capsule white matter

structure and autistic phenotypes in 22q11.2 CNV carriers is required to understand this

mechanism.

The 22qDel is also one of the greatest genetic risk factors for psychosis(4,80). A previous

neuroimaging study has found evidence for lower axonal density in commissural and association

fibers and higher axonal density and free water in the anterior thalamic radiation, predominated

by projection fibers, among individuals with first episode psychosis(56). The differences in

thalamic radiation converge with our results of higher ICVF and ISO in the PTR among 22qDel

carriers compared to controls. A recent study reported thalamocortical axonal overgrowth in

22qDel thalamocortical organoids, likely due to elevated *FOXP2* expression as a trans-effect of

the 22qDel(21). In addition, hyperconnectivity in thalamic circuitry has been reported among

22qDel carriers compared to controls(81–83), which has also been hypothesized as a potential

biomarker for psychosis(84). It is also noteworthy that the 22qDup, carriers of a putative

protective genetic factor for psychosis(4,8,9), showed the opposite pattern in our study, reflected

by lower ICVF and ISO in the PTR. This may suggest that a higher axonal density of thalamic

projection fibers, possibly due to thalamocortical axonal overgrowth, is implicated in the

etiology of psychosis.

Finally, despite widespread group differences between 22q11.2 CNV carriers and controls, we

did not find strong support for 22q11.2 CNV-dependent differences in the developmental

trajectories of NODDI measures. Thus, the results do not support alterations in late maturational

changes in axonal architecture, such as cumulative axonal diameter expansion. These findings

contrast with the notably altered developmental pattern of resting-state functional connectivity

observed in 22qDel carriers compared to controls(81,85). Concomitant with these findings, a

parallel investigation in a mouse model of 22qDel found evidence for developmental changes in

dendritic spine density(85). Taken together, these results suggest that changes in synapses and

spine density may reshape brain circuits during adolescence, while altered axonal packing, as

reflected by ICVF, may be established earlier. Still, it is important to note that several factors

may contribute to these contrasting findings, including limited statistical power to detect

interactions, high within-group variability, and/or imperfect biophysical specificity of the

NODDI measures. Moreover, the observed alterations in axonal density may instead be reflective

of atypical prenatal axonogenesis or lack of early axonal pruning in 22qDel and 22qDup carriers.

This is in line with previous research on 22qDel-derived cortical spheroids exhibiting excessive

prenatal overgrowth of thalamocortical axons(21) and neural progenitors that stay longer in the

cell cycling state(86), which may be smaller in size, as indicated by reduced neurospheres

derived from human induced pluripotent stem cells from 22qDel carriers(87). To speculate, the

gene dosage effect of 22q11.2 genomic region on axonal density observed in our study may be

reflective of a differential effect on gene expression levels of genes that are involved in

axonogenesis, possibly through trans-regulatory mechanisms such as the mediating effect of

FOXP2 gene expression levels on 22qDel thalamic neurons(21). However, to our knowledge, no

studies have utilized cellular models of the reciprocal 22qDup; such future studies are warranted

to determine differential gene dosage effects on axonal development.

Strengths and limitations

The current study includes a unique sample across a wide age-range consisting of both 22qDel

and 22qDup carriers and a large typically developing control sample with multi-shell dMRI data,

allowing for gene dosage and developmental analyses on metrics derived from advanced

neuroimaging techniques. The use of advanced dMRI measures provides novel insight into the

axonal architecture of the white matter microstructure in 22q11.2 CNV carriers, insight that

cannot be achieved through conventional DTI measures(18,51). We also performed several

sensitivity analyses demonstrating the robustness of the results of the group level analyses by

analyzing individuals derived from the same scanner site only and adjusting for brain volumetric

differences. The results revealed largely the same pattern, albeit with some group differences

becoming attenuated due to reduction in sample size. It is also important to note that small

sample sizes—due to the population prevalence of these CNVs—limit our ability to statistically

detect small effects. Despite this, previous studies have found robust differences in white matter

microstructure with smaller sample sizes than our study(20), which remains the largest study on

biophysical white matter measures in 22q11.2 CNV carriers to date. Here, we also utilized

GAMMs to capture nonlinear age effects, account for repeated measures and avoid biased group

comparisons, especially important in instances where sample sizes are modest and age

distributions differ between groups. Although we did not find evidence for altered developmental

trajectories in 22q11.2 CNV carriers, we urge caution when interpreting these results, as the

interaction effects may be more subtle in magnitude than the corresponding main effects and thus

require greater power to detect. Thus, we cannot exclude the possibility of subtly altered

developmental trajectories in NODDI measures among 22q11.2 CNV carriers compared to

controls. However, because robust group differences in white matter microstructure between

22qDel carriers and controls are already evident in early adolescence(20), and neither our

analyses nor a previous multi-site DTI study of 22qDel vs controls(13) show notable age-by-

group interactions, there is currently no clear evidence that these group differences are driven by

large maturational shifts during adolescence. Future studies with larger sample sizes, dense

longitudinal data and methods that examine differences in smoothed confidence intervals(e.g.,

15,86) may shed new light on possible age-dependent differences across the lifespan in 22q11.2

CNV carriers. Finally, despite the advantages of NODDI, it is still subject to the limitations

inherent to dMRI, which is used to infer tissue microstructure by measuring water diffusion

within a voxel of tissue. In addition, although NODDI measures have been shown to reflect their

histological counterparts(53–55), it is important to note that biophysical models are emerging

techniques that remain under active development(51,89–93). Our results also raise new

questions. For instance, it is unclear if the altered axonal density in 22q11.2 CNV carriers is a

consequence of an absolute difference in total neurons in the brain, altered axonal diameter

distribution, or other mechanisms that alter space requirements in white matter tissue. Future

studies utilizing both in-vitro cellular models and post-mortem examinations of 22q11.2 CNV

carriers are warranted.

Conclusions

The results of the current study provide new insights into the underlying neurobiology of the

altered white matter microstructure in 22q11.2 CNV carriers, linking copy numbers at the

22q11.2 genomic locus to altered axonal density. However, we do not find evidence for altered

age-related changes in axonal density or dispersion across childhood and adolescence, possibly

reflecting atypical axonogenesis and/or axonal pruning during early neurodevelopment in

22q11.2 CNV carriers. Future studies are warranted to characterize the morphological features of

white matter axons across fetal development, and to connect these developmental alterations to

neuropsychiatric phenotypes.

Acknowledgments**:**

We thank Sergiu Pasca, M.D., for contributing to the study. This work used computational and

storage services associated with the Hoffman2 Cluster which is operated by the UCLA Office of

Advanced Research Computing's Research Technology Group and servers operated by the USC

Mark and Mary Stevens Neuroimaging and Informatics Institute (INI). Data from the UCLA and

Stanford cohorts were supported by grants from the MCHRI Uytengsu-Hamilton 22q11

Neuropsychiatry Research Award (UH22QEXTFY22-04) and NIH 1R21MH116473-01A1,

R37MH085953, 1U01MH119736-01 (to CEB), R01MH100900 (to JH), and R01MH129858 (to

CEB, PMT, IES, OAA). Research reported in this publication was supported by the National

Institute Of Mental Health of the National Institutes of Health under Award Number

U01MH109589/ U01AG052564 and by funds provided by the McDonnell Center for Systems

Neuroscience at Washington University in St. Louis. The HCP-Development 2.0 Release data

used in this report came from DOI: 10.15154/1520708. The HCP-Aging 2.0 Release data used in

this report came from DOI: 10.15154/1520707.

Conflict of Interest:

OAA is a consultant for Cortechs.ai and Precision Health and has received speaker's honoraria

from Lundbeck, Janssen, Otsuka, Lilly and Sunovion. The remaining authors have no conflicts

of interest to declare.

Supplementary information is available at MP's website.

References

1.Olsen L, Sparsø T, Weinsheimer SM, Dos Santos MBQ, Mazin W, Rosengren A, et al.
Prevalence of rearrangements in the 22q11.2 region and population-based risk of
neuropsychiatric and developmental disorders in a Danish population: a case-cohort study.
Lancet Psychiatry. 2018 July 1;5(7):573–80.

2. Schneider M, Debbané M, Bassett AS, Chow EWC, Fung WLA, van den Bree MBM, et al.
Psychiatric Disorders From Childhood to Adulthood in 22q11.2 Deletion Syndrome: Results
From the International Consortium on Brain and Behavior in 22q11.2 Deletion Syndrome. Am
J Psychiatry. 2014 June;171(6):627–39.

3. Hoeffding LK, Trabjerg BB, Olsen L, Mazin W, Sparsø T, Vangkilde A, et al. Risk of
Psychiatric Disorders Among Individuals With the 22q11.2 Deletion or Duplication: A Danish
Nationwide, Register-Based Study. JAMA Psychiatry. 2017 Mar 1;74(3):282–90.

4. Marshall CR, Howrigan DP, Merico D, Thiruvahindrapuram B, Wu W, Greer DS, et al.
Contribution of copy number variants to schizophrenia from a genome-wide study of 41,321
subjects. Nat Genet. 2017 Jan;49(1):27–35.

5. Emanuel BS, McDonald-McGinn D, Saitta SC, Zackai EH. CHAPTER 2 - The 22q11.2
Deletion Syndrome. Adv Pediatr. 2001 Jan 1;48(1):39–73.

6. Boot E, Óskarsdóttir S, Loo JCY, Crowley TB, Orchanian-Cheff A, Andrade DM, et al.
Updated clinical practice recommendations for managing adults with 22q11.2 deletion
syndrome. Genet Med [Internet]. 2023 Mar 1 [cited 2025 Oct 11];25(3). Available from:
https://www.gimjournal.org/article/S1098-3600(22)01028-0/fulltext

7. McDonald-McGinn DM, Sullivan KE, Marino B, Philip N, Swillen A, Vorstman JAS, et al.
22q11.2 deletion syndrome. Nat Rev Dis Primer. 2015 Nov 19;1(1):15071.

8. Rees E, Kirov G, Sanders A, Walters JTR, Chambert KD, Shi J, et al. Evidence that
duplications of 22q11.2 protect against schizophrenia. Mol Psychiatry. 2014 Jan;19(1):37–40.

9. Owen MJ, Legge SE, Rees E, Walters JTR, O'Donovan MC. Genomic findings in
schizophrenia and their implications. Mol Psychiatry. 2023 Sept;28(9):3638–47.

10.   Ou Z, Berg JS, Yonath H, Enciso VB, Miller DT, Picker J, et al. Microduplications of
22q11.2 are frequently inherited and are associated with variable phenotypes. Genet Med.
2008 Apr 1;10(4):267–77.

11.   Portnoï MF. Microduplication 22q11.2: A new chromosomal syndrome. Eur J Med
Genet. 2009 Mar 1;52(2):88–93.

12.   Lin A, Vajdi A, Kushan-Wells L, Helleman G, Hansen LP, Jonas RK, et al. Reciprocal
Copy Number Variations at 22q11.2 Produce Distinct and Convergent Neurobehavioral
Impairments Relevant for Schizophrenia and Autism Spectrum Disorder. Biol Psychiatry.
2020 Aug 1;88(3):260–72.

13.   Villalón-Reina JE, Martínez K, Qu X, Ching CRK, Nir TM, Kothapalli D, et al. Altered
white matter microstructure in 22q11.2 deletion syndrome: a multisite diffusion tensor
imaging study. Mol Psychiatry. 2020 Nov;25(11):2818–31.

14.   Sun D, Ching CRK, Lin A, Forsyth JK, Kushan L, Vajdi A, et al. Large-scale mapping of
cortical alterations in 22q11.2 deletion syndrome: Convergence with idiopathic psychosis and
effects of deletion size. Mol Psychiatry. 2020 Aug;25(8):1822–34.

15.   Ching CRK, Gutman BA, Sun D, Villalon Reina J, Ragothaman A, Isaev D, et al.
Mapping Subcortical Brain Alterations in 22q11.2 Deletion Syndrome: Effects of Deletion
Size and Convergence With Idiopathic Neuropsychiatric Illness. Am J Psychiatry. 2020 July
1;177(7):589–600.

16.   Schleifer CH, O'Hora KP, Fung H, Xu J, Robinson TA, Wu AS, et al. Effects of gene
dosage and development on subcortical nuclei volumes in individuals with 22q11.2 copy
number variations. Neuropsychopharmacology. 2024 May;49(6):1024–32.

17.   Mori S, Zhang J. Principles of Diffusion Tensor Imaging and Its Applications to Basic
Neuroscience Research. Neuron. 2006 Sept 7;51(5):527–39.

18.   Pierpaoli C, Jezzard P, Basser PJ, Barnett A, Di Chiro G. Diffusion tensor MR imaging
of the human brain. Radiology. 1996 Dec;201(3):637–48.

19.   Beaulieu C. The basis of anisotropic water diffusion in the nervous system - a technical
review. NMR Biomed. 2002;15(7–8):435–55.

20.   Raven EP, Veraart J, Kievit RA, Genc S, Ward IL, Hall J, et al. In vivo evidence of
microstructural hypo-connectivity of brain white matter in 22q11.2 deletion syndrome. Mol
Psychiatry. 2023 July 26;1–11.

21.   Shin D, Kim CN, Ross J, Hennick KM, Wu SR, Paranjape N, et al. Thalamocortical
organoids enable *in vitro* modeling of 22q11.2 microdeletion associated with neuropsychiatric
disorders. Cell Stem Cell. 2024 Mar 7;31(3):421-432.e8.

22.   Wu P, Teot L, Murdoch G, Monaghan-Nichols AP, McFadden K. Neuropathology of
22q11 Deletion Syndrome in an Infant. Pediatr Dev Pathol. 2014 Sept 1;17(5):386–92.

23.   Seitz-Holland J, Lyons M, Kushan L, Lin A, Villalon-Reina JE, Cho KIK, et al.
Opposing white matter microstructure abnormalities in 22q11.2 deletion and duplication
carriers. Transl Psychiatry. 2021 Nov 10;11(1):1–11.

24.   Mukai J, Tamura M, Fénelon K, Rosen AM, Spellman TJ, Kang R, et al. Molecular
Substrates of Altered Axonal Growth and Brain Connectivity in a Mouse Model of
Schizophrenia. Neuron. 2015 May 6;86(3):680–95.

25.   Mukai J, Dhilla A, Drew LJ, Stark KL, Cao L, MacDermott AB, et al. Palmitoylation-
dependent neurodevelopmental deficits in a mouse model of 22q11 microdeletion. Nat
Neurosci. 2008 Nov;11(11):1302–10.

26.   Hiramoto T, Sumiyoshi A, Yamauchi T, Tanigaki K, Shi Q, Kang G, et al. Tbx1, a gene
encoded in 22q11.2 copy number variant, is a link between alterations in fimbria myelination
and cognitive speed in mice. Mol Psychiatry. 2022 Feb;27(2):929–38.

27.   Budel S, Padukkavidana T, Liu BP, Feng Z, Hu F, Johnson S, et al. Genetic Variants of
Nogo-66 Receptor with Possible Association to Schizophrenia Block Myelin Inhibition of
Axon Growth. J Neurosci. 2008 Dec 3;28(49):13161–72.

28.   Fernandez A, Meechan DW, Karpinski BA, Paronett EM, Bryan CA, Rutz HL, et al.
Mitochondrial Dysfunction Leads to Cortical Under-Connectivity and Cognitive Impairment.
Neuron. 2019 June 19;102(6):1127-1142.e3.

29.   Khan TA, Revah O, Gordon A, Yoon SJ, Krawisz AK, Goold C, et al. Neuronal defects
in a human cellular model of 22q11.2 deletion syndrome. Nat Med. 2020 Dec;26(12):1888–
98.

30.   Stark KL, Xu B, Bagchi A, Lai WS, Liu H, Hsu R, et al. Altered brain microRNA
biogenesis contributes to phenotypic deficits in a 22q11-deletion mouse model. Nat Genet.
2008 June;40(6):751–60.

31.   Lin A, Forsyth JK, Hoftman GD, Kushan-Wells L, Jalbrzikowski M, Dokuru D, et al.
Transcriptomic profiling of whole blood in 22q11.2 reciprocal copy number variants reveals
that cell proportion highly impacts gene expression. Brain Behav Immun - Health. 2021 Dec
1;18:100386.

32.   Nehme R, Pietiläinen O, Artomov M, Tegtmeyer M, Valakh V, Lehtonen L, et al. The
22q11.2 region regulates presynaptic gene-products linked to schizophrenia. Nat Commun.
2022 June 27;13(1):3690.

33.   Huang H, Zhang J, Wakana S, Zhang W, Ren T, Richards LJ, et al. White and gray
matter development in human fetal, newborn and pediatric brains. NeuroImage. 2006 Oct
15;33(1):27–38.

34.   Kochunov P, Williamson DE, Lancaster J, Fox P, Cornell J, Blangero J, et al. Fractional
anisotropy of water diffusion in cerebral white matter across the lifespan. Neurobiol Aging.
2012 Jan 1;33(1):9–20.

35.   Wilson S, Pietsch M, Cordero-Grande L, Price AN, Hutter J, Xiao J, et al. Development
of human white matter pathways in utero over the second and third trimester. Proc Natl Acad
Sci U S A. 2021 May 18;118(20):e2023598118.

36.   Lebel C, Beaulieu C. Longitudinal Development of Human Brain Wiring Continues from
Childhood into Adulthood. J Neurosci. 2011 July 27;31(30):10937–47.

37.   Huang H, Xue R, Zhang J, Ren T, Richards LJ, Yarowsky P, et al. Anatomical
Characterization of Human Fetal Brain Development with Diffusion Tensor Magnetic
Resonance Imaging. J Neurosci. 2009 Apr 1;29(13):4263–73.

38.   Vasung L, Raguz M, Kostovic I, Takahashi E. Spatiotemporal Relationship of Brain
Pathways during Human Fetal Development Using High-Angular Resolution Diffusion MR
Imaging and Histology. Front Neurosci [Internet]. 2017 July 11 [cited 2024 Dec 16];11.
Available from:
https://www.frontiersin.org/journals/neuroscience/articles/10.3389/fnins.2017.00348/full

39.   Horgos B, Mecea M, Boer A, Szabo B, Buruiana A, Stamatian F, et al. White Matter
Dissection of the Fetal Brain. Front Neuroanat. 2020 Sept 25;14:584266.

40.   Rakic P, Yakovlev PI. Development of the corpus callosum and cavum septi in man. J
Comp Neurol. 1968;132(1):45–72.

41.   Ren T, Anderson A, Shen WB, Huang H, Plachez C, Zhang J, et al. Imaging, anatomical,
and molecular analysis of callosal formation in the developing human fetal brain. Anat Rec A
Discov Mol Cell Evol Biol. 2006;288A(2):191–204.

42.   Ouyang M, Dubois J, Yu Q, Mukherjee P, Huang H. Delineation of early brain
development from fetuses to infants with diffusion MRI and beyond. NeuroImage. 2019 Jan
15;185:836–50.

43.   Takahashi E, Folkerth RD, Galaburda AM, Grant PE. Emerging Cerebral Connectivity in
the Human Fetal Brain: An MR Tractography Study. Cereb Cortex. 2012 Feb 1;22(2):455–64.

44.   Kostović I, Jovanov-Milošević N. The development of cerebral connections during the
first 20–45 weeks' gestation. Semin Fetal Neonatal Med. 2006 Dec 1;11(6):415–22.

45.   LaMantia AS, Rakic P. Axon overproduction and elimination in the corpus callosum of
the developing rhesus monkey. J Neurosci. 1990 July 1;10(7):2156–75.

46.   LaMantia AS, Rakic P. Axon overproduction and elimination in the anterior commissure
of the developing rhesus monkey. J Comp Neurol. 1994 Feb 15;340(3):328–36.

47.   Paus T. Growth of white matter in the adolescent brain: Myelin or axon? Brain Cogn.
2010 Feb 1;72(1):26–35.

48.   Benes FM. Myelination of Cortical-hippocampal Relays During Late Adolescence.
Schizophr Bull. 1989 Jan 1;15(4):585–93.

49.   Kulikova S, Hertz-Pannier L, Dehaene-Lambertz G, Buzmakov A, Poupon C, Dubois J.
Multi-parametric evaluation of the white matter maturation. Brain Struct Funct. 2015 Nov
1;220(6):3657–72.

50.   Jones DK, Knösche TR, Turner R. White matter integrity, fiber count, and other fallacies:
The do's and don'ts of diffusion MRI. NeuroImage. 2013 June 1;73:239–54.

51.   Zhang H, Schneider T, Wheeler-Kingshott CA, Alexander DC. NODDI: Practical in vivo
neurite orientation dispersion and density imaging of the human brain. NeuroImage. 2012 July
16;61(4):1000–16.

52.   Lee HH, Yaros K, Veraart J, Pathan JL, Liang FX, Kim SG, et al. Along-axon diameter
635   variation and axonal orientation dispersion revealed with 3D electron microscopy:
636   implications for quantifying brain white matter microstructure with histology and diffusion
637   MRI. Brain Struct Funct. 2019 May;224(4):1469–88.

638   53.   Grussu F, Schneider T, Tur C, Yates RL, Tachrount M, Ianuș A, et al. Neurite dispersion:
639   a new marker of multiple sclerosis spinal cord pathology? Ann Clin Transl Neurol.
640   2017;4(9):663–79.

641   54.   Jespersen SN, Bjarkam CR, Nyengaard JR, Chakravarty MM, Hansen B, Vosegaard T, et
642   al. Neurite density from magnetic resonance diffusion measurements at ultrahigh field:
643   Comparison with light microscopy and electron microscopy. NeuroImage. 2010 Jan
644   1;49(1):205–16.

645   55.   Sato K, Kerever A, Kamagata K, Tsuruta K, Irie R, Tagawa K, et al. Understanding
646   microstructure of the brain by comparison of neurite orientation dispersion and density
647   imaging (NODDI) with transparent mouse brain. Acta Radiol Open. 2017 Apr
648   1;6(4):2058460117703816.

649   56.   Kraguljac NV, Guerreri M, Strickland MJ, Zhang H. Neurite Orientation Dispersion and
650   Density Imaging in Psychiatric Disorders: A Systematic Literature Review and a Technical
651   Note. Biol Psychiatry Glob Open Sci. 2022 Jan 21;3(1):10–21.

652   57.   Genc S, Malpas CB, Holland SK, Beare R, Silk TJ. Neurite density index is sensitive to
653   age related differences in the developing brain. NeuroImage. 2017 Mar 1;148:373–80.

654   58.   Genc S, Raven EP, Drakesmith M, Blakemore SJ, Jones DK. Novel insights into axon
655   diameter and myelin content in late childhood and adolescence. Cereb Cortex N Y NY. 2023
656   Jan 4;33(10):6435–48.

657   59.   Fan Q, Tian Q, Ohringer NA, Nummenmaa A, Witzel T, Tobyne SM, et al. Age-related
658   alterations in axonal microstructure in the corpus callosum measured by high-gradient
659   diffusion MRI. NeuroImage. 2019 May 1;191:325–36.

660   60.   Benes FM, Turtle M, Khan Y, Farol P. Myelination of a Key Relay Zone in the
661   Hippocampal Formation Occurs in the Human Brain During Childhood, Adolescence, and
662   Adulthood. Arch Gen Psychiatry. 1994 June 1;51(6):477–84.

663   61.   Somerville LH, Bookheimer SY, Buckner RL, Burgess GC, Curtiss SW, Dapretto M, et
664   al. The Lifespan Human Connectome Project in Development: A large-scale study of brain
665   connectivity development in 5–21 year olds. NeuroImage. 2018 Aug 22;183:456.

666   62.   Bookheimer SY, Salat DH, Terpstra M, Ances BM, Barch DM, Buckner RL, et al. The
667   Lifespan Human Connectome Project in Aging: An overview. NeuroImage. 2019 Jan
668   15;185:335–48.

669   63.   Harms MP, Somerville LH, Ances BM, Andersson J, Barch DM, Bastiani M, et al.
670   Extending the Human Connectome Project across ages: Imaging protocols for the Lifespan
671   Development and Aging projects. NeuroImage. 2018 Dec;183:972–84.

672   64.   Garyfallidis E, Brett M, Amirbekian B, Rokem A, Van Der Walt S, Descoteaux M, et al.
673   Dipy, a library for the analysis of diffusion MRI data. Front Neuroinformatics [Internet]. 2014
674   [cited 2023 Aug 24];8. Available from:
675   https://www.frontiersin.org/articles/10.3389/fninf.2014.00008

676   65.   Manjón JV, Coupé P, Concha L, Buades A, Collins DL, Robles M. Diffusion Weighted
677   Image Denoising Using Overcomplete Local PCA. PLOS ONE. 2013 Sept 3;8(9):e73021.

678   66.   Andersson JLR, Skare S, Ashburner J. How to correct susceptibility distortions in spin-
679   echo echo-planar images: application to diffusion tensor imaging. NeuroImage. 2003
680   Oct;20(2):870–88.

681   67.   Andersson JLR, Sotiropoulos SN. An integrated approach to correction for off-resonance
682   effects and subject movement in diffusion MR imaging. NeuroImage. 2016 Jan 15;125:1063–
683   78.

684   68.   Jahanshad N, Kochunov P, Sprooten E, Mandl RC, Nichols TE, Almassy L, et al. Multi-
685   site genetic analysis of diffusion images and voxelwise heritability analysis: A pilot project of
686   the ENIGMA–DTI working group. NeuroImage. 2013 Nov 1;81:455–69.

687   69.   Mori S, Oishi K, Jiang H, Jiang L, Li X, Akhter K, et al. Stereotaxic white matter atlas
688   based on diffusion tensor imaging in an ICBM template. NeuroImage. 2008 Apr 1;40(2):570–
689   82.

690   70.   Beer JC, Tustison NJ, Cook PA, Davatzikos C, Sheline YI, Shinohara RT, et al.
691   Longitudinal ComBat: A method for harmonizing longitudinal multi-scanner imaging data.
692   NeuroImage. 2020 Oct 15;220:117129.

693   71.   Pakkenberg B, Pelvig D, Marner L, Bundgaard MJ, Gundersen HJG, Nyengaard JR, et al.
694   Aging and the human neocortex. Exp Gerontol. 2003 Jan 1;38(1):95–9.

695   72.   Wood S. Generalized additive models: an introduction with R. Chapman and Hall/CRC;
696   2017.

697   73.   Garic D, Yeh FC, Graziano P, Dick AS. In vivo restricted diffusion imaging (RDI) is
698   sensitive to differences in axonal density in typical children and adults. Brain Struct Funct.
699   2021 Nov 1;226(8):2689–705.

700   74.   Aboitiz F, Scheibel AB, Fisher RS, Zaidel E. Fiber composition of the human corpus
701   callosum. Brain Res. 1992 Dec 11;598(1):143–53.

702   75.   Liewald D, Miller R, Logothetis N, Wagner HJ, Schüz A. Distribution of axon diameters
703   in cortical white matter: an electron-microscopic study on three human brains and a macaque.
704   Biol Cybern. 2014 Oct 1;108(5):541–57.

705   76.   Meechan DW, Tucker ES, Maynard TM, LaMantia AS. Diminished dosage of 22q11
706   genes disrupts neurogenesis and cortical development in a mouse model of 22q11
707   deletion/DiGeorge syndrome. Proc Natl Acad Sci. 2009 Sept 22;106(38):16434–45.

708   77.   Lin A, Ching CRK, Vajdi A, Sun D, Jonas RK, Jalbrzikowski M, et al. Mapping 22q11.2
709   Gene Dosage Effects on Brain Morphometry. J Neurosci. 2017 June 28;37(26):6183–99.

710   78.   Hegarty JP, Monterrey JC, Tian Q, Cleveland SC, Gong X, Phillips JM, et al. A Twin
711   Study of Altered White Matter Heritability in Youth With Autism Spectrum Disorder. J Am
712   Acad Child Adolesc Psychiatry. 2024 Jan;63(1):65–79.

713   79.   Saaybi S, AlArab N, Hannoun S, Saade M, Tutunji R, Zeeni C, et al. Pre- and Post-
714   therapy Assessment of Clinical Outcomes and White Matter Integrity in Autism Spectrum
715   Disorder: Pilot Study. Front Neurol. 2019 Aug 13;10:877.

716   80.   Singh T, Poterba T, Curtis D, Akil H, Al Eissa M, Barchas JD, et al. Rare coding variants
717   in ten genes confer substantial risk for schizophrenia. Nature. 2022 Apr;604(7906):509–16.

718   81.   Schleifer C, Lin A, Kushan L, Ji JL, Yang G, Bearden CE, et al. Dissociable Disruptions
719   in Thalamic and Hippocampal Resting-State Functional Connectivity in Youth with 22q11.2
720   Deletions. J Neurosci. 2019 Feb 13;39(7):1301–19.

721   82.   Mancini V, Zöller D, Schneider M, Schaer M, Eliez S. Abnormal Development and
722   Dysconnectivity of Distinct Thalamic Nuclei in Patients With 22q11.2 Deletion Syndrome
723   Experiencing Auditory Hallucinations. Biol Psychiatry Cogn Neurosci Neuroimaging. 2020
724   Sept 1;5(9):875–90.

725   83.   Moreau CA, Kumar K, Harvey A, Huguet G, Urchs SGW, Schultz LM, et al. Brain
726   functional connectivity mirrors genetic pleiotropy in psychiatric conditions. Brain. 2022 Sept
727   5;146(4):1686–96.

728   84.   Steullet P. Thalamus-related anomalies as candidate mechanism-based biomarkers for
729   psychosis. Schizophr Res. 2020 Dec 1;226:147–57.

730   85.   Alvino FG, Gini S, Minetti A, Pagani M, Sastre-Yagüe D, Barsotti N, et al. Synaptic-
731   dependent developmental dysconnectivity in 22q11.2 deletion syndrome. Sci Adv. 2025 Mar
732   12;11(11):eadq2807.

733   86.   Rao SB, Brundu F, Chen Y, Sun Y, Zhu H, Shprintzen RJ, et al. Aberrant pace of cortical
734   neuron development in brain organoids from patients with 22q11.2 deletion syndrome and
735   schizophrenia [Internet]. bioRxiv; 2023 [cited 2024 Feb 5]. p. 2023.10.04.557612. Available
736   from: https://www.biorxiv.org/content/10.1101/2023.10.04.557612v1

737   87.   Toyoshima M, Akamatsu W, Okada Y, Ohnishi T, Balan S, Hisano Y, et al. Analysis of
738   induced pluripotent stem cells carrying 22q11.2 deletion. Transl Psychiatry. 2016
739   Nov;6(11):e934–e934.

88.   Jalbrzikowski M, Lin A, Vajdi A, Grigoryan V, Kushan L, Ching CRK, et al.
Longitudinal trajectories of cortical development in 22q11.2 copy number variants and
typically developing controls. Mol Psychiatry. 2022 Oct;27(10):4181–90.

89.   Jelescu IO, Budde MD. Design and Validation of Diffusion MRI Models of White
Matter. Front Phys [Internet]. 2017 Nov 28 [cited 2025 Nov 18];5. Available from:
https://www.frontiersin.org/journals/physics/articles/10.3389/fphy.2017.00061/full

90.   Guerrero JM, Adluru N, Bendlin BB, Goldsmith HH, Schaefer SM, Davidson RJ, et al.
Optimizing the intrinsic parallel diffusivity in NODDI: An extensive empirical evaluation.
PLOS ONE. 2019 Sept 25;14(9):e0217118.

91.   Jelescu IO, Veraart J, Fieremans E, Novikov DS. Degeneracy in model parameter
estimation for multi-compartmental diffusion in neuronal tissue. NMR Biomed.
2016;29(1):33–47.

92.   Lampinen B, Szczepankiewicz F, Mårtensson J, van Westen D, Sundgren PC, Nilsson M.
Neurite density imaging versus imaging of microscopic anisotropy in diffusion MRI: A model
comparison using spherical tensor encoding. NeuroImage. 2017 Feb 15;147:517–31.

93.   Kamiya K, Hori M, Aoki S. NODDI in clinical research. J Neurosci Methods. 2020 Dec
1;346:108908.

Figure 1. Point estimates of group differences in intracellular volume fraction (ICVF),
orientation dispersion index (ODI) and isotropic volume fraction (ISO). The control group and
the 22qDel group are used as reference groups for the group comparisons. Gene dosage effects
are based on a continuous variable of copy number (i.e., 22qDel = 1, control = 2, and 22qDup =
3). E.g., Significant negative gene dosage effects indicate that an increased number of copies of
genes in the 22q11.2 genomic locus is associated with lower values of the diffusion measures.
ACR = anterior corona radiata, ALIC = anterior limb of internal capsule, BCC = body of corpus
callosum, CC = corpus callosum, CGC = cingulum cingulate gyrus, CGH = cingulum
hippocampus, CR = corona radiata, CST = corticospinal tract, EC = extreme/external capsule,
FX = fornix. FXST = fornix (crus)/stria terminalis, GCC = genu of corpus callosum, IC =
internal capsule, PCR = posterior corona radiata, PLIC = posterior limb of the internal capsule,
PTR = posterior thalamic radiation, RLIC = retrolenticular part of the internal capsule, SCC =
splenium of corpus callosum, SCR = superior corona radiata, SFO = superior fronto-occipital
fasciculus, SLF = superior longitudinal fasciculus, SS = sagittal stratum, TAP = tapetum of the
corpus callosum, UNC = uncinate fasciculus.

Figure 2. Developmental trajectories of intracellular volume fraction (ICVF) in global and
regional white matter microstructure for 22q11.2 deletion carriers (red), controls (black) and
22q11.2 duplication carriers (blue). ACR = anterior corona radiata, ALIC = anterior limb of
internal capsule, BCC = body of corpus callosum, CC = corpus callosum, CGC = cingulum
cingulate gyrus, CGH = cingulum hippocampus, CR = corona radiata, CST = corticospinal tract,
EC = extreme/external capsule, FX = fornix. FXST = fornix (crus)/stria terminalis, GCC = genu
of corpus callosum, IC = internal capsule, PCR = posterior corona radiata, PLIC = posterior limb
of the internal capsule, PTR = posterior thalamic radiation, RLIC = retrolenticular part of the
internal capsule, SCC = splenium of corpus callosum, SCR = superior corona radiata, SFO =
superior fronto-occipital fasciculus, SLF = superior longitudinal fasciculus, SS = sagittal stratum,
TAP = tapetum of the corpus callosum, UNC = uncinate fasciculus.

Figure 3. Developmental trajectories of orientation dispersion index (ODI) in global and
regional white matter microstructure for 22q11.2 deletion carriers (red), controls (black) and
22q11.2 duplication carriers (blue). ACR = anterior corona radiata, ALIC = anterior limb of
internal capsule, BCC = body of corpus callosum, CC = corpus callosum, CGC = cingulum
cingulate gyrus, CGH = cingulum hippocampus, CR = corona radiata, CST = corticospinal tract,
EC = extreme/external capsule, FX = fornix. FXST = fornix (crus)/stria terminalis, GCC = genu
of corpus callosum, IC = internal capsule, PCR = posterior corona radiata, PLIC = posterior limb
of the internal capsule, PTR = posterior thalamic radiation, RLIC = retrolenticular part of the
internal capsule, SCC = splenium of corpus callosum, SCR = superior corona radiata, SFO =
superior fronto-occipital fasciculus, SLF = superior longitudinal fasciculus, SS = sagittal stratum,
TAP = tapetum of the corpus callosum, UNC = uncinate fasciculus.

Figure 4. Developmental trajectories of isotropic volume fraction (ISO) in global and regional
white matter microstructure for 22q11.2 deletion carriers (red), controls (black) and 22q11.2
duplication carriers (blue). ACR = anterior corona radiata, ALIC = anterior limb of internal
capsule, BCC = body of corpus callosum, CC = corpus callosum, CGC = cingulum cingulate
gyrus, CGH = cingulum hippocampus, CR = corona radiata, CST = corticospinal tract, EC =
extreme/external capsule, FX = fornix. FXST = fornix (crus)/stria terminalis, GCC = genu of
corpus callosum, IC = internal capsule, PCR = posterior corona radiata, PLIC = posterior limb of
the internal capsule, PTR = posterior thalamic radiation, RLIC = retrolenticular part of the
internal capsule, SCC = splenium of corpus callosum, SCR = superior corona radiata, SFO =
superior fronto-occipital fasciculus, SLF = superior longitudinal fasciculus, SS = sagittal stratum,
TAP = tapetum of the corpus callosum, UNC = uncinate fasciculus.

Table 1. Participant baseline demographics

	Control (N=890)	22q11.2 Deletion Carriers (N=50)	22q11.2 Duplication Carriers (N=24)	P-value
Scans				
n	901	69	34	-
Age				
Mean (SD)	21.9 (13.5)	20.7 (9.4)	21.6 (14.1)	.538
Median [Min, Max]	15.4 [7.8, 51.1]	20.0 [7.4, 51.1]	14.5 [8.3, 49.4]	
Sex				
Female	485 (54.5%)	32 (64.0%)	13 (54.2%)	.420

Notes. Kruskal-Wallis rank sum test and Pearson Chi-squared test were used to compare group differences in baseline age and sex characteristics, respectively.