1   Gene dosage effects of 22q11.2 copy number variants on in-vivo measures of white matter
2   axonal density and dispersion
3

Rune Boen[1], Julio E. Villalón-Reina[2], Leila Kushan[1], Kathleen P. O'Hora[1], Hoki Fung[1], Nadine

Parker[3], Ibrahim A. Akkouh[4], Dag Alnæs[3], Ruth O'Hara[5], Matthew John Marzelli[5], Lara Foland-

Ross[5], Christina French Chick[5], Isabelle Cotto[5], Allan L. Reiss[5], Joachim Hallmayer[5], Paul M.

Thompson[2], Ole A. Andreassen[3], Ida E. Sønderby[3,4,6], Carrie E. Bearden[1,7]

[1] Semel Institute for Neuroscience and Human Behavior, University of California Los Angeles,

Los Angeles, CA, USA.

[2] Imaging Genetics Center, Mark and Mary Stevens Neuroimaging and Informatics Institute,

University of Southern California, Los Angeles, CA, USA

[3] Centre for Precision Psychiatry, Division of Mental Health and Addiction, Oslo University

Hospital & University of Oslo, Oslo, Norway

[4] Department of Medical Genetics, Oslo University Hospital, Oslo, Norway

[5] Stanford University School of Medicine, Stanford, CA, USA

[6] KG Jebsen Centre for Neurodevelopmental Disorders, University of Oslo, Oslo, Norway.

[7] Department of Psychology, University of California Los Angeles, Los Angeles, CA, USA.

Correspondence: Carrie E. Bearden
A7-460 Semel Institute, Los Angeles, CA 90095, USA
cbearden@mednet.ucla.edu
23

24   Keywords: 22q11.2 copy number variant, NODDI, white matter, axonal density, axonal
25   dispersion

27   Abstract

22q11.2 deletion (22qDel) and duplication (22qDup) carriers have an increased risk of

neurodevelopmental disorders and exhibit altered brain structure, including white matter

microstructure. However, the underlying cellular architecture and age-related changes

contributing to these white matter alterations remain poorly understood. Neurite orientation

dispersion and density imaging (NODDI) was used on mixed cross-sectional and longitudinal

data to examine group differences and age-related trajectories in measures of axonal density (i.e.,

intracellular volume fraction; ICVF), axonal orientation (orientation dispersion index; ODI) and

free water diffusion (isotropic volume fraction; ISO) in 50 22qDel (n scans = 69, mean age =

21.7, age range = 7.4-51.1, 65.2% female) and 24 22qDup (n scans = 34, mean age = 23.3, age

range = 8.3-49.4, 55.0% female) carriers, and 890 controls (n scans = 901, mean age = 21.9, age

range = 7.8-51.1, 54.5%). The results showed widespread gene dosage effects, with higher ICVF

in 22qDel and lower ICVF in 22qDup compared to controls, and region-specific effects of the

22qDel and 22qDup on ODI and ISO measures. However, 22qDel and 22qDup carriers did not

exhibit an altered age-related trajectory relative to controls. Observed differences in ICVF

suggest higher white matter axonal density in 22qDel and lower axonal density in 22qDup

compared to controls. Conversely, differences in ODI are highly localized, indicating region-

specific effects on axonal dispersion in white matter. We do not find evidence for altered

developmental trajectories of axonal density or dispersion among 22q11.2 CNV carriers,

suggesting stable disruptions to neurodevelopmental events before childhood.

48   Introduction

22q11.2 copy number variants (CNVs) are rare, recurrent structural genetic variants that have

profound impacts on neurodevelopment and confer increased risk for several developmental

neuropsychiatric disorders, including autism spectrum disorder (ASD) and attention

deficit/hyperactivity disorder (ADHD)(1–4) The 22q11.2 deletion (22qDel) is estimated to occur

in one in ~3700 live births(1) and confers a markedly increased risk of psychosis(1–4).The

22q11.2 duplication is estimated to occur in one in ~1600 live births(1) and, conversely, confers

a putative reduced risk of psychosis(4–6). Thus, in contrast to research on individuals with

behaviorally defined psychiatric disorders, studying a molecularly confirmed CNV like the

22q11.2 locus provides a genetically traceable and hypothesis-driven probe of brain alterations in

individuals with a high risk of developing brain disorders. Moreover, reciprocal CNVs (i.e., gain

and loss of genomic material at the same locus) allows for investigation of converging and

diverging neurodevelopmental pathways of brain alterations by examining gene dosage effects

on brain phenotypes.

22q11.2 CNVs have been associated with medium to large differences in several measures of

brain structure, including cortical thickness, surface area, subcortical volume and white matter

microstructure(7–10), indicating altered cellular architecture. However, the complex neuronal

composition and dispersion of the cortical mantle of the brain makes it challenging to study the

cellular architecture in MRI-derived measures of grey matter, such as cortical thickness. In

contrast, the white matter consists of predominantly axons, which can be probed using diffusion

MRI (dMRI) in vivo. As water diffusion is restricted or hindered by biological tissue, dMRI can

be used to generate measures of white matter microstructure that is influenced by the cellular

architecture along white matter tracts(11–13). For instance, water molecules in an environment

of dense and coherent white matter axons will diffuse along the primary direction of the axons,

which has been commonly measured by diffusion tensor imaging (DTI)-derived metrics such as

fractional anisotropy (FA, i.e., standardized value of anisotropic diffusion) and mean diffusivity

(i.e., measure of diffusion in all directions).

A previous consortium-based DTI-study of 22qDel carriers reported an overall pattern of higher

fractional anisotropy (FA) and lower diffusivity across most white matter tracts in 22qDel

carriers compared to controls(7), which could reflect an increase in the cumulative cellular

membrane circumference, increase in smaller tortuous axons(7), and/or higher density of axons

with disproportionately small diameters(7, 14) compared to controls. Indeed, other studies have

suggested other lines of evidence for increased axonal density in 22qDel carriers, specifically

excessive prenatal overgrowth of thalamocortical axons in a cellular model of the 22qDel(15)

and a higher frequency of deep layer cortical neurons in a post-mortem examination of a 3-

month-old infant with 22qDel(16), suggesting increased axonal density in 22qDel carriers

compared to controls. In contrast to 22qDel, only one prior small dMRI study has included

22qDup carriers, which found lower FA and higher diffusivity in white matter in 22qDup

carriers compared to controls(17). This opposing pattern in 22qDel and 22qDup carriers suggests

the existence of a gene dosage effect, reflecting divergent effects of the 22q11.2 CNV on the

cellular architecture of white matter tracts. Indeed, the 22q11.2 locus is a hotspot for genes

critical to white matter development, including axonal morphology and function in white matter

such as regulation of axonal diameter, growth and branching (e.g., *ZDHHC8*(18,19)*, TBX1*(20)*,*

RTN4R/Nogo-66(21)*, TXNRD2*(22)*),* as well as dendritic morphology and neuronal excitability

(e.g., DGCR8(23,24)). In addition, both 22qDel and 22qDup affect expression of genes outside

the 22q11.2 locus(15,23,25,26), which can include genes that influence axonal morphology(15).

However, to our knowledge, no previous studies have directly compared axonal density in white

matter tracts between 22qDel and 22qDup carriers to determine if a gene dosage effect exists.

Moreover, such effects may be region-specific due to the unique developmental trajectories of

white matter fibers. For example, limbic, commissural and projection fibers develop and mature

earlier than association fibers(27–30). Indeed, it has been postulated that the nature of the white

matter alterations in 22qDel carriers may differ between commissural and association fibers(7),

potentially influenced by the maturational patterns of these fiber tracts. Specifically, the presence

of commissural fibers are observed in the first trimester to second trimester(31–36), whereas the

emergence of association fibers are found later during the second trimester(32,33,36,37). The

number of axons peak around birth for commissural fibers(38,39), followed by axonal

elimination(39,40), axon growth and myelination from birth through adolescence (36,41,42). The

association fibers are the least mature fibers during infancy(43) and are the last to mature, as

indicated by developmental changes in DTI measures well into adulthood(30). Thus, it is

plausible that the 22q11.2 CNV may have maturation-specific effects on commissural and
association tracts.

Conventional DTI measures are inherently nonspecific to the underlying white matter cellular
architecture(12,44,45) and can be influenced by alterations to axonal density and/or dispersion
that can be driven by early (e.g., axonal proliferation or axonal pruning) or later (e.g., axonal
diameter expansion) neurodevelopmental events. In contrast, neurite orientation dispersion and
density imaging (NODDI) is a dMRI method that is better suited to test the hypothesis of altered
axonal density in 22q11.2 CNV carriers, as it provides measures of the biophysical properties of
white matter tracts, including axonal density (i.e., intracellular volume fraction, ICVF), axonal
dispersion (i.e., orientation dispersion index, ODI), and free water diffusion/cerebrospinal fluid
(i.e., isotropic water diffusion, ISO). Moreover, NODDI-derived measures have been found to
correlate with their histological counterparts(46–48) and to be altered in several neuropsychiatric
disorders, including psychosis and ASD(49). Thus, examining NODDI measures in 22q11.2
CNV carriers may be informative for detecting converging neurobiological patterns with
idiopathic developmental neuropsychiatric disorders. In addition, ICVF has been found to be
more sensitive to age-related differences in white matter microstructure compared to DTI-
derived metrics during development(50). As such, this metric can yield novel insight into age-
specific neurobiological events that underlie changes in white matter microstructure(51–53),
such as prenatal proliferation of axons(38,39) and prolonged periods of axonal diameter
expansion across childhood and adolescence(51). Thus, by mapping the neurodevelopmental
trajectories of the axonal architecture across white matter regions, we can obtain important
insight into the maturational mechanisms that underlie altered white matter observed in 22qDel
and 22qDup carriers.

In the current study, we aim to characterize the effects of 22q11.2 CNVs on axonal density and
axonal dispersion in white matter regions. We hypothesize gene-dosage effects on axonal density
in white matter regions such that 22qDel carriers will exhibit higher and 22qDup carriers will
exhibit lower axonal density, respectively. We also aim to investigate spatiotemporal effects of
the 22q11.2 CNV on axonal architecture by examining age-related trajectories using mixed
cross-sectional and longitudinal data, and by leveraging a large dataset of typically developing

controls from the Human Connectome Project (HCP)(54,55). We reasoned that alterations to

prenatal neurodevelopmental processes will manifest as robust group differences in NODDI

measures, whereas alterations to later neurodevelopmental processes will be reflected by altered

age-related trajectories in NODDI measures across childhood and adolescence.

<div align="center">Methods</div>

Participants

A total of 964 participants with 1,004 scans were included in the analysis, including 50 22qDel

carriers, 24 22qDup carriers and 890 age and sex matched typically developing controls.

Kruskal-Wallis rank sum test and Pearson Chi-squared test were used to compare group

differences in baseline age and sex characteristics, respectively (Table 1). The 22qDel carriers

were recruited from either the University of California, Los Angeles (UCLA) or Stanford

University, whereas the 22qDup carriers were recruited from UCLA only. The control group

included unrelated healthy control participants recruited from UCLA and Stanford University,

and healthy control participants derived from the Human Connectome Project (HCP, including

HCP-Development(54) and HCP-Aging(55) to establish robust developmental trajectories from

childhood to adulthood (see sTable 1-2 for baseline characteristics across scanner sites).

Diffusion MRI acquisition, preprocessing and quality control

Multi-shell dMRI was acquired using a 3T Siemens Prisma scanner with the following

parameters: TE = 89.2ms, TR = 3230ms, flip angle = 78°, slice thickness = 1.5mm, voxel size =

1.5mm isotropic obtained with 7 b = 0s/mm^2 images, and 3 b = 200s/mm^2, 6 b = 500s/mm^2, 46 b

= 1500s/mm^2 and 46 b = 3000s/mm^2 diffusion weighted volumes. The scans were acquired with

both anterior to posterior and posterior to anterior phase encoding directions, resulting in a total

of 216 volumes, following the HCP acquisition protocol(56). Manual inspections and statistical

analyses were conducted to exclude participants with excessive head movement (see

supplementary note 1 for details). We also included dMRI data derived from the HCP-D

(https://www.humanconnectome.org/study/hcp-lifespan-development) and HCP-A

(https://www.humanconnectome.org/study/hcp-lifespan-aging) as described elsewhere(54,55).

All the raw dMRIs were denoised with LPCA and corrected for Gibbs ringing artifacts using

DiPy (57,58); susceptibility induced distortions, head movement, and eddy currents using

TOPUP and EDDY in FSL (59,60). Afterwards, we ran the diffusion preprocessing pipeline

from HCP (https://github.com/Washington-University/HCPpipelines). NODDI was fitted using

the dMIPY package (https://github.com/AthenaEPI/dmipy). The NODDI derived scalar maps

were registered to the ENIGMA-DTI template(61) and average skeletonized ROI-based

measures were calculated based on the Johns Hopkins University White Matter atlas(62)

following the ENIGMA-DTI protocol (https://enigma.ini.usc.edu/protocols/dti-protocols/)(61),

including anterior corona radiata (ACR), anterior limb of internal capsule (ALIC), body of

corpus callosum (BCC), corpus callosum (CC), cingulum cingulate gyrus (CGC), cingulum

hippocampus (CGH), corona radiata (CR), corticospinal tract (CST), extreme/external capsule

(EC), fornix (FX) , fornix (crus)/stria terminalis (FXST), genu of corpus callosum (GCC),

internal capsule (IC), posterior corona radiata (PCR), posterior limb of the internal capsule

(PLIC), posterior thalamic radiation (PTR), retrolenticular part of the internal capsule (RLIC),

splenium of corpus callosum (SCC), superior corona radiata (SCR), superior fronto-occipital

fasciculus (SFO), superior longitudinal fasciculus (SLF), sagittal stratum (SS), tapetum of the

corpus callosum (TAP), uncinate fasciculus (UNC) (see sFigure 1 for a brain map). Due to

repeated measures across scanner sites, we harmonized all of the NODDI measures using

longCombat(63). Due to the focus on white matter regions, which mostly consist of myelinated

axons(64), we define ICVF as a measure of axonal density, ODI as a measure of axonal

dispersion and ISO as a measure of free water diffusivity(45).

Statistical analyses

Group comparisons

We used generalized additive mixed models (GAMM) in R(65) to test for group differences in

the NODDI measures (i.e., ICVF, ISO and ODI) across 27 variables (24 regions of interest

(ROIs) and 3 average measures). Here, we compared 22qDel carriers to controls, 22qDup

carriers to controls, and 22qDel carriers to 22qDup carriers. In addition, we ran gene dosage

analyses to test for associations between the number of copies of the 22q11.2 genomic locus and

the NODDI measures by treating copy number as a continuous variable (22qDel = 1, control = 2,

and 22qDup = 3). All group analyses were adjusted for the smoothed age effect using cubic

regression splines, sex, and repeated measures through the inclusion of participant ID as random

intercept. All GAMM analyses were fitted using default arguments, where k = -1 is set for the

smoothed age terms and the use of restricted maximum likelihood when fitting all models. The

standardized beta is reported and used as a measure of effect size. To account for multiple

testing, we applied FDR correction across 75 (i.e., dependent variables) * 4 (i.e., three group

analyses and one gene dosage analysis) = 300 comparisons. Finally, we conducted several

sensitivity analyses to test the robustness of the results by including 1) participants from the

same scanner site only, and by adjusting for 2) cerebral white matter volume, 3) intracranial

volume or 4) cerebrospinal fluid volume (see supplementary note 2 for details).

Developmental trajectories

To examine the developmental trajectories of the NODDI measures, we examine the smoothed

effect of age, with cubic regression splines using GAMM, in controls, 22qDel carriers and

22qDup carriers. All age-related analyses were conducted on site-harmonized data, and adjusted

for the fixed effect of sex and the random effect of subject ID. First, we present the smoothed

effect of age for the control group, along with the corresponding FDR-corrected p-values specific

to the control group analysis, to establish typical developmental trajectories across white matter

regions. The effective degrees of freedom (edf) represent the complexity of the smoothed age

term, where values close to 1 indicate a linear relationship and higher values indicate non-linear

relationships between chronological age and the response variable. We then compared the

developmental trajectories of 22qDel and 22qDup carriers to the control group. A significant

interaction term between the smoothed-age term and 22qDel or 22qDup indicates a difference in

developmental trajectory compared to controls. To account for multiple testing, we applied FDR-

correction across all the interaction terms (i.e., age*22qDel and age*22qDup).

227   Results

Gene dosage effects of the 22q11.2 CNV on intracellular volume fraction

Mean ICVF was significantly greater in 22qDel carriers compared to controls, and significantly

greater across all regions except for FX and CGH. In contrast, 22qDup carriers showed

significantly lower mean ICVF, and lower ICVF across all white matter regions compared to

controls. Reflecting a similar pattern, there a negative gene dosage effect of the 22q11.2 CNV

(i.e., decreasing ICVF with increasing copy number) on mean ICVF and regional ICVF across

all white matter regions (Figure 1, left panel, sTable 3). Sensitivity analyses revealed that

scanner site, cerebral white matter volume, intracranial volume, and cerebrospinal fluid volume

did not influence these results (sFigure 2, sTable 4-7).

Higher mean, but regionally variable effects, of 22q11.2 CNV on orientation dispersion

index

Both 22qDel and 22qDup carriers showed greater mean ODI compared to controls; however,

there were region-specific differences. Relative to controls, 22qDel carriers had higher ODI in

limbic (i.e., FXST, CGH), commissural (i.e., GCC) and association (i.e., SS, SLF, SFO, EC)

fiber tracts, but also directionally variable effects on projection tracts (i.e., higher ODI in ACR

and PTR; lower ODI in IC, PLIC, ALIC and CST) compared to controls.

22qDup carriers showed higher ODI in projection (i.e., IC, PLIC, ALIC, CST), commissural

(i.e., BCC) and association (i.e., CGC, UNC, SFO) fiber tracts, but also lower ODI in PCR,

where projections fibers predominate, compared to controls (Figure 1, middle panel, sTable 3).

The sensitivity analyses revealed the same pattern, albeit with fewer significant results; opposing

effects of the 22qDel and 22qDup on ODI in projection regions (i.e., PTR, PLIC, IC) and

association regions (i.e., SS and SLF) yielded the most robust effects (sFigure 3, sTable 4-7).

Reduced free water diffusion in 22qDup carriers

There was no significant difference in mean isotropic water diffusion (ISO) between 22qDel and

controls. Regionally, however, 22qDel carriers exhibited higher ISO in a few projection tracts

(i.e., PCR, ACR), and directionally variable effects on limbic (i.e., higher in FX; lower in FXST)

commissural fibers (i.e., higher in TAP; lower in SCC) compared to controls (Figure 1, right

panel). Sensitivity analyses revealed only higher ISO in the FX and PTR in 22qDel carriers

compared to controls after adjusting for intracranial volume (still significant for both FX and

PTR), cerebrospinal fluid and white matter volume (significant for FX; sFigure 3, sTable 3).

22qDup carriers showed lower mean ISO compared to controls, primarily driven by lower ISO in

white matter regions predominated by projection fibers (IC, RLIC, CR, PCR, SCR, ACR, PTR)

and association fibers (UNC, SS, SLF, SFO, Figure 1, right panel). However, similar to the

22qDel carriers, 22qDup carriers also exhibited higher ISO in FX and FXST. Overall, the

sensitivity analyses yielded the same pattern of significant group-level differences between

22qDup and controls, except for the UNC, RLIC and IC that were no longer significant (sFigure

4, sTable 4-7).

22q11.2 CNV carriers show typical developmental trajectories in axonal density and

dispersion

In the control group, there were significant age effects on ICVF, ODI and ISO across all white

matter regions from childhood to adulthood, largely involving nonlinear age-associated

increases, except for the ODI in the tapetum of the corpus callosum (sTable 8). Most of the

measures showed a non-linear age effect across white matter regions, except for ODI in ALIC,

CST, EC, SFO, SLF and SS, which showed a linear age effect indicated by their effective

degrees of freedom (i.e., edf = 1). The age-related trajectories of 22qDel and 22qDup carriers did

not statistically differ from the age-related trajectories of the control group for the global and

regional measures of ICVF (Figure 2), ODI (Figure 3) or ISO (Figure 4) after adjusting for

multiple comparisons (sTable 9).

284   Discussion
285

Hitherto, this is the largest study to examine biophysical dMRI-derived measures in brain white

matter among 22q11.2 CNV carriers. The results showed widespread higher intracellular volume

fraction (ICVF) in 22qDel and lower ICVF in 22qDup compared to controls, where gene dosage

at the 22q11.2 locus was associated with lower ICVF across all white matter regions, suggesting

a strong effect of the 22q11.2 CNV on axonal density. Further, we found a higher mean axonal

dispersion (i.e., orientation dispersion index; ODI) for both 22qDel and 22qDup compared to

controls, but also directionally variable effects of the 22q11.2 CNV on ODI across white matter

regions; overall, having more copies of the 22q11.2 genomic locus was associated with lower

ODI in regions where association, projection and limbic fibers predominate. However, we also

observed that more copies of the 22q11.2 genomic locus were associated with higher ODI in a

few regions where projection fibers predominate, including the internal capsule and corticospinal

tract. In addition, we also observed lower free water diffusion (ISO) in 22qDup carriers

compared to controls, in several white matter tracts where projection and association fibers

predominate. Finally, we find that 22qDel and 22qDup carriers showed similar developmental

trajectories as controls in all NODDI measures across childhood and adolescence.

The current study extends on the previous findings of higher FA in 22qDel(7) and lower FA in

22qDup(17) relative to typically developing controls, by focusing on measures of two of the

main contributors to FA: axonal density and axonal dispersion(45). Across regions, we found

robust evidence for higher axonal density in 22qDel carriers and lower axonal density in 22qDup

carriers compared to controls, suggesting that axonal density is the main contributor to

differences in FA alterations in white matter microstructure among 22q11.2 CNV carriers. It

should be noted, however, that some white matter regions have previously been found to show

lower (i.e., fornix (cres)/stria terminalis, superior longitudinal fasciculus, extreme/external

capsule) or no difference in FA (i.e., sagittal stratum, uncinate fasciculus, posterior thalamic

radiation, superior fronto-occipital fasciculus) in 22qDel carriers compared to controls(7). In the

current study, we find that all white matter regions, except for the UNC, were characterized by

both higher axonal density and higher axonal dispersion in 22qDel carriers compared to controls.

Thus, it seems that a higher axonal dispersion counteracts the influence of a higher ICVF,

yielding lower or similar FA relative to controls in 22qDel carriers.

The greatest dosage effects of the 22q11.2 genomic region on ICVF were found in commissural

tracts and the corona radiata, which consist of fibers in the corpus callosum and projection fibers

from motor cortex, respectively. Such strong opposing effects on white matter architecture may

be driven by mechanisms that are directly related to the deletion and duplication of genes within

the 22q11.2 locus. However, the exact mechanisms underlying the dosage effect on these tracts

are unknown, but it should be noted that the corpus callosum has been found to be larger in

children with the 22qDel, which contrasts with the overall pattern of lower brain volume in

22qDel carriers(8,9). As corpus callosum is made up by subregions where the size of the axons

typically differ in humans, e.g., the relative density of thick and thin axons (i.e., the axon

diameter distribution) differs across the corpus callosum(66–68), it seems plausible that axonal

overgrowth or axonal diameter expansion may underlie alterations to axonal density of

commissural fibers and corpus callosum volume. Indeed, deletion of genes within the 22q11.2

locus (e.g., *ZDHHC8* and *RTN4R/Nogo-66)* has been linked to axonal growth and branching in

mice models of the 22qDel (18,21). Others have shown that heterozygous deletion of the *TBX1*

among mice can result in relatively more axons with small diameters and no large diameters in

the fimbria compared to wild-type mice(20), which is in line with the predictions from previous

diffusion MRI studies on 22q11.2 deletion carriers(7,14).

In addition to higher ICVF, our results also revealed higher ODI in 22qDel carriers for a few

white matter tracts that are predominated by association fibers (i.e., connections within the same

hemisphere), including the fornix (crus)/stria terminalis, extreme/external capsule and superior

longitudinal fasciculus. Lower FA has previously been found in these white matter regions in

22qDel compared to controls(7). While lower FA can indicate both lower axonal density and/or

higher axonal dispersion/crossing fibers(45), our result of increased ODI indicates that this is

primarily driven by higher axonal dispersion in association tracts in 22qDel carriers relative to

controls. Moreover, long association fibers typically originate from upper cortical layers and

show protracted development compared to limbic and projection fibers(27–30), which may

indicate that the white matter fibers originating from upper cortical layers show different

morphology compared to fibers originating from deeper cortical layers. This interpretation aligns

with several lines of previous research, including a mouse model of 22qDel showing fewer

neural progenitors(69), shorter axons, and less dendritic branching in neurons from the upper

cortical layers relative to wild type mice(22). The latter finding was observed after deletion of

the *TXNRD2* gene*,* which also resulted in fewer long distance cortical connections and more

local connections, which may yield higher axonal dispersion for long distance/association fibers

in 22qDel carriers (e.g., lower alignment and/or rerouting through compensation of alternative or

less direct paths). In addition, a post-mortem study reported lower frequency of neurons in the

upper cortical layers in a 22qDel carrier(16), and a neuroimaging study reported altered axonal

morphology (i.e., utilizing ultra-strong dMRI gradients sensitive to axonal morphology) and

lower white matter volume in association tracts in 22qDel carriers compared to controls(14),

supporting this interpretation.

Overall, the results point towards a divergent pattern of white matter microstructure between

22qDel and 22qDup carriers; however, there were a few points of convergence. Such diverging

and converging effects are of particular interest in research on genomic structural variants, where

gene dosage effects are often observed for neuroimaging derived features (i.e.,

divergence)(10,17,70) but where the clinical phenotypic profile may converge, (e.g., increased

risk of ASD for both 22qDel and 22qDup carriers(71)). In the current study, we observed that a

few regions exhibit similar directions of effect for both 22qDel and 22qDup, including higher

ODI in extreme/external capsule, higher ISO in fornix and lower ISO in fornix (crus)/stria

terminalis. These results indicate higher axonal dispersion in the extreme/external capsule in

both 22qDel and 22qDup, whereas the differences in fornix and fornix (crus)/stria terminalis may

be driven by CSF partial volume in the voxels at the fornix and adjacent areas. White matter

alterations in the external capsule, containing mostly corticocortical association fibers, have also

been associated with idiopathic ASD diagnosis (i.e., lower FA in twins with autism compared to

control twins)(72) and symptom severity, and it is one of the white matter regions most

susceptible to environmental factors (i.e., as estimated by variance attributed to environmental

effects in twin studies) in individuals with ASD(72). Moreover, a recent pilot study found

within-subject white matter changes in the external capsule among toddlers with ASD after a

behavioral intervention that improved verbal and communication skills(73), possibly implicating

a role of the external capsule in individuals with or at risk for speech and communication

problems. Our results suggest altered axonal circuitry in the external capsule may be implicated

in the behavioral phenotypes of ASD commonly observed in 22q11.2 CNV carriers, although

future research on the relationship between external capsule white matter structure and autistic

phenotypes in 22q11.2 CNV carriers is required to understand this mechanism.

The 22qDel is also one of the greatest genetic risk factors for psychosis (4,74). A previous

neuroimaging study has found evidence for lower axonal density in commissural and association

fibers and higher axonal density and free water in the anterior thalamic radiation, predominated

by projection fibers, among individuals with first episode psychosis(49). The differences in

thalamic radiation converge with our results of higher axonal density and free water in the

posterior thalamic radiation among 22qDel carriers compared to controls. A recent study

reported thalamocortical axonal overgrowth in 22qDel thalamocortical organoids, likely due to

elevated *FOXP2* expression as a trans-effect of the 22qDel(15). In addition, hyperconnectivity in

thalamic circuitry has been reported among 22qDel carriers compared to controls(75–77), which

has also been hypothesized as a potential biomarker for psychosis(78). It is also noteworthy that

the 22qDup, carriers of a putative protective genetic factor for psychosis(4–6), showed the

opposite pattern in our study, reflected by lower axonal density and free water in the posterior

thalamic radiation. This may suggest that a higher axonal density of thalamic projection fibers,

possibly due to thalamocortical axonal overgrowth, is implicated in the etiology of psychosis.

Finally, despite widespread moderate to large group differences between 22q11.2 CNV carriers

and controls, we did not observe any significant interactions between the NODDI measures and

age, indicating typical age-related changes in 22q11.2 CNV carriers from childhood to

adulthood. Thus, the results do not support alterations to late maturational changes in axonal

architecture, such as cumulative axonal diameter expansion. As such, the observed alterations in

axonal density may instead be reflective of atypical prenatal axonogenesis or lack of early axonal

pruning in 22qDel and 22qDup carriers. This is in line with previous research on 22qDel-

derived cortical spheroids exhibiting excessive prenatal overgrowth of thalamocortical axons(15)

and neural progenitors that stay longer in the cell cycling state(79), which may be smaller in size,

as indicated by reduced neurospheres (i.e., neural stem or progenitor cells) derived from human

induced pluripotent stem cells from 22qDel carriers(80). To speculate, the gene dosage effect of

22q11.2 genomic region on axonal density observed in our study may be reflective of a

differential effect on gene expression levels of genes that are involved in axonogenesis, possibly

through trans-regulatory mechanisms such as the mediating effect of *FOXP2* gene expression

levels on 22qDel thalamic neurons(15). However, to our knowledge, no studies have utilized

cellular models of the reciprocal 22qDup; such future studies are warranted to determine

differential gene dosage effects on axonal development.

Strengths and limitations

The current study includes a unique sample across a wide age-range consisting of both 22qDel

and 22qDup carriers and a large typically developing control sample with multi-shell dMRI data,

allowing for gene dosage and developmental analyses on metrics derived from advanced

neuroimaging techniques. The use of advanced dMRI measures provides novel insight into the

axonal architecture of the white matter microstructure in 22q11.2 CNV carriers, insight that

cannot be achieved through conventional DTI measures(12,45). Furthermore, the inclusion of the

large HCP dataset allowed us to estimate robust typical developmental trajectories from

childhood into adulthood. We also performed several sensitivity analyses demonstrating the

robustness of the results of the group level analyses by analyzing individuals derived from the

same scanner site only, as well as adjusting for volumetric differences in brain structure. The

results revealed largely the same pattern, albeit with some group differences becoming

attenuated. However, this was largely due to an increase in the standard error terms due to

reduction in sample size, whereas the point estimates and the directions of the effect sizes were

consistent with the main analyses. Thus, the results appear to be robust to any linear relationship

between the white matter measures and volumetric differences in brain structure between

22qDel, 22qDup and controls. In addition, the sensitivity analyses also showed that the main

results are not attributable to differences in scanner site or acquisition in our sample. It is also

important to note that small sample size is common in CNV research, limiting our ability to

detect small effects. Despite this, previous studies have found robust differences in white matter

microstructure with smaller sample sizes than our study(14), which remains the largest study on

biophysical white matter measures in 22q11.2 CNV carriers to date. Finally, although NODDI

measures can provide important insight into axonal density and axonal dispersion, it is important

to note that these are indirect measures of the cellular architecture, as estimated through a

multicompartment model(45). Our results also raise new questions. For instance, it is unclear if

the altered axonal density in 22q11.2 CNV carriers is a consequence of an absolute difference in

total neurons in the brain, altered axonal diameter distribution, or other mechanisms that alter

space requirements in white matter tissue. Future studies utilizing both in-vitro cellular models

and post-mortem examinations of 22q11.2 CNV carriers are warranted.

Conclusions

The results of the current study provide new insights into the underlying neurobiology of the

altered white matter microstructure in 22q11.2 CNV carriers, linking copy numbers at the

22q11.2 genomic locus to altered axonal density. However, we do not find evidence for altered

age-related changes in axonal density or dispersion across childhood and adolescence, possibly

reflecting atypical axonogenesis and/or axonal pruning during early neurodevelopment in

22q11.2 CNV carriers. Future studies are warranted to characterize the morphological features of

white matter axons across fetal development, and to connect these developmental alterations to

neuropsychiatric phenotypes.

Acknowledgments**:**

We thank Sergiu Pasca, M.D., for contributing to the study. This work used computational and

storage services associated with the Hoffman2 Cluster which is operated by the UCLA Office of

Advanced Research Computing's Research Technology Group and servers operated by the USC

Mark and Mary Stevens Neuroimaging and Informatics Institute (INI). Data from the UCLA and

Stanford cohorts were supported by grants from the MCHRI Uytengsu-Hamilton 22q11

Neuropsychiatry Research Award (UH22QEXTFY22-04) and NIH 1R21MH116473-01A1,

R37MH085953, 1U01MH119736-01 (to CEB), R01MH100900 (to JH), and R01MH129858 (to

CEB, PMT, IES, OAA). Research reported in this publication was supported by the National

Institute Of Mental Health of the National Institutes of Health under Award Number

U01MH109589/ U01AG052564 and by funds provided by the McDonnell Center for Systems

Neuroscience at Washington University in St. Louis. The HCP-Development 2.0 Release data

used in this report came from DOI: 10.15154/1520708. The HCP-Aging 2.0 Release data used in

this report came from DOI: 10.15154/1520707.

Conflict of Interest:

OAA is a consultant for Cortechs.ai and Precision Health and has received speaker's honoraria

from Lundbeck, Janssen, Otsuka, Lilly and Sunovion. The remaining authors have no conflicts

of interest to declare.

References

1. Olsen L, Sparsø T, Weinsheimer SM, Dos Santos MBQ, Mazin W, Rosengren A, et al.

Prevalence of rearrangements in the 22q11.2 region and population-based risk of

neuropsychiatric and developmental disorders in a Danish population: a case-cohort study.

Lancet Psychiatry. 2018 Jul 1;5(7):573–80.

2. Schneider M, Debbané M, Bassett AS, Chow EWC, Fung WLA, van den Bree MBM, et al.

Psychiatric Disorders From Childhood to Adulthood in 22q11.2 Deletion Syndrome: Results

From the International Consortium on Brain and Behavior in 22q11.2 Deletion Syndrome. Am

J Psychiatry. 2014 Jun;171(6):627–39.

3. Hoeffding LK, Trabjerg BB, Olsen L, Mazin W, Sparsø T, Vangkilde A, et al. Risk of

Psychiatric Disorders Among Individuals With the 22q11.2 Deletion or Duplication: A Danish

Nationwide, Register-Based Study. JAMA Psychiatry. 2017 Mar 1;74(3):282–90.

4. Marshall CR, Howrigan DP, Merico D, Thiruvahindrapuram B, Wu W, Greer DS, et al.

Contribution of copy number variants to schizophrenia from a genome-wide study of 41,321

subjects. Nat Genet. 2017 Jan;49(1):27–35.

5. Rees E, Kirov G, Sanders A, Walters JTR, Chambert KD, Shi J, et al. Evidence that

duplications of 22q11.2 protect against schizophrenia. Mol Psychiatry. 2014 Jan;19(1):37–40.

6. Owen MJ, Legge SE, Rees E, Walters JTR, O'Donovan MC. Genomic findings in

schizophrenia and their implications. Mol Psychiatry. 2023 Sep;28(9):3638–47.

7. Villalón-Reina JE, Martínez K, Qu X, Ching CRK, Nir TM, Kothapalli D, et al. Altered white

matter microstructure in 22q11.2 deletion syndrome: a multisite diffusion tensor imaging

study. Mol Psychiatry. 2020 Nov;25(11):2818–31.

8. Sun D, Ching CRK, Lin A, Forsyth JK, Kushan L, Vajdi A, et al. Large-scale mapping of

cortical alterations in 22q11.2 deletion syndrome: Convergence with idiopathic psychosis and

effects of deletion size. Mol Psychiatry. 2020 Aug;25(8):1822–34.

9. Ching CRK, Gutman BA, Sun D, Villalon Reina J, Ragothaman A, Isaev D, et al. Mapping

Subcortical Brain Alterations in 22q11.2 Deletion Syndrome: Effects of Deletion Size and

Convergence With Idiopathic Neuropsychiatric Illness. Am J Psychiatry. 2020 Jul

1;177(7):589–600.

10.   Schleifer CH, O'Hora KP, Fung H, Xu J, Robinson TA, Wu AS, et al. Effects of gene

dosage and development on subcortical nuclei volumes in individuals with 22q11.2 copy

number variations. Neuropsychopharmacology. 2024 May;49(6):1024–32.

11.   Mori S, Zhang J. Principles of Diffusion Tensor Imaging and Its Applications to Basic

Neuroscience Research. Neuron. 2006 Sep 7;51(5):527–39.

12.   Pierpaoli C, Jezzard P, Basser PJ, Barnett A, Di Chiro G. Diffusion tensor MR imaging

of the human brain. Radiology. 1996 Dec;201(3):637–48.

13.   Beaulieu C. The basis of anisotropic water diffusion in the nervous system - a technical

review. NMR Biomed. 2002;15(7–8):435–55.

14.   Raven EP, Veraart J, Kievit RA, Genc S, Ward IL, Hall J, et al. In vivo evidence of

514   microstructural hypo-connectivity of brain white matter in 22q11.2 deletion syndrome. Mol

515   Psychiatry. 2023 Jul 26;1–11.

516   15.   Shin D, Kim CN, Ross J, Hennick KM, Wu SR, Paranjape N, et al. Thalamocortical

517   organoids enable *in vitro* modeling of 22q11.2 microdeletion associated with neuropsychiatric

518   disorders. Cell Stem Cell. 2024 Mar 7;31(3):421-432.e8.

519   16.   Wu P, Teot L, Murdoch G, Monaghan-Nichols AP, McFadden K. Neuropathology of

520   22q11 Deletion Syndrome in an Infant. Pediatr Dev Pathol. 2014 Sep 1;17(5):386–92.

521   17.   Seitz-Holland J, Lyons M, Kushan L, Lin A, Villalon-Reina JE, Cho KIK, et al.

522   Opposing white matter microstructure abnormalities in 22q11.2 deletion and duplication

523   carriers. Transl Psychiatry. 2021 Nov 10;11(1):1–11.

524   18.   Mukai J, Tamura M, Fénelon K, Rosen AM, Spellman TJ, Kang R, et al. Molecular

525   Substrates of Altered Axonal Growth and Brain Connectivity in a Mouse Model of

526   Schizophrenia. Neuron. 2015 May 6;86(3):680–95.

527   19.   Mukai J, Dhilla A, Drew LJ, Stark KL, Cao L, MacDermott AB, et al. Palmitoylation-

528   dependent neurodevelopmental deficits in a mouse model of 22q11 microdeletion. Nat

529   Neurosci. 2008 Nov;11(11):1302–10.

530   20.   Hiramoto T, Sumiyoshi A, Yamauchi T, Tanigaki K, Shi Q, Kang G, et al. Tbx1, a gene

531   encoded in 22q11.2 copy number variant, is a link between alterations in fimbria myelination

532   and cognitive speed in mice. Mol Psychiatry. 2022 Feb;27(2):929–38.

21.   Budel S, Padukkavidana T, Liu BP, Feng Z, Hu F, Johnson S, et al. Genetic Variants of

Nogo-66 Receptor with Possible Association to Schizophrenia Block Myelin Inhibition of

Axon Growth. J Neurosci. 2008 Dec 3;28(49):13161–72.

22.   Fernandez A, Meechan DW, Karpinski BA, Paronett EM, Bryan CA, Rutz HL, et al.

Mitochondrial Dysfunction Leads to Cortical Under-Connectivity and Cognitive Impairment.

Neuron. 2019 Jun 19;102(6):1127-1142.e3.

23.   Khan TA, Revah O, Gordon A, Yoon SJ, Krawisz AK, Goold C, et al. Neuronal defects

in a human cellular model of 22q11.2 deletion syndrome. Nat Med. 2020 Dec;26(12):1888–

98.

24.   Stark KL, Xu B, Bagchi A, Lai WS, Liu H, Hsu R, et al. Altered brain microRNA

biogenesis contributes to phenotypic deficits in a 22q11-deletion mouse model. Nat Genet.

2008 Jun;40(6):751–60.

25.   Lin A, Forsyth JK, Hoftman GD, Kushan-Wells L, Jalbrzikowski M, Dokuru D, et al.

Transcriptomic profiling of whole blood in 22q11.2 reciprocal copy number variants reveals

that cell proportion highly impacts gene expression. Brain Behav Immun - Health. 2021 Dec

1;18:100386.

26.   Nehme R, Pietiläinen O, Artomov M, Tegtmeyer M, Valakh V, Lehtonen L, et al. The

22q11.2 region regulates presynaptic gene-products linked to schizophrenia. Nat Commun.

2022 Jun 27;13(1):3690.

27.   Huang H, Zhang J, Wakana S, Zhang W, Ren T, Richards LJ, et al. White and gray

553   matter development in human fetal, newborn and pediatric brains. NeuroImage. 2006 Oct

554   15;33(1):27–38.

555   28.   Kochunov P, Williamson DE, Lancaster J, Fox P, Cornell J, Blangero J, et al. Fractional

556   anisotropy of water diffusion in cerebral white matter across the lifespan. Neurobiol Aging.

557   2012 Jan 1;33(1):9–20.

558   29.   Wilson S, Pietsch M, Cordero-Grande L, Price AN, Hutter J, Xiao J, et al. Development

559   of human white matter pathways in utero over the second and third trimester. Proc Natl Acad

560   Sci U S A. 2021 May 18;118(20):e2023598118.

561   30.   Lebel C, Beaulieu C. Longitudinal Development of Human Brain Wiring Continues from

562   Childhood into Adulthood. J Neurosci. 2011 Jul 27;31(30):10937–47.

563   31.   Huang H, Xue R, Zhang J, Ren T, Richards LJ, Yarowsky P, et al. Anatomical

564   Characterization of Human Fetal Brain Development with Diffusion Tensor Magnetic

565   Resonance Imaging. J Neurosci. 2009 Apr 1;29(13):4263–73.

566   32.   Vasung L, Raguz M, Kostovic I, Takahashi E. Spatiotemporal Relationship of Brain

567   Pathways during Human Fetal Development Using High-Angular Resolution Diffusion MR

568   Imaging and Histology. Front Neurosci [Internet]. 2017 Jul 11 [cited 2024 Dec 16];11.

569   Available from:

570   https://www.frontiersin.org/journals/neuroscience/articles/10.3389/fnins.2017.00348/full

571   33.   Horgos B, Mecea M, Boer A, Szabo B, Buruiana A, Stamatian F, et al. White Matter

572   Dissection of the Fetal Brain. Front Neuroanat. 2020 Sep 25;14:584266.

573   34.   Rakic P, Yakovlev PI. Development of the corpus callosum and cavum septi in man. J

574   Comp Neurol. 1968;132(1):45–72.

575   35.   Ren T, Anderson A, Shen WB, Huang H, Plachez C, Zhang J, et al. Imaging, anatomical,

576   and molecular analysis of callosal formation in the developing human fetal brain. Anat Rec A

577   Discov Mol Cell Evol Biol. 2006;288A(2):191–204.

578   36.   Ouyang M, Dubois J, Yu Q, Mukherjee P, Huang H. Delineation of early brain

579   development from fetuses to infants with diffusion MRI and beyond. NeuroImage. 2019 Jan

580   15;185:836–50.

581   37.   Takahashi E, Folkerth RD, Galaburda AM, Grant PE. Emerging Cerebral Connectivity in

582   the Human Fetal Brain: An MR Tractography Study. Cereb Cortex. 2012 Feb 1;22(2):455–64.

583   38.   Kostović I, Jovanov-Milošević N. The development of cerebral connections during the

584   first 20–45 weeks' gestation. Semin Fetal Neonatal Med. 2006 Dec 1;11(6):415–22.

585   39.   LaMantia AS, Rakic P. Axon overproduction and elimination in the corpus callosum of

586   the developing rhesus monkey. J Neurosci. 1990 Jul 1;10(7):2156–75.

587   40.   LaMantia AS, Rakic P. Axon overproduction and elimination in the anterior commissure

588   of the developing rhesus monkey. J Comp Neurol. 1994 Feb 15;340(3):328–36.

589   41.   Paus T. Growth of white matter in the adolescent brain: Myelin or axon? Brain Cogn.

590   2010 Feb 1;72(1):26–35.

591   42.   Benes FM. Myelination of Cortical-hippocampal Relays During Late Adolescence.

592   Schizophr Bull. 1989 Jan 1;15(4):585–93.

43.   Kulikova S, Hertz-Pannier L, Dehaene-Lambertz G, Buzmakov A, Poupon C, Dubois J.

Multi-parametric evaluation of the white matter maturation. Brain Struct Funct. 2015 Nov

1;220(6):3657–72.

44.   Jones DK, Knösche TR, Turner R. White matter integrity, fiber count, and other fallacies:

The do's and don'ts of diffusion MRI. NeuroImage. 2013 Jun 1;73:239–54.

45.   Zhang H, Schneider T, Wheeler-Kingshott CA, Alexander DC. NODDI: Practical in vivo

neurite orientation dispersion and density imaging of the human brain. NeuroImage. 2012 Jul

16;61(4):1000–16.

46.   Grussu F, Schneider T, Tur C, Yates RL, Tachrount M, Ianuş A, et al. Neurite dispersion:

a new marker of multiple sclerosis spinal cord pathology? Ann Clin Transl Neurol.

2017;4(9):663–79.

47.   Jespersen SN, Bjarkam CR, Nyengaard JR, Chakravarty MM, Hansen B, Vosegaard T, et

al. Neurite density from magnetic resonance diffusion measurements at ultrahigh field:

Comparison with light microscopy and electron microscopy. NeuroImage. 2010 Jan

1;49(1):205–16.

48.   Sato K, Kerever A, Kamagata K, Tsuruta K, Irie R, Tagawa K, et al. Understanding

microstructure of the brain by comparison of neurite orientation dispersion and density

imaging (NODDI) with transparent mouse brain. Acta Radiol Open. 2017 Apr

1;6(4):2058460117703816.

49.   Kraguljac NV, Guerreri M, Strickland MJ, Zhang H. Neurite Orientation Dispersion and

613   Density Imaging in Psychiatric Disorders: A Systematic Literature Review and a Technical

614   Note. Biol Psychiatry Glob Open Sci. 2022 Jan 21;3(1):10–21.

615   50.   Genc S, Malpas CB, Holland SK, Beare R, Silk TJ. Neurite density index is sensitive to

616   age related differences in the developing brain. NeuroImage. 2017 Mar 1;148:373–80.

617   51.   Genc S, Raven EP, Drakesmith M, Blakemore SJ, Jones DK. Novel insights into axon

618   diameter and myelin content in late childhood and adolescence. Cereb Cortex N Y NY. 2023

619   Jan 4;33(10):6435–48.

620   52.   Fan Q, Tian Q, Ohringer NA, Nummenmaa A, Witzel T, Tobyne SM, et al. Age-related

621   alterations in axonal microstructure in the corpus callosum measured by high-gradient

622   diffusion MRI. NeuroImage. 2019 May 1;191:325–36.

623   53.   Benes FM, Turtle M, Khan Y, Farol P. Myelination of a Key Relay Zone in the

624   Hippocampal Formation Occurs in the Human Brain During Childhood, Adolescence, and

625   Adulthood. Arch Gen Psychiatry. 1994 Jun 1;51(6):477–84.

626   54.   Somerville LH, Bookheimer SY, Buckner RL, Burgess GC, Curtiss SW, Dapretto M, et

627   al. The Lifespan Human Connectome Project in Development: A large-scale study of brain

628   connectivity development in 5–21 year olds. NeuroImage. 2018 Aug 22;183:456.

629   55.   Bookheimer SY, Salat DH, Terpstra M, Ances BM, Barch DM, Buckner RL, et al. The

630   Lifespan Human Connectome Project in Aging: An overview. NeuroImage. 2019 Jan

631   15;185:335–48.

632   56.   Harms MP, Somerville LH, Ances BM, Andersson J, Barch DM, Bastiani M, et al.

633   Extending the Human Connectome Project across ages: Imaging protocols for the Lifespan

634   Development and Aging projects. NeuroImage. 2018 Dec;183:972–84.

635   57.   Garyfallidis E, Brett M, Amirbekian B, Rokem A, Van Der Walt S, Descoteaux M, et al.

636   Dipy, a library for the analysis of diffusion MRI data. Front Neuroinformatics [Internet]. 2014

637   [cited 2023 Aug 24];8. Available from:

638   https://www.frontiersin.org/articles/10.3389/fninf.2014.00008

639   58.   Manjón JV, Coupé P, Concha L, Buades A, Collins DL, Robles M. Diffusion Weighted

640   Image Denoising Using Overcomplete Local PCA. PLOS ONE. 2013 Sep 3;8(9):e73021.

641   59.   Andersson JLR, Skare S, Ashburner J. How to correct susceptibility distortions in spin-

642   echo echo-planar images: application to diffusion tensor imaging. NeuroImage. 2003

643   Oct;20(2):870–88.

644   60.   Andersson JLR, Sotiropoulos SN. An integrated approach to correction for off-resonance

645   effects and subject movement in diffusion MR imaging. NeuroImage. 2016 Jan 15;125:1063–

646   78.

647   61.   Jahanshad N, Kochunov P, Sprooten E, Mandl RC, Nichols TE, Almassy L, et al. Multi-

648   site genetic analysis of diffusion images and voxelwise heritability analysis: A pilot project of

649   the ENIGMA–DTI working group. NeuroImage. 2013 Nov 1;81:455–69.

650   62.   Mori S, Oishi K, Jiang H, Jiang L, Li X, Akhter K, et al. Stereotaxic white matter atlas

651   based on diffusion tensor imaging in an ICBM template. NeuroImage. 2008 Apr 1;40(2):570–

652   82.

63.   Beer JC, Tustison NJ, Cook PA, Davatzikos C, Sheline YI, Shinohara RT, et al.

Longitudinal ComBat: A method for harmonizing longitudinal multi-scanner imaging data.

NeuroImage. 2020 Oct 15;220:117129.

64.   Pakkenberg B, Pelvig D, Marner L, Bundgaard MJ, Gundersen HJG, Nyengaard JR, et al.

Aging and the human neocortex. Exp Gerontol. 2003 Jan 1;38(1):95–9.

65.   Wood S. Generalized additive models: an introduction with R. Chapman and Hall/CRC;

2017.

66.   Garic D, Yeh FC, Graziano P, Dick AS. In vivo restricted diffusion imaging (RDI) is

sensitive to differences in axonal density in typical children and adults. Brain Struct Funct.

2021 Nov 1;226(8):2689–705.

67.   Aboitiz F, Scheibel AB, Fisher RS, Zaidel E. Fiber composition of the human corpus

callosum. Brain Res. 1992 Dec 11;598(1):143–53.

68.   Liewald D, Miller R, Logothetis N, Wagner HJ, Schüz A. Distribution of axon diameters

in cortical white matter: an electron-microscopic study on three human brains and a macaque.

Biol Cybern. 2014 Oct 1;108(5):541–57.

69.   Meechan DW, Tucker ES, Maynard TM, LaMantia AS. Diminished dosage of 22q11

genes disrupts neurogenesis and cortical development in a mouse model of 22q11

deletion/DiGeorge syndrome. Proc Natl Acad Sci. 2009 Sep 22;106(38):16434–45.

70.   Lin A, Ching CRK, Vajdi A, Sun D, Jonas RK, Jalbrzikowski M, et al. Mapping 22q11.2

Gene Dosage Effects on Brain Morphometry. J Neurosci. 2017 Jun 28;37(26):6183–99.

71.   Lin A, Vajdi A, Kushan-Wells L, Helleman G, Hansen LP, Jonas RK, et al. Reciprocal

Copy Number Variations at 22q11.2 Produce Distinct and Convergent Neurobehavioral

Impairments Relevant for Schizophrenia and Autism Spectrum Disorder. Biol Psychiatry.

2020 Aug 1;88(3):260–72.

72.   Hegarty JP, Monterrey JC, Tian Q, Cleveland SC, Gong X, Phillips JM, et al. A Twin

Study of Altered White Matter Heritability in Youth With Autism Spectrum Disorder. J Am

Acad Child Adolesc Psychiatry. 2024 Jan;63(1):65–79.

73.   Saaybi S, AlArab N, Hannoun S, Saade M, Tutunji R, Zeeni C, et al. Pre- and Post-

therapy Assessment of Clinical Outcomes and White Matter Integrity in Autism Spectrum

Disorder: Pilot Study. Front Neurol. 2019 Aug 13;10:877.

74.   Singh T, Poterba T, Curtis D, Akil H, Al Eissa M, Barchas JD, et al. Rare coding variants

in ten genes confer substantial risk for schizophrenia. Nature. 2022 Apr;604(7906):509–16.

75.   Schleifer C, Lin A, Kushan L, Ji JL, Yang G, Bearden CE, et al. Dissociable Disruptions

in Thalamic and Hippocampal Resting-State Functional Connectivity in Youth with 22q11.2

Deletions. J Neurosci. 2019 Feb 13;39(7):1301–19.

76.   Mancini V, Zöller D, Schneider M, Schaer M, Eliez S. Abnormal Development and

Dysconnectivity of Distinct Thalamic Nuclei in Patients With 22q11.2 Deletion Syndrome

Experiencing Auditory Hallucinations. Biol Psychiatry Cogn Neurosci Neuroimaging. 2020

Sep 1;5(9):875–90.

77.   Moreau CA, Kumar K, Harvey A, Huguet G, Urchs SGW, Schultz LM, et al. Brain

693   functional connectivity mirrors genetic pleiotropy in psychiatric conditions. Brain. 2022 Sep

694   5;146(4):1686–96.

695   78.   Steullet P. Thalamus-related anomalies as candidate mechanism-based biomarkers for

696   psychosis. Schizophr Res. 2020 Dec 1;226:147–57.

697   79.   Rao SB, Brundu F, Chen Y, Sun Y, Zhu H, Shprintzen RJ, et al. Aberrant pace of cortical

698   neuron development in brain organoids from patients with 22q11.2 deletion syndrome and

699   schizophrenia [Internet]. bioRxiv; 2023 [cited 2024 Feb 5]. p. 2023.10.04.557612. Available

700   from: https://www.biorxiv.org/content/10.1101/2023.10.04.557612v1

701   80.   Toyoshima M, Akamatsu W, Okada Y, Ohnishi T, Balan S, Hisano Y, et al. Analysis of

702   induced pluripotent stem cells carrying 22q11.2 deletion. Transl Psychiatry. 2016

703   Nov;6(11):e934–e934.

Figure 1. Point estimates of group differences in intracellular volume fraction (ICVF),
orientation dispersion index (ODI) and isotropic volume fraction (ISO). The control group and
the 22qDel group are used as reference groups for the group comparisons. Gene dosage effects
are based on a continuous variable of copy number (i.e., 22qDel = 1, control = 2, and 22qDup =
3). E.g., Significant negative gene dosage effects indicate that an increased number of copies of
genes in the 22q11.2 genomic locus is associated with lower values of the diffusion measures.
ACR = anterior corona radiata, ALIC = anterior limb of internal capsule, BCC = body of corpus
callosum, CC = corpus callosum, CGC = cingulum cingulate gyrus, CGH = cingulum
hippocampus, CR = corona radiata, CST = corticospinal tract, EC = extreme/external capsule,
FX = fornix. FXST = fornix (crus)/stria terminalis, GCC = genu of corpus callosum, IC =
internal capsule, PCR = posterior corona radiata, PLIC = posterior limb of the internal capsule,
PTR = posterior thalamic radiation, RLIC = retrolenticular part of the internal capsule, SCC =
splenium of corpus callosum, SCR = superior corona radiata, SFO = superior fronto-occipital
fasciculus, SLF = superior longitudinal fasciculus, SS = sagittal stratum, TAP = tapetum of the
corpus callosum, UNC = uncinate fasciculus.

Figure 2. Developmental trajectories of intracellular volume fraction (ICVF) in global and
regional white matter microstructure for 22q11.2 deletion carriers (red), controls (black) and
22q11.2 duplication carriers (blue). ACR = anterior corona radiata, ALIC = anterior limb of
internal capsule, BCC = body of corpus callosum, CC = corpus callosum, CGC = cingulum
cingulate gyrus, CGH = cingulum hippocampus, CR = corona radiata, CST = corticospinal tract,
EC = extreme/external capsule, FX = fornix. FXST = fornix (crus)/stria terminalis, GCC = genu
of corpus callosum, IC = internal capsule, PCR = posterior corona radiata, PLIC = posterior limb
of the internal capsule, PTR = posterior thalamic radiation, RLIC = retrolenticular part of the
internal capsule, SCC = splenium of corpus callosum, SCR = superior corona radiata, SFO =
superior fronto-occipital fasciculus, SLF = superior longitudinal fasciculus, SS = sagittal stratum,
TAP = tapetum of the corpus callosum, UNC = uncinate fasciculus.

Figure 3. Developmental trajectories of orientation dispersion index (ODI) in global and
regional white matter microstructure for 22q11.2 deletion carriers (red), controls (black) and
22q11.2 duplication carriers (blue). ACR = anterior corona radiata, ALIC = anterior limb of
internal capsule, BCC = body of corpus callosum, CC = corpus callosum, CGC = cingulum
cingulate gyrus, CGH = cingulum hippocampus, CR = corona radiata, CST = corticospinal tract,
EC = extreme/external capsule, FX = fornix. FXST = fornix (crus)/stria terminalis, GCC = genu
of corpus callosum, IC = internal capsule, PCR = posterior corona radiata, PLIC = posterior limb
of the internal capsule, PTR = posterior thalamic radiation, RLIC = retrolenticular part of the
internal capsule, SCC = splenium of corpus callosum, SCR = superior corona radiata, SFO =
superior fronto-occipital fasciculus, SLF = superior longitudinal fasciculus, SS = sagittal stratum,
TAP = tapetum of the corpus callosum, UNC = uncinate fasciculus.

Figure 4. Developmental trajectories of isotropic volume fraction (ISO) in global and regional
white matter microstructure for 22q11.2 deletion carriers (red), controls (black) and 22q11.2
duplication carriers (blue). ACR = anterior corona radiata, ALIC = anterior limb of internal
capsule, BCC = body of corpus callosum, CC = corpus callosum, CGC = cingulum cingulate
gyrus, CGH = cingulum hippocampus, CR = corona radiata, CST = corticospinal tract, EC =
extreme/external capsule, FX = fornix. FXST = fornix (crus)/stria terminalis, GCC = genu of
corpus callosum, IC = internal capsule, PCR = posterior corona radiata, PLIC = posterior limb of
the internal capsule, PTR = posterior thalamic radiation, RLIC = retrolenticular part of the
internal capsule, SCC = splenium of corpus callosum, SCR = superior corona radiata, SFO =
superior fronto-occipital fasciculus, SLF = superior longitudinal fasciculus, SS = sagittal stratum,
TAP = tapetum of the corpus callosum, UNC = uncinate fasciculus.

Table 1. Participant baseline demographics

	Control (N=890)	22q11.2 Deletion Carriers (N=50)	22q11.2 Duplication Carriers (N=24)	P-value
Scans				
n	901	69	34	-
Age				
Mean (SD)	21.9 (13.5)	21.7 (9.8)	23.3 (14.9)	.334
Median [Min, Max]	15.4 [7.8, 51.1]	20.6 [7.4, 51.1]	15.4 [8.3, 49.4]	
Sex				
Female	485 (54.5%)	30 (65.2%)	11 (55.0%)	.362

22Q11.2 GENE DOSAGE EFFECTS ON WHITE MATTER