Transcriptomic biomarkers reveal jasmonic and salicylic acid state under field herbivory

Atsuki Tomita[1,2], Taro Maeda[1], Natsumi Mori-Moriyama[3], Yasuyuki Nomura[3], Yuko Kurita[4],

Makoto Kashima[5], Shigeyuki Betsuyaku[6], Yasuhiro Sato[7*], Atsushi J. Nagano[1, 8*]

[1] Institute for Advanced Biosciences, Keio University, Tsuruoka, Yamagata 997-0052, Japan

[2] Graduate School of Media and Governance, Keio University, Fujisawa, Kanagawa 252-0882, Japan

[3] Research Institute for Food and Agriculture, Ryukoku University, Otsu, Shiga 520-2194, Japan

[4] Graduate School of Agricultural and Life Sciences, The University of Tokyo, Bunkyo-ku, Tokyo 113-8657, Japan

[5] Faculty of Science, Toho University, Funabashi, Chiba, 274-8510, Japan

[6] Faculty of Agriculture, Ryukoku University, Otsu, Shiga 520-2194, Japan

[7] Faculty of Environmental Earth Science, Hokkaido University, Sapporo, Hokkaido 060-0810, Japan

[8] Bioscience and Biotechnology Center, Nagoya University, Nagoya, Aichi, 464-8601, Japan

Corresponding Author:

Atsushi J. Nagano

Bioscience and Biotechnology Center, Nagoya University, Nagoya, Aichi, 464-8601, Japan

Email: anagano@agr.nagoya-u.ac.jp

Yasuhiro Sato

Faculty of Environmental Earth Science, Hokkaido University, Sapporo, Hokkaido 060-0810, Japan

Email: yassato@ees.hokudai.ac.jp

Keywords: Salicylic acid, Jasmonic acid, transcriptome, biomarker, field environments

Running head: Transcriptomic biomarkers reveal SA/JA state under field herbivory

Abstract

Plants coordinate defence against pathogens and herbivores through the crosstalk between salicylic acid (SA) and jasmonic acid (JA) signalling pathways. However, little is known about how plants integrate the phytohormone concentration profiles to shape defence states in complex field environments. Here we show that systematically varying combinatorial SA and JA concentrations in *Arabidopsis thaliana* reveals 43 distinct combination-specific transcriptional signatures beyond SA–JA antagonism. Leveraging these data, we developed and validated machine-learning-based transcriptomic biomarkers that enabled independent quantification of SA and JA response states. By applying these biomarkers to field-grown *A. thaliana*, we revealed that SA/JA response states were dynamically shaped by abundant and co-occurrence of herbivores on individual plants. Our study provides a strategy to decode hormonal responses to biotic environments from field transcriptomes, bridging the gap between molecular signalling and ecological dynamics.

Introduction

In natural environments, plants are challenged by the community of diverse attackers such as herbivores and pathogens. This complex biotic environment necessitates dynamic integration and processing of stress signals to optimize defence strategies. Interactions between the salicylic acid (SA) and jasmonic acid (JA) signalling pathways have long been characterized by reciprocal antagonism[1,2]: SA promotes resistance to biotrophic pathogens[3], whereas JA mediates defence against insects and necrotrophic pathogens[4], partly through glucosinolate (GSL)-based anti-herbivore metabolites in Brassicaceae[5,6]. The SA-JA crosstalk, often described as reciprocal antagonism, allows plants to prioritise defence against attackers but can also compromise resistance to others[1,7]. Genetic and pharmacological studies have also shown the varying states of SA and JA, such that SA–JA crosstalk can turn into antagonistic, synergistic or additive relationships depending on phytohormone dose and timing[8,9]. However, we still lack a high-resolution quantification of the SA–JA interaction landscape that is necessary to understand plant responses to diverse attackers.

The plant SA/JA response to diverse biotic stresses is inherently context-dependent and highly dynamic[2,10,11]. The states of SA and JA varies not only among species and genotypes but also within a single plant–herbivore interaction process through different life stages. In *Arabidopsis thaliana*, for example, oviposition by *Pieris rapae* induces SA-dependent defences[12], whereas feeding by the hatched larvae subsequently triggers JA-dependent responses[13]. SA and JA signalling are not only host-regulated but are also actively manipulated by pathogens, herbivores, and associated organisms[14,15]. This dynamic defence states create a feedback loop that alters herbivore oviposition and host choice[15]. Furthermore, in natural settings, the simultaneous or sequential attack by diverse herbivore communities[10,11] drives a far greater diversification of SA/JA balances and downstream defence outputs than single-attacker experiments can simulate[16]. Therefore, developing robust method to quantify

the SA/JA signalling levels is pivotal for bridging the gap between laboratory and field conditions. To this end, it is essential to incorporate multiple-gene markers, which are themselves subject to SA-JA crosstalk.

To quantify plant responses to the diverse SA/JA balances, a systematic transcriptomic analysis using a wide range of SA and JA concentration combinations is essential[8,9]. Such an approach is crucial for comprehensively mapping the antagonistic, synergistic, and dose-dependent interactions between these pathways. However, it is challenging to systematically analyse the various combinations of SA and JA concentrations because large-scale RNA-seq experiments are required. To overcome this issue, many researchers have so far developed high-throughput RNA-Seq library preparation methods[17]. For instance, our previous study developed a Low-cost and eASY RNA-Seq method named Lasy-Seq[18]. This method has been applied to the systematic measurement of transcriptomic responses to multiple environmental inputs [19] but not to hormonal treatments and biotic stresses yet.

Here we show transcriptomic profiles of responses to 64 combinations of SA and JA concentrations in *A. thaliana* seedlings and derive transcriptomic biomarkers that quantify SA and JA response states. We applied these biomarkers to field transcriptomes from 2,381 *A. thaliana* individuals growing in distinct field in Otsu, Japan, and Zurich, Switzerland. By integrating these estimated response states with herbivore census data, we aimed to explore how variation in SA/JA signalling is closely linked to the specific assemblages and co-occurrence patterns of major herbivores on individual plants.

Results

Expression topography of SA/JA responses revealed by high-throughput RNA-seq.

To evaluate how SA/JA concentration altered the interaction between the SA and JA signalling pathways, we performed transcriptome profiling of *A. thaliana* seedlings treated with 64 different SA/JA concentrations combinations for six hours (Fig. 1a,b; Supplementary Table 1). We employed an assay system that allows the simultaneous processing of 96 samples using 96-well plates (Fig. 1a,b), yielding data of 370 samples with 12,963 expressed genes (Supplementary Fig. 1). PCA analysis indicated that the first principal component explained 19.3% of variance and coordinated with the SA treatment concentrations (Fig. 1c; Supplementary Fig. 2a), while the second principal component explained 6.7% of variance and coordinated with the JA treatment concentrations (Fig. 1d; Supplementary Fig. 2b). These variations were unlikely attributable to experimental bias as no apparent differences were observed among the 96 well plates (Supplementary Fig. 2c–f). The number of DEGs between control and phytohormone-treated samples increased in a dose-dependent manner for SA and JA (Supplementary Fig. 3a–c). For both SA and JA, there was a larger number of up-regulated genes than down-regulated genes in most conditions from the 0 mM to 0.33 mM treatments (Supplementary Fig. 3d).

The systematic data then enabled us to depict a gene expression pattern in response to SA/JA concentration, which was designated as an expression topography hereafter (Fig. 1e–h). We evaluated the expression topography of well-known marker genes involved in the SA/JA response in addition to two genes associated with JA metabolism, such as *PR1*, *VSP1*, *LOX3* and *CYP94B3*. The SA response marker gene, *PR1*, exhibited an expression threshold at 0.033 mM SA with an optimal concentration of 0.33 mM, expression of which was completely suppressed under 1 mM JA (Fig. 1e; Supplementary Fig. 4a). The JA response marker gene, *VSP1*, exhibited an expression threshold at 0.001 mM JA and an optimum at 0.1 mM, expression of which was reduced under SA conditions (Fig. 1f; Supplementary Fig. 4b).

These results are consistent with the qualitative understandings of SA/JA responses and SA-JA antagonism in previous studies[8,20]. Regarding JA metabolism, we detected expression changes of two key genes *LOX3* and *CYP94B3* in response to the SA/JA concentration. The expression of *LOX3* increased in a JA-concentration-dependent manner up to 0.33 mM JA (Fig. 1g; Supplementary Fig. 4c). This increase is consistent with previous evidence for autoregulation, since LOX3 encodes the enzyme for the first step of JA biosynthesis and accordingly LOX3-initiated pathway is known to exhibit a positive feedback loop[21]. Interestingly, *LOX3* expression was suppressed under 1 mM SA/JA treatment, suggesting that the positive feedback loop of *LOX3* is terminated once SA/JA levels reach substantial accumulation. In contrast, the expression of *CYP94B3*, a key regulator in the suppression of the JA signalling cascade, showed a slight increasing trend in a JA concentration-dependent manner (Fig. 1h; Supplementary Fig. 4d). Since CYP94B3 oxidizes the bioactive form of jasmonoyl-L-isoleucine, this oxidation by CYP94B3 regulates JA signalling levels and could affect SA/JA signalling levels [22]. Notably, *CYP94B3* expression was maximally induced under high SA conditions, showing a 4.6-fold increase between 0 mM and 1 mM SA in the presence of 1 mM JA. These results indicate that these genes fine-tune the SA/JA crosstalk depending on the specific combination of concentrations.

Diverse functional responses to the SA/JA concentrations in the expression topography

To characterize the diversity of expression topographies, we focused on 3,001 SA/JA-responsive genes (Supplementary Fig. 1) and performed affinity propagation clustering. This classification revealed 43 distinct gene clusters characterized by specific SA/JA concentration-dependent regulation patterns (Fig. 2a) and hierarchically aggregated into broad groups A–H to capture higher-order trends (Supplementary Fig. 5). These clusters contained 23 to 299 genes that shared highly coherent expression topographies (Fig. 2b).

Cluster A-02 expression showed JA-dependent activation and suppression by SA. Conversely,

cluster C-01 exhibited SA-dependent activation and JA-dependent suppression. To identify

putative cis-regulatory elements among the expression topographies, we performed motif

enrichment analysis on the upstream sequences of genes in each cluster (Fig. 2c;

Supplementary Table 2). Clusters A-02 and E-03, which peaked at intermediate

concentrations of both SA and JA, were strongly enriched for a GT3a type trihelix motif,

suggesting that GT element-binding factors preferentially tune gene expression under

moderate combinatorial phytohormone concentrations. By contrast, the SA dominated cluster

C-01 was characterized by pronounced enrichment of TGA-class bZIP motifs, consistent with

canonical NPR1-TGA-dependent SA signalling. Additional clusters were marked by motifs

for other stress-associated transcription factors (e.g. other members of TGA, WRKY, ERF and

MYB). Collectively, these results suggest that distinct *cis*-regulatory architectures provide the

basis for complex SA/JA concentration-dependent outputs.

To examine association between expression topographies and biological functions,

we also conducted a GO enrichment analysis on 43 clusters. The analysis showed that 18

clusters were significantly enriched with 1 to 103 GO terms (adjusted $P < 0.05$, Fig. 3a;

Supplementary Table 3). Genes in cluster A-02 were enriched in GO terms "response to

jasmonic acid" (GO:0009753, adjusted $P = 4.09 \times 10^{-4}$). This cluster contained many JA-

responsive genes reported in previous studies, such as *VSP2*, *JAZ8*, *JAZ10* and *JOX3* [23].

Cluster C-01 showed the expression induced by SA and suppressed by JA. The *PR1* gene was

contained in this cluster. Multiple GO terms were enriched in cluster C-01, including

"response to bacterium" (GO:0009617, adjusted $P = 1.18 \times 10^{-7}$) and "response to salicylic

acid" (GO:0009751, adjusted $P = 3.55 \times 10^{-6}$). When both SA and JA were treated in high

concentrations, genes in cluster B-01 were up-regulated and genes in clusters F-07 and F-08

were down-regulated. In cluster B-01, GO terms such as "protein folding" (GO:0006457,

adjusted $P = 2.68 \times 10^{-5}$) and "response to reactive oxygen species" (GO:0000302, adjusted P

$= 1.61 \times 10^{-3}$) were enriched. The cluster B-01 mainly comprised genes encoding heat shock

proteins such as HSP17.4 and HSP17.6. Clusters F-07 and F-08 were enriched in GO terms

such as "ribosome biogenesis" (GO: 0042254, adjusted $P = 7.71 \times 10^{-30}$ and 5.31×10^{-4}) and

"translation" (GO: 0006412, adjusted $P = 1.52 \times 10^{-75}$ and 3.28×10^{-2}). GO terms associated

with "photosynthesis" (GO:0015979, adjusted $P = 1.57 \times 10^{-4}$ and 4.22×10^{-3}) were enriched

in clusters F-05 and F-06. Genes in the clusters F-05 and F-06 were down-regulated in a dose-

dependent and additive manner by SA/JA co-treatment (Fig. 2a). These results were

consistent with previous studies, which demonstrated that co-treatment with SA and JA

reduced the expression of photosynthesis-related genes [23].

We found that two subtypes of GSLs, such as indole and aliphatic GSL, were

induced by different concentrations of JA. In cluster E-01, the GO term "glucosinolate

biosynthetic process" (GO:0019761, adjusted $P = 5.32 \times 10^{-3}$) was significantly enriched (Fig.

3a). Interestingly, "glucosinolate biosynthetic process" was also significantly enriched in

cluster F-02 (adjusted $P = 8.22 \times 10^{-3}$). The cluster E-01 contained genes related to indole

glucosinolate biosynthesis, *e.g. SUR1, CYP83B1, CYP79B3, MAM*. In contrast, the cluster F-

02 contained genes related to aliphatic GSL biosynthesis, *e.g. MAM3, CYP79F2, BCAT4,*

SOT18. The expression peak of indole GSL biosynthesis genes was found at lower

concentration of JA than that of aliphatic GSL (Fig. 3b). In low SA concentrations (0 mM to

0.01 mM), the expression peaks of indole GSL biosynthesis genes were found around 0.1 mM

JA, whereas those of aliphatic GSL biosynthesis genes were found around 0.003 mM JA (Fig.

3b). The expression peaks of indole GSL biosynthesis genes were found at approximately 30-

fold higher JA concentrations than that of aliphatic GSL-related genes. This response was

specific to low SA concentrations (Fig. 3c, Supplementary Fig. 6). Taken together, the ~30-

fold difference in JA sensitivity between these pathways, combined with their suppression by

SA, suggests a defence strategy finely tuned by the quantitative balance of SA and JA.

Multiple genes are required to account for the SA/JA response

Our comprehensive dataset allowed us to systematically evaluate how the expression of a single gene correlated with treated concentrations of SA and JA (Fig. 4a). Specifically, we focused on *PR1* and *VSP1* as they have long been used as major expression markers of SA and JA response[23]. Correlation coefficients between *PR1* and treatment concentrations of SA/JA were 0.63 and -0.30, respectively (Fig. 4a), expression of which had the 29th highest correlation with SA concentration. In contrast, *VSP1* had the 4,695th highest correlation with JA concentration. Correlation coefficients between *VSP1* and SA/JA treatment concentrations were -0.20 and 0.07, respectively (Fig. 4a). The genome-wide pattern of correlations indicates that correlations with SA and JA are associated with each other (Fig. 4a). We also focused on the other genes with high correlations. For example, *HRG2* exhibited the highest positive correlation with SA treatment concentration ($r = 0.74$), while that with JA was 0.14 (Fig. 4b). Meanwhile, UL4Z exhibited the highest negative correlation with SA, showing correlations of -0.70 and -0.22 with SA and JA, respectively (Fig. 4c). In contrast, the gene with the highest positive correlation with JA treatment concentration was *AT2G47950*. The correlation with JA was 0.85 and that with SA was 0.16 (Fig. 4d). The gene with the highest negative correlation with JA was *SAUR6*, with a correlation of -0.68 with JA and -0.44 with SA (Fig. 4e). These four genes responded to SA/JA at a specific threshold level (Fig. 4b–e), indicating that the expression level of a single gene alone cannot be markers to predict quantitative phytohormone levels These results showed that multiple genes responded to SA/JA treatments in a complex manner, thereby leading us to develop a quantitative response marker using multiple genes.

Development of a transcriptomic biomarker of SA/JA response state.

To develop a transcriptomic biomarker capable of estimating the quantitative state of

SA/JA responses, we employed machine learning-based regressions on the expression levels

of multiple genes. Specifically, we used an ensemble-learning approach with L1-regularized

linear regression (least absolute shrinkage and selection operator, LASSO) to construct

predictive models from the combinatorial SA/JA transcriptome dataset (Fig. 5a;

Supplementary Fig. 7; see Methods for training/test data details). The resulting biomarkers

demonstrated high predictive accuracy in an independent test dataset, yielding correlation

coefficients of 0.87 for SA and 0.95 for JA between predicted and actual treatment

concentrations (Fig. 5d,e). Notably, these values exceeded those of the best-correlated single

genes (Fig. 4), although some underestimation was observed in the high-concentration range.

These biomarkers also exhibited high specificity, such that the predicted SA and JA

concentrations were weakly correlated with each other ($r = 0.18$; Fig. 5f) and vice versa ($r =$

0.38; Fig. 5g). To infer the functional basis of these markers, we determined the contribution

of individual genes by calculating their selection frequency within the ensemble model

(Supplementary Fig. 8). The distributions of selection frequencies indicated that limited

number of genes were frequently used in the marker, but identified predominant contributors

(selection frequency > 0.95) for the JA marker including canonical signaling components such

as *JAZ1, JAZ2, JAZ3, JAZ5, JAZ6,* and *LOX2*. Similarly, predominant contributors for the SA

marker included known SA-responsive genes, including *ZAR1, WRKY18, WRKY70, ROXY18,*

ECS1 and *ARR4* (Supplementary Table 4). Thus, our multi-gene transcriptomic markers

provide a prospective predictor for the SA/JA response states.

To demonstrate the utility of our biomarkers for quantitative phenotyping, we applied

them to investigate SA/JA response states in *npr3/4* double mutants (*npr3-1npr4-3* and *npr3-*

2npr4-2), which are deficient in SA perception[24,25]. We profiled the transcriptomes of these

mutants under 25 combinatorial SA/JA conditions (total 194 samples; Supplementary Fig. 7)

and estimated their response states using our transcriptomic biomarkers (Fig. 5h,i). These

mutants showed significantly higher predicted SA concentration than that in Col-0 for all SA

treatment conditions except 1 mM SA (Fig. 5h), indicating that the mutants seemed to have a

higher sensitivity to SA than Col-0. This result suggested that NPR3/4 suppressed the SA

signalling pathway in conditions with lower than 1mM SA, which was consistent with

previous studies showing that NPR3 and NPR4 acted as transcriptional co-repressors of SA-

responsive genes[24,25]. For JA responses, the mutants showed significant differences from Col-

0 under several conditions, although these effects were not consistent among mutant alleles

(Fig. 5i). Collectively, these results demonstrate that our transcriptomic biomarkers provide a

robust, quantitative framework for decoding complex phytohormone response states, and

suggest their applicability to phenotyping of diverse mutants.

Application of biomarkers to transcriptome dataset under field herbivory.

Lastly, we asked whether our biomarkers were able to capture JA and SA responses

to herbivores in complex field environment. To address this question, we applied our

biomarkers to field data including 2,381 *A. thaliana* individuals and their insect communities

(1,195 samples in Otsu, and 1,186 samples in Zurich). SA and JA responses were

uncorrelated in Zurich, whereas a significant positive correlation was detected in Otsu (Fig.

6a). We then tested whether these responses were associated with insect communities, which

considerably differed between the two sites[26] (Fig. 6b). The vast majority of herbivores in

Zurich were caterpillars of diamondback moths ('Px') and flea beetles ('Ps' and 'Pa') that

could jump between plants[27], and those in Otsu were small white butterflies *Pieris rapae*

('Pr') and Px. While the variability of JA and SA responses was not directly explained by

insect species richness (Supplementary Fig. 9), we detected specific responses to abundant

herbivores (Fig. 6b). Specifically, in Zurich, the individual number of Px and wingless

Brevicoryne brassicae ('Bb') were positively correlated with the JA response and negatively

correlated with the SA response, consistent with a canonical SA-JA crosstalk pattern[1,2].

Conversely, Pa and Ps were positively associated with the SA response and negatively with

the JA response. In Otsu, both eggs ('Pr_e') and larvae (Pr_l') of small white butterflies

correlated positively with SA and JA responses. These results suggest that herbivore

communities may differentiate JA and SA responses through the species-specific influences

between the two sites.

To further narrow down the herbivore-specific responses, we performed LASSO

regression of herbivore abundance on the JA and SA responses (Fig. 6c). In Zurich, Px

showed the strongest effects on both hormonal responses ($\beta = 1.15$ for JA, $\beta = -0.40$ for SA).

Bb was also selected as a predictor of the JA response ($\beta = 0.45$). This result is consistent with

the general pattern that chewing herbivores induce JA signaling [28]. Pa exhibited the only

positive coefficient for the SA response ($\beta = 0.10$). In Otsu, Pr_e and Pr_l emerged as the

strongest predictors of the SA ($\beta = 0.58$) and JA ($\beta = 0.63$) responses, respectively, indicating

life-stage-specific signatures within the same species. These patterns are consistent with SA

induction by oviposition and JA induction by larval feeding[12,13].

To corroborate these findings, we analysed herbivore co-occurrence networks on

individual plants (Fig. 6d). Species selected by LASSO occupied central positions within their

respective networks. In Zurich, Pa and Px were significantly negatively associated (r = -0.25;

Supplementary Fig. 10a), whereas the Otsu network was cohesive with a strong co-occurrence

between Pr_e and Pr_l (r = 0.59; Supplementary Fig. 10b). Taken together, these results

indicate that the presence of major herbivores, coupled with their co-occurrence or exclusion

patterns, shapes site-specific JA and SA responses.

Discussion

Our systematic approach combining combinatorial phytohormone treatments with large-scale transcriptomics provides quantitative evidence for SA/JA responses. For example, our analysis revealed that genes encoding the biosynthesis of antimicrobial indole GSLs peaked at high JA concentrations, whereas those mediating aliphatic GSLs, which can deter a wide variety of herbivores[29], peaked at lower JA levels (Fig. 3b). Given that high JA accumulation is characteristic of wounded sites[30] while lower levels are typical of systemic tissues[31], this differential regulation reflects an effective strategy whereby plants deploy localized antimicrobial defences at wound sites while priming systemic tissues against subsequent herbivory. As demonstrated by the concentration-specific regulation of GSL biosynthesis genes, our experimental system successfully captured fine-tuned transcriptomic changes triggered in response to different SA/JA concentrations. These findings provide testable hypotheses based on large-scale transcriptome data.

We successfully quantified the response of whole seedlings to a wide range of SA/JA concentrations and combinations, but this might obscure tissue-specific and cell-type-specific responses could be obscured. Our measurement of total RNA amounts of shoot and root indicated that approximately 70% of total RNA of seedling was derived from shoot and 30% from root (Supplementary Fig. 12a). Responses of root-specific highly expressed genes were successfully measured (Supplementary Fig. 12b). These findings suggest that tissue-specific gene expressions can be detected unless they exhibit opposite patterns and are thus offset among tissues. While capturing specific responses in minor cell types remains challenging, highly expressed genes in specialized cells (e.g., *TGG2* in myrosin cells[32]; Supplementary Fig. 12c) may still be detectable. Considering the rapid advances of single-cell RNA-Seq technologies, future integration of systematic SA/JA treatment with single-cell RNA-Seq analysis will be a promising direction to fully resolve these cell-type-specific responses.

The transcriptomic biomarker that we developed in this study offer a robust

framework for predicting SA/JA response states across diverse contexts. Advances in

methodology have enabled sensitive and accurate quantification of phytohormones[33].

However, these measured levels do not always correspond to response states[33]. The

complementary use of our biomarkers with phytohormone quantification will help elucidate

the mechanisms underlying the discrepancies. Furthermore, our biomarkers will be applicable

to large-scale transcriptome data in emerging research areas, such as single-cell and field

transcriptomics. A study combining single-nucleus RNA-Seq and spatial transcriptomics to

analyse plant immune responses has been reported recently[34]. Applying the transcriptomic

biomarkers to such datasets would reveal SA/JA response dynamics during plant immune

responses at a single-cell resolution.

In field transcriptomics, massive transcriptome data acquired under natural

conditions are modelled with environmental data to reveal the responses of organisms in real-

world environments[35]. While laboratory studies have frequently reported apparent trade-offs

between SA and JA response[2,8,9,36], our field transcriptomics data reveal a more complex

reality shaped by community composition. We show that the balance between SA and JA

differs between the two field sites, exhibited significant differences in the insect community

composition as shown by our previous fieldwork[26]. Specifically, in the Otsu site, the Pr was

abundant, egg deposition and larval feeding of which are known to respectively induce SA

and JA-mediated defence responses[12,13]. This simultaneous induction may account for why

SA and JA are positively correlated in the Otsu site. Contrarily, Pa and Px were abundant in

the Zurich site. Pa can jump between neighbouring plants while chewing leaves[26,27], which

may be associated with the plant's SA response states. Importantly, integrating transcriptomic

biomarkers with herbivore data provided testable hypotheses about plant-herbivore

relationships. For instance, we detected the positive correlation between Pa abundance and

SA responses, along with its segregation from Px. This suggests the selective colonization of

highly mobile Pa onto plants that have already induced an SA response. Collectively,

contrasting SA/JA balance between the two field sites might be accounted by herbivore

specificity and mobility, which can be tested by excluding insect herbivores from the field.

In conclusion, our findings highlight the utility of combining transcriptomics and

herbivore community data to understand phytohormonal responses to complex biotic stress in

complex field environments. Previous field transcriptomic studies have primarily focused on

analyses using meteorological data[19,37,38]. While these efforts have successfully elucidated

plant responses to abiotic environmental factors, the responses to biotic factors remain largely

unexplored. By integrating the transcriptome biomarkers developed in this study with field

transcriptomics, it will deepen our understanding of the interplay between plant defence

responses and meteorological factors in real-world environments.

Materials and methods

Plant materials and SA/JA treatment

We used *A. thaliana* (Col-0: CS70000, *npr3-1npr4-3*: CS72351 and *npr3-2npr4-2*: CS72352) to analyse the responses to SA & JA concentrations. The seeds of these lines were obtained from the Arabidopsis Biological Resource Center. Methyl jasmonate (135-14411, FUJIFILM Wako, Osaka, Japan) and sodium salicylate (S3007, Merck, Darmstadt, Germany) were dissolved to 1 M in ethanol or water, respectively. 100× SA or 100× JA stock solutions were then diluted from 1 M stock to 0 mM, 0.1 mM, 0.33 mM, 1 mM, 3.3 mM, 10 mM, 33 mM, and 100 mM in ethanol or water, respectively. Sixty-four variations of 50× SA/JA mixed stock solutions were prepared by combining eight concentrations of 100× SA and 100× JA, and mixing equal amounts of each. These 64 stock solutions for SA/JA treated samples, 50% EtOH in 1/2 MS medium for mock samples, and only 1/2 MS medium for untreated samples were transferred to a 96-well plate, and stored at −20°C until use. A single sterilized seed was sown in each well of 96-well plates and grown in 50 µL of liquid 1/2 MS medium. After 3 days of treatment at 4°C in the dark, the plants were cultivated at 20 °C for 5 days under continuous light. SA/JA treatments were performed by adding 2 µL of 50× SA/JA mixed stock solutions and 48 µL of liquid 1/2 MS medium to each well containing 5-day-old seedlings. This allowed the 5-day-old seedlings to be incubated for 6 h under 64 variations of SA and JA concentration conditions (Fig. 1a). To monitor the broad spectrum of early defence responses, we treated with SA/JA for 6 h[39]. Mock and untreated samples were treated for same time as above using 1% ethanol in 1/2 MS medium and only 1/2 MS medium, respectively. Four replicates of the plant were treated by each SA & JA condition, respectively. We removed the medium and added 100 µL of direct-RT buffer [40] and a zirconia bead (YTZ-4, AS-ONE, Japan) to each well. Samples were ground using the TissueLyser II (QIAGEN, MD, USA) by 2 rounds for 1.5 minutes of 2.5 Hz. The homogenates were centrifuged at 3,000 rpm for 10 min, and 50 µL of the supernatants were stored as lysates at -

20 °C. For test data of the transcriptomic biomarkers of SA/JA response state, we performed

the same procedure using Col-0, *npr3-1npr4-3* and *npr3-2npr4-2* under twenty-five SA & JA

concentration conditions made by combining five concentrations: 0 mM, 0.001 mM, 0.01 mM,

0.1 mM and 1 mM (Supplementary Fig. 7).

RNA-Seq experiment and basic data analysis

The 5 μL of lysate was used by RNA-Seq library preparation with the Lasy-seq

version 1.1 method [18]. The library was sequenced by HiSeqX (Illumina, San Diego, CA) with

the paired-end 150+150bp mode. Trimomatic version 0.3.3 [41] was used for trimming read data,

and bowtie version 1.1.2 [42] and RSEM version 1.3.0 [43] were used for expression

quantification. For the following analysis, we used $\log_2(RPM+1)$ values calculated by the

same procedure of the previous study[18]. The expression topography of all SA/JA responsive

genes is shown in Supplementary Fig. 13.

We excluded samples with fewer than $10^{5.5}$ total reads from the analysis

(Supplementary Fig. 1b). We defined genes with more than 1.5 average $\log_2 (RPM+1)$ as the

expressed genes (Supplementary 1c). The consistency of the dataset across different 96-well

plates was verified by PCA (Supplementary Fig. 2c-f). We calculated the standard deviation

(SD) of the SA/JA-treated and control samples using the following procedure. To eliminate

the effect of differences in sample numbers, 63 samples were randomly selected from each of

the SA/JA-added samples and control samples. We calculated standard deviations among the

63 SA/JA-added samples as $SD_{treated}$ and standard deviations among the 63 control samples as

$SD_{control}$. In the expressed genes, SA/JA responsive genes (3,001 genes in 370 samples) were

selected based on the ratio of the $SD_{treated}$ to $SD_{control}$ (Supplementary Fig. 1d).

We detected differentially expressed genes between the control and each

concentration treatment using a likelihood ratio test of the edgeR package [44]. 0 mM SA and 0

mM JA, mock and untreated conditions were used as control samples. The FDR was

controlled using Benjamini and Hochberg's method[45].

Expression Topography

The average expression levels were calculated for each condition. The average

expression levels of 8×8 conditions data for each gene were designated as an expression

topography. To extract key characteristics of the expression topography, smoothing by a 3×3

moving average filter was applied to the expression data. Z-scores of the smoothed expression

data were calculated using the scale function of R. Using the apcluster version 1.4.8 package

[46], clustering was performed by the affinity propagation method based on Z-scores of the

smoothed expression data of SA/JA-responsive genes. Pearson's correlation coefficient (*r*)

was calculated using the Z-scores averaged over each cluster of affinity propagation. Groups

of clusters were determined by hierarchical clustering. The correlation coefficients were as the

distance between clusters, and the optimal number of groups was estimated by the gap

statistic.

Gene Ontology analysis

Gene Ontology annotations were obtained from The Arabidopsis Information

Resource (TAIR) on June 29, 2021. Fisher's exact test was used to test the enrichment of GO

terms. The multiple testing correction was performed with a FDR[45]. For these analyses, R

3.6.0 and 4.2.0 were used.

Transcription Factor Binding Motif Enrichment Analysis

Transcription factor binding motif enrichment analysis was performed using the

Simple Enrichment Analysis (SEA) tool from the MEME Suite version 5.5.0[47]. Promoter

sequences (1,000 bp upstream of the transcription start site) were extracted for each gene

identified through affinity propagation clustering. A background sequence set was constructed

from all promoter sequences of expressed genes. The analysis utilized the Arabidopsis DAP-

seq motif database[48]. SEA was run with default settings, and multiple testing was controlled

using the FDR[45]. To ensure robustness, we focused on motifs satisfying stringent criteria:

false discovery rate (FDR) < 0.05, enrichment ratio ≥ 2, and presence in >10% of target

sequences (Supplemental Table 2).

Transcriptomic biomarker of SA/JA response state

To create transcriptomic biomarkers correlated specifically with SA or JA, we

employed the method using ensemble learning and L1-regularized linear regression.

Expression data of the 12,963 expressed genes from 370 samples in 64 concentration

conditions were used as training data. The 1,296 genes, 10% of the expressed genes, were

randomly selected and used as input variables for LASSO[49]. The hyperparameter λ of LASSO

was determined by cross-validation. The selection of subset of genes and LASSO were

repeated 1,000 times. The median of the predicted values of the 1,000 regression models was

used as the transcriptomic biomarker. Gene selection frequency was defined as the proportion

of trials in which the gene had non-zero coefficient out of trials in which the gene was used as

input variable (Supplementary Fig. 8; Supplemental Table 4). The test data of regression

models were transcriptome data at 25 conditions of SA and JA combining 5 concentrations: 0,

0.001, 0.01, 0.1, and 1 mM.

Field experiment under naturally occurring herbivores

In this study, we reused field data deposited by Sato et al. (2024) who observed

naturally occurring herbivores on 199 *A. thaliana* accessions in the field. Briefly, 199

accessions were planted nearly 600 plants (= 199 accessions x 3 blocks) at outdoor gardens

located in Zurich, Switzerland (47°23'N, 8°33'E; alt. ca. 500 m) and Otsu, Japan (35°06'N,

134°56'E; alt. ca. 200 m). Plants were initially cultivated under short-day conditions and then

transferred to the field sites during early summer in 2017 and 2018. The 200 potted plants

(199 accessions and an additional Col-0) were randomly placed within a 260 cm x 64 cm

block in a checkered manner. Insect herbivores on individual plants were counted every two

to three days throughout the growing season. The original phenotypic data utilized in this

study are available in the Zenodo repository (doi:10.5281/zenodo.7945318).

RNA-Seq experiment of Field samples

Leaf samples for transcriptomic analysis were harvested three weeks post-

transplantation, corresponding to the endpoint of the herbivore observations. To minimize

circadian variations in gene expression, all samples were collected within a two-hour window

cantered around solar noon at each site. Harvested leaves were immediately immersed in an

RNA preservation buffer containing 5.3 M (NH4)2SO4, 20 mM EDTA, and 25 mM trisodium

citrate dihydrate (pH 5.2). Samples were incubated at $4^{\circ}C$ overnight and subsequently stored

at $-80^{\circ}C$ until processing. Total RNA was extracted using the Maxwell 16 Lev Plant RNA Kit

(Promega Japan) according to the manufacturer's protocol. RNA-seq libraries were prepared

using the Lasy-Seq version 1.1[18]. These libraries were sequenced by Illumina HiSeq 2500 and

HiSeq X platforms to generate 50-bp and 150-bp reads, from the 3' end, respectively. Output

fastq files were processed and mapped in the same way as described above. We excluded

samples with fewer than $10^{5.5}$ total reads from the analysis. We defined genes with more than

1.5 average $\log_2(RPM+1)$ as the expressed genes. Expressed genes in all three independent

datasets (training, validation, and field datasets from Zurich and Otsu) were retained for

biomarkers. Following the procedure for the transcriptomic biomarker developed in this study,

the SA/JA response states were estimated for field-grown *A. thaliana* plants.

Prediction of SA/JA response states from herbivore abundances

To quantify which herbivore species were most strongly associated with SA/JA

response states, we tested whether the predicted SA/JA scores could be explained by

herbivore community composition. To minimize the influence of strongly tissue- or stage-

specific genes and to enrich for core components of SA/JA signalling, we used as biomarkers

only expressed genes ($\log_2$(RPM + 1) > 1.5) in both the seedling training dataset and the field

leaf dataset. We constructed four models in total, corresponding to each combination of field

site (Zurich or Otsu) and response variable (SA or JA). In each model, the response variable

was the predicted SA or JA response state, and the predictors were the abundances of 14

herbivore species recorded on individual plants. We fitted LASSO[49] regression models using

the glmnet package in R (family = "gaussian", alpha = 1). The regularization parameter λ was

selected by cross-validation using the default procedures implemented in glmnet.

Analysis of co-occurrence patterns among herbivore species

$\log_2$-transformed count data were used to assess the association of herbivore species

abundances on individual plants. We calculated Pearson's correlation coefficients between all

species pairs. Pearson's correlation was tested using cor.test in R (Supplementary Fig. 10).

Species pairs showing significant associations ($|r| > 0.15$ and $p < 0.05$) were retained and used

to construct the abundance-based co-occurrence network (Fig. 6d). The network was

visualized using the Fruchterman–Reingold algorithm implemented in the igraph package in

R, such that species with stronger associations ($|r|$) were positioned closer to each other.

Herbivore species that were extremely rare (maximum co-occurrence frequency < 0.01 across

all pairs; Supplementary Fig. 11) were excluded from the network visualizations.

Data availability

The RNA-Seq data of combinatorial treatments with SA and JA were submitted to

the NCBI Sequence Read Archive repository under the BioProject PRJNA1232845. The

RNA-Seq data from the field herbivory experiments were submitted to the NCBI Sequence

Read Archive repository under the BioProject PRJNA1055060, PRJNA1055104,

PRJNA1055317, PRJNA1055424, PRJNA1055734, PRJNA1055736, PRJNA1056126, and

PRJNA1055939. R scripts and data used in the analysis (count data, etc.) are available via the

figshare repository (https://doi.org/10.6084/m9.figshare.29323187). (These data are currently

under embargo and will be made publicly available upon acceptance of the manuscript.)

References

1.   Hou, S. & Tsuda, K. Salicylic acid and jasmonic acid crosstalk in plant immunity. *Essays Biochem.* 66, 647–656 (2022).

2.   Pieterse, C. M. J., Van Der Does, D., Zamioudis, C., Leon-Reyes, A. & Van Wees, S. C. M. Hormonal modulation of plant immunity. *Annu. Rev. Cell Dev. Biol.* 28, 489–521 (2012).

3.   Maruri-López, I., Aviles-Baltazar, N. Y., Buchala, A. & Serrano, M. Intra and Extracellular Journey of the Phytohormone Salicylic Acid. *Front. Plant Sci.* 10, 423 (2019).

4.   Howe, G. A. & Jander, G. Plant immunity to insect herbivores. *Annu. Rev. Plant Biol.* 59, 41–66 (2008).

5.   Hopkins, R. J., Van Dam, N. M. & Van Loon, J. J. A. Role of glucosinolates in insect-plant relationships and multitrophic interactions. *Annu. Rev. Entomol.* 54, 57–83 (2009).

6.   Abuyusuf, M. *et al.* Glucosinolates and Biotic Stress Tolerance in Brassicaceae with Emphasis on Cabbage: A Review. *Biochem. Genet.* 61, 451–470 (2023).

7.   Pieterse, C. M. J., Ton, J. & Van Loon, L. C. Cross-talk between plant defence signalling pathways: boost or burden? *AgBiotechNet* 3, 1–8 (2001).

8.   Mur, L. A. J., Kenton, P., Atzorn, R., Miersch, O. & Wasternack, C. The outcomes of concentration-specific interactions between salicylate and jasmonate signaling include synergy, antagonism, and oxidative stress leading to cell death. *Plant Physiol.* 140, 249–262 (2006).

9.   Robert-Seilaniantz, A., Grant, M. & Jones, J. D. G. Hormone crosstalk in plant disease and defense: More than just JASMONATE-SALICYLATE antagonism. *Annu Rev Phytopathol* 49, 317–343 (2011).

10.   Stam, J. M. *et al.* Plant interactions with multiple insect herbivores: From community to genes. *Annu. Rev. Plant Biol.* 65, 689–713 (2014).

11.   Fernández de Bobadilla, M., Vitiello, A., Erb, M. & Poelman, E. H. Plant defense strategies against attack by multiple herbivores. *Trends Plant Sci.* 27, 528–535 (2022).

12.   Hilfiker, O. *et al.* Insect eggs induce a systemic acquired resistance in Arabidopsis.

Plant J. 80, 1085–1094 (2014).

13.　Vos, I. A. *et al.* Onset of herbivore-induced resistance in systemic tissue primed for
jasmonate-dependent defenses is activated by abscisic acid. *Front. Plant Sci.* 4, 539
(2013).

14.　Zheng, X. Y. *et al.* Coronatine promotes pseudomonas syringae virulence in plants by
activating a signaling cascade that inhibits salicylic acid accumulation. *Cell Host*
Microbe 11, 587–596 (2012).

15.　Zhu, F., Poelman, E. H. & Dicke, M. Insect herbivore-associated organisms affect plant
responses to herbivory. *New Phytol.* 204, 315–321 (2014).

16.　Nguyen, D., Rieu, I., Mariani, C. & van Dam, N. M. How plants handle multiple
stresses: hormonal interactions underlying responses to abiotic stress and insect
herbivory. *Plant Mol. Biol.* 91, 727–740 (2016).

17.　Stark, R., Grzelak, M. & Hadfield, J. RNA sequencing: the teenage years. *Nat. Rev.*
Genet. 20, 631–656 (2019).

18.　Kamitani, M., Kashima, M., Tezuka, A. & Nagano, A. J. Lasy-Seq: a high-throughput
library preparation method for RNA-Seq and its application in the analysis of plant
responses to fluctuating temperatures. *Sci. Rep.* 9, 7091 (2019).

19.　Hashida, Y. *et al.* Field-crop transcriptome models are enhanced by measurements in
systematically   controlled   environments.   Preprint   at   bioRxiv
https://doi.org/10.1101/2024.09.21.614268 (2024).

20.　Thaler, J. S., Humphrey, P. T. & Whiteman, N. K. Evolution of jasmonate and salicylate
signal crosstalk. *Trends Plant Sci.* 17, 260–270 (2012).

21.　Yang, T. H., Lenglet-Hilfiker, A., Stolz, S., Glauser, G. & Farmer, E. E. Jasmonate
precursor biosynthetic enzymes LOX3 and LOX4 control wound-response growth
restriction. *Plant Physiol.* 184, 1172–1180 (2020).

22.　Kitaoka, N. *et al.* CYP94B3 activity against jasmonic acid amino acid conjugates and
the elucidation of 12-O-β-glucopyranosyl-jasmonoyl-l-isoleucine as an additional
metabolite. *Phytochemistry* 99, 6–13 (2014).

23.　Zhang, N., Zhou, S., Yang, D. & Fan, Z. Revealing Shared and Distinct Genes
Responding to JA and SA Signaling in Arabidopsis by Meta-Analysis. *Front. Plant Sci.*

11, 908 (2020).

24. Ding, Y. *et al.* Opposite Roles of Salicylic Acid Receptors NPR1 and NPR3/NPR4 in
Transcriptional Regulation of Plant Immunity. *Cell* 173, 1454–1467 (2018).

25. Fu, Z. Q. *et al.* NPR3 and NPR4 are receptors for the immune signal salicylic acid in
plants. *Nature* 486, 228–232 (2012).

26. Sato, Y. *et al.* Reducing herbivory in mixed planting by genomic prediction of neighbor
effects in the field. *Nat. Commun.* 15, 8467 (2024).

27. Takimoto, H., Sato, Y., Nagano, A. J., Shimizu, K. K. & Kanagawa, A. Using a two-
stage convolutional neural network to rapidly identify tiny herbivorous beetles in the
field. *Ecol. Inform.* 66, 101466 (2021).

28. Onkokesung, N., Reichelt, M., van Doorn, A., Schuurink, R. C. & Dicke, M.
Differential costs of two distinct resistance mechanisms induced by different herbivore
species in Arabidopsis. *Plant Physiol.* 170, 891–906 (2016).

29. Jeschke, V., Gershenzon, J. & Vassão, D. G. A mode of action of glucosinolate-derived
isothiocyanates: Detoxification depletes glutathione and cysteine levels with
ramifications on protein metabolism in Spodoptera littoralis. *Insect Biochem. Mol. Biol.*
71, 37–48 (2016).

30. Glauser, G. *et al.* Velocity estimates for signal propagation leading to systemic
jasmonic acid accumulation in wounded Arabidopsis. *J. Biol. Chem.* 284, 34506–34513
(2009).

31. Ohnmeiss, T. E., McCloud, E. S., Lynds, G. Y. & Baldwin, I. T. Within-plant
relationships among wounding, jasmonic acid, and nicotine: Implications for defence in
Nicotiana sylvestris. *New Phytol.* 137, 441–452 (1997).

32. Maeda, T. *et al.* Single-Cell RNA Sequencing of Arabidopsis Leaf Tissues Identifies
Multiple Specialized Cell Types: Idioblast Myrosin Cells and Potential Glucosinolate-
Producing Cells. *Plant Cell Physiol.* 64, 234–247 (2023).

33. Vrobel, O. & Tarkowski, P. Can plant hormonomics be built on simple analysis? A
review. *Plant Methods* 19, 107 (2023).

34. Nobori, T. Exploring the untapped potential of single-cell and spatial omics in plant
biology. *New Phytol.* 247, 1098–1116 (2025).

35.   Nagano, A. J. *et al.* Deciphering and prediction of transcriptome dynamics under
fluctuating field conditions. *Cell* 151, 1358–1369 (2012).

36.   Zhang, P. J. *et al.* Phloem-feeding whiteflies can fool their host plants, but not their
parasitoids. *Funct. Ecol.* 27, 1304–1312 (2013).

37.   Kashima, M. *et al.* Genomic basis of transcriptome dynamics in rice under field
conditions. *Plant Cell Physiol.* 62, 1436–1445 (2021).

38.   Nagano, A. J. *et al.* Annual transcriptome dynamics in natural environments reveals
plant seasonal adaptation. *Nat. Plants* 5, 74–83 (2019).

39.   Goda, H. *et al.* The AtGenExpress hormone and chemical treatment data set:
experimental design, data evaluation, model data analysis and data access. *Plant J.* 55,
526–542 (2008).

40.   Kashima, M. *et al.* DeLTa-Seq: direct-lysate targeted RNA-Seq from crude tissue lysate.
Plant Methods 18, 99 (2022).

41.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 30, 2114–2120 (2014).

42.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat.
Methods* 9, 357–359 (2012).

43.   Li, B. & Dewey, C. N. RSEM: Accurate transcript quantification from RNA-Seq data
with or without a reference genome. *BMC Bioinformatics* 12, 323 (2011).

44.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: A Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* 26, 139–
140 (2009).

45.   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *J. R. Stat. Soc. B* 57, 289–300 (1995).

46.   Bodenhofer, U., Kothmeier, A. & Hochreiter, S. Apcluster: An R package for affinity
propagation clustering. *Bioinformatics* 27, 2463–2464 (2011).

47.   Bailey, T. L., Williams, N., Misleh, C. & Li, W. W. MEME: discovering and analyzing
DNA and protein sequence motifs. *Nucleic Acids Res.* 34, W369–W373 (2006).

48. O'Malley, R. C. *et al.* Cistrome and Epicistrome Features Shape the Regulatory DNA
Landscape. *Cell* 165, 1280–1292 (2016).

49. Tibshirani, R. Regression Shrinkage and Selection Via the Lasso. *J. R. Stat. Soc. B* 58,
267–288 (1996).

Acknowledgements

We thank Dynacom Co., Ltd. (Chiba, Japan) for technical assistance with the analysis of

RNA-Seq data.

Author contributions

AJN and SB conceived the study. YN, YK and MK developed the assay system. AT, YS and

NMM conducted the RNA-Seq experiments. AT, TM, YS and AJN analysed the data. AT, YS

and AJN wrote the manuscript with input from all co-authors.

Funding

This work was supported by JST CREST JPMJCR15O2, JST FOREST JPMJFR210B, JSPS

(JP20H00423, JP23H00386, JP23K18156) and MEXT (JP23H04967) for AJN; and by JST

PRESTO JPMJPR17Q4 and JSPS JP23K14270 for YS.

Competing Financial Interests

The authors declare that they have no competing interests.

Figure1 Treatments of 64 combinations of SA/JA concentrations

(a) SA/JA treatment for *A. thaliana* seedling using 96-well plates. Each sample was treated with SA and JA at 64 conditions that were systematically combined among 8 concentrations: 0 mM, 0.001 mM, 0.0033 mM, 0.01 mM, 0.033 mM, 0.1 mM, 0.33 mM, and 1 mM. (b) A 96-well plate with seedling just prior to the SA/JA treatment. (c. d) Principal component analysis of samples. Each point represents samples. The color of each point indicates the log-transformed SA concentration (c), JA concentration (d). (e-h) Expression topography of genes related SA/JA responses. The horizontal axis of the heatmap corresponds to the concentration of SA, and the vertical axis to the concentration of JA. The color of each cell indicates the average $\log_2(\text{RPM} + 1)$ for each concentration condition, n = 3~11. *PR1* is a marker gene for SA responses. (e). *VSP1* is a marker gene for JA responses. (f). *LOX3* is encoding enzymes involved in JA synthesis (g). *CYP94B3* is involved in the downregulation of the JA response (h).

Figure 2 Forty-three gene clusters of expression topography represent various SA/JA responses

(a) Average expression topography of each cluster. The horizontal axis is the concentration of SA, and the vertical axis is the concentration of JA. The colors represent the relative expression values of the genes in each cluster, with lighter colors indicating higher expression levels.

(b) Upper 2 plots indicate concentration of SA/JA. Lower 2 plots show the relative expression levels of the cluster A-02 and cluster C-01. Expression profiles of 40 genes in A-02 and 64 genes in C-01 are indicated as gray points and lines. Mean expression is indicated as red lines. (c) Transcription factor binding motifs enriched in gene clusters. Bubbles represent motifs significantly enriched (FDR < 0.05, enrichment ratio ≥ 2, TP% > 10%) in each cluster. Bubble size indicates the percentage of target sequences containing the motif, and bubble color represents the statistical significance (-log10(FDR)). The top 3 most significant motifs per cluster are shown. A full list of enriched motifs was indicated in Supplemental Table 2.

Figure 3 Genes with similar expression topographies were involved in same biological prosses.

(a) GO enrichment in the gene clusters. The number of genes contained in each cluster is indicated in parentheses. The size of the circles indicates the number of genes. The color scale indicates the adjusted P-value (FDR) in the GO enrichment analysis. A full list of enriched GO terms was indicated in Supplemental Table 3. (b, c) Relative expression levels of glucosinolate (GSL) synthesis genes. Mean expressions for 0 mM to 0.01 mM SA conditions (b) or 0.033 mM to 1 mM SA conditions (c) are plotted. The horizontal axis indicates JA concentration (mM), and the vertical axis indicates relative expression level. Aliphatic GSL synthesis genes are shown in red. Indole GSL synthesis genes are shown in blue. Bold lines indicate LOESS curves drawn for each glucosinolate type. (d) Expression topographies of aliphatic GSL synthesis genes. (e) Expression topographies of indole GSL synthesis genes. Data used in the panel (b) are indicated by green boxes.

Figure 4 Dose-responsive gene expression and Transcriptomic biomarker of SA/JA response state

(a,b) Histogram and scatter plot of correlation coefficients between gene expression levels and SA/JA concentrations (a), Genes with the highest or lowest correlation coefficient are indicated as color points. (b–e) Expression topographies of genes with the highest positive and negative correlation. b, H2O2 response gene *HRG2*, showing the highest positive correlation with SA. (c) Ribosomal protein gene *UL4Z*, showing the highest negative correlation with SA. (d) Uncharacterized protein gene *AT2G47950*, showing the highest positive correlation with JA. (e) Auxin-inducible gene *SAUR6*, showing the highest negative correlation with JA.

Figure 5 Transcriptomic biomarker of SA/JA response state

(a) Scheme of development of a transcriptomic biomarker of SA/JA response state from RNA-seq data. See Materials and Methods for

further details. (b,c) Predictions of the transcriptomic biomarkers for training data. Predicting of the marker for SA response state and treated

SA concentration (b). Predicting of the marker for JA response state and treated JA concentration (c). (d-g) Predicting of the transcriptomic

biomarkers for test data of Col-0. Prediction of the SA marker and treated SA concentration for test data (d). Prediction of the JA marker and

treated JA concentration for test data (e). Prediction of the SA marker and treated JA concentration for test data (f). Prediction of the JA

marker and treated SA concentration for test data (g). The horizontal axis is the predicted SA/JA concentration by the transcriptomic

biomarker, and the vertical axis is the treated SA/JA concentration. The result of liner regression is shown in the blue line. (h, i) Predictions of

the SA/JA marker for npr3/4 double mutants. Data obtained from 25 concentration conditions were grouped by the treated concentrations of

SA (h) and JA (i). The vertical axis indicates the predicted concentrations by the transcriptomic biomarker. Results of Mann-Whitney's U test

were indicated as *: P < 0.05, **: P < 0.01, ***: P < 0.001 and NS: not significant.

Figure 6. SA/JA response states of field transcriptomes and their association with herbivore communities.

(a) Relationship between predicted SA and JA concentrations in Zurich and Otsu populations. Blue lines and r indicate linear regression fits and Pearson correlation coefficients, respectively. (b) Heatmap of Pearson correlation coefficients between predicted SA/JA concentrations and insect abundance. Side bars indicate total insect counts per population. Colors represent correlation strength (red: positive; blue: negative). (c) LASSO regression coefficients estimating insect effects on SA/JA responses. Only species with non-zero coefficients are shown. Colors represent the sign and magnitude of coefficients (red: positive; blue: negative). (d) Insect co-occurrence networks based on significant correlations (|r| > 0.15, p < 0.05). Nodes represent insect species. Square: LASSO-selected predictors. Circle: others. Edges represent correlations (green: positive/co-occurrence; pink: negative/segregation), with width proportional to correlation strength (|r|).