**Prolonged Single Neuron Voltage Imaging in Behaving Mammals**

Yangyang Wang[1]†, Cara Ravasio [1]†, Yuxin Zhou[1], Xue Han[1]*

**Affiliations:**
[1]Department of Biomedical Engineering, Boston University; Boston, MA 02215, USA.

*Corresponding author. Email: xuehan@bu.edu
†These authors contributed equally to this work

**Abstract**

Long-term recording of single-neuron membrane voltage dynamics during behavior is highly
desirable. Using ElectraOFF, a fully genetically encoded, photostable, high-performance
fluorescence voltage indicator, we achieve routine cellular-resolution imaging over tens of minutes,
occasionally up to eighty minutes, in behaving mice with minimal signal loss across neuron types
and brain regions. This extended recording capability reveals plasticity changes throughout
intracranial electrical neuromodulation, highlighting the novel insights enabled by prolonged
voltage imaging.

**Main Text**

The new generation of genetically encoded voltage indicators (GEVIs) enables high
spatiotemporal resolution imaging of membrane voltage (Vm) from individual neurons in the
mammalian brain[1–9]. In addition to capturing spiking, GEVIs can detect subthreshold dynamics
from genetically defined neurons, which has not been possible with traditional extracellular
recording or calcium imaging methods[10–13]. However, most GEVIs still suffer from poor
photostability, generally limiting recording duration to about 10-20 minutes[1–9], but success for
longer recordings lasting tens of minutes has been demonstrated with GEVI JEDI-2P (~40 mins)[7]
and ASAP5 (~1 hour)[6] using two-photon microscopy. The newly developed green GEVI
ElectraOFF[14], exhibits high voltage sensitivity, excellent photostability, and is compatible with a
widefield microscope that permits simultaneous high-speed recording of many cells. Furthermore,
ElectraOFF is fully genetically encoded, without needing any chemical cofactors, greatly
improving its applicability *in vivo*. We tested the performance of ElectraOFF in different types of
neurons across the hippocampal CA1, visual cortex and striatum in voluntarily locomoting mice,
and used ElectraOFF to probe cellular plasticity during chronic electrical stimulation widely used
in clinical neuromodulation.

To characterize the performance of ElectraOFF in the brain of awake mice, we transduced
hippocampal CA1 pyramidal neurons with AAV9-CaMKII-ElectraOFF and installed a glass cranial
window for chronic optical access (see **Methods**). ElectraOFF-expressing neurons were imaged
using a custom widefield fluorescence microscope at a frame rate of 667 Hz, while mice were
head-fixed and voluntarily locomoting. Image frames were collected for 30 s every minute to allow
for data streaming, and the recorded image stacks were motion corrected and processed offline

to obtain the Vm traces and spike timing for each neuron. We started imaging with a low LED
power at ~20 mW/mm$^2$ to minimize photobleaching and gradually increased the power every 5-
20 min to compensate for the loss of fluorescence (up to ~200 mW/mm$^2$). Using this experimental
protocol, we were able to obtain recordings that lasted for up to 80 min without compromising
spike quality (**Fig. 1a-c and Fig. S1**).

To quantify ElectraOFF's photostability, we computed raw Vm fluorescence (F), normalized spike
amplitude (dF/F; with dF defined as the fluorescence difference between the spike peak and the
minimum value within the preceding 5 ms), spike signal-to-baseline ratio (SBR; defined as spike
dF divided by variance of spike-removed Vm fluctuations during each 30-second trial), and spike
rate (defined as the mean rate in each 10-second bin; see **Methods**). Photobleaching rates were
estimated by linearly fitting across all data collected at the same LED power, except spike rate
which was fit across the entire recording because little relationship was detected between LED
power and spike rate (**Fig. S2m**). We found that for the example neuron shown in **Fig. 1a-c**, raw
F declined at 0.21 ± 0.18% per minute (mean ± s.d. here and throughout unless otherwise stated;
n=5 testing periods with varying LED power) throughout the 80 min recording time (**Fig. 1d**), spike
rate decreased at 0.5% per minute ($R^2$=0.14, p <0.001) (**Fig. 1g**), while spike dF/F,  and SBR
remained remarkably stable with decay rates of 0.12 ± 0.76 and 0.04 ± 0.61% per minute
respectively (**Fig. 1e, f and Fig. S2a**). Thus, while ElectraOFF exhibits some photobleaching of
raw F and a decrease in detectable spikes, compensatory LED adjustments can maintain stable
spike dF/F and SBR, ensuring spike quality and enabling reliable long-term Vm recordings in CA1
of awake animals.

To systematically evaluate the performance of ElectraOFF across cell types and brain regions in
behaving mice, we expressed ElectraOFF in striatal cholinergic (ChI) neurons of Chat-Cre mice
and in visual cortical parvalbumin (PV) interneurons of PV-Cre mice using AAV9-CAG-DIO-
ElectraOFF. ElectraOFF supported stable voltage imaging that routinely lasted for about 10 min
in striatal ChIs and for tens of minutes in cortical PVs (**Fig 2a, b, n**). For the three example neurons,
the bursting ChI, the tonically firing ChI, and the PV neuron, we detected minimal photobleaching
of raw F at 2.9 ± 1.0%, 2.3 ± 5.0% and 1.3 ± 0.55% per minute respectively (**Fig. 2c, o, Fig. S2e**),
moderate decay in spike rate at 7.2%, 8.9%, and 2.6% per minute ($R^2$=0.78, p<0.001; $R^2$=0.52,
p<0.001; $R^2$=0.23, p<0.001), and negligible decay in spike dF/F, and spike SBR (**Fig. 2d-f, p-r**
**and Fig. S2f-h**). Across the population of recorded CA1 pyramidal neurons, striatal ChIs and
cortical PVs, raw F photobleached at 0.72 ± 0.30%, 1.3 ± 0.89%, 2.7 ± 2.7% per minute
respectively (**Fig. 2t and Fig. S2i**), and spike rate decayed at 3.4 ± 2.4%, 8.5 ± 1.7%, 5.0 ± 6.2%
per minute respectively (**Fig. 2w, Fig.S2j**), while dF/ F, spike SBR showed little decay (**Fig. 2u, v**
**and Fig. S2k,l**). Furthermore, ElectraOFF reliably distinguished striatal ChIs exhibiting spike
bursts (**Fig. 2g**) versus tonic firing (**Fig. 2h).** Aligning Vm to spikes revealed prominent Vm delta
oscillation in bursting ChIs (**Fig. 2i, k**), but not in tonically firing ChIs (**Fig. 2j, l**), underscoring
ElectraOFF's ability to detect subthreshold Vm dynamics beyond spike timing.

To test the ability of ElectraOFF in capturing behaviorally relevant stimuli, we presented mice with
4 Hz visual flickers for 10 s in the middle of each 30-second-long imaging trial (**Fig. 2m, n**). We
found that visual flicker evoked a transient Vm depolarization in visual cortical PV neurons that
lasted for about 1 s, which was followed by sustained hyperpolarization throughout the stimulation

period and a pronounced rebound after stimulation offset (**Fig. 2s**). Interestingly, while spiking
increased during the initial Vm depolarization, spike rate quickly returned to pre-stimulation levels
even though Vm remained hyperpolarized, highlighting ElectraOFF's ability to capture long-lasting
synaptic inhibition during behavior.

Finally, we investigated the sustained effects of intracranial electrical stimulation on CA1 neurons
to understand the impact of prolonged stimulation on cellular plasticity. We delivered biphasic
square-wave pulses (100 µs/phase) at 40 Hz that have been shown to entrain CA1 neurons[15].
Stimulation was delivered locally for 1 min in the middle of each 90 s trial (**Fig. 3a**) at a current
intensity sufficient to evoke a neural response without inducing detectable behavioral changes
(**Methods**). Each neuron was tested for 5 trials with an intertrial interval of 1.5 min. Consistent
with previous observations, of the 49 neurons recorded, 27 (55%) were significantly entrained
(**Fig. 3f**), exhibiting greater Vm spectral power at 40 Hz during stimulation than baseline. Across
all recorded neurons, stimulation did not change population firing rate, but it evoked a pronounced
Vm hyperpolarization immediately after stimulation onset that lasted for ~1 s, which was followed
by Vm depolarization that stabilized after ~15 s (**Fig. 3b**). To further evaluate the time-course of
the evoked responses, we divided the stimulation period into four consecutive 15-second epochs
and compared the 40 Hz spectral power of Vm across epochs for every trial (**Fig. 3c**). We found
that entrainment strength across the entrained neurons declined overtime within each trial, with
the highest 40 Hz power observed during the first 15 s (**Fig. 3d, e**). Furthermore, we detected a
gradual reduction in entrainment strength across successive trials (**Fig. 3g**). Thus, prolonged
electrical stimulation-evoked Vm changes progressively decay over time, underscoring the
capacity of ElectraOFF to uncover plasticity mechanisms in neural circuits.

**Discussion**

ElectraOFF exhibits minimal photobleaching in the brains of behaving mice, with raw fluorescence
intensity decaying at ~0.5-1.3% per minute. This modest reduction allows for routine imaging for
tens of minutes across cell types and brain regions. By progressively increasing LED power over
time during prolonged imaging to minimize excitation light exposure, we were able to image most
neurons for tens of minutes, and up to 80 minutes in CA1, without compromising spike quality. As
spike detection depends on spike peak fluorescence, we noted a spike rate decay at a low rate
of 6% per minute, while maintaining spike dF/F and spike SBR. Photobleaching is intrinsic to
fluorescence imaging, and it is important to include control analysis when examining long term
changes to rule out photobleaching effects. Voltage-dependent ElectraOFF fluorescence can be
readily recorded with a high-speed widefield microscope, allowing for simultaneous recording
from multiple neurons which cannot be easily achieved with high-speed two-photon or confocal
microscopy techniques that have limited field-of-view sizes[9,16,17].

Using ElectraOFF, we captured prominent Vm hyperpolarization upon visual flicker presentation
in visual cortical interneurons and upon electrical stimulation in CA1 pyramidal neurons. These
results highlight the unique advantage of voltage imaging in detecting synaptic inhibition, not
possible using conventional extracellular electrodes or calcium imaging. As Vm determines spike
timing, the ability to record Vm over an extended time unlocks new frontiers for analyzing the

subthreshold decision-making processes of individual neurons during behavior and pathology. As
an example, we demonstrated that upon sustained stimulation, either through natural visual flicker
presentation or artificial intracranial electrical current delivery, Vm exhibits complex temporal
changes, switching between depolarization and hyperpolarization which was often not
accompanied by spiking changes. In particular, during chronic intracranial electrical stimulation,
evoked Vm entrainment progressively decayed over time, revealing robust plasticity mechanisms
related to clinical electrical neuromodulation.

**Figure 2. ElectraOFF is photostable across cell types and brain regions during both intrinsic and evoked activity. a**) left, Vm from an example striatal cholinergic interneuron (ChI) with bursty firing in a voluntarily locomoting mouse. right, zoomed-in views of Vm for the periods i-iv shown on the left. **b**) Similar to a, but for a ChI with tonic firing. **c**) Raw F of the example bursting ChI over 14 min of the recording. The adjustments of LED power are indicated by gaps. Each dot represents the mean raw F of a 30 s trial. The black lines are linear fits with periods having constant LED power. **d**) Similar to c), but for normalized spike dF/F. Each dot corresponds to a spike. **e**) Similar to d), but for spike SBR. **f)** Spike rate over time. LED power shifts are indicated by vertical black lines and the red line is the linear fit across the entire recording. Each dot corresponds to a 10 s bin of the recording. **g**) The inter-spike interval (ISI) return map of all spikes recorded for the example bursting ChI shown in a) (n = 1411 spikes). **h**) Similar to g), but for the example tonic ChI shown in b) (n = 1203 spikes). **i**) Spike-triggered Vm for example bursting ChI. Black line is the mean, and shaded areas indicate 95% confidence interval. **j**) Similar to i), but for the example tonic ChI. **k**) Spike-triggered Vm power spectrogram for the bursting ChI. **l**) Similar to k), but for the tonic ChI. **m**) Experimental schematic for visual flicker presentation to a head-fixed voluntarily locomoting mouse, while recording visual cortical PV interneurons. Visual flickers were delivered at 4 Hz, for 10 s during each 30 s trial. **n**) Vm traces of an example PV neuron across 35 min recording. **o-r**) Similar to c-f), but for the example PV. **s**) Vm (black) and spike rate (red) changes aligned to visual flicker onset. Shading areas indicate standard error of mean (n = 70 trials). **t**) The photobleaching rates of raw F for CA1 neurons (n=21 neurons), striatal ChIs (n=8 neurons), and visual cortical PV neurons (n=3 neurons). Each dot represents a neuron. Red lines are the mean, and the boxes mark the interquartile range. **u-w**) Similar to t), but for spike dF/F, SBR, and spike rate respectively (n=21 for CA1 neurons, n=8 for ChIs, and n=3 for PVs).

**Figure 3. Sustained ElectraOFF recordings revealed intracranial electrical stimulation induced cellular plasticity effects. a)** Top, schematic of the experimental setup. Bottom,

stimulation protocols. Stimulation was delivered locally in CA1, while recording Vm of CA1 pyramidal neurons. Each field of view underwent five stimulation trials with a 1.5 min inter-trial interval. Each trial included 15 s pre-stimulation baseline, 60 s electrical stimulation (40 Hz, biphasic pulses), and 15 s post-stimulation periods. **b**) Stimulation evoked change in Vm (black) and spike rate change (red). Vm change is normalized to mean spike height and shifted by the mean pre-stimulation baseline Vm. Shaded areas represent standard error of mean (SEM) (n = 49 neurons). **c**) Trial-wise average Vm traces with SEM shown as shading (n=49). The stimulation periods were subdivided into 15-second epochs as highlighted on top). Vertical lines denote stimulation onset and offset (solid) and 15-second subdivisions (dotted). **d**) Power spectral density (PSD) of Vm traces, computed for each 15-second epoch and normalized to the baseline PSD for each trial. Gray lines indicate mean PSD across neurons; shaded regions indicate SEM (n=49). **e**) Violin plots of 40 Hz PSD across stimulation epochs for entrained neurons (n=27) (Statistical details in Supplemental Table 1). **f**) Vm spectrograms of entrained (left, n=27) and not-entrained (right, n=22) neurons aligned to stimulation onset. **g**) Similar to e) but compared across trials for each epoch. (Statistical details in Supplemental Table 1). Violin plots depict the kernel density overlaid with box plots showing the interquartile range (1x, 1.5x). The white lines in the boxes are the median. All tests are two-tailed. $*p < 0.05$, $**p < 0.01$, $***p < 0.005$.

192    **Supplemental Table 1:** Statistical Details for Fig. 3

| Stats Table | | | | | |
|---|---|---|---|---|---|
| | | | | | |
| Fig 3e p-values (Friedman test with Bonferroni multiple comparisons correction) | | | | | |
| | Trial 1 | Trial 2 | Trial 3 | Trial 4 | Trial 5 |
| 0-15s   vs 15-30s | $4.36 \times 10^{-6}$ | 0.161 | 0.027 | 1 | $1.26 \times 10^{-5}$ |
| 0-15s   vs 30-45s | 0.009 | $4.36 \times 10^{-6}$ | 0.122 | 0.001 | 0.013 |
| 0-15s   vs 45-60s | $9.27 \times 10^{-5}$ | $8.16 \times 10^{-7}$ | 0.161 | 0.271 | $2.52 \times 10^{-6}$ |
| 15-30s vs 30-45s | 0.439 | 0.037 | 1 | 0.027 | 0.550 |
| 15-30s vs 45-60s | 1 | 0.013 | 1 | 1 | 1 |
| 30-45s vs 45-60s | 1 | 1 | 1 | 0.550 | 0.271 |
| | | | | | |
| Fig 3g p-values (Friedman test with Bonferroni multiple comparisons correction) | | | | | |
| | 0-15s | 15-30s | 30-45s | 45-60s | |
| Trial 1 vs Trial 2 | 1 | 0.160 | 0.707 | 1 | |
| Trial 1 vs Trial 3 | 0.478 | 1 | 1 | 1 | |
| Trial 1 vs Trial 4 | $8.10 \times 10^{-8}$ | 1 | $5.08 \times 10^{-5}$ | 0.707 | |
| Trial 1 vs Trial 5 | $5.08 \times 10^{-5}$ | 0.020 | 0.008 | 0.002 | |
| Trial 2 vs Trial 3 | 1 | 0.002 | 1 | 1 | |
| Trial 2 vs Trial 4 | $2.21 \times 10^{-5}$ | 0.001 | 0.059 | 1 | |
| Trial 2 vs Trial 5 | 0.004 | $3.62 \times 10^{-7}$ | 1 | 0.059 | |
| Trial 3 vs Trial 4 | 0.002 | 1 | $7.52 \times 10^{-4}$ | 1 | |
| Trial 3 vs Trial 5 | 0.098 | 0.707 | 0.059 | 0.020 | |
| Trial 4 vs Trial 5 | 1 | 1 | 1 | 0.478 | |

**Figure S1. An example 80-minute long voltage recording from two adjacent CA1 neurons in an awake mouse. a**) The same imaging field of view as in Fig. 1a, showing three neurons expressing ElectraOFF. The black arrow indicates the neuron whose Vm is shown in Fig. 1a, and the blue arrow indicates a neuron whose Vm is shown in b). Spikes are indicated by red dots. **b**) Vm throughout the 80-minute-long recording for the neuron indicated by the blue arrow in a. **c**) Zoomed in view of Vm from the two simultaneously recorded neurons during periods highlighted by the corresponding boxes in b). The blue and black traces are from the neurons indicated by the blue and black arrows in a) respectively.

**Figure S2. Detailed characterization of ElectraOFF photostability across cell types. a**) The relationship between dF/F and spike SBR of the example CA1 neuron shown in Fig. 1a. Each dot corresponds to a spike. The black line indicates the linear fit (n=24365 spikes, $R^2$=0.362, k=0.002, p < 0.001). **b-d**) Similar to a), but for the example PV neuron (n=1435 spikes, $R^2$=0.004, k=-0 0004, p < 0.05), bursting ChI (n=1411 spikes, $R^2$=0.122, k=0.003, p < 0.001), and tonic ChI (n=1208 spikes, $R^2$=0.155, k = 0002, p<0.001) from Fig. 2. **e**) Raw F of the example tonic ChI from Fig. 2 over 14 min of the recording. The adjustments of LED power are indicated by gaps. Each dot represents the mean raw F of a 30 s trial. The black lines are linear fits with periods having constant LED power. **f)** Similar to e), but for normalized spike dF/F. Each dot corresponds to a spike. **g)** Similar to f), but for spike SBR. **h)** Spike rate over time. LED power shifts are indicated by vertical black lines and the red line is the linear fit across the entire recording. Each dot corresponds to a 10 s bin of the recording. **i)** The average raw F over time for all neurons. Each dot corresponds to the mean for each 30 s long recording period normalized to the first point of the given neuron. Cell type is color coded (Blue: 21 CA1 neurons (n=1174 recording periods); Green: 3 visual cortical PV neurons (n=120 recording periods); Brown: 8 striatal ChIs (n=126 recording periods). **j-l)** Similar to i), but for the spike rate, normalized dF/F, and SBR. The colored lines indicate the linear fit trendline for each cell type. **m)** Similar to j) but spike rate vs raw F (proxy for LED power). The black line is the linear fit across all neurons regardless of cell type and indicates there is a very weak correlation between the spike rate and raw F (n = 1420, k=0.007, $R^2$=0.003, p=0.03).

**Methods:**

**Mouse Surgical Preparation**

All animal experiments were performed in accordance with the National Institute of Health Guide for Laboratory Animals and approved by the Boston University Institutional Animal Care and Use and Biosafety Committees. Same-sex littermates were housed together prior to surgery and singly housed post-surgery. Cages were enriched with igloos or running wheels. Animal facilities were maintained around 70° F and 50% humidity and mice were kept on a 12 h light/dark cycle. A total of 3 adult mice, 1 ChAT-Cre female, 1 PV-Cre male (JAX stock #017320, B6.129P2-Pvalb$^{tm1(cre)Arbr}$/J, Jackson Laboratory), and 1 C57BL/6 male (Charles River Laboratories Inc.), 16-24 weeks old at the start of the experiment were used in this study.

Under isoflurane general anesthesia, mice were surgically implanted with a sterilized recording ydevice for hippocampal or striatal recordings, or a glass coverslip for cortical recordings. Mice were provided with sustained release (SR) buprenorphine (3.25 mg/kg) pre-operatively that provides 72 hours of analgesia. Mice were singly housed to avoid damage to the recording apparatus after surgery.

For CA1 recordings, the recording device consisted of a custom imaging window attached to a guide infusion cannula and two stimulation electrodes. For striatal recordings, the recording device consisted of a custom imaging window attached to a guide infusion cannula. The imaging window consisted of a stainless-steel cannula (outer diameter 0.317 cm, inner diameter 0.236 cm, height 1.75 mm for CA1 window and 2 mm for striatal window; Item 3ADD3; Grainger), adhered to a circular coverslip (size 0, outer diameter 3 mm; No. CS-3R-0; Warner Instruments) using an ultraviolet-curable adhesive (Norland Products). The infusion guide cannula (8 mm, 26 gauge; P1 Technologies) was soldered at a 60° angle relative to the coverslip with the end flush to the base of the imaging cannula. The two stimulation electrodes (polyamide-coated steel core wire, diameter 127 μm, No. 7N003736501F; P1) were soldered onto dip pins (No. ED85100-ND; DigiKey), and secured to the infusion cannula with adhesive gel (Loctite 454; Loctite SF 713). The electrodes were bent so that the male end of the dip pin was perpendicular to the infusion cannula, while the electrode wire was parallel to the infusion cannula. The tips of the electrodes were about 3 mm apart and terminated below the imaging window. The recording apparatus was implanted over CA1 (centered at the stereotaxic coordinates: AP −2.0 mm, ML 1.7 mm, DV −1.5 mm from pia), or over the striatum (centered at the stereotaxic coordinates: AP 0.5 mm, ML 1.8 mm, DV −2 mm from pia) after gently aspirating the overlying cortical tissue and thinning the corpus callosum. A custom aluminum head-plate was attached to the skull posterior to the window via dental cement (Parkell, S380 and Stoelting, 5145). After a full recovery from the surgery (about 10 days), mice with a recording apparatus were infused with 1.5 μL of AAV9-CaMKII-ElectraOFF or AAV9-CAG-DIO-ElectraOFF (>10$^{13}$ GC/mL, Shanghai Sunbio Medical Biotechnology and Charles River Laboratories), at a speed of 200 nL per minute, using a 10 μL syringe (World Precision Instruments, NANOFIL 10 μL syringe) controlled by a micro-infusion pump (Micro-2T, World Precision Instruments) attached to 28 gauge PTFE tubing (No. STT-28; Component Supply Co.) with a 33 gauge infusion cannula (8 mm with 1 mm projection; P1 Technologies). The infusion cannula was removed 10 min after the infusion had been completed to improve AAV spread.

For cortical recordings, a small craniotomy was created over the visual cortex (centered at the
stereotactic coordinate: AP -3.6 mm, ML 2.3 mm), and a total of 1.5 µL of AAV9-CAG-DIO-
ElectraOFF was injected via a 10 µL syringe (World Precision Instruments, NANOFIL 10 µL
syringe) controlled by a micro-infusion pump (MicroPump4, World Precision Instruments) in 3
locations within the craniotomy, avoiding major blood vessels, ~150 µm below pia. After viral
injection the syringe was left in place for 10 min at each location, and then the syringe was
removed. A circular glass coverslip (size 0, outer diameter 3 mm; No. CS-3R-0; Warner
Instruments) was then pressed onto the cortex to ensure stable optical recordings and was
secured to the skull with ultraviolet-curable cement (Tetric EvoFlow; Ivoclar). A custom aluminum
head-plate was attached to the skull posterior to the window via dental cement (Parkell, S380 and
Stoelting, 5145).

## Voltage Imaging

Animals were imaged starting 2 weeks post viral infusion in the CA1, 3 weeks in the striatum, and
4 weeks in the visual cortex, to ensure sufficient ElectraOFF expression. ElectraOFF voltage
imaging was performed with a custom wide-field fluorescence microscope equipped with a
sCMOS camera (Hamamatsu Photonics, C13440-20CU) and a 40X objective (Nikon, CFI Apo
NIR 40×/0.8 NA W). A blue 470 nm LED (Thorlabs, M470L4) controlled by a T-Cube LED driver
(LEDD1B, low gain; Thorlabs Inc.) was sequentially coupled to a 505 nm long-pass filter (Thorlabs,
DMLP505R), excitation filter (FF01-390/482/532/640-25), quadband dichromatic mirror (Semrock
Di03-R405/488/532/635), a multi-band dichromatic mirror (Chroma Technology Corp.,
ZT405/514/635rpc), and a 525 ± 22.5 nm bandpass emission filter (Semrock, FF01-525/45)
emission filter.

ElectraOFF fluorescence images were collected using HCImage Live (Hamamatsu Photonics),
triggered via a TTL pulse generated by a NI DAQ (National Instruments, BNC-2110 and PCIe-
6323) controlled by MATLAB. Similarly, the electrical stimulator or visual flicker LED, if applicable,
was also triggered by TTL pulses. HCImage Live collected up to 90 s of data (20000 or 60000
frames) with an exposure time set at 1.5 ms. HCImage was set to capture 128x512 pixels with
2x2 binning, corresponding to 41x164 µm$^2$. Each trial was saved in a separate DCAM file. The
files were analyzed offline with MATLAB. Camera frame exposure from the Hamamatsu camera,
electrical stimulation pulses, and visual flicker pulses were simultaneously recorded as TTL pulses
with an Open Ephys acquisition board (Open Ephys, SKU: OEPS-9030) at a rate of 30 kHz to
ensure proper alignment during further analysis offline[18].

## Electrical Stimulation

Electrical stimulation was delivered by a constant current stimulator (Model 4100; A-M Systems).
The stimulation waveform was a cathode-leading biphasic square wave pulse delivered at 40 Hz
with a 200 µs total pulse width (100 µs/phase). The current amplitude used was empirically
determined by delivering a short burst of stimulation with a current amplitude starting at 200 µA
and increasing in 25 µA steps until we observed an obvious ElectraOFF response but not any
behavioral changes. Electrical stimulation delivery was triggered using a MATLAB generated TTL
pulse for 60 s, which started 15 s after triggering the first camera frame.

**Visual Flicker Presentation**

Visual flickers were presented via a 5 mm white LED (Amazon, MCIGICM) positioned in front of the animal. LED was pulsed at 4 Hz with a 50% duty cycle, for a total of 10 s, at an intensity of 20.5 μW in a dimly lit room. LED was triggered using a NI-DAQ (National Instruments, BNC-2110 and PCIe-6323), 10 s after the first camera frame.

**ElectraOFF fluorescence pre-processing to obtain Vm traces**

All data analysis was performed offline using custom MATLAB (2024a, Mathworks Inc.). ElectraOFF fluorescence Videos were motion-corrected using a pairwise rigid motion correction algorithm as described previously[19]. Briefly, all trials for a single experimental recording were first concatenated. The shift of each frame was identified by calculating the maximum cross-correlation coefficient between each frame and the reference image. Regions of interest (ROIs) corresponding to individual neurons were subsequently manually identified using a max-minus-min projection image of the motion corrected data and the drawPolygon function (MATLAB). The ElectraOFF fluorescence trace was extracted by averaging all the pixels within the ROI to obtain the ROI trace. We then estimate the background trace for each ROI trace by averaging across all pixels of a manually selected area within the same imaging field of view, free of neuropil. The correlation coefficients of the background trace and the ROI trace were then obtained using the linear fit (y=mx+b), where m and b were found by minimizing the error between the background trace, x, and the ROI trace, y. The background trace is then scaled by plugging back in for x using the newly estimated coefficients to calculate a new y representing the portion of the ROI trace explainable by background fluctuations. ElectraOFF trace was then computed as the ROI trace minus the scaled background trace. ElectraOFF traces were further denoised by removing values greater than 10 standard deviations from the mean in the positive going direction. We then subtracted the median value of the trace, yielding the Vm trace for each neuron.

**Spike Identification, spike amplitude and spike-to-baseline ratio (SBR) calculation**

To identify spikes, we first computed the baseline Vm fluctuations by averaging two traces yielded by (1) applying a moving average filter (window = 150 ms) of the Vm trace and, separately, (2) applying a second-order Savitsky-Golay filter (window = 150 ms) to the Vm trace. Baseline Vm fluctuations were then subtracted from the Vm trace to isolate the high frequency components ($Vm_{high}$). Spikes were then detected using $Vm_{high}$ as peaks greater than are 5 (CamKs) or 6 (ChIs and PVs) standard deviations away from the mean. Spike amplitude was calculated using the Vm trace as the difference between the spike peak and the minimum fluorescence value within the 3 frames (~5 ms) preceding the spike peak. Identified spikes with an amplitude greater than 35 were considered artifacts and excluded from analysis. The SBR of each spike was computed as the spike amplitude divided by the approximated baseline noise. Baseline noise was defined for each trial as the standard deviation of the positive-half of $Vm_{high}$ to exclude contributions from the negative-going spikes and slow baseline Vm fluctuations.

**Photostability Analysis**

To assess the photostability of ElectraOFF *in vivo* we measured four parameters over time (1)
mean ElectraOff raw fluorescence (F) for each 30 s trial, (2) normalized spike amplitude (dF/F)
for each spike, (3) spike SBR for each spike, and (4) spike rate calculated for each 10 s bin.
Because the excitation LED power was occasionally adjusted to optimize recording length, raw
F, dF/F and SBR changes were computed for each period of constant LED power. Within each
period with constant LED power, data were linearly fit across time, and the percent-change per
minute was calculated as the slope of the fit line divided by the starting y-value of the fit. Since
spike rate was independent of LED power, changes in spike rate were estimated by linearly
fitting over the full length of the recording.

**Spike triggered Vm Analysis.**

The spike-triggered Vm waveform for a neuron was calculated by first taking a 1-second window
centered around each detected spike. Each window was scaled by its spike height, and then
averaged across all spikes for the neuron. For visualization, final traces were flipped to follow the
convention that spikes are positive-going.

When calculating the spike-triggered average Vm power spectra, spikes were first removed by
replacing the 5 data points centered around each spike peak with linearly interpolated values to
minimize broadband power changes at spike peaks. The power spectrum of the spike removed
trace was then obtained for each 30 s trial using Matlab's continuous wavelet transform function
(cwt) (Morlet Wavelet, VoicesPerOctave = 12). The spectral trace was then windowed around
each spike (1 s windows) and averaged across all spikes to obtain the spike triggered average
Vm power spectra for a neuron.

**Visual flicker and electrical stimulation evoked responses.**

To assess the visual flicker and electrical stimulation evoked responses, Vm traces were first
flipped, normalized to spike height, detrended, referenced relative to the median baseline Vm,
and averaged across neurons. Briefly, we first obtained a scaled trace by normalizing every flipped
trial trace to its average spike height. The scaled trace was then lowpass filtered (fifth order
butterworth with 50 Hz cutoff) and fit with a linear trendline. This linear trendline was then
subtracted from the scaled trace to account for photobleaching. A trial average trace was then
calculated for each neuron by averaging these detrended traces. The median of the baseline
period was then subtracted from the average trace, and the traces were then smoothed for
visualization with a moving median filter (150 ms; 100 frames). The final reported mean trace and
standard error were calculated using these smoothed trial-averaged traces.

To assess the effect of neuromodulation on firing rate, a binarized spike raster (each frame with
a spike is 1) was generated for each trial, and the trial-average raster then calculated for each
neuron. A moving mean filter (300 ms; 200 frames) was applied to calculate the firing rate trace.
For each trial-average trace, the average baseline firing rate was calculated and subtracted to
estimate the change in firing rate (Hz) relative to baseline, and the traces were then smoothed for
visualization with a moving mean filter (150 ms; 100 frames) The final reported mean trace and
standard error were calculated using these smoothed trial-averaged traces.

To classify neurons as entrained or not-entrained by 40 Hz electrical stimulation, each cell's
normalized trial traces were broken into 15 s chunks as seen in **Fig. 3c** and the power spectral
density (PSD) of each of these chunks was calculated individually using Matlab's fast-Fourier
transform function (fft). The PSD around 40 Hz (39.5-40.51 Hz window) for each of the 15-second
stimulation epochs was then Z-scored relative to the baseline PSD distribution in that frequency
window. Neurons were classified as entrained if the normalized PSD at 40 Hz during the first 15
s of stimulation in any trial was greater than 3. Otherwise, neurons were classified as not-
entrained.
391

**Acknowledgments:** We thank members of Han Lab for their help throughout the study. We also
thank Dr. Kiryl Piatkevich for providing the ElectraOFF plasmids and AAV9 particles. X. H.
acknowledges funding from the NIH (1RF1NS129520, 1R01MH122971, and 1R01NS115797)
and NSF (2002971-DIOS, and 1955981-CIF). C.R. acknowledges funding from NSF Graduate
Research Fellowship 2021324226.

**Author contributions:** Y.W., C.R. and X.H. designed the study. Y.W. collected experimental data.
C.R. analyzed the data. Y.Z. contributed some experimental data. Y.W., C.R. and X.H. prepared
the manuscript, and all authors edited the manuscript. X.H. oversaw all aspects of the project and
supervised the study.

**Declaration of interests:** The authors declare no competing interests.

**Reference**

1. Piatkevich, K. D. *et al.* Population imaging of neural activity in awake behaving mice. *Nature* **574**, 413–417 (2019).

2. Tian, H. *et al.* Video-based pooled screening yields improved far-red genetically encoded voltage indicators. *Nat Methods* **20**, 1082–1094 (2023).

3. Kannan, M. *et al.* Dual-polarity voltage imaging of the concurrent dynamics of multiple neuron types. *Science* **378**, eabm8797 (2022).

4. Abdelfattah, A. S. *et al.* A general approach to engineer positive-going eFRET voltage indicators. *Nat Commun* **11**, 3444 (2020).

5. Abdelfattah, A. S. *et al.* Sensitivity optimization of a rhodopsin-based fluorescent voltage indicator. *Neuron* **111**, 1547-1563.e9 (2023).

6. Hao, Y. A. *et al.* A fast and responsive voltage indicator with enhanced sensitivity for unitary synaptic events. *Neuron* **112**, 3680-3696.e8 (2024).

7. Liu, Z. *et al.* Sustained deep-tissue voltage recording using a fast indicator evolved for two-photon microscopy. *Cell* **185**, 3408-3425.e29 (2022).

8. Platisa, J. *et al.* High-speed low-light in vivo two-photon voltage imaging of large neuronal populations. *Nat Methods* **20**, 1095–1103 (2023).

9. Xiao, S. *et al.* Large-scale deep tissue voltage imaging with targeted-illumination confocal microscopy. *Nat Methods* **21**, 1094–1102 (2024).

10. Dana, H. *et al.* High-performance calcium sensors for imaging activity in neuronal populations and microcompartments. *Nat Methods* **16**, 649–657 (2019).

11. Zhang, Y. *et al.* Fast and sensitive GCaMP calcium indicators for imaging neural populations. *Nature* **615**, 884–891 (2023).

12. Kodandaramaiah, S. B. *et al.* Multi-neuron intracellular recording in vivo via interacting autopatching robots. *eLife* **7**, e24656 (2018).

13. Tao, C., Zhang, G., Xiong, Y. & Zhou, Y. Functional dissection of synaptic circuits: in vivo patch-clamp recording in neuroscience. *Front Neural Circuits* **9**, 23 (2015).

14. Zhang, H. *et al.* Bright and Photostable Voltage Sensors Derived from mBaoJin. 2025.05.30.657123 Preprint at https://doi.org/10.1101/2025.05.30.657123 (2025).

15. Lowet, E. *et al.* Deep brain stimulation creates informational lesion through membrane depolarization in mouse hippocampus. *Nat Commun* **13**, 7709 (2022).

16. Villette, V. *et al.* Ultrafast Two-Photon Imaging of a High-Gain Voltage Indicator in Awake Behaving Mice. *Cell* **179**, 1590-1608.e23 (2019).

17. Mertz, J. *Introduction to Optical Microscopy*. (Cambridge University Press, 2019).

441    18.    Siegle, J. H. *et al.* Open Ephys: an open-source, plugin-based platform for multichannel
442    electrophysiology. *J Neural Eng* **14**, 045003 (2017).

443    19.    Pnevmatikakis, E. A. & Giovannucci, A. NoRMCorre: An online algorithm for piecewise
444    rigid motion correction of calcium imaging data. *Journal of Neuroscience Methods* **291**, 83–94
445    (2017).