CONTEXT-DEPENDENT VARIABILITY OF HIF HETERODIMERS INFLUENCES INTERACTIONS WITH MACROMOLECULAR AND SMALL MOLECULE PARTNERS

Joseph D. Closson[1,2], Xingjian Xu[1,2], Meiling Zhang[1], Tarsisius T. Tiyani[1,2], Leandro Pimentel Marcelino[1,3], Eta A. Isiorho[1], Jason S. Nagati[4,5], Joseph A. Garcia[4,5], Kevin H. Gardner[1,3,6,*]

*: direct correspondence to kgardner@gc.cuny.edu

[1]: Structural Biology Initiative, CUNY Advanced Science Research Center, New York, NY 10031
[2]: Ph.D. Program in Biochemistry, The Graduate Center – City University of New York, New York, NY 10016
[3]: Department of Chemistry and Biochemistry, City College of New York, New York, NY 10031
[4]: Department of Medicine, Columbia University Irving Medical Center, New York, NY 10032
[5]: Departments of Medicine and Research & Development, James J. Peters Veterans Affairs Medical Center, Bronx, NY 10468
[6]: Ph.D. Programs in Biochemistry, Biology, and Chemistry, The Graduate Center – City University of New York, New York, NY 10016

ORCIDs:
Joseph D. Closson: 0000-0002-4618-3119 / Xingjian Xu: 0000-0002-9609-4755 / Meiling Zhang: 0000-0002-0670-6018 / Tarsisius T. Tiyani: 0000-0002-2806-4079 / Leandro Pimentel Marcelino: 0000-0001-6067-0725 / Eta A. Isiorho: 0000-0002-6242-9297 / Jason S. Nagati: 0000-0002-4507-6581 / Joseph A. Garcia: 0000-0002-5621-7538 / Kevin H. Gardner: 0000-0002-8671-2556

Keywords: Hypoxia inducible factors (HIFs), aryl hydrocarbon receptor nuclear translocator (ARNT), clear cell renal cell carcinoma (ccRCC), hypoxia, transcription factors, small molecule binding, higher order complex formation

Running title: Variability of HIF complexes influences intermolecular interactions

Abstract

Hypoxia inducible factors (HIFs) are transcription factors that coordinate cellular

responses to low oxygen levels, functioning as an α/β heterodimer which binds a short hypoxia

response element (HRE) DNA sequence. Prior studies suggest HIF/HRE complexes are

augmented by the binding of additional factors nearby, but those interactions are not well

understood. Here, we integrated structural and biochemical approaches to investigate several

functionally relevant HIF assemblies with other protein, small molecule, and DNA partners.

First, we used cryo-electron microscopy (cryo-EM) to establish HIF-1 and HIF-2 self-assemble

to form novel "dimer-of-heterodimers" (DoHD) complexes on extended human EPO enhancer

sequences, with one heterodimer bound at a canonical HRE site and the second binding in an

inverted fashion to an HRE-adjacent sequence (HAS) 8 bp away. Consistent with ARNT PAS-B

domains predominating interactions within a DoHD, we found HIF-1 and HIF-2 co-assemble

mixed DoHD complexes on the same DNA. Second, we saw that despite the increased

complexities of the larger complexes, ligands for the isolated ARNT or HIF-2α PAS-B domains

are still capable of binding and disrupting both the heterodimer and DoHD complexes, albeit

with variable potencies depending on the ligand. Finally, we combined cryo-EM and hydrogen-

deuterium exchange by mass spectrometry (HDX-MS) to show how HIF-1 and HIF-2

heterodimers engage the transforming acidic coiled-coil containing protein 3 (TACC3)

coactivator via both ARNT and HIF-α subunits, though this was unseen in the larger DoHD. Our

findings highlight the importance of both molecular context and dynamics in biomolecular

complex formation, adding to the complexities of potential regulation.

Significance Statement

Hypoxia inducible factors (HIFs) are transcription factors that regulate oxygen-dependent

cellular processes with implications in certain types of cancers. Current molecular structures of

HIFs bound to short DNA fragments provide insights into their function, but leave open

questions about how they bind longer natural DNA fragments and interact with small molecules

and protein coactivators. Integrating structural and biochemical techniques, we discovered a

novel assembly in which two HIFs bind together on a single extended DNA fragment, forming a

"dimer-of-heterodimers", which exhibits some differences in ligand and coactivator binding than

heterodimers or isolated PAS domains. Our studies highlight how functional contexts can shift

structural paradigms and provide greater insight into the mechanisms by which HIFs and similar

bHLH-PAS transcription factors operate.

Background

The hypoxia inducible factors (HIFs) are basic helix-loop-helix Per-ARNT-Sim (bHLH-

PAS) transcription factors that regulate oxygen dependent processes for higher eukaryotes under

hypoxic conditions, controlling the expression of hundreds of genes in erythropoiesis, cell

growth, metabolism, and other pathways involved in adaptation to low oxygen levels (3, 11).

HIFs are heterodimers comprised of one of three isoforms of an oxygen-sensitive α-subunit

(HIF-1, 2, 3α) and a constitutively expressed β-subunit, aryl hydrocarbon nuclear translocator

(ARNT, also known as HIF-1β) (3, 12). Under normoxia, HIF-α subunits are degraded and

inactivated through O_2-dependent enzymatic hydroxylation of specific proline and asparagine

residues in the HIF-α C-terminus (4-6). These hydroxylation markers recruit the Von Hippel-

Lindau (VHL) E3 ubiquitin ligase (proline hydroxylation) and block recruitment of p300/CREB-

binding protein (CBP) transcriptional coactivators (asparagine hydroxylation), coordinately

repressing HIF activity via independent mechanisms (3). These hydroxylation events cannot

occur under hypoxia, allowing HIF-α to accumulate, enter the nucleus, and heterodimerize with

ARNT, resulting in a HIF-α/ARNT complex (HIF complex, HIF) bound to a 5 bp hypoxia

response element (HRE) box (5'-RCGTG) in the promoter or enhancer regions of oxygen-

regulated genes to regulate transcription (**Fig. 1A**) (3, 11, 13, 14). DNA-bound HIF complexes

recruit other regulatory components such as p300 or CBP coactivators, as well as a class of

coiled-coil coactivators (CCCs), as integral parts of transcriptional regulation (**Fig. 1**) (10, 11,

15).

Functionally, HIF-1 and 2 play major roles in natural cellular hypoxic responses, with

their dysfunction implicated in a range of cancers and other diseases (3, 11, 16-18). Though their

expression profiles tend to overlap, HIF-1 is typically associated with expression of glycolytic

Figure 1. Schematic of HIF pathway regulation and components. (A) Under normoxic conditions, HIF-α is inactivated in the cytosol through parallel oxygen-dependent post-translational hydroxylations of residues in the protein C-terminus. These include a proline-directed mechanism (HIF-1α: Pro402 and 562, HIF-2α: Pro405 and 531) which leads to the recruitment of the von Hippel-Lindau E3 ubiquitin ligase, leading to poly-ubiquitination and proteosomal degradation (3, 4). A second route focuses on an asparagine residue (HIF-1α: Asn803, HIF-2α: Asn851) in the C-terminal transactivation domain (CTAD), preventing recruitment of CBP/p300 coactivators essential for HIF-driven transcriptional activation (5, 6). Under hypoxic conditions, neither of these HIF-α hydroxylations occur, allowing the fully functional protein to accumulate, translocate into to the nucleus, and heterodimerize with ARNT. The HIF-α-ARNT complex then associates with specific HRE sequences in the promoter or enhancer regions of oxygen-dependent genes and recruits coactivators, leading to transcriptional activation. (B) Schematic depicting the domain arrangements of human HIF-2α and ARNT, as well as indicating which domains interact with HRE DNA as well as CCC- and CBP/p300-type transcriptional coactivators.

enzymes ubiquitously, whereas HIF-2 is associated with expression of erythropoietin and

angiogenic genes localized to the kidney, lung, heart and small intestine (19, 20). Conversely,

HIF-3 has been implicated in fatty acid metabolism and potentially as a negative regulator of

hypoxic gene expression (21, 22). Constitutively high levels of HIFs can result in increased

levels of cellular growth, leading to tumorigenesis and specific types of cancers such as renal cell

carcinoma (HIF-2) and breast cancer (HIF-1), while the naturally hypoxic environment of tumors

can also upregulate HIF activity, exacerbating tumor progression through activation of hypoxic

pathways (11, 16-18, 23). Therefore, HIFs are an attractive target in the development of

anticancer agents with one such method of inhibition being the disruption of HIF-α-ARNT

complex formation, the proposed mechanism of action for the FDA approved HIF-2 inhibitor

and renal cell carcinoma/Von Hippel Lindau syndrome treatment, belzutifan (24-26).

Several protein/DNA crystal structures of bHLH-PAS transcription factors provide

structural insights into HIF/DNA binding interactions, most relevantly structures of murine HIF-

1 and HIF-2 bound to artificial 20 bp DNA fragments centered on canonical 5 bp HRE boxes

(5'-RCGTG) (9). Collectively, these structures show expected bHLH/DNA interactions, as well

as extensive association between the bHLH and PAS domains between both protein subunits

(**Fig. S1A,B**) (9). More broadly, this structural arrangement is highly conserved among

crystallographic and cryo-EM structures of various HIF complexes, as swapping HIF-1/2/3α

isoforms or ARNT homologs (to brain and muscle ARNT-like, BMAL1), leads to only minor

perturbations (**Fig. S1**) (9, 27, 28). Substantial changes in organization are only seen with

changes in α subunits, as seen between HIF-2α/BMAL1 and CLOCK/BMAL1 (28, 29)

complexes, which reorganize quaternary arrangements of PAS domains within the heterodimer

assembly (**Fig. S1**).

While these structural studies improved the understanding of protein/protein and

protein/DNA interactions in HIF complexes, several fundamental gaps in knowledge remain.

Foremost among these is the inability to explain long-standing data showing the importance of

DNA sequences adjacent to canonical HREs for HIF-driven transcriptional regulation. Initially

identified by deletion/mutation experiments of an enhancer region 3' to the human EPO gene,

full HIF-1 activity depends on the presence not only the HRE-box, but also an additional short

sequence (5'-CAC-3') 8 bp downstream, dubbed the HRE adjacent sequence (HAS) (14, 30).

Deletion or mutagenesis of the HAS markedly reduced HIF activity in cells (30); recent follow-

up studies showed that even minor changes to the length of the 8 bp HRE/HAS intervening

sequence similarly ablates HIF activity in cells, strongly implying cooperativity between these

regions (14, 30). While initially suggested that HAS sequences might function by recruiting a

separate transcription factor complex such as CREB-1/ATF-1, more recent studies suggest the

HAS might bind a second HIF heterodimer due to its partial complementarity to the HRE (14,

31, 32). Such higher order complexes have not been seen in existing structures of HIFs bound to

short HRE-only DNA fragments, leaving the arrangements of such higher-order HIF complexes

an open topic.

A second open area relates to the binding of small molecules within bHLH-PAS

complexes, which can naturally or artificially modulate their function. Most of such compounds

bind within the PAS domains themselves, particularly into substantially-sized internal cavities

can exist (*e.g.* 100-500 $Å^3$ in HIF-α PAS-B domains (27, 33)), providing sites for specific

binding to disrupt protein/protein interactions. This strategy has been successfully used for the

HIF-2α specific inhibitor belzutifan, now clinically used to treat various forms of clear cell renal

cell carcinoma (19, 21, 24-26, 34). Additional binding sites are on PAS domain surfaces (1),

providing alternative routes to such regulation. However, almost all binding information on these

compounds have been determined on isolated PAS domains (1, 7, 35) or DNA-free bHLH-PAS

heterodimers (9, 24, 27). As such, substantive questions remain open about the impact of HIF

complex state on small molecule binding.

Finally, information remains sparse regarding interactions between the HIF-α/ARNT

heterodimer and other proteins involved in transcriptional regulation, such as transcriptional

coactivators. While some of these, such as p300/CBP, are well-known to interact with C-terminal

disordered regions in HIF-1α and HIF-2α (36), other coactivators are thought to directly interact

with the ordered bHLH/PAS core. These include the CCCs, three of which are known

components of ARNT-containing transcriptional complexes: transforming acidic coiled-coil

containing protein 3 (TACC3), thyroid hormone receptor/retinoblastoma interacting protein 230

(TRIP230) and coiled-coil coactivator (CoCoA) (2, 15, 37-39). Prior biochemical and structural

studies from our group suggested that ARNT/CCC interactions are driven by the ARNT PAS-B

domain binding to a C-terminal conserved "TACC box" in TACC3, CoCoA, and TRIP230 (10,

15, 37, 40). However, studies on CCC/ARNT PAS-B complexes (including binding, NMR, and

crystallographic analyses (10, 15)) and crystal structures of CCC-free bHLH-PAS heterodimers

(9) have conflicting views of the accessibility of proposed binding sites, necessitating

characterization of CCC-bound bHLH/PAS complexes to resolve these issues.

With these gaps in knowledge, we assert that current structural models of HIF

heterodimers may not fully explain existing data about larger or variant HIF complexes that form

in different functional contexts with additional DNA, proteins, or small molecules bound. To

address this possibility, we explored different HIF complexes using cryo-electron microscopy

(cryo-EM) together with a variety of biochemical and biophysical approaches. Specifically, we

compared cryo-EM models of human HIF-2 bound to either a synthetic 20 bp HRE fragment or

extended DNAs from the human EPO enhancer region which naturally contain both the HRE and

HAS boxes. While we observed a very similar HIF-2/20 bp HRE cryo-EM structure (3.61 Å) as

previously seen by X-ray crystallography, the HIF-2/51 bp HRE/HAS cryo-EM structure (3.74

Å) adopted a novel larger configuration, with two individually assembled HIF-2 heterodimers

binding each other symmetrically while simultaneously engaging the HRE and HAS boxes of a

single DNA fragment. These larger "dimer of heterodimer" (DoHD) complexes are chiefly

mediated by ARNT-ARNT interactions between the adjacent HIF-2 heterodimers, implying that

they could form in other HIF complexes as well. Consistent with this, we observed longer

HRE/HAS DNAs efficiently forming similar large complexes for HIF-1 and even mixed HIF-

1/HIF-2 complexes. Additionally, we investigated the ability of both HIF-2 complexes to bind

small molecules known to bind isolated HIF-2α and ARNT PAS-B domains, finding these

ligands can still bind and disrupt the larger complexes, despite the apparent inaccessibility of

their binding sites in static structures. Finally, we propose a binding interface for CCCs on HIF-1

and HIF-2 bound to 20 bp DNA fragments through cryo-EM modeling and independently

validated by hydrogen-deuterium exchange mass spectrometry (HDX-MS). This binding, while

observed in heterodimers, is seemingly impaired in the larger DoHD complex, again suggesting

differences in dynamics among the complexes. Taken together, our findings suggest that the

molecular context of HIF complexes can substantially modulate their structures, which in turn

impacts their ability to function and associate with other functionally important components.

Results

Cryo-EM of HIF-2/HRE complexes show similar structures, but more dynamics, than

crystallographically

187   Initially, we set out to determine a cryo-EM structure of the human HIF-2 bHLH/PAS

heterodimer bound to a synthetic 20 bp HRE DNA fragment previously used for crystallographic

studies (9), mixed with C-terminal fragments of human TACC3 or CoCoA. Such a structure

would allow direct comparisons between these human complexes and prior structures of murine

proteins (**Fig. S1**) (9), while also giving insights into CCC binding modes. The resulting datasets

showed a high degree of heterogeneity, including a substantial number of particles that appeared

to be HIF-2 complexes without bound CCCs. Focusing first on these HIF-2/HRE complexes, we

used WARP (41) to pick ~1,980,000 particles and cryoSPARC (42, 43) to progressively filter

out undesired ones through iterative processes of 2D classification, *ab initio* reconstruction, and

3D refinement (**Fig. S3**). We observed a mix of 2D class averages from this analysis, including

those clearly corresponding to HIF-2/HRE complexes like the prior crystal structures (**Fig. 2A**).

Intriguingly, we also saw classes that exhibited greater conformational dynamics than previously

reported, including those that appeared to be DNA-free (with correspondingly disordered bHLH

domains) and DNA-bound complexes with a detached or blurred portion similar in size to a PAS

domain (**Fig. 2B**). Given the organization of HIF bHLH/PAS heterodimers, we interpret this as a

ARNT PAS-B domain which has separated from the rest of the HIF-2α/ARNT heterodimer.

Using ~262,000 particles (13.3% of the total particles picked) from the HIF-2/DNA class

noted above, we obtained a 3.61 Å EM map which we successfully fit with an AlphaFold-

generated model of human HIF-2α-ARNT bound to 20 bp HRE DNA (**Fig. S4A**) without

requiring any large-scale rearrangements (**Fig. 2C**), save for the termini of the HRE fragment

which were left unmodeled as a result. Similar to HIF crystal structures (9, 27), several long

protein loops were unresolved, with the most notable being the ARNT PAS-A/PAS-B linker

(ARNT residues 346-359), likely due to high flexibility in these regions.

Notably, while our cryo-EM structure superimposed well with the murine crystal

structure, as expected from the > 95% sequence identities of the HIF-2α and ARNT bHLH/PAS

regions, we observed that the structures slightly differ in the position of the ARNT PAS-B

Figure 2. Cryo-EM structural analysis of human HIF-2 bound to 20 bp HRE fragments. (A) Cryo-EM 2D classes of HIF-2 complexes, with individual PAS and bHLH domains and DNA visible. Scale bar represents 100 Å. (B) Some 2D classes of HIF-2 complexes showed either an apparent ARNT PAS-B detached from the rest of the HIF-2α-ARNT core (top row, left) or destabilized bHLH domains lacking bound DNA (top row, right). 2D projections generated through backprojection of the final volume are shown in the bottom row for comparison. (C) 3.61 Å EM map (262,466 particles, 13.3% of all collected particles) was generated from refinement of the most populated *ab initio* volume into which an AlphaFold 2 model of human HIF-2 was fit. (D) Overlay of the mouse HIF-2 crystal structure (PDB code: 4ZPK, (9)) and human HIF-2 cryo-EM structure show highly similar domain architectures. (E) Local resolutions mapped onto the HIF-2 cryo-EM density map show a region of particularly low local resolution/high flexibility in the ARNT PAS-B (Fα-helix, Gβ-strand, Hβ strand) domain and bHLH domains.

domains. This difference seems to be a ~2.5 Å domain-level shift of the cryo-EM ARNT PAS-B

domain away from the adjacent HIF-2α PAS-B domain and the rest of the heterodimer (**Fig.**

2D). Supporting this, local resolution analysis of the cryo-EM map estimated most regions had

resolutions between 3.2-4.0 Å, with the lowest protein-associated resolutions corresponding to

the ARNT PAS-B domain and in some of the extended loops, again consistent with local

dynamics in these regions (**Fig. 2E**). Together with our observation of even larger movements of

ARNT PAS-B in the 2D class averages (**Fig. 2B**), we suggest that this domain has markedly

more dynamics than previously appreciated, with potential functional links we explore below.

HIF-2α-ARNT forms a heterotetrameric structure on EPO enhancer DNAs

224   As noted above, previous studies indicated that additional DNA sequences adjacent to

HRE boxes are functionally important for full activation of certain oxygen-regulated gene

promoters (14, 30). To explore the structural basis of this observation, we used cryo-EM to

examine HIF-2 complexes bound to longer ~50 bp human EPO enhancer DNA sequences,

including both a HRE and HAS box previously demonstrated as necessary for full activation.

Using DNAs including both HRE and HAS boxes with additional flanking sequences on either

the 5' (dubbed BS1 HRE/HAS, 51 bp) or 3' (dubbed BS2 HRE/HAS, 52 bp) ends, we collected

cryo-EM datasets of human HIF-2 complexes bound to these longer DNAs, mixed with C-

terminal TACC3 fragments. As with our studies of the shorter 20 bp HRE complexes, most

particles appeared absent of TACC3, so we focus here on complexes of HIF-2 bound to the 51

bp BS1 HRE/HAS fragments.

235   Early in our analysis, we observed 2D class averages that were markedly different than

seen for HIF-2 bound to 20 bp HRE (**Fig. 2A,3A**). Instead, HIF-2 bound to 51 bp HRE/HAS

showed classes with much larger protein density arranged more symmetrically about the DNA,

prompting further structural examination of these complexes. Final refinement of data from

164,000 particles (25.0% of all particles picked) (**Fig. S5**), led to the determination of a 3.74 Å

structure which neatly fit two individually assembled HIF-2α-ARNT heterodimers as well as the

51 bp BS1 HRE/HAS fragment into a "dimer-of-heterodimers" (DoHD) conformation within our

density map without needing any large scale domain rearrangement (**Fig. 3B**). We underscore

DoHD complexes were observed by size exclusion chromatography before freezing (**Fig. S6**)

and we did not see any comparable higher-order assemblies in our HIF-2/20 bp HRE cryo-EM

results, supporting that formation of this larger complex can occur *in vitro*, is dependent on the

bound DNA component, and is not an artifact of cryo-EM processing. Conversely, we did not

see any smaller assemblies of single heterodimers bound to 51 bp HRE/HAS in cryo-EM,

implying the DoHD complex is stable enough to be isolated *in vitro* and remain intact during

cryo-EM (**Fig. S6**).

250   This intriguing higher-order complex orients the two HIF-2 heterodimers around a

pseudo two-fold rotation axis, with one heterodimer rotated ~180° from the other. The first of

these heterodimers (1° heterodimer) bound to the HRE box through the HIF-2α and ARNT

bHLH domains in a nearly identical fashion to the minimal complexes with 20 bp HRE-only

DNA fragments. The 2° heterodimer bound the HAS box through its respective bHLH domains

(**Fig. 3B,S6**), positioned in opposition to the 1° heterodimer, with the spacing between the HRE

and HAS boxes facilitating direct interactions between the heterodimers. Protein/protein

interactions between the two heterodimers were chiefly mediated through the ARNT PAS-B

domains, utilizing residues in the Fα-helix and Gβ-strand with potential interactions between

Leu407, Gln414, and Met426 of both ARNT PAS-B domains that could be unambiguously

Figure 3. Structural and biochemical analyses of HIF dimer-of-heterodimer complexes. (A) 2D cryo-EM class averages from analyses of data collected on HIF-2α-ARNT heterodimers bound to 51 bp BS1 HRE/HAS DNA fragments, showing markedly different (and often, more symmetrical) arrangements than comparable 2D class averages of HIF-2 bound to 20 bp HRE DNAs. Scale bar represents 100 Å. (B) A 3.74 Å EM map (164,352 particles, 25.0% of all collected particles) was generated from refinement of the most populated *ab initio* volume into which two individually assembled human HIF-2 heterodimers were fit, along with the corresponding 51 bp BS1 HRE/HAS DNA fragment. (C) Local resolution data mapped onto the HIF-2 dimer-of-heterodimers cryo-EM map showed that the ARNT PAS-B Fα-helix and Gβ-strand were higher relative resolution than in corresponding sections of the 20 bp HRE-bound HIF-2 EM map, suggestive of ordering between the heterodimer and dimer-of-heterodimer complexes. (D) The 3.90 Å cryo-EM map (71,802 particles, 27.3% of all collected particles) of HIF-1 bound to a 52 bp BS2 HRE/HAS DNA fragment showed a highly similar domain arrangement to HIF-2, though the ARNT PAS-A domains were less resolved. (E) Highlighted residues were suspected to form interactions between the opposing ARNT PAS-B domains in the HIF-2 supercomplex (basic residues in blue, acidic residues in red, uncharged residues in magenta) (F) SEC-MALS data showed expected differences in the molecular weights of HIF-2 on 20 bp HRE DNA fragments (96.7±6.3 kDa) and 51 bp BS1 HRE/HAS DNA fragments (214.0±14.1 kDa), but revealed an intermediate value (144.0±6.9 kDa) for a HIF-2 variant with mutations in four key ARNT PAS-B Fα-helix and Gβ-strand residues which are at the ARNT/ARNT interface essential to the dimer-of-heterodimers complex. (G) Luciferase reporter assays containing the wildtype (WT) or mutant versions of the human EPO enhancer with changes in the upstream GC-rich region (GC-rich-mut), HRE (HRE-mut), or HAS (HAS-mut) site were performed in HEK293T cells in conjunction with ectopic expression of oxygen-independent (PPN) HIF-1α or HIF-2α. Empty pIRES vector expression was measured to ensure empty vector would not contribute to reporter response. Mutations to either the HRE or HAS site substantially blunted reporter induction by either PPN HIF-1α and PPN HIF-2α.

 placed at the protein/protein interface despite the limited resolution of the cryo-EM map.

Additional interactions may come from the HIF-2α PAS-A EF loops as well (**Fig. S6**). Cryo-EM

data for the central 27 basepairs of the 51 bp fragment were clearly resolved to allow us to

confidently place both the HRE and HAS boxes into the cryo-EM map, with slight upward

curvature 3' to the HAS-box before losing resolution at the 3' terminus, possibly to support

binding of the 2° heterodimer to the DNA.

We also performed local resolution analysis of this dimer-of-heterodimers cryo-EM map,

which spanned 3.2-4.2 Å resolution (**Fig. 3C**). While similar overall to our analysis from the 20

bp HRE-bound HIF-2, we observed two substantial changes. First, we saw higher resolution

density in the dimer-of-heterodimers map corresponding to the ARNT PAS-B domains

compared to the 20 bp HRE-bound HIF-2 map, specifically around the β-sheet surface and Fα-

helix located at the dimer/dimer interface, implying a domain-level stabilization. In contrast, the

dimer-of-heterodimers map appears to show relatively low resolution in the ARNT PAS-A

domains of both heterodimers, likely due to being distal to the protein/protein and protein/DNA

interactions which stabilize the complex.

Implications and validation of dimer-of-heterodimer assembly on other HIFs

The central role of ARNT in mediating these dimer-of-heterodimer complexes has

several implications. First, analogous HIF-1 DoHD complexes should be able to form on longer

DNAs, given the shared role of ARNT in both HIF-1 and HIF-2. Second, point mutations at the

ARNT PAS-B domain, which provides the majority of protein/protein interface in the cryo-EM

structure, should destabilize dimer-of-heterodimer formation. Finally, "mixed" dimer-of-

heterodimers should be evident *in vitro*, with HIF-1 and HIF-2 heterodimers co-complexing to

the same DNA fragment. We address each of these implications in turn below.

284   Starting with the ability of HIF-1 to form dimer-of-heterodimer structures, we collected a

285 cryo-EM dataset of human HIF-1 bHLH/PAS fragments bound to 52 bp BS2

286 HRE/HASfragments, as initial characterization of HIF-1/DNA complexes appeared more

287 homogenous with BS2 (**Fig. S7**). 2D classes from this dataset were very similar to those from

288 our prior HIF-2/HRE/HAS studies above, suggesting similar structure formation (**Fig. S7B**).

289 Further data refinement from 71,800 particles led to a 3.90 Å cryo-EM map, into which we could

290 fit two individually assembled HIF-1 AlphaFold-generated models using the inverted repeat-type

291 arrangement as in the HIF-2 dimer-of-heterodimers (**Fig. 3B,D; Fig. S6,S7**). As before, the

292 ARNT PAS-B Fα-helix and Gβ-strand elements appear to provide the bulk of protein/protein

293 interaction surface within the HIF-1 complex, consistent with the ARNT PAS-B domain playing

294 a role in dimerizing HIF-1 as well as HIF-2. The HIF-1α PAS-A EF loops again appear to

295 potentially play a further role in stabilizing the DoHD complex, notably with apparent EM

296 density between nearby cysteines (Cys118) in these loops (**Fig. S7H**). Interestingly, the ARNT

297 PAS-A domains of this complex appeared to be of a much worse resolution than in the HIF-2

298 DoHD complex, leading to fewer residues of this domain being confidently fit into the EM map

299 (**Fig. S7F**).

301 **The ARNT PAS-B domain serves as a major interface for HIF-2α-ARNT**

302 **homodimerization**

303   Our cryo-EM structures and local resolution analyses of the HIF dimer-of-heterodimers

304 support the ARNT PAS-B domain as a dimerization interface between the two HIF-α-ARNT

305 heterodimers (**Fig. 3B-D**). Independently, the ARNT PAS-B domain has been suggested to

306 participate in higher order HIF assemblies via analyses of intermolecular contacts in HIF crystal

structures (14), albeit in a different arrangement not supported by our cryo-EM structure. To test

the role of ARNT PAS-B as a potential dimerization interface, we mutated specific residues at

the ARNT PAS-B/ARNT PAS-B interface of our dimer-of-heterodimer structure. These

residues, located in the ARNT PAS-B Fα-helix and neighboring β-strands, are proximal and

show improved local resolution in the dimer-of-heterodimers compared to the simpler

heterodimer structure, implying these regions are stabilized through direct interactions (**Fig. 3C**).

At this interface, we observed several sets of salt-bridges (Glu362 to Gln414/Lys417;

Asp410/Ser 411 to Arg440) and nonpolar clusters (Leu423 to Leu423; Met426 to

Leu407/Val425) (**Fig. 3E**) which are evolutionarily conserved among vertebrate ARNTs (**Fig.**

S8). To disrupt these interactions, we mutated Glu362 and Arg440 to alanine and Leu423 and

Met426 to lysine, developing a quad mutant ARNT (ARNT E362A/L423K/M426K/R440A)

which we predicted to have lower dimer-of-heterodimer stability than wildtype. Using size

exclusion chromatography coupled with multi-angle light scattering (SEC-MALS), we examined

the average molecular weights of the wildtype and mutant complexes with the 51 bp BS1

HRE/HAS DNA fragment. For wildtype HIF-2, this complex had an average molecular weight

of 214.0±14.1 kDa, similar to the expected 202.7 kDa for two HIF-2 heterodimers bound to a 51

bp DNA duplex and effectively double the 96.7±6.3 kDa observed for a HIF-2 heterodimer

bound to a 20 bp HRE fragment (**Fig. 3F**). In contrast, HIF-2 containing the quad

E362A/L423K/M426K/R440A mutant ARNT bound to 51 bp BS1 DNA showed a later-eluting

SEC peak with an average 144.0±6.9 kDa molecular weight (**Fig. 3F**). Though higher than the

expected 117.0 kDa for a single HIF-2 heterodimer stably bound to a 51 bp DNA duplex, our

SEC and MALS data strongly support an equilibrium shift towards a single heterodimer,

implying disruption of DoHD formation. Combining these results to the cryo-EM structure and

local resolution analysis suggest the importance of the ARNT PAS-B Fα-helix and adjacent β-

strands as an interface in the formation of dimer-of-heterodimers.

ARNT-mediated formation of mixed HIF-1/HIF-2 dimers-of-heterodimers

As the last of the three implications of our dimer-of-heterodimer structures, we asked if

the commonality of ARNT in mediating comparable HIF-1 and HIF-2 dimers-of-heterodimers

could facilitate HIF-1 and HIF-2 complexing together to form a "heterodimer-of-heterodimers"

complex on extended DNAs. To test this, we created a fusion protein of HIF-1α tagged on the N-

terminus with *E. coli* maltose binding protein (MBP) to increase its molecular weight by about

45 kDa, allowing us to distinguish HIF-1 and HIF-2 complexes by SDS-PAGE and size

exclusion chromatography.

After copurifying this MBP-tagged HIF-1 with untagged HIF-2, we bound them to 51 bp

BS1 HRE/HAS DNA fragments, and ran this mixed sample over SEC. We observed two peaks

in the resulting chromatogram (**Fig. S9A**), with SDS-PAGE analysis showing all three subunits

(peak 1, MBP-HIF-1α, HIF-2α, ARNT) in the first, earlier-eluting peak and only HIF-2α and

ARNT in the later peak (peak 2) (**Fig. S9D**). The elution volume of the later peak (peak 2) was

similar to that of independently isolated HIF-2 DoHD complex while the earlier eluting peak

(peak 1) corresponded to an elution volume between those of independently isolated MBP-HIF-1

DoHD and HIF-2 DoHD complexes (**Fig.S9A**). This supports the formation of a tandem HIF-

1/HIF-2 *heterodimer*-of-heterodimers on an extended DNA fragment. Intriguingly, we did not

see a peak corresponding to MBP-HIF-1/MBP-HIF-1 heterodimers in the mixed sample, which

we would expect to form based on our cryo-EM data and have confirmed through SEC of MBP-

HIF-1 heterodimers bound to 51 bp HRE/HAS DNA fragments (**Fig. S9A**).

To explore the DNA sequence requirements for forming these higher order structures, we

repeated these experiments using 51 bp BS1 DNA fragments with mutations into either the HRE

or HAS-box to reduce protein binding to those sites (**Fig. S9B**). As expected, mutation of the

HRE-box prevented almost any heterodimer formation, suggesting the HAS box alone is

insufficient to stably recruit either single heterodimers or dimer-of-heterodimers; protein that is

not bound to DNA cannot stably form heterodimers, and therefore elutes in the void (**Fig.**

S9C,S9E). In contrast, mutation of the HAS-box increased the presence of a later elution peak

we attribute to a single heterodimer on DNA, while DoHD formation was reduced but still

evident (**Fig. S9C,S9E**).

Coupled with the inability to observe stable dimers-of-heterodimers on 20 bp HRE-only

DNAs (by either cryo-EM or X-ray crystallography), we interpret these data as giving us insights

into the coupled protein/protein and protein/DNA interactions needed to assemble higher-order

HIF complexes: HIF/HRE interactions are key for 1° heterodimer recruitment, with additional

DNA interactions – ideally from an appropriately-spaced HAS box – needed to stabilize the

recruitment of a 2° heterodimer. This is consistent with prior experiments which scanned the

transcriptional potency of HIF-driven promoters with HRE/HAS spacings of 2 to 12 bp (in 2 bp

steps) showing maximal activity specifically at 8 bp(14).

Formation of HIF dimer-of-heterodimers is important to cellular function

Building on our observation of HIF dimers-of-heterodimers *in vitro*, we conducted

further studies to determine if these complexes would assemble in cells, and if they are

functionally relevant. We generated luciferase reporter plasmids driven by EPO enhancer region

sequences or variants containing mutations in the HRE, HAS or 5' GC-rich regions, and

cotransfected these into HEK293T cells together with expression plasmids for oxygen-invariant

"PPN" variants of human HIF-1α or HIF-2α. Examining luciferase expression from these cells,

we observed robust expression of the reporter genes driven by either the WT promoter or one

containing point mutations in the GC-rich region. In contrast, expression from plasmids with

HRE mutations was ablated by approximately 90%, as expected from the key role that HREs

play in recruiting HIFs. Notably, expression from HAS-mutated promoters was also substantially

decreased compared to WT controls, with approximately 70-80% decreased activity (**Fig. 3G**).

As neither mutation completely ablated HIF-dependent transcription of the reporter, we suggest

that either the HRE or HAS sequence on its own may still recruit a single heterodimeric

complex. Indeed, many HIF-regulated genes do not contain an HAS-box, and therefore must

function by recruiting HIF only to the HRE-box (14, 44).

Structural differences between HIF-2α-ARNT complexes impacts ligand binding

Our work demonstrates that differences in protein and DNA fragments of HIF complexes

can markedly impact the quaternary structures of these components. To explore the functional

impact of these changes on the binding of small molecules to various HIF complexes, we

measured the efficacies of several HIF-2α or ARNT PAS-B binding compounds to disrupt

various HIF-2 complexes. Such compounds, which have been sought as leads for disruptors of

protein/protein interactions in HIFs, have typically been identified using biophysical or

biochemical approaches focused on ligand binding to isolated PAS-B domains (2, 7, 35). These

compounds range in affinity from nanomolar (e.g. compound **2** binding to HIF-2α PAS-B, K_d 81

nM (7)) to high micromolar (e.g. ARNT PAS-B binding KG-548 and KG-279 (1)) (**Fig. 4A**), but

comparable measurements have typically not been available on larger complexes, some of which

occlude obvious access to binding sites on or inside of the PAS domain targets. To address this

shortcoming, we used amplified luminescent proximity homogenous assay screen (AlphaScreen,

Revvity) to measure the ability of several HIF-2α or ARNT targeting ligands to disrupt the HIF-2

heterodimer and the dimer-of-heterodimers (FLAG-tagged ARNT subunits) bound to

biotinylated 20 bp HRE or 51 bp BS1 HRE/HAS DNA fragments. AlphaScreen detects the

proximity between two biomolecules separately tagged for affinity to either a donor

(streptavidin-coated bead for biotinylated HRE or HRE/HAS) or an acceptor bead (anti-FLAG

coated bead for FLAG-tagged ARNT) by emitting luminescence in response to diffusion of

singlet oxygen between beads. As such, disruption of the HIF-2 complexes would result in

displacement of ARNT from the DNA, leading to a detectable decrease in luminescence that can

be used to determine the potency of small molecules to disrupt either complex (45, 46).

The first of these compounds, KG-548, disrupted the single HIF-2 heterodimer with a lower

potency than the affinity to the isolated ARNT PAS-B(K_d 348 µM ARNT PAS-B; ~ 1 mM IC_{50}

HIF-2 heterodimer) despite its binding interface on the ARNT PAS-B surface being blocked by

the HIF-2α PAS-B domain in the heterodimeric complex (**Fig. 4B**). This observation suggests

some flexibility in the ARNT PAS-B conformation with respect to the rest of the heterodimer, as

supported by some cryo-EM 2D classes of the HIF-2 complex with the ARNT PAS-B domain

apparently detached from the other core domains (**Fig. 2B**), as well as the lower local resolution

of the ARNT PAS-B domain within the 3D heterodimeric complex (**Fig. 2E**). Interestingly, KG-

548 was also capable of disrupting the HIF-2 DoHD complex with a much higher potency (IC_{50}

= 74 µM), despite the apparent stabilization of the ARNT PAS-B domains in this complex (**Fig.**

4C). The greatly increased potency of KG-548 against the DoHD complex may suggest new

disrupting interfaces for KG-548, possibly between the heterodimers, or a modification of the

Figure 4. Contextual differences can impact HIF-2 small molecule binding and disruption potency. (A) Known small molecule binders of isolated ARNT (KG-548 (1, 2), KG-279 (1, 2)) and HIF-2α PAS-B (compound 2 (7), PT2385 (8)) domains are shown along with their literature affinities for the specific PAS-B targets, obtained by ITC and NMR. (B) Small molecule disruption of the HIF-2 heterodimer is seen across PAS-B binding compounds, as assayed by AlphaScreen probing interactions between a FLAG-tagged ARNT subunit of HIF-2 and a biotinylated 20 bp HRE fragment. (C) Small molecule disruption of the HIF-2 DoHD complex is also seen across the tested ligands, as detected with AlphaScreen probing interactions between a FLAG-tagged ARNT subunit of HIF-2 and a biotinylated 51 bp HRE/HAS fragment. In both (B) and (C), we observe some differences in disruption potency in the larger contexts compared to the isolated PAS-B domain binding affinity, particularly for the ARNT PAS-B binders.

existing inhibitory interface that increases potency of the inhibitor. We observed KG-279

disrupting both HIF-2 bound to 20 bp HRE DNA fragments (**Fig. 4B**) and HIF-2 DoHD on 51 bp

BS1 HRE/HAS DNA fragments (**Fig. 4C**). Though binding to a single heterodimer would be

expected due to observed flexibility of the ARNT PAS-B domain, binding to the larger dimer-of-

heterodimers suggests the ARNT PAS-B internal cavity is accessible regardless of quaternary

structure. The very substantial increase in potency to the HIF-2 heterodimer ($IC_{50} = 25$ μM) in

comparison to the affinity to the isolated ARNT PAS-B ($K_d = 1.2$ mM) suggests novel binding

sites for KG-279 on the larger complexes. Additionally, compound 2 and PT2385 were able to

disrupt both the HIF-2 heterodimer and DoHD complexes, suggesting the HIF-2α PAS-B cavity

remains equally accessible despite the changes in quaternary structure (**Fig. 4C**).

TACC3 interacts with loops of both HIF-α and ARNT subunits

Though we can assemble dimer-of-heterodimers complexes on HRE/HAS DNA

fragments, we understand that only a subset of HIF-driven promoters contain both binding motifs

(**Table S1**). Therefore, gene promoters containing only HRE binding motifs, we suspect that

HIFs may form varying higher-order complexes with other parts of the transcriptional regulatory

machinery. One well studied candidate for HIF binding partners are the CCCs. While CCCs are

clearly involved in transcriptional initiation from HIF-driven promoters, and that this is mediated

by direct interaction between HIF and CCC proteins, the location of such interaction remains

unclear. NMR and other biophysical data from our lab suggested that CCCs bind to isolated

ARNT PAS-B domains at a site (2, 15, 39) inconsistent with subsequently-determined crystal (9)

and cryo-EM (*vide infra*) structures of larger HIF-2/HRE complexes.

To begin addressing this incompatibility, we sought to characterize larger HIF-2

complexes bound to CCCs. We purified C-terminal fragments of two CCCs, CoCoA and TACC3

(10, 15), confirmed that these were dimeric and coiled coil (**Fig. S10A,B**), and used MST to

measure the binding of these fragments to isolated ARNT PAS-B domains. We observed

micromolar affinities for these interaction, K_d 80 µM for CoCoA and 3 µM for TACC3 (**Fig.**

5A), consistent with prior results (10, 15). Repeating these measurements with the same CCC

fragments binding to HRE-bound HIF-2 complexes, however, showed K_d ~10 µM interactions of

both coactivators with HIF-2 (**Fig. 5B**). These shifts in affinity between ARNT PAS-B and the

intact bHLH/PAS heterodimers show the involvement of additional regions outside of ARNT

PAS-B in the HIF-2α/ARNT heterodimer, regardless of whether this is an improvement

(CoCoA) or worsening (TACC3).

Figure 5. Coiled-coil coactivator binding to HIF heterodimers. (A, B) MST measurements of C-terminal fragments of CoCoA(407-535) and TACC3(758-838) binding to the (A) isolated ARNT PAS-B domains and (B) HIF-2α-ARNT heterodimers (HIF-2) bound to 20 bp HRE fragments. The two highest points of the hCoCoA to HIF-2 binding curve showed evidence of aggregation and were removed. TACC3 binding to HIF-2 showed a decrease at high concentration, suggesting a second binding mode or aggregation. (C) A truncated TACC3(788-838, "shTACC3") was designed based on our TACC3(758-838) crystal structure showing highly elevated B-factors and poor electron density in residues 758-788. This CCC fragment bound a 20 bp HRE-bound HIF-2 with a minimal affinity drop from the longer TACC3 (K$_d$ 26 μM by MST), and without the secondary binding mode seen for longer fragments to HIF-2. (D) 2D classes of shTACC3 bound to 20 bp HRE-bound HIF-2 showed visible core domains with excess blurred density extending out of the central region. (E) *Ab initio* reconstruction of 20 bp HRE-bound HIF-2 bound to shTACC3. HIF-2 fit into the volume without major rearrangement, leaving excess density along the ARNT PAS-B/HIF-2α PAS-A interface for most of shTACC3 to be fit. Fitting was solely based on excess density in cryo-EM map of HIF/TACC3 complexes, with the sole assumption that the TACC3 C-terminus is closest to ARNT PAS-B per prior deletion experiments (10). (F) Schematic of interactions between TACC3(788-838) and HIF-2 shows three of the four PAS domains interacting with the TACC3 coiled coil. Residues indicated on TACC3 originate from either chain. (G) Cryo-EM of HIF-1 bound to the longer TACC3(758-858) fragment showed similar excess density as HIF-2/TACC3(788-838), consistent with TACC3 binding similarly to HIF-1 and HIF-2. TACC3 residues 793-838 were fit into the excess density for both HIF-1 and HIF-2. (H) HDX-MS deuterium uptake plots showed several HIF-2α and ARNT loops with increased protection from solvent in the presence of TACC3. (I) Mapping peptides with TACC3-induced HDX changes onto shTACC3-bound HIF-2 model (increased protection in green) showed these protected loops along the proposed TACC3 binding interface from cryo-EM.

To obtain more structural insight into CCC/HIF-2 interactions, we added a TACC3

fragment (residues 758-838) to cryo-EM samples of HIF-2 bound to 20 bp HRE fragments. 2D

classes of this complex failed to resolve the HIF-2 protein domains, instead showing blurred

circular densities with "stick-like" objects protruding from them, which are of the right

dimensions to be TACC3 (**Fig. S10C**). Notably, of these classes showed a bend or break in the

stick-like density we attribute to TACC3, suggesting flexibility or dynamics within the

coactivator. This hypothesis was supported by a crystal structure of TACC3(758-838) (**Fig. 5C,**

S10D), which had low quality electron density and high B-factors for most residues N-terminal

of Lys788 (**Fig. 5C, Fig. S10D**), consistent with the break point seen in the cryo-EM classes.

Given this, we made a shorter TACC3(788-838) fragment ("shTACC3") which we utilized

forour subsequent biochemical and structural analyses. MST binding analysis of shTACC3

fragment titrated into HIF-2/HRE complex showed a 26 µM affinity (**Fig. 5C**), slightly

weakened from the longer 758-838 construct.

Having confirmed *in vitro* binding on shTACC3 to HIF-2/HRE complexes, we collected

a cryo-EM dataset to obtain structural information on this activator/coactivator complex. Using

3,320,000 particles from this dataset, we proceeded with 2D classification and observed classes

with visible core domains of HIF-2 as well as blurred excess density which we attributed to

shTACC3 (**Fig. 5D**). *Ab initio* reconstruction from 222,000 particles led to a low resolution cryo-

EM density map (**Fig. S11**) in which we were able to loosely fit our cryo-EM structure of HIF-2

bound to 20 bp HRE (**Fig. 5E**). This fitting left additional unoccupied density alongside the

ARNT PAS-B and HIF-2α PAS-A domains (**Fig. 5E**). Into this region, we fit an AlphaFold

structure of the C-terminal 46 residues of shTACC3 as a dimeric coiled coil (**Fig. S5C**) with the

C-termini facing the ARNT PAS-B domain, as suggested by our prior work (10, 15). This

density became more diffuse towards the shTACC3 N-termini, consistent with the apparent

flexibility seen in 2D class averages of longer TACC3/HIF-2 complexes and our TACC3 crystal

structure. This structural model places shTACC3 adjacent to the HIF-2 PAS domains, suggesting

shTACC3 interacts via three sets of interactions with HIF-2 through unresolved loops in the

ARNT PAS-B and HIF-2α PAS-A domains (**Fig. 5F**) and explaining the differences in affinity

we observed for TACC3 binding to ARNT PAS-B and larger HIF-2 complexes. Parallel cryo-

EM studies of complexes of the longer TACC3 (758-838) bound to HIF-1/HRE showed TACC3-

associated density in a comparable spot relative to the HIF heterodimer, suggesting comparable

binding interfaces between the two isoforms (**Fig. 5G**). We underscore that while the low

resolution of this map implies some inherent motion or flexibility in the HIF and TACC3

orientations, this model explains independently-acquired data from HIF-1/TACC3 and HIF-

2/TACC3 samples as well as MST data suggesting interactions of TACC3 with ARNT PAS-B

and additional parts of the HIF heterodimers.

To further bolster our HIF-2/TACC3 binding model, we acquired HDX-MS data on HIF-

2/HRE complexes in the absence and presence of shTACC3. These experiments revealed several

HIF-2α and ARNT peptides with notably less deuteration in the presence of shTACC3, most of

which were along the TACC3-binding interfaces seen in cryo-EM. Integrating the cryo-EM and

HDX-MS results, we found four loops in the proposed CCC binding interface supported by both

methods (**Fig. 5H,I; Table S4**): ARNT PAS-A/B linker (357-366), HIF-2α PAS-A Fα/Gβ loop

(157-163), HIF-2α PAS-A Hβ/Iβ loop (201-207) and HIF-2α PAS-B C-terminal loop (346-355).

We also observed several HIF-2α peptides with TACC3-induced protection from HDX exchange

at sites remote from the proposed binding site, suggesting long-range stabilization of the HIF-

2α/ARNT heterodimer upon CCC binding (**Fig. 5I; Table S4**).

As previously mentioned, we did not see TACC3 binding to the HIF-2 dimer-of-

heterodimers via cryo-EM, despite the presence of the CCCs within samples used in these

analyses. We performed microscale thermophoresis to further probe interactions between

TACC3 and the HIF-2 dimer-of-heterodimers complex, but were again unable to see binding,

prompting us to believe TACC3 cannot bind the larger complexes. Notably, the HIF-2α and

ARNT loops we saw as critical for binding appear to be unobstructed in the HIF-2 dimer-of-

heterodimers complex, leading us to suspect that differences in dynamics across the complex

(such as those suggested by differences in cryo-EM local resolution) may contribute to this

finding.

Discussion

HIFs have been implicated in specific types of cancer and the progression of

tumorigenesis, making them a target for anticancer therapeutics (16, 17, 25, 26). Development of

such drugs, including the HIF-2α inhibitor belzutifan, required a mix of approaches applied to

reductionist models of HIF-α and ARNT components typically restricted to the isolated PAS

domains, though recently advanced to DNA-bound bHLH/PAS heterodimers. While these

approaches were successful at informing both basic science and applications of HIF biology,

they left unexplained prior findings in three areas – larger DNAs, small molecule binding, and

CCC binding – which we advance here.

The formation of higher-ordered HIF complexes was first suggested by Semenza and

Wang alongside their discovery of HIF-driven gene expression, and has been recently bolstered

by cellular and biochemical work from Rosell-Garcia *et al.* demonstrating that HRE or HAS

mutations within HIF-driven promoters drastically reduce HIF activity in cells, similar to our

results in **Fig. 3G** (14, 30, 47). Using a split-enzyme reconstitution assay, these authors also

nicely demonstrate the likely formation of DoHD complexes in cells. Taken together, HIF-driven

transcription clearly occurs from HRE-only or HRE/HAS-containing promoter/enhancer

sequences, with selectivity likely stemming from differences in HIF concentrations. Rosell-

Garcia *et al.* proposed that larger HIF complexes would form through interactions between

ARNT PAS-B and bHLH domains of interacting heterodimers, as suggested through HIF crystal

contacts, though such an arrangement places the two sets of bHLH DNA binding domains

relatively far from each other, requiring substantial distortion to bind a single DNA containing

HRE and HAS boxes (14).

In contrast, our DoHD complex places two HIF heterodimers in a symmetric arrangement

via the ARNT PAS-B β-sheet surfaces, with two sets of DNA binding domains ideally placed to

contact HRE and HAS boxes with the optimal 8 bp spacer, and not ones with shorter or longer

separations as shown by Rosell-Garcia *et al.* (14). Additionally, the sequences of the HRE-

(RCGTG) and HAS-boxes (CAC) being imperfect inverse repeats likely explains how HIFs can

recognize this second binding site and bind with a 180° rotation relative to the 1° heterodimer.

Notably, little rearrangement of the HIF heterodimer is required to assemble these larger dimer-

of-heterodimer assemblies, in stark contrast to CLOCK/BMAL1 in which heterodimers undergo

a dramatic 90° intramolecular pivot as they go from isolated heterodimers to a nucleosome-

bound complex (**Fig. S12**). Our model also explains mutation data from Rosell-Garcia showing

that intact ARNT PAS-B domains are required for higher-order HIF complexes to form *in vitro*

or in cells. Further, the high conservation of the ARNT PAS-B domain across vertebrates,

including residues involved in DoHD interactions, suggests that such complexes are likely to be

found in a wide variety of HIF species variants (**Fig. 8C**). As such, we suggest that our dimer-of-

heterodimers model best explains a variety of biochemical and molecular data on interactions

and functional importance of these higher-order HIF structures.

Categorization of different HIF-associated promoters containing HRE-only or HRE/HAS

motifs reveals that the latter group are largely involved in cell proliferation, metabolism and

angiogenesis, while HRE-only containing genes share these functions along with others

including transcription regulation, pH regulation, signaling and drug resistance (**Table S1**) (3,

44). HIFs have about a 2-3-fold higher affinity for the HRE-box alone in comparison to the

HRE/HAS-boxes together (14), suggesting a dose dependence in which the formation of higher-

ordered complexes, and therefore the expression of HRE-only or HRE/HAS-containing genes,

depends upon HIF concentration (**Fig. 6**). This is supported by data from transfection-based

control of HIF levels for cells grown in culture (14), but has yet to be demonstrated *in vivo* or

with natural or artificial regulators of HIF complexes.

The dimer-of-heterodimers concept also strongly suggests the potential for mixed HIF-

1/HIF-2 complexes controlling transcription alongside the more canonical HIF-1/HIF-1 or HIF-

2/HIF-2 complexes. Since we have determined both HIF-1 and HIF-2 can co-complex to form

mixed dimer-of-heterodimer complexes together on the same DNA fragment, the two isoforms

likely engage in crosstalk to regulate overlapping gene expression. Mixed dimer-of-heterodimer

formation may also serve as a mechanism of rescue in cases where either HIF isoform is

depleted, either through natural or artificial means (*e.g.* belzutifan inhibition of HIF-2). Rescue

of either HIF isoform has been seen in cases where HIF-1 is inactivated, HIF-2 activity will

increase to compensate (48, 49); we propose that part of this mechanism may be in the

substitution of heterodimers in dimer-of-heterodimer complexes.

This observation raises the potential of other bHLH/PAS heterodimers to join in higher

order complexes. Recent cryo-EM studies of a HIF-2α/BMAL1 complex show a similar

heterodimeric structure as HIF-2α/ARNT, exposing a similar surface on the BMAL1 PAS-B as

used in ARNT PAS-B in the formation of DoHD complexes (50) (**Fig. 6**). Sequence alignment

of the human ARNT, BMAL1 and ARNT2 PAS-B domains shows that three residues (E362,

Q414, R440) integral to DoHD formation are conserved (**Fig. 3E,F;Fig. S8A,B**). With this, we

suggest that dimer-of-heterodimer formation may be plausible for other bHLH-PAS transcription

factors provided they adopt a similar quaternary structure to HIF. For example, CLOCK-BMAL1

heterodimers do not appear to be compatible with joining HIF/ARNT heterodimers based on

existing structures, although some evidence of CLOCK/BMAL1 higher order complexes have

been seen in structural studies of nucleosome-associated structures (28, 51, 52).

Switching from the self-assembly of HIF-2α and ARNT to their ability to bind small

molecule ligands, our data clearly indicates that small molecule binding and their abilities to

disrupt dimerization is conserved across complex assemblies. We have previously found a slate

of small molecules which can bind the isolated HIF-2α or ARNT PAS-B domains (e.g.

compound 2 (7), PT2385 (8), KG-548 (1, 2), KG-279 (1, 2)) as assessed by NMR, ITC, or MST.

Notably, all small molecule compounds we tested disrupted DNA-bound HIF-2α-ARNT

heterodimers. Our examinations of KG-548, a small molecule that binds the beta-sheet surface of

ARNT PAS-B, shows that it clearly binds and disrupts single HIF-2α-ARNT heterodimers

bound to a short HRE-only DNA, despite the fact that its binding surface on ARNT PAS-B is

occluded in both our cryo-EM and prior crystallographic structures of these complexes (1, 9).

We view this result as strongly suggesting the ARNT PAS-B domain having more flexibility

than evident in these static structures, as supported by the relatively low local resolution of this

Figure 6. Proposed promoter-dependent formation of higher-order HIF and their downstream effects. (A) Under hypoxic conditions, we propose HIFs form different complexes based on the sequence of the promoter/enhancer region of the gene of interest. For HRE-only containing genes, we propose HIFs form a single heterodimer on the HRE-box, recruiting other components such as CCCs. For genes containing an HRE-box followed by an 8 bp spacer and an HAS-box, we propose that HIFs instead form a dimer-of-heterodimers. We suggest the specification of which genes are activated by HIF are dose dependent, in that lower cellular levels of HIF will activate HRE-only genes while higher cellular levels of HIF will activate HRE/HAS containing genes. Additionally, HIF-α can form noncanonical complexes with alternative class II bHLH-PAS proteins, including BMAL1, as well as mixed HIF-1/HIF-2 dimer-of-heterodimer complexes, each under varying conditions. We suggest that the similar quaternary structure of the HIF-2α-BMAL1 complex lends itself to the potential formation of dimer-of-heterodimers structures, though this remains to be experimentally validated.

domain in the single heterodimer/DNA complex and our observation of 2D classes where the

ARNT PAS-B seems to have detached from the result of the heterodimer. More broadly, studies

of other bHLH/PAS transcription factors show that the ARNT PAS-B domain (or its analog in

BMAL1) adopt a very diverse set of conformations with respect to different Class I

heterodimerization partners (HIF-α, AHR, NPAS, CLOCK), supporting this hypothesis (9, 29,

53, 54). Despite the higher complexity and perceived stability of the larger HIF-2 DoHD

complexes, we were still able to see disruption of these complexes from all tested small molecule

inhibitors, which is especially interesting in the case of KG-279 and KG-548, which had not yet

been reported as capable of disrupting HIF-2 dimerization, even in single heterodimers (1).

Finally, our work here sheds some much-needed light on the nature of interactions

between CCCs and the HIF-α/ARNT heterodimers. While past work established the importance

of these interactions in HIF-driven transcriptional activation via ARNT (10, 40, 55-59), prior

work from our lab and others led to different, and conflicting, views as to the structural basis of

activator/coactivator recruitment (2, 9, 15, 39). Our complexes here of HIF-1 and HIF-2 bound to

different TACC3 fragments resolve this dilemma, showing TACC3 alongside the length of either

HIF complex, with what we believe to be the C-terminus of TACC3 directly interacting with the

ARNT PAS-B domain based on prior work from our lab (2).. Additional interactions are

indicated by cryo-EM (and HDX-MS for HIF-2), showing the involvement of loops on the HIF-

α subunit. Notably, the HDX-MS data shows a clear stabilization of the ARNT PAS-A/PAS-B

linker upon TACC3 binding; a similar stabilization has been observed with a HIF-2 binding

small molecule activator (24), raising the question about similar activation modes despite the

differences in activator binding to the HIF heterodimer. The recruitment of TACC3 to a single

HIF heterodimer bolsters the idea that HIFs may form markedly different complexes depending

on the functional context; in a case where HIFs cannot form dimer-of-heterodimer complexes,

we believe that other components, such as CCCs, are recruited to augment HIF function.

We close by underscoring that structural biology – like all experimental methods – has its

fundamental strengths and weaknesses. Within HIF biology, the strengths are clear as evidenced

by mechanistic understanding of how natural O_2 regulation is achieved, how disease-associated

disruptions have their effects, and how small molecules can be used therapeutically to target

those diseases. However, we equally see reminders that single structures of complexes, while

extremely useful on their own, may not entirely represent protein assemblies in their true

functional contexts, due to experimental liabilities (*e.g.* presence or absence of protein dynamics)

or experimental choice (*e.g.* selection of proteins, DNAs, small molecules). Our work here

underscores the importance of the latter; with the rich variety of HIF interacting partners and

promoters known and yet to be discovered, many discoveries remain ahead.

Materials and Methods

Cloning and Expression of HIF Complexes

6xHis-tagged wildtype human HIF-2α (residues 5-361) or HIF-1α (residues 4-361) and untagged

human ARNT (residues 91-470) were cloned into a pETDuet-1 vector. HIF-α (6xHis-tagged)

and ARNT were coexpressed in BL21-CodonPlus (DE3) *E. coli* (Agilent) in Lysogeny Broth

with 0.1 mg/mL ampicillin and 0.034 mg/mL chloramphenicol. Cells were grown at 37°C and

induced with 0.1 mM isopropyl-β-D-thiogalactopyranoside (IPTG) upon reaching an OD_{600} of

0.8-1.0, then brought to 16°C to induce overnight (16-18 hr). Cells were harvested by spinning at

4°C and 4658xg for 45 min, resuspended in lysis buffer (50 mM HEPES (pH 7.4), 300 mM

NaCl, 3 mM β-mercaptoethanol (BME), 1 mM phenylmethylsulfonyl fluoride (PMSF), 0.1

mg/mL lysozyme and 0.01 U/mL benzonase nuclease), flash frozen and stored at -80°C.

Cloning and Expression of Coiled-Coil Coactivators

Wildtype human TACC3 (residues 758-838 or residues 788-838 [shTACC3]) or wildtype human

CoCoA (residues 407-535) were cloned into a pHis-GB1 vector. CCCs were then expressed in

BL21 (DE3) *E. coli* (Agilent) in Lysogeny Broth with 0.1 mg/mL ampicillin. Cells were grown

at 37°C and induced with 1 mM isopropyl-β-D-thiogalactopyranoside (IPTG) upon reaching an

OD_{600} of 0.8-1.0, then brought to 17°C to induce overnight (16-18 hr). Cells were harvested by

spinning at 4°C and 4658xg for 45 min, resuspended in lysis buffer (50 mM Tris (pH 7.4), 150

mM NaCl, 1 mM PMSF 0.1 mg/mL lysozyme and 0.01 U/mL benzonase nuclease), flash frozen

and stored at -80°C.

Cloning and Expression of ARNT PAS-B

Wildtype human ARNT PAS-B (residues 356-470) was cloned into a pHis-GB1 expression

vector. ARNT PAS-Bs were then expressed in BL21 (DE3) *E. coli* cells (Agilent) in Lysogeny

Broth with 0.1 mg/mL ampicillin. Cells were grown at 37°C and induced with 1 mM isopropyl-

β-D-thiogalactopyranoside (IPTG) upon reaching an OD_{600} of 0.8-1.0, then brought to 17°C to

induce overnight (16-18 hr). Cells were harvested by spinning down at 4°C and 4658xg for 45

minutes, resuspended in lysis buffer (50 mM Tris (pH 7.4), 150mM NaCl, 2mM BME, 1mM

PMSF), flash frozen and stored at -80°C.

Purification of HIFα-ARNT Heterodimers

Frozen HIF-1 or HIF-2 cell pellets were thawed, lysed by sonication and centrifuged at 47850xg

for 45 min. HIF-α-ARNT was purified using an AKTA Püre (Cytiva) by flowing lysate over a 5

mL HisTrap HP (Cytiva), followed by wash buffer (50 mM HEPES (pH 7.4), 300 mM NaCl, 20

mM imidazole, 2 mM BME) with stepwise increasing concentrations of imidazole (20-260 mM)

to remove nonspecifically bound protein. 6xHis-tagged HIF-α (bound to ARNT) was eluted with

elution buffer (50 mM HEPES (pH 7.4), 300 mM NaCl, 500 mM imidazole, 2 mM BME), mixed

with about a 20x molar excess of annealed DNA (20 bp, 51 bp BS1 or 52 bp BS2, IDT) and 5

mM MgCl$_2$ and allowed to bind for at least one hour at 4°C. Protein was then concentrated about

10x by dialyzing (3.5 kDa MWCO dialysis tubing, ThermoFisher) against dry polyethylene

glycol 35000. HRE-bound HIF-α-ARNT complexes were then further purified through a

Superdex 200 increase 10/300 analytical size exclusion column (Cytiva) with the final analytical

buffer (50 mM HEPES (pH 7.4), 250 mM NaCl, 5 mM MgCl$_2$). Protein samples were used

immediately or stored at 4°C for a maximum of 48 hr.

20 bp HRE: GGCTGCGTACGTGCGGGTCGT

51 bp BS1 HRE:

AGGGGTGGAGGGGGCTGGGCCCTACGTGCTGTCTCACACAGCCTGTCTGAC

52 bp BS2 HRE:

TGGGCCCTACGTGCTGTCTCACACAGCCTGTCTGACCTCTCGACCCTACCGG

Purification of Coiled-Coil Coactivators

Frozen cell pellets were thawed, lysed by sonication and centrifuged at 47850xg for 45 min.

TACC3 or CoCoA were purified using an AKTA Püre (Cytiva) by flowing lysate over a 5 mL

HisTrap HP (Cytiva), followed by wash buffer (50 mM Tris (pH 7.4), 150 mM NaCl, 20 mM

imidazole) with stepwise increasing concentrations of imidazole (39-106 mM) to remove

nonspecifically bound protein. 6xHis-tagged-GB1-CCC (TACC3 or CoCoA) was eluted with

elution buffer (50 mM Tris (pH 7.4), 150 mM NaCl, 500 mM imidazole) and diluted 1:10 in

TEV buffer (50 mM Tris (pH 7.4), 0.1 mM EDTA) with a 3:1000 addition of TEV protease to

cleave the His and GB1-tags from the protein. Cleaved tags were then removed by flowing the

protein over a 5 mL HisTrap HP to separate them from the protein of interest. CCC-containing

fractions were then concentrated about 15x using a 10 kDa MWCO Millipore centrifugal filter

unit (Sigma-Aldrich) at 1200xg and 4°C. Concentrated protein was further purified through a

Superdex 200 increase 16/40 HiScale size exclusion column (Cytiva) with the final analytical

buffer (50 mM HEPES (pH 7.4), 50 mM NaCl, 5 mM MgCl$_2$). Protein samples were used

immediately or stored at 4°C for a maximum of 7 days.

Purification of ARNT PAS-B

Frozen cell pellets were thawed, lysed by sonication and centrifuged at 47850xg for 45 min.

ARNT PAS-B was purified using an AKTA Püre (Cytiva) by flowing lysate over a 5 mL

HisTrap HP (Cytiva), followed by wash buffer (50 mM Tris (pH 7.4), 150 mM NaCl, 20 mM

imidazole, 1 mM DTT) with stepwise increasing concentrations of imidazole (39-106 mM) to

remove nonspecifically bound protein. 6xHis-tagged-GB1-ARNT PAS-B was eluted with elution

buffer (50 mM Tris (pH 7.4), 150 mM NaCl, 500 mM imidazole, 1 mM DTT) and diluted 1:10

in TEV buffer (50 mM Tris (pH 7.4), 0.1 mM EDTA) with a 3:1000 addition of TEV protease to

cleave the His and GB1-tags from the protein. Cleaved tags were then removed by flowing the

protein over a 5 mL HisTrap HP to separate them from the protein of interest. ARNT PAS-B-

containing fractions were then concentrated about 15x using a 10 kDa MWCO Millipore

centrifugal filter unit (Sigma-Aldrich) at 1200xg and 4°C. Concentrated protein was further

purified through a Superdex 75 increase 16/40 HiScale size exclusion column (Cytiva) with the

final analytical buffer (50 mM HEPES (pH 7.4), 15 mM NaCl, 1 mM DTT). Protein samples

were used immediately or stored at -80°C.

Cryogenic Electron Microscopy

Purified HIF complexes were concentrated to a final concentration of 10-20 µM (1-2 mg/mL)

using 10 kDa Pierce Concentrators (ThermoFisher) at 4°C and 1000xg. TACC3 (residues 788-

838) was mixed with the HIF-2 complexes to a final concentration of 70-150 µM and allowed to

bind for ~20 min on ice. 3.5 µL of sample was applied to Quantifoil Au 300 R1.2/1.3 with

Ultrathin Carbon EM grids (Electron Microscopy Sciences) and plunge frozen using a Vitrobot 3

(ThermoFisher) at 100% humidity and 4°C. Frozen grids were then analyzed at the New York

Structural Biology Center (NYSBC) Members Electron Microscopy Center (MEMC) using a

TFS Titan Krios transmission electron microscope (ThermoFisher). Micrographs were collected

at a pixel size of 0.844 Å/px at an exposure rate of 58.94 $e^-/Å^2$/frame with a defocus range of 0.8-

2.0 µm. Particles were picked from micrographs and extracted to a box size of 400x400 pixels

using Warp (41) and analyzed using CryoSPARC (42). Particles were typically sorted through

iterative rounds of 2D classification, *ab initio* reconstruction, and heterogeneous, homogenous

and non-uniform refinement.

Amplified Luminescent Proximity Homogenous Assay (Alpha) Screen

HIF-2 complexes with C-terminally FLAG-tagged (DYKDDDDK) ARNT subunits were

purified and bound to either 20 bp HRE or 51 bp HRE/HAS biotinylated DNA fragments (IDT).

300 nM of the resulting complexes were incubated with 20 µg/mL streptavidin Alpha donor

beads (Revvity), 20 µg/mL anti-FLAG Alpha acceptor beads (Revvity), and varying

concentrations of KG-548, KG-279, compound 2 and PT2385 with a final DMSO concentration

of 1% (v/v) and a final tween-20 concentration of 0.05% (v/v). Automation of sample

preparation was accomplished using an Opentrons OT-2 liquid handler (Opentrons Labworks

Inc.). Samples were allowed to incubate one hour at 25°C in the dark before being analyzed on a

SpectraMax i3 plate reader (Molecular Devices) equipped with an AlphaScreen 384 STD

cartridge (Molecular Devices). Data was collected using SoftMax Pro (Molecular Devices) and

plotted with Grace. All samples were run in triplicate.

Crystallization and Crystallographic Structure Determination of TACC3(758-838)

12.5 mg/mL of purified human TACC3(758-838) and 9.3 mg/mL of human ARNT PAS-B(356-

470) were mixed in 50 mM HEPES, pH 7.4, 50 mM NaCl, 50 mM $MgCl_2$, and 5% glycerol

crystallized in 1 M imidazole, pH 7 and 50% (v/v) MPD via the sitting-drop vapor diffusion

method. The condition was found via high-throughput screening using NeXtal Protein Complex

sparse matrix crystallization screen (NeXtal Technologies; condition H9, the drop contained

equal volumes of protein solution and mother liquor). Crystals were harvested directly from the

screen and cryoprotected with LV CryoOil (MiTeGen) and flash-cooled in liquid N_2 prior to data

collection. Data were collected at the National Synchrotron Light Source II on the AMX (17-ID-

1) beamline at Brookhaven National Laboratory. Data were processed using the autoPROC tool

box (60) resulting in a 2.28 Å data set. While electron density for ARNT PAS-B was not

observed, sufficient density for TACC3 residues 780-836 was visible to allow the structure to be

determined by molecular replacement with Phaser (61) using PDB entry 4PKY (chains B and C)

as the starting search model. Several cycles of refinement were conducted using Coot (62) and

Phenix.refine (63). Data collection, processing and refinement statistics are provided in **Table**

S2. Coordinates for TACC3(758-838) have been deposited in the RCSB with accession number

9YDY, including residues 780-836 on one chain and 782-836 on the other.

Microscale Thermophoresis

Purified HIF complexes bound to carboxyfluorescein (FAM) labeled HRE (IDT) were diluted to

a final concentration of 1μM fluorophore, with an addition of 0.05% tween-20. ARNT PAS-B

proteins (20 μM) were instead labeled with fluorophore at free amino sites using an ALEXA

Fluor 488 N-hydroxysuccinimide (NHS)-ester (Thermo Fisher) by first exchanging into pH 8.2

analytical buffer, allowing incubation with 50 μM of ALEXA 488 NHS ester for 1 hour at 4°C in

the dark, then exchanging back into analytical buffer to remove excess dye. Final protein

concentration is calculated using the following equation (where A_{280} is the absorbance reading at

280 nm, A_{495} is the absorbance reading at 495 nm, and ε_{280} is the extinction coefficient of the

protein at 280 nm) and adjusted to 1 μM.

$$\frac{A_{280} - \left(\left(\frac{A_{495}}{73000}\right) * 12400\right)}{\varepsilon_{280}}$$

A serial dilution of shTACC3 was generated in TACC3 buffer (50 mM HEPES (pH 7.4), 50 mM

NaCl, 5 mM $MgCl_2$). MST samples were generated as a 1:1 (v/v) mix of HIF complex (fixed 500

nM final concentration) to each concentration of the TACC3 serial dilution. Samples were drawn

into standard MST capillaries (NanoTemper) and loaded into a Monolith NT.115 (NanoTemper)

for binding analyses. Data were obtained using MO.Control (NanoTemper) and analyzed using

PALMIST and GUSSI (64).

Structural modeling of HIF complexes

AlphaFold-generated structures were flexibly fit into refined cryo-EM density maps using

ChimeraX (65) and ISOLDE (66) to apply distance restraints. For the HIF-2 dimer of

heterodimers, we flexibly fit our cryo-EM structure of human HIF-2 into the obtained cryo-EM

density. Structures were then further refined using PHENIX (63) to perform global real space

refinement and Coot (62) to perform atomic refinement and remove unresolved regions.

Structural validation was performed using MolProbity and comprehensive validation in PHENIX

(63). For HIF-1 and HIF-2/TACC3 complexes, shTACC3 fragments were fit into corresponding

cryo-EM maps (with the HIF complex docked and fixed) using the Dock in Map function of

PHENIX (63) and assuming that the TACC3 C-terminus is closest to ARNT PAS-B (10). We

emphasize that the HDX-MS data collected on HIF-2/TACC3 complex was not used to guide

this model generation, as it was saved for post-modeling validation. Coordinates for HIF-1 and

HIF-2 complexes were deposited with RCSB under the following accession numbers: 9OF0

(HIF-2 heterodimer/HRE; EMDB EMD-70416), 9OF2 (HIF-2 dimer of heterodimers/BS1

HRE/HAS; EMDB EMD-70418), 9OFU (HIF-1 dimer of heterodimers/BS2 HRE/HAS; EMDB

EMD-70443). Coordinates for models of TACC3/HIF-1/HRE and TACC3/HIF-2/HRE

complexes are deposited at PDB-IHM under the accession numbers 9A9Z (TACC3/HIF-1) and

9A9Y (TACC3/HIF-2).

Size Exclusion Chromatography Coupled with Multi-Angle Light Scattering (SEC-MALS)

Purified HIF complexes were concentrated to a final concentration of 15-20 µM (1-2 mg/mL)

using 10 kDa Pierce Concentrators (ThermoFisher) at 4°C and 1000xg. Samples were then

filtered through an Ultrafree 0.1 µm centrifugal filter (Millipore) at 4°C and 1000xg. 500 µL of

sample were passed through a Superdex 200 10/300 size exclusion column (Cytiva) with the HIF

analytical buffer (50 mM HEPES (pH 7.4), 250 mM NaCl, 5 mM $MgCl_2$) before passing through

a DAWN HELEOS II MALS detector and an Optilab T-rEX Refractive Index Detector

(Wyatt/Waters) using a P-920/UPC-900 AKTA FPLC System (Cytiva). SEC-MALS data was

obtained and analyzed using ASTRA (Wyatt/Waters) and plotted in R.

Size Exclusion Chromatography for HIF-1/HIF-2 Complex Determination

Human HIF-1α (residues 4-361) and ARNT (residues 91-470) were cloned into a pHis-MBP

vector (67) and expressed as detailed above for HIF-complexes. MBP-tagged HIF-1 and HIF-2

were separately lysed through sonication and centrifuged for 45 min at 47850xg and 4°C.

Lysates were then combined and purified together using an AKTA Püre (Cytiva) by flowing

lysate over a 5 mL HisTrap HP (Cytiva), followed by wash buffer (50 mM HEPES, 300 mM

NaCl, 20 mM imidazole, 2 mM BME) with stepwise increasing concentrations of imidazole (20

mM, 68 mM, 164 mM, 260 mM) to remove nonspecifically bound protein. Mixed proteins were

then eluted with elution buffer (50 mM HEPES (pH 7.4), 300 mM NaCl, 500 mM imidazole, 2

mM BME), then mixed with about a 20x molar excess of 51 bp BS1 HRE, 51 bp BS1 DNA with

a mutated HRE-box, or 51 bp BS1 DNA with a mutated HAS-box (IDT) and 5 mM $MgCl_2$ and

allowed to bind at 4°C overnight (10-16 hr). Protein was then concentrated about 10x by

dialyzing (3.5 kDa MWCO dialysis tubing, ThermoFisher) against dry polyethylene glycol

35000. DNA-bound MBP-HIF-1/HIF-2 complexes were further purified through a Superdex 200

increase 16/40 HiScale (Cytiva) with the final analytical buffer (50 mM HEPES, (pH 7.4) 250

mM NaCl, 5 mM $MgCl_2$) to separate complexes via size exclusion chromatography. Complex

components were then visualized using gel electrophoresis. SEC profiles were plotted in R.

Mutant HRE-box BS1 DNA:

AGGGGTGGAGGGGGCTGttaaaTgatatCTGTCTCACACAGCCTGTCTGAC

Mutant HAS-box BS1 DNA:

AGGGGTGGAGGGGGCTGGGCCCTACGTGCTGTCgatatgtcgaCtTCTGAC

Hydrogen Deuterium Exchange Coupled with Mass Spectrometry (HDX-MS)

Purified 20 bp HRE-bound HIF-2 was incubated with shTACC3 for one hour in about a 3:1

molar ratio of TACC3 to HIF-2 (74 μM shTACC3 and 24 μM HIF-2) or an equivalent volume of

TACC3 buffer (50 mM HEPES, 50 mM NaCl, 5 mM $MgCl_2$). Using a LEAP HDX Automation

Platform (Trajan Automation), the protein mixture was deuterated in D_2O buffer (50 mM

HEPES, 50 mM NaCl, 5 mM $MgCl_2$, solvated in 100% D_2O) for varying timepoints (0 s, 50 s,

300 s, 1000 s, 3000 s, 5000 s) after which the deuterated protein mixture was mixed with quench

buffer (3 M guanidinium HCl, 3% acetonitrile, 1% formic acid) at 0°C. Samples were then

loaded onto a Waters Enzymate BEH Pepsin Column for proteolysis followed by a C18

analytical column (Hypersil Gold, 50 mm length × 1 mm diameter, 1.9 μm particle size,

ThermoFisher Scientific) before being loaded onto a maXis-II ETD ESI-qTOF mass

spectrometer (Bruker) for fragment mass analysis. Timepoints both with and without TACC3

present were completed in triplicate. Raw HDX data was analyzed using Bruker Compass Data

Analysis 5.3 and Biotools 3.2 software and further analyzed with HDexaminer (Sierra

Analytics/Trajan Automation) to calculate relative exchange rates before being plotted in R.

Samples were independently run in triplicate.

Reporter assays

Luciferase reporter assays were performed with transiently transfected HEK293T cells. Cells

were plated onto 48-well plates at 100,000 cells/well. Twenty-four hours later, cells were

transfected in triplicate with Lipofectamine 2000 (Invitrogen). Each well received 33 ng reporter

plasmid under control of wild-type or mutant human 3' erythropoietin (Epo) enhancer fused to a

firefly luciferase cassette (pGL3-Basic, Promega), 200 ng empty expression plasmid (pIRES-

hrGFP-2a, Stratagene), and 100 ng plasmid expressing oxygen insensitive "PPN" HIF-1α or

HIF-2α variants, which contain mutations in two proline and one asparagine residue that are

normally subject to O_2-dependent post-translational modifications (68). At 24 hr post-

transfection, cells were rinsed one time with PBS (ThermoFisher), followed by lysis with 50 µL

core luciferase buffer (30 mM tricine (pH 7.8), 8 mM magnesium acetate, 0.2 mM EDTA, 1%

Triton X-100, and 100 mM 2-mercaptoethanol). Luciferase assays were performed by adding 50

µL substrate buffer (core buffer supplemented with 2.5 mM $MgCl_2$, 1.5 mM ATP, 0.5 mM

coenzyme A, and 0.5 mM D-luciferin) to 10 µL cell lysate and immediately reading

luminescence on a CLARIOstar Plus plate reader (BMG Labtech). Long oligos with Nhe I and

Xho I restriction sites at the 5' and 3' ends, respectively, were used to position the 3' human Epo

enhancer upstream of a minimal mouse Epo promoter. Sequences of the 3' human Epo enhancer

sequences used in this study are indicated below (mutations and restriction sites in lowercase):

<u>WT:</u>

gctagcAGATCTGGGAAACGAGGGGTGGAGGGGGCTGGGCCCTACGTGCTGTCTCACA

CAGCCTGTCTGACCTCTCGACCCTACCGGGCCTGAGGCCACAAGCTCTGCCTACGCT

GGTCAATAAGGTGGCTCCATTCAAGGCCTCACCGCAGTAAGGCAGCTGCCAACCCTc

gag

<u>GC-rich mutant:</u>

gctagcAGATCTGGGAAACGAGGctaGaAttcctCTGGGCCCTACGTGCTGTCTCACACAGCC

TGTCTGACCTCTCGACCCTACCGGGCCTGAGGCCACAAGCTCTGCCTACGCTGGTCA

ATAAGGTGGCTCCATTCAAGGCCTCACCGCAGTAAGGCAGCTGCCAACCCTcgag

<u>HRE mutant:</u>

gctagcAGATCTGGGAAACGAGGGGTGGAGGGGGCTGttaaaTgatatCTGTCTCACACAGCC

TGTCTGACCTCTCGACCCTACCGGGCCTGAGGCCACAAGCTCTGCCTACGCTGGTCA

ATAAGGTGGCTCCATTCAAGGCCTCACCGCAGTAAGGCAGCTGCCAACCCTcgag

<u>HAS mutant:</u>

gctagcAGATCTGGGAAACGAGGGGTGGAGGGGGCTGGGCCCTACGTGCTGTCgatatgtcg

aCtTCTGACCTCTCGACCCTACCGGGCCTGAGGCCACAAGCTCTGCCTACGCTGGTCA

ATAAGGTGGCTCCATTCAAGGCCTCACCGCAGTAAGGCAGCTGCCAACCCTcgag

Acknowledgements

We thank Dan Lee and Julia Simon for technical assistance, Soumendu Boral and other members

of the Gardner laboratory as well as Rick Bruick, Uttam Tambar, Qing Zhang, and Derek Tan for

helpful discussions. This work was supported by NIH grants U54 CA132378 and U54

CA137788 (to K.H.G.), Mathers Foundation grant MF-2112-02288 (to K.H.G. and J.A.G.), and

funds from the Department of Veterans Affairs I01BX000446 (to J.A.G.). Cryo-EM data

collection was performed at the Members Electron Microscopy Center at the New York

Structural Biology Center, with major support from the Simons Foundation (SF349247). This

research used the AMX and FMX beamlines of the National Synchrotron Light Source II, a U.S.

References

1. X. Xu *et al.*, Identification of Small Molecule Ligand Binding Sites On and In the ARNT PAS-B
Domain. *Journal of Biological Chemistry* **300**, 107606 (2024).
2. Y. Guo *et al.*, Regulating the ARNT/TACC3 axis: Multiple approaches to manipulating
protein/protein interactions with small molecules. *ACS chemical biology* **8**, 626-635 (2013).
3. G. L. Semenza, Hypoxia-inducible factor 1 (HIF-1) pathway. *Science's STKE: signal
transduction knowledge environment* **2007**, cm8 (2007).
4. M. Strowitzki, E. Cummins, C. Taylor, Protein Hydroxylation by Hypoxia-Inducible Factor (HIF)
Hydroxylases: Unique or Ubiquitous? *Cells* **8**, 384 (2019).
5. D. Lando *et al.*, FIH-1 is an asparaginyl hydroxylase enzyme that regulates the transcriptional
activity of hypoxia-inducible factor. *Genes & Development* **16**, 1466-1471 (2002).
6. D. Lando, D. J. Peet, D. A. Whelan, J. J. Gorman, M. L. Whitelaw, Asparagine Hydroxylation of
the HIF Transactivation Domain: A Hypoxic Switch. *Science* **295**, 858-861 (2002).
7. T. H. Scheuermann *et al.*, Allosteric Inhibition of Hypoxia Inducible Factor-2 with Small
Molecules. *Nature chemical biology* **9**, 271-276 (2013).
8. P. M. Wehn *et al.*, Design and Activity of Specific Hypoxia-Inducible Factor-2α (HIF-2α)
Inhibitors for the Treatment of Clear Cell Renal Cell Carcinoma: Discovery of Clinical Candidate
(S)-3-((2,2-Difluoro-1-hydroxy-7-(methylsulfonyl)-2,3-dihydro-1H-inden-4-yl)oxy)-5-
fluorobenzonitrile (PT2385). *Journal of Medicinal Chemistry* **61**, 9691-9721 (2018).
9. D. Wu, N. Potluri, J. Lu, Y. Kim, F. Rastinejad, Structural integration in hypoxia-inducible
factors. *Nature* **524**, 303-308 (2015).
10. C. L. Partch, K. H. Gardner, Coactivators necessary for transcriptional output of the hypoxia
inducible factor, HIF, are directly recruited by ARNT PAS-B. *Proceedings of the National
Academy of Sciences* **108**, 7739-7744 (2011).
11. D. C. Bersten, A. E. Sullivan, D. J. Peet, M. L. Whitelaw, bHLH-PAS proteins in cancer. *Nature
Reviews Cancer* **13**, 827-841 (2013).
12. N. Albadari, S. Deng, W. Li, The transcriptional factors HIF-1 and HIF-2 and their novel
inhibitors in cancer therapy. *Expert Opinion on Drug Discovery* **14**, 667-682 (2019).
13. S. Kaluz, M. Kaluzová, E. J. Stanbridge, Regulation of gene expression by hypoxia: integration
of the HIF-transduced hypoxic signal at the hypoxia-responsive element. *Clinica chimica acta;
international journal of clinical chemistry* **395**, 6-13 (2008).
14. T. Rosell-Garcia *et al.*, Multimerization of HIF enhances transcription of target genes containing
the hypoxia ancillary sequence. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory
Mechanisms* **1866**, 194963 (2023).
15. Y. Guo, T. H. Scheuermann, C. L. Partch, D. R. Tomchick, K. H. Gardner, Coiled-coil
Coactivators Play a Structural Role Mediating Interactions in Hypoxia-inducible Factor
Heterodimerization. *The Journal of Biological Chemistry* **290**, 7707-7721 (2015).
16. J. C. Jun, A. Rathore, H. Younas, D. Gilkes, V. Y. Polotsky, Hypoxia-Inducible Factors and
Cancer. *Current sleep medicine reports* **3**, 1-10 (2017).
17. S. Mazumder, P. J. Higgins, R. Samarakoon, Downstream Targets of VHL/HIF-α Signaling in
Renal Clear Cell Carcinoma Progression: Mechanisms and Therapeutic Relevance. *Cancers* **15**,
1316 (2023).
18. E. E. Wicks, G. L. Semenza, Hypoxia-inducible factors: cancer progression and clinical
translation. *The Journal of Clinical Investigation* **132**, e159839.
19. C.-J. Hu, L.-Y. Wang, L. A. Chodosh, B. Keith, M. C. Simon, Differential roles of hypoxia-
inducible factor 1alpha (HIF-1alpha) and HIF-2alpha in hypoxic gene regulation. *Molecular and
Cellular Biology* **23**, 9361-9374 (2003).
20. E. B. Rankin *et al.*, Hypoxia-inducible factor–2 (HIF-2) regulates hepatic erythropoietin in vivo.
The Journal of Clinical Investigation **117**, 1068-1077 (2007).

21. S. Hara, J. Hamada, C. Kobayashi, Y. Kondo, N. Imura, Expression and Characterization of
Hypoxia-Inducible Factor (HIF)-3α in Human Kidney: Suppression of HIF-Mediated Gene
Expression by HIF-3α. *Biochemical and Biophysical Research Communications* **287**, 808-813
(2001).
22. A. M. Fala *et al.*, Unsaturated fatty acids as high-affinity ligands of the C-terminal Per-ARNT-
Sim domain from the Hypoxia-inducible factor 3α. *Scientific Reports* **5**, 12698 (2015).
23. Z.-j. Liu, G. L. Semenza, H.-f. Zhang, Hypoxia-inducible factor 1 and breast cancer metastasis.
Journal of Zhejiang University Science B **16**, 32-43 (2015).
24. D. Wu *et al.*, Bidirectional Modulation of HIF-2 Activity through Chemical Ligands. *Nature*
chemical biology **15**, 367-376 (2019).
25. E. Jonasch *et al.*, Belzutifan for Renal Cell Carcinoma in von Hippel–Lindau Disease. *New*
England Journal of Medicine **385**, 2036-2046 (2021).
26. T. K. Choueiri *et al.*, Inhibition of hypoxia-inducible factor-2α in renal cell carcinoma with
belzutifan: a phase 1 trial and biomarker analysis. *Nature medicine* **27**, 802-805 (2021).
27. X. Diao *et al.*, Identification of oleoylethanolamide as an endogenous ligand for HIF-3α. *Nature*
Communications **13**, 2529 (2022).
28. W. Ruan *et al.*, BMAL1–HIF2A heterodimer modulates circadian variations of myocardial
injury. *Nature* **641**, 1017-1028 (2025).
29. N. Huang *et al.*, Crystal Structure of the Heterodimeric CLOCK:BMAL1 Transcriptional
Activator Complex. *Science (New York, NY)* **337**, 189-194 (2012).
30. G. L. Semenza, G. L. Wang, A nuclear factor induced by hypoxia via de novo protein synthesis
binds to the human erythropoietin gene enhancer at a site required for transcriptional activation.
Molecular and Cellular Biology **12**, 5447-5454 (1992).
31. B. L. Ebert, H. F. Bunn, Regulation of Transcription by Hypoxia Requires a Multiprotein
Complex That Includes Hypoxia-Inducible Factor 1, an Adjacent Transcription Factor, and
p300/CREB Binding Protein. *Molecular and Cellular Biology* **18**, 4089-4096 (1998).
32. H. Kimura *et al.*, Identification of Hypoxia-inducible Factor 1 Ancillary Sequence and Its
Function in Vascular Endothelial Growth Factor Gene Induction by Hypoxia and Nitric Oxide *.
Journal of Biological Chemistry **276**, 2292-2298 (2001).
33. F. Corrêa, J. Key, B. Kuhlman, Kevin H. Gardner, Computational Repacking of HIF-2α Cavity
Replaces Water-Based Stabilized Core. *Structure* **24**, 1918-1927 (2016).
34. X. Ren, X. Diao, J. Zhuang, D. Wu, Structural basis for the allosteric inhibition of hypoxia-
inducible factor (HIF)-2 by belzutifan. *Molecular Pharmacology* 10.1124/molpharm.122.000525,
MOLPHARM-AR-2022-000525 (2022).
35. T. H. Scheuermann *et al.*, Artificial ligand binding within the HIF2α PAS-B domain of the HIF2
transcription factor. *Proceedings of the National Academy of Sciences* **106**, 450-455 (2009).
36. S. A. Dames, M. Martinez-Yamout, R. N. De Guzman, H. J. Dyson, P. E. Wright, Structural basis
for Hif-1α/CBP recognition in the cellular hypoxic response. *Proceedings of the National*
Academy of Sciences **99**, 5271-5276 (2002).
37. T. V. Beischlag *et al.*, Recruitment of Thyroid Hormone Receptor/Retinoblastoma-interacting
Protein 230 by the Aryl Hydrocarbon Receptor Nuclear Translocator Is Required for the
Transcriptional Response to Both Dioxin and Hypoxia *. *Journal of Biological Chemistry* **279**,
54620-54628 (2004).
38. M. P. Labrecque *et al.*, A TRIP230-Retinoblastoma Protein Complex Regulates Hypoxia-
Inducible Factor-1α-Mediated Transcription and Cancer Cell Invasion. *PLOS ONE* **9**, e99214
(2014).
39. C. L. Partch, P. B. Card, C. A. Amezcua, K. H. Gardner, Molecular Basis of Coiled Coil
Coactivator Recruitment by the Aryl Hydrocarbon Receptor Nuclear Translocator (ARNT)*.
Journal of Biological Chemistry **284**, 15184-15192 (2009).
40. J. H. Kim, M. R. Stallcup, Role of the Coiled-coil Coactivator (CoCoA) in Aryl Hydrocarbon
Receptor-mediated Transcription*. *Journal of Biological Chemistry* **279**, 49842-49848 (2004).

41. D. Tegunov, P. Cramer, Real-time cryo-electron microscopy data preprocessing with Warp.
Nature Methods **16**, 1146-1152 (2019).
42. A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nature Methods* **14**, 290-296 (2017).
43. A. Punjani, H. Zhang, D. J. Fleet, Non-uniform refinement: adaptive regularization improves
single-particle cryo-EM reconstruction. *Nature Methods* **17**, 1214-1221 (2020).
44. V. L. Dengler, M. Galbraith, J. M. Espinosa, Transcriptional Regulation by Hypoxia Inducible
Factors. *Critical reviews in biochemistry and molecular biology* **49**, 1-15 (2014).
45. A. Yasgar, A. Jadhav, A. Simeonov, N. P. Coussens, AlphaScreen-Based Assays: Ultra-High-
Throughput Screening for Small-Molecule Inhibitors of Challenging Enzymes and Protein-
Protein Interactions. *Methods in molecular biology (Clifton, NJ)* **1439**, 77-98 (2016).
46. J. L. Rogers *et al.*, Development of Inhibitors of the PAS-B Domain of the HIF-2α Transcription
Factor. *Journal of medicinal chemistry* **56**, 1739-1747 (2013).
47. G. L. Wang, G. L. Semenza, General involvement of hypoxia-inducible factor 1 in transcriptional
response to hypoxia. *Proceedings of the National Academy of Sciences* **90**, 4304-4308 (1993).
48. J. Contenti *et al.*, HIF-1 inactivation empowers HIF-2 to drive hypoxia adaptation in aggressive
forms of medulloblastoma. *Cell Death Discovery* **10** (2024).
49. M. Y. Koh, G. Powis, Passing the baton: the HIF switch. *Trends in Biochemical Sciences* **37**,
364-372 (2012).
50. R. M. Mello *et al.* (2024) BMAL1-HIF2α heterodimers contribute to ccRCC. (Cold Spring
Harbor Laboratory).
51. A. K. Michael *et al.*, Cooperation between bHLH transcription factors and histones for DNA
access. *Nature* **619**, 385-393 (2023).
52. D. Sharma *et al.*, Tandem association of CLOCK:BMAL1 complexes on DNA enables
recruitment of CBP/p300 through multivalent interactions. *BioRxiv* (2025).
53. D. Wu, X. Su, N. Potluri, Y. Kim, F. Rastinejad, NPAS1-ARNT and NPAS3-ARNT crystal
structures implicate the bHLH-PAS family as multi-ligand binding transcription factors. *eLife* **5**,
e18790.
54. X. Diao *et al.*, Structural basis for the ligand-dependent activation of heterodimeric AHR-ARNT
complex. *Nature Communications* **16** (2025).
55. J. H. Kim, H. Li, M. R. Stallcup, CoCoA, a Nuclear Receptor Coactivator which Acts through an
N-Terminal Activation Domain of p160 Coactivators. *Molecular Cell* **12**, 1537-1549 (2003).
56. J. H. Kim, C. K. Yang, M. R. Stallcup, Downstream signaling mechanism of the C-terminal
activation domain of transcriptional coactivator CoCoA. *Nucleic Acids Research* **34**, 2736-2750
(2006).
57. J. H. Nahm *et al.*, Transforming acidic coiled-coil-containing protein 3 (TACC3) overexpression
in hepatocellular carcinomas is associated with "stemness" and epithelial-mesenchymal
transition-related marker expression and a poor prognosis. *Tumour Biology: The Journal of the
International Society for Oncodevelopmental Biology and Medicine* **37**, 393-403 (2016).
58. Y.-L. Wen *et al.*, Transforming acidic coiled-coil protein-3: a novel marker for differential
diagnosis and prognosis prediction in endocervical adenocarcinoma. *Molecular Medicine* **27**, 60
(2021).
59. C. L. Partch, K. H. Gardner, Coactivator recruitment: A new role for PAS domains in
transcriptional regulation by the bHLH-PAS family. *Journal of Cellular Physiology* **223**, 553-557
(2010).
60. C. Vonrhein *et al.*, Data processing and analysis with the autoPROC toolbox. *Acta
Crystallographica Section D Biological Crystallography* **67**, 293-302 (2011).
61. A. J. McCoy *et al.*, Phaser crystallographic software. *Journal of Applied Crystallography* **40**,
658-674 (2007).
62. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta
Crystallographica Section D, Biological Crystallography* **66**, 486-501 (2010).

63.   D. Liebschner *et al.*, Macromolecular structure determination using X-rays, neutrons and
1049   electrons: recent developments in Phenix. *Acta Crystallographica Section D: Structural Biology*
1050   **75**, 861-877 (2019).
1051   64.   T. H. Scheuermann, S. B. Padrick, K. H. Gardner, C. A. Brautigam, On the acquisition and
1052   analysis of microscale thermophoresis data. *Analytical Biochemistry* **496**, 79-93 (2016).
1053   65.   E. C. Meng *et al.*, UCSF ChimeraX: Tools for structure building and analysis. *Protein Science*
1054   **32**, e4792 (2023).
1055   66.   T. I. Croll, ISOLDE: a physically realistic environment for model building into low-resolution
1056   electron-density maps. *Acta Crystallographica Section D: Structural Biology* **74**, 519-530 (2018).
1057   67.   P. Sheffield, S. Garrard, Z. Derewenda, Overcoming Expression and Purification Problems of
1058   RhoGDI Using a Family of "Parallel" Expression Vectors. *Protein Expression and Purification*
1059   **15**, 34-39 (1999).
1060   68.   E. M. Dioum *et al.*, Regulation of Hypoxia-Inducible Factor 2α Signaling by the Stress-
1061   Responsive Deacetylase Sirtuin 1. *Science* **324**, 1289-1293 (2009).
1062

A

100 Å

B

Detached/blurred
ARNT PAS-B

Missing DNA/
Unresolved bHLH

20-bp HRE GGCTGCG**TACGTG**CGGGTCGT
HRE Box

C

HIF-2α-ARNT-20-bp HRE
PAS-B
PAS-A
PAS-B
PAS-A
bHLH
HRE
bHLH

D

Mouse xtal
HIF-2α-ARNT-
20-bp HRE
Human cryo-EM
HIF-2α-ARNT-
20-bp HRE

E

ARNT
PAS-B

Hβ   Gβ   Fα

3.5 3.6 3.7 3.8 3.9 Å

B

HIF-2α-ARNT/HIF-2α-ARNT
-51-bp HRE/HAS

BS1 51-bp HRE/HAS AGGGGTGGAGGGGGGCTGGGCCC**TACGTG**CTGTCTCA**CACA**GCCTGTCTGAC
BS2 52-bp HRE/HAS   TGGGCCC**TACGTG**CTGTCTCA**CACA**GCCTGTCTGACCTCTCGACCCTACCGG
                          HRE Box   HAS Box

D

HIF-1α-ARNT/HIF-1α-ARNT
-51-bp HRE/HAS