Modulation of error sensitivity during motor learning across time, space and environment variability

Dmitrii Todorov[1,2], Maelys Moulin[1,3,4], Ethan Buch[5], Romain Quentin[1]

1. EDUWELL Team, Lyon Neuroscience Research Center (CRNL), INSERM U1028, CNRS UMR5292, Université Claude Bernard Lyon 1, Lyon, France;
2. Sorbonne Université, CNRS, Inserm, Laboratoire d'Imagerie Biomédicale, LIB, F-75006, Paris, France;
3. IDH Team, LIRMM, CNRS, University of Montpellier, Montpellier, France;
4. EuroMov Digital Health in Motion, University of Montpellier, IMT Mines Alès, Montpellier, France;
5. Human Cortical Physiology and Neurorehabilitation Section, NINDS, NIH, Bethesda, Maryland 20892

Keywords: Motor learning, Error sensitivity, Movement variability, Reaching movements, Generalization

Abstract

We never experience the exact same situation twice. In our dynamic and constantly changing environment, we continuously need to adapt our behavior, either due to external (e.g., a change of wind) or internal factors (e.g., muscle noise). Motor adaptation is the process of recalibration of movements in response to such perturbation. It activates when an agent needs to change their movement in response to a perceived error. It has been proposed that in order to perform motor adaptation, the brain continuously updates an error sensitivity signal that controls how much is learned from a past error. Such error sensitivity reflects the speed of learning, and understanding its dynamics is crucial to promoting efficient learning. In our experiment, healthy participants performed a visuomotor adaptation task requiring reaching multiple targets using a joystick. They experienced two periods of stable perturbations and two periods of random perturbations. In agreement with past research, we found that error sensitivity is lower in unstable environments. Moreover, error sensitivity is higher in a stable environment without perturbation than in a stable environment with perturbations. We observed a continuous increase of the error sensitivity within a stable perturbation environment. Finally, continuous spatial (across target locations) and temporal (across trials with same target) generalization of learning is present not only in stable but also in the random environment.

NEW & NOTEWORTHY

We investigated how error sensitivity evolves and generalizes across different environmental stability settings. We show that error sensitivity increases over time when perturbations are stable and that error sensitivity is higher in environments without extrinsic perturbations than in environments where they are present. Finally, we evidence a generalization pattern present independently of environment predictability.

KEYWORDS

Motor adaptation, Reaching movements, Visuomotor rotation, Error sensitivity, Adaptation rate, Generalization, error-based adaptation

Introduction

Motor learning corresponds to the improvement of motor skills through practice. It is an essential part of human behavior. Motor adaptation is a particular type of motor learning where an agent adjusts their movements through trial and error and in response to a change in the environment(Malone, Vasudevan, and Bastian 2011). In humans performing reaching movements, the motor adaptation process is largely driven by the perceived error(Morehead and Orban de Xivry 2021). The degree to which such errors are used to update the next movement is determined by so-called *error sensitivity* (ES)(Herzfeld et al. 2014) -- that is, an error-and-time-dependent adaptation rate. While many studies have initially assumed that the error sensitivity is constant (Donchin, Francis, and Shadmehr 2003; Jordan and Rumelhart 1995; Kawato, Furukawa, and Suzuki 1987; Thoroughman and Shadmehr 2000), more recent studies have investigated its dynamics and propose computational models that describe how error sensitivity is updated based on past behavior. For example, it was posited that the update of error sensitivity depends on the sign consistency between two consecutives errors (if two consecutives errors are of the same sign, error sensitivity will increase) (Herzfeld et al. 2014). Another model suggested that error sensitivity is dynamically modulated, depending on the mean and standard deviation across a temporal window of previous errors (Tan, Jenkinson, and Brown 2014).

Another important phenomenon in motor adaptation is *generalization*. When adapting to a new environment, we rarely start from zero. Generalization is the transfer of knowledge from one context to another and it is a fundamental part of the motor adaptation process (Krakauer et al. 2006). We could consider different types of generalization: across space (adaptation uses information about the experience that happened on the immediately preceding trial even if it involved a target located at a different place in space) and across time (adaptation uses information about the experience that happened during the last encounter of the same target as currently visible, even if it did not happen on the previous trial). Prior studies of spatial generalization have investigated this phenomenon from many different angles but always assuming static learning rate (Krakauer et al. 2006; Tanaka and Sejnowski 2014). The generalization of learning between two spatially distinct targets is modulated by the distance between them (Donchin et al. 2003). The closer the two targets are, the greater the extent of adaptation generalization. Decay of generalization as a function of distance between targets is not solely due to muscle patterns being different for different targets for force field (Berniker et al. 2014) and digitizing tablet (Brayanov, Press, and Smith 2012). To our knowledge, very few studies have investigated temporal generalization (Donchin et al. 2003). We studied generalization of error sensitivity across time (trials) and space (target position).

Only few studies investigated such trial-resolved dynamics of error sensitivity and only a handful of perturbation schedules were explored. In our study, twenty participants performed a visuomotor adaptation task (Fig 1) in which they had to reach one out of four possible targets by controlling a cursor with a joystick at each trial. We specifically studied how error sensitivity evolves within environmental conditions with predictable ("stable") perturbation and with completely unpredictable ("random") perturbations. Error sensitivity was larger in stable than in random environment but only when including zero-perturbation trials (Fig 2, C and D). Interestingly, error sensitivity in a stable environment was the highest right before the perturbation was introduced, dropped at the onset of a new perturbation (Fig 2C) and grew during the perturbation or washout, similarly to (Herzfeld et al. 2014) (Fig 3). Such an increase of error sensitivity during a stable environment survives even when the previous error size is controlled for. Our results also demonstrate that participants generalize their learning from errors across space and time with a monotonous generalization curve, both in stable and in random environments (Fig 4). We found that, in our data, error sensitivity values are not fully consistent with two well-known models: error-sign-consistency model (Herzfeld et al. 2014) and error sensitivity being ratio of squared mean and variance inside a temporal window of 20 trials (Tan et al. 2014)). We did however find that error

sensitivity is related to another windowed statistic: the inverse of the windowed standard deviation with history lengths between 8 and 22 trials.

Overall these results improve our understanding of motor adaptation, which in turn could help developing improved motor rehabilitation protocols for conditions such as stroke in the future. (Krakauer 2006; Krakauer and Carmichael 2022).

Methods

Participants and experimental sessions.

Twenty healthy and right-handed volunteers (15 women, 5 men, mean age of 25.9 +/- 4.1 years) participated in the study after providing their written informed consent to participate in the project, which was approved by the Combined Neuroscience Institutional Review Board of the National Institute of Health (NIH). They all had normal physical and neurological examinations and normal or corrected-to-normal vision.

Visuomotor adaptation task.

Visual stimuli were displayed using pygame Python 3 package (https://www.pygame.org). The visual stimuli were back-projected on a translucent screen in front of the participants. The screen was positioned at a distance of 80 cm from both eyes of the participants. Each trial started with a black circle (called the home, radius = 1.5 cm) at the center of the screen (Fig 1). Participants were instructed to position a white cursor (filled circle, radius = 0.6 cm) inside the home position using a joystick fiber optic response pad (Current Designs, Inc., USA) with their right hand. After maintaining the cursor position inside the home position perimeter for 464 ms, a green target (filled circle, radius = 0.9 cm) appeared at one out of 4 possible positions (22.5°, 67.5°, 112.5° and 157.5 on a unit circle centered on the home position with a radius of 20 cm), while the white cursor simultaneously disappeared. Participants then used the joystick to move the cursor towards the target position. The (unperturbed) cursor position coordinates were a simple linear transformation of the two joystick outputs (horizontal and vertical angles) provided by the joystick driver. The visual feedback of the cursor was absent during the joystick movement – only reappearing as static endpoint feedback once the home-to-cursor distance reached 20 cm (i.e., – passed through the invisible target array circle), regardless of whether the subject hit or missed the target. The movement lasted on average 750+/-190 ms. Participants used this endpoint feedback, which was presented for a total duration of 232ms, to perceive their error (i.e. the target-to-end-point cursor feedback distance). After an inter-trial interval of 1500 ms, participants returned the cursor to the home position with the joystick. The cursor had to remain again within the home position perimeter for 464 ms in order to trigger the next trial target presentation. The entire trial lasted on average 3.729 +/- 0.332 s. The task was composed of four blocks of 192 trials each (768 trials in total) and lasted on average 50.3+/-5.07 minutes. A stable environment block (Fig 1) consisted of 24 trials without perturbation, followed by 60 trials with a +30° (for half of the participants) or -30° (for another half) visual perturbation, followed by 24 trials without perturbation, followed by 60 trials with a -30° (or +30° if the first perturbation was -30°) and followed by 24 trials without perturbation. Participants were pseudo-randomly assigned to start with a +30° or -30° perturbation. In the second and the fourth blocks, the perturbation was applied pseudo-randomly over the 192 trials. More precisely, errors presented during these blocks were randomly permuted replays of the errors experienced by the participant in the previous stable block. Thus, end-point cursor feedback presented on each trial during the random perturbation block had no fixed relationship to the actual movement performed (such trials

are often called "error clamp" trials (Morehead and Orban de Xivry 2021) but are usually used differently). This design approach ensured that the overall distribution of errors was controlled and kept consistent between the stable and random environments..

Data analysis

Fig 1. Experimental design. Participants controlled a cursor with a joystick to reach targets appearing on the screen. On some trials, visual feedback was perturbed by a visuomotor rotation of 30 degrees clockwise or counterclockwise.

Half of participants started with a +30° perturbation and another half started with a -30° perturbation. To simplify group analysis, we inverted the signs of errors, reach angles and target angles for participants who started with the -30° perturbation and further conducted the analyses only with these values.

We define error sensitivity at the beginning of trial n using the following formula

$$ES(n) = \frac{\left(RA_{n-1} - TA_{n-1}\right) - \alpha \cdot \left(RA_n - TA_n\right)}{e_{n-1}}, \tag{1}$$

where RA_{n-1} and RA_n are the angles between the endpoints of the subject movement and the origin *before visuomotor perturbation is applied* at trials $n-1$ and n, and TA_{n-1} and TA_n are the target angles at these trials (target location is never perturbed by the visuomotor perturbation but the target locations generally change from trial to trial), e_{n-1} is the perceived visual error at the feedback onset of trial n-1 and α is the retention factor. We emphasize that here, ES(n) represents the value of error sensitivity at the start of the trial, though it is determined post-hoc by analyzing the reach angle at the trial's end. We used the value $\alpha = 0.924$ following (Avraham, Keizman, and Shmuelof 2020).

Hit counts

A trial is considered a "hit" when, at the end of the trial, the displayed cursor circle intersects with the target circle. Participants hit the target on average in 15.7%, ± 2.9% of all trials. Mean hit proportions across environments are identical because errors during random blocks were drawn from the preceding stable block.

We remove trials for which previous trials were hit trials from consideration because there is no error to correct in the next movement. We also remove trials where error sensitivity exceeds 5 times the mean standard deviations within participants (corresponding to 0.85% of remaining trials after excluding trials with a preceding hit.

Results

Error Sensitivity is higher when no *external* perturbation is present

Fig 2. **A** Error dynamics. **B** Error sensitivity dynamics (mean over participants, shading represents standard deviation, purple line represents perturbation, text on top denotes experimental conditions). **C** Average error sensitivity in stable and random environments (averaged within participant). **D** same as **C** but with a more fine-grained stable environment, colors are consistent with the text colors in panel A. On both plots **C** and **D** all values shown are significantly positive. The p-values codes are after Holm-Bonferroni correction applied.

Error sensitivity is predominantly positive in all environments (Fig 2D, one-sample t-tests against zero, all p<0.05, during perturbation: error sensitivity = 0.19 ± 0.22, t=3.99, p=3.92e-04; during no perturbation condition: error sensitivity = 0.99 ± 0.24, t=18.29, p=3.22e-13, during washout condition: error sensitivity = 0.48 ± 0.18, t=12.18, p=3.01e-10; during random condition: error sensitivity = 0.19 ± 0.12, t=7.19 p=7.87e-07). All p-values are Holm-Bonferroni corrected for multiple comparisons. It demonstrates that participants compensate for their errors even in the random environment. Negative error sensitivity values are occasionally observed at the trial level, which can be attributed to movement noise when participants use a joystick to control a cursor.

Error sensitivity is significantly higher in stable (mean=0.33±0.16) than random (mean=0.19±0.12) environment (two sample t-test, t=-5.19, p=5.24e-05) (Fig. 2C).

To compare error sensitivity in sub-conditions, repeated measures ANOVA revealed a significant effect of condition on error sensitivity (F(3,57)=85.70, p<1e-3, generalized eta-squared=0.749). We then conducted post-hoc pairwise comparisons using t-tests with Holm-Bonferroni corrections. Error sensitivity sharply increases when no perturbation is present. Specifically, as visible on Fig 2B, error sensitivity is higher during the no perturbation period (mean=0.99±0.24) compared to periods with stable perturbations (mean=0.19±0.22) (t=10.55, p=1.11e-08). Error sensitivity is also higher during washout periods (mean=0.48±0.18) compared to periods with stable perturbations (t=5.38, p=6.88e-0.5). Surprisingly, error sensitivity during stable perturbations is not significantly different from the error sensitivity in the random condition (t=0.05, p=9.59e-01).

The results underscore the complexity of motor adaptation, highlighting the influence of environmental predictability on the learning process.

Error sensitivity grows in stable environment

Fig 3. Error sensitivity dynamics in different experiment stages A: Correlations between error sensitivity and trial number within environments, B: Partial correlations between error sensitivity and trial number, controlled for the absolute error magnitude on the same trial. The corrected trial number corresponds to the residuals of the correlation between trial number and absolute error size and the corrected error sensitivity corresponds to the

residuals of the correlation between error sensitivity and absolute error size. Solid blue line represents the mean over participants and conditions, shading represents standard deviations. No perturbation, Washout and Random have two sets of data representing two separate appearances of the condition for each subject, Perturbation has four such sets (see Fig 2A). Dashed lines are linear fits. The mean spearman rho across participants and the significance of the one-sample t-test against zero is annotated for each panel. These results show that error sensitivity grows during perturbation and washout periods. Colors correspond to colors in Fig 2A,D.

Our results demonstrate that error sensitivity grows with time during perturbation and washout periods of stable environment (one-sample t-test against zero on the distribution of spearman rho across participants, for the perturbation condition: rho=0.15+/-0.08, p=8.38e-08, for the washout condition: r_{mean}=0.27+/-0.14, p=1.12e-07). To account for the potential influence of error magnitude on these correlations, partial correlations conditioned on the absolute error size of the previous trial error were also examined and results were similar (for the perturbation condition: rho=0.24+/-0.08, p=9.95e-11, for the washout condition: rho=0.28+/-0.14, p=4.88e-08). All p-values in this paragraph are Holm-Bonferroni corrected.

The perturbation condition exhibited a stronger positive correlation compared to the random condition (paired two-sample t-test comparing rho values: mean T = 9.30, p-value = 1.01e-07, suggesting a more pronounced increase in error sensitivity as participants adjusted back to a non-perturbed state. For partial correlation this held as well (T=12.76, p-value=5.49e-10). In washout the correlation and partial correlation were larger than in no perturbation condition (for regular correlation T=3.53 p=1.12e-02, for partial correlation T=4.34 p=1.76e-03). Unsurprisingly rho values in washout are larger than in the random condition as well (for regular correlation T=8.19 p=6.53e-07, for partial correlation T=8.63 p=2.94e-07). All p-values in this paragraph are Holm-Bonferroni corrected.

Generalization across time and space in stable and random environments

Fig 4. Generalization patterns. A and C. Spatial generalization in stable and random environments. Error sensitivities are measured separately depending on the distance between targets in two consecutive trials. B and D. Temporal generalization in stable and random environments. Error sensitivities are measured as the correction of either the preceding error (n-1) or the error at n-2, n-3 or n-4 trials. Error sensitivities are averaged within participants and then across participants. Solid line shows a polynomial fit to means. The top line of significance codes are for comparison of error sensitivity with zero. All significance values were computed using Holm-Bonferroni correction.

Characterizing how error sensitivity is influenced by the temporal distance between trials and the spatial distance between targets provides insights into the nature of motor memory and adaptation across different conditions. Spatial generalization is a phenomenon of transfer adaptation across nearby movements and more specifically, across nearby reaching targets. To conduct spatial distance analysis, we grouped trials by the distance between the current and the previous trial target. Error sensitivity decreases with spatial distance both in stable and random environment (in stable: rho = -0.77+/-0.28, p-value = 1.15e-10; in random: rho = -0.77+/-0.36, p-value = 6.08e-09), indicating that errors made while reaching for spatially distant targets influence less the learning than errors associated with proximal targets. (Note that for the spatial distance, the average error sensitivity within distances are computed using different numbers of trials because targets were uniformly distributed and fewer trials have larger target jumps between consecutive trials).

Temporal generalization assessed sensitivity not only to the most recent error but also to preceding errors. To conduct temporal generalization analysis, we can define an extension to formula (1): instead of dividing by the most recent error we can divide the error correction by the k-preceding error in the following way

$$ES_k(n) = \frac{\left(RA_{n-1} - TA_{n-1}\right) - \alpha \cdot \left(RA_n - TA_n\right)}{e_{n-k}}. \tag{2}$$

Interestingly, these results reveal that generalization of error sensitivity occurs even in the random environment, albeit at lower magnitudes compared to the stable environment: Spearman correlation of error sensitivity versus spatial distance is notably negative (mean r = -0.71, std = 0.44, p-value = 3.46e-07), as is the correlation of error sensitivity versus temporal distance (mean r = -0.48, std = 0.30, p-value = 4.25e-07).

This quantity shows how correction on trial n is modulated by error history of depth k. There are two ways one can analyze the quantity (2): either buy looking at all trials or buy looking at the subset of trials for which the target at trial n-k was the same as during trial k. We report below the latter but the qualitative results remain the same for the former (except for the error sensitivity at the temporal distance k=2 in the random condition that is not statistically different from zero in that case). We observed a continuous decline in error sensitivity with increasing temporal distance both in stable and random environment (in stable: rho = -0.41 +/- 0.4, p-value = 9.43e-05; in random: mean rho = -0.50+/-0.29, p-value = 1.21e-07), indicating that errors occurring in earlier trials contribute less to learning than more recent errors, regardless of the target's consistency.

Comparative analysis between environments (comparison of Spearman coefficient within participant within condition) did not find significant difference between error sensitivity decay slopes for conditions neither for spatial, neither for temporal distance (p-values = 5.51e-01, 8.32e-01 respectively). This suggests a consistent generalization decay pattern across conditions.

Error sensitivity grows when recent errors have consistent signs and drops when they don't

The paper that coined the term "error sensitivity" (Herzfeld et al. 2014) proposed a formula for its trial to trial evolution. We can test it directly by checking how our empirically defined quantity from formula (1) changes between consecutive trials in relation to consistency of consecutive error signs. Formally we define the error sensitivity change with the following formula:

$$ES_{change}(n) = ES(n) - ES(n-1), \tag{3}$$

Fig 5. Changes of error sensitivity as a function of error sign consistency between consecutive trials, averaged within participant. Top line of significance codes corresponds to two-sided comparison against 0. Allo p-values are Holm-Bonferroni corrected.

In this analysis, trials for which either previous or the pre-previous trial was a hit trial were removed (otherwise error sensitivity values are too high and error sensitivity change is not informative). In average, 15% ± 3% of trials per participant were removed (in addition to those above 5*std threshold removed for all analyses in the beginning). In the stable environment, when consecutive errors are inconsistent in sign, there is a decrease in error sensitivity, with a mean change of -1.09 (standard deviation = 0.26), and this change is significantly different from zero (p-value = 6.33e-14). Conversely, for consistent consecutive errors, error sensitivity increases, evidenced by a mean change of 0.29 (standard deviation = 0.12), also significantly different from zero (p-value = 1.07e-09). This corroborates findings of Herzfeld et al (Herzfeld et al. 2014). However, in the random environment, there is no increase of the error sensitivity when consecutive errors are consistent in sign or decrease of error sensitivity when consecutive errors are inconsistent in sign. These results imply that in the unpredictable conditions of the random environment, the consistency of error feedback does not seem to influence error sensitivity adjustments. Nevertheless, learning persists in this unpredictable setting, as indicated by the above-zero average error sensitivity (Fig. 2D)

It is possible that such results depend on motor variability. Maybe error sensitivity should not be expected to change if the inconsistency of error signs happens only due to motor noise. We checked that by repeating the analysis in this paragraph excluding trials whether either previous or pre-previous errors

were smaller than the standard deviation of the error in the no-perturbation stage. Although such trials constitute a large proportion of all trials within participants (mean = 64 %, std = 13 %), after excluding them overall, results remained qualitatively the same (stable inconsistent is negative with p-value=1.13e-11, and stable consistent is positive p-value=1.06e-06, see also Fig S2) .

Error sensitivity relation to statistics of recent errors

In (Tan et al. 2014) authors suggested that one can use a nonlinear transformation of the perceived errors to compute adaptation rate (we don't call it "error sensitivity" because they did not explicitly compare their values with outputs of formula 1). Specifically in (Tan et al. 2014) the history of 20 last trials was considered and the following formula was used:

$$AR_{Tan}(n) = c\frac{\mu_{20}^2(n)}{\sigma_{20}^2(n)}, \tag{4}$$

where $\mu_{20}^2(n)$, $\sigma_{20}^2(n)$ are square of the mean and variance of the observed error in the window of last 20 trials preceding (but not including) trial n, and the coefficient c was fitted using a computational model (note that if one does not ask for the precise value of the coefficient c, the proportionality can be checked without using a computational model). The authors of (Tan et al. 2014) also considered other history lengths and other powers for μ and σ but they were deemed less appropriate.

We tested whether a similar quantity is related to adaptation rate in our data. We compared error sensitivity with the AR_{Tan}. We employed mixed linear models to systematically test whether different statistics of previous errors relate to error sensitivity across the four conditions (zero perturbation, perturbation, washout, random). Our mixed linear models include these four conditions without interactions but with random effects. The error statistics types that we investigated were
- the standard deviation of previous errors,
- the square of the mean of previous errors,
- the inverse standard deviation of previous errors,
- the ratio of mean to standard deviation,
- the ratio of mean to variance and
- the ratio of mean squared to variance (akin to formula 2),

each time for history lengths between 4 and 24 trials. When fitting a linear mixed model for each of the error statistics, we removed trials where the value of the windows statistic was higher than 5 times its standard deviation. It led to removal of less than 3% of the trials, except for std and square of the mean with histories longer than 20 trials, where it went up to 7%.

We selected models that had a significant positive slope and we excluded models that converged but had the slope less than 1e-3, judging the effect too weak. Only models utilizing the *inverse of the standard deviation* with history lengths between 8 and 20 trials remained after this filtering. The largest slope = 0.032 (p-value = 9.18e-06) was for history of length 16. For this statistic the intercept at baseline condition (stable perturbation) was not statistically different from zero. Differences between the intercepts during washout and during no perturbation conditions were significantly positive (coefficients 0.75 and 0.31 with p-values 9.25e-26 and 1.19e-10)

Overall this analysis suggests that the exact quantity suggested by (Tan et al. 2014) does not seem to influence error sensitivity in our data. However, the inverse of the standard deviation within the 8-20 previous trial window shows a significant relationship with error sensitivity, although the effect size (slope in the linear mixed model) is relatively small. In other words, our data does only weakly supports the hypothesis that error sensitivity is determined by the previous error statistics (Tan et al. 2014). Incorporating interactions with experimental conditions in our model does not alter the overall outcome. (See Supplementary).

Discussion

Summary of results:

To investigate the dynamics of error sensitivity during a motor adaptation task, we measured error sensitivity during several conditions, including stable and random environments. We report four main findings. First, error sensitivity is higher during no perturbation than during perturbation, and it grows within a stable environment. Second, error sensitivity is still above zero even in a random environment. It means that participants continue to correct their errors even when nothing can be learned from them. Third, there is spatial and temporal generalization both in stable and random environments. Fourth, neither the error consistency (Herzfeld et al. 2014) nor windows error statistics-based (Tan et al. 2014) models for error sensitivity dynamics do not fully agree with our data.

Dynamics of error sensitivity in stable versus random environments

Error sensitivity during no perturbation conditions is significantly higher than during perturbation and washout conditions (Fig. 2D). This increase, visible at the trial level each time the perturbation disappears (Fig. 2B), could be explained because participants are more eager to compensate for their own motor noise (motor variability present physiologically in the absence of any perturbation), maybe because they are more familiar with it, than with an external perturbation. The error sensitivity also grows within stable perturbation, which one could explain in two ways: either i) towards the end of the perturbation the internal model adaptation stabilizes and then the same explanation as for no perturbation explains higher error sensitivity or ii) the perturbation onset induces a drop in error sensitivity and then the error sensitivity slowly increases until the next change in stability. The latter is closer to the hypothesis in (Herzfeld et al. 2014).

The washout phase, marked by the removal of perturbations, seems to facilitate a recovery or enhancement in error sensitivity, with a significant positive correlation between trial number and error sensitivity. In contrast, during random conditions, error sensitivity gradually decreases over time, although this decrease is not statistically significant. Such a negative trend suggests that unpredictable conditions may hinder the adaptive process.

Lower error sensitivity in random environment agrees with previous results in (Gonzalez Castro et al. 2014). Note that in general variability in movement outcomes can be both extrinsic (e.g. random condition in our experiment) and intrinsic (e.g. due to physiologic tremor). Our study focuses on the effect of the former one. The effect of intrinsic motor variability on learning rate is unclear (He et al. 2016).

Memory of Error Model

In stable condition our results agree with the idea that error consistency sign modulates error sensitivity, offered as a part of "Memory of Error" model in (Herzfeld et al. 2014). However, the lack of error consistency dependence in the random condition (Fig 5) challenges the Memory of Error model (Albert et al. 2021; Herzfeld et al. 2014). A possible explanation for these differences is the presence of multiple targets and partial generalization in our study (Fig. 4). In contrast, the Memory of Error model does not account for target location and therefore assumes perfect generalization across targets, regardless of their spatial distance. Another possible explanation for these differences is the use of a joystick as controller in our study, which may potentially involve different adaptation mechanisms compared to the robotic arm used in (Herzfeld et al. 2014).

Implicit vs explicit learning

Motor adaptation involves two key processes: implicit learning and explicit strategy application. Implicit learning refers to the unconscious adjustment to a perturbation, aiming to partially compensate for it. Explicit strategy application, on the other hand, involves the conscious use of strategies such as re-aiming (Tsay et al. 2023).

Disentangling these two learning processes is challenging and requires careful experimental design (Tsay et al. 2023) Even with such efforts, the results can remain ambiguous (Maresch, Mudrik, and Donchin 2021). However, due to a large body of literature on the topic, some theoretical understanding starts to crystallize as well and certain learning phenomena are considered to be a feature of one process but not another.

Which mechanism is more prevalent in our setup is not exactly clear, but there are more reasons for it to be predominantly implicit, even though endpoint-only feedback often involves explicit learning (Tsay et al. 2023). The absence of savings (see next section) suggests lower involvement of explicit learning. One might argue that explicit learning has been partially disrupted by the random period. However, we observe evidence of spatial generalization during both the stable and random periods, and we find no differences in the shapes of the generalization curves. Given the unlikely role of explicit strategy in a random environment, the similarity in the shapes of the generalization curves leads us to hypothesize that implicit learning predominates in both the random and stable environments. Finally in general, a localized (unimodal, decaying to zero for large errors) spatial generalization profile (Fig S3, leftmost panel) is a feature of implicit learning (Tsay et al. 2023).

Note that spatial generalization in the context of environment stability was studied earlier (Fernandes, Stevenson, and Kording 2012) and was found to be similar in both high and low variance perturbation conditions. However the perturbation schedule there was different than in our experiment: first only one target was presented and then others. This way the possibility of forgetting what was learned for the training direction was not addressed.

Besides that in (Avraham et al. 2020) it was already shown that error sensitivity is lower in case of low environmental consistency and the authors provided evidence that modulation was due to explicit learning. However a more recent study (Albert et al. 2021) challenged it showing that this effect is rather due to implicit learning and suggested that results in (Avraham et al. 2020) can be attributed to an unusual feature in experimental design: relatively long (600ms+) delay between the movement termination and endpoint feedback presentation.

Consistent with (Albert et al. 2021) we find that lower error consistency in the random environment does not reduce error sensitivity but rather suppresses its growth (Fig 2C, Fig 3). Dependence of error sensitivity on previous error has a familiar shape (Fig S3, first column) reported

previously in (Herzfeld et al. 2014). We have not found a statistically significant difference between generalization curves in random and stable environments (Fig S3, first column).

Savings

In our experiment the participants perceived the same perturbation twice: the first and last stable perturbation values are equal and also the second and third constant perturbation values are equal. In such situations, a phenomenon called "savings" may be observed (Smith, Ghazizadeh, and Shadmehr 2006): when re-experiencing a perturbation, adaptation could be faster. We did not find savings in our experiment. Error sensitivities were not statistically different between the first and last perturbation (two sample t-test, t=-1.90, p=1.47e-01, Bonferroni corrected) nor between the second and the third perturbation (t=1.33, p=4.01e-01, Bonferroni corrected). We discuss below why it could be the case.

Although still debated (Tsay et al. 2023) there is a hypothesis that savings occur exclusively in explicit strategy learning. In our data, however, no evidence of savings was observed, despite being expected. Error sensitivity values during the first and last perturbations were not significantly different (t = -1.90, p = 0.0733), nor were those during the second and third perturbations (t = 1.33, p = 0.200). This suggests that implicit learning was not the primary mechanism used by our participants.

There is evidence that savings can survive adaptation in the opposite direction (Malone et al. 2011; Sarwary, Selen, and Medendorp 2013). However, this evidence comes from the gait adaptation, where savings might be supported by different mechanisms. Moreover, the 2nd and 3rd perturbations in our experiment, which were not separated by an opposite consistent perturbation, still showed no evidence of savings in our analysis.

Savings suppression after a long washout of a special kind was observed earlier in (Kitago et al. 2013; Kojima, Iwamoto, and Yoshida 2004). In particular, (Kitago et al. 2013) report that long series of error clamp trials can suppress savings, and in our experiment the random environment can be considered a sort of random clamp environment, just that clamp direction changes on every trial. In (Albert et al. 2021) authors suggest a mechanism for savings suppression: error sensitivity update weight decay in Memory of Error model. Potentially, it could apply for our experiment as well because although the distribution of errors during random periods equaled the distribution of errors during the precedent stable period, the error consistency pattern was destroyed by trial permutation. However, since the Memory of Error model does not fully align with our data (see corresponding section above), it is unlikely to fully explain the lack of savings.

In addition to the random perturbations, there were three pauses lasting approximately one minute each. However, their influence on savings is likely less significant than that of the random period, which lasted nearly 15 minutes. The average pause duration was 44.7 seconds (SD = 13.4 seconds).

Another important consideration is connected to the results of (Hadjiosif, Morehead, and Smith 2023). The authors of that paper showed that motor adaptation can be split into a temporally persistent and temporary volatile component, the latter being defined as the learning that decays quickly (within 15-25 seconds). Somewhat counterintuitively it is the volatile component that they show to be responsible for savings (the persistent one is responsible for long term retention). In our experiment, given that the targets appear in a pseudorandom order, in most cases the same target does not appear twice in a row. This creates a delay of multiple trials between the same target trials with mean time difference = 14.9 seconds (mean trail distance = 4 trials). While this number is slightly lower than 15-25 seconds mentioned in (Hadjiosif et al. 2023; Shadmehr Lab 2024), since the volatile adaptation decay is exponential and we have a relatively high motor noise in our experiment, we expect that this effect contributes to the absence of savings. The decay pattern of temporal generalization we observe in Fig 2B,D supports this idea (recall that we computed the quantity (2) looking at the subset of trials for which the target at trial n-k was the same as during trial k).

We would like to underline that this study was not designed to study savings specifically and further behavioral and modeling research needed to shed more light on the absence of savings in our setup.

Trial dependence of Error Sensitivity

Trial dependence of error sensitivity is a matter of debate. For example in (Heald, Lengyel, and Wolpert 2021) authors proposed COIN model that gave an alternative explanation for results of (Herzfeld et al. 2014), based on context sequence switches.

In (Wang et al. 2023) authors targeted the question of necessity of trial dependence of error sensitivity specifically and offered a model alternative to Memory of Error that is simpler and is able to explain their data as well as (Albert et al. 2021) data. However it is not consistent with our data because in our experiment error distributions between stable and random environments are designed to coincide perfectly and yet error sensitivity distributions between them are very different.

Continuous monotonous increase of error sensitivity within a stable environment was reported earlier in Experiment 6 of (Albert et al. 2021) using a different experimental controller and implicit learning-only design (achieved by reducing reacting time) .

Controller type

The comparison of different motor adaptation studies should always be exercised with care, as various factors can influence the results. These include the type of controller used to guide the cursor (e.g., manipulandum, mouse, joystick, tablet, hand tracking), the type of perturbations (e.g., visuomotor rotations, visuomotor mirror reversal, visuomotor scaling, force field), the perturbation schedule (e.g., gradual, abrupt, random every N-th trial), the feedback presentation (e.g., online, online with delay, endpoint-only), and task timing (e.g., the presence of motor preparation and limits on the speed of the movement). Compared to manipulandums and hand tracking, joysticks are less commonly used as controllers in research. Our motivation for using the joystick and visuomotor rotation was its compatibility with non-invasive neural recordings.

Moreover, spatial generalization patterns seem to strongly depend on the controller type, probably due to different relationships between proprioception-informed error and visual error and active vs passive behavior of the apparatus itself (Fleury, Prablanc, and Priot 2019). However the amount of decrease of error sensitivity with increasing target distance can strongly depend on the training type, in particular whether training was focused (involved similar movements) and with visual feedback, or not (see Discussion in (Berniker et al. 2014)).

Future directions

Recently in (Sugiyama, Schweighofer, and Izawa 2023) another model was suggested that connects adaptation rate with reward. Their reward was given explicitly to modulate adaptation rate. It would be interesting to try to adapt that approach to the case when no explicit reward is given and participants self-reinforce themselves somehow. Modeling such rewards however is a rather intricate task (Shmuelof et al. 2012; Todorov et al. 2019).

Funding

The intramural National Institute of Neurological Disorders and Stroke, Division of Intramural Research, NIH, as well as the National Institute of Mental Health Magnetoencephalography (MEG) Core Facility and Functional Magnetic Resonance Facility facilities on the NIH campus (Bethesda, MD) contributed to this research. This project received funding from the ATIP-AVENIR program. D. Todorov received support from ANR-23-CPJ1-0137-01. We gratefully acknowledge the computing time granted by EBRAINS and provided on the JUSUF system at Jülich Supercomputing Centre (JSC), project icei-hpb-2022–0017.

Data Availability Statement

Data (subject to participant consent) used to generate the findings of this study are freely available at https://osf.io. Custom code generated during this study is available at github.com.

Acknowledgements

We thank D. Herzfeld, O. Abdoun and G. Marrelec for their valuable advice and the Eduwell team at the Lyon Neuroscience Research Center (CRNL) for general support.

References

Albert, Scott T., Jihoon Jang, Hannah R. Sheahan, Lonneke Teunissen, Koenraad Vandevoorde, David J. Herzfeld, and Reza Shadmehr. 2021. "An Implicit Memory of Errors Limits Human Sensorimotor Adaptation." *Nature Human Behaviour* 5(7):920–34. doi:10.1038/s41562-020-01036-x.

Avraham, Guy, Matan Keizman, and Lior Shmuelof. 2020. "Environmental Consistency Modulation of Error Sensitivity during Motor Adaptation Is Explicitly Controlled." *Journal of Neurophysiology* 123(1):57–69. doi:10.1152/jn.00080.2019.

Berniker, Max, David W. Franklin, J. Randall Flanagan, Daniel M. Wolpert, and Konrad Kording. 2014. "Motor Learning of Novel Dynamics Is Not Represented in a Single Global Coordinate System: Evaluation of Mixed Coordinate Representations and Local Learning." *Journal of Neurophysiology* 111(6):1165–82. doi:10.1152/jn.00493.2013.

Brayanov, Jordan B., Daniel Z. Press, and Maurice A. Smith. 2012. "Motor Memory Is Encoded as a Gain-Field Combination of Intrinsic and Extrinsic Action Representations." *Journal of Neuroscience* 32(43):14951–65. doi:10.1523/JNEUROSCI.1928-12.2012.

Donchin, Opher, Joseph T. Francis, and Reza Shadmehr. 2003. "Quantifying Generalization from Trial-by-Trial Behavior of Adaptive Systems That Learn with Basis Functions: Theory and Experiments in Human Motor Control." *Journal of Neuroscience* 23(27):9032–45. doi:10.1523/JNEUROSCI.23-27-09032.2003.

Fernandes, Hugo L., Ian H. Stevenson, and Konrad P. Kording. 2012. "Generalization of Stochastic Visuomotor Rotations." *PloS One* 7(8):e43016. doi:10.1371/journal.pone.0043016.

Fleury, Lisa, Claude Prablanc, and Anne-Emmanuelle Priot. 2019. "Do Prism and Other Adaptation Paradigms Really Measure the Same Processes?" *Cortex* 119:480–96. doi:10.1016/j.cortex.2019.07.012.

Gonzalez Castro, Luis Nicolas, Alkis M. Hadjiosif, Matthew A. Hemphill, and Maurice A. Smith. 2014. "Environmental Consistency Determines the Rate of Motor Adaptation." *Current Biology* 24(10):1050–61. doi:10/f54fbf.

Hadjiosif, Alkis M., J. Ryan Morehead, and Maurice A. Smith. 2023. "A Double Dissociation between Savings and Long-Term Memory in Motor Learning." *PLOS Biology* 21(4):e3001799. doi:10.1371/journal.pbio.3001799.

He, Kang, You Liang, Farnaz Abdollahi, Moria Fisher Bittmann, Konrad Kording, and Kunlin Wei. 2016. "The Statistical Determinants of the Speed of Motor Learning." *PLOS Computational Biology* 12(9):e1005023. doi:10.1371/journal.pcbi.1005023.

Heald, James B., Máté Lengyel, and Daniel M. Wolpert. 2021. "Contextual Inference Underlies the Learning of Sensorimotor Repertoires." *Nature* 600(7889):489–93. doi:10.1038/s41586-021-04129-3.

Herzfeld, David J., Pavan A. Vaswani, Mollie K. Marko, and Reza Shadmehr. 2014. "A Memory of Errors in Sensorimotor Learning." *Science* 345(6202):1349–53. doi:10.1126/science.1253138.

Jordan, Michael I., and David E. Rumelhart. 1995. "Forward Models: Supervised Learning with a Distal Teacher." in *Backpropagation*. Psychology Press.

Kawato, M., Kazunori Furukawa, and R. Suzuki. 1987. "A Hierarchical Neural-Network Model for Control and Learning of Voluntary Movement." *Biological Cybernetics* 57(3):169–85. doi:10.1007/BF00364149.

Kitago, Tomoko, Sophia Ryan, Pietro Mazzoni, John Krakauer, and Adrian Haith. 2013. "Unlearning versus Savings in Visuomotor Adaptation: Comparing Effects of Washout, Passage of Time, and Removal of Errors on Motor Memory." *Frontiers in Human Neuroscience* 7.

Kojima, Yoshiko, Yoshiki Iwamoto, and Kaoru Yoshida. 2004. "Memory of Learning Facilitates Saccadic Adaptation in the Monkey." *Journal of Neuroscience* 24(34):7531–39. doi:10.1523/JNEUROSCI.1741-04.2004.

Krakauer, John W. 2006. "Motor Learning: Its Relevance to Stroke Recovery and Neurorehabilitation." *Current Opinion in Neurology* 19(1):84–90. doi:10.1097/01.wco.0000200544.29915.cc.

Krakauer, John W., and S. Thomas Carmichael. 2022. *Broken Movement: The Neurobiology of Motor Recovery after Stroke*. MIT Press.

Krakauer, John W., Pietro Mazzoni, Ali Ghazizadeh, Roshni Ravindran, and Reza Shadmehr. 2006. "Generalization of Motor Learning Depends on the History of Prior Action." *PLOS Biology* 4(10):e316. doi:10.1371/journal.pbio.0040316.

Malone, Laura A., Erin V. L. Vasudevan, and Amy J. Bastian. 2011. "Motor Adaptation Training for Faster Relearning." *Journal of Neuroscience* 31(42):15136–43. doi:10.1523/JNEUROSCI.1367-11.2011.

Maresch, Jana, Liad Mudrik, and Opher Donchin. 2021. "Measures of Explicit and Implicit in Motor Learning: What We Know and What We Don't." *Neuroscience & Biobehavioral Reviews* 128:558–68. doi:10.1016/j.neubiorev.2021.06.037.

Morehead, J. Ryan, and Jean-Jacques Orban de Xivry. 2021. *A Synthesis of the Many Errors and Learning Processes of Visuomotor Adaptation*. *preprint*. Neuroscience. doi:10.1101/2021.03.14.435278.

Sarwary, A. M. E., L. P. J. Selen, and W. P. Medendorp. 2013. "Vestibular Benefits to Task Savings in Motor Adaptation." *Journal of Neurophysiology* 110(6):1269–77. doi:10.1152/jn.00914.2012.

Shadmehr Lab, dir. 2024. *Maurice Smith: Cerebellum and Human Motor Memory*.

Shmuelof, Lior, Vincent S. Huang, Adrian M. Haith, Raymond J. Delnicki, Pietro Mazzoni, and John W. Krakauer. 2012. "Overcoming Motor 'Forgetting' through Reinforcement of Learned Actions." *Journal of Neuroscience* 32(42):14617–21. doi:10.1523/jneurosci.2184-12.2012.

Smith, Maurice A., Ali Ghazizadeh, and Reza Shadmehr. 2006. "Interacting Adaptive Processes with Different Timescales Underlie Short-Term Motor Learning" edited by J. Ashe. *PLoS Biology* 4(6):e179. doi:10.1371/journal.pbio.0040179.

Sugiyama, Taisei, Nicolas Schweighofer, and Jun Izawa. 2023. "Reinforcement Learning Establishes a Minimal Metacognitive Process to Monitor and Control Motor Learning Performance." *Nature Communications* 14(1):3988. doi:10.1038/s41467-023-39536-9.

Tan, Huiling, Ned Jenkinson, and Peter Brown. 2014. "Dynamic Neural Correlates of Motor Error Monitoring and Adaptation during Trial-to-Trial Learning." *Journal of Neuroscience* 34(16):5678–88. doi:10.1523/JNEUROSCI.4739-13.2014.

Tanaka, Hirokazu, and Terrence J. Sejnowski. 2014. "Motor Adaptation and Generalization of Reaching Movements Using Motor Primitives Based on Spatial Coordinates." *Journal of Neurophysiology* 113(4):1217–33. doi:10/f62679.

Thoroughman, Kurt A., and Reza Shadmehr. 2000. "Learning of Action through Adaptive Combination of Motor Primitives." *Nature* 407(6805):742–47. doi:10.1038/35037588.

Todorov, Dmitrii I., Robert A. Capps, William H. Barnett, Elizaveta M. Latash, Taegyo Kim, Khaldoun C. Hamade, Sergey N. Markin, Ilya A. Rybak, and Yaroslav I. Molkov. 2019. "The Interplay between Cerebellum and Basal Ganglia in Motor Adaptation: A Modeling Study." *Plos One* 14(4):e0214926.

Tsay, Jonathan, Richard Ivry, John W. Krakauer, Adrian M. Haith, Guy Avraham, and Hyosub E. Kim. 2023. "Strategy Processes in Sensorimotor Learning: Reasoning, Refinement, and Retrieval."

Wang, Tianhe, Guy Avraham, Jonathan S. Tsay, Sabrina J. Abram, and Richard B. Ivry. 2023. "Perturbation Variability Does Not Influence Implicit Sensorimotor Adaptation." 2023.01.27.525949.