Investigating milk derived extracellular vesicles as mediators of maternal stress and environmental intervention

Running title: Extracellular vesicles and maternal stress

Julia Martz[1&], Baila Hammer[2&], Tristen J. Langen[1], Benjamin Berkowitz[2], Benzion Berkowitz[2], Jasmyne A. Storm[3], Jueqin Lu[3], Deepali Lehri[4], Sanoji Wijenayake[3], Jordan Marrocco,[2,5,6]*, Amanda C. Kentner[1]*

[1]School of Arts & Sciences, Health Psychology Program, Massachusetts College of Pharmacy
and Health Sciences, 179 Longwood Avenue, Boston Massachusetts, United States 02115.
[2]Department of Biology, Touro University, 3 Times Sq, New York, New York, United States 10036.
[3]Department of Biology, The University of Winnipeg, 515 Portage Ave, Winnipeg, Manitoba, Canada.
[4]Genomics Core Laboratory, New York Medical College, 15 Dana Road, Valhalla, New York, United States 10595.
[5]Department of Physiology, New York Medical College, 15 Dana Road, Valhalla, New York, United States 10595.
[6]Feil Family Brain and Mind Research Institute, Weill Cornell Medicine, 1300 York Ave, New York, New York, United States, 10065.

[&]Shared first authors

*Corresponding authors (all email correspondence should include both authors):

Jordan Marrocco
jmarrocc@touro.edu
Office #646-981-4708

Amanda Kentner
amanda.kentner@mcphs.edu
Office #617-274-3360
Fax # 617-732-2959

Total Word Count: 4970
Abstract: 244
Introduction: 808 words
Figures: 6
Tables: 0
Supplementary Materials:
1. Supplemental Methods and Experimental Overview
2. Supplemental Figure 1
3. Supplemental Table 1 – *MIA Reporting Guidelines*
4. Supplemental Table 2 – *overlap of P12 and P74 miRNA-mRNA interactions*
5. Supplemental Table 3 – *statistical results*

Abstract

Parental communication signals are transmitted through nursing and critically shape neurodevelopmental trajectories. Mirroring some well characterized effects of gestational challenges in rodents, maternal immune activation (MIA) during the lactational period disrupts maternal physiology, decreases milk quality, and is associated with adverse neurobehavioral outcomes in offspring. This occurs without MIA significantly affecting maternal care. While gestational MIA models are responsive to environmental interventions, which beneficially alter maternal milk composition and associated offspring outcomes, the milk-borne mediators underlying resilience remain poorly understood. Given their ability to transport and deposit biologically active cargos, we propose that milk-derived extracellular vesicles (MEVs) are vehicles that deliver environmental programming signals (e.g., miRNAs) from nursing mothers to their offspring. Using a rat model, we show that lactational MIA altered MEV miRNA cargo and expression of hippocampal miRNAs in offspring. Several miRNAs in MEVs were also found in the hippocampus of matching offspring. Remarkably, the miRNA dysregulation observed in MEVs and hippocampus was rescued when dams were raised in an enriched environment, suggesting environmental enrichment protected from the effects of MIA, as also observed in the behavioral phenotype. RNA-seq of adult offspring hippocampus showed long-term transcriptional changes associated with the gene targets of early-life regulated miRNAs. Our results position MEV miRNA cargos as dynamic programming signals by which maternal experience is communicated to offspring, encoding both stress-induced and protective cues that influence development. This suggests that breastfeeding interventions can regulate the genetic cargo of the milk, programming the life of developing infants.

Introduction

Intergenerational transmission of biological cues shape offspring neurodevelopment and long-term behavioral outcomes. Beyond direct caregiving and nutritional transfer through nursing, recent research has identified extracellular vesicles (EVs) as a mechanism through which parental signals may be transmitted to offspring[1-7]. EVs are membrane-bound nanovesicles containing microRNAs (miRNAs), messenger RNAs (mRNAs), proteins, peptides, and lipids, capable of influencing gene expression and cellular function in target tissues[8,9]. While the mechanisms underlying parental contributions to offspring development via reproductive tract-derived EVs and sperm are well described[2,3], maternal EV signaling during lactation remains less understood.

Breast milk is a biologically active medium; it supplies nutrition and other regulatory information to offspring[10]. Indeed, breastfeeding is associated with numerous benefits for infant development, including enhanced immune function, cognitive outcomes, and reduced risk for metabolic diseases[10-13]. Milk contains a substantial population of milk-derived extracellular vesicles (MEVs) that can survive the digestive tract, enter systemic circulation, and reportedly accumulate in peripheral tissues and the brain, including the hippocampus of nursing neonatal mice[5,7,14-19]. Complementary evidence from a blood brain barrier (BBB) cell-based model demonstrates that peripherally derived EVs transport their cargos across brain microvascular endothelial cells. This is most apparent under inflammatory conditions when barrier integrity is reduced[20]. Moreover, although the BBB is established in neonates, some brain associated barriers are more permeable to small molecules during this developmental stage[21-23], a time when the brain shows increased bioavailability to peripherally administered EVs compared to adults[24]. These findings highlight that early life contexts characterized by reduced BBB permeability, whether developmental or inflammatory driven, may facilitate brain exposure to MEVs, providing a pathway for maternal signals to shape neurodevelopment.

EVs are increasingly recognized as important mediators of intercellular communication within the central nervous system (CNS). For example, EVs derived from both the periphery and CNS (e.g., neurons, astrocytes) can interact with and regulate microglia function[10,25-26]. A recent *in vitro* study demonstrated that MEVs are readily taken up by homeostatic and polarized human microglia, influencing the abundance and enzymatic activity of a critical epigenetic modulator, the DNA methyltransferase 1[27]. The study also illustrated a mechanistic association between DNMT1 regulation and MEV-miRNA 148a-5P.

miRNA are small noncoding RNAs that regulate post-transcriptional gene expression and affect neural development, primarily by acting as repressors of target messenger RNA[28,29]. Specific miRNA trafficked by MEVs are known to regulate synaptic development [see 18-19,30] and inflammatory responses, alongside other physiological functions[31]. Additionally, MEVs have beneficial effects in the gut through interactions with epithelial cells[16,32] and may influence offspring health and development by directly influencing the microbiome via their miRNA cargos[7,14,33]. These findings suggest that MEV's have multiple points of contact with CNS target sites by which they can modify offspring neurodevelopment through postnatal transcriptional signaling.

Maternal infections during pregnancy are relatively common[34] and have been linked to an increased risk of neurodevelopmental disorders in offspring[35]. To investigate the mechanisms underlying this association, animal models of maternal immune activation (MIA), where maternal immune stimulation during gestation induces neurodevelopmental impairments in offspring, are widely used and validated[36-37]. While these gestational MIA models are supported by epidemiological evidence and offer strong translational relevance, comparatively little is known about the prevalence or risks of infections during the lactational

period[38]. Recently, animal models of MIA during lactation have been evaluated[39-41], demonstrating that maternal immune challenges during the nursing period impact offspring neurodevelopmental outcomes similarly to the gestational models. Importantly, these behavioral effects occur despite minimal alterations in maternal care behaviors[40-43]. This suggests that other variables, like milk-borne factors, may mediate these outcomes. While stressors experienced across lactation can influence maternal physiology and milk composition[10,44], comprehensive analyses of traditional milk factors such as nutritional content, immunoglobulins, inflammatory cytokines, and even the maternal immunogen used to stimulate lactational MIA have not fully explained the effects on offspring neurodevelopment[41]. Therefore, alternative mechanisms must be considered.

Given the capacity of MEVs to deliver regulatory molecules to recipient cells and impart biological effects[7,14-15,18], and their demonstrated involvement in the CNS[27], MEVs are a plausible candidate to mediate the effects of lactational MIA. Indeed, environmental experiences such as stress and illness modify the miRNA cargo of MEVs[6,45-47], supporting the hypothesis that MEVs are sensitive to maternal environmental conditions and may relay this information to nursing offspring. Based on these findings, we hypothesize that lactational MIA alters the cargo of MEVs, leading to changes in the transcriptomic profile of the developing brain. These molecular changes may drive long-term alterations in gene expression, contributing to behavioral abnormalities observed in adulthood. Furthermore, because environmental enrichment (EE) has been shown to mitigate the behavioral consequences of gestational MIA[48-50] and to improve milk quality in healthy laboratory rats[51], we propose that an EE intervention may stabilize MEV cargos, protecting against the effects of lactational MIA. Understanding these mechanisms will provide insight into how the maternal environment shapes offspring neurodevelopment via milk-derived signaling pathways and may suggest strategies to enhance resilience in at-risk populations.

Methods and Experimental Overview

All animal protocols were approved by the MCPHS Institutional Animal Care and Use Committee and complied with AAALAC guidelines. The experimental timeline is illustrated in **Figure 1A**. Complete methodological details are provided in **Supplementary Methods**, **Supplementary Table 1**[37], and DeRosa et al.[41,51].

Animals and Housing: Sprague Dawley rats (Charles River, Wilmington) were bred and housed in either environmentally enriched (EE) or standard (SD) laboratory conditions as previously described[48,51]. On postnatal day (P)10, lactating dams were administered either 100 µg/kg lipopolysaccharide (LPS; *Escherichia coli* O26:B6) or pyrogen-free saline via intraperitoneal injection (n = 7-8 litters/group). Dams were separated from their litters to allow milk to accumulate for 2 hours post-injection while litters were kept warm on a heating pad[41,51]. Based on previous work, this period of separation would not induce a strong corticosterone response in neonates[52].

Maternal Assessments: Sickness behavior was evaluated at baseline, 30-, 120-, and 300-minutes post-injection using a composite score of ptosis, piloerection, and lethargy (0 = none, 1 = mild, 2 = severe), by blinded investigators, as adapted from prior studies[49,53-54]. Maternal care was observed on P9–P11 during AM and PM sessions. Behavioral metrics included pup licking/grooming frequency and time spent on the nest across six 1-minute intervals[41,51].

Milk Collection and Analysis: Two hours after injection, dams were anesthetized with isoflurane and administered oxytocin (0.2 mL; 20 USP/mL, i.p.) to induce milk letdown[41,51]; based on their pharmacokinetic profiles, these drugs are not absorbed by offspring[55-57]. Approximately 2.0 mL of milk was manually expressed and collected over 50–60 minutes.

Dams were returned to their litters once recovered from anesthesia. Milk was collected again 24 hours later on P11. Creamatocrit was determined following procedures previously outlined (n = 7-8 litters/group)[41,51,58], and 100 µL of milk was stored at –80 °C for corticosterone assays (n = 7 litters/group). Samples were thawed and rotated overnight, then analyzed using a small sample ELISA protocol ((#ADI-900–097, Enzo Life Sciences; 1:40 dilution)[41,51]. Values below detection thresholds, commonly observed in saline controls, were imputed as zero for analysis[59].

Milk-Derived Extracellular Vesicles (MEVs): Following centrifugation to remove the cream layer, MEVs were isolated from the supernatant using differential ultracentrifugation and filtration (DUC), as previously outlined[2-3,27,60] (**Figure 2A)** and compliant with MISEV 2023 guidelines[61]. Casein was precipitated[62], and whey fractions were ultracentrifuged to pellet MEVs, which were washed and resuspended in PBS. MEVs were characterized per MISEV 2023 guidelines using nanoparticle tracking analysis (NTA), western blotting, and transmission electron microscopy (TEM). NTA determined MEV size and concentration (**Figure 2B**). Western immunoblotting confirmed the presence of markers CD-9, Alix, Flotillin-1, and the absence of Calnexin (**Figure 2C**); these protein markers were assessed qualitatively to verify MEV identity[27,60-61]. TEM verified MEV morphology (**Figure 2D**).

Offspring Assessments: The offspring of saline and MIA dams were evaluated on neonatal huddling behavior (a proxy for neonatal thermoregulation; n = 7-8 litters/group), two hours after reuniting with their dams on P10 and P11[41,63]. Litters were weaned into same-sex pairs on P21, maintaining their original housing assignments. Starting on P70, one male and one female offspring from each litter were assessed on the open field and social preference tests (n = 7-8 litters/group/sex)[41,48, 64]. One male and one female from each litter was euthanized with isoflurane, and whole (at P12) or ventral (at P74[65]; hippocampus dissected, frozen on dry ice and stored at –80°C.

RNA Extraction and Sequencing: Total RNA was extracted from hippocampal tissues using the RNeasy Lipid Tissue Mini Kit (74804, Qiagen) and from MEV samples using the Exosomal RNA Isolation Kit (58000, Norogen Biotek), according to the manufacturer's instructions. Samples were quantified using Qubit 2.0 Fluorometer (ThermoFisher Scientific) and RNA integrity was checked with 4200 TapeStation (Agilent Technologies).

MEV small RNA-seq (n = 6 litters/group): Libraries were prepared using NEB Small RNA Library Prep Kit (New England Biolabs), validated on TapeStation, quantified using Qubit 2.0 Fluorometer, and sequenced on Illumina NovaSeq (2x150bp PE), and de-multiplexed using Illumina's bcl2fastq 2.20 software.

Neonatal hippocampus small RNA-seq (n = 6 litters/group): Libraries were prepared using Illumina TruSeq Small RNA Library Prep Kit, cDNA constructs purified via BluePippin, validated on TapeStation, quantified using Qubit 2.0 Fluorometer, and sequenced on Illumina NextSeq 2000 (1x50bp).

Adult hippocampus RNA-seq (n = 6 litters/group): Strand-specific libraries were generated using NEBNext Ultra II Directional RNA Library Prep Kit (Illumina), validated on TapeStation, quantified using Qubit 2.0 Fluorometer, and sequenced on Illumina NovaSeq (2x150bp PE), and de-multiplexed using Illumina's bcl2fastq 2.20 software.

Sequencing Analysis: Differentially expressed miRNAs and genes were identified based on a $p<0.01$, (FC)>2.0. Heatmaps were generated using Multiple Experiment Viewer. Venn diagrams were generated using DeepVenn (https://www.deepvenn.com). Gene ontology was determined using the DAVID functional annotation cluster tool (https://david.ncifcrf.gov/), while miRNA Enrichment Analysis and Annotation Tool (miEAA) was used to functionally annotate the different sets of miRNAs. Bubble plots and pie charts were generated in GraphPad Prism 10.4.2.

Statistical Analyses: Data were analyzed using Prism (GraphPad) or SPSS (IBM). ANOVAs (Housing x MIA x Time or Housing x MIA) were used to evaluate milk and behavioral endpoints, unless there were violations to the assumption of normality (Shapiro-Wilk test) in which case Kruskal-Wallis tests were employed (expressed as X^2). Partial eta-squared (n_p^2) is reported as an index of effect size for the ANOVAs[66]. Both male and female animals were included, and separate analyses were run for each sex[41,51,67,68]. All data are expressed as mean ± SEM.

Results

EE protected against lactational MIA-induced changes in offspring behavior

Maternal immune challenge during lactation significantly increased sickness behaviors in dams, regardless of housing assignment, validating the model. A MIA by time interaction identified an increased severity of sickness behavior in LPS treated dams 5 hours post immune challenge, compared to saline dams (**Figure 1B**). A housing by time interaction indicated that SD dams appeared less healthy than EE dams at baseline, consistent with prior findings[51]. SD dams, confined to smaller cages, nurse more due to a limited ability to escape from pups. SD housing is associated with displays of "pressing" behavior where dams press their ventral side against the cage, seemingly to hide their teats and take a break from the energy-intensive nursing task[51].

Huddling behavior was significantly reduced in SD-LPS neonatal offspring compared to neonates from SD-saline dams (MIA by housing interaction; average number of clusters **Figure 1C**). Diminished neonatal huddling behavior was buffered by EE (SD-LPS versus EE-LPS).

In the open field test, MIA by housing interactions were observed for both adult male and female offspring in terms of distance traveled (**Supplementary Figure 1A, B**). However, follow up tests were not significant in males; female EE-LPS rats traveled less than SD-LPS females. While males were not affected, SD-LPS female offspring spent a reduced percentage of time in the center of an open field compared to SD-Saline females (MIA by housing interaction: **Figure 1D, E**). This MIA induced avoidance behavior was prevented by EE housing. In contrast, male SD-LPS adult offspring had a decreased social preference index compared to SD-Saline males (MIA by housing interaction: **Figure 1F, G**). Complex EE housing also protected against the MIA associated alteration in male social behavior.

Maternal care differences do not appear to account for these effects since there were no significant group differences in terms of total time on nest or the number of pup-directed licking or grooming bouts (**Supplementary Figure 1C, D**). While previous work has shown housing differences in the maternal care of untreated rats[51], stress induced by the LPS/Saline injections may have normalized the amount of maternal care displayed across SD and EE groups in the present study. **Statistics reported in Supplemental Table 3**.

Housing condition regulates lactational MIA-induced changes in MEV miRNA cargos

Building on our previous findings that MIA reduces milk quality[41] while EE improves it[51], we investigated whether EE housing could mitigate MIA-induced changes to milk composition. Percent creamatocrit was decreased in the milk of SD-LPS dams compared to SD-saline, which was mitigated by EE housing (MIA x Housing interaction; main effect of time; **Figure 1H**). However, the body weights of nursing pups were not affected by either MIA or housing across P10-P12 (p>0.05; **Figure 1I**) nor at weaning (p>0.05; **Supplemental Figure 1E**). Milk corticosterone, known to influence offspring development and temperament[69-71], was elevated by MIA on P10 which was not reduced by EE (main effect of

MIA; **Figure 1J**). Therefore, it is unlikely that attenuation of corticosterone underlies the protective effects of EE on behavior. **Statistics reported in Supplemental Table 3**.

Although the total yield of MEVs recovered during the isolation process was reduced by MIA on P11 (MIA x Time interaction; **Figure 2E**) this was also not protected by EE in this model. Given the characteristics of MEVs to carry bioactive cargos such as miRNAs[18], we turned our interests towards the effects of MIA and EE on MEV cargo composition. The cargo of MEVs isolated from SD-LPS dams showed 66 differentially expressed miRNAs compared to MEVs from SD-Saline dams. Specifically, 22 miRNAs were differentially expressed at P10, while 46 miRNAs were differentially expressed at P11, indicating that miRNA expression was mostly affected the day after the LPS challenge (**Figure 3A,C**). Curiously, in the cargo of MEVs isolated from EE-LPS dams only 29 miRNAs, of which 16 at P10 and 14 at P11, were differentially expressed compared to MEVs from EE-Saline dams (**Figure 3B,D**). This means that, when dams were raised in EE, the number of LPS-induced miRNAs did not vary the day after the LPS challenge (P11) as opposed to what was observed in dams raised in SD. In addition, we found that novel miRNA-629 was downregulated both at P10 and P11 in SD-LPS MEVs compared to SD-Saline MEVs, while the regulation of novel miRNA-375 diverged between P10 and P11. Finally, novel miRNA-1347 was upregulated in the MEVs of LPS-treated dams compared to Saline-treated dams raised in EE, regardless of timepoint (**Figure 3C,D**).

The composition of miRNAs in the nursing offspring hippocampus correlates with MEV miRNA cargo as a function of housing

We then sought to explore the expression of miRNAs in the hippocampus of the nursing offspring (P12) of dams whose MEV miRNA cargo was significantly affected by MIA as a function of EE. LPS caused differential expression of 50 miRNAs (45 miRNAs, adj p-value <0.05) in SD males and 4 miRNAs (miR-1247-5p, adj p-value<0.05) in SD females (**Figure 4A**). In contrast, as also observed in MEV from EE dams, EE dramatically reduced the number of differentially expressed LPS-related miRNAs in the hippocampus compared to miRNAs induced by LPS in SD rats. Namely, LPS regulated 2 miRNAs in EE males and 5 miRNAs in EE females (2 miRNAs, adj p-value <0.05) compared to their respective Saline-treated EE controls (**Figure 4B**).

We were interested in understanding the genomic function of differentially expressed miRNAs regulated by LPS as a function of housing across groups. To predict the miRNA-mRNA targets, we used miRDB (threshold target score>80) combined with DAVID to cluster predicted genes according to their gene ontology. In selected pathways of interest, we found that miRNA-targeted genes in LPS rats (120 in SD rats and 103 in EE rats) were involved in several cellular functions including transcriptional and translational regulation, signal transduction, neurotransmission, as well as epigenetic regulation, regardless of their housing condition. Notably, glutamatergic/GABAergic signaling, immune response, growth factors, and development-related pathways, were targeted in LPS rats raised in SD, but not in EE (**Figure, 4C-F**).

Building on the evidence that LPS caused behavioral alterations in adulthood that depended on early housing condition, we investigated the genomic profile in the ventral hippocampus of the adult offspring. The downregulation of *Thbs1*, a gene involved in autism spectrum disorder risk[72-73], in the hippocampus of male rats whose mothers were raised in SD and challenged with LPS, was consistent with the upregulation of the miRNA regulator, miR-709, in the hippocampus at P12. The same was true for the upregulation of the gene *Naa11*, also involved in autism susceptibility[74], whose regulator, miR-128-1-5p, was downregulated at P12 in the hippocampus of male rats whose mothers were raised in SD and challenged with LPS. We found that miRNA targets at P12 predicted the regulation of several genes that

belonged to the same gene superfamily of the ones that were differentially expressed in the adult ventral hippocampus (**Supplemental Table 2**).

To assess whether miRNAs differentially expressed in MEVs were also affected in the hippocampus of the nursing offspring at P12, we compared the miRNA counts in both miRNA-seq datasets. While the overall pattern of common expression between MEV and hippocampus did not depend on treatment, housing, or sex, there were differences in the genomic function of miRNA sets across groups. Using miRNA Enrichment Analysis and Annotation Tool, we found that, out of the top five pathways, integrin and dopamine signaling, as well as adrenalin and noradrenalin biosynthesis, were regulated across all experimental groups (**Figure 5A-H**). However, the regulation of the glutamate signaling pathway was exclusive to LPS-treated groups raised in EE (**Figure 5G,H**). Additionally, the drug metabolism P450 pathway was regulated in all groups except EE-LPS. Again, this suggests that EE has unique effects on LPS-induced transcriptional regulation.

Discussion

Postpartum infections are known to affect breastfeeding people, however the consequences of illness on lactation and associated offspring outcomes are understudied[38]. Despite awareness of the importance of maternal health in the postpartum period, interventions aimed at supporting caregivers are often underutilized and undervalued. Compounding this gap is a limited understanding of the biological mechanisms through which such interventions may exert their effects. In this context, we identify MEV miRNAs as a dynamic signaling pathway capable of conveying both stress-induced and protective environmental cues, from mothers to their offspring, during the lactational period. This highlights a potential mechanistic link between maternal experience and early developmental programing (see **Figure 6** for summary of proposed mechanisms).

Consistent with prior work, we found that MIA increased corticosterone levels in maternal milk[41], a glucocorticoid known to influence stress responsivity and emotional regulation in offspring[69-71]. However, EE housing did not attenuate MIA-induced elevations in milk corticosterone. This suggests that corticosterone is unlikely to mediate the protective effects of EE observed in this lactational MIA model. Moreover, while MIA caused a reduction in creamatocrit that was rescued by EE, offspring body weights were unaffected during the neonatal and weaning periods. Adult EE males did weigh more than their adult SD counterparts; however, the absence of the early weight differences, in combination with the early transcriptional and behavioral changes, supports the idea that other milk constituents may play a more prominent role in developmental programming.

Instead, our findings point to MEVs and their miRNA cargos as key mediators of environmentally driven signaling between mothers and nursing offspring. Here, MIA induced robust alterations in MEV miRNA cargo, particularly in mothers housed under standard laboratory conditions. In contrast, dams reared in EE displayed fewer MIA-induced changes in MEV miRNA profiles, supporting the notion that enrichment stabilizes maternal biological outputs under stress. Notably, in MEVs from standard housed dams, novel miRNA-629 was consistently downregulated by MIA across both sampling days (P10 and P11), while novel miRNA-375 showed a biphasic expression pattern. These miRNAs predicted by miRDeep 2.0 and clustered by DAVID targeted genes involved in transcriptional regulation, ion channel function, and signal transduction, pathways critical to neurodevelopment and synaptic plasticity[75,76]. In MEVs from enriched dams, MIA upregulated novel miRNA-1347, which is predicted to target the expression of Tbl1xr1, a gene implicated in developmental

disorders whose regulation confers protection[77-80]. This suggests that EE-modified MEV cargo programs developmental adaptation in the offspring via transcriptional processes.

The regulatory potential of MEV miRNAs was consistent with changes observed in the neonatal hippocampus on P12. The differential expression of MIA-induced hippocampal miRNAs in SD offspring was dramatically greater in male than in females, aligning with previous work showing sex specific vulnerability to early life stressors[81]. Similar to MIA-induced differentially expressed miRNAs in MEVs, enrichment reduced the number of differentially expressed hippocampal miRNAs, indicating that EE can buffer against stress-induced transcriptomic modifications. Several miRNAs were expressed in both MEVs and the neonatal hippocampus, suggesting potential transfer of regulatory signals from milk to brain. Moreover, miR-709 and miR-128-1-5p were differentially expressed in neonatal hippocampus and showed the expected directional changes in their predicted gene targets (*Thbs1* and *Naa11*) in the adult hippocampal transcriptome. Cross-tissue miRNA-mRNA targets provide compelling support to the hypothesis that MEVs are sensitive to maternal environmental conditions and may relay this information to the developing offspring brain[6].

Lactational MIA induced behavioral alterations in offspring that were also buffered by enrichment. In female offspring, MIA reduced the percent of time animals explored the center of the open field arena, indicative of heightened anxiety-like behavior; in male offspring, social preference was diminished by the MIA challenge. Importantly, EE exposure prevented these adverse outcomes in both sexes, consistent with prior studies demonstrating the protective effects of enriched housing on offspring stress regulation and social engagement[48-50,82]. Interestingly, differences in maternal care were not observed across groups, suggesting that EE protection was not mediated by changes in overt maternal behaviors, such as time spent on the nest or frequency of licking/grooming. This is noteworthy, as enriched housing can modulate maternal care under non-stress conditions[51], but the added MIA challenge during lactation may have masked group differences. Instead, it appears that the behavioral buffering effects of EE may involve more subtle or molecular-level changes in maternal signaling via MEVs in this model.

We acknowledge that there are limitations one should take into account when interpreting some of the findings. First, the dynamic nature of milk composition across the lactational period limits the temporal generalizability of our results. Our study only sampled across two days, and it remains unclear whether certain phases of lactation may be more or less responsive to either MIA or EE. Second, although we identified associations between miRNAs in MEVs and neonatal hippocampus, direct mechanistic evidence for MEV transfer and functional integration in the offspring brain is lacking. It is not known whether neonatal hippocampal miRNA profiles were directly reflective of deposited maternal MEV cargo or rather due to indirect actions between MEVs and the microbiome, influencing the hippocampal transcriptome, for example. Future studies using fluorescently labeled MEVs and individual miRNA cargo, in addition to MEV depletion and supplementation approaches are needed to directly test the uptake and influence of MEVs in brain. MEVs themselves carry multiple types of cargo, including lipids and peptides that can influence neurodevelopment, in addition to the miRNAs evaluated here. Finally, while our study highlights MEVs as plausible mediators of environmental signaling, we do not exclude interactive effects with other milk constituents such as glucocorticoids or fats[41]. The complexity of milk composition argues against a single-pathway explanation, and it is likely that additive or synergistic interactions across multiple milk factors are shaping neurodevelopmental trajectories.

Conclusions

Breastfeeding remains the recommended source of nutrition during most maternal illnesses, as many pathogens including influenza, COVID-19, and foodborne agents are not transmitted directly through breast milk, yet antibodies against these pathogens are transmitted to offspring[83]. Critically, illness and stress can alter milk composition, including the regulatory cargo of MEVs. Our findings demonstrate that MEV miRNAs are sensitive to the maternal environment and can shape offspring neurodevelopment, highlighting the need for increased support for breastfeeding individuals. This includes a need for improved education and communication pathways to ensure breastfeeding individuals are informed and supported[84]. Indeed, enrichment exposure stabilized MEV cargo and mitigated the effects of MIA on offspring, positioning MEV miRNAs as dynamic programming signals by which both stress-induced and protective cues are relayed. Given evidence linking breastfeeding to reduced neurodevelopmental disorder risk[85], our study positions MEVs as a key mechanism by which maternal experiences influence infant brain development, with important implications for maternal-infant health policy and care.

Figure Captions

Figure 1. Environmental enrichment protects against lactational maternal immune activation (MIA) induced changes in offspring behavior. (**A**) Timeline of experimental procedures. (**B**) Maternal sickness behavior scores. (**C**) Neonatal huddling behavior (average number of pup clusters) across P10 and P11. Adult (**D**) male and (**E**) female percent (%) time in the center of the open field. Adult (**F**) male and (**G**) female social preference. (**H**) Percent (%) creamatocrit from P10-P11 and (**I**) the full litter body weights of nursing pups (g) across P10-P12. (**J**) Milk corticosterone (ng/mL) after MIA on P10; ND: non detectable values that fell below level of assay detection. Data are expressed as mean ± SEM; SD-Saline: n = 8; SD-LPS: n = 8; EE-Saline: n = 7; EE-LPS: n = 8. *$p < 0.05$, **$p <0.01$, ***$p <0.001$, SD-Saline versus SD-LPS; #$p < 0.05$, ##$p <0.01$, ###$p <0.001$, SD-LPS versus EE-LPS; [a]$p < 0.05$, [aa]$p < 0.01$, [aaa]$p <0.001$, main effect of drug; [b]$p < 0.05$, [bb]$p < 0.01$, [bbb]$p <0.001$, main effect of housing; [c]$p < 0.05$, [cc]$p < 0.01$, [ccc]$p <0.001$, main effect of time; LPS: lipopolysaccharide; P: postnatal day. Created in BioRender. Kentner, A. (2024) https://BioRender.com/z10e449.

Figure 2. Characterization of milk derived extracellular vesicles (MEVs) isolated from rat milk following maternal immune activation (MIA). (**A**) The differential ultracentrifugation (DUC) and serial filtration protocol followed for MEV isolation. (**B**) Size (nm) and concentration (particles/mL) of MEVs as determined by nanoparticle tracking analysis (NTA) with a mean size of 115.9 ± 31.5 nm. (**C**) The presence or absence of positive (Alix, Flotillin-1, CD9; cellular control: human MEVs) and negative (Calnexin; cellular control: Hep G2 cells) protein markers for MEVs as determined by western immunoblotting. (**D**) Transmission electron microscopy (TEM) of a representative MEV pellet. Scale bar is 200 nm with some individual MEVs identified in sizes ranging from 70.5 to 126 nm. (**E**) Total yield of MEVs recovered. Data are expressed as mean ± SEM, n = 6. [aa]$p < 0.01$, main effect of drug; LPS: lipopolysaccharide; P: postnatal day; SD: standard housing. Created in BioRender. Kentner, A. (2025) https://BioRender.com/x9rk66v.

Figure 3. Lactational maternal immune activation (MIA) alters milk derived extracellular vesicles (MEV) miRNA cargos, which are buffered by environmental

enrichment (EE) housing. miRNA clustered in the heatmaps display the Log_2FC p<0.01, FC>2 in (**A**) SD-LPS versus SD-Saline and (**B**) EE-LPS versus EE-Saline MEVs at P10 and P11. Venn diagrams show the number of differentially expressed miRNA in P10 (blue) and P11 (red), and their overlapping miRNAs and direction of change (↑ upregulated, ↓ downregulated) for (**C**) SD-LPS versus SD-Saline and (**D**) EE-LPS versus EE-Saline. LPS: lipopolysaccharide; P: postnatal day; SD: standard housing; n = 6. *If significant at P10 and P11.

Figure 4. Lactational maternal immune activation (MIA) alters miRNA expression in the hippocampus of nursing offspring, which is buffered by environmental enrichment (EE) housing. miRNA clustered in the heatmaps display the Log_2FC p<0.01, FC>2 in (**A**) SD-LPS versus SD-Saline and (**B**) EE-LPS versus EE-Saline neonatal hippocampus on P12 for male and female offspring. (**C,D**) The bubble plots show the enrichment score of gene pathways generated using DAVID by clustering genes predicted as miRNA-mRNA targets (mirdb.org, target score >80). Gene counts per cluster are represented as bubble size. Gene pathways highlighted are top unique clusters from (**C**) Figure A, and (**D**) Figure B. (**E,F**) Pie charts to visualize gene count distribution of pathways highlighted in (**E**) Figure C and (**F**) Figure D. LPS: lipopolysaccharide; P: postnatal day; SD: standard housing; n = 6; *if FDR<0.05.

Figure 5. Overlapping miRNAs matched between maternal MEVs and the hippocampus of their nursing male and female neonatal offspring. (**A, C, E, G**) Venn diagrams showing the number of miRNAs found in milk exosomes at P11 (yellow) and the hippocampus of male (blue) and female (red) offspring at P12, and their overlapping miRNA in groups (**A**) SD-Saline, (**C**) SD-LPS, (**E**) EE-Saline, and (**G**) EE-LPS. (**B, D, F, H**) The bubble plots show the -Log□□ (p-value) of the top 5 enriched miRWalk pathways identified for the overlapping miRNA in (**B**) Figure A, (**D**) Figure C, (**F**) Figure E, and (**H**) Figure G, using the miRNA Enrichment Analysis and Annotation Tool (ccb-compute2.cs.uni-saarland.de/mieaa). miRNA count per pathway is indicated by bubble size. LPS: lipopolysaccharide; P: postnatal day; SD: standard housing; EE: environmental enrichment; n = 6.

Figure 6. Overarching hypothesis of milk derived extracellular vesicles (MEVs) as a mechanism of postnatal neuroimmune programming. (**A**) Nursing mammals pass MEVs through their milk to their feeding offspring. (**B**) Insults such as inflammatory stressors change the composition of MEV cargo. (**C**) Data suggest that when absorbed by nursing offspring, MEVs can interact with the gut microbiome and/or cross into and accumulate in the offspring brain. (**D**) We propose that maternal immune activation (MIA) alters the composition of MEV cargo, which then infiltrates the feeding offspring. Through either direct (brain) or indirect (e.g., gut microbiome) interactions, the altered MEV cargo (e.g., miRNAs), lead to transcriptional changes in brain regions like the hippocampus, affecting later life behavior in offspring. Environmental enrichment interventions can neutralize these changes, conserving behavior. Created in BioRender. Kentner, A. (2025) https://BioRender.com/50zpt07.

Supplemental Table 1. Maternal immune activation reporting guidelines table.

Supplemental Table 2. Validation of miRNA-mRNA target predictions from nursing offspring and associated mRNA in the adult hippocampus. The table shows the overlap of differentially expressed (p<0.01, FC>2) gene families (column) that are common in the

hippocampus of male and female rats at P12 and present at P74 in their respective groups (rows). LPS: lipopolysaccharide; P: postnatal day; SD: standard housing; EE: environmental enrichment; n = 6.

Supplemental Table 3. Statistical Reporting of Results.

Supplemental Figure 1. Maternal care and offspring outcomes following lactational maternal immune activation (MIA) and environmental enrichment (EE) housing. Total distance traveled (meters) for (**A**) male and (**B**) female adult offspring. Maternal total time on nest (seconds) and (**D**) frequency of licking and grooming behaviors across P9 to P11. Adult body weights (g) for (**E**) male and (**F**) female offspring. Data are expressed as mean ± SEM; ##p <0.01, SD-LPS versus EE-LPS; LPS: lipopolysaccharide; P: postnatal day; SD: standard housing; n = 7-8.

Funding and Disclosures
This project was funded by NIMH under Award Number R15MH114035 (to ACK), Tuoro University Presidential Research Development Grant (to JM), the Manitoba Medical Service Foundation Grant (to SW), the Massachusetts College of Pharmacy and Health Sciences (MCPHS) Center for Undergraduate Research (TJL & ACK), a MCPHS Summer Undergraduate Research Fellowship (SURF) to TJL, an MCPHS New Discovery Award (ACK), and the Massachusetts Life Sciences Center – Capital Grants, which provided financial and equipment support. Electron Microscopy imaging, consultation, and/or services were performed in the HMS Electron Microscopy Facility. We thank the Genomics Core Laboratory and its late Director, Dr. Brahmaraju Mopidevi, at the New York Medical College for the miRNA sequencing of the neonatal hippocampus. The authors would also like to thank Azenta Life Sciences for their support with the miRNA sequencing of MEVs and adult hippocampal RNA sequencing. Figures 1, 2 and 6 were made with BioRender.com and an earlier version of this manuscript was posted on the preprint server bioRvix. The content is solely the responsibility of the authors and does not necessarily represent the official views of any of the financial supporters. The sequencing datasets are being deposited to GEO.

Author Contributions
Julia M, TJL, BH, DL, JS, JL, Jordan M, and ACK ran the experiments; Julia M., BH, Jordan M., BB, BB, BM, SW, & ACK analyzed and interpreted the data; ACK wrote the manuscript; Julia M, BH, TJL, BM, JS, JL, SW, Jordan M, & ACK edited the manuscript. Jordan M, SW, & ACK were involved in project design. Jordan M, SW, and ACK supervised trainees. ACK conceptualized the project.

Conflict of Interest
The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Supplementary information is available at https://www.nature.com/mp/

References

1. Aplin JD, Myers JE, Timms K, Westwood M. Tracking placental development in health and disease. Nat Rev Endocrinol. 2020 Sep;16(9):479-494. doi: 10.1038/s41574-020-0372-6. Epub 2020 Jun 29. PMID: 32601352.
2. Morgan CP, Chan JC, Bale TL. Driving the Next Generation: Paternal Lifetime Experiences Transmitted via Extracellular Vesicles and Their Small RNA Cargo. Biol Psychiatry. 2019 Jan 15;85(2):164-171. doi: 10.1016/j.biopsych.2018.09.007. Epub 2018 Sep 27. PMID: 30580777; PMCID: PMC6309802.
3. Chan JC, Morgan CP, Adrian Leu N, Shetty A, Cisse YM, Nugent BM, Morrison KE, Jašarević E, Huang W, Kanyuch N, Rodgers AB, Bhanu NV, Berger DS, Garcia BA, Ament S, Kane M, Neill Epperson C, Bale TL. Reproductive tract extracellular vesicles are sufficient to transmit intergenerational stress and program neurodevelopment. Nat Commun. 2020 Mar 20;11(1):1499. doi: 10.1038/s41467-020-15305-w. PMID: 32198406; PMCID: PMC7083921.
4. Sánchez-Rubio M, Abarzúa-Catalán L, Del Valle A, Méndez-Ruette M, Salazar N, Sigala J, Sandoval S, Godoy MI, Luarte A, Monteiro LJ, Romero R, Choolani MA, Wyneken Ú, Illanes SE, Bátiz LF. Maternal stress during pregnancy alters circulating small extracellular vesicles and enhances their targeting to the placenta and fetus. Biol Res. 2024 Sep 28;57(1):70. doi: 10.1186/s40659-024-00548-4. PMID: 39342314; PMCID: PMC11438166.
5. Zhou F, Ebea P, Mutai E, Wang H, Sukreet S, Navazesh S, Dogan H, Li W, Cui J, Ji P, Ramirez DMO, Zempleni J. Small Extracellular Vesicles in Milk Cross the Blood-Brain Barrier in Murine Cerebral Cortex Endothelial Cells and Promote Dendritic Complexity in the Hippocampus and Brain Function in C57BL/6J Mice. Front Nutr. 2022 May 6;9:838543. doi: 10.3389/fnut.2022.838543. PMID: 35600828; PMCID: PMC9121399.
6. Bozack AK, Colicino E, Rodosthenous R, Bloomquist TR, Baccarelli AA, Wright RO, Wright RJ, Lee AG. Associations between maternal lifetime stressors and negative events in pregnancy and breast milk-derived extracellular vesicle microRNAs in the programming of intergenerational stress mechanisms (PRISM) pregnancy cohort. Epigenetics. 2021 Apr;16(4):389-404. doi: 10.1080/15592294.2020.1805677. Epub 2020 Aug 25. PMID: 32777999; PMCID: PMC7996083.
7. Zempleni J, Sukreet S, Zhou F, Wu D, Mutai E. Milk-Derived Exosomes and Metabolic Regulation. Annu Rev Anim Biosci. 2019 Feb 15;7:245-262. doi: 10.1146/annurev-animal-020518-115300. Epub 2018 Oct 4. PMID: 30285461.
8. Yáñez-Mó M, Siljander PR, Andreu Z, Zavec AB, Borràs FE, Buzas EI, Buzas K, Casal E, Cappello F, Carvalho J, Colás E, Cordeiro-da Silva A, Fais S, Falcon-Perez JM, Ghobrial IM, Giebel B, Gimona M, Graner M, Gursel I, Gursel M, Heegaard NH, Hendrix A, Kierulf P, Kokubun K, Kosanovic M, Kralj-Iglic V, Krämer-Albers EM, Laitinen S, Lässer C, Lener T, Ligeti E, Linē A, Lipps G, Llorente A, Lötvall J, Manč ek-Keber M, Marcilla A, Mittelbrunn M, Nazarenko I, Nolte-'t Hoen EN, Nyman TA, O'Driscoll L, Olivan M, Oliveira C, Pállinger É, Del Portillo HA, Reventós J, Rigau M, Rohde E, Sammar M, Sánchez-Madrid F, Santarém N, Schallmoser K, Ostenfeld MS, Stoorvogel W, Stukelj R, Van der Grein SG, Vasconcelos MH, Wauben MH, De Wever O. Biological properties of extracellular vesicles and their physiological functions. J Extracell Vesicles. 2015 May 14;4:27066. doi: 10.3402/jev.v4.27066. PMID: 25979354; PMCID: PMC4433489.

9. EL Andaloussi S, Mäger I, Breakefield XO, Wood MJ. Extracellular vesicles: biology and emerging therapeutic opportunities. Nat Rev Drug Discov. 2013 May;12(5):347-57. doi: 10.1038/nrd3978. Epub 2013 Apr 15. PMID: 23584393.

10. Wijenayake S, Martz J, Lapp HE, Storm JA, Champagne FA, Kentner AC. The contributions of parental lactation on offspring development: It's not udder nonsense! Horm Behav. 2023 Jul;153:105375. doi: 10.1016/j.yhbeh.2023.105375. Epub 2023 Jun 1. PMID: 37269591; PMCID: PMC10351876.

11. Ames SR, Lotoski LC, Azad MB. Comparing early life nutritional sources and human milk feeding practices: personalized and dynamic nutrition supports infant gut microbiome development and immune system maturation. Gut Microbes. 2023 Jan-Dec;15(1):2190305. doi: 10.1080/19490976.2023.2190305. PMID: 37055920; PMCID: PMC10114993.

12. Victora CG, Bahl R, Barros AJ, França GV, Horton S, Krasevec J, Murch S, Sankar MJ, Walker N, Rollins NC; Lancet Breastfeeding Series Group. Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. Lancet. 2016 Jan 30;387(10017):475-90. doi: 10.1016/S0140-6736(15)01024-7. PMID: 26869575.

13. de Weerth C, Aatsinki AK, Azad MB, Bartol FF, Bode L, Collado MC, Dettmer AM, Field CJ, Guilfoyle M, Hinde K, Korosi A, Lustermans H, Mohd Shukri NH, Moore SE, Pundir S, Rodriguez JM, Slupsky CM, Turner S, van Goudoever JB, Ziomkiewicz A, Beijers R. Human milk: From complex tailored nutrition to bioactive impact on child cognition and behavior. Crit Rev Food Sci Nutr. 2023;63(26):7945-7982. doi: 10.1080/10408398.2022.2053058. Epub 2022 Mar 30. PMID: 35352583.

14. Melnik BC, Stremmel W, Weiskirchen R, John SM, Schmitz G. Exosome-Derived MicroRNAs of Human Milk and Their Effects on Infant Health and Development. Biomolecules. 2021 Jun 7;11(6):851. doi: 10.3390/biom11060851. PMID: 34200323; PMCID: PMC8228670.

15. Khanam A, Ngu A, Zempleni J. Bioavailability of orally administered small extracellular vesicles from bovine milk in C57BL/6J mice. Int J Pharm. 2023 May 25;639:122974. doi: 10.1016/j.ijpharm.2023.122974. Epub 2023 Apr 25. PMID: 37105241; PMCID: PMC10175213.

16. Hock A, Miyake H, Li B, Lee C, Ermini L, Koike Y, Chen Y, Määttänen P, Zani A, Pierro A. Breast milk-derived exosomes promote intestinal epithelial cell growth. J Pediatr Surg. 2017 May;52(5):755-759. doi: 10.1016/j.jpedsurg.2017.01.032. Epub 2017 Jan 29. PMID: 28188035.

17. Manca S, Upadhyaya B, Mutai E, Desaulniers AT, Cederberg RA, White BR, Zempleni J. Milk exosomes are bioavailable and distinct microRNA cargos have unique tissue distribution patterns. Sci Rep. 2018 Jul 27;8(1):11321. doi: 10.1038/s41598-018-29780-1. PMID: 30054561; PMCID: PMC6063888.

18. Jiang X, You L, Zhang Z, Cui X, Zhong H, Sun X, Ji C, Chi X. Biological Properties of Milk-Derived Extracellular Vesicles and Their Physiological Functions in Infant. Front Cell Dev Biol. 2021 Jun 25;9:693534. doi: 10.3389/fcell.2021.693534. PMID: 34249944; PMCID: PMC8267587

19. Lönnerdal B. Human Milk MicroRNAs/Exosomes: Composition and Biological Effects. Nestle Nutr Inst Workshop Ser. 2019;90:83-92. doi: 10.1159/000490297. Epub 2019 Mar 13. PMID: 30865991.

20. Chen CC, Liu L, Ma F, Wong CW, Guo XE, Chacko JV, Farhoodi HP, Zhang SX, Zimak J, Ségaliny A, Riazifar M, Pham V, Digman MA, Pone EJ, Zhao W. Elucidation of Exosome Migration across the Blood-Brain Barrier Model In Vitro. Cell Mol Bioeng. 2016 Dec;9(4):509-529. doi: 10.1007/s12195-016-0458-3. Epub 2016 Jul 7. PMID: 28392840; PMCID: PMC5382965.

21. Saunders NR, Habgood MD, Dziegielewska KM. Barrier mechanisms in the brain, II. Immature brain. Clin Exp Pharmacol Physiol. 1999 Feb;26(2):85-91. doi: 10.1046/j.1440-1681.1999.02987.x. PMID: 10065326.

22. Mallard C, Ek CJ, Vexler ZS. The myth of the immature barrier systems in the developing brain: role in perinatal brain injury. J Physiol. 2018 Dec;596(23):5655-5664. doi: 10.1113/JP274938. Epub 2018 Apr 16. PMID: 29528501; PMCID: PMC6265562.

23. Stolp H, Neuhaus A, Sundramoorthi R, Molnár Z. The Long and the Short of it: Gene and Environment Interactions During Early Cortical Development and Consequences for Long-Term Neurological Disease. Front Psychiatry. 2012 Jun 12;3:50. doi: 10.3389/fpsyt.2012.00050. PMID: 22701439; PMCID: PMC3372875.

24. Haney MJ, Klyachko NL, Harrison EB, Zhao Y, Kabanov AV, Batrakova EV. TPP1 Delivery to Lysosomes with Extracellular Vesicles and their Enhanced Brain Distribution in the Animal Model of Batten Disease. Adv Healthc Mater. 2019 Jun;8(11):e1801271. doi: 10.1002/adhm.201801271. Epub 2019 Apr 18. PMID: 30997751; PMCID: PMC6584948.

25. Li Y, Yang YY, Ren JL, Xu F, Chen FM, Li A. Exosomes secreted by stem cells from human exfoliated deciduous teeth contribute to functional recovery after traumatic brain injury by shifting microglia M1/M2 polarization in rats. Stem Cell Res Ther. 2017 Sep 29;8(1):198. doi: 10.1186/s13287-017-0648-5. PMID: 28962585; PMCID: PMC5622448.

26. Paolicelli RC, Bergamini G, Rajendran L. Cell-to-cell Communication by Extracellular Vesicles: Focus on Microglia. Neuroscience. 2019 May 1;405:148-157. doi: 10.1016/j.neuroscience.2018.04.003. Epub 2018 Apr 13. PMID: 29660443.

27. Wijenayake S, Eisha S, Purohit MK, McGowan PO. Milk derived extracellular vesicle uptake in human microglia regulates the DNA methylation machinery : Short title: milk-derived extracellular vesicles and the epigenetic machinery. Sci Rep. 2024 Nov 19;14(1):28630. doi: 10.1038/s41598-024-79724-1. PMID: 39562680; PMCID: PMC11576889.

28. Zhang H, Shykind B, Sun T. Approaches to manipulating microRNAs in neurogenesis. Front Neurosci. 2013 Jan 17;6:196. doi: 10.3389/fnins.2012.00196. PMID: 23335878; PMCID: PMC3547386.

29. Saliminejad K, Khorram Khorshid HR, Soleymani Fard S, Ghaffari SH. An overview of microRNAs: Biology, functions, therapeutics, and analysis methods. J Cell Physiol. 2019 May;234(5):5451-5465. doi: 10.1002/jcp.27486. Epub 2018 Nov 23. PMID: 30471116.

30. Liao Y, Du X, Li J, Lönnerdal B. Human milk exosomes and their microRNAs survive digestion in vitro and are taken up by human intestinal cells. Mol Nutr Food Res. 2017 Nov;61(11). doi: 10.1002/mnfr.201700082. Epub 2017 Aug 15. PMID: 28688106.

31. Alsaweed M, Lai CT, Hartmann PE, Geddes DT, Kakulas F. Human Milk Cells Contain Numerous miRNAs that May Change with Milk Removal and Regulate Multiple Physiological Processes. Int J Mol Sci. 2016 Jun 17;17(6):956. doi: 10.3390/ijms17060956. PMID: 27322254; PMCID: PMC4926489.

32. Zhang Y, Belaid M, Luo X, Daci A, Limani R, Mantaj J, Zilbauer M, Nayak K, Vllasaliu D. Probing milk extracellular vesicles for intestinal delivery of RNA therapies. J Nanobiotechnology. 2023 Nov 3;21(1):406. doi: 10.1186/s12951-023-02173-x. PMID: 37924132; PMCID: PMC10623793.

33. Seo M, Anderson G. Gut-Amygdala Interactions in Autism Spectrum Disorders: Developmental Roles via regulating Mitochondria, Exosomes, Immunity and

microRNAs. Curr Pharm Des. 2019;25(41):4344-4356. doi: 10.2174/1381612825666191105102545. PMID: 31692435.

34. Collier SA, Rasmussen SA, Feldkamp ML, Honein MA; National Birth Defects Prevention Study. Prevalence of self-reported infection during pregnancy among control mothers in the National Birth Defects Prevention Study. Birth Defects Res A Clin Mol Teratol. 2009 Mar;85(3):193-201. doi: 10.1002/bdra.20540. PMID: 19086018.

35. Brown AS, Meyer U. Maternal Immune Activation and Neuropsychiatric Illness: A Translational Research Perspective. Am J Psychiatry. 2018 Nov 1;175(11):1073-1083. doi: 10.1176/appi.ajp.2018.17121311. Epub 2018 Sep 17. PMID: 30220221; PMCID: PMC6408273.

36. Estes ML, McAllister AK. Maternal immune activation: Implications for neuropsychiatric disorders. Science. 2016 Aug 19;353(6301):772-7. doi: 10.1126/science.aag3194. PMID: 27540164; PMCID: PMC5650490.

37. Kentner AC, Bilbo SD, Brown AS, Hsiao EY, McAllister AK, Meyer U, Pearce BD, Pletnikov MV, Yolken RH, Bauman MD. Maternal immune activation: reporting guidelines to improve the rigor, reproducibility, and transparency of the model. Neuropsychopharmacology. 2019 Jan;44(2):245-258. doi: 10.1038/s41386-018-0185-7. Epub 2018 Aug 21. PMID: 30188509; PMCID: PMC6300528.

38. PRGLAC [Task Force on Research Specific to Pregnant Women and Lactating Women], 2018. Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD): https://www.nichd.nih.gov/sites/default/files/2018-09/PRGLAC_Report.pdf.

39. Vilela FC, Antunes-Rodrigues J, Elias LL, Giusti-Paiva A. Corticosterone synthesis inhibitor metyrapone preserves changes in maternal behavior and neuroendocrine responses during immunological challenge in lactating rats. Neuroendocrinology. 2013;97(4):322-30. doi: 10.1159/000346354. Epub 2013 Feb 6. PMID: 23295343.

40. Nascimento AF, Alves GJ, Massoco CO, Teodorov E, Felicio LF, Bernardi MM. Lipopolysaccharide-induced sickness behavior in lactating rats decreases ultrasonic vocalizations and exacerbates immune system activity in male offspring. Neuroimmunomodulation. 2015;22(4):213-21. doi: 10.1159/000363350. Epub 2014 Aug 19. Erratum in: Neuroimmunomodulation. 2015;22(4):278. PMID: 25139475.

41. DeRosa H, Caradonna SG, Tran H, Marrocco J, Kentner AC. Got milk? Maternal immune activation during the mid-lactational period affects nutritional milk quality and adolescent offspring sensory processing in male and female rats. Mol Psychiatry. 2022 Dec;27(12):4829-4842. doi: 10.1038/s41380-022-01744-y. Epub 2022 Sep 2. PMID: 36056174; PMCID: PMC9771965.

42. Aubert A, Goodall G, Dantzer R, Gheusi G. Differential effects of lipopolysaccharide on pup retrieving and nest building in lactating mice. Brain Behav Immun. 1997 Jun;11(2):107-18. doi: 10.1006/brbi.1997.0485. PMID: 9299060.

43. Merengueli, J., Kentner, A.C. (2025). Maternal immune activation during the lactational period alters offspring behavior, reproductive development, and immune function in mice. [preprint]. bioRxiv 2025.05.26.656156; doi: https://doi.org/10.1101/2025.05.26.656156.

44. Edwards PD, Lavergne SG, McCaw LK, Wijenayake S, Boonstra R, McGowan PO, Holmes MM. Maternal effects in mammals: Broadening our understanding of offspring programming. Front Neuroendocrinol. 2021 Jul;62:100924. doi: 10.1016/j.yfrne.2021.100924. Epub 2021 May 14. PMID: 33992652.

45. Wang Y, Fang J, Zeng HF, Zhong JF, Li HX, Chen KL. Identification and bioinformatics analysis of differentially expressed milk exosomal microRNAs in milk

exosomes of heat-stressed Holstein cows. Funct Integr Genomics. 2022 Feb;22(1):77-87. doi: 10.1007/s10142-021-00814-8. Epub 2021 Nov 27. PMID: 34839400.

46. Ren W, La Y, Ma X, Wu X, Guo X, Chu M, Yan P, Lan X, Liang C. Comparative Analysis of miRNA Expression Profiles of Yak Milk-Derived Exosomes at Different Altitudes. Animals (Basel). 2025 Jan 2;15(1):87. doi: 10.3390/ani15010087. PMID: 39795030; PMCID: PMC11718820.

47. Golan-Gerstl R, Ben Ya'acov A, Musseri M, Goldenberg R, Chammah Y, Cherki T, Reif S, Bar Gil Shitrit A. Expression Profile of MicroRNAs in Breast Milk of Women With Inflammatory Bowel Disease: Correlation With Disease Activity and Medical Treatments. Inflamm Bowel Dis. 2025 Apr 10;31(4):912-922. doi: 10.1093/ibd/izae290. PMID: 39820274; PMCID: PMC11985391.

48. Núñez Estevez KJ, Rondón-Ortiz AN, Nguyen JQT, Kentner AC. Environmental influences on placental programming and offspring outcomes following maternal immune activation. Brain Behav Immun. 2020 Jan;83:44-55. doi: 10.1016/j.bbi.2019.08.192. Epub 2019 Sep 4. PMID: 31493445; PMCID: PMC6906258.

49. Connors EJ, Shaik AN, Migliore MM, Kentner AC. Environmental enrichment mitigates the sex-specific effects of gestational inflammation on social engagement and the hypothalamic pituitary adrenal axis-feedback system. Brain Behav Immun. 2014 Nov;42:178-90. doi: 10.1016/j.bbi.2014.06.020. Epub 2014 Jul 7. PMID: 25011058.

50. Zhao X, Mohammed R, Tran H, Erickson M, Kentner AC. Poly (I:C)-induced maternal immune activation modifies ventral hippocampal regulation of stress reactivity: prevention by environmental enrichment. Brain Behav Immun. 2021 Jul;95:203-215. doi: 10.1016/j.bbi.2021.03.018. Epub 2021 Mar 23. PMID: 33766701; PMCID: PMC8187276.

51. DeRosa H, Caradonna SG, Tran H, Marrocco J, Kentner AC. Milking It for All It's Worth: The Effects of Environmental Enrichment on Maternal Nurturance, Lactation Quality, and Offspring Social Behavior. eNeuro. 2022 Aug 25;9(4):ENEURO.0148-22.2022. doi: 10.1523/ENEURO.0148-22.2022. PMID: 35995560; PMCID: PMC9417599.

52. Rosenfeld P, Wetmore JB, Levine S. Effects of repeated maternal separations on the adrenocortical response to stress of preweanling rats. Physiol Behav. 1992 Oct;52(4):787-91. doi: 10.1016/0031-9384(92)90415-x. PMID: 1409954.

53. Kentner AC, James JS, Miguelez M, Bielajew C. Investigating the hedonic effects of interferon-alpha on female rats using brain-stimulation reward. Behav Brain Res. 2007 Feb 12;177(1):90-9. doi: 10.1016/j.bbr.2006.10.033. Epub 2006 Nov 28. PMID: 17126922.

54. Hayley S, Kelly O, Anisman H. Murine tumor necrosis factor-alpha sensitizes plasma corticosterone activity and the manifestation of shock: modulation by histamine. J Neuroimmunol. 2002 Oct;131(1-2):60-9. doi: 10.1016/s0165-5728(02)00259-x. PMID: 12458037.

55. Drugs and Lactation Database (LactMed) [Internet]. Bethesda (MD): National Library of Medicine (US) 200g; Isoflurane. [Updated 2020 Nov 16]. Bookshelf. Available from: https://www.tandfonline.com/doi/abs/10.1080/15424065.2012.735134?casa_token=_4T1848mJgwAAAAA:qSoL3T_p88u53bA6gskHc6ZnJY_0n2etdDL5InvnIB6DZ19xhGPoUI1Lxvob0u2AWVBGcrpOowBOEQ

56. Par Pharmaceutical, Inc, Pitocin [Label]. Chestnut Ridge (NY): Par Pharmaceutical; [2020]. Available

from: https://dailymed.nlm.nih.gov/dailymed/fda/fdaDrugXsl.cfm?setid=6d4b2c25-2e5d-49b5-93bc-2ae8a20916d1&type=display

57. Lee JJ, Rubin AP. Breast feeding and anaesthesia. Anaesthesia. 1993 Jul;48(7):616-25. doi: 10.1111/j.1365-2044.1993.tb07130.x. PMID: 8346780.

58. Paul HA, Hallam MC, Reimer RA. Milk Collection in the Rat Using Capillary Tubes and Estimation of Milk Fat Content by Creamatocrit. J Vis Exp. 2015 Dec 16;(106):e53476. doi: 10.3791/53476. PMID: 26709708; PMCID: PMC4694028.

59. Barnett HY, Geys H, Jacobs T, Jaki T. Methods for non-compartmental pharmacokinetic analysis with observations below the limit of quantification. Stat Biopharm Res. 2021;13:59–70. https://doi.org/10.1080/19466315.2019.1701546.

60. Wijenayake S, Eisha S, Tawhidi Z, Pitino MA, Steele MA, Fleming AS, McGowan PO. Comparison of methods for pre-processing, exosome isolation, and RNA extraction in unpasteurized bovine and human milk. PLoS One. 2021 Sep 30;16(9):e0257633. doi: 10.1371/journal.pone.0257633. PMID: 34591894; PMCID: PMC8483318.

61. Welsh JA, Goberdhan DCI, O'Driscoll L, Buzas EI, Blenkiron C, Bussolati B…Zimmerman AJ; MISEV Consortium; Théry C, Witwer KW. Minimal information for studies of extracellular vesicles (MISEV2023): From basic to advanced approaches. J Extracell Vesicles. 2024 Feb;13(2):e12404. doi: 10.1002/jev2.12404.

62. Morozumi M, Izumi H, Shimizu T, Takeda Y. Comparison of isolation methods using commercially available kits for obtaining extracellular vesicles from cow milk. J Dairy Sci. 2021 Jun;104(6):6463-6471. doi: 10.3168/jds.2020-19849. Epub 2021 Mar 11. PMID: 33714584.

63. Naskar S, Narducci R, Balzani E, Cwetsch AW, Tucci V, Cancedda L. The development of synaptic transmission is time-locked to early social behaviors in rats. Nat Commun. 2019 Mar 13;10(1):1195. doi: 10.1038/s41467-019-09156-3. PMID: 30867422; PMCID: PMC6416358.

64. Scarborough J, Mueller F, Arban R, Dorner-Ciossek C, Weber-Stadlbauer U, Rosenbrock H, Meyer U, Richetto J. Preclinical validation of the micropipette-guided drug administration (MDA) method in the maternal immune activation model of neurodevelopmental disorders. Brain Behav Immun. 2020 Aug;88:461-470. doi: 10.1016/j.bbi.2020.04.015. Epub 2020 Apr 9. PMID: 32278850.

65. Nguyen HB, Bagot RC, Diorio J, Wong TP, Meaney MJ. Maternal care differentially affects neuronal excitability and synaptic plasticity in the dorsal and ventral hippocampus. Neuropsychopharmacology. 2015 Jun;40(7):1590-9. doi: 10.1038/npp.2015.19. Epub 2015 Jan 19. PMID: 25598429; PMCID: PMC4915255.

66. Miles J, Shevlin M. (2001). Applying regression and correlation: a guide for students and researchers. London: Sage.

67. Clayton JA. Applying the new SABV (sex as a biological variable) policy to research and clinical care. Physiol Behav. 2018 Apr 1;187:2-5. doi: 10.1016/j.physbeh.2017.08.012. Epub 2017 Aug 17. PMID: 28823546.

68. Ordoñes Sanchez E, Bavley CC, Deutschmann AU, Carpenter R, Peterson DR, Karbalaei R, Flowers J 2nd, Rogers CM, Langrehr MG, Ardekani CS, Famularo ST, Bongiovanni AR, Knouse MC, Floresco SB, Briand LA, Wimmer ME, Bangasser DA. Early life adversity promotes resilience to opioid addiction-related phenotypes in male rats and sex-specific transcriptional changes. Proc Natl Acad Sci U S A. 2021 Feb 23;118(8):e2020173118. doi: 10.1073/pnas.2020173118. Erratum in: Proc Natl Acad Sci U S A. 2022 Apr 26;119(17):e2204210119. doi: 10.1073/pnas.2204210119. PMID: 33593913; PMCID: PMC7923376.

69. Sullivan EC, Hinde K, Mendoza SP, Capitanio JP. Cortisol concentrations in the milk of rhesus monkey mothers are associated with confident temperament in sons, but not daughters. Dev Psychobiol. 2011 Jan;53(1):96-104. doi: 10.1002/dev.20483. PMID: 20730788; PMCID: PMC3188439.

70. Hinde K, Skibiel AL, Foster AB, Del Rosso L, Mendoza SP, Capitanio JP. Cortisol in mother's milk across lactation reflects maternal life history and predicts infant temperament. Behav Ecol. 2015 Jan-Feb;26(1):269-281. doi: 10.1093/beheco/aru186. Epub 2014 Oct 31. PMID: 25713475; PMCID: PMC4309982.

71. Dettmer AM, Murphy AM, Guitarra D, Slonecker E, Suomi SJ, Rosenberg KL, Novak MA, Meyer JS, Hinde K. Cortisol in Neonatal Mother's Milk Predicts Later Infant Social and Cognitive Functioning in Rhesus Monkeys. Child Dev. 2018 Mar;89(2):525-538. doi: 10.1111/cdev.12783. Epub 2017 Mar 29. PMID: 28369689; PMCID: PMC6528802.

72. Lu L, Guo H, Peng Y, Xun G, Liu Y, Xiong Z, Tian D, Liu Y, Li W, Xu X, Zhao J, Hu Z, Xia K. Common and rare variants of the THBS1 gene associated with the risk for autism. Psychiatr Genet. 2014 Dec;24(6):235-40. doi: 10.1097/YPG.0000000000000054. PMID: 25304225.

73. Paketçi C, Ermiş Ç, Şişman AR, Hız S, Baykara B, Yiş U. Blood neurofilament light chain and thrombospondin-1 levels of patients with autism spectrum disorder. Turk J Med Sci. 2022 Aug;52(4):1041-1049. doi: 10.55730/1300-0144.5406. Epub 2022 Aug 10. PMID: 36326357; PMCID: PMC10388028.

74. Lin F, Jun Li, Ziqi Wang, Zhang T, Lu T, Jiang M, Yang K, Jia M, Zhang D, Wang L. Replication of previous autism-GWAS hits suggests the association between *NAA1, SORCS3,* and *GSDME* and autism in the Han Chinese population. Heliyon. 2023 Dec 22;10(1):e23677. doi: 10.1016/j.heliyon.2023.e23677. PMID: 38234914; PMCID: PMC10792458.

75. Bagni C, Zukin RS. A Synaptic Perspective of Fragile X Syndrome and Autism Spectrum Disorders. Neuron. 2019 Mar 20;101(6):1070-1088. doi: 10.1016/j.neuron.2019.02.041. PMID: 30897358; PMCID: PMC9628679.

76. Joo Y, Benavides DR. Local Protein Translation and RNA Processing of Synaptic Proteins in Autism Spectrum Disorder. Int J Mol Sci. 2021 Mar 10;22(6):2811. doi: 10.3390/ijms22062811. PMID: 33802132; PMCID: PMC8001067.

77. Zhu J, Liu H, Hu Y, Liu J, Dai C, Liang J, Cheng B, Tan M, Zhang Y, Cao Q, Lai X. Mechanistic insights into retinoic-acid treatment for autism in the improvement of social behavior: Evidence from a multi omics study in rats. Neuropharmacology. 2025 Mar 1;265:110244. doi: 10.1016/j.neuropharm.2024.110244. Epub 2024 Dec 4. PMID: 39643238.

78. Li JY, Daniels G, Wang J, Zhang X. TBL1XR1 in physiological and pathological states. Am J Clin Exp Urol. 2015 Apr 25;3(1):13-23. PMID: 26069883; PMCID: PMC4446378.

79. Mastrototaro G, Zaghi M, Massimino L, Moneta M, Mohammadi N, Banfi F, Bellini E, Indrigo M, Fagnocchi G, Bagliani A, Taverna S, Rohm M, Herzig S, Sessa A. TBL1XR1 Ensures Balanced Neural Development Through NCOR Complex-Mediated Regulation of the MAPK Pathway. Front Cell Dev Biol. 2021 Feb 23;9:641410. doi: 10.3389/fcell.2021.641410. PMID: 33708771; PMCID: PMC7940385.

80. Quan Y, Zhang Q, Chen M, Wu H, Ou J, Shen Y, Li K, Xun G, Zhao J, Hu Z, Xia K, Guo H. Genotype and Phenotype Correlations for TBL1XR1 in Neurodevelopmental Disorders. J Mol Neurosci. 2020 Dec;70(12):2085-2092. doi: 10.1007/s12031-020-01615-7. Epub 2020 Jun 10. PMID: 32524419.

81. Bale TL. Sex differences in prenatal epigenetic programming of stress pathways. Stress. 2011 Jul;14(4):348-56. doi: 10.3109/10253890.2011.586447. PMID: 21663536.

82. Kentner AC, Cryan JF, Brummelte S. Resilience priming: Translational models for understanding resiliency and adaptation to early life adversity. Dev Psychobiol. 2019 Apr;61(3):350-375. doi: 10.1002/dev.21775. Epub 2018 Oct 11. PMID: 30311210; PMCID: PMC6447439.

83. CDC [Centers for Disease Control]. 2025. Breastfeeding special circumstances: illness or conditions and breastfeeding: https://www.cdc.gov/breastfeeding-special-circumstances/hcp/illnesses-conditions/index.html; influenza: https://www.cdc.gov/breastfeeding-special-circumstances/hcp/illnesses-conditions/flu.html; COVID-19: https://www.cdc.gov/breastfeeding-special-circumstances/hcp/illnesses-conditions/flu.html; foodborne and waterborne illness: https://www.cdc.gov/breastfeeding-special-circumstances/hcp/illnesses-conditions/food-water-borne-illness.html

84. NICHD Inaugural stakeholder meeting for the prioritization of therapeutic research needs for pregnant, postpartum, and lactating (PPL) persons. 2024. Inaugural Stakeholder Meeting for the Prioritization of Therapeutic Research Needs for Pregnant, Postpartum, and Lactating (PPL) Persons Meeting Summary: https://www.nichd.nih.gov/sites/default/files/inline-files/PPL_Persons_Therapeutic_Needs_Mtng_Summ.pdf

85. Tseng PT, Chen YW, Stubbs B, Carvalho AF, Whiteley P, Tang CH, Yang WC, Chen TY, Li DJ, Chu CS, Yang WC, Liang HY, Wu CK, Yen CF, Lin PY. Maternal breastfeeding and autism spectrum disorder in children: A systematic review and meta-analysis. Nutr Neurosci. 2019 May;22(5):354-362. doi: 10.1080/1028415X.2017.1388598. Epub 2017 Oct 18. PMID: 29046132.

Figure 1

A DUC

Thaw milk on ice

3,000 x g - 10 min (2x)
1,200 x g - 10 min (2x)
21,500 x g - 30 min (2x)
21,500 x g - 1 hr

Filter the supernatant
0.45 μm syringe filter

Casein precipitation
with acetic acid +
4,500 x g - 30 min

Filter the supernatant
0.22 μm syringe filter

100,000 x g - 120 min (2x)

Re-suspend pellet in 1X PBS

B Nanoparticle tracking analysis: Size and Concentration

Results

Stats: Merged Data

Mean:	115.9 nm
Mode:	108.0 nm
SD:	31.5 nm
D10:	81.0 nm
D50:	111.4 nm
D90:	157.6 nm

C Western Blotting

Cellular Control   Rat MEVs

Alix   ~ 100kDa
Calnexin   ~80kDa
Flotillin-1   ~49kDa
CD9   ~28kDa

D Transmission Electron Microscopy (TEM)

88.1nm   70.5nm
126nm
77.3nm
92.2nm

200 nm

E MEVs Recovered

○ SD-Saline
● SD-LPS
□ EE-Saline
■ EE-LPS

* = SD-LPS versus SD-Saline
# = SD-LPS versus EE-LPS
a = main effect of drug
b = main effect of housing
c = main effect of time

aa

Figure 2

Figure 3

Figure 4

Figure 5

A Mammary Gland

Milk derived extracellular vesicles (MEVs) contain bioactive cargo such as lipids, proteins, miRNA, mRNA, and DNA.

B Mammary Gland

Stress

MEV cargo are altered following environmental insults like stress or inflammation.

C

MEV cargo are absorbed by nursing offspring, which then interact with gut microbiome and/or directly with brain

D

| Saline | Lactational MIA | |
| Standard Housing | Standard Housing | Environmental Enrichment |

Stress

small RNA NGS

sociability

Figure 6