The genetic architecture of G-protein coupled receptor signaling

Taylor L. Mighell[1] and Ben Lehner[1, 2, 3, 4, *]

[1]Centre for Genomic Regulation (CRG), The Barcelona Institute of Science and

Technology, Barcelona, Spain

[2]Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, United

Kingdom

[3]Universitat Pompeu Fabra (UPF), Barcelona, Spain

[4]Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain

14
15 *e-mail: bl11@sanger.ac.uk
16

Abstract

G-protein coupled receptors (GPCRs) are the most abundant class of

human receptors and drug targets. The vast majority of GPCR drugs bind the

conserved orthosteric pocket, which can lead to non-specificity and toxicity.

Precision modalities such as biased signaling and allosteric modulation could

lead to GPCR therapeutics with improved efficacy and safety profiles. The

rational design of precision therapies is, however, limited by the lack of high-

resolution functional understanding of GPCRs. Here we present a general

approach to chart high-resolution functional maps of GPCRs and apply the

approach to the β2 adrenergic receptor. Across ~150,000 measurements of five

different phenotypes, we construct an unprecedented picture of signaling

properties of this model receptor. A contiguous core activation network of

residues required for signaling links the ligand binding site to the transducer

binding site. Residues controlling signaling parameters connect to this core

network, with determinants of signaling bias located more peripherally. Dose-

response curves for nearly all possible amino acid substitutions, a total of

>7,500 different proteins, reveal the relationship between expression level and

potency; and the spatial distribution of potency changing residues. The multi-

modal maps rationalize previous serendipitously discovered allosteric sites and

prospectively annotate novel sites to target. Application of this approach across

the superfamily of GPCRs will unlock deep understanding of these critical

molecules and enable rational design of precision GPCR therapeutics.

Introduction

G-protein coupled receptors (GPCRs) are the largest family of

transmembrane proteins that receive and transmit signals from the outside to

the inside of cells. These signals are very diverse and include

neurotransmitters, metabolites, immune system molecules, pH, and photons of

light[1]. The topology of GPCRs is highly conserved and consists of seven

transmembrane (TM) helices connected by alternating intra- and extracellular

loops (ECLs and ICLs). Binding of an agonist to the extracellular portion of the

receptor results in an allosteric conformational change to the intracellular

portion of the receptor[2], which enables binding and activation of one or more

members of the family of heterotrimeric G-proteins, which subsequently activate

diverse downstream signaling pathways[3]. Receptor activation also results in

phosphorylation of the receptor C-terminal tail and recruitment of β-arrestin,

which terminates G-protein signaling by internalizing the receptor, and also

scaffolds G-protein independent signaling[4]. Because of their role as master

signal transducers and their availability on the cell surface, GPCRs are the most

important class of drug targets, with one third of clinical therapeutics targeting

GPCRs[5,6].

The vast majority of established GPCR drugs act by binding to the

orthosteric site and activating or inactivating the receptor[7]. However, alternative

modalities for modulating GPCRs exist, such as biased agonism[8] (also called

"functional selectivity") and allosteric modulation[9]. Biased agonists are defined

in opposition to balanced agonists; a biased agonist preferentially signals

through one downstream pathway over another (e.g. G-protein over β-arrestin),

compared to balanced agonists. In cases where signaling through one pathway

leads to an unwanted side effect, biased agonists may achieve improved

therapeutic efficacy. A therapeutic benefit for biased agonism has been

proposed for at least thirty GPCRs[10].

Allosteric drugs are those that bind the receptor at a spatially distinct site

from the orthosteric site. The high structural conservation of orthosteric sites

makes it difficult to target one receptor without also engaging multiple family

members, frequently causing toxicity and other side-effects that result in clinical

failure[7]. In contrast, allosteric drugs targeting non-conserved sites can have

high specificity and reduced toxicity. Further, since allosteric drugs modulate

responses to physiological ligands, their effects temporally and spatially match

physiological signaling processes[11]. Due to these major advantages, increasing

effort is being put toward development of biased and allosteric modulators[12].

However, with our current understanding of GPCR function, these molecules

are typically only found by blind screening or iteration upon previously

discovered molecules[13,14].

In order to design better GPCR drugs, an improved understanding of the

rich functional and allosteric architecture linking agonist binding with intracellular

signaling is required. Much work has focused on using the structures of active

and inactive states of receptors to try and understand this allosteric

mechanism[15,16]. This has provided important mechanistic insights, but these

structures are only the starting point for assigning a functional role for the

hundreds of individual amino acids in each highly dynamic receptor.

A powerful complementary approach for understanding how proteins

work is the use of mutagenesis followed by functional characterization[17]. Serial

experiments have measured the effects of hundreds of alanine substitutions on

GPCR signaling through one[18] or two[19,20] pathways, revealing a rich diversity of

signaling responses to individual mutations. Massively multiplexed approaches

have been used to characterize surface expression[21–24], ligand binding[23,25], and

signaling of human GPCRs[26–28]. However, these approaches to quantifying

signaling relied on linking transcriptional reporters to endogenous signaling

pathways, which can be non-linear and subject to feedback circuits, adding

noise to measurements. Additionally, these systems have only been

established for G-protein mediated signaling pathways. Finally, these

multiplexed studies were not extended to formal dose-response analysis.

We construct and deploy here GPCR-MAPS (GPCR multiplexed assays

with pathway specificity), an experimental and computational framework for

generating comprehensive functional maps of GPCRs. We used the beta-2

adrenergic receptor (β2AR) as a model receptor. This extensively studied

receptor is expressed in the cardiovascular system where it modulates smooth

muscle relaxation and bronchodilation; β2AR agonists are therefore used for

treating asthma as well as chronic obstructive pulmonary disease[29]. First, we

measure the surface expression level and signaling response of all possible

variants in the receptor to saturating concentration of agonist through G-protein

and β-arrestin pathways. We model the relationship between expression and

signaling to account for indirect effects of changes in expression on signaling

and identify a structurally contiguous network of residues linking the ligand

binding site to the transducer binding site. By directly comparing G-protein and

β-arrestin signaling, we then identify residues involved in biased signaling

enriched at the extracellular loops (also known as the "extracellular vestibule")

and near the conserved NPxxY motif in the 7[th] transmembrane (TM) helix. We

then extend the platform to enable, for the first time, massively multiplexed

dose-response analysis for GPCRs. A 12-point dose-response experiment,

encompassing nearly 100,000 measurements, enables highly accurate

quantification of the dose-response curves of 5,588 different receptor variants.

The data provide numerous functional insights, including identifying intracellular

loop 3 as a hotspot for mutations that increase potency. Finally, we use the rich

data generated in this study to explain the allosteric activity of known

modulators and prospectively annotate new structural pockets to therapeutically

target. Application of the GPCR-MAPS platform across the important

superfamily of GPCRs will advance our understanding of these proteins and

enable rational, precision drug design.

Results

GPCR-MAPS: multiplexed, multimodal assay with pathway specificity

An ideal platform for measuring GPCR signaling activity at scale would

enable the readout of the two canonical signaling pathways (G-protein- and β-

arrestin-mediated) and would rely on a signal detector proximal to the receptor,

thereby avoiding feedback or non-linearity of intracellular signaling pathways. A

previously described experimental platform[30] uses three genetic constructs to

report on GPCR signaling: 1) a GPCR of interest with a fusion of the exogenous

transcription factor Gal4, and linked by sequence encoding an evolved light,

oxygen, and voltage sensitive (eLOV) domain as well as tobacco etch virus

protease cleavage site (TEVcs), 2) a GPCR binding partner (e.g. β-arrestin)

fused to NanoLuciferase (NLuc) and tobacco etch virus protease (TEV), and 3)

the fluorescent protein Citrine under the control of the UAS promoter. When co-

expressed in cells, agonist-mediated activation of the GPCR causes recruitment

of the transducer construct to the receptor. In the absence of light, the eLOV

domain cages the TEVcs and prevents cleavage; however, co-addition of the

NLuc substrate furimazine results in bioluminescence, and when the transducer

brings NLuc into close proximity with the eLOV domain, the steric hindrance is

relieved and TEVp can cleave at TEVcs. This cleavage results in liberation of

Gal4, which can move to the nucleus and drive transcription of the UAS-Citrine

reporter construct.

While this assay faithfully reports on GPCR activation, it is limited to

serial experimental designs and has only been demonstrated with one

downstream signaling partner. Therefore, we extensively re-engineered the

system to enable massively multiplexed experiments. We combined the GPCR

construct and the reporter construct onto a single plasmid and added a DNA

barcode (26 nucleotides) in the 3' untranslated region (UTR) of the Citrine

reporter so that the GPCR variant on any given plasmid can be identified

(Figure 1a and Supplementary Figure 1d). We also added the attB

recombination site to enable integration into HEK293T landing pad cells[31],

which ensures copy number of exactly one in each cell. We used the active-

state-specific nanobody Nb80[32] as the binding partner to optimize the assay.

While the original assay used transient transfection to introduce transgenic

material and took place over the course of minutes, we found that substantially

increased times were necessary, presumably due to the reduction of copy

number of the receptor and reporter constructs (24 hours, Supplementary

Figure 1a). Further, we tested evolved variants of both TEV and the LOV

domain, uTEV1Δ[33] and hLOV1[34]; the combination that yielded the highest

signal to noise ratio was uTEV1Δ and eLOV (Supplementary Figure 1b). Next,

we also validated the use of β-arrestin and miniG$_s$[35] as the binding partner in

this system (Supplementary Figure 1b). Finally, in order to account for

differences in barcode abundance in the library, as well as barcode-specific

effects on RNA abundance, we also include a condition in which Gal4 is

expressed alone (Supplementary Figure 1d-e); this condition should maximally

stimulate all barcodes and provides normalization for the experimental

conditions[26]. While this platform enables multiplexed measurement of GPCR

signaling, it is expected that many variants that exhibit decreased signaling will

do so because of a lack of surface expression[27]. We therefore implemented in

parallel a FACS-based method to measure the surface expression of library

members[36] (Figure 1a, Materials and Methods).

Surface expression and signaling through both canonical pathways

We generated a saturation mutagenesis library of β2AR in a vector also

containing: an N-terminal HA epitope tag and C-terminal fusions of eLOV,

TEVcs, and Gal4, and additionally the UAS-Citrine module (attB-HA-b2AR-

eLOV-TEVcs-Gal4-UAS-Citrine, Supplementary Figure 1d) including all

missense and nonsense substitutions, synonymous mutations where possible

(i.e. the encoded amino acid can be encoded by alternate codons), and single

residue deletions. We then introduced short degenerate nucleotide barcodes

into the 3' UTR of Citrine and used long read sequencing to make 212,840

β2AR genotype-barcode links (Supplementary Figure 1c, Materials and

Methods). These represented 8,970 out of an intended 9,046 unique genotypes

(99.2%).

We introduced this library into HEK293T landing pad cells[31], ensuring a

single variant is expressed per cell. Then, we performed a FACS-based

multiplexed surface expression experiment. Namely, cells were immunostained

with anti-HA antibody, then sorted into four bins based on fluorescence intensity

(Supplementary Figure 1d). DNA from the cells sorted into each bin was purified

and converted into sequencing libraries, and the barcodes were sequenced with

short-read sequencing (Figure 1a, Materials and Methods).

All counts from barcodes representing the same variant were tallied and

combined with the fluorescence value of each FACS bin to arrive at surface

expression estimates. Variants with low frequency in the library were filtered to

yield 8,740 high confidence surface expression estimates (Materials and

Methods, Supplementary Table 1). These measurements were highly

reproducible across three replicates (Pearson r=0.94-0.96, Figure 1c). Scores

were normalized between the median of premature stop codons, set to 0, and

the wild-type surface expression level, set to 1. A heatmap representation of the

data shows nonsense mutations are uniformly damaging, except those

occurring within intracellular loop 1 (ICL1) and those in the C-terminus,

suggesting the receptor can be stably expressed as either a single-pass

transmembrane protein, or as a C-terminal tail truncation. The transmembrane

helices are more mutation sensitive than extra- or intra-cellular loops, consistent

with previous work[27,36] (Figure 1d). Synonymous wild-type and nonsense

variants have well separated expression distributions, with missense variants

spanning the full range between those two classes (Figure 1e).

Next, we performed the GPCR-MAPS signaling assay using Nb80 (as a

surrogate for G-protein, as Nb80 returns higher signal to noise than miniG$_s$

(Supplementary Figure 1b)) as well as β-arrestin as readouts for the two

canonical GPCR signaling pathways. Namely, we transiently transfected one of

the transducer constructs (pUC19-NLuc-[Nb80/βArr]-uTEV1Δ) or the Gal4

normalization construct (pUC19-CMV-Gal4, Supplementary Figure 1e) into cells

bearing the β2AR receptor construct. The next day, saturating concentration of

isoprenaline (10 µM) was added to the transducer samples to induce signaling,

and 24 hours later, those cells were collected (Figure 1b, Supplementary Figure

1f). The Gal4 maximum induction samples were collected 30 hours after Gal4

transfection. RNA was collected from all samples, reverse transcribed,

converted to short-read sequencing libraries and sequenced (Supplementary

Figure 1f-g, Materials and Methods).

Signaling scores were calculated using the variant frequency in the

experimental samples compared to variant frequency in the Gal4 condition and

re-scaled with 0 representing no signaling activity and 1 representing wild-type

signaling. Low frequency variants were filtered to yield 7,850 high confidence

Nb80 and β-arrestin signaling scores (Materials and Methods, Supplementary

Table 1). Replicates were well correlated (Pearson's r=0.9-0.91 for Nb80

replicates, r=0.88-0.89 for β-arrestin replicates, Figure 1c). Heatmap

representations of the data reveal similar overall trends between Nb80 and β-

arrestin (as well as with the surface expression data) defined by

transmembrane regions being most sensitive to mutation (Figure 1 f, h).

Synonymous wild-type variants are again well separated from nonsense

variants, and missense variants again span the whole distribution from

nonsense to wild-type-like (Figure 1 g, i). The Nb80 and β-arrestin signaling

scores are correlated with each other (Spearman's ρ = 0.87), as well as with the

surface expression scores (ρ=0.55, 0.6, respectively, Supplementary Figure

1h), consistent with changes in expression being a very important cause of

changes in signaling.

While the overall patterns of mutational tolerance are similar between

surface expression and signaling datasets, there are positions with stark

differences. For example, missense mutations at the eight positions making up

the isoprenaline binding pocket have minimal effect on stability (Figure 1j), while

these same mutations are strongly deleterious for signaling through both

binding partners (Figure 1k-l).

Residues contributing to signaling beyond expression changes

While the effects of mutations substantially manifest through expression

changes, we sought to identify mutation effects that specifically affect signaling.

So, we directly compared signaling and surface expression scores of missense

variants. These demonstrate a non-linear, correlated relationship (ρ = 0.55, 0.6

for surface expression compared with Nb80 and β-arrestin, respectively, Figure

2a-b). We fit a LOWESS regression to the surface expression and signaling

data, which acts as a null expectation of signaling level produced by a given

surface expression level. This null expectation explains a significant amount of

variance (R^2 = 0.37 and 0.42 for Nb80 and β-arrestin, respectively).

We next calculated signaling residuals and subsequently z-scores of all

variants to the global model (i.e. residual divided by standard error of the mean

for each variant, Figure 2c-e). Comparing the group of z-scores at each position

with the z-scores from all other positions with a two-sided Mann-Whitney U

allows us to identify residues where mutations have greater or lesser signaling

activity than would be expected based on their surface expression level (Figure

2e, Supplementary Table 2).

272   For both Nb80 and β-arrestin, more residues confer loss-of-function

(LoF) than gain-of-function (GoF) when mutated, and the magnitude of loss is

greater than the magnitude of gain (Figure 2 f-g). In total, 49 and 73 residues

are LoF for Nb80 and β-arrestin, respectively (false discovery rate, FDR = 0.1,

Supplementary Table 2). Of these, 47 are shared (Figure 2h). An example of a

strong LoF residue is D113^{3x32} (Figure 2i, GPCRdb numbering[37] in superscript),

which participates directly in ligand binding. There are 31 and 53 GoF residues

for Nb80 and β-arrestin (FDR=0.1), respectively, of which 20 are shared.

E268^{6x30} is the most significant GoF (Figure 2j), consistent with previous work

demonstrating constitutively activating mutations at this site[38]. Finally, we

extended this analysis to highlight residues with highly variable effects on

signaling. Namely, after excluding the LoF and GoF residues, we compared the

interquartile range (IQR) of z-scores at all positions (Supplementary Figure 2a-

b). We categorized those residues with IQR above the 90th percentile as

change-of-function (CoF): 34 and 31 residues for Nb80 and β-arrestin,

respectively, with 8 shared (Figure 2h). An example of a strong CoF residue is

S120^{3x39} (Figure 2k). This residue repacks upon β2AR activation[18] and is

thought to modulate the conformation of helix 7[39].

290   We prioritized for further analysis residues that were identified in the

analyses of both transducers. We note that average position z-scores and

position IQRs are well correlated (r = 0.73 and 0.56, respectively,

Supplementary Figure 2c-d), indicating that many residues contribute similarly

to signaling through both pathways. Plotting the residues identified for both

transducers onto the structure of isoprenaline-stimulated β2AR (PDB ID: 7DHR)

reveals that LoF residues are predominantly within the core of the receptor

(Figure 2l), whereas GoF and CoF residues are dispersed on the periphery

(Figure 2m-n).

300   Structural basis of loss-, gain-, and change-of-function residues

Next, we analyzed in more detail the identified functional residues. We

examined the LoF residues in an inactive (PDB ID: 2RH1) and an active (PDB

ID: 3SN6) structure of β2AR. Strikingly, the majority of these residues (39 out of

47) form an interconnected path linking the ligand binding site all the way to the

transducer binding site in the active state (Figure 3a, hereafter referred to as the

"core activation network"). All 8 isoprenaline-binding residues are identified

(D113^{3x32}, V117^{3x36}, F193^{45x52}, S203^{5x43}, S207^{5x461}, F290^{6x52}, N293^{6x55}, and

N312^{7x38}) along with many residues in the second shell of the ligand binding

pocket (i.e. residues that contact residues that contact isoprenaline, Figure 3b).

These residues include W286^{6x48} of the CWxP motif and I121^{3x40} of the PIF

motif[40,41]. The core activation network then connects to N322^{7x49} of the critical

NPxxY motif[40] via two pathways: first, W286^{6x48} contacts N318^{7x45} which then

connects to N322^{7x49}. Second, a pathway from A119^{3x38} goes through A78^{2x49}

and D79^{2x50} of the allosteric sodium binding pocket which then contacts

N322^{7x49}. Of the NPxxY motif, N322^{7x49}, I325^{7x52}, Y326^{7x53} are identified as LoF.

Finally, in the active state structure, Y326^{7x53} of the NPxxY motif then makes

contact with R131^{3x50} (of the DRY motif) and I127^{3x46} of helix 3. Y326^{7x53} is also

thought to make an important water-mediated hydrogen bond with Y219$^{5.58}$,[42]

which is also identified as LoF and makes a direct contact with R131^{3x50}.

Of the eight LoF residues not connected to the core activation network,

three likely directly disrupt transducer binding[32,43] (P138^{34x50}, F139^{34x51}, and

Y141^{34x53}), while three more take part in critical disulfide bonds between ECL2

and TM3 (C106^{3x25}, C190^{ECL2}, C191^{45x50}). F193^{45x52} is a disconnected LoF

residue that makes direct contact with bound isoprenaline. Finally, P88^{2x58} in

TM2 is the last disconnected residue.

We then overlaid the GoF residues onto the inactive and active

structures (Supplementary Figure 3a). Two of the GoF residues, D130^{3x49} and

E268^{6x30}, participate in the highly conserved ionic lock (with R131^{3x50} of the

DRY motif[44]). While R131^{3x50} of the DRY motif is a LoF residue that contacts

Nb80[32] and is predicted to contact β-arrestin,[43] mutations at D130^{3x49} of the

DRY motif have been shown to constitutively activate the receptor[45]. The other

GoF residues are predominantly in two clusters: one consisting of TMs 3, 5, and

6, near the NPxxY and DRY motifs (Supplementary Figure 3b). L124 sits in the

center of the receptor, near the I of PIF motif, DRY motif, and NPxxY. Mutations

at L124^{3x43} are known to cause constitutive activity[46]. The other cluster is near

the ligand binding site and in the extracellular vestibule (Supplementary Figure

3c). E187 and T189 are in ECL2, where they may affect ligand binding or

activation[47]. W313^{7x39} and V87^{2x57} form two van der waals interactions in the

inactive (PDB:2RH1), but only one in the active state (PDB:3SN6,

GetContacts); our data suggest that disrupting this interaction leads to

increased signaling. Finally, there are 7 GoF residues in the C-terminal tail

(Supplementary Figure 3g), consistent with previous findings[26].

Plotting the CoF residues onto the inactive and active state structures

(Supplementary Figure 3d) reveals that they cluster near important LoF

residues that make structural changes during activation: for instance, $S120^{3x39}$

and $E122^{3x41}$ are adjacent to $I121^{3x40}$ of the dynamic PIF motif. Likewise,

$F321^{7x48}$ is adjacent to $N322^{7x49}$ of the dynamic NPxxY motif and identified LoF,

while $F332^{8x50}$ is nearby (Supplementary Figure 3e). This proximity to essential

switch residues suggests that CoF residues may specifically stabilize or

destabilize the active or inactive receptor state depending on the amino acid

substitution.

Genetic basis of biased signaling

Next, we sought to directly compare mutational effects on Nb80 and β-

arrestin signaling to understand the molecular basis of biased signaling (Figure

4a). Missense mutational effects on signaling through Nb80 and β-arrestin are

well correlated (Pearson's r=0.89, Figure 4b). However, single mutants

previously shown to bias the receptor toward G-protein signaling (F193A[48],

Y219A[49], and Y199A[50]) are indeed biased towards Nb80 in our experiments

(Figure 4c). To find residues with stronger effects on one pathway compared to

the other, we implemented a similar strategy as before: we fit a LOWESS

regression curve to the data, and calculated residuals and subsequently z-

scores (Figure 4c) for all variants in the dataset. Then, we compared z-scores of

mutations at each position with z-scores of mutations at all other positions. This

identifies 19 residues where mutations have significantly greater effects on β-

arrestin signaling and 7 positions where mutations have significantly greater

effects on Nb80 signaling (FDR=0.1, Figure 4d, Supplementary Table 3). As

before, we then compared the IQR of all remaining residues; here, to more

closely match the number of biased residues, we prioritized the top 5% of

residues as bias switches (Supplementary Figure 4a).

Examples of strongly biased residues are shown (Figure 4e-f) as well as

a strong bias switching residue (Figure 4g). One expectation is that mutations at

residues making direct contacts with either Nb80 or β-arrestin could result in

biased signaling effects. Nb80 and β-arrestin have overlapping but distinct

binding interactions with the intracellular portion of β2AR. Nb80 forms direct

contacts with 16 residues in TMs 3, 5, 6, and 7[32]. Since there is no

experimentally determined structure of β2AR bound by β-arrestin, we defined

this interface from a simulation study[43]. The β-arrestin binding interface is

predicted to be larger, encompassing 28 residues across ICLs 1, 2, 3, and TMs

6 and 7, with 10 residues overlapping the Nb80 interface.

Three out of the seven residues biased toward β-arrestin make direct

binding interactions with both Nb80 in the solved structure and with β-arrestin in

the simulation (P138^{34x50}, F139^{34x51}, and Y326^{7x53}), suggesting a more critical

role for these residues in binding Nb80. The other four β-arrestin-biased

residues are either in ICL3 (G252 and G255) or the C-terminal tail (E373 and

C406, Supplementary Figure 4b). The only residue biased toward Nb80 within

either binding interface is P330^{8x48}, which is predicted to interact with Nb80 but

not β-arrestin. There is also one bias switch residue, R328^{7x55}, that is predicted

to bind β-arrestin. This relative sparsity of bias effects suggests that few

residues at the transducer interface contribute significantly more to signaling

through either pathway, at least for these two transducers (Supplementary

Figure 4c).

In contrast, there are many residues biased toward Nb80 located in the

extracellular vestibule: W99^{23x50}, F101^{23x52}, G102^{3x21}, W173^{ECL2}, A181^{ECL2},

C184^{ECL2}, Y185^{ECL2}, N196^{5x36}, and E306^{7x32} (Figure 4h, k). The most strongly

Nb80 biased residue, W99^{23x50}, was previously identified as particularly

mutationally intolerant[26]; however only by measuring both canonical signaling

pathways could we uncover its strongly biased effects (Figure 4e). Considered

together, the extracellular loops (ECLs) are enriched in Nb80 biased residues

compared to the whole receptor (OR=6.4, p=1.0x10^{-3}, one-sided Fisher exact

test), and in particular, ECL2 is enriched for Nb80 biased residues (OR=5.6,

p=0.014, one-sided Fisher exact test).

The importance of the extracellular vestibule for G-protein biased

signaling revealed in our mutational data is also consistent with structural data.

Salmeterol is a G-protein biased β2AR agonist that binds the orthosteric site in

a similar way as the endogenous ligand (epinephrine) except that it has a long

aryloxyalkyl tail that extends into the extracellular vestibule, where it interacts

extensively with ECL2[51]. Also, comparative structural analyses of other GPCRs

have suggested that the extracellular vestibule adopts a more constricted

conformation in the β-arrestin-bound state compared to the G-protein-bound

state.[52] Taken together, these data point to the extracellular vestibule, and in

particular ECL2, as hotspots for G-protein-biased agonism at the β2AR.

Among the bias switching residues, the second strongest is the vestibule

residue, Y308^{7x34} (Supplementary Figure 3d). This residue has been proposed

to be necessary for G-protein signaling[53], as well as forming a hydrophobic gate

with F193^{45x52} that interacts with and dehydrates ligand molecules as they

traverse the vestibule and approach the orthosteric binding site[54]. In the active

state, these two residues form a "lid" over the ligand, with F193^{45x52} making a

direct contact with isoprenaline (Figure 4i). Based on the alanine mutation,

F193^{45x52} was postulated to coordinate β-arrestin signaling[48]. Interestingly, while

this single variant does have a stronger effect on β-arrestin signaling, other

variants at the same site have the opposite effect (Supplementary Figure 4e).

We examined the effects of mutations at these two sites and found that

signaling bias is correlated with mutant hydrophobicity for Y308^{7x34} (ρ = 0.67, p

= 4.3x10^{-3}, Figure 4j), but not for F193^{45x52} (ρ = -0.36, p = 0.16, Supplementary

Figure 4f). This suggests that the role of Y308^{7x34} in determining bias is directly

tied to its hydrophobicity, while the relationship of F193^{45x52} to bias is more

nuanced.

Besides the vestibule, there are multiple bias-functional residues (Nb80-

or β-arrestin-biased or bias switches) in or near the NPxxY motif of TM7(Figure

4l). In fact, TM7 is enriched for bias-functional residues compared to the whole

receptor (OR=3.7, p=0.011, one-sided Fisher exact test, Supplementary Figure

4g). P323^{7x50} is not predicted to contact directly β-arrestin[43], suggesting that the

Nb80 bias of the residue cannot be explained simply by disrupting a direct

contact. N318^{7x45}, F332^{8x50}, and R328^{7x55} are all bias switches in close proximity

to the NPxxY motif (Figure 4k). A role for TM7 in determining bias is consistent

with prior nuclear magnetic resonance experiments demonstrating changes in

TM7 conformation upon binding of biased agonists[55].

Quantifying >7,500 dose-response curves

The data generated thus far represents β2AR signaling with saturating

levels of isoprenaline, which convolutes the contribution of different

pharmacological parameters, such as potency (the amount of agonist needed to

elicit a certain response) and efficacy (the maximum signaling response elicited

by saturating concentrations of agonist). Therefore, we built a platform for

measuring full dose-response curves for thousands of β2AR variants in parallel.

First, we performed the GPCR-MAPS signaling assay with 12 serially diluted

concentrations of isoprenaline, ranging from 10 pM to 10 µM (Figure 5a). In

order to compare the magnitude of signaling across experimental conditions,

following RNA isolation, we performed quantitative PCR (qPCR) to estimate the

total amount of reporter transcription (Figure 5b, Supplementary Figure 5a,

Materials and Methods). Then, by deep sequencing the barcodes and

normalizing the frequencies by the estimated total reporter RNA, we arrived at

normalized scores across the 12 concentrations (Figure 5c and Supplementary

Figure 5b). This allowed us to quantify dose-response relationships for 7,680

different proteins, increasing data of this type for β2AR or any GPCR >10-fold[18–

20].

Heatmap and correlation representations of the data reveals induction of

signaling across the ~92,000 measurements (Figure 5d and Supplementary

Figure 5c). Clustering the induction dataset revealed three predominant

clusters- these can be assigned as wild-type-like, moderately damaging, and

strong loss-of-function (Figure 5e).

Next, we fit a four-parameter Hill curve to the wild-type genotype,

normalizing E_{min} and E_{max} to 0 and 1, respectively, and deriving a half maximal

response estimate of 26.2 nM and slope (Hill coefficient) of 0.84 (Figure 5f).

Then we fit curves to all variants; after quality control (Supplementary Figure

5e-f, Materials and Methods), this resulted in 5,588 high quality dose response

curves with well-estimated parameters (Figure 5. g-I). We note that variants with

reduced signaling level are more poorly fit and therefore likelier to be filtered.

Genetic determinants of potency

We sought to use the fitted dose response curves to understand the

contribution of residues to potency, quantified as the amount of agonist required

to reach a half maximal response (EC_{50}, Figure 6a). To validate our EC_{50}

measurements, we compared them to a set of individually assayed alanine

substitutions[18]; these two datasets are well correlated (Pearson r=0.74, n=317

variants, Figure 6b). Next, we examined correlations between the parameters of

the dose-response curves, as well as their dependence upon surface

expression. Expression level has been predicted to be an important influence

on signaling parameters in theoretical work[56–58], and by experimental work on a

plant hormone receptor[59]. The strongest relationships we found were between

slope and E_{max} (ρ = -0.46) and between surface expression and EC_{50} (ρ = -0.41,

Figure 6c-d). Surface expression level is thus an important influence on potency

in the β2AR, consistent with theoretical predictions[56–58]. In order to isolate

mutational effects on EC_{50} beyond those caused by changes in expression we

again used LOWESS regression to model the relationship between surface

expression and EC_{50} (Supplementary Figure 6a) and used residuals to that

model to identify positions with consistently higher or lower residuals than

expected. 29 residues were identified as GoF (i.e. they had EC_{50} lower than

expected) and 59 were identified as LoF (i.e. higher EC_{50} than expected,

FDR=0.1, Figure 6e). There is substantial overlap between these sets and the

sets identified from the saturating concentration experiment (23/59 LoF shared,

4/29 GoF shared). Finally, the interquartile range was used again to identify 29

EC_{50} CoF residues (one shared with saturating concentration experiment,

Supplementary Figure 5b, Supplementary Table 4).

In principle, mutations could affect potency beyond changes in

expression by changing the affinity of isoprenaline binding, or by affecting the

probability of isoprenaline-bound receptor to adopt the active conformation[60].

While mutations at residues making direct contact with isoprenaline would be

expected to reduce affinity, mutations at these sites mostly totally abolish

signaling; dose-response curves could only be fit to 32 out of 132 missense

variants at these sites. However, these 32 variants do have over an order of

magnitude reduced potency (median EC_{50}=477 nM, compared to wild-type 26.2

nM). Interestingly, EC_{50} LoF residues are enriched in the second shell (15/59

LoF sites (25.4%, OR=4.12, $p=1.9 \times 10^{-4}$, Fisher exact test). While not

statistically enriched, a higher fraction of EC_{50} CoF sites (5/24, 20.8%) are in the

second shell compared to only 1 out of 29 GoF residues (3.4%) in the second

shell (Figure 6i). This shows that whereas mutations in residues directly

contacting the ligand have very strong effects on affinity and so completely

inactivate signaling, mutations in second shell residues frequently have more

moderate effects and modulate the response. These effects are, by definition,

allosteric, either indirectly altering ligand binding affinity or altering the

consequences of ligand engagement. This also suggests that mutations in the

second shell are likelier to reduce affinity than increase it.

Beyond changes in ligand affinity mutations can also affect signaling by

altering the likelihood of ligand-bound receptor adopting the active

conformation.[60] One way to disrupt the activation pathway could be for

mutations to interfere with the core activation network identified in the maximum

signaling experiment. We therefore quantified the distance of EC_{50}-affecting

residues to this network and found that LoF and CoF residues are closer in 3D

space than GoF residues (p=7.4x10^{-4}, 1.3x10^{-3}, respectively, Mann Whitney U

test, Figure 6j). Taken together, as with the maximum signaling analysis, these

data suggest that mutations that reduce potency do so by perturbing the core

activation pathway whereas the mechanism-of-action of GoF mutations is likely

distinct.

Three of the four most significant EC_{50} GoF residues are E268^{6x30}

(Figure 6g), D130^{3x49}, and L272^{6x34}. E268^{6x30} and D130^{3x49} were already

identified in the maximum signaling analysis and make up the conserved ionic

lock while L272^{6x34} is in close proximity to the ionic lock, where mutations are

also known to destabilize the inactive state.[60] Nearby E268^{6x30} in intracellular

loop 3 (ICL3), there are nine EC_{50} GoF residues, which is enriched compared to

the whole receptor (OR=3.38, p=7.3x10^{-3}, one-sided Fisher exact test). This

loop connects TM5 and TM6, which make dramatic movements upon receptor

activation (Figure 6k). These GoF residues could act in a similar way as

D130^{3x49}, E268^{6x30}, and L272^{6x34}, weakening the interaction between TM3 and

TM6 or other helices. Alternatively, previous studies suggest that ICL3 may

have an autoinhibitory role that sterically occludes transducer binding in the

inactive state, and that some mutations can abrogate this autoinhibitory effect[61].

Further experiments would be necessary to determine the exact mechanism of

the EC_{50} GoF phenotypes in ICL3. Additionally, there are two GoF residues in

the N-terminus that coincide with the two N-glycosylation sites[62] (N5 and N15,

Supplementary Figure 6d) suggesting that loss of these glycosylation

modifications can increase receptor potency.

F282^{6x44} of the PIF motif is the strongest EC_{50} CoF residue consistent

with previous data showing multiple constitutive activating mutations and at

least one attenuating mutation.[63] Likewise, S120^{3x39} and L212^{5x51} are identified

as CoF and are directly adjacent to the other two members of the PIF motif

(I121^{3x40} and P211^{5x50}) reinforcing that some perturbations to this key switch

motif can result in gains or loss of signaling. There are also several EC_{50} CoF

residues in TMs 5 and 6 (4 in each, Supplementary Figure 6f), reflecting the key

role of these helices in activation.

Rationalization of known allosteric modulators and prospective pocket

prioritization

We next brought to bear our rich datasets to rationalize the activity of

known allosteric modulators and prospectively prioritize new pockets for

therapeutic discovery. Cmpd-15 is a negative allosteric modulator (NAM) but its

mechanism is likely simple steric interference with transducer binding.[64] AS408

is a NAM that binds away from the orthosteric and transducer binding sites,

between TMs 3 and 5[65] (Figure 7a). Across the three analyses (maximum

signaling, biased signaling, and EC_{50}), the 7 residues making up this pocket

have 7 total annotations, compared to 96 annotations for all 220 structurally

resolved surface exposed residues (>10% relative solvent accessible surface

area in 2RH1), indicating a functionally important pocket. In particular, V129^{3x48}

is LoF for maximum signaling as well as EC_{50}, while E122^{3x41} is CoF for

maximum signaling as well as EC_{50} (Figure 7b-c). In addition, C125^{3x44} is CoF

for EC_{50} and $I214^{5x53}$ is LoF for EC_{50}. Mutations at $P211^{5x50}$ are biased toward

Nb80 signaling, though AS408 is not thought to be a biased NAM.

While the AS408 pocket is replete with functional residues, Cmpd-6, a

positive allosteric modulator[66] (PAM), binds in a pocket formed by 9 residues

between TMs 2, 3, 4, and ICL2 (Figure 7d) with only 4 annotations. However,

one of the residues is $D130^{3x49}$, where mutations cause strong GoF for both

maximum signaling and EC_{50} (Figure 7e-f), potentially explaining the PAM

activity of Cmpd-6. Additionally, while Cpmd-6 mostly forms hydrophobic

interactions with β2AR, it does form one hydrogen bond with $K149^{4x41}$, which is

an EC_{50} CoF residue (Figure 7e-f). Taken together, our functional annotations

are consistent with the activities of the two known β2AR allosteric modulators.

To explore prospective functional annotation, we used Fpocket[67] to

predict pockets in the inactive β2AR (PDB ID:2RH1). One additional pocket,

between TMs 5 and 6 (Figure 7g), has 8 functional annotations among 10

residues, with most being maximum signaling or EC_{50} LoF or CoF, but also one

GoF residue (Figure 7h-i), suggesting a highly active pocket. Indeed, an

allosteric agonist was recently found to bind and activate another GPCR,

GPR88, at this same site[68]. On the other hand, a nearby and partially

overlapping pocket also between TMs 5 and 6 (Supplementary Figure 7a) has

only two annotations among 9 residues (Supplementary Figure 7b-c),

deprioritizing this pocket as a therapeutic target.

Discussion

GPCRs are the most successful class of drug targets, yet nearly all

GPCR drugs act orthosterically, which often comes with severe drawbacks

related to specificity and toxicity[7]. Precision modalities such as biased signaling

and allosteric modulation could lead to GPCR drugs with improved efficacy and

safety, but these drugs have historically only been found by serendipity. Despite

a revolution in GPCR structural biology, we lack sufficient functional

understanding of GPCR signaling to design biased or allosteric drugs. To

address that shortcoming, we have developed and implemented GPCR-MAPS,

a new framework for high resolution functional mapping of GPCRs.

Our method makes multiple technical advancements over existing

technologies for multiplexed GPCR assessment[26–28]. First, our approach

detects receptor activation via protein-protein interaction, instead of via

endogenous signaling pathways. This reduces experimental noise and enables

more flexible experimental design. Second, GPCR-MAPS is highly modular - we

demonstrate here the successful use of a nanobody, a mini-G protein, and β-

arrestin as signal transducers. The receptor-proximal readout also facilitates

comparisons across these signal transducers, which can otherwise be

confounded by differential signal amplification and crosstalk[69]. Validation of

additional transducers will continue to enrich the functional readouts possible

with the method. To demonstrate generalizability, we also show the vasopressin

2 receptor signaling through miniGs and β-arrestin in response to the

endogenous ligand arginine vasopressin (Supplementary Figure 6e). Finally, we

extended the platform to enable massively multiplexed GPCR dose-response

modeling for the first time (Supplementary Figure 7).

Across >150,000 measurements, we have charted here the first

complete map of how mutations affect diverse signaling properties of a human

GPCR. After normalizing for expression, residues that specifically disrupt

signaling form a contiguously connected core activation network of residues

linking the ligand binding site to the transducer coupling site (Figure 8a).

Residues important for biased signaling are enriched in the extracellular

vestibule, particularly ECL2, as well as TM7 near the NPxxY motif (Figure 8b).

>7,500 dose response curves enabled accurate EC_{50} estimation of >5,500

unique receptors. EC_{50} is related to expression level, validating theoretical

models[56–58] and agreeing with recent work on plant hormone receptors[59].

Adjusting for expression level, there is a strong enrichment of residues

decreasing potency in the second shell of isoprenaline binding, and residues

increasing potency in ICL3 (Figure 8c). Residues that decrease potency are in

closer proximity to the core activation network and so likely change potency by

directly affecting the core activation network. In contrast, residues increasing

potency are far from the core activation network, likely affecting potency via

mechanisms independent of the core activation network. This distinction in the

location and likely mechanism of LoF and GoF variants parallels observations in

the Src kinase, where mutations that decrease kinase activity are enriched

close to the active site but mutations that increase it are not[70].

We believe that GPCR-MAPS – and related approaches[18–20,27,28] – will

provide the foundations for designing new classes of precision GPCR drugs.

Residues biased toward Nb80 signaling are enriched in ECL2, which forms part

of the exosite of salmeterol, a β2AR agonist strongly biased toward G-protein

signaling, suggesting mutant behavior is related to drug activity binding at the

same sites. Likewise, known allosteric modulators bind in pockets with high

impact mutations that match their activities. AS408, a NAM, binds in a pocket

with loss- and change- of function residues, while Cmpd-6FA, a PAM, binds in a

pocket with gain- and change- of function residues. In addition to these known

allosteric sites, GPCR-MAPS prioritizes an additional – and, to our knowledge,

novel – allosteric pocket to target in β2AR and deprioritizes one where

mutations only have limited effects.

The GPCR-MAPS approach is modular and scalable, and its application

across the GPCR superfamily will enable a better understanding of the largest

family of human receptors. To date GPCR drug discovery has proceeded from

an era of blind high-throughput screening to an era of structure-based drug

discovery. We believe we are now entering a third era of GPCR drug discovery,

where high-resolution functional maps empower the rational design of highly

specific biased drugs and allosteric modulators.

Materials and Methods

Plasmids

Plasmids encoding SPARK2 components were gifts from Christina Kim and Alice Ting and were obtained from Addgene (Addgene plasmid # 125228, 104839, 104840, 104841). From these plasmids we generated new plasmids using Gibson cloning. attB-HA-b2AR-eLOV-TEVcs-Gal4-UAS-Citrine encodes the receptor fusion followed by the bGH poly(A) signal, then the UAS sequence followed by Citrine fluorescent protein. A 26 nucleotide barcode was inserted in the 3' UTR of the Citrine sequence. To make cloning more efficient, we transferred the fusion coding sequence from the NanoLuc-15aa linker-βarrestin2-HA-GS linker-TEVp plasmid from the pAAV vector (which contains GC-rich inverted terminal repeats) to the pUC19 vector. We then used site directed mutagenesis to introduce the S153N variant to TEV, generating uTEV1Δ). We then made versions of this plasmid with Nb80 and miniG$_s$ sequence instead of β-Arrestin. We also modified the miniG$_s$ vector so that uTEV1Δ was fused N-terminally (NanoLuc-15aa linker-GS linker-TEVp-miniG$_s$). pCAG-NLS-HA-Bxb1 was a gift from Kenny Matreyek and Doug Fowler.

Saturation mutagenesis library creation

We designed six pools of mutation-bearing oligonucleotides, each pool representing mutated coding sequence for 66-70 amino acids of the full-length (413 amino acids) b2AR with flanking wild-type sequence for amplification from the pool (using primers dialout_tile[1-6]_[F/R], Supplementary Table 5). For each pool we designed primers to amplify the whole plasmid except for the mutated segment (using primers b2AR_satmut_tile[1-6]_[F/R]), then used Gibson assembly to introduce the mutagenic oligos. Oligos were designed to

encode all possible single amino acid changes (using the human optimized

codon for each alternative amino acid), all premature stop codons, and all

synonymous codons when possible (using the most optimal codon, excluding

the wild-type codon).

Barcode integration and variant-barcode association

10 µg of the attB-HA-b2AR-eLOV-TEVcs-Gal4-UAS-Citrine saturation

mutagenesis plasmid was digested with 40 units PpuMI (New England Biolabs)

overnight at 37°, then in the morning 10 units more PpuMI were added and 30

minutes more incubation at 37°. Then the cut band was gel purified. The

barcode construct (oTLM_257) was made double stranded by PCR amplifying

with oTLM_258 and oTLM_259 (Supplementary Table 5). Double stranded

barcode was incorporated into the cut plasmid backbone by Gibson assembly

and transformed into electrocompetent bacteria. Following recovery in SOC

media, we plated 0.1% of the transformation to estimate the number of clones,

and inoculated the rest into LB media with ampicillin for overnight growth.

Estimated number of clones was 300,000. In the morning the bacteria was

pelleted and plasmid was midi-prepped (Qiagen Plasmid Plus Midi Kit).

We used PacBio long read sequencing to associate each variant with the

barcode. We digested the plasmid pool with BsiWI (New England Biolabs) to

release the coding sequence and barcode and SMRT bell adapters were ligated

according to manufacturer's instructions. The library was sequenced on a

PacBio Sequel IIe. We used alignparse[71] to align reads, quality filter, and call

sequences for each barcode (Supplementary Figure 1).

Landing pad integration

HEK 293T LLP-iCasp9-Blast[31] cells were a gift from Kenny Matreyek.

Cells were grown in DMEM with GlutaMAX (Gibco), with Tetracyline-free fetal

bovine serum. 14 million cells were plated onto a T225 flask. The next day,

transfection was done with Lipofectamine 3000 (Invitrogen) with 17.5 μg of

pCAG-NLS-HA-Bxb1 and 17.5 μg of the mutated and barcoded attB-HA-b2AR-

eLOV-TEVcs-Gal4-UAS-Citrine. The cells were allowed to incubate for two

days. Then, the media was removed and replaced with media containing 2

μg/mL doxycycline and incubated for one day. Then, the media was changed

for media containing 2 μg/mL doxycycline and 10 nM rimiducid

(MedChemExpress) and incubated for one day. Then media was changed to

include just 2 μg/mL doxycycline and outgrowth of cells was allowed for ~one

week, passaging cells when they reached 90% confluency.

GPCR-MAPS signaling experiment

10 million cells were plated onto each T175 flask and allowed to incubate

overnight. The next day, the cells were transfected with 36 μg of transducer

plasmids (pUC19-NanoLuc-β-arrestin-uTEVp1 or pUC19-NanoLuc-Nb80-

uTEVp1), using Lipofectamine 3000 and allowed to incubate overnight. The

next morning, cells were treated with media containing furimazine and either

drug or DMSO: [13.05 mL DMEM, 126.56 μL furimazine, 270 μL 1M HEPES,

and 13.5 μL isoproterenol or DMSO]. Furimazine was obtained from the Nano-

Glo Live Cell Assay System (Promega). Isoprenaline was obtained from Sigma

(catalog number 15627). Cells were incubated in the dark (covered with

aluminum foil) at 37° for 24 hours. Then, cells were lifted with Trypsin-EDTA,

pelleted and stored in -80 until RNA extraction. For the Gal4 maximum

stimulation condition, after 10 million cells were plated and grown overnight, 36

µg of pUC19-CMV-Gal4 was transfected with Lipofectamine 3000. The cells

were allowed to incubate for 30 hours, then cells were pelleted and stored at -

80°.

RNA isolation and sequencing library preparation

For the first set of experiments (Nb80 and BArr signaling with 10 µM

isoprenaline), cell pellets were thawed and resuspended in buffer RLT with

added beta-mercaptoethanol. Then, they were homogenized by centrifuging

through QIAshredder columns. RNA was then purified from the homogenate

with RNeasy Midi-prep kit (Qiagen) and eluted with 400 µL of water. Between

120 and 160 µg of total RNA was used as input for reverse transcription (RT)

reactions (6-8 µg per reaction, 20 reactions per sample) using SuperScript III

reverse transcriptase (Invitrogen). For the dose response experiment,

Dynabeads mRNA direct purification kit (ThermoFisher Scientific) was used

according to manufacturers instructions and mRNA was eluted in 80 µL of Tris-

HCl. Total mRNA yields were between 11.5 µg and 16 µg. Between 5.75 and 8

µg mRNA was used as input for RT (1.15-1.6 µg per reaction, 10 reactions per

sample). For both sets of reactions, RT was carried out with 2 pmol of primer

(oTLM_266, Supplementary Table 5) and incubated at 55° for one hour then

75° for 15 minutes. Following this, RNase A was added to a final concentration

of 100 µg/mL and reactions incubated at 37° for 30 minutes. Then, reactions

were cleaned up with Nucleospin PCR cleanup columns (Macherey Nagel) and

eluted in 50 µL of elution buffer. 0.5 µL of the cDNA eluate was used in a qPCR

reaction (Q5 High-Fidelity polymerase, New England Biolabs, and EvaGreen

fluorescent dye, Biotium) to determine the number of cycles required to reach

near-plateau phase. Then, 40 µL of cDNA was amplified across 4x 50 µL PCR

reactions with the program 98° for 30s, followed by 13-16 cycles of [98° for 15s,

64° for 30s, 72° for 30s].

Reactions were column purified and 4% of this was used as input for a

second round PCR, which added the rest of the Illumina adapter as well as

unique index sequences. The second round PCR was 98° for 30s, followed by 5

cycles of [98° for 15s, 64° for 30s, 72° for 30s]. Frameshifting primers were

used to introduce variability in the first few cycles of the sequencing reaction

(oTLM_277-282, Supplementary Table 5). Reactions were then gel purified and

analyzed on TapeStation (Agilent Technologies). Sequencing was done with

Illumina NovaSeq with 2x50 paired end reads.

An additional step was taken for the dose response experiment.

Following RT, we performed triplicate qPCR reactions with each cDNA sample,

along with a set of serial dilutions of the original library plasmid, in order to

estimate the relative quantity of Citrine RNA produced from each experiment.

We used primers oTLM_021 and oTLM_024, which amplify the 5' UTR of

Citrine. We chose to amplify this region instead of the barcode (as was done for

the library prep) to avoid idiosyncratic effects of amplification efficiency caused

by diverse and potentially different barcode identities across different samples.

Following qPCR we fit a standard curve (Supplementary Fig. 5b) then used this

to estimate RNA abundance for the dose response samples (Figure 5b).

Surface expression experiment cell culture and FACS

Cells were cultured in 245x245mm cell culture flasks until ~80%

confluent. Then, cells were lifted with Trypsin-EDTA, pelleted, supernatant

removed, and resuspended in blocking buffer (1% bovine serum albumin in

phosphate buffered saline). Cells were counted and 50 M cells were transferred

to a 5 mL conical tube with 3.5 mL of blocking buffer, and incubated on ice,

covered from light, with rocking for 30 minutes. Then, anti-HA antibody (Cell

Signaling HA-Tag 6E2 Alexa Fluor 647 Conjugate) was added at dilution of

1:100 and incubated on ice, covered from light, with rocking for 60 more

minutes. Then, cells were pelleted, supernatant removed, and resuspended in

blocking buffer. Sorting was done on the BD FACSAria instrument. Bins were

drawn so that each represented ~25% of cells. In total, ~20M cells were sorted

for replicate 1, and ~30M cells were sorted for replicates 2 and 3. 30-45M 2x50

basepair reads were collected on an Illumina NovaSeq instrument for each

replicate.

Surface expression sequencing library preparation

Sorted cells were pelleted and DNA extracted using Qiagen DNeasy

Blood & Tissue kit, using 2 columns for each sample. 200 µL elution buffer was

used to elute from each column, then eluates for each sample were combined.

Then, 320 µL of each sample was was used as input for 16x 50 µL PCRs with

primers adjacent to the barcode that include partial Illumina adapters and also

varying lengths of degenerate bases to increase nucleotide complexity in the

first few cycles of the sequencing run (oTLM277-oTLM282, Supplementary

Table 5). Cycling was 98° for 30s, followed by 25 cycles of [98° for 15s, 64° for

30s, 72° for 30s]. Reactions were column purified and 4% of this was used as

input for a second round PCR, which added the rest of the Illumina adapter as

well as unique index sequences. The second round PCR was 98° for 30s,

followed by 5 cycles of [98° for 15s, 64° for 30s, 72° for 30s]. The products of

the second PCR were gel purified and analyzed on TapeStation, then

sequenced on Illumina NovaSeq with 2x50 paired end reads.

Sequencing data processing

807   Paired-end sequencing reads were first merged with vsearch[72] and then

808 adapters were trimmed with cutadapt[73]. This leaves each read representing the

809 barcode sequence, which was then matched with barcodes identified in the

810 variant-barcode association with a custom script.

811 **Calculation of surface expression scores**

812   Surface expression scores were calculated as previously described[36].

Briefly, the frequency of each variant in each bin is enumerated by combining

counts of all barcodes associated with that variant. The frequency, along with

the total number of cells sorted into each bin, is used to estimate the number of

cells in each bin. Then, the log10 transformed geometric mean fluorescence

value of all cells sorted into each bin is combined with the estimated cell counts

to arrive at the raw surface expression estimate. These estimates were then

normalized between 0 (represented by the median of presumed loss-of-function

variants: e.g. premature stop codons before the 300^{th} amino acid) and 1, the

WT score. Surface expression scores were considered high confidence if the

estimated number of cells sorted was >50.

Calculation of signaling scores

824   First, barcode read counts for all barcodes representing the same variant

were combined and then the frequency of that variant was calculated by

dividing that number by the total number of barcode counts in the sample.

Then, to arrive at a raw score, we divided the variant frequency in the

experimental sample by the variant frequency in the Gal4 sample, and take the

log_2 of this. The Gal4 sample represents a maximum stimulation condition in

which the barcode frequency in the library, as well as any idiosyncratic effects

of the barcode sequence on RNA stability should be captured. Raw scores are

then scaled between 0 and 1, with the median of nonsense mutations set as 0

and the wild-type score set as 1. Signaling scores were considered high

confidence if the frequency of that variant in the Gal4 condition was >$2x10^{-5}$.

LOWESS modeling and identifying outlier positions

The python package statsmodels.nonparametric.smoothers_lowess[74]

was used with the following parameter values: frac=0.5, it=3, delta=0.0.

Following curve fitting, residuals were calculated as the distance in the y-axis

between the data point and the line, and z-scores were calculated as the

residual / standard error of the mean. In the case of Nb80 versus β-arrestin, the

average standard error of the mean between the two measurements was used.

Structural analysis

Residue contacts were defined by GetContacts

(https://getcontacts.github.io/index.html). The binding interface of β2AR with

Nb80 was defined by GetContacts analysis of structure. The predicted inactive

and active structures in Figure 6k were taken from GPCRdb and were produced

with AlphaFold-Multistate[37]. These structures were used to enable visualization

of the entire ICL3, which is not resolved in experimentally solved structures.

Dose response curve fitting

In the package neutcurve[75], we used the neutcurve.hillcurve.HillCurve

function to fit curves with the following options:

infectivity_or_neutralized='neutralized', fix_slope_first=True, fs_stderr=None,

fixbottom=False, fixtop=False, fixslope=False, fitlogc=False,

use_stderr_for_fit=False, no_curve_fit_first=False. Following the fitting

procedure, we then filtered curves based on heuristics: R^2 > 0.9, EC50 within

the measured range (i.e. between 10pM and 10uM), and $E_{max} < 1.5x$ the

maximum measured signaling value. See Supplementary Figure 5e for the

number of variants filtered.

Fpocket pocket prediction

We used PDB ID 2RH1 as the structural template for pocket prediction

and used the Fpocket web server with default settings to predict pockets. We

chose the two pockets to illustrate simply based on having many or few

annotated residues.

Acknowledgments

We thank Christina Kim for advice regarding the SPARK2 system. We

thank the CRG/UPF Flow Cytometry Unit for technical assistance. We thank

Jana Selent and lab members for helpful discussion. We thank all members of

the Lehner lab for helpful discussion.

Data availability

Files needed to reproduce analyses can be found at zenodo

(https://zenodo.org/records/15363336). Raw sequencing reads can be found at

Sequence Read Archive (accession number PRJNA1260170).

Code availability

Custom code to reproduce analyses can be found at github

(https://github.com/lehner-lab/GPCR-MAPS).

Competing interests

CRG and the Wellcome Sanger Institute have filed a patent application

for the use of the methods described in this study for mapping allostery in

GPCRs; T.M. and B.L. are listed as inventors. B.L. is a founder and shareholder

of ALLOX.

Figure 1. GPCR-MAPs

a Schematic of the experimental framework. A mutagenized GPCR (in this case β2AR) is encoded in a plasmid that also includes the attB recombination site, an N-terminal HA epitope tag, C-terminal fusions of a light, oxygen, and voltage sensitive (LOV) domain, the cleavage site of TEV protease (TEVcs), and Gal4 transcription factor. Downstream, on the same plasmid, is the fluorescent protein Citrine, driven by Gal4-responsive UAS promoter, with a degenerate DNA barcode included in the 3' untranslated region. This construct is recombined into HEK 293T LLP-iCasp9-Blast landing pad cells. Then, two assays are performed to measure the surface expression (FACS-based) and signaling activity (transcriptional reporter) of variants *en masse***.** The outputs of these experiments (DNA for FACS and RNA for transcriptional reporter) are sequenced with short read technology. Estimated surface expression and signaling scores are then modeled to understand the contribution of each of these to the variant effects. **b** Detailed schematic of the signaling assay. The receptor construct is expressed from landing pad cells while the transducer construct is transiently transfected. Addition of agonist (isoprenaline in this case) and furimazine results in recruitment of transducer to the receptor fusion and subsequent cleavage of the TEVcs, liberating Gal4 which moves to the nucleus and drives transcription of the UAS-Citrine-barcode construct. **c** Replicate Pearson correlations for the surface expression, Nb80 signaling, and βArr signaling. $p < 10^{-100}$ in each case. **d** Heatmap representation of the surface expression data. β2AR positions on the x-axis, mutation on the y-axis. Gray cells indicate expression similar to wild-type while blue and red cells indicate reduced and increased expression, respectively. **e** Histogram of surface

expression variant scores; nonsense variants in red, synonymous wild-type in

green, and missense in grey. **f** Heatmap representation of the Nb80 signaling

data. Axes and color scale as in (d). **g** Histogram of Nb80 signaling scores;

colors as in (e). **h** Heatmap representation of β-Arrestin signaling scores. Axes

and color scale as in (d). **i** Histogram of β-Arrestin signaling scores; colors as in

(e). **j** Surface expression scores for residues that directly contact isoprenaline in

the solved structure (PDB ID: 7DHR). **k** Nb80 signaling scores for residues that

directly contact isoprenaline. **l** β-Arrestin signaling scores for residues that

directly contact isoprenaline.

Figure 2. Deconvoluting expression and signaling effects

a Surface expression versus Nb80 signaling scores for missense variants.

$p<10^{-100}$. **b** Surface expression versus β-Arrestin signaling scores for missense

variants. $p<10^{-100}$. **c** Schematic of the method for identifying residues

contributing to signaling beyond what can be explained by surface expression.

First, the signaling data is related to surface expression with a LOWESS

regression. Then, for each variant, residuals to this fit are calculated. Then,

residues with significantly biased residuals are identified as loss-of-function or

gain-of-function via Mann Whitney U test. Finally, positions where variants have

a wide range of effects, quantified using the interquartile range of variant z-

scores, are identified as change-of-function. **d** LOWESS fit of surface

expression versus Nb80 signaling, with variant points colored by residual to the

fit. **d** Same as (e) but for β-Arrestin signaling. **f** Volcano plot showing the

position-average z-score and the minus log10 transformed p-value of all

residues for Nb80 signaling. Residues with significantly biased residuals are

colored according to the average z-score. **g** Same as in (f) but for β-Arrestin

signaling. **h** Venn diagrams showing the number of residues called loss-, gain-,

and chage-of-function for Nb80 and β-Arrestin signaling. **i** Surface expression

and signaling effects of variants at D113^{3x32}, plotted over the population of

variant effects, for Nb80 and β-Arrestin versus surface expression. Error bars

represent standard error of the mean. **j** Same as (i) but with variants at position

E268^{6x30}. **k** Same as (i) and (j) but with variants at position S120^{3x39}. **l** Loss-of-

function residues highlighted in purple on the isoprenaline-bound β2AR

structure (PDB ID: 7DHR, isoprenaline in yellow). **m** Gain-of-function residues

highlighted in green on β2AR. **n** Change-of-function residues highlighted in red

on β2AR

Figure 3. A core activation network links ligands to transducers

a Inactive and active state structures of β2AR with LoF residues colored purple.

Those LoF residues that make a direct contact with other LoF residues are

visualized as pseudobonds connecting them, while disconnected LoF residues

are shown with spheres. **b** Enlarged view of the core activation network.

Figure 4. Genetic basis of biased signaling

a Schematic showing β2AR signaling through β-Arrestin and G-protein (Nb80)

mediated pathways in response to isoprenaline. **b** Nb80 versus β-Arrestin

signaling for missense variants. $p<10^{-100}$, n=6,791. **c** Nb80 versus β-Arrestin

signaling with a LOWESS fit and variant scores colored by residual to the fit

line. Individual variant scores, with error bars representing standard error of the

mean, shown for three variants previously reported to cause biased signaling. **d**

Volcano plot showing the position-average z-score and the minus log10

transformed p-value of all residues for Nb80 versus β-Arrestin signaling.

Residues with significantly biased residuals are colored according to the

average z-score. **e** Nb80 and β-Arrestin signaling effects of variants at W99^{23x50},

plotted over the population of variant effects, error bars represent standard error

of the mean. **f** Same as in (e) but with variants at P138^{34x50}. **g** Same as in (e)

but with variants at F332^{8x50}. **h** β-Arrestin biased, Nb80 biased, or bias switch

residues highlighted on the solved structure, 7DHR. Isoprenaline show in

yellow. **i** F193^{45x52} and Y318^{7x34} form a "lid" over isoprenaline in the solved

structure (7DHR). **j** Hydrophobicity (Kyte-Doolittle scale) of the mutant amino

acid compared with the Nb80 versus β-Arrestin signaling bias. **k** Enlarged view

of the extracellular vestibule. Colored residues are significantly Nb80-biased

(orange), β-Arrestin-biased (blue), or bias switches (magenta). **l** Enlarged view

of NPxxY motif and TM7, colored residues same as in (k).

Figure 5. Quantifying >7,500 dose-response relationships

a Experimental design: serial dilutions of isoprenaline were used to stimulate

landing pad cells expressing the β2AR library. Then, the GPCR-MAPS signaling

assay was performed and RNA was collected from the stimulated cells. **b** The

RNA abundance from each sample was first quantified with qPCR, then the

variant frequencies were estimated with short-read sequencing. **c** Per-variant

normalized scores were then calculated by normalizing variant frequencies in

each condition by the RNA abundance of each condition. **d** Heatmap

representations of the 8 highest concentration drug treatment conditions. β2AR

positions on the x-axis, mutation identities on the y-axis. One y-axis is blown up;

the rest of the heatmaps follow this same scheme. Grey cells indicate wild-type-

like signaling, blue and red indicate reduced and increased signaling,

respectively. One colorbar is blown up; the rest of the heatmaps follow this

same scheme. **e** Clustered heatmap of signaling intensities. Colorbar on top.

Three main clusters correspond to wild-type-like, moderately damaging, and

complete loss-of-function curves. **f** The curve fit to the wild-type data, with E_{max},

E_{min}, EC_{50}, and slope indicated. **g** Fit curves from the wild-type-like cluster in

black, wild-type curve in red. **h** Fit curves from the moderately damaging cluster

in black, wild-type curve in red. **i** Fit curves from the loss-of-function cluster in

black, wild-type curve in red.

Figure 6. Genetic determinants of potency

a Schematic showing example curves for increased and decreased potency. **b**

Comparison of variant EC_{50} estimates from this study with previously published

estimates[18]. Red dashed line is x=y line. **c** Spearman correlations of curve

fitting parameter estimates and surface expression scores. **d** Surface

expression scores compared with log-transformed EC_{50} estimates. $p < 10^{-100}$. **e**

Volcano plot showing the position-average z-score and the minus log10

transformed p-value of all residues for log EC_{50} versus surface expression. **f**

Log-transformed EC_{50} versus surface expression for variants at $R328^{7x55}$,

plotted over the population of variant effects, error bars represent standard error

of the mean. **g** Same as in (f) but with variants at $E268^{6x30}$. **h** Same as in (f) but

with variants at $F282^{6x44}$. **i** EC_{50} loss-of-function, gain-of-function, or change-of-

function residues highlighted on the solved structure, 7DHR. Isoprenaline show

in yellow. **j** Violin plots showing minimum distance between LoF, GoF, or CoF

residues from the core activation network in AlphaFold-Multistate active state

model. Distances capped at 20Å. **k** Portions of TMs 3, 5, and 6, as predicted by

AlphaFold-Multistate and downloaded from GPCRdb[37]. LoF residues colored

aquamarine, GoF residues colored brown, and CoF residues colored salmon.

1015

Figure 7. Rationalization and prioritization of allosteric pockets

a The binding pocket of known negative allosteric modulator, AS408, shown in

yellow. PDB ID: 6OBA. Residues making direct contact with the NAM are

colored beige. **b** Expression normalized maximum signaling through Nb80,

signaling bias, and expression normalized potency of the mutations at sites

making up the binding pocket of AS408. Shown below the heatmaps are the

categorical annotations of significant residues: purple, green, and red indicate

LoF, GoF, and CoF residues for maximum signaling. Orange, blue, and

magenta indicate Nb80 biased, β-Arrestin, biased, or bias switches.

Aquamarine, brown, and salmon indicate LoF, GoF, and CoF for potency. **c** The

binding pocket of AS408 with annotated residues colored. For residues

annotated in both maximum signaling and potency analyses, the color from

maximum signaling was used. **d, e, f** Same as in (a, b, c) but for the positive

allosteric modulator Cmpd-6FA, with structure 6N48. **g, h, i** Same as in (a, b, c)

but for a predicted active pocket in structure 2RH1. **j, k, l** Same as in (a, b, c)

but for a predicted inactive pocket in structure 2RH1.

1032

Figure 8. Summary of findings.

a Selected findings from the maximum signaling analysis. Core activation

network shown as purple lines, residues or topological domains colored

according to annotation. **b** Selected findings from biased signaling analysis.

NPxxY motif collectively illustrated as "bias switch" as there are multiple bias

functional residues in and near this motif. **c** Selected findings from the potency

analysis. Ligand binding and second shell residues illustrated as triangle.

References

References

1. Bockaert, J. & Philippe Pin, J. Molecular tinkering of G protein-coupled receptors: an

evolutionary success. *EMBO J.* **18**, 1723–1729 (1999).

2. Farrens, D. L., Altenbach, C., Yang, K., Hubbell, W. L. & Khorana, H. G. Requirement

of Rigid-Body Motion of Transmembrane Helices for Light Activation of Rhodopsin.

Science **274**, 768–770 (1996).

3. Rosenbaum, D. M., Rasmussen, S. G. F. & Kobilka, B. K. The structure and function of

G-protein-coupled receptors. *Nature* **459**, 356–363 (2009).

4. Jean-Charles, P.-Y., Kaur, S. & Shenoy, S. K. GPCR signaling via β-arrestin-dependent

mechanisms. *J. Cardiovasc. Pharmacol.* **70**, 142–158 (2017).

5. Overington, J. P., Al-Lazikani, B. & Hopkins, A. L. How many drug targets are there?

Nat. Rev. Drug Discov. **5**, 993–996 (2006).

6. Sriram, K. & Insel, P. A. G Protein-Coupled Receptors as Targets for Approved Drugs:

How Many Targets and How Many Drugs? *Mol. Pharmacol.* **93**, 251–258 (2018).

7. Wold, E. A., Chen, J., Cunningham, K. A. & Zhou, J. Allosteric Modulation of Class A

GPCRs: Targets, Agents, and Emerging Concepts. *J. Med. Chem.* **62**, 88–127 (2019).

8. Wootten, D., Christopoulos, A., Marti-Solano, M., Babu, M. M. & Sexton, P. M.

Mechanisms of signalling and biased agonism in G protein-coupled receptors. *Nat.*

Rev. Mol. Cell Biol. **19**, 638–653 (2018).

9. May, L. T., Leach, K., Sexton, P. M. & Christopoulos, A. Allosteric modulation of G

protein-coupled receptors. *Annu. Rev. Pharmacol. Toxicol.* **47**, 1–51 (2007).

10.   Kenakin, T. Biased Receptor Signaling in Drug Discovery. *Pharmacol. Rev.* **71**,

267–315 (2019).

11.   Wootten, D., Christopoulos, A. & Sexton, P. M. Emerging paradigms in GPCR

allostery: implications for drug discovery. *Nat. Rev. Drug Discov.* **12**, 630–644

(2013).

12.   Lorente, J. S. *et al.* GPCR drug discovery: new agents, targets and indications.

Nat. Rev. Drug Discov. 1–22 (2025) doi:10.1038/s41573-025-01139-y.

13.   Bermudez, M., Nguyen, T. N., Omieczynski, C. & Wolber, G. Strategies for the

discovery of biased GPCR ligands. *Drug Discov. Today* **24**, 1031–1037 (2019).

14.   Allosteric Modulator Discovery: From Serendipity to Structure-Based Design. *J.*

Med. Chem. **62**, 6405–6421 (2019).

15.   Venkatakrishnan, A. J. *et al.* Diverse activation pathways in class A GPCRs

converge near the G-protein-coupling region. *Nature* **536**, 484–487 (2016).

16.   Zhou, Q. *et al.* Common activation mechanism of class A GPCRs. *eLife* **8**, e50279

(2019).

17.   Ward, W. H. J., Timms, D. & Fersht, A. R. Protein engineering and the study of

structure—function relationships in receptors. *Trends Pharmacol. Sci.* **11**, 280–284

(1990).

18.   Heydenreich, F. M. *et al.* Molecular determinants of ligand efficacy and potency

in GPCR signaling. *Science* **382**, eadh1859 (2023).

19.   Morales-Pastor, A. *et al.* Multiple triggers converge to preferential effector

coupling in the CB2R through a complex allosteric communication network.

2024.02.27.582279 Preprint at https://doi.org/10.1101/2024.02.27.582279 (2024).

20.   Cao, Y. *et al.* Unraveling allostery within the angiotensin II type 1 receptor for

Gαq and β-arrestin coupling. *Sci. Signal.* **16**, eadf2173 (2023).

21.   Penn, W. D. *et al.* Probing biophysical sequence constraints within the

transmembrane domains of rhodopsin by deep mutational scanning. *Sci. Adv.* **6**,

eaay7505 (2020).

22.   McKee, A. G. *et al.* Systematic profiling of temperature- and retinal-sensitive

rhodopsin variants by deep mutational scanning. *J. Biol. Chem.* **297**, 101359 (2021).

23.   Heredia, J. D. *et al.* Mapping Interaction Sites on Human Chemokine Receptors

by Deep Mutational Scanning. *J. Immunol.* **200**, 3825–3839 (2018).

24.   Park, J. *et al.* Structural architecture of a dimeric class C GPCR based on co-

trafficking of sweet taste receptor subunits. *J. Biol. Chem.* **294**, 4759–4774 (2019).

25.   Yoo, J. I., Daugherty, P. S. & O'Malley, M. A. Bridging non-overlapping reads

illuminates high-order epistasis between distal protein sites in a GPCR. *Nat.*

Commun. **11**, 690 (2020).

26.   Jones, E. M. *et al.* Structural and functional characterization of G protein–

coupled receptors with deep mutational scanning. *eLife* **9**, e54895 (2020).

27.   Howard, M. K. *et al.* Molecular basis of proton sensing by G protein-coupled

receptors. *Cell* **188**, 671-687.e20 (2025).

28.   Howard, C. J. *et al.* High resolution deep mutational scanning of the

melanocortin-4 receptor enables target characterization for drug discovery. *eLife*

13, (2024).

29.   Ippolito, M. & Benovic, J. L. Biased agonism at β-adrenergic receptors. *Cell.*

Signal. **80**, 109905 (2021).

30.   Kim, C. K., Cho, K. F., Kim, M. W. & Ting, A. Y. Luciferase-LOV BRET enables

versatile and specific transcriptional readout of cellular protein-protein interactions.

eLife **8**, e43826 (2019).

31.   Matreyek, K. A., Stephany, J. J., Chiasson, M. A., Hasle, N. & Fowler, D. M. An

improved platform for functional assessment of large protein libraries in

mammalian cells. *Nucleic Acids Res.* **48**, e1 (2020).

32.   Rasmussen, S. G. F. *et al.* Structure of a nanobody-stabilized active state of the

β2 adrenoceptor. *Nature* **469**, 175–180 (2011).

33.   Sanchez, M. I. & Ting, A. Y. Directed evolution improves the catalytic efficiency

of TEV protease. *Nat. Methods* **17**, 167–174 (2020).

34.   Kim, M. W. *et al.* Time-gated detection of protein-protein interactions with

transcriptional readout. *eLife* **6**, e30233 (2017).

35.   Carpenter, B. & Tate, C. G. Engineering a minimal G protein to facilitate

crystallisation of G protein-coupled receptors in their active conformation. *Protein*

Eng. Des. Sel. **29**, 583–594 (2016).

36.   Mighell, T. L. & Lehner, B. A pharmacological chaperone stabilizer rescues the

expression of the vast majority of pathogenic variants in a G protein-coupled

receptor. 2024.11.28.625821 Preprint at

https://doi.org/10.1101/2024.11.28.625821 (2024).

37.   Pándy-Szekeres, G. *et al.* GPCRdb in 2023: state-specific structure models using

AlphaFold2 and new ligand resources. *Nucleic Acids Res.* **51**, D395–D402 (2023).

38.   Valentin-Hansen, L. *et al.* The Arginine of the DRY Motif in Transmembrane

Segment III Functions as a Balancing Micro-switch in the Activation of the β2-

Adrenergic Receptor. *J. Biol. Chem.* **287**, 31973–31982 (2012).

39.   Sanchez-Reyes, O. B. *et al.* G Protein-Coupled Receptors Contain Two

Conserved Packing Clusters. *Biophys. J.* **112**, 2315–2326 (2017).

40.   Nygaard, R., Frimurer, T. M., Holst, B., Rosenkilde, M. M. & Schwartz, T. W.

Ligand binding and micro-switches in 7TM receptor structures. *Trends Pharmacol.*

Sci. **30**, 249–259 (2009).

41.   Yang, D. *et al.* G protein-coupled receptors: structure- and function-based drug

discovery. *Signal Transduct. Target. Ther.* **6**, 1–27 (2021).

42.   Ragnarsson, L., Andersson, Å., Thomas, W. G. & Lewis, R. J. Mutations in the

NPxxY motif stabilize pharmacologically distinct conformational states of the α1B-

and β2-adrenoceptors. *Sci. Signal.* **12**, eaas9485 (2019).

43.   Zhao, L., He, X., Jiang, H. & Cheng, X. Computational characterization of

transducer recognition of β2 adrenergic receptor. *Biochem. Biophys. Res. Commun.*

592, 67–73 (2022).

44.   Rovati, G. E., Capra, V. & Neubig, R. R. The highly conserved DRY motif of class

A G protein-coupled receptors: beyond the ground state. *Mol. Pharmacol.* **71**, 959–

964 (2007).

45.   Rasmussen, S. G. F. *et al.* Mutation of a Highly Conserved Aspartic Acid in the

*β*2 Adrenergic Receptor: Constitutive Activation, Structural Instability, and

Conformational Rearrangement of Transmembrane Segment 6. *Mol. Pharmacol.* **56**,

175–184 (1999).

46.   Tao, Y. X., Abell, A. N., Liu, X., Nakamura, K. & Segaloff, D. L. Constitutive

activation of G protein-coupled receptors as a result of selective substitution of a

conserved leucine residue in transmembrane helix III. *Mol. Endocrinol. Baltim. Md*

14, 1272–1282 (2000).

47.   Wheatley, M. *et al.* Lifting the lid on GPCRs: the role of extracellular loops. *Br. J.*

Pharmacol. **165**, 1688–1703 (2012).

48.   Ippolito, M., De Pascali, F., Inoue, A. & Benovic, J. L. Phenylalanine 193 in

Extracellular Loop 2 of the β2-Adrenergic Receptor Coordinates β-Arrestin

Interaction. *Mol. Pharmacol.* **101**, 87–94 (2022).

49.   Choi, M. *et al.* G protein–coupled receptor kinases (GRKs) orchestrate biased

agonism at the β2-adrenergic receptor. *Sci. Signal.* **11**, eaar7084 (2018).

50.   Sanchez-Soto, M. *et al.* A structural basis for how ligand binding site changes

can allosterically regulate GPCR signaling and engender functional selectivity. *Sci.*

Signal. **13**, eaaw5885 (2020).

51.   Masureel, M. *et al.* Structural insights into binding specificity, efficacy and bias

of a β2AR partial agonist. *Nat. Chem. Biol.* **14**, 1059–1066 (2018).

52.   Bock, A. & Bermudez, M. Allosteric coupling and biased agonism in G protein-

coupled receptors. *FEBS J.* **288**, 2513–2528 (2021).

53.   Woo, A. Y.-H. *et al.* Tyrosine 308 is necessary for ligand-directed Gs protein-

biased signaling of β2-adrenoceptor. *J. Biol. Chem.* **289**, 19351–19363 (2014).

54.   Dror, R. O. *et al.* Pathway and mechanism of drug binding to G-protein-coupled

receptors. *Proc. Natl. Acad. Sci.* **108**, 13118–13123 (2011).

55.   Liu, J. J., Horst, R., Katritch, V., Stevens, R. C. & Wüthrich, K. Biased Signaling

Pathways in β2-Adrenergic Receptor Characterized by 19F-NMR. *Science* **335**, 1106–

1110 (2012).

56.   Chure, G. *et al.* Predictive shifts in free energy couple mutations to their

phenotypic consequences. *Proc. Natl. Acad. Sci.* **116**, 18275–18284 (2019).

57.   Razo-Mejia, M. *et al.* Tuning Transcriptional Regulation through Signaling: A

Predictive Theory of Allosteric Induction. *Cell Syst.* **6**, 456-469.e10 (2018).

58.   Steiner, P. J. *et al.* A Closed Form Model for Molecular Ratchet-Type Chemically

Induced Dimerization Modules. *Biochemistry* **62**, 281–291 (2023).

59.   Maximilian R Stammnitz & Ben Lehner. The genetic architecture of an allosteric

hormone receptor. *bioRxiv* (2025).

60.   Weis, W. I. & Kobilka, B. K. The Molecular Basis of G Protein–Coupled Receptor

Activation. *Annu. Rev. Biochem.* **87**, 897–919 (2018).

61.   Sadler, F. *et al.* Autoregulation of GPCR signalling through the third intracellular

loop. *Nature* **615**, 734–741 (2023).

62.   Li, X., Zhou, M., Huang, W. & Yang, H. N-glycosylation of the β2 adrenergic

receptor regulates receptor function by modulating dimerization. *FEBS J.* **284**, 2004–

2018 (2017).

63.   Chen, S. *et al.* Mutation of a Single TMVI Residue, Phe282, in the β2-Adrenergic

Receptor Results in Structurally Distinct Activated Receptor Conformations.

Biochemistry **41**, 6045–6053 (2002).

64.   Liu, X. *et al.* Mechanism of intracellular allosteric β2AR antagonist revealed by

X-ray crystal structure. *Nature* **548**, 480–484 (2017).

65.   Liu, X. *et al.* An allosteric modulator binds to a conformational hub in the β2

adrenergic receptor. *Nat. Chem. Biol.* **16**, 749–755 (2020).

66.   Liu, X. *et al.* Mechanism of β2AR regulation by an intracellular positive allosteric

modulator. *Science* **364**, 1283–1287 (2019).

67.   Le Guilloux, V., Schmidtke, P. & Tuffery, P. Fpocket: An open source platform

for ligand pocket detection. *BMC Bioinformatics* **10**, 168 (2009).

68.   Chen, G. *et al.* Activation and allosteric regulation of the orphan GPR88-Gi1

signaling complex. *Nat. Commun.* **13**, 2375 (2022).

69.   Onaran, H. O. *et al.* Systematic errors in detecting biased agonism: Analysis of

1205   current methods and development of a new model-free approach. *Sci. Rep.* **7**,

1206   44247 (2017).

1207   70.   Beltran, A., Faure, A. J. & Lehner, B. The allosteric landscape of the Src kinase.

1208   2024.04.26.591297 Preprint at https://doi.org/10.1101/2024.04.26.591297 (2024).

1209   71.   Crawford, K. H. d & Bloom, J. D. alignparse: A Python package for parsing

1210   complex features from high-throughput long-read sequencing. *J. Open Source*

1211   *Softw.* **4**, 1915 (2019).

1212   72.   Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. VSEARCH: a versatile

1213   open source tool for metagenomics. *PeerJ* **4**, e2584 (2016).

1214   73.   Martin, M. Cutadapt removes adapter sequences from high-throughput

1215   sequencing reads. *EMBnet.journal* **17**, 10–12 (2011).

1216   74.   Seabold, S. & Perktold, J. Statsmodels: Econometric and Statistical Modeling

1217   with Python. *Proc. 9th Python Sci. Conf.* 92–96 (2010) doi:10.25080/Majora-

1218   92bf1922-011.

1219   75.   Loes, A. N. *et al.* High-throughput sequencing-based neutralization assay

1220   reveals how repeated vaccinations impact titers to recent human H1N1 influenza

1221   strains. *J. Virol.* **98**, e00689-24 (2024).

1222

a

Inactive (2RH1)　　　　　　Active (3SN6)

b

Ligand binding (D113^{3x32}, V114^{3x33}, V117^{3x36}, T118^{3x37}, A119^{3x38}, I121^{3x40}, S165^{4x57}, Y199^{5x39}, F193^{45x52}, S203^{5x43}, S207^{5x461}, F208^{5x47}, Y209^{5x48}, F290^{6x52}, N293^{6x55}, I294^{6x56}, F289^{6x51}, N312^{7x38}, G315^{7x41}

M82^{2x53}

Sodium binding pocket (A76^{2x47}, A78^{2x49}, D79^{2x50})

I127^{3x46}

NPxxY (N322^{7x49}, I325^{7x52}, Y326^{7x53})

CWxP (W286^{6x48})

PIF (I121^{3x40})

N318^{7x45}

Y219^{5x58}

V129^{3x48}

A128^{3x47}

Transducer binding
I135^{3x54}, T136^{3x55}, C327^{7x54}, S329^{8x47}, P330^{8x48}

DRY (R131^{3x50})

a 12x serial isoP dilutions

10 µM
2.8 µM
816 nM
233 nM
66 nM
19 nM
5.4 nM
1.5 nM
444 pM
126 pM
36 pM
10 pM

GPCR-MAPS

Collect RNA

b Estimate RNA abundance per condition (qPCR)

c Per-variant normalized score: variant freq / condition RNA abun.

Illumina sequence for per-variant frequencies

Signaling magnitude

d

A V I L M F Y W S T N Q C G P R H K D E * del
Median

10 uM
2.8 uM
816 nM
233 nM
66 nM
19 nM
5.4 nM
1.5 nM

N-term ICL1 TM1 TM2 ECL1 TM3 ICL2 TM4 ECL2 TM5 ICL3 TM6 ECL3 TM7 C-term

e

1. Wild-type-like
2. Mod. damaging
3. Loss-of-function

10 pM → 10 µM

f Wild-type

$E_{max} = 1$
$E_{min} = 0$
$EC_{50} = 26.2$ nM
slope = 0.84

g Wild-type-like

n = 4,886 vars

h Mod. damaging

n = 674 vars

i Loss-of-function

n = 28 vars