From Morphology to Computation: How Synaptic Organization Shapes Place Fields in CA1 Pyramidal Neurons

Simone Tasciotti[1,2], Daniel Maxim Iascone[3], Spyridon Chavlis[2], Luke Hammond[3,#], Yardena Katz[3], Attila Losonczy[3,4], Franck Polleux[3,4], Panayiota Poirazi[2,5]

[1] Department of Biology, University of Crete, Heraklion, Greece

[2] Institute of Molecular Biology and Biotechnology, Foundation for Research and Technology Hellas, Heraklion, Greece

[3] Mortimer B. Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY, USA

[4] Department of Neuroscience, Columbia University, New York, NY, USA

[5] Lead contact

[#] Present address: Department of Neurology, The Ohio State University, Wexner Medical School, Columbus, Ohio, USA

* Corresponding author

Abstract

The synaptic mechanisms driving feature selectivity in specific neuron types remains a fundamental and unresolved challenge in neuroscience. In hippocampal CA1 pyramidal neurons (PNs), the development of place selectivity, manifested as place fields, is believed to result from dendritic integration of spatially distributed inputs combined with behavioral time scale plasticity (BTSP). BTSP involves dendritic spikes that temporally regulate synaptic potentiation and depotentiation. However, the role of excitatory (E) and inhibitory (I) synaptic distributions in the emergence of place specificity in CA1 PNs remains unclear, due to the lack of detailed synaptic reconstructions in vivo. Here, we present full synaptic reconstructions from individual CA1 PNs in the mouse hippocampus, revealing that these neurons receive approximately 10,000–15,000 E synapses and 900–1,400 I synapses. Computational modeling of biologically relevant E and I synaptic distributions shows that spatial tuning is preserved with co-tuned, spatially clustered E synapses, but disrupted when E distributions are randomized. Moreover, synapse clustering has a different contribution to spatial tuning in apical vs. basal domains. Our results reveal a complex and finely-tuned interplay between presynaptic input patterns and the spatial organization of their postsynaptic targets in dictating neuronal output in CA1 PNs.

Keywords

dendritic computation, synaptic clustering, CA1, pyramidal neuron, biophysical model, morphology, dendritic nonlinearity, dendritic domains

Introduction

The hippocampus and particularly its CA1 region plays a crucial role in spatial navigation. CA1 pyramidal neurons (PNs), specifically the spatially tuned 'place cells', are widely hypothesized

to construct cognitive maps of the environment by combining spatial information and

contextual cues[1-3]. Studies have shown that place cells are highly dynamic, undergoing

remapping when an animal encounters a new or altered environment[2,4]. During remapping,

CA1 place cells reorganize their spatial firing patterns in response to environmental changes,

effectively encoding a new cognitive map[5]. In familiar settings, place cells exhibit stable firing

fields, supporting the retrieval of established memories, while in novel contexts, rapid

remapping enables the encoding of new spatial information.

This flexibility stems from the integrative properties of CA1 PNs, which allow them to adjust

the salience of specific input streams based on the environmental context[6,7]. However, the

electrophysiological and structural variability of these neurons hinders our in-depth

understanding of their computational capabilities[8]. Numerous studies have attempted to

characterize how these properties, along with the organization of synaptic inputs, collectively

shape the input-output dynamics of CA1 PNs[8-14]. Yet, the impact of the spatial distribution of

E and I synaptic inputs on the formation and functional features of place cells remain largely

unexplored. Synaptic organization along the dendritic arbor heavily influences the integrative

properties of neurons. For example, projections from the CA3 region predominantly target

proximal perisomatic dendrites, while long-range inputs from the entorhinal cortex (EC)

innervate the distal apical dendrites[15,16]. This anatomical segregation allows both the

independent processing of distinct information streams and their conditional integration, thus

dynamically shaping the neuron's feature selectivity[17-19]. Additionally, differential ion channel

distributions and morphological features of dendrites across these CA1 layers modulate

synaptic efficacy, such that identical inputs can have variable impacts on neuronal output

depending on their dendritic location[20,21]. A comprehensive exploration of how different E and

I incoming inputs shape the local dendritic properties is essential for elucidating the

mechanisms underlying neuronal information processing.

Previous results suggested that rat CA1 PNs are characterized by non-uniform synaptic

distributions at the sub-branch level[9]. This type of synaptic arrangement can greatly affect the

dendritic input-output function[22,23]. For example, synchronous activation of synapses within

the same branch was shown to drive strong nonlinear dendritic responses[24–26], a property that

generalizes to other cell types[27–29]. These nonlinear dendritic responses were eliminated when

the same synapses were distributed across different dendrites[24,26], highlighting the importance

of synaptic distribution on CA1 input-output mapping.

The abovementioned effects are underlined by voltage-dependent ionic conductances in

dendrites, which enable the induction of local regenerative events, turning the neuron into a

complex processing system with enhanced computational capabilities[22,24,26,30–32]. As the

spatio-temporal organization of synaptic inputs is essential for triggering these local events,

synapse location truly matters[24,26,32]. Localized within dendritic compartments that host

numerous nonlinear mechanisms, synapses engage in dynamic interactions, amplifying,

attenuating, or gating signals based on their dendritic positions, which suggests that the

formation and maintenance of synaptic contacts is not a randomized process but an actively

regulated mechanism[33–38]. Recent findings further reveal that co-tuned inputs frequently

converge onto spatially clustered synapses, creating dendritic "hotspots" that can more easily

produce non-linear dendritic events[39,40], as suggested by earlier theoretical work[26,32,41].

Therefore, it is clear that to fully understand how CA1 PNs compute a place field, it is essential

to consider the spatiotemporal organization of their incoming inputs.

In this study, we combined experimental and computational modelling approaches to map the

spatial distribution of all E and I synapses in a subset of mouse CA1 PNs in order to investigate

how the spatial distribution of E and I synapses affects the output of spatially tuned CA1 PNs.

We first employed an advanced sparse labeling technique to reconstruct the complete

morphologies of CA1 PNs with high precision. We meticulously traced the dendritic

arborizations and mapped the locations and morphological features of both inhibitory and

excitatory synapses. We found a non-uniform distribution of excitatory synapses across the

dendritic arbor, in contrast to a more homogeneous arrangement of inhibitory synapses.

Specifically, terminal dendrites in both apical and basal dendritic domains had higher spine

density than intermediate ones, which resulted in a higher probability of finding large spines

located in close proximity (i.e., spine clustering) within terminal dendritic compartments. These

data were used to build biophysical models that were extensively validated against

electrophysiological data and used to investigate how the spatial distribution of synaptic inputs

influences the output of place cells during simulated exploration of a linear track. Models

predicted that the spatial allocation of excitatory inputs strongly influences both the spatial

tuning and the place cell properties of simulated neurons. Specifically, spatial tuning that was

present in simulations using co-tuned and spatially clustered excitatory synapses but was

disrupted when target locations were randomized. Moreover, our models suggest that there

might be a set of optimal, domain-specific input patterns that impinge on the reconstructed

synapses and turn these neurons into high quality place cells. Specifically, we predict that

inputs to the distal apical tuft are more effective when activated in large clusters whereas

inputs to the basal dendrites are more effective when synaptic clusters are distributed over

multiple branches. Together, our findings reveal a complex and finely-tuned interplay between

presynaptic input patterns and the spatial organization of their postsynaptic targets in

specifying the output properties of CA1 PNs.

Results

Reconstruction of CA1 Pyramidal Neuron Morphology

To reconstruct the dendritic morphology and synaptic distribution of all excitatory and inhibitory

synapses of mouse dorsal CA1 PNs at single-cell resolution *in vivo*, we deployed a

combination of sparse reporter expression, imaging, and computational tracing methods that

we previously developed[42]. We labeled dorsal CA1 PNs through *in utero* electroporation of

plasmids expressing Cre-dependent Flex-tdTomato (red cell filler) Flex-EGFP-Gephyrin

(inhibitory synaptic marker) along with low levels of a Cre recombinase plasmid for targeted,

sparse labeling[43,44] (**Figure 1A**). After high-resolution confocal imaging (**Figure 1B**), we used

open-source reconstruction software Vaa3D[45] to obtain a trace of the dendritic tree (**Figure**

1C). Through the Vaa3D Synapse Detector toolkit[42], we annotated all excitatory and inhibitory

synapses located within each segment of the dendritic tree. Given that the majority of

excitatory synapses in CA1 PNs occur on spines[9], we used them as a structural proxy of all

excitatory synapses in our neurons. Inhibitory synapses were identified by co-localizing EGFP-

Gephyrin puncta with cytosolic tdTomato (**Figure 1B**). Using this method, we obtained detailed

information about each synapse's position and shape, including features like volume, spine

head diameter, and spine neck length. We also identified the relative location of inhibitory

synapses, whether on the dendritic shaft or a dually innervated spine. Finally, Vaa3D

algorithms linked all synaptic features to the corresponding branch segment node, and neuron

trace fragments of the same neuron from serial tissue sections were stitched together to create

complete morphological reconstructions. Through this method, we obtained the dendritic arbor

morphology and excitatory (**Figure 1D**) and inhibitory (**Figure 1E**) synapse mapping of six

CA1 PNs (with one that lacks the inhibitory synapse map). After implementing the synaptic

reconstruction pipeline, we applied a custom smoothing algorithm to dendritic diameter

measurements to eliminate minor artifacts introduced by the reconstruction software (**see**

Methods, Figure S1). From the structural analysis of these neurons, we observed, on

average, 11,796±2,212 (mean±std, n=6 neurons, std: standard deviation) spines and 741±

545 inhibitory synapses, distributed over 7,164±982 μm of average total dendritic length.

Consistent with previous studies[46], the average spine density was estimated to be 1.41±0.18

while the inhibitory synapse density was 0.10±0.07[47] (**Figure 1F**).

Figure 1: Reconstruction and analysis of all excitatory and inhibitory synapses across CA1 pyramidal neurons. A. Sparsely labeled pyramidal neurons expressing tdTomato (red) and EGFP-Gephyrin (green) in hippocampal CA1. Grey: tissue autofluorescence using 488nm excitation. Scale bar: 300 μm. **B**. Single-synapse resolution image of a CA1 pyramidal neuron expressing tdTomato (red) and EGFP-Gephyrin (green). Scale bar: 60 μm. **C**. Dendritic arbor of whole CA1 PN shown in panel B, annotated to highlight apical tuft dendrites (magenta), apical oblique and apical trunk dendrites (blue) and basal dendrites (yellow). **D-E**. All dendritic spines (excitatory synapses; D, green) and Gephyrin+ inhibitory synapses (E, purple) annotated throughout the dendritic arbor (grey) of the example CA1 PN reconstructed within insets from (B-C). Scale bars: 50 μm. **F.** Quantitative analysis of all CA1 PNs reconstructed showing dendritic structure and synaptic distribution. From left to right: total dendritic length

155 distributions (μm); total number of excitatory synapses and their density (syn/μm); total number of inhibitory
156 synapses and their density (syn/μm); percentage of inhibitory synapses localized on a spine (dually innervated
157 spines).

Domain-Specific Analysis of Synaptic Organization in CA1 PNs

CA1 PNs exhibit significant variability in the structural and electrophysiological properties of
their dendritic arbors, which are also targeted by different presynaptic sources[8,9,48,49].
Consequently, we opted to analyze the synaptic organization by subdividing the dendrites into
four distinct domains, namely *Basal*, *Apical Trunk*, *Oblique*, and *Tuft* dendritic domains. Within
each domain, we further distinguished intermediate and terminal branches based on their
relative branch order, referred to as *subdomains* (**Figure 2A**). We found that peri-somatic and
intermediate dendritic regions exhibit a substantially lower spine density than more terminal
branches (**Figure 2B**), which is in line with the literature[9,46] while inhibitory synapses show no
significant variation across subdomains (**Figure 2B**).

Within these dendritic subdomains, we also characterized the size of synaptic connections in
order to infer the degree of potentiation as a function of the dendritic subdomain. Synaptic
volume corresponds to spine head size or postsynaptic density size for excitatory and
inhibitory synapses, respectively, and is linearly correlated with synaptic strength[22,50,51]. As
described in **Iascone et al.** (**2020**)[42] for layer 2/3 PNs in the mouse primary somatosensory
cortex, we used the Synapse Detector tool to map the distribution of synaptic strengths across
all different subdomains. Large synapses were defined as those exceeding the 80th percentile
of the synaptic volume distribution of each cell. We found similar distributions of large
synapses across dendritic subdomains, with the exception of the terminal Tuft, which exhibited
a greater percentage of large spines compared to its intermediate counterpart (**Figure 2C**).

Another structural characteristic that may indicate an important function is the increase of
synaptic density in a branch-specific way. During learning, new spines tend to form in proximity
to already existing spines, or appear in small clusters, and to be more stable, marking

clustered potentiation at branch level[38,52–56]. To check this possibility, we compared the

experimentally observed variation (standard deviation, in short STD) in E and I synapse

density within dendritic subdomains to the synaptic densities of a subdomain-wise null

distribution (**see Methods**) for each neuron. Dendritic subdomains with STD higher than that

of the null distribution exhibit this branch-specific enrichment. We found branch-specific

enrichment of excitatory synapses in the Trunk and Intermediate Basal domains, whereas

inhibitory synapses were overall uniformly distributed (**Figure 2D**).

Figure 2: Analysis of Dendritic Subdomains and Synaptic Distributions. A. Left: pie charts showing the

percentage of excitatory (E) and inhibitory (I) synapses located in different domains. Right: schematic

representation of a CA1 PN, with distinct dendritic domains highlighted in different colors. These include the Basal

(green), Tuft (pink), Trunk, and Oblique (blue), further subdivided into intermediate and terminal subdomains using darker and lighter colors, respectively. **B.** Density of E and I synapses across different subdomains. **C.** Percentage of large spines (i.e., spines with a volume above the 80th percentile of the size distribution) for each subdomain. **D.** Variation (standard deviation, STD) in E and I synapse densities for each subdomain. Experimentally observed STDs (green/purple) are compared to those expected from a random redistribution of synapses within their specific subdomains (grey). Specifically, synapses were randomly shuffled across branches within the same subdomain, and excitatory/inhibitory synaptic density variation was recalculated. Excitatory -but not inhibitory- synapse variability is significantly different from the shuffled distribution (2-way ANOVA for spines density std: distribution: $F_{(1, 4228)}=374.74$, $p=4.63e-80$; domain: $F_{(6, 4228)}=794.07$, $p=0.00e+00$; distribution x domain: $F_{(6, 4228)}=10.66$, $p=8.71e-12$). **E.** Schematic illustration of the hierarchical clustering algorithm used to identify spine clusters. The algorithm detects groups of large spines (with volumes above the 80th percentile) that are located within 10 micrometers of each other. The dashed line indicates the 10 μm distance cutoff. Red dendrogram strokes represent a spine cluster. **F.** Density of spine clusters across subdomains. Left: experimentally observed distribution (green) vs. a domain-wise shuffled distribution (grey). Right: experimentally observed distribution vs. a branch-wise shuffled distribution. Statistically significant differences are found only between the experimentally observed distribution and the shuffled-per-domain distribution. (2-way ANOVA for cluster density between `Exp. Data` and `Shuffled`: protocol: $F_{(1, 775761)}=136.83$, $p<10^{-30}$; domain: $F_{(6, 775761)} = 3412.82$, $p<10^{-80}$; protocol x domain: $F_{(6, 775761)}=58.33$, $p<10^{-70}$). When shuffling within subdomains, the two distributions substantially overlap, with the exception of the Trunk (2-way ANOVA for cluster density between `Exp. Data` and `Shuffled per subdomain`: protocol: $F_{(1, 775761)}=1.06$, $p=0.303$; domain: $F_{(6, 775761)}=82383.08$, $p<1e^{-6}$; protocol x domain: $F_{(6, 775761)} = 4.58$, $p=-1.19e-04$). Pair-wise two-sided t-tests are shown in all plots. $^*p < 0.05$, $^{**}p < 0.005$, $^{***}p < 0.001$, and $^{****}p < 0.0001$, corrected for multiple comparisons using Bonferroni's method. Error-bands in **B**, **C**, **D**, and **F** represent 95% confidence interval (CI) of the mean.

Given that excitatory synapses are non-uniformly distributed, we next asked whether there is a finer patterning of synaptic connections within the different subdomains. Over the past decades, several studies demonstrated that the volume of spines tends to increase with the strengthening of the synapse[57–59]. Spines, however, do not grow in isolation but rather interact with neighboring synapses, often resulting in the formation of the so-called synaptic clusters. Indeed, several studies investigated how non-linear dendritic mechanisms promote plasticity

of nearby excitatory synapses[24,54,55,60,61]. For this reason, we investigated the presence of

structural clusters in the different dendritic subdomains. To do so, we first defined a putative

structural cluster as a group of three or more large (as defined previously) synapses within a

branch segment of ~10 μm[62]. This approach is based on the assumption that heterosynaptic

plasticity drives co-tuned nearby synapses to grow together and deviate from the average

size[60,63–66]. We used a hierarchical clustering algorithm for the identification of these putative

structural clusters (**Figure 2E**). Our analysis revealed a higher density of structural clusters in

the terminal domains, a result that was significantly different from the density distribution

obtained through domain-wise shuffling of synapses (**Figure 2F**). However, when comparing

these observations with a subdomain-wise null distribution, the differences were no longer

significant (**Figure 2F**). This finding suggests that the observed variations in cluster density

between domains are primarily driven by differences in synaptic density within the

corresponding subdomains.

Taken together, our results show that the spatial distribution of excitatory synapses in CA1

PNs is not random, showing a clear enrichment for terminal branches in basal and apical

oblique dendrites, but not in apical tuft dendrites. However, this fine, domain-specific

organization is not found for inhibitory synapses, creating an asymmetric E/I balance.

From Reconstruction of Synaptic Distribution to Modeling of Neuronal

Functional Properties

To investigate how E and I synaptic distribution may influence the emergence of spatial tuning

in CA1 pyramidal neurons, we constructed a biophysical model for each reconstructed

neuronal morphology (**Figure 3A**) and simulated their activity during a virtual linear track

exploration task (**Figure 3B**). The models were implemented using the NEURON simulation

environment[67], adhering to the channel distribution parameters established by **Poirazi et al.**

2003[26] **(see Methods)**.

Given the high synaptic density in our reconstructed neurons, we employed a fine spatial

discretization by increasing the number of segments in the models. This ensured enhanced

spatial resolution, enabling us to capture the detailed biophysical interactions between

adjacent synapses (**Figure 3C**). To ensure the accuracy and biological relevance of our

models, we validated both their passive and active properties against experimentally observed

values. Passive properties, including input resistance (R_{in}), membrane time constant (τ_m),

Sag Ratio, and dendritic attenuation, were within physiological regimes[8,68,69] (**Figure 3D-4A;**

see Methods). Similarly, the active properties, such as somatic and dendritic spiking activity,

Frequency-Current (F-I) curve, and dendritic nonlinearities, were validated with widely used

protocols[8,24] (**Figure 4B-E; see Methods**).

Figure 3: From morphologies to biophysical models. A. Reconstructions of dendritic morphology of the five

CA1 pyramidal neurons used to generate the computational models. **B.** Somatic voltage traces produced by a

place-tuned stimulation protocol used on the corresponding neuronal reconstruction. **C.** Left: schematic illustration

of the cell's segmentation used in the NEURON simulation environment (obtained using DendroTweaks[70]). Right:

example of a branch structure from the data and its corresponding segmentation in the simulator. **D.** Passive

membrane properties of CA1 PN models match experimental values. From left to right: input resistance, membrane

time constant, and sag ratio. Experimental barplots redrawn from Masurkar et al., 2020[8].

The excitatory synaptic weights were calibrated based on experimental findings to ensure

physiological relevance[71]. Specifically, we implemented the principle of synaptic democracy,

which posits that synaptic inputs across the dendritic tree contribute equally to somatic

depolarization, regardless of their location[72]. To achieve this, proximal synapses were

parameterized to elicit a 0.2 mV depolarization at the soma[20]. For distal synapses, the α-

amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptor conductance was

progressively increased as a function of the distance from the soma[71,73]. This adjustment

ensured that while distal synapses generated larger dendritic excitatory postsynaptic

potentials (EPSPs), the resulting somatic depolarization remained constant at 0.2 mV (**Figure**

4F; **see Methods**). Moreover, considering the large variability of spine sizes observed in our

analysis, we decided to account for it in the calculation of the synapse strength by linearly

varying the validated weight according to the spine head volume[74] (**see Methods**).

Inhibitory synapses were validated differentially based on their presynaptic sources to ensure

an accurate representation of their functional characteristics. For this study, inhibitory

synapses were assigned to six distinct locomotion active interneuron types, neurogliaform

(NGF), oriens-lacunosum moleculare (OLM), parvalbumin-expressing basket (PVbasket),

apical dendrite targeting cholecystokinin-expressing (CCK), Bistratified, and Ivy cells, based

on their known post-synaptic target distributions across the dendritic tree[75–77]. Synaptic

weights, as well as rise and decay times, were calibrated by replicating experimental protocols

reported in the literature[78–82] (**Figure 4G**). Similar to their excitatory counterparts, inhibitory

synapses were also scaled linearly with the synaptic volume. These rigorous validations

enhanced the reliability of the models and ensured they accurately reflected the physiological

characteristics of CA1 PNs.

Figure 4: Electrophysiological Properties of CA1 PN Models Match Experimental Data. A. Dendritic

attenuation of somatically injected negative current in control conditions (red) or simulating extracellular cesium

(grey, 80% reduction of the h-current). Experimental data redrawn from Golding et al., 2005 are similar to model

outputs. **B.** Examples of somatic voltage traces induced by a 300 pA current injection in the soma (left) or in a

dendrite (right) located at ~250 μm from the soma. **C.** The input-output (Frequency-current (F-I)) curves of the

models resemble experimental ones from Masurkar et al., 2020[8]. **D.** Peak amplitude of EPSPs measured at the

soma as a function of the expected arithmetic sum of individual EPSPs. Experimental example curve redrawn from

Losonczy and Magee, 2006[24]. **E.** Somatic depolarization upon simultaneous excitation of a progressively increasing

number of synapses. Left: stimulation of 1 to 30 simulated synapses on an oblique dendrite. Right: spines

stimulated through glutamate uncaging, redrawn from Losonczy and Magee, 2006[24]. **F.** Dendritic and somatic

excitatory postsynaptic potentials (EPSPs). Synaptic AMPA conductance strength increases with path distance

from the soma, ensuring invariant somatic depolarization. **G.** Inhibitory postsynaptic currents (IPSCs) originating

from various interneurons, including NGF (neurogliaform cells), OLM (oriens-lacunosum moleculare cells),

PVbasket (parvalbumin-expressing basket cells), CCKbasket (cholecystokinin-expressing cells), Bistratified and

Ivy cells. Validated to match experimentally observed data. Errorbars in **C**, and **F** represent 95% confidence interval

(CI) of the mean.

After validating the passive and active properties of the individual PNs, we constructed the

inputs to reliably reproduce the dynamics of incoming synaptic activity under a linear track

exploration task. To achieve this, we produced spike trains that simulate excitatory inputs

originating from CA3 and the entorhinal cortex (EC), capturing the neural patterns associated

with spatial navigation. Specifically, we modeled convergent EC inputs as multiple grid

representations covering a virtual linear track, from which CA3 place-like inputs were derived[6]

(**see Methods**). The two resulting input streams were subdivided into a total of 12 place fields,

each containing hundreds of co-tuned spike trains. In addition, background synaptic noise was

simulated by introducing independent Poisson-distributed, and Gaussian-filtered inputs to

CA1 PNs (**Figure 5B**).

Inhibitory inputs, conversely, were modeled as theta-filtered, phase-shifted Poisson spike

trains, with each presynaptic source tailored to match the specific characteristics reported in

the literature[76] (**Figure 5C; see Methods**). All inputs were distributed across synapses located

in the anatomically relevant regions of the dendritic tree, ensuring that the simulated activity

closely approximated the complex synaptic interactions observed during spatial navigation

(**Figure 5A**). In particular, each interneuron targeted a synapse based on a probability derived

from a non-linear function that depends on the distance from the soma. (**Figure S2**).

Figure 5: Patterns of Excitatory and Inhibitory Inputs to CA1 Pyramidal Cells. A. Schematic representation of

the model neuron depicting excitatory and inhibitory input sources that are organized according to anatomical

findings. **B.** Different types of excitatory inputs to the model. From top to bottom: Heatmap of EC grid-like, CA3

place-like and random noise spike trains. Each row represents a different presynaptic input neuron. The colormap

is the same across heatmaps and brighter colors denote higher firing rates. **C.** Population activity of simulated

interneuron spike trains compared to the Excitatory Theta Oscillation. Firing frequencies and theta phases are

based on experimental findings.

338 Functional Role of Input Clustering in Place Cell Computation

After confirming that our model accurately reproduces the biophysical and electrophysiological

properties of CA1 PNs given realistic synaptic input dynamics, we set out to investigate how

spatial organization of excitatory synaptic inputs influences place-tuning properties.

Specifically, we investigated how the clustering of excitatory synaptic inputs alters the

computational capabilities of the neuron in a simulated navigational task.

A recent glutamate imaging study has characterized the spatiotemporal patterns of synaptic

inputs to CA1 place cells during navigation on familiar linear tracks[39]. This study revealed that

synapses receive inputs across all track locations, but the excitatory drive is significantly

higher within the somatic place field than outside it. Additionally, the synapses that are co-

tuned with the soma exhibit a greater tendency to form spatial clusters. Given these findings,

we generated 1,000 simulated connectivity patterns for each neuronal reconstruction

designed to emulate the spatial organization of synaptic inputs that have been observed

experimentally. Specifically, each connectivity pattern included a subset of structural clusters,

as defined previously, which were preferentially targeted by place-tuned inputs. Inputs

associated with a selected place field targeted a higher number of clustered synapses, which

determined the somatic place-tuning, while the remaining synapses either received place-

tuned or noisy inputs based on a uniform distribution.

This allocation protocol resulted in two distinct synaptic populations: (1) co-tuned clustered

synapses, referred to as *Functional Clusters*, which were exclusively place-tuned, and (2)

dispersed synapses, which were either place-tuned or targeted by noisy inputs. This

configuration represented our Clustered condition (**Figure 6A, Control**). To assess the role of

synaptic clustering in neuronal computation, we generated alternative synaptic distributions

by shuffling the input targets under three different constraints. First, we generated a

configuration where the total number of activated synapses was preserved, but inputs were

randomly reassigned, disregarding the original preference for clustered synapses, which are

larger and stronger (**Figure 6A, Shuffled** – Same Synapse Number (SSN)). Second, since

synaptic strength scales with size, we generated a configuration in which the total synaptic

volume rather than the number of inputs was preserved. This ensured a closer approximation

of the excitatory drive comparable to the Clustered configuration but caused a much larger

number of synapses to get activated (**Figure 6A, Shuffled** – Same Synapse Volume (SSV)).

Lastly, we generated a configuration where the cumulative synaptic weight was preserved.

This configuration considers the volume while also accounting for the "synaptic democracy"

adjustment where synaptic conductance scales with the distance from the soma (**Figure 6A,**

Shuffled – Same Synapse Weight (SSW)). These three shuffled configurations allowed us to

disentangle the influence of synaptic clustering, size, and weight in shaping neuronal place

computations.

To verify that the shuffling protocols disrupted input clustering, we quantified the number of

functional synaptic clusters across dendritic domains for each input allocation scheme. As

expected, the Clustered configuration exhibited a significantly higher number of functional

clusters across all dendritic domains, particularly when compared to the SSN condition, which

preserved the number of inputs but disrupted their spatial organization (**Figure 6B**). The SSV

protocol, while maintaining a relatively high number of functional clusters due to the increased

number of activated synapses, resulted in clusters with a smaller average size and lower total

synaptic volume (**Figure S3**). These findings confirm that random shuffling alters the structural

properties of synaptic clustering.

Next, we evaluated how these different input allocation protocols influenced the simulated

activity of CA1 PNs. We found that neurons in the Clustered condition exhibited the

characteristic place-field activity, while all shuffling protocols led to its disruption (**Figure 6C**).

To quantify the extent of disruption, we assessed specific place field metrics such as in-field

and out-field firing frequency, field size and spatial information[83]. We found that only the

Clustered configuration generated place-field features, such as In-Field Firing Frequency

(**Figure 6D**) and Spatial Information (**Figure 6E**), that aligned qualitatively with experimental

results from **Mizuseki et al.** (**2012**)[84]. The SSN configuration resulted in substantially lower

firing activity, leading to a marked reduction in Spatial Information. In contrast, both the SSV

and SSW configurations led to a much higher firing frequency, with increased spike

occurrences outside the place field. This widespread activity diminished the spatial selectivity

of neuronal firing, ultimately reducing the overall Spatial Information. These results suggest

that synaptic clustering plays a critical role in maintaining the spatial tuning of place cells and

perturbing clustering disrupts single-cell encoding of spatial information.

Figure 6: Effect of Synapse Clustering on Spatial Tuning. A. Schematic representation of four different synaptic configurations. Colors represent the tuning of synaptic inputs. Clustered: Synapses within pre-identified structural clusters are more likely to receive co-tuned inputs. Shuffled - Same Number: Inputs are randomly distributed while maintaining the same number of activated synapses as in the Clustered configuration. Shuffled - Same Total Weight: Inputs are randomly distributed while preserving the total synaptic strength of the activated inputs as in the Clustered configuration. Shuffled - Same Total Volume: Inputs are randomly distributed but the total volume of activated synapses remains the same as in the Clustered configuration. **B.** Number of activated functional synaptic clusters (co-tuned synapses located within a 10 μm spatial window) across dendritic domains (Basal, Apical, and Tuft) for each configuration. Statistically significant differences are indicated (2-way ANOVA for the number of Functional Clusters in the Tuft domain: cluster protocol: $F_{(3, 16)}=5.16$, $p=1.10e-02$; Apical domain: cluster protocol: $F_{(3, 16)}=142.96$, $p=9.19e-12$; Basal domain: cluster protocol: $F_{(3, 16)}=19.16$, $p=1.52e-05$). Pair-wise t-test between Control and Shuffled protocols are shown in the plot. $^*p < 0.05$, $^{**}p < 0.005$, $^{***}p < 0.001$, and $^{****}p < 0.0001$, corrected for multiple comparisons. Error-bars represent 95% confidence interval (CI) of the mean. **C.** Average activity heatmaps showing place tuning of a population of CA1 PN models stimulated under the different

synaptic configurations. Input shuffling significantly disrupts the output place fields. **D.** Distribution of in-field firing

frequencies. Top: The Clustered configuration qualitatively replicates the experimentally observed spatial firing

patterns of CA1 PNs as reported by Mizuseki et al. (2012)[84]. Bottom: Firing frequency distributions of the shuffling

configurations. **E.** Distribution of spatial information content (measured in bits per spike). Top: The Clustered

configuration aligns qualitatively with the experimentally observed spatial firing patterns described by Mizuseki et

al. (2012)[84]. Bottom: Spatial information distribution for the shuffling configurations.

Dendritic domain-specific clustering differentially contributes to spatial tuning

While global synaptic clustering alters the place-specific properties of our model CA1 PNs, its

role in domain-specific integration remains unclear. To address this question, we probed the

contribution of domain-specific clustering features to neuronal output. Specifically, we asked

how synaptic clusters which are co-tuned with the soma ("in-field" clusters), and are located

in the Basal, Apical or Tuft dendrites, contribute to neuronal spatial tuning. We focused on

three clustering features: the *Total Cluster Volume* (the cumulative volume of all synapses

within in-field-tuned clusters), the *Average Cluster Size* (the mean number of synapses of

these clusters), and the *% Dendrites with In-field-Tuned Clusters*, thereby capturing both the

strength and spatial distribution of synaptic inputs (**Figure 7A**).

We then applied a machine-learning regression algorithm to evaluate how well these features

can predict the spatial tuning properties previously mentioned (**Figure 7B**). While the model

accurately predicted in-field and out-field firing rates, suggesting that the structural properties

of synaptic clusters strongly influence firing frequency, it poorly predicted field size, indicating

that this property may involve more complex mechanisms (**Figure 7C**).

Notably, analysis of feature importance revealed domain-specific strategies for influencing

somatic output. In the Tuft domain, *Total Cluster Volume* was the most critical predictor of

spatial tuning quality. This suggests that larger clusters or bigger spines may be required to

drive local spikes, thus overcoming signal attenuation caused by a large electrotonic distance

from the soma (**Figure 7D**) and exert a significant influence on somatic firing[85–87]. In contrast,

for the Basal domain, the *% Dendrites with In-field-Tuned Clusters* was the most important

predictor of spatial tuning quality. These findings suggest that parallel input streams across

multiple branches may represent an optimal strategy for reliably modulating somatic output,

highlighting the integration advantages conferred by proximity to the soma (**Figure 7D**). This

interpretation is further supported by the application of the SHAP (SHapley Additive

exPlanations) framework to explain the output of our Random Forest results[88]. This approach

provides a per-feature and per-sample attribution to each individual predictor, highlighting the

contribution of each input to the corresponding output value (**Figure S4**).

Interestingly, none of the three clustering features could reliably predict somatic output for the

Apical domain, a region that in our analysis encompassed both Trunk and Oblique domains.

This suggests that the apical compartment may rely on alternative mechanisms to shape

spatial tuning that are not captured by the abovementioned clustering features.

Figure 7: Cluster features Predict Spatial Tuning Quality *A.* Cluster features used as input to the multivariate random forest model to predict spatial tuning quality. The latter is assessed by the spatial information, out-field firing frequency, Infield firing frequency and field-size metrics. For each dendritic domain (Basal, Apical, and Tuft), cluster features include: the volume of in-field clusters, the size of in-field clusters, and the percentage of branches containing at least one in-field cluster. **B.** Schematic representation of the random forest methodology. The model combines predictions from multiple decision trees, each trained on subsets of the data (800 simulations used for training, 200 used for testing), to enhance robustness and accuracy in predicting model outputs. **C.** Mean Squared Error (MSE) of the output predictions on the testing set, providing a quantitative measure of the model's prediction accuracy. **D.** Feature importance in predicting simulation outputs for the Tuft, Apical, and Basal dendritic domains. The dashed line indicates the chance level of feature importance (~1.11). Error-bars represent the 95% confidence interval (CI) of the mean.

Together, these findings suggest that synaptic integration in spatially-tuned CA1 PNs is not

uniform across dendritic domains. The Tuft appears to capitalize on large, high-volume

clusters to initiate powerful nonlinear events, whereas the Basal domain favors a strategy of

parallelized integration across multiple dendrites to collectively drive somatic depolarization.

The apical domain's mechanisms remain more enigmatic, hinting at a rich, multidimensional

mode of integration that may involve other variables such as temporal coordination or dynamic

gating of the inhibitory inputs. These findings advance our understanding of how the

organization of synaptic inputs shapes neuronal processing and underscore the importance

of dendritic compartmentalization in spatial navigation.

Discussion

In this study, we combined high-resolution anatomical mapping with biophysical computational

modeling to investigate how the spatial distribution of excitatory synapses influences the

integrative properties of spatially-tuned CA1 PNs. Although the structural and

electrophysiological characteristics of these neurons have been well-documented *ex vivo*[8–10],

linking detailed anatomical properties of both E and I synapses to functional outcomes has

remained challenging. Our study reveals that excitatory synapses onto CA1 PNs are

organized in a non-uniform, domain-specific manner across the dendritic tree, and the

patterned activation of these synapses has a marked impact on the place-tuning properties of

these neurons.

Specifically, our comprehensive mapping of morphological and synaptic features reveals

marked differences in spine density between dendritic domains. In line with prior work, we

found that terminal dendrites exhibit a higher spine density compared to intermediate

dendrites, whereas inhibitory synapses are distributed uniformly throughout the dendritic tree[9].

As a result, the probability of finding spine clusters is much higher in terminal dendritic

compartments as opposed to intermediate ones. This disparity raises important questions

about the mechanisms that enable the preferential formation of spines toward terminal

regions. We propose that the smaller diameter and sealed ends of terminal dendrites lead to

elevated depolarization, thereby increasing the likelihood of synaptic plasticity events and

subsequent spine formation, a specialization that may enhance the neuron's ability to encode

spatial information[10,89,90]. This hypothesis is supported by prior modelling and experimental

work, whereby activation of inputs in terminal dendrites facilitates the induction of NMDA

spikes, resulting in persistent firing in L5 prefrontal cortical neuron models[91]. Similarly, in L2/3

pyramidal neurons of the primary visual cortex, NMDA spikes in the tuft dendrites facilitate

orientation tuning[92]. Models predict that such NMDA spikes are more effectively induced by

co-activated spines that are organized in clusters[93].

Previous studies inferred patterns of synaptic organization by analyzing E and I synaptic

density in individual branches and extrapolating respective distributions by multiplying

synaptic density to cumulated dendritic length[9]. Instead, our approach provides, for the first

time, a complete, neuron-wide reconstruction of E and I synapses in multiple mouse CA1 PNs,

as previously done for mouse layer 2/3 cortical PNs[42]. This unprecedented level of detail

allowed us to capture the full statistical distribution of synapses across entire dendritic trees,

revealing organizational principles that might have been overlooked in more localized

analyses. Prior work has focused primarily on identifying synaptic clusters by observing newly

formed spines during behavior[33,35,38]. These studies, while more relevant for linking synaptic

clustering to behavioral outcomes, lack accessibility to synaptic clustering at the whole-cell

level, as imaging techniques allow visualizing just a small proportion of dendrites *in vivo*. Our

dataset provides access to high-resolution anatomical features of all E and I synapses found

within all dendritic compartments of a CA1 PN. While providing only a static snapshot of

synaptic structure, it allows us to probe the functional impact of synaptic structure on single-

neuron computations, something not possible without this detailed information.

In addition to the structural mapping of E and I synaptic distribution across the entire dendritic

arbor, our work has generated heavily validated computational models of 5 different

reconstructed CA1 PNs using a common set of biophysical parameters. These models

account for both E and I synapse distributions and their respective input dynamics and serve as powerful tools for investigating questions that go beyond place cell tuning. Several important predictions came out of our models: first, we predict that place cells are more efficiently generated (higher quality place tuning) when co-tuned inputs are delivered to clusters of large spines. Indeed, randomizing the allocation of co-activated synapses -while keeping their number, total volume or total strength fixed-, resulted in degraded place field activity. This finding is likely due to enhanced signal propagation, reduced signal-to-noise and increased energy efficiency by activating spines in clusters[41,94,95]. Achieving the same, spatially-tuned, somatic output with small and randomly distributed synapses would require more contacts, increasing energy consumption and reducing available space for additional synapses[96,97]. This strategy aligns with principles of economical wiring and metabolic efficiency, suggesting that neurons may have evolved to optimize both their structural and functional features in order to better solve complex cognitive tasks[98].

Second, our models predict that the influence of synaptic clustering on neuronal spatial tuning varies depending on the dendritic domain. While tuft dendrites may require larger clustered synapses to impact somatic activity, in basal dendrites, distributed integration of multiple clustered inputs plays a more important role. These observations are in line with studies demonstrating strong compartmentalization of dendritic processing in CA1 PNs both *in vitro*[68,85,99] and *in vivo*[14,100]. In particular, clusters of strong, NMDA-rich synapses primarily found in the apical tuft, which are capable of generating plateau potentials, are more likely to contribute to hippocampal feature selectivity during navigation than basal synapses[99–101]. The activation of these large events is also known to initiate global plasticity events like the newly found BTSP[102,102,103].

Notably, the proportion of basal dendrites containing in-field clusters emerged as the most critical factor influencing spatial tuning. Previous studies have reported a strong correlation between basal dendritic and somatic activity, suggesting that neurons parallelize basal inputs to more efficiently regulate somatic output[14,104]. Additionally, the strong coupling between

basal dendrites and the soma may allow backpropagating action potentials to more effectively

induce plastic changes, aligning basal dendritic tuning with somatic responses[14,105,106].

In sum, our results provide an integrated approach whereby detailed anatomical mapping and

computational modeling are combined to unravel how E and I synaptic organization and its

driving forces, which operate at the subcellular scale, shape cellular computations underlying

spatial navigation. Specifically, it highlights the complex interplay between co-tuned activation

of excitatory synapse clusters and dendritic domain properties in modulating place cell tuning.

By linking such morphological features to computational outcomes, this work advances our

understanding of the fundamental mechanisms through which structural and functional

synaptic distributions modulate neuronal processing.

Resource availability

Lead contact

Further information and requests should be directed to and will be fulfilled by the lead contact,

Panayiota Poirazi poirazi@imbb.forth.gr

Materials availability

This study did not generate new, unique reagents.

Data and code availability

The source code that generates all simulations and Figures, as well as the data that support

this study will be freely available upon publication on GitHub.

Acknowledgments

We thank all members of the PoiraziLab for their valuable feedback on the project. This project
has received funding from the European Union's Horizon 2020 research and innovation
programme under the Marie Skłodowska-Curie grant agreement No.860949; Stavros
Niarchos Foundation (SNF) and the Hellenic Foundation for Research and Innovation
(H.F.R.I.) under the 5th Call of "Science and Society" Action Always strive for excellence –
Theodoros Papazoglou" (Project Number: 28056); H.F.R.I call "Basic research Financing
(Horizontal support of all Sciences)" under the National Recovery and Resilience Plan "Greece
2.0" funded by the European Union – NextGenerationEU (Project Number: 014941); COFLEX
14941 HFRI P.N. 80147/17.1.24; NIH grant No 1R01MH124867 to P.P.; NIH grants
F31NS101820 and F32MH125600 to D.M.I.; NIH-NINDS (R35 NS127232) (FP), and an award

580 from the NOMIS Foundation (FP). A.L. is supported by National Institute of Mental Health
581 (NIMH) R01MH124047 and R01MH124867; National Institute on Aging (NIA) RF1AG080818;
582 National Institute of Neurological Disorders and Stroke (NINDS) Brain Initiative
583 U01NS115530; NINDS R01NS121106, NINDS R01NS131728, and NINDS Brain Initiative
584 R01NS133381

Author contributions

P.P., A.L., F.P., S.C., D.M.I., and S.T. conceived the study. D.M.I. performed the in-utero
electroporation, prepared the tissue, and acquired the confocal images. L.H. developed a
script enabling image deconvolution of the raw PN images. Y.K. and D.M.I. manually edited
semi-automated synaptic reconstructions and participated in data analysis.

S.T. refined the morphological traces, developed and validated the model, implemented the
simulation code, and analyzed the data. S.C. and S.T. generated the input patterns. S.T. wrote
the manuscript, with editing support from P.P., A.L., F.P., S.C., and D.M.I.

Declaration of interests

596 The authors declare no competing interests.

Declaration of generative AI and AI-assisted technologies in the writing process

While preparing this work, the authors used large language models (ChatGPT-4o) for spelling

and grammar checks. The author(s) reviewed and edited the content as needed and take full

responsibility for the final publication.

Materials and Methods

Mice

All animals were handled according to protocols approved by the Institutional Animal Care and

Use Committee at Columbia University, New York. Postnatal day 98 C57BL/6 mice (strain

code: 000664; Jackson Laboratory) were used for all experiments. Timed-pregnant female

mice were maintained in a 12-hour light/dark cycle and obtained by overnight breeding with

males of the same strain. For timed-pregnant mating, noon after mating is considered E0.5.

Constructs

The tdTomato reporter was previously generated by the Polleux lab[42]. Briefly, the tdTomato

insert was subcloned into the pAAV-Ef1a-DIO eNpHR 3.0-EYFP plasmid (Addgene 26966)

between the AscI and NheI cloning sites. The inhibitory synapse reporter EGFP-Gephyrin

(clone P1) was obtained from H. Cline (TSRI, La Jolla, USA) and subcloned into the pUBC-

DIO Teal-Gephyrin plasmid (Addgene 73918) between the AscI and NheI cloning sites.

In utero **electroporation**

In utero electroporation targeting the dorsal hippocampus was performed using a triple-

electrode setup as previously described in E15.5 mouse embryos[13,107,108]. Endotoxin-free DNA

was injected using a glass pipette into one ventricle of the mouse embryos. Electroporation

was performed at E15.5 using a square wave electroporator (ECM 830, BTX) with two anodes

(positively charged) placed laterally on either side of the head and one cathode (negatively

charged) rostrally at a 0° angle to the horizontal plane. The electroporation settings were: 5

pulses of 40V for 50 ms with 500 ms intervals. Plasmids were used at the following

concentrations: Flex-tdTomato reporter plasmid: 1 µg/µl; Flex-EGFP-Gephyrin 0.45 µg/µl;

NLS-Cre recombinase: 150 pg/µl.

Tissue preparation

631   Animals at the indicated age were anaesthetized with isoflurane before intracardiac perfusion

632   with PBS and 4% PFA (Electron Microscopy Sciences). As previously described[42], 130 µm

633   coronal brain sections were obtained using a vibrating microtome (Leica VT1200S). Sections

634   were mounted on slides and briefly dehydrated at room temperature to reduce section

635   thickness before being coverslipped in Fluoromount-G (SouthernBiotech).

Confocal imaging

Confocal imaging was performed as previously described[42]. Briefly, images of electroporated

neurons in slices were acquired with a Nikon AXR confocal microscope equipped with a 100x,

1.49 NA Apochromat oil-immersion objective and controlled by NIS-Elements (Nikon

Instruments Inc., Melville, NY). Image volumes were acquired using galvano scanning with a

1024x1024 scan region, corresponding to a lateral pixel size of 110 nm and a z-step size of

100 nm. To compensate for signal attenuation caused by tissue scattering and spherical

aberration with increasing depth, laser power was linearly increased as a function of depth to

normalize the mean fluorescent intensity of pixels from image planes throughout the stack[42].

Following acquisition, image stacks were deconvolved using the GPU-based Richardson-Lucy

deconvolution algorithm within NIS-Elements. Deconvolution was performed with a theoretical

point spread function based on imaging acquisition parameters and used 20 iterations.

Individual z-stacks were subsequently stitched together using Fiji's Grid/Collection stitching

tool[109] to generate a single three-dimensional volume representing the entire neuron. Dendritic

spines and inhibitory synapses were quantified based on tdTomato fluorescence and EGFP-

Gephyrin puncta fluorescence, respectively. All quantifications were performed in CA1

hippocampus in sections of comparable rostro-caudal position.

Neuron trace reconstruction

Neuron traces, digital reconstructions of the dendritic tree, are an effective representation of

neuronal topology and geometry. The traces described using a tree graph and consist of 3-D

point coordinates, diameters, and connectivity between points. This representation enables

an extensive quantitative analysis of the geometrical organization of the neurons they

represent including total length, branching angles, distribution statistics and cumulative

distance from the soma[110]. In this paper, the initial reconstructions are obtained using the

automatic tracing methods built into the open-source 3D visualization and analysis tool

Vaa3D[45]. Then, experts manually proofread the traces and make adjustments with the built-in

proof-editing tools.

Synapse detection

Excitatory and inhibitory synapses were annotated with the Spine Detector and IS Detector

toolkits within Vaa3D software as previously described[42]. Briefly, excitatory synapses are

identified as dendritic spines segmented by the Spine Detector toolkit using the image and

neuron trace as inputs[111,112]. Because the dendrite traces represent the dendrites with a series

of overlapping nodes[113], information about the volume and distance from the dendrite of each

spine can be associated with its nearest node to assign a location within the spatial context of

the dendritic arbor. Users have the ability to accept or reject potential spines and adjust their

volume.

We labeled inhibitory synapses using the scaffolding protein Gephyrin tagged with EGFP as

a marker[114]. Because these synapses can only occur on the dendrites or the spines of neurons

of interest, IS Detector uses the image from the cell-fill channel containing the dendrites and

the spines as a mask image to extract the relevant region for the inhibitory synaptic marker.

Then, signal beyond user-input parameters for minimum/maximum voxel count and distance

from the trace is excluded and potential inhibitory synapses from the resulting image are

identified based on a user-input intensity threshold. Users have the ability to accept or reject

potential inhibitory synapses, adjust their volume, and assign them as dendrite-targeting or

spine-targeting.

Dendritic diameter correction

To improve the accuracy of the morphological data and ensure the physiological plausibility of

the neuronal reconstructions, a custom smoothing algorithm was applied to correct artifactual

fluctuations in dendritic diameters (**Figure S1**). This algorithm utilized a weighted moving

average filter to smooth out large changes in diameter while preserving the overall structural

integrity of the neuron. The filter was designed to account for local variations in diameter by

applying larger weight to nearby points, reducing the influence of outliers and providing a

consistent tapering pattern along the dendrites.

Prior to smoothing algorithm, a minimum diameter threshold was applied to all diameters to

address the occurrence of bottlenecks. These bottlenecks could significantly impact

simulations by increasing the axial resistance and altering signal propagation. The minimum

diameter threshold was domain-specific, reflecting the physiological constraints of different

neuronal compartments.

Synaptic distribution analysis

Synaptic null distribution: within-domain and within-subdomain randomization

For each reconstructed morphology, synapses were randomly reassigned within branches of

the same domain or sub-domain, ensuring a uniform probability distribution. Specifically, we

first extracted the nodes of interest from the reconstructed arbor trace and then reassigned

the annotations describing spine features to new, randomly selected locations within the same

domain. This process preserved the overall structure of the arbor while randomizing synaptic

positions. We repeated this procedure 1,000 times to create the null distribution, conducting

the necessary analysis on each randomized dataset.

Structural cluster analysis

Putative structural clusters were identified using a hierarchical clustering algorithm applied

solely to large spines, i.e., those above the 80th percentile of the spine volume distribution.

For each branch, we selected the large spines and clustered them based on their relative

distance from the soma (i.e., path length). Specifically, we used the SciPy hierarchical

clustering linkage with *complete* method to calculate the Euclidian distance. A threshold of 10

micrometers was then used to define clusters, after which we collected their characteristics

(e.g., spine count, total volume). Since the clustering was performed separately for each

branch, any potential clusters spanning branching points were excluded from the analysis.

Computational Modeling

The CA1 PN models, implemented using the Python version of the NEURON simulation

environment[67], and were inspired by prior work by using the ionic mechanism and their

distribution throughout the dendrosomatic axis[26]. The ionic channels, written using NMODL

language, used for this project include: inward transient sodium current (I_{Na}), outward delayed

rectifier potassium current (I_{Kdr}), A-type potassium current (I_A), Sodium persistent current (I_{Nap}),

Hyperpolarization-activated current (I_h), Slowly activating muscarinic potassium current (I_m),

Calcium current from T-type channel (I_{CaT}), Calcium current from L-type channel (I_{CaL}), Calcium

current from R-type channel (I_{CaR}), Slow calcium-dependent potassium current (I_{sAHP}), Medium

calcium-dependent potassium current (I_{mAHP}). Minor modifications were implemented in the

kinetics and distribution of some channels. To facilitate the rapid and reproducible generation

of models, we developed a Python Class capable of processing neuronal morphologies in

.swc file format and automatically incorporating the appropriate biophysical properties and

channel distributions.

To ensure high spatial resolution in the simulations, we determined the number of segments

(n_{seg}) for each neuronal section using the lambda rule, which is based on the Alternating

Current (AC) length constant (λ). A ten-fold higher frequency was used in the calculation to

obtain a finer spatial discretization of each section. Specifically, we employed the following

equation (Eq. 1):

735
$$n_{seg} = 2 \cdot \left\lfloor \frac{\frac{L}{d_\lambda \cdot \lambda_f} + 0.999}{2} \right\rfloor + 1 \ (1)$$

where L is the length of the section, d_λ the fraction of the AC length constant (typically 0.1),

$\lambda_f f$ the AC length constant at the chosen frequency f, and $\lfloor \cdot \rfloor$ denotes a function that converts

a specified value into an integer number. This approach ensured that all segments were small

enough to disentangle inputs coming from most of nearby synapse couples, providing

accurate spatial and biophysical representation of the model.

Modeling Synapses

We incorporated both glutamatergic (excitatory) and GABAergic (inhibitory) synapses. All

dendritic spines identified in the reconstructed neurons were assumed to be excitatory

synapses containing both AMPA and NMDA receptors, while inhibitory synapses were

modeled as GABA$_A$-mediated inputs. Synaptic strength was adjusted based on the synapse

size measured from the original reconstruction and, for a subset of excitatory synapses, by

the distance from the soma. Specifically, for excitatory synapses, validated conductance

values for both NMDAR and AMPAR components were first calculated. The AMPA

conductance was then scaled by a volume-dependent scaling factor (Eq. 2). Both AMPA and

NMDA components for synapses in Basal, Trunk and Oblique compartments were further

adjusted by a distance-dependent scaling factor[71,73]. In contrast, inhibitory synapses were

scaled solely by the volume-dependent factor (Eq. 3).

754
$$AMPA\ Volume\ scaling = 0.6 \cdot \frac{Spine\ volume_i}{Average\ spine\ volume} + 0.4\ (2)$$

755
$$GABA\ Volume\ scaling = \frac{PSD\ volume_i}{Average\ PSD\ volume}\ (3)$$

AMPA & GABA$_A$ definition

758 AMPA and GABA$_A$ synapses were simulated with a two-state kinetic model, defined by rise

759 and decay times. The synaptic current is calculated according to the following equations (Eqs.

760 4, 5):

761
$$I_{syn} = g_{syn}(t) \cdot (V_m - E_{syn}), syn\ \in \{\text{AMPA}, \text{GABA}_A\}\ (4)$$

762
763
764
$$g_{syn}(t) = f_{norm} \cdot \bar{g}_{peak} \cdot \left(\exp\left(-\frac{t}{\tau_d}\right) - \exp\left(-\frac{t}{\tau_r}\right)\right)\ (5)$$

765 where I_{syn} is the synaptic current, V_m the membrane potential, g_{syn} the receptor conductance,

766 g_{peak} the peak conductance and E_{syn} its reverse potential. τ_r and τ_d represent the rise and

767   decay time constants, respectively. The factor f_{norm} normalizes the difference of exponentials

768   so that the peak of g_{syn} equals $\bar{g}_{peak}$ and its calculated using the formula below (Eq. 6).

$$f_{norm} = \frac{1}{g_{syn}(t_{peak})}, t_{peak} = \frac{\tau_d \tau_r}{\tau_d - \tau_r} \ln \frac{\tau_d}{\tau_r} \ (6)$$

772   ***NMDA definition***

773   The NMDA component of the excitatory synapses was also simulated as a two-state kinetic

774   model but with an additional factor that accounts for the voltage dependent removal of the

775   Magnesium blockage. The mechanism is described by the following equations (Eqs. 7, 8):

$$I_{NMDA} = g_{NMDA}(t) \cdot h(V_m) \cdot (V_m - E_{NMDA}) \ (7)$$

$$h(V_m) = \frac{1}{1 + \eta [Mg^{2+}]_o \exp\left(-\gamma (V_m - V_{half})\right)} \ (8)$$

where $h(\cdot)$ is a sigmoid function, η denotes the Magnesium sensitivity of unblock, $[Mg^{2+}]_o$ the

Magnesium extracellular concentration, γ the slope factor that determines how steep the

transition is around the midpoint, and V_{half} the membrane potential of the midpoint of the

sigmoid. $g_{NMDA}(t)$ is calculated with the same equations as the other synaptic currents (Eqs.

5, 6).

Pyramidal neuron model validation

The parameters that defined the biophysical properties and channel distributions were

carefully validated to match the cell's behavior with experimental data. We ensured that using

the same values for all morphologies would still adhere to the experimental values for each

cell, while also introducing variability expected in a heterogeneous population of pyramidal

neurons. We used hyperpolarizing step current sweeps to validate passive properties as the

input resistance (R_{in}), membrane time constant (τ_m), sag ratio and the dendritic attenuation,

following the protocols illustrated in **Magee (1998)**[68]**, Golding et al. (2005)**[69] for the voltage

attenuation and **Masurkar et al. (2020)**[8] for R_{in}, τ_m and sag ratio. We then used a depolarizing

step current sweeps to validate the frequency-input (F-I) curve and the adaptation from

Masurkar et al. (2020)[8]. For the synaptic strength validation, we followed the protocol used in

Magee and Cook, 2000[71], stimulating average size synapses at increasing distance from the

soma and maintaining the 0.2 mV somatic depolarization observed experimentally.

Inputs design

Grid-like and Place-like inputs

In accordance with the formulation described in **Shuman et al. (2020)**[115], entorhinal cortex

layer III (ECIII) grid-like inputs were modeled as the summation of three sinusoidal wave

components, following the approach of **Solstad et al. (2006)**[116]. In contrast, place-like inputs

from CA3 were generated by integrating these grid-like inputs. Specifically, grid-cell activity

was represented as the sum of three two-dimensional sinusoidal gratings, each defined by a

distinct wave vector. The resulting grid-like function was computed as (Eq. 9):

807
$$g_w(x,y) = \frac{2}{3}\left(\frac{1}{3}\sum_{i=1}^{3}\cos\left(k_i\left(r - r_0\right)\right) + \frac{1}{2}\right) \quad (9)$$

808 where the parameter set $w = (\lambda, \theta, r_0)$ defines the grid spacing λ, orientation θ, and the spatial

809 location of a theoretical place cell $r_0 = [x_0, y_0]^{\mathrm{T}}$, while the variable $r = [x,y]^{\mathrm{T}}$ represents the

810 current position, and k denotes the wave vectors arranged at angular intervals of $\frac{\pi}{3}$ and $\frac{2\pi}{3}$,

811 respectively (Eq. 10).

812
$$k_i = \frac{1}{\lambda}\left[\cos\left(\theta + (i-1)\frac{\pi}{3}\right), \sin\left(\theta + (i-1)\frac{\pi}{3}\right)\right]^{\mathrm{T}} \quad (10)$$

813 These grid-like functions were used to compute the probability of spike generation as a

814 function of the animal's position and the theoretical place fields. To incorporate theta rhythmic

815 modulation, a sinusoidal filter was applied to these probabilities (Eq 11):

816
$$p(t) = \mathrm{g}_w(x,y)\frac{\sin\left(2\pi \cdot f_\theta \cdot \left(\frac{t}{1000}\right) + \varphi_{theta}\right) + 1}{2} \quad (11)$$

where $f_\theta = 8\mathrm{Hz}$ and $\phi_\theta = 0^o$ correspond to the theta-cycle frequency and phase,

respectively, and t is expressed in milliseconds. A spike was generated for a given grid cell

whenever $p(t)$ exceeded a random value sampled from a normal distribution with mean

$\mu = 0.7$ and standard deviation $\sigma = 0.05$ or a random value from a uniform distribution used

to introduce baseline noise. This process was repeated across all grid-like cells to

construct the entire set of theoretical place fields.

To model CA3 place-like inputs, we combined the generated grid-like fields from EC inputs,

ensuring proper gamma filtering and theta-phase locking. Similarly to EC inputs, a spike was

generated when the probability $p(t)$ was greater than a normally distributed threshold value

or a noise baseline. Additionally, a 17 ms delay was introduced to account for the natural

temporal delay between EC and CA3 input streams. Finally, each spike train was filtered to a

function that simulate the effect of synaptic failure based on published data[117].

Noisy inputs

To account for non-place-tuned inputs, we generated a set of noisy inputs modeled as

Poisson-distributed spike trains with low baseline firing rates. To introduce temporal

fluctuations, these spike trains were modulated by applying a Gaussian filter at randomly

selected time points, transiently increasing the firing rate. In our approach, the probability for

a spike to occur was (Eq. 12):

836
$$p(t) = O + H \cdot \exp\left(\frac{\left(t - t_{peak}\right)^2}{2\sigma^2}\right) (12)$$

where O represents the baseline offset of the gaussian curve, H is the peak amplitude, t is the

time at which the spike occurred and t_{peak} is a randomly chosen value corresponding to the

mean value of the Gaussian filter. The standard deviation σ is also chosen randomly in a fixed

window of values (400 ms, 720 ms) and represents the width of the window with slightly higher

firing frequency.

This process ensured that noisy inputs exhibited sporadic increases of activity while

maintaining an overall unstructured firing pattern, effectively capturing the background

synaptic activity present in the hippocampal CA1 place cells. Similarly to place tuned inputs,

the noisy spike trains were further processed to account for synaptic failure.

Inhibitory inputs

Inhibitory inputs were modeled as theta-modulated, gamma-filtered Poisson-distributed spike

trains, with firing properties tailored to different interneuron subtypes as described in **Bezaire**

et al., 2016[76]. To generate these spike trains, we set baseline firing rates according to known

physiological values for each interneuron type. These spike trains were then filtered using a

theta-modulated function, ensuring that their activity exhibited phase locking to the theta cycle

at subtype-specific phases (Eq. 11).

Additionally, a gamma-frequency filtering step was applied to refine the temporal structure of

the remaining spikes. The level of randomness was controlled by introducing a noise

parameter, which varied across interneuron types to reflect their intrinsic firing variability.

Delays of 300 ms were incorporated to approximate transmission latencies. Specifically, inputs

were tuned to replicate the activity of different interneurons neurogliaform (NGF) cells, oriens-

lacunosum moleculare (OLM) cells, parvalbumin-expressing basket (PVbasket) cells, apical

dendrite targeting cholecystokinin-expressing (CCK) cells, Bistratified cells, and Ivy cells with

peak firing rates ranging from 2.8 Hz to 30 Hz, distinct theta-phase preferences, and varying

degrees of noise, capturing the diversity of inhibitory influences in the hippocampal network.

Input distribution

Synaptic inputs were distributed in an anatomically accurate manner. For inhibitory synapses,

their connection probabilities to each interneuron subtype were based on estimates from

Bezaire and Soltesz (2013)[118], with slight upward adjustments to account for missing

interneuron types. While NGF and OLM inputs were uniformly distributed across tuft synapses,

the distribution of other interneurons followed a nonlinear function of the distance from the

soma. This distribution pattern was informed by anatomical observations while maintaining

symmetry between the apical and basal dendritic domains. Specifically, PV basket synapses

were concentrated near the soma, while CCK and Ivy cells targeted more distal synapses in

the apical and basal domains, respectively. Bistratified cell synapses were predominantly

located in the middle regions of both domains (**Figure S2**).

ECIII grid-like inputs were directed to synapses on the tuft dendrites in the stratum lacunosum-

moleculare, whereas CA3 place-like inputs primarily targeted synapses in the trunk, oblique,

and basal dendritic domains, spanning the stratum oriens and stratum radiatum. At the branch

level, place-tuned inputs were assigned to both clustered and dispersed synapses. Using

GluSnFR-based experimental data from **Adoff et al. (2021)**[39] as a reference, we determined

the probability of each synapse to receive place-tuned inputs, guided by the observed ratios

of clustered vs. dispersed synapses and the relative distributions in the apical vs. basal

domains.

The total number of activated clustered synapses, tuned to the somatic place field, was

randomly determined using a transformation of a Gaussian-distributed variable (Eq. 13):

$$n_{in} = r \cdot \frac{n_{cs}}{n_{pf}}, r = 0.9 + \frac{z^{1.5}}{3} \ (13)$$

Where n_{in} represents the number of clustered in-field tuned synapses, n_{cs} and n_{pf} represent

the number of clustered synapses and the number of place fields, respectively. z is a normally

distributed random variable with mean 0 and standard deviation 1, so that $z \sim \mathcal{N}(0,1)$, keeping

only the positive values. The range r is limited to values lower than 1.6.

The Gaussian shape was chosen to replicate the distribution of place field properties observed

at the population level by **Mizuseki et al. (2012)**[84]. Clustered synapses assigned to the

remaining 11 place fields were selected randomly with a fixed probability of (Eq. 14):

$$n_{out} = 0.4 \cdot \frac{n_{cs}}{n_{pf}} \ (14)$$

894 Where n_{out} is the number of clustered out-field tuned synapses. Dispersed synapses were

895 also randomly selected with fixed probabilities, determining whether they would receive place-

896 tuned or noisy inputs. To account for biological variability, we generated a set of 1,000 distinct

897 connectivity patterns for each of the five morphologies, ensuring a realistic representation of

898 network heterogeneity.

Inputs Shuffling

To investigate the role of synaptic clustering in shaping neuronal responses, we implemented

three distinct synaptic input shuffling protocols that redistributed the incoming place-tuned

inputs while preserving specific structural or functional properties of the original configuration.

In the Shuffled – Same Number (SSN) condition, for each dendritic domain, we first computed

the number of synapses activated in the control (i.e., clustered) configuration. The same input

patterns were then reassigned to an equal number of randomly selected synapses within the

same domain, thereby preserving synapse count but disrupting spatial clustering.

In the Shuffled – Same Volume (SSV) protocol, we calculated the total volume of all synapses

activated per place field in the control configuration. Synaptic locations were then randomly

selected until their cumulative volume matched that total volume. As this procedure typically

resulted in a greater number of activated synapses, due to the inclusion of smaller synapses,

the set of input spike trains used was expanded by sampling from the pool of inputs applied

in the original, clustered configuration.

The Shuffled – Same Weight (SSW) protocol further refined this approach by summing the

total synaptic weight from the control configuration, accounting for both the synaptic volume

and the synaptic democracy principle (i.e., equalization of synaptic impact regardless of

location). Synapses were then randomly selected until their combined weights matched the

original total. Although this method also led to an increased number of activated synapses

compared to the control, the number was lower than that observed in the SSV condition.

Place cell quantification

To quantify the spatial tuning properties of the simulated neurons, we computed the firing rate

maps for each cell across the linear track, and used them to calculate the various place field

features. The Field Size identifies the spatial extent of contiguous bins around the peak firing

location where the firing rate remains above a given relative threshold. Once the Place Field

width was characterized, we determined the mean firing rate within and outside the identified

field region, respectively named In-field and Out-field Firing Frequency, respectively. In

addition, spatial selectivity was assessed using a modified version of the spatial information

index[83]. This metric calculates the information content of firing in bits per second by comparing

the firing rate in each spatial bin to the mean firing rate across the track, weighted by

occupancy probability, i.e., time spent in each spatial bin. The time spent in each bin was used

to normalize firing rates, and the spatial information was computed (Eq. 15) as:

933
$$Spatial\ Info = \sum_{i=1}^{N_{bins}} p_i \cdot \frac{\lambda_i}{\lambda} \cdot \log_2 \left(\frac{\lambda_i}{\lambda}\right) \ (15)$$

934   Where p_i is the occupancy probability of bin i, λ_i is the firing rate in that bin and λ is the mean

935   firing rate across all bins, N_{bins}.

Multivariate Random Forest

To assess whether the structural features of synaptic clusters could predict the functional

output of the simulated neurons, we employed a supervised machine learning approach using

a Random Forest regression model. Specifically, we trained a multi-output regression model

to predict place field properties (i.e., Spatial Information, In-field and Out-of-field Firing

Frequency, and Field Size) from cluster-derived input features (i.e., total in-field cluster

volume, average in-field cluster size, and the proportion of dendritic branches containing in-

field clusters). These features were normalized to [0,1] range using a Min-Max Scaler to

ensure comparability across dimensions. The target output matrix comprised the

aforementioned spatial metrics, which were likewise scaled prior to training. For the analysis

we used simulations conducted under the Control allocation protocol, excluding those that

exhibited overflow or contained missing values.

We used a Multi-Output Regressor wrapping a Random Forest Regressor (100 estimators per

model). Models were trained and evaluated over 10 independent runs using an 80/20 train-

test split, with mean squared error used to assess performance across both individual outputs

and overall predictions. Control protocols were validated against two baselines: one involving

permutation of the input features (Random training inputs, not shown) and another involving

selective removal of key input features (Removal input feature, not shown) to confirm the

robustness and informativeness of the cluster descriptors. Feature importance values were

computed as the average importance across all estimators within the ensemble.

SHAP-Based Feature Attribution Analysis

To investigate the contribution of synaptic cluster features to the spatial coding properties of

simulated neurons, we employed SHAP (*SHapley Additive exPlanations*), a game-theory

approach for interpreting machine learning models[88]. We computed SHAP values using the

TreeExplainer provided by the SHAP Python library, which is optimized for tree-based models.

SHAP values were calculated on the training set to ensure consistency with the model fitting.

The *summary_plot()* function was used to visualize both the magnitude and direction of each

feature's effect. Feature values were color-coded to indicate their relative magnitude (red for

high values, blue for low values), enabling intuitive interpretation of whether increasing a

specific feature tended to increase or decrease a given output.

To complement the qualitative visual inspection, we also computed the Pearson correlation

between feature values and their corresponding SHAP values. This provided a numerical

estimate of the directionality and strength of influence each feature exerted on the model

output.

Simulation environment and statistical analysis

All simulations were performed on our High-Performance Computing Cluster (Rocks 7.0) with

624 cores and 3.328 TB of shared RAM under CentOS 7.4 OS[119], through the NEURON

simulator[120] using Python via an Anaconda virtual environment. Neuronal models have been

simulated at a sampling rate of 40 kHz (dt = 0.025 ms). For all standard statistical tests

(detailed in all relevant figure legends), the α was chosen to be 0.05 for statistical significance.

To correct for multiple comparisons, the α was divided by the number of tests according to the

Bonferroni's procedure. Throughout the figures, p values are denoted by * ($p < 0.05$), ** ($p <$

0.01), and *** ($p < 0.001$).

References

1.   O'Keefe, J., and Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary
evidence from unit activity in the freely-moving rat. Brain Res *34*, 171–175.
https://doi.org/10.1016/0006-8993(71)90358-1.

2.   Muller, R.U., and Kubie, J.L. (1987). The effects of changes in the environment on the
spatial firing of hippocampal complex-spike cells. J Neurosci *7*, 1951–1968.
https://doi.org/10.1523/JNEUROSCI.07-07-01951.1987.

3.   Quirk, G.J., Muller, R.U., and Kubie, J.L. (1990). The firing of hippocampal place cells
in the dark depends on the rat's recent experience. J Neurosci *10*, 2008–2017.
https://doi.org/10.1523/JNEUROSCI.10-06-02008.1990.

4.   Bostock, E., Muller, R.U., and Kubie, J.L. (1991). Experience-dependent modifications
of hippocampal place cell firing. Hippocampus *1*, 193–205.
https://doi.org/10.1002/hipo.450010207.

5.   Leutgeb, S., Leutgeb, J.K., Barnes, C.A., Moser, E.I., McNaughton, B.L., and Moser,
M.-B. (2005). Independent codes for spatial and episodic memory in hippocampal
neuronal ensembles. Science *309*, 619–623. https://doi.org/10.1126/science.1114037.

6.   Turi, G.F., Li, W.-K., Chavlis, S., Pandi, I., O'Hare, J., Priestley, J.B., Grosmark, A.D.,
Liao, Z., Ladow, M., Zhang, J.F., et al. (2019). Vasoactive Intestinal Polypeptide-
Expressing Interneurons in the Hippocampus Support Goal-Oriented Spatial Learning.
Neuron *101*, 1150-1165.e8. https://doi.org/10.1016/j.neuron.2019.01.009.

7.   Zutshi, I., Valero, M., Fernández-Ruiz, A., and Buzsáki, G. (2022). Extrinsic control and
intrinsic computation in the hippocampal CA1 circuit. Neuron *110*, 658-673.e5.
https://doi.org/10.1016/j.neuron.2021.11.015.

8.   Masurkar, A.V., Tian, C., Warren, R., Reyes, I., Lowes, D.C., Brann, D.H., and
Siegelbaum, S.A. (2020). Postsynaptic integrative properties of dorsal CA1 pyramidal
neuron subpopulations. J Neurophysiol *123*, 980–992.
https://doi.org/10.1152/jn.00397.2019.

9.   Megías, M., Emri, Z., Freund, T.F., and Gulyás, A.I. (2001). Total number and
distribution of inhibitory and excitatory synapses on hippocampal CA1 pyramidal cells.
Neuroscience *102*, 527–540. https://doi.org/10.1016/s0306-4522(00)00496-6.

10.   Spruston, N. (2008). Pyramidal neurons: dendritic structure and synaptic integration.
Nat Rev Neurosci *9*, 206–221. https://doi.org/10.1038/nrn2286.

11.   Dougherty, K.A., Islam, T., and Johnston, D. (2012). Intrinsic excitability of CA1
pyramidal neurones from the rat dorsal and ventral hippocampus. J Physiol *590*, 5707–
5722. https://doi.org/10.1113/jphysiol.2012.242693.

12.   Sheffield, M.E.J., Adoff, M.D., and Dombeck, D.A. (2017). Increased Prevalence of
Calcium Transients across the Dendritic Arbor during Place Field Formation. Neuron
96, 490-504.e5. https://doi.org/10.1016/j.neuron.2017.09.029.

13.   Blockus, H., Rolotti, S.V., Szoboszlay, M., Peze-Heidsieck, E., Ming, T., Schroeder, A.,
Apostolo, N., Vennekens, K.M., Katsamba, P.S., Bahna, F., et al. (2021). Synaptogenic

activity of the axon guidance molecule Robo2 underlies hippocampal circuit function.
Cell Rep *37*, 109828. https://doi.org/10.1016/j.celrep.2021.109828.

14. O'Hare, J.K., Gonzalez, K.C., Herrlinger, S.A., Hirabayashi, Y., Hewitt, V.L., Blockus,
H., Szoboszlay, M., Rolotti, S.V., Geiller, T.C., Negrean, A., et al. (2022).
Compartment-specific tuning of dendritic feature selectivity by intracellular Ca2+
release. Science *375*, eabm1670. https://doi.org/10.1126/science.abm1670.

15. Amaral, D.G., and Witter, M.P. (1989). The three-dimensional organization of the
hippocampal formation: a review of anatomical data. Neuroscience *31*, 571–591.
https://doi.org/10.1016/0306-4522(89)90424-7.

16. Suh, J., Rivest, A.J., Nakashiba, T., Tominaga, T., and Tonegawa, S. (2011).
Entorhinal cortex layer III input to the hippocampus is crucial for temporal association
memory. Science *334*, 1415–1420. https://doi.org/10.1126/science.1210125.

17. Larkum, M. (2013). A cellular mechanism for cortical associations: an organizing
principle for the cerebral cortex. Trends Neurosci *36*, 141–151.
https://doi.org/10.1016/j.tins.2012.11.006.

18. Xu, N., Harnett, M.T., Williams, S.R., Huber, D., O'Connor, D.H., Svoboda, K., and
Magee, J.C. (2012). Nonlinear dendritic integration of sensory and motor input during
an active sensing task. Nature *492*, 247–251. https://doi.org/10.1038/nature11601.

19. Takahashi, H., and Magee, J.C. (2009). Pathway interactions and synaptic plasticity in
the dendritic tuft regions of CA1 pyramidal neurons. Neuron *62*, 102–111.
https://doi.org/10.1016/j.neuron.2009.03.007.

20. Magee, J.C. (2000). Dendritic integration of excitatory synaptic input. Nat Rev Neurosci
1, 181–190. https://doi.org/10.1038/35044552.

21. Tzilivaki, A., Kastellakis, G., Schmitz, D., and Poirazi, P. (2022). GABAergic
Interneurons with Nonlinear Dendrites: From Neuronal Computations to Memory
Engrams. Neuroscience *489*, 34–43.
https://doi.org/10.1016/j.neuroscience.2021.11.033.

22. Katz, Y., Menon, V., Nicholson, D.A., Geinisman, Y., Kath, W.L., and Spruston, N.
(2009). Synapse distribution suggests a two-stage model of dendritic integration in CA1
pyramidal neurons. Neuron *63*, 171–177. https://doi.org/10.1016/j.neuron.2009.06.023.

23. Branco, T., and Häusser, M. (2010). The single dendritic branch as a fundamental
functional unit in the nervous system. Curr Opin Neurobiol *20*, 494–502.
https://doi.org/10.1016/j.conb.2010.07.009.

24. Losonczy, A., and Magee, J.C. (2006). Integrative properties of radial oblique dendrites
in hippocampal CA1 pyramidal neurons. Neuron *50*, 291–307.
https://doi.org/10.1016/j.neuron.2006.03.016.

25. Gasparini, S., and Magee, J.C. (2006). State-dependent dendritic computation in
hippocampal CA1 pyramidal neurons. J Neurosci *26*, 2088–2100.
https://doi.org/10.1523/JNEUROSCI.4428-05.2006.

26. Poirazi, P., Brannon, T., and Mel, B.W. (2003). Arithmetic of subthreshold synaptic
summation in a model CA1 pyramidal cell. Neuron *37*, 977–987.
https://doi.org/10.1016/s0896-6273(03)00148-x.

27. Polsky, A., Mel, B.W., and Schiller, J. (2004). Computational subunits in thin dendrites
of pyramidal cells. Nat Neurosci *7*, 621–627. https://doi.org/10.1038/nn1253.

28. Cichon, J., and Gan, W.-B. (2015). Branch-specific dendritic Ca(2+) spikes cause
persistent synaptic plasticity. Nature *520*, 180–185.
https://doi.org/10.1038/nature14251.

29. Tzilivaki, A., Kastellakis, G., and Poirazi, P. (2019). Challenging the point neuron
dogma: FS basket cells as 2-stage nonlinear integrators. Nat Commun *10*, 3664.
https://doi.org/10.1038/s41467-019-11537-7.

30. Mel, B. (1991). The Clusteron: Toward a Simple Abstraction for a Complex Neuron. In
Advances in Neural Information Processing Systems (Morgan-Kaufmann).

31. Mel, B.W., Ruderman, D.L., and Archie, K.A. (1998). Translation-invariant orientation
tuning in visual "complex" cells could derive from intradendritic computations. J
Neurosci *18*, 4325–4334. https://doi.org/10.1523/JNEUROSCI.18-11-04325.1998.

32. Poirazi, P., and Mel, B.W. (2001). Impact of active dendrites and structural plasticity on
the memory capacity of neural tissue. Neuron *29*, 779–796.
https://doi.org/10.1016/s0896-6273(01)00252-5.

33. Fu, M., Yu, X., Lu, J., and Zuo, Y. (2012). Repetitive motor learning induces
coordinated formation of clustered dendritic spines in vivo. Nature *483*, 92–95.
https://doi.org/10.1038/nature10844.

34. Oh, W.C., Parajuli, L.K., and Zito, K. (2015). Heterosynaptic structural plasticity on local
dendritic segments of hippocampal CA1 neurons. Cell Rep *10*, 162–169.
https://doi.org/10.1016/j.celrep.2014.12.016.

35. Frank, A.C., Huang, S., Zhou, M., Gdalyahu, A., Kastellakis, G., Silva, T.K., Lu, E.,
Wen, X., Poirazi, P., Trachtenberg, J.T., et al. (2018). Hotspots of dendritic spine
turnover facilitate clustered spine addition and learning and memory. Nat Commun *9*,
422. https://doi.org/10.1038/s41467-017-02751-2.

36. Takahashi, N., Kitamura, K., Matsuo, N., Mayford, M., Kano, M., Matsuki, N., and
Ikegaya, Y. (2012). Locally synchronized synaptic inputs. Science *335*, 353–356.
https://doi.org/10.1126/science.1210362.

37. Kastellakis, G., Silva, A.J., and Poirazi, P. (2016). Linking Memories across Time via
Neuronal and Dendritic Overlaps in Model Neurons with Active Dendrites. Cell Rep *17*,
1491–1504. https://doi.org/10.1016/j.celrep.2016.10.015.

38. Sehgal, M., Filho, D.A., Kastellakis, G., Kim, S., Lee, J., Shen, Y., Huang, S., Lavi, A.,
Fernandes, G., Davila Mejia, I., et al. (2025). Compartmentalized dendritic plasticity in
the mouse retrosplenial cortex links contextual memories formed close in time. Nat
Neurosci *28*, 602–615. https://doi.org/10.1038/s41593-025-01876-8.

39. Adoff, M.D., Climer, J.R., Davoudi, H., Marvin, J.S., Looger, L.L., and Dombeck, D.A.
(2021). The functional organization of excitatory synaptic input to place cells. Nat
Commun *12*, 3558. https://doi.org/10.1038/s41467-021-23829-y.

40. Hwang, F.-J., Roth, R.H., Wu, Y.-W., Sun, Y., Kwon, D.K., Liu, Y., and Ding, J.B.
(2022). Motor learning selectively strengthens cortical and striatal synapses of motor

engram neurons. Neuron *110*, 2790-2801.e5.
https://doi.org/10.1016/j.neuron.2022.06.006.

41. Kastellakis, G., Cai, D.J., Mednick, S.C., Silva, A.J., and Poirazi, P. (2015). Synaptic
clustering within dendrites: an emerging theory of memory formation. Prog Neurobiol
126, 19–35. https://doi.org/10.1016/j.pneurobio.2014.12.002.

42. Iascone, D.M., Li, Y., Sümbül, U., Doron, M., Chen, H., Andreu, V., Goudy, F., Blockus,
H., Abbott, L.F., Segev, I., et al. (2020). Whole-Neuron Synaptic Mapping Reveals
Spatially Precise Excitatory/Inhibitory Balance Limiting Dendritic and Somatic Spiking.
Neuron *106*, 566-578.e8. https://doi.org/10.1016/j.neuron.2020.02.015.

43. Atasoy, D., Aponte, Y., Su, H.H., and Sternson, S.M. (2008). A FLEX switch targets
Channelrhodopsin-2 to multiple cell types for imaging and long-range circuit mapping. J
Neurosci *28*, 7025–7030. https://doi.org/10.1523/JNEUROSCI.1954-08.2008.

44. Schnütgen, F., Doerflinger, N., Calléja, C., Wendling, O., Chambon, P., and
Ghyselinck, N.B. (2003). A directional strategy for monitoring Cre-mediated
recombination at the cellular level in the mouse. Nat Biotechnol *21*, 562–565.
https://doi.org/10.1038/nbt811.

45. Peng, H., Bria, A., Zhou, Z., Iannello, G., and Long, F. (2014). Extensible visualization
and analysis for multidimensional images using Vaa3D. Nat Protoc *9*, 193–208.
https://doi.org/10.1038/nprot.2014.011.

46. Perez-Cruz, C., Nolte, M.W., van Gaalen, M.M., Rustay, N.R., Termont, A., Tanghe, A.,
Kirchhoff, F., and Ebert, U. (2011). Reduced spine density in specific regions of CA1
pyramidal neurons in two transgenic mouse models of Alzheimer's disease. J Neurosci
31, 3926–3934. https://doi.org/10.1523/JNEUROSCI.6142-10.2011.

47. Horton, S., Mastrolia, V., Jackson, R.E., Kemlo, S., Pereira Machado, P.M., Carbajal,
M.A., Hindges, R., Fleck, R.A., Aguiar, P., Neves, G., et al. (2024). Excitatory and
inhibitory synapses show a tight subcellular correlation that weakens over
development. Cell Rep *43*, 114361. https://doi.org/10.1016/j.celrep.2024.114361.

48. Bannister, N.J., and Larkman, A.U. (1995). Dendritic morphology of CA1 pyramidal
neurones from the rat hippocampus: I. Branching patterns. J Comp Neurol *360*, 150–
160. https://doi.org/10.1002/cne.903600111.

49. Cavalieri, D., Angelova, A., Islah, A., Lopez, C., Bocchio, M., Bollmann, Y., Baude, A.,
and Cossart, R. (2021). CA1 pyramidal cell diversity is rooted in the time of
neurogenesis. Elife *10*, e69270. https://doi.org/10.7554/eLife.69270.

50. Lamprecht, R., and LeDoux, J. (2004). Structural plasticity and memory. Nat Rev
Neurosci *5*, 45–54. https://doi.org/10.1038/nrn1301.

51. Pennacchietti, F., Vascon, S., Nieus, T., Rosillo, C., Das, S., Tyagarajan, S.K., Diaspro,
A., Del Bue, A., Petrini, E.M., Barberis, A., et al. (2017). Nanoscale Molecular
Reorganization of the Inhibitory Postsynaptic Density Is a Determinant of GABAergic
Synaptic Potentiation. J Neurosci *37*, 1747–1756.
https://doi.org/10.1523/JNEUROSCI.0514-16.2016.

52. Losonczy, A., Makara, J.K., and Magee, J.C. (2008). Compartmentalized dendritic
plasticity and input feature storage in neurons. Nature *452*, 436–441.
https://doi.org/10.1038/nature06725.

53. Lai, C.S.W., Adler, A., and Gan, W.-B. (2018). Fear extinction reverses dendritic spine
formation induced by fear conditioning in the mouse auditory cortex. Proc Natl Acad Sci
U S A *115*, 9306–9311. https://doi.org/10.1073/pnas.1801504115.

54. Leighton, A.H., Cheyne, J.E., and Lohmann, C. (2024). Clustered synapses develop in
distinct dendritic domains in visual cortex before eye opening. Elife *12*, RP93498.
https://doi.org/10.7554/eLife.93498.

55. Hedrick, N.G., Wright, W.J., and Komiyama, T. (2024). Local and global predictors of
synapse elimination during motor learning. Sci Adv *10*, eadk0540.
https://doi.org/10.1126/sciadv.adk0540.

56. Wright, W.J., Hedrick, N.G., and Komiyama, T. (2025). Distinct synaptic plasticity rules
operate across dendritic compartments in vivo during learning. Science *388*, 322–328.
https://doi.org/10.1126/science.ads4706.

57. Nusser, Z., Lujan, R., Laube, G., Roberts, J.D., Molnar, E., and Somogyi, P. (1998).
Cell type and pathway dependence of synaptic AMPA receptor number and variability
in the hippocampus. Neuron *21*, 545–559. https://doi.org/10.1016/s0896-
6273(00)80565-6.

58. Yuste, R., and Bonhoeffer, T. (2001). Morphological changes in dendritic spines
associated with long-term synaptic plasticity. Annu Rev Neurosci *24*, 1071–1089.
https://doi.org/10.1146/annurev.neuro.24.1.1071.

59. Bourne, J.N., and Harris, K.M. (2008). Balancing structure and function at hippocampal
dendritic spines. Annu Rev Neurosci *31*, 47–67.
https://doi.org/10.1146/annurev.neuro.31.060407.125646.

60. Harvey, C.D., and Svoboda, K. (2007). Locally dynamic synaptic learning rules in
pyramidal neuron dendrites. Nature *450*, 1195–1200.
https://doi.org/10.1038/nature06416.

61. Larkum, M.E. (2022). Are Dendrites Conceptually Useful? Neuroscience *489*, 4–14.
https://doi.org/10.1016/j.neuroscience.2022.03.008.

62. Yadav, A., Gao, Y.Z., Rodriguez, A., Dickstein, D.L., Wearne, S.L., Luebke, J.I., Hof,
P.R., and Weaver, C.M. (2012). Morphologic evidence for spatially clustered spines in
apical dendrites of monkey neocortical pyramidal cells. J Comp Neurol *520*, 2888–
2902. https://doi.org/10.1002/cne.23070.

63. Engert, F., and Bonhoeffer, T. (1997). Synapse specificity of long-term potentiation
breaks down at short distances. Nature *388*, 279–284. https://doi.org/10.1038/40870.

64. Schuman, E.M., and Madison, D.V. (1994). Locally distributed synaptic potentiation in
the hippocampus. Science *263*, 532–536. https://doi.org/10.1126/science.8290963.

65. Kastellakis, G., Tasciotti, S., Pandi, I., and Poirazi, P. (2023). The dendritic engram.
Front Behav Neurosci *17*, 1212139. https://doi.org/10.3389/fnbeh.2023.1212139.

66. Fu, M., Yu, X., Lu, J., and Zuo, Y. (2012). Repetitive motor learning induces
coordinated formation of clustered dendritic spines in vivo. Nature *483*, 92–95.
https://doi.org/10.1038/nature10844.

67.   Hines, M.L., and Carnevale, N.T. (1997). The NEURON simulation environment. Neural
1189   Comput *9*, 1179–1209. https://doi.org/10.1162/neco.1997.9.6.1179.

1190   68.   Magee, J.C. (1998). Dendritic hyperpolarization-activated currents modify the
1191   integrative properties of hippocampal CA1 pyramidal neurons. J Neurosci *18*, 7613–
1192   7624. https://doi.org/10.1523/JNEUROSCI.18-19-07613.1998.

1193   69.   Golding, N.L., Mickus, T.J., Katz, Y., Kath, W.L., and Spruston, N. (2005). Factors
1194   mediating powerful voltage attenuation along CA1 pyramidal neuron dendrites. J
1195   Physiol *568*, 69–82. https://doi.org/10.1113/jphysiol.2005.086793.

1196   70.   Makarov, R., Chavlis, S., and Poirazi, P. (2024). DendroTweaks: An interactive
1197   approach for unraveling dendritic dynamics. eLife *13*.
1198   https://doi.org/10.7554/eLife.103324.1.

1199   71.   Magee, J.C., and Cook, E.P. (2000). Somatic EPSP amplitude is independent of
1200   synapse location in hippocampal pyramidal neurons. Nat Neurosci *3*, 895–903.
1201   https://doi.org/10.1038/78800.

1202   72.   Häusser, M. (2001). Synaptic function: dendritic democracy. Curr Biol *11*, R10-12.
1203   https://doi.org/10.1016/s0960-9822(00)00034-8.

1204   73.   Nicholson, D.A., Trana, R., Katz, Y., Kath, W.L., Spruston, N., and Geinisman, Y.
1205   (2006). Distance-dependent differences in synapse number and AMPA receptor
1206   expression in hippocampal CA1 pyramidal neurons. Neuron *50*, 431–442.
1207   https://doi.org/10.1016/j.neuron.2006.03.022.

1208   74.   Asrican, B., Lisman, J., and Otmakhov, N. (2007). Synaptic strength of individual spines
1209   correlates with bound Ca2+-calmodulin-dependent kinase II. J Neurosci *27*, 14007–
1210   14011. https://doi.org/10.1523/JNEUROSCI.3587-07.2007.

1211   75.   Klausberger, T., and Somogyi, P. (2008). Neuronal diversity and temporal dynamics:
1212   the unity of hippocampal circuit operations. Science *321*, 53–57.
1213   https://doi.org/10.1126/science.1149381.

1214   76.   Bezaire, M.J., Raikov, I., Burk, K., Vyas, D., and Soltesz, I. (2016). Interneuronal
1215   mechanisms of hippocampal theta oscillations in a full-scale model of the rodent CA1
1216   circuit. Elife *5*, e18566. https://doi.org/10.7554/eLife.18566.

1217   77.   Dudok, B., Klein, P.M., Hwaun, E., Lee, B.R., Yao, Z., Fong, O., Bowler, J.C., Terada,
1218   S., Sparks, F.T., Szabo, G.G., et al. (2021). Alternating sources of perisomatic inhibition
1219   during behavior. Neuron *109*, 997-1012.e9.
1220   https://doi.org/10.1016/j.neuron.2021.01.003.

1221   78.   Price, C.J., Scott, R., Rusakov, D.A., and Capogna, M. (2008). GABA(B) receptor
1222   modulation of feedforward inhibition through hippocampal neurogliaform cells. J
1223   Neurosci *28*, 6974–6982. https://doi.org/10.1523/JNEUROSCI.4673-07.2008.

1224   79.   Szabadics, J., Tamás, G., and Soltesz, I. (2007). Different transmitter transients
1225   underlie presynaptic cell type specificity of GABAA,slow and GABAA,fast. Proc Natl
1226   Acad Sci U S A *104*, 14831–14836. https://doi.org/10.1073/pnas.0707204104.

1227   80.   Maccaferri, G., Roberts, J.D., Szucs, P., Cottingham, C.A., and Somogyi, P. (2000).
1228   Cell surface domain specific postsynaptic currents evoked by identified GABAergic

neurones in rat hippocampus in vitro. J Physiol *524 Pt 1*, 91–116.
https://doi.org/10.1111/j.1469-7793.2000.t01-3-00091.x.

81.   Lee, S.-H., Marchionni, I., Bezaire, M., Varga, C., Danielson, N., Lovett-Barron, M.,
Losonczy, A., and Soltesz, I. (2014). Parvalbumin-positive basket cells differentiate
among hippocampal pyramidal cells. Neuron *82*, 1129–1144.
https://doi.org/10.1016/j.neuron.2014.03.034.

82.   Fuentealba, P., Begum, R., Capogna, M., Jinno, S., Márton, L.F., Csicsvari, J.,
Thomson, A., Somogyi, P., and Klausberger, T. (2008). Ivy cells: a population of nitric-
oxide-producing, slow-spiking GABAergic neurons and their involvement in
hippocampal network activity. Neuron *57*, 917–929.
https://doi.org/10.1016/j.neuron.2008.01.034.

83.   Skaggs, W., McNaughton, B., and Gothard, K. (1992). An Information-Theoretic
Approach to Deciphering the Hippocampal Code. In Advances in Neural Information
Processing Systems (Morgan-Kaufmann).

84.   Mizuseki, K., Royer, S., Diba, K., and Buzsáki, G. (2012). Activity dynamics and
behavioral correlates of CA3 and CA1 hippocampal pyramidal neurons. Hippocampus
22, 1659–1680. https://doi.org/10.1002/hipo.22002.

85.   Jarsky, T., Roxin, A., Kath, W.L., and Spruston, N. (2005). Conditional dendritic spike
propagation following distal synaptic activation of hippocampal CA1 pyramidal neurons.
Nat Neurosci *8*, 1667–1676. https://doi.org/10.1038/nn1599.

86.   Bittner, K.C., Grienberger, C., Vaidya, S.P., Milstein, A.D., Macklin, J.J., Suh, J.,
Tonegawa, S., and Magee, J.C. (2015). Conjunctive input processing drives feature
selectivity in hippocampal CA1 neurons. Nat Neurosci *18*, 1133–1142.
https://doi.org/10.1038/nn.4062.

87.   Stuart, G.J., and Spruston, N. (2015). Dendritic integration: 60 years of progress. Nat
Neurosci *18*, 1713–1721. https://doi.org/10.1038/nn.4157.

88.   Lundberg, S.M., Erion, G., Chen, H., DeGrave, A., Prutkin, J.M., Nair, B., Katz, R.,
Himmelfarb, J., Bansal, N., and Lee, S.-I. (2020). From Local Explanations to Global
Understanding with Explainable AI for Trees. Nat Mach Intell *2*, 56–67.
https://doi.org/10.1038/s42256-019-0138-9.

89.   Wei, D.S., Mei, Y.A., Bagal, A., Kao, J.P., Thompson, S.M., and Tang, C.M. (2001).
Compartmentalized and binary behavior of terminal dendrites in hippocampal pyramidal
neurons. Science *293*, 2272–2275. https://doi.org/10.1126/science.1061198.

90.   Segev, I., and Rall, W. (1998). Excitable dendrites and spines: earlier theoretical
insights elucidate recent direct observations. Trends Neurosci *21*, 453–460.
https://doi.org/10.1016/s0166-2236(98)01327-7.

91.   Sidiropoulou, K., and Poirazi, P. (2012). Predictive features of persistent activity
emergence in regular spiking and intrinsic bursting model neurons. PLoS Comput Biol
8, e1002489. https://doi.org/10.1371/journal.pcbi.1002489.

92.   Smith, S.L., Smith, I.T., Branco, T., and Häusser, M. (2013). Dendritic spikes enhance
stimulus selectivity in cortical neurons in vivo. Nature *503*, 115–120.
https://doi.org/10.1038/nature12600.

93.   Ujfalussy, B.B., and Makara, J.K. (2020). Impact of functional synapse clusters on
neuronal response selectivity. Nat Commun *11*, 1413. https://doi.org/10.1038/s41467-
020-15147-6.

94.   Larkum, M.E., and Nevian, T. (2008). Synaptic clustering by dendritic signalling
mechanisms. Curr Opin Neurobiol *18*, 321–331.
https://doi.org/10.1016/j.conb.2008.08.013.

95.   Kastellakis, G., and Poirazi, P. (2019). Synaptic Clustering and Memory Formation.
Front Mol Neurosci *12*, 300. https://doi.org/10.3389/fnmol.2019.00300.

96.   Chirillo, M.A., Waters, M.S., Lindsey, L.F., Bourne, J.N., and Harris, K.M. (2019). Local
resources of polyribosomes and SER promote synapse enlargement and spine
clustering after long-term potentiation in adult rat hippocampus. Sci Rep *9*, 3861.
https://doi.org/10.1038/s41598-019-40520-x.

97.   Harris, K.M. (2020). Structural LTP: from synaptogenesis to regulated synapse
enlargement and clustering. Curr Opin Neurobiol *63*, 189–197.
https://doi.org/10.1016/j.conb.2020.04.009.

98.   Karbowski, J., and Urban, P. (2024). Cooperativity, Information Gain, and Energy Cost
During Early LTP in Dendritic Spines. Neural Comput *36*, 271–311.
https://doi.org/10.1162/neco_a_01632.

99.   Bittner, K.C., Andrasfalvy, B.K., and Magee, J.C. (2012). Ion channel gradients in the
apical tuft region of CA1 pyramidal neurons. PLoS One *7*, e46652.
https://doi.org/10.1371/journal.pone.0046652.

100. O'Hare, J.K., Wang, J., Shala, M.D., Polleux, F., and Losonczy, A. (2025). Distal tuft
dendrites predict properties of new hippocampal place fields. Neuron, S0896-
6273(25)00249-1. https://doi.org/10.1016/j.neuron.2025.03.029.

101. Bloss, E.B., Cembrowski, M.S., Karsh, B., Colonell, J., Fetter, R.D., and Spruston, N.
(2018). Single excitatory axons form clustered synapses onto CA1 pyramidal cell
dendrites. Nat Neurosci *21*, 353–363. https://doi.org/10.1038/s41593-018-0084-6.

102. Milstein, A.D., Li, Y., Bittner, K.C., Grienberger, C., Soltesz, I., Magee, J.C., and
Romani, S. (2021). Bidirectional synaptic plasticity rapidly modifies hippocampal
representations. eLife *10*, e73046. https://doi.org/10.7554/eLife.73046.

103. Magee, J.C., and Grienberger, C. (2020). Synaptic Plasticity Forms and Functions.
Annual Review of Neuroscience *43*, 95–117. https://doi.org/10.1146/annurev-neuro-
090919-022842.

104. Rolotti, S.V., Blockus, H., Sparks, F.T., Priestley, J.B., and Losonczy, A. (2022).
Reorganization of CA1 dendritic dynamics by hippocampal sharp-wave ripples during
learning. Neuron *110*, 977-991.e4. https://doi.org/10.1016/j.neuron.2021.12.017.

105. Golding, N.L., Kath, W.L., and Spruston, N. (2001). Dichotomy of action-potential
backpropagation in CA1 pyramidal neuron dendrites. J Neurophysiol *86*, 2998–3010.
https://doi.org/10.1152/jn.2001.86.6.2998.

106. Zhou, W.-L., Yan, P., Wuskell, J.P., Loew, L.M., and Antic, S.D. (2008). Dynamics of
action potential backpropagation in basal dendrites of prefrontal cortical pyramidal

neurons. Eur J Neurosci *27*, 923–936. https://doi.org/10.1111/j.1460-
9568.2008.06075.x.

107. Meyer-Dilhet, G., and Courchet, J. (2020). In Utero Cortical Electroporation of Plasmids
in the Mouse Embryo. STAR Protoc *1*, 100027.
https://doi.org/10.1016/j.xpro.2020.100027.

108. Virga, D.M., Hamilton, S., Osei, B., Morgan, A., Kneis, P., Zamponi, E., Park, N.J.,
Hewitt, V.L., Zhang, D., Gonzalez, K.C., et al. (2024). Activity-dependent
compartmentalization of dendritic mitochondria morphology through local regulation of
fusion-fission balance in neurons in vivo. Nat Commun *15*, 2142.
https://doi.org/10.1038/s41467-024-46463-w.

109. Preibisch, S., Saalfeld, S., and Tomancak, P. (2009). Globally optimal stitching of tiled
3D microscopic image acquisitions. Bioinformatics *25*, 1463–1465.
https://doi.org/10.1093/bioinformatics/btp184.

110. Ascoli, G.A. (2006). Mobilizing the base of neuroscience data: the case of neuronal
morphologies. Nat Rev Neurosci *7*, 318–324. https://doi.org/10.1038/nrn1885.

111. Harris, K.M., and Kater, S.B. (1994). Dendritic spines: cellular specializations imparting
both stability and flexibility to synaptic function. Annu Rev Neurosci *17*, 341–371.
https://doi.org/10.1146/annurev.ne.17.030194.002013.

112. Arellano, J.I., Benavides-Piccione, R., DeFelipe, J., and Yuste, R. (2007). Ultrastructure
of Dendritic Spines: Correlation Between Synaptic and Spine Morphologies. Front
Neurosci *1*, 131–143. https://doi.org/10.3389/neuro.01.1.1.010.2007.

113. Peng, H., Long, F., and Myers, G. (2011). Automatic 3D neuron tracing using all-path
pruning. Bioinformatics *27*, i239-247. https://doi.org/10.1093/bioinformatics/btr237.

114. Chen, J.L., Villa, K.L., Cha, J.W., So, P.T.C., Kubota, Y., and Nedivi, E. (2012).
Clustered dynamics of inhibitory synapses and dendritic spines in the adult neocortex.
Neuron *74*, 361–373. https://doi.org/10.1016/j.neuron.2012.02.030.

115. Shuman, T., Aharoni, D., Cai, D.J., Lee, C.R., Chavlis, S., Page-Harley, L., Vetere,
L.M., Feng, Y., Yang, C.Y., Mollinedo-Gajate, I., et al. (2020). Breakdown of spatial
coding and interneuron synchronization in epileptic mice. Nat Neurosci *23*, 229–238.
https://doi.org/10.1038/s41593-019-0559-0.

116. Solstad, T., Moser, E.I., and Einevoll, G.T. (2006). From grid cells to place cells: a
mathematical model. Hippocampus *16*, 1026–1031. https://doi.org/10.1002/hipo.20244.

117. Krächan, E.G., Fischer, A.U., Franke, J., and Friauf, E. (2017). Synaptic reliability and
temporal precision are achieved via high quantal content and effective replenishment:
auditory brainstem versus hippocampus. J Physiol *595*, 839–864.
https://doi.org/10.1113/JP272799.

118. Bezaire, M.J., and Soltesz, I. (2013). Quantitative assessment of CA1 local circuits:
knowledge base for interneuron-pyramidal cell connectivity. Hippocampus *23*, 751–785.
https://doi.org/10.1002/hipo.22141.

119. Papadopoulos, P.M., Katz, M.J., and Bruno, G. (2001). NPACI: rocks: tools and
techniques for easily deploying manageable Linux clusters. In Proceedings 2001 IEEE

1353   International Conference on Cluster Computing, pp. 258–267.
1354   https://doi.org/10.1109/CLUSTR.2001.959986.

1355   120. Hines, M.L., and Carnevale, N.T. (2001). NEURON: a tool for neuroscientists.
1356   Neuroscientist *7*, 123–135. https://doi.org/10.1177/107385840100700207.

1357

1358 **Supplementary figures**

1359

1360 ***Figure S1: Dendritic diameter smoothing.*** *Radius profiles of three example dendritic branches before (A) and*

1361 *after (B) application of the smoothing algorithm. Profiles are shown for Tuft (left), Apical (center), and Basal (right)*

1362 *branches, with radius plotted as a function of the distance from the soma.*

1363

1364

1365

1366 ***Figure S2: Probability of inhibitory input sources targeting synapses in CA1 pyramidal cells.*** *Distribution of*

1367 *inhibitory presynaptic input probabilities for synapses located on the Tuft (A), Apical (B), and Basal (C) dendrites.*

1368 *The probability varies with dendro-somatic path distance, capturing how different interneuron types preferentially*

1369 *target specific dendritic regions. Histograms show the number of synapses as a function of distance from the soma,*

1370 *while the overlaid curves represent the probability that a synapse is targeted by each interneuron type.*

1371

1372

1373

1374

1375

Figure S3: Characterization of synaptic clusters under alternative input allocation protocols. *Cluster volume*

(A) and size (B) were significantly reduced following synaptic shuffling across all protocols. Statistically

significant differences were observed between the control configuration and each of the shuffled conditions,

with variation maintained across dendritic domains. Pairwise t-tests between control and shuffled protocols

are indicated in the plots (*p < 0.05, **p < 0.005, ***p < 0.001, and ****p < 0.0001), with corrections for

multiple comparisons. Error-bars represent the 95% confidence interval (CI) of the mean.

Figure S4. SHAP summary plots for feature impact on spatial coding metrics. *SHAP value distributions for*

the prediction of four output features: (A) spatial information (bits/s), (B) in-field firing frequency (Hz), (C) out-of-

field firing frequency (Hz), and (D) place field size (cm). Each dot represents a SHAP value for a given sample,

with its color indicating the relative value of the corresponding input feature (blue = low, red = high). Features are

ranked by their mean absolute SHAP value, reflecting their overall contribution to model output (Feature

Importance). The numerical values on the right of each feature represent the Pearson correlation between the input

feature values and their associated SHAP values, providing an estimate of the direction and linearity of the feature's

effect on each output metric.