Preserved reptile scales retain microscopic features, revealing a new instance of convergent evolution

Calvin A. Riiska[1], Gordon W. Schuett[2,3],

Joseph R. Mendelson III[4,5], Jennifer M. Rieser[1]

[1]Emory University Department of Physics, Atlanta, GA

[2]Department of Biology — Neuroscience Institute, Georgia State University, Atlanta, GA

[3]Chiricahua Desert Museum, Rodeo, NM

[4]Department of Research, Zoo Atlanta, Atlanta, GA

[5]School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA

May 30, 2025

Abstract

Small-scale structures on biological surfaces can profoundly impact how animals move, appear, and interact with their environments. Such textures may be especially important for limbless reptiles, such as snakes and legless lizards, because their skin serves as the primary interface with the world around them. Here, we examine ventral microstructures of several limbless reptiles, which are hypothesized to be highly specialized to aid locomotion via frictional interactions. Inspired by prior studies that investigated potential links between microtextures, phylogeny, habitat, and locomotion—but that were limited by their reliance on shed skins—we characterized the structures present on preserved museum specimens and

found that they are quantitatively similar to those found on shed skins. Using this result, we confirmed a previously hypothesized—but untested due to the lack of shed skins—third independent evolution of sidewinding-specific isotropic microtexture. Specifically, we examined a museum-preserved *Bitis peringueyi* specimen and identified a new instance of convergent evolution in sidewinding viper microstructures: the loss of micro-spikes (present on many snake species) and the appearance of micro-pits with a characteristic spacing. Our results reveal that museum-preserved specimens retain intact microtextures, greatly expanding the availability of samples for evolutionary studies.

1   Introduction

The diverse textures present on animal skins or exoskeletons often play crucial roles in mediating interactions with the environment. Remarkably, microtextures with features on the micron or nanometer scale, often requiring the use of sub-diffraction-limit imaging techniques to fully characterize, can influence environmental interactions at macroscopic scales by modulating interactions with light, sound, fluids, and other biological matter [1]. For example, reflective microstructures on birds and butterflies can enhance coloration for mating displays [2], reduce visibility for predator avoidance [3, 4], and aid in thermoregulation [5]. Sound absorbing micro-scales on moth wings provide acoustic camouflage from predatory bats [6–8]. Hydrophobic textures help shed water from insect wings [9, 10], and when combined with hydrophilic structures, can enable water harvesting in both desert beetles [11] and rattlesnakes [12] as well as adhesion to air-water interfaces [13]. Micron-sized denticles on shark skins help to streamline flows around their bodies [14, 15], enhance thrust [16], and protect delicate tissues like gill flaps [17]. Microstructures can also interlock (as in bird feathers [18], enhancing lift during flight) or prevent adhesion (as in carnivorous pitcher plants [19], aiding in prey capture by destabilizing insect footholds).

Microstructures can also directly enhance locomotion through specialized interactions with other surfaces. For instance, setae on gecko feet utilize van der Waals forces to allow

these animals to adhere to both smooth and rough surfaces [20–22]. Similar structures are found on the feet of flies and beetles and, when combined with a sticky fluid secretion, these insects can cling to a variety of surfaces as well [23–26]. Other insects such as ants, stick insects, and katydids have smooth pads with internal fibers that distort around surface features to increase contact-area-dependent forces that are further enhanced by the presence of a fluid secretion [23, 24, 26–28].

For limbless animals moving across terrestrial substrates, the ventral surface serves as the primary contact with the environment during locomotion. Consequently, it has been hypothesized that specialized ventral microstructures likely evolved to enhance frictional interactions. Many snake species have rearward-pointing micro-spikes that are thought to aid movement through the creation of anisotropic friction—providing low friction for forward sliding, high friction to prevent backward sliding [29], and high lateral friction to reduce sideways slip and enhance forward locomotion [30, 31]. Intriguingly, these structures also appear to be linked to habitat; qualitative observations suggest that arboreal snakes have longer spikes than terrestrial species, potentially enhancing friction forces that are crucial for traversing trees, while aquatic and marine species tend to have shorter spikes, perhaps due to the decreased importance of interactions with solid substrates [32]. Notably, in some desert-dwelling sidewinding vipers, micro-spikes are either greatly reduced or completely absent. Instead, they possess an isotropic array of pits, found to have evolved independently at least twice in distantly related species of vipers [33]. Computational modeling revealed that sidewinding, a periodic pattern of movement in which a planar wave of body curvature is coupled to a vertical wave of body lifting (where lifted segments are moved forward and non-lifted segments are pressed against the substrate) that produces an overall sideways motion [34, 35], is enhanced by isotropic friction, which these isotropic structures are hypothesized to produce [33].

While these results strongly suggest an important interplay between microstructures, movement strategies, habitats, and phylogeny, establishing definitive links has been con-

67   strained by limited sample availability; previous studies have largely relied on shed skins,

68   which can be difficult to obtain for rare or elusive species [33, 36].

Figure 1: The ventral scales of snakes possess microscopic textures. (a) A corn-snake (*Pantherophis guttatus*). (b) Zoomed-in image of several ventral scales, which serve as the primary animal-substrate interface during locomotion. (c) Microstructures within a 20 μm $\times$ 20 μm region of a single ventral scale from the shed skin of a cornsnake, imaged using atomic force microscopy. Many snake species have similar textures, composed of repeating rows of head-to-tail oriented spike-like features, that are thought to enhance frictional interactions during movement.

In contrast, natural history museums house extensive collections of formalin-fixed and

ethanol-preserved reptile specimens, samples of which are made available to researchers upon

request. These samples have been widely used for biological research [37, 38], including bio-

metric (e.g., limb sizes, bone shapes, and allometry) [39] and histological (e.g. cell arrange-

ments and tissue strength) [40] measurements, information on basic natural history and life

habits [41], and DNA extraction [42] for genome sequencing to understand the origin of

traits [43] or to recover phylogenetic relationships [44]. While the preservation process is

known to have some effects on reptile specimens, e.g., shortened snout-vent lengths, lower

body masses [45], and reduced pigmentation [38], the effects of preservation on fine-scale sur-

face microstructures are not known. Some previous studies have included measurements from

preserved as well as shed snake skins [32, 46], however, whether the geometry of preserved

structures differs from those on sheds was not explored or discussed. Given the stiffness of

snake skin, which is composed of keratin [47–49], we hypothesize that these microstructures

may remain intact through the processes of specimen fixation and preservation. If true, this

would open extensive museum collections to quantitative analysis of keratin microstructures across diverse species [50], thus alleviating the limitation of sampling based on availability of shed skins.

In this study, we used atomic force microscopy (AFM) to demonstrate that formalin-fixed, ethanol-preserved museum specimens retain microstructures comparable to those present on shed skins. Unlike the more widely used scanning electron microscopy (SEM) [51–53], AFM provides nanometer-resolution topographical measurements that enable precise quantification of in-plane geometry, vertical height profiles, and spatial anisotropy. Using the approach described below, we quantitatively assess both the height distributions and dominant spacings of features within skin microstructures. We first show that textures on shed skins of cornsnakes *Pantherophis guttatus* are similar within an individual (i.e., from distinct regions within a single scale), across individuals (i.e., from sheds collected from two different snakes), as well as for two preserved museum specimens. We then demonstrate that this structural similarity holds for multiple snake species as well as for a legless lizard (*Ophisaurus attenuatus*), and conclude that microstructures are retained over decades throughout the preservation and storage processes. Finally, we leverage these results to confirm a previously untested hypothesis [33], identifying a new, fourth instance of the pitted microstructure associated with sidewinding locomotion from a preserved Peringuey's adder *Bitis peringueyi*, a species known to use sidewinding locomotion. This finding reveals a third independent evolutionary origin of sidewinding behavior accompanied by the loss of micro-spikes in favor of an isotropic pitted microstructure. Further, this result provides a concrete example of how museum collections can expand sample availability to enable hypothesis testing and broader comparative studies of the qualities and evolution of specialized ventral microstructures, ultimately generating unique insights that could establish new links between behavior, habitat, and phylogeny.

2 Results

We used AFM to image 20 μm $\times$ 20 μm sections on snake ventral scales (see Methods). These images revealed that the surface features and detailed morphologies of microstructures on preserved specimens closely resembled those on shed skin samples (Figure 2). We examined museum samples and shed samples from four snake species—the colubrids (*Pantherophis guttatus* (cornsnake) and *Thamnophis saurita* (ribbonsnake)), and the viperids (*Crotalus polystictus* (Mexican lance-headed rattlesnake) and *Crotalus atrox* (western diamond-backed rattlesnake)). In each case, the ventral microstructures consisted of periodic rows of raised, rearward-pointing, overlapping spikes, typically $1-5$ μm in length. AFM enabled precise height measurements at the nanometer scale, revealing that spikes in *P. guttatus*, for example, were raised about 100 nm above the surface. Secondary features such as ridges and small pits are observed between spikes or between rows of spikes. These microstructures are remarkably well preserved, even in decades-old specimens; our oldest sample, from *C. polystictus*, was collected and preserved in 1939.

We also observed similar keratinous microstructures on another reptile. In ventral scales from both shed skin and museum samples of the legless slender glass lizard *Ophisaurus atten-uatus*, the shapes of cell borders and sub-micron-scale pits appear similar in both samples. These observations suggest that a wide range of keratin surface structures can be studied using preserved specimens across a broad taxonomic diversity of reptiles, greatly expanding sample availability for comparative and evolutionary studies.

2.1 Quantitative similarity within and across shed skins

To determine the extent to which ventral microstructures are similar within and across sam-ples, as well as across sample types, we developed a robust quantification method inspired by previous work [33]. All analyses began with AFM-acquired height maps of 60 μm $\times$ 60 μm regions. We started by quantifying features on a shed skin sample (s1) from *P. guttatus*,

Figure 2: Microstructure topographies for shed skins and preserved samples.
AFM scans of 20 μm x 20 μm regions on shed skins (top row) and museum-preserved samples
(bottom row). Structures appear similar for four snake species (*P. guttatus*, *C. polystictus*,
C. atrox, and *T. saurita*, left four columns) and a glass lizard (*O. attenuatus*, right column).
Microstructures are retained even for older samples—the *C. polystictus* sample was preserved
in 1939. Scale bars: 5 μm, and the top of each image is toward the head of the animal.

chosen because of its common use in locomotion studies (e.g., [54–58])and characterizing its

microstructure may elucidate links to movement. To quantify the geometric arrangement

of microstructural features, we first Gaussian filtered each image to remove depressions and

bulges occurring over tens of microns caused by irregularities in the sample (see Figures

S2, S3, S5, and S6). These variations occur over much longer length scales (at least tens

of microns) than the spiked structures we are interested in characterizing; removing these

larger-scale variations allows for direct comparisons of the heights and spacings of the spiked

features of interest. The resulting mean-subtracted image displays height variations asso-

ciated with in-plane microstructures that can be compared across sample types. To assess

within-sample variability, we imaged six distinct regions (denoted a–f, spaced about 5 mm

apart) along a single ventral scale from a shed skin sample of *P. guttatus*(Figure 3a). To

compare height distributions across images, we used the Jensen-Shannon (JS) divergence,

which is bounded between zero and one and measures the distinguishability of two distri-

butions. A JS divergence value of zero indicates that two distributions are identical while

a value of one indicates that the distributions are perfectly distinguishable. We found that the height distributions across these sites were similar (Figure 3b), with low JS divergence values (≤ 0.04) between each pair of distributions (see Table S1).

For in-plane structure, we expanded previous analyses [33] to quantify two key spatial metrics: the spacing between the rows of spikes, r, and the lateral spacing between adjacent spikes, s (see labels in Figure 3a). Starting with our Gaussian filtered, mean-subtracted height map (Figure S3), we first computed the 2D power spectrum (Figure 3c). The frequencies associated with the two brightest peaks in the power spectrum correspond to high lateral periodicity (the spike spacing, s), while secondary peaks above and below the primary peaks reflect vertical periodicity (the row spacing, r). The broadness of the primary and secondary peaks as well as the slight rotational offset—caused by imperfections in sample mounting—made extracting dominant peak locations and orientations from these images challenging; therefore, we used a Radon transform, which projects the intensity of a 2D image onto lines of varying orientation that pass through the center of the image, to identify the directions associated with dominant peaks.

The Radon transform of the power spectrum (Figure 3d) is brightest for the orientation, θ_r, that projects the primary intensity peaks in the power spectrum onto the origin. In the slice at θ_r, the secondary peaks above and below the primary power spectrum peaks collapse, creating smaller off-center peaks that identify the spatial frequency associated with the vertical spacing, r, between rows of spikes. The orientation, θ_s, with the largest spacing between primary peaks in the Radon transform corresponds to the spatial frequency associated with the lateral spacing, s, between adjacent spikes.

Vertical slices through the Radon transform, taken at θ_r and θ_s, are shown in Figure 3e. Row and spike spatial frequencies and uncertainties were estimated by locally fitting Gaussians to relevant peaks (smaller off-center peaks for rows and off-center peaks for spikes; see Methods Section 4.4 and Figure S1 for full details). At site s1c, the row spacing is $r = 3.2 \pm 0.3$ μm, and the spike spacing is $s = 1.0 \pm 0.1$ μm. This approach allows for sys-

tematic comparisons across sites, and across six sites within one sample, differences between

sites fell within measurement uncertainty of one site, (Figure 3f, g), indicating structural

similarity along the scale.

To quantify the directional variability of the microstructure, we measured the anisotropy

index, a quantity introduced in a previous study [33]. The anisotropy index of a region is

defined as the JS divergence between vertical slices of the Radon transform (of the power

spectrum) separated by 90°. An anisotropy index close to zero indicates a uniform structure

with little direction dependence while larger values indicate higher directionality. In our cal-

culation, we divided each image into four quadrants and calculated the average and standard

deviation of the anisotropy index over all quadrants. The anisotropy index at site s1c was

0.51 ± 0.01 indicating a large degree of directionality. While anisotropy uncertainties do not

always overlap, the anisotropy values are comparable across sites, providing a baseline for

structural variability in like samples.

A second shed skin sample (s2) from *P. guttatus* showed that the height distribution and

spacing of structures do not vary between individuals (Figure 4c-e). This sample had a row

spacing of $r = 3.3 \pm 0.3$ μm and a spike spacing of $s = 1.1 \pm 0.1$ μm, nearly identical to s1c.

Structures from different individuals were also similarly anisotropic—s2 had a calculated

anisotropy of 0.62 ± 0.02—confirming that microstructural features are consistent between

individuals of the same species (Figure 4f).

2.2 Microstructures are retained on preserved samples

To assess similarities and differences across sample types, we compared microstructures from

two shed skins and two preserved samples of *P. guttatus* (Figure 4a,b). Distributions of

feature heights (Figure 4c) were consistent across all samples, with JS divergences ≤ 0.04,

comparable to within-sample variation. Use of spatial frequency analyses described above

showed that preserved specimens retained row- and spike-spacing values similar to those from

shed skins (Figure 4d,e). While one museum specimen (m1) showed slightly higher average

Figure 3: Quantification of microscopic features reveals structural similarity across a single scale. **(a)** A zoomed-in view of microstructures on a cornsnake shed skin (s1). Two quantities, r and s (the average row and spike spacing, respectively), are used to characterize and compare samples. Scale bar: 2 μm. **(b)** A third quantity, the relative height variations within a sample, characterized through distributions of mean-subtracted and rescaled feature heights, provides an additional comparative measurement. Distributions of relative height variations from evenly spaced sites within a single scale (as illustrated by the inset) indicate similarities across sites. **(c)** Power spectrum of zoomed-out version of (a), with dominant and secondary peaks associated with in-plane structural periodicities along and across spikes. Scale bar: 1 μm^{-1}. **(d)** Radon transform of (c). The angular direction, θ_r, where the two primary peaks collapse is identified as the angle with maximal intensity. **(e)** Slices of the Radon transform used to calculate r and s. Secondary peaks in the dominant direction slice (blue) correspond to the spatial frequency of rows of spikes. Peaks in the slice where the primary peaks collapse furthest apart (θ_s, red) correspond to the lateral spatial frequency of the spikes. The **(f)** row spacing, **(g)** spike spacing, and **(h)** anisotropy indices are all similar across the six sites in the sample.

Figure 4: Quantification of structural similarities across samples and sample types. AFM images of cornsnake microstructures of **(a)** a shed skin and **(b)** a preserved museum sample are visually similar. Scale bars: 5 μm. **(c)** Distributions of height variations from two different individuals each for shed skins (s1 and s2) and museum samples (m1 and m2). The **(d)** row spacing and **(e)** spike spacing are quantitatively similar across sample types. **(f)** The anisotropy indices are slightly more variable but exhibit no systematic trend.

spacings ($r = 4.1 \pm 0.5$ μm; $s = 1.2 \pm 0.2$ μm), these values fell within the range of natural variation. The second museum specimen (m2) closely matched the shed skin measurements ($r = 3.3 \pm 0.4$ μm; $s = 0.9 \pm 0.1$ μm). It is worth noting that we also examined whether body size might influence structural spacing. Using ventral scale width as a proxy, we ranked specimens from largest to smallest: s1 (31.7 mm), m1 (25.0 mm), s2 (17.9 mm), and m2 (12.6 mm). No trend emerged linking individual size to microstructural spacing.

Extending our analysis beyond *P. guttatus*, we applied the same quantification technique to preserved samples from three additional snake species—*C. polystictus*, *C. atrox*, and *T. saurita*—and a glass lizard, *O. attenuatus.* In all snake species, preserved samples had row and spike spacings and anisotropy indices consistent with those of shed skins (Figure 5). In

Figure 5: Quantitative comparison of structural measurements across sample types for several species. (a) The preserved vs shed row spacings for four snake species and one lizard species all lie close to the dashed gray $y = x$ line. **(b)** The preserved vs shed spike spacings also lie close to $y = x$, indicating similarity of different sample types and multiple species. **(c)** The anisotropy indices are also similar across sample types (including a frozen sample for *C. atrox*, blue triangle). Note that *O. attenuatus* only has a dominant structural frequency in the head-to-tail direction.

O. attenuatus, which only showed a dominant periodicity in the head-to-tail direction, both preserved and shed samples exhibited similar cell row dimensions and anisotropy indices, supporting the conclusion that keratin-based microtextures are preserved across diverse reptilian taxa.

2.3   A third independent origin of pitted microstructure in sidewinding snakes

Using our result that microstructures are preserved on museum specimens, we analyzed a museum-preserved sample of *Bitis peringueyi*, a sidewinding true viper (Viperinae) native to the Namib Desert of southern Africa [59]. We found that its ventral microstructure closely resembles those of other sidewinding viper species. Previous work [33] identified a distinctive pitted microstructure—markedly different from and lacking the micro-spikes typical of other snakes—in three sidewinding species from two phylogenetically distant genera: the pitviper (Crotalinae) *Crotalus cerastes*, and the true vipers (Viperinae) *Cerastes cerastes*, and *Cerastes vipera*. However, that study was limited to shed skin samples, and no shed skins were available for *Bitis peringueyi*. By leveraging our results from this study and us-

ing a museum-preserved specimen, we were able to include this fourth viperid species and expand the evolutionary context of sidewinding microstructures.

This result reveals a third evolutionary origin of the pitted sidewinding microstructure, suggesting strong selective pressure for this trait among sidewinding vipers. Sidewinding has evolved independently in desert-dwelling members of the Viperinae and Crotalinae sub-families without notable changes in body morphology [60, 61]. Among the species studied here—three in Viperinae and one in Crotalinae—all exhibit similar pitted structures while micro-spikes are absent (Figure 6a,b). Based on the phylogeny in Figure 6a, which was adapted from [44], our results here and those found previously [33] indicate three instances of convergent evolution of ventral microstructures linked to sidewinding locomotion, likely in response to similar environmental challenges of sandy substrates [62].

Using power spectra and Radon transforms (Figure S12), we measured the characteristic spatial frequency of the pitted textures finding that all four sidewinding species, including *Bitis peringueyi*, had an average pit spacing between 0.9 μm and 1.5 μm with overlapping uncertainties up to ±0.6 μm (Figure 6c). This overlap suggests convergence not only in structure type, but possibly in pit spacing as well. Additionally, all sidewinding species show low anisotropy indices (Figure 6d), reflecting the isotropic nature of the pitted texture. Notably, *Bitis peringueyi* conforms to the isotropic pitted pattern, providing another example of convergent evolution in both behavior and potentially beneficial ventral microstructures.

3   Discussion and conclusions

In this work, we used AFM to demonstrate that microstructures on the keratin-based skins of several preserved snakes and one lizard are each topographically similar to shed skin samples from the same species. Few studies have systematically quantified the in-plane geometry of snake microstructures; most have provided qualitative observations [52, 53] or hand measured the size of features on the surface [33]. By collecting height profile data and calculating feature spacings, we developed robust analyses that enabled comparisons of

Figure 6: A preserved *B. peringueyi* sample reveals a third origin of convergently evolved microstructure in sidewinding vipers. **(a)** A phylogeny of viperids adapted from [44] shows the three branches represented in our sidewinder microstructure images. It is most parsimonious to conclude three independent origins of these microstructures, and sidewinding locomotion, than to propose a singular origin in the clade Viperidae with dozens of independent losses. **(b)** A similar pitted structure appears in each of the four sidewinding species imaged. Scale bars: 5 μm.**(c)** The spacing between pits in each species is similar, suggesting this structure may have evolved with a characteristic spacing. **(d)** The microstructure of each sidewinder species is similarly isotropic.

ventral microstructures of reptiles from both shed skins and those found on preserved skins.

Our analyses revealed that preserved specimens, including those preserved for over 85 years, retained their fine-scale surface structures. Specifically, we found that surface heights, spike spacings, and anisotropy indices were similar across sample types, with inter-sample variability falling within intra-sample variability. This validation greatly expands the range of specimens available for microstructural analysis beyond the limited supply and availability of shed skins and opens the door for broad studies linking structure to habitat, phylogeny, and locomotion strategy.

We leveraged our results to investigate an untested hypothesis [33] which discovered two evolutionarily independent losses of micro-spikes in favor of isotropic pitted microstructures found on three sidewinding viper species. A computational model demonstrated that such isotropic structures may enhance frictional interactions that benefit sidewinding movement, and therefore, the authors predicted similar microstructures on a fourth sidewinding species, *B. peringueyi*. While we were also unable to obtain a shed skin for this species, our results here indicate that preserved museum specimens can be used instead of shed skins. Indeed, AFM images of a museum specimen revealed that *B. peringueyi* has the predicted pitted microstructure observed on the other sidewinding vipers. Comparisons of quantified structures demonstrated that all sidewinding species had a similar spacing between pits, suggesting that this structure may have evolved with a characteristic geometry.

We note that although the morphology of snake microstructures was retained in museum samples, the effects of preservation on mechanical properties—e.g., elasticity, friction, and wear—of the skin are not known. Though some prior work has measured elasticity [63,64], microscopic friction [29,30,36,65], and microstructure wear and durability [64] for shed skins from a few species, how structures relate to mechanical properties is not well understood. Future work should explore these connections further, as well as investigate how chemical fixatives used in preservation may or may not affect mechanical properties. Either way, linking microstructures to mechanics, even if only shed skin samples can be used, would

not only advance our understanding of biological surface function but could also inspire the design of engineered materials with tunable anisotropic friction [66,67].

While differences in mechanical properties across sample types remain to be explored, our results show that preserved specimens may be used in the place of shed skins to study structural properties of reptile skins. This finding greatly expands opportunities for high-resolution comparative studies across a wide range of species, including rare or elusive taxa. As the environment affects both the body shape [68] and skin thickness [69] of snakes, it is reasonable to hypothesize that the form and function of microstructures can vary as well. For example, dorsal microstructures in snakes often are hydrophobic and show strong phylo-genetic signal [53,70]. In contrast, ventral structures appear more closely tied to locomotion and ecological niche, such as observations of longer spikes on arboreal snakes (which may enhance frictional interactions) and shorter or smoother structures that may reduce drag or debris accumulation for aquatic and fossorial species [32,71,72]. Museum collections can provide the phylogenetic diversity needed to gain new insights into drivers of the evolution of structures that are crucial for environmental interactions.

4   Methods

4.1   Sample preservation

The preservation process involves injecting the animal with formalin (3.7% formaldehyde), positioning it, and wrapping it in formalin-soaked cloth to cross-link proteins, position the specimen in a desired configuration, and prevent decomposition [37,73]. Samples are then preserved indefinitely in 70% ethanol, which is relatively non-toxic and prevents microbial and fungal growth [74]. Some specimens are preserved directly in ethanol, which better preserves DNA, though ethanol concentrations must remain high to prevent decay [37]. Frozen samples were collected freshly deceased and kept on ice for two days, then submerged in water and placed in a freezer so ice would encase the specimen. Shed skins were collected

from specimens housed at various locations, including the Rieser lab at Emory University and Zoo Atlanta in Atlanta GA (see Table S2), and stored in sealed Ziplock bags at room temperature.

4.2   Sample preparation

For this study, we examined three different types of snake skin samples: those from shed skins, preserved samples, and frozen samples. For shed skins: sample s2 (*P. guttatus*) and the sample from *O. attenuatus* were collected from our lab population after each animal had naturally shed; sample s1 (*P. guttatus*) was collected at Zoo Atlanta, Georgia, along with samples from *Crotalus polystictus* and *Crotalus cerastes*; samples from *Crotalus atrox* were obtained from a captive population in the Chiricahua Desert Museum in New Mexico. Sections of 2–3 ventral scales were cut out of the sheds for imaging.

Preserved samples were obtained from either the Field Museum of Natural History in Chicago, Illinois, or the Amphibian and Reptile Diversity Research Center, University of Texas at Arlington. These samples came from specimens that had been formalin fixed and preserved in $\geq 70\%$ ethanol. Samples from undamaged sections of freshly dead road-killed adult specimens of *C. atrox* from Cochise County, Arizona, were formalin-fixed and ethanol-preserved using standard museum protocols; additional samples of skin from those specimens also were frozen for subsequent attempts at AFM imaging. To obtain skin from these samples for imaging, the top layer of skin on individual ventral scales was peeled off of the specimens.

Initially, we attempted to image frozen and preserved samples while the skin was still attached to the scale tissue. This approach proved to be difficult because the sample would relax during imaging and the AFM tip would lose contact. This problem could potentially be remedied by letting the samples dry overnight or by imaging them while partially submerged in a bath of water (but with top surface exposed to air) but we found that these techniques worked inconsistently. The uppermost dermal layer, the stratum corneum, of skin would also occasionally become detached. As a result, we removed this layer and imaged it directly.

Some museum samples that we obtained, however, did not have the top layer of skin present. We initially assumed that it was not removable on some samples, but after imaging the top layer of tissue on these samples, no microstructures were found. We concluded that the top layer of skin was absent in these cases; it is common to find the top layers of reptile scales detached from museum specimens and accumulated at the bottom of specimen jars. In order to accurately capture microstructures on museum or frozen samples, the stratum corneum layer of the epidermis must be present.

Before imaging, all skin samples were cleaned, using a previously established protocol [33], to remove dirt and debris that may cover microstructures. The cut out or peeled off ventral scales were placed in a solution of 5% Micro-90 mixed with deionized water. We then used a Cole Parmer ultrasonic cleaner to sonicate the samples in the solution for 30 minutes. After sonicating, the samples were rinsed with tap water to remove the Micro-90 solution. The samples were then laid flat on a wire screen and left to air dry in the lab for at least 12 hours to ensure they were completely dry before imaging. We used two-sided scotch tape to adhere the scales to glass microscope slides. After this, the samples were ready for imaging. We ensured that the cleaning and drying process does not affect the microstructure itself by imaging samples that were clean without washing. Nevertheless, small amounts of residual debris are expected and do appear in some of the analyzed images.

Our study does not capture variations in the structure based on the location on the body (reviewed in [32], [46], and [52]), nor any potential ontogenetic changes in ventral microstructures that may occur. Related specifically to museum specimens, detailed history of chemical methods of fixation and preservation often are not available for all specimens. Because fixation in approximately 10% formalin solution and preservation in 70% ethanol is traditional, we assume that our specimens borrowed from museums were prepared in this way. Our sampling and information available does not allow us to determine if microstructures persist under all possible methods of specimen preparation (e.g., fixation and preservation in isopropanol). All samples were taken from ventral scales in the middle third of each snake's

body.

4.3   Data collection: AFM imaging

All samples were imaged using a Jupiter XR large-sample AFM from Asylum Research at Oxford Instruments. We used tapping mode imaging with either a 160AC or 240AC integrated silicon nitride probe. 20μm $\times$ 20μm and 60μm $\times$ 60μm sections of each sample were imaged with 512 pixel resolution at 1 Hz and 0.5 Hz speeds respectively.

To identify structural variation across the same sample, we imaged the same sample of shed skin from *P. guttatus* at six locations across the same scale roughly spaced 5-mm apart laterally. All other images in this study were taken in a central region of a scale. To show the variation between different samples of from the same species, we imaged a second shed skin sample from *P. guttatus*. We then compared structures across sample types by comparing these images to those taken from two different preserved samples of *P. guttatus*. All images except those comparing variation across the same sample were taken in the central region of a scale. A similar series of images was taken for *C. atrox* samples including frozen specimens and analyzed in Figure S4 to show the viability of frozen samples.

4.4   Analysis: Quantifying the structures

Images from the AFM were flattened, meaning an offset was removed from each scan line so that they all have the same average value, and the overall slope of each scan line is removed. This algorithm is built in to the Igor-based AFM imaging software provided by Asylum Research. It is not a plane fit, instead acting on each scan line. The order of flattening can be selected where order one removes a linear fit, order two removes a quadratic, and so on. For each image, order one was chosen. Flattening accounts for thermal drift and sudden changes in tip behavior, both of which can introduce nonphysical offsets in height between consecutive scan lines and slopes in the sample. Flattening also corrects for small imperfections in sample mounting where it may be impossible to get the sample perfectly flat

to the nanoscale. Flattened images were taken to be raw images for our purposes. The raw images still had larger scale depressions and bulges occurring over tens of microns, a larger scale than the spikes and pits we wanted to characterize. To bring the features of interest into the same plane, we applied a Gaussian filter with a standard deviation of 10 pixels to each image and subtracted the filter from the original image. The final step in processing was mean-subtracting each image. (See Figures S2, S3, S5, and S6 for the results of filtering images. Distributions of raw heights are shown in Figure S7.)

We first compared the distribution of heights in each of the filtered and mean subtracted images. By calculating the Jensen-Shannon (JS) divergence between pairs of distributions, we quantified how similar the heights of features and roughness are across images and sample types.

To calculate the periodicity of in-plane surface features, we built off a previous analysis technique [33]. We calculated a power spectrum of the height values in each image to identify dominant spatial frequencies. Using a fast Fourier transform rather than a power spectrum yielded similar results (Figure S13). Images may be slightly tilted from their desired orientation, so we calculated a Radon transform with an angular increment of $0.25°$ from these power spectra to more easily measure the location of dominant frequencies independent of the axes. In order to accurately capture the spatial frequency of the structure, imaged large regions that contained at least 15-20 rows of micro-spikes. For sidewinders, 20 μm × 20 μm images were deemed to have enough features to be adequate for this analysis. However, 20 μm × 20 μm images from non-sidewinding species only captured a few rows of spikes, so 60 μm ×60 μ m images with several rows of spikes were used for this analysis. (Power spectra and Radon transforms for each of our samples are shown in Figures S8, S9, S10, S11, and S12.)

For spiked structures (all those on non-sidewinding snake species), the Radon transform slice with the highest peak has two secondary peaks whose frequency values correspond to the spacing between rows of spikes in the image. The dominant peaks in the slice with two

dominant peaks furthest apart corresponds to the lateral spacing between spikes (see Figure S1a–b). In each case, the data was smoothed to identify the desired peaks and a linear model was fit to the minima on either side of the peak (Figure S1c,e). Subtracting the linear model from the slice allows us to fit a Gaussian curve to the peak and measure its width (Figure S1d,f). The mean of this distribution corresponds to the spatial frequency and the standard deviation gives a measure of the variation in the spatial frequency across the image. From these values we are able to calculate the row spacing, spike spacing, and uncertainties associated with each.

Images from sidewinding species produced a much more uniform Radon transform. By taking an average of all slices in the Radon transform, we were able to identify mirrored peaks on either side of 0 Hz corresponding to a singular dominant spatial frequency. We subtracted the average Radon value at 0 Hz from the data and fit a Gaussian distribution to each side. We calculated the spacing between pits and the associated uncertainty using the mean and standard deviation of these distributions.

We also adapted a previous technique [33] to calculate an anisotropy index from each sample. We first divided each image into quadrants. We then calculated a power spectrum and Radon transform for each. From each Radon transform, we took a slice in the dominant θ_d direction containing the maximum intensity, and a second slice at $\theta_d \pm 90°$. Note that these slices are different from those used to calculate the row and spike spacing. Those slices tended to be $\approx 90°$ apart but were found independently. The JS divergence between the two slices serves as the anisotropy index for that quadrant. Finding the mean and standard deviation of the anisotropy indices gives a measure of the anisotropy index and its uncertainty for the whole image.

Author contributions

C.A.R.: conceptualization, data curation, data interpretation, formal analysis, investigation, methodology, software, visualization, writing–original draft, writing–review and edit-

ing. G.W.S.: sample acquisition, writing–review and editing. J.R.M.III: conceptualiza-
tion, resources, sample acquisition, writing–review and editing. J.M.R.: conceptualization,
data interpretation, methodology, project administration, resources, software, supervision,
writing–review and editing.

Acknowledgments

The authors thank the Field Museum of Natural History The University of Texas at Arlington
for loans of specimens; Jessica Tingle and Gordon Berman for helpful discussions.

5 References

[1] Calvin A. Riiska, Chantal Nguyen, Orit Peleg, and Jennifer M. Rieser. The Physics of Animal Behavior: Form, Function, and Interactions. *Annual Review of Condensed Matter Physics*, 15(1):325–350, March 2024.

[2] Dakota E. McCoy, Allison J. Shultz, Charles Vidoudez, Emma Van Der Heide, Jacqueline E. Dall, Sunia A. Trauger, and David Haig. Microstructures amplify carotenoid plumage signals in tanagers. *Scientific Reports*, 11(1):8582, April 2021.

[3] Pete Vukusic and J Roy Sambles. Photonic structures in biology. *Nature*, 424:5, 2003.

[4] Aline Vieira-Silva, Gabriel B. Evora, André V. L. Freitas, and Paulo S. Oliveira. The Relevance of Flash Coloration Against Avian Predation in a *Morpho* Butterfly: A Field Experiment in a Tropical Rainforest. *Ethology*, 130(12):e13517, December 2024.

[5] Anirudh Krishna, Xiao Nie, Andrew D. Warren, Jorge E. Llorente-Bousquets, Adriana D. Briscoe, and Jaeho Lee. Infrared optical and thermal properties of microstructures in butterfly wings. *Proceedings of the National Academy of Sciences*, 117(3):1566–1572, January 2020.

[6] Zhiyuan Shen, Thomas R. Neil, Daniel Robert, Bruce W. Drinkwater, and Marc W. Holderied. Biomechanics of a moth scale at ultrasonic frequencies. *Proceedings of the National Academy of Sciences*, 115(48):12200–12205, November 2018.

[7] Thomas R. Neil, Zhiyuan Shen, Daniel Robert, Bruce W. Drinkwater, and Marc W. Holderied. Moth wings are acoustic metamaterials. *Proceedings of the National Academy of Sciences*, 117(49):31134–31141, December 2020.

[8] Thomas R. Neil, Zhiyuan Shen, Daniel Robert, Bruce W. Drinkwater, and Marc W. Holderied. Moth wings as sound absorber metasurface. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 478(2262):20220046, June 2022.

[9] Shubham Sharma, R. Uma Shaanker, and Anand Kumar Subramaniyan. Anti-wetting wing surface characteristics of a water bug, Diplonychus annulatus. *BioSystems*, 248:105402, February 2025.

[10] Jiyu Sun and Bharat Bhushan. The structure and mechanical properties of dragonfly wings and their role on flyability. *Comptes Rendus. Mécanique*, 340(1-2):3–17, December 2011.

[11] Andrew R. Parker and Chris R. Lawrence. Water capture by a desert beetle. *Nature*, 414(6859):33–34, November 2001.

[12] Akshay Phadnis, Kenneth C Manning, Gordon W Schuett, and Konrad Rykaczewski. Role of scale wettability on rain-harvesting behavior in a desert-dwelling rattlesnake. *ACS omega*, 4(25):21141–21147, 2019.

[13] Chiaki Suzuki, Yasuharu Takaku, Hiroshi Suzuki, Daisuke Ishii, Tateo Shimozawa, Shuhei Nomura, Masatsugu Shimomura, and Takahiko Hariyama. Hydrophobic-hydrophilic crown-like structure enables aquatic insects to reside effectively beneath the water surface. *Communications Biology*, 4(1):708, June 2021.

[14] Meagan Popp, Connor F. White, Diego Bernal, Dylan K. Wainwright, and George V. Lauder. The denticle surface of thresher shark tails: Three-dimensional structure and comparison to other pelagic species. *Journal of Morphology*, 281(8):938–955, August 2020.

[15] George V. Lauder, Dylan K. Wainwright, August G. Domel, James C. Weaver, Li Wen, and Katia Bertoldi. Structure, biomimetics, and fluid dynamics of fish skin surfaces. *Physical Review Fluids*, 1(6):060502, October 2016.

[16] Benjamin S. Savino and Wen Wu. Thrust generation by shark denticles. *Journal of Fluid Mechanics*, 1000:A80, December 2024.

[17] Molly K Gabler-Smith, Dylan K Wainwright, Greta A Wong, and George V Lauder. Dermal Denticle Diversity in Sharks: Novel Patterns on the Interbranchial Skin. *Integrative Organismal Biology*, 3(1):obab034, December 2021.

[18] Laura Y. Matloff, Eric Chang, Teresa J. Feo, Lindsie Jeffries, Amanda K. Stowers, Cole Thomson, and David Lentink. How flight feathers stick together to form a continuous morphing wing. *Science*, 367(6475):293–297, January 2020.

[19] David Labonte, Adam Robinson, Ulrike Bauer, and Walter Federle. Disentangling the role of surface topography and intrinsic wettability in the prey capture mechanism of Nepenthes pitcher plants. *Acta Biomaterialia*, 119:225–233, January 2021.

[20] K. Autumn and Anne M. Peattie. Mechanisms of Adhesion in Geckos. *Integrative and Comparative Biology*, 42(6):1081–1090, December 2002.

[21] G Huber, S Gorb, N Hosoda, R Spolenak, and E Arzt. Influence of surface roughness on gecko adhesion. *Acta Biomaterialia*, 3(4):607–610, July 2007.

[22] Yu Tian, Noshir Pesika, Hongbo Zeng, Kenny Rosenberg, Boxin Zhao, Patricia McGuiggan, Kellar Autumn, and Jacob Israelachvili. Adhesion and friction in gecko toe attachment and detachment. *Proceedings of the National Academy of Sciences*, 103(51):19320–19325, December 2006.

[23] R. G. Beutel and S. N. Gorb. Ultrastructure of attachment specializations of hexapods (Arthropoda): evolutionary patterns inferred from a revised ordinal phylogeny. *Journal of Zoological Systematics and Evolutionary Research*, 39(4):177–207, December 2001.

[24] Jan-Henning Dirks and Walter Federle. Fluid-based adhesion in insects – principles and challenges. *Soft Matter*, 7(23):11047, 2011.

[25] S.N. Gorb. The design of the fly adhesive pad: distal tent setae are adapted to the delivery of an adhesive secretion. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 265(1398):747–752, May 1998.

[26] Thomas B. H. Schroeder, Jared Houghtaling, Bodo D. Wilts, and Michael Mayer. It's Not a Bug, It's a Feature: Functional Materials in Insects. *Advanced Materials*, 30(19):1705322, May 2018.

[27] David Labonte and Walter Federle. Functionally Different Pads on the Same Foot Allow Control of Attachment: Stick Insects Have Load-Sensitive "Heel" Pads for Friction and Shear-Sensitive "Toe" Pads for Adhesion. *PLoS ONE*, 8(12):e81943, December 2013.

[28] Constanze Grohmann, Miriam Judith Henze, Thomas Nørgaard, and Stanislav N. Gorb. Two functional types of attachment pads on a single foot in the Namibia bush cricket *Acanthoproctus diadematus* (Orthoptera: Tettigoniidae). *Proceedings of the Royal Society B: Biological Sciences*, 282(1809):20142976, June 2015.

[29] Martina J. Baum, Alexander E. Kovalev, Jan Michels, and Stanislav N. Gorb. Anisotropic Friction of the Ventral Scales in the Snake Lampropeltis getula californiae. *Tribology Letters*, 54(2):139–150, May 2014.

[30] Martina J. Benz, Alexander E. Kovalev, and Stanislav N. Gorb. Anisotropic frictional properties in snakes. In Akhlesh Lakhtakia, editor, *Bioinspiration, Biomimetics, and Bioreplication 2012*, volume 8339, page 83390X. International Society for Optics and Photonics, SPIE, 2012.

[31] David L Hu, Jasmine Nirody, Terri Scott, and Michael J Shelley. The mechanics of slithering locomotion. *Proceedings of the National Academy of Sciences*, 106(25):10081–10085, 2009.

[32] C.V. Schmidt and S. Gorb. *Snake Scale Microstructure: Phylogenetic Significance and Functional Adaptations.* Zoologica (E. Schweizerbart'sche Verlagsbuchhandlung). Schweizerbart, 2012.

[33] Jennifer M. Rieser, Tai-De Li, Jessica L. Tingle, Daniel I. Goldman, and Joseph R. Mendelson. Functional consequences of convergently evolved microscopic skin features on snake locomotion. *Proceedings of the National Academy of Sciences*, 118(6):e2018264118, February 2021.

[34] Carl Gans and Kim Hyung L. KINEMATIC DESCRIPTION OF THE SIDEWINDING LOCOMOTION OF FOUR VIPERS. *Israel Journal of Ecology and Evolution*, 38(1):9 – 23, 1992. Place: Leiden, The Netherlands Publisher: Brill.

[35] B. C. Jayne. Kinematics of Terrestrial Snake Locomotion. *Copeia*, 4:915–927, 1986. Journal: Copeia. 1986(4). 915-927.

[36] J Hazel, M Stone, M.S Grace, and V.V Tsukruk. Nanoscale design of snake skin for reptation locomotions via friction anisotropy. *Journal of Biomechanics*, 32(5):477–484, May 1999.

[37] Steve W. Gotte, Jeremy F. Jacobs, and George R. Zug. Preserving reptiles for research. In C. Kenneth Dodd, editor, *Reptile Ecology and Conservation*, pages 73–86. Oxford University Press, May 2016.

[38] Elliott R Jacobson, Harvey B Lillywhite, and Daniel G Blackburn. Overview of biology, anatomy, and histology of reptiles. *Diseases and Pathology of Reptiles*, 1:1–214, 2021.

[39] Robert R Sokal and F James Rohlf. *Biometry*. Macmillan, 1995.

[40] Giuseppe Musumeci. Past, present and future: overview on histology and histopathology. *J Histol Histopathol*, 1(5):1–3, 2014.

[41] Randall S Reiserer, Gordon W Schuett, and Harry W Greene. Seed ingestion and germination in rattlesnakes: overlooked agents of rescue and secondary dispersal. *Proceedings of the Royal Society B: Biological Sciences*, 285(1872):20172755, 2018.

[42] Claus M. Zacho, Martina A. Bager, Ashot Margaryan, Peter Gravlund, Anders Galatius, Arne R. Rasmussen, and Morten E. Allentoft. Uncovering the genomic and metagenomic research potential in old ethanol-preserved snakes. *PLOS ONE*, 16(8):e0256353, August 2021.

[43] Chandra Shekhar Pareek, Rafal Smoczynski, and Andrzej Tretyn. Sequencing technologies and genome sequencing. *Journal of Applied Genetics*, 52(4):413–435, November 2011.

[44] Laura R.V. Alencar, Tiago B. Quental, Felipe G. Grazziotin, Michael L. Alfaro, Marcio Martins, Mericien Venzon, and Hussam Zaher. Diversification in vipers: Phylogenetic relationships, time of divergence and shifts in speciation rates. *Molecular Phylogenetics and Evolution*, 105:50–62, December 2016.

[45] Bart Vervust, Stefan Van Dongen, and Raoul Van Damme. The effect of preservation on lizard morphometrics – an experimental study. *Amphibia-Reptilia*, 30(3):321–329, 2009.

[46] R. A. Berthé, G. Westhoff, H. Bleckmann, and S. N. Gorb. Surface structure and frictional properties of the skin of the Amazon tree boa Corallus hortulanus (Squamata, Boidae). *Journal of Comparative Physiology A*, 195(3):311–318, March 2009.

[47] Alberto Ripamonti, Lorenzo Alibardi, Giuseppe Falini, Simona Fermani, and Massimo Gazzano. Keratin-lipid structural organization in the corneous layer of snake. *Biopolymers*, 91(12):1172–1181, December 2009.

[48] Mattia Toni and Lorenzo Alibardi. Alpha- and beta-keratins of the snake epidermis. *Zoology*, 110(1):41–47, February 2007.

[49] J. McKittrick, P.-Y. Chen, S. G. Bodde, W. Yang, E. E. Novitskaya, and M. A. Meyers. The Structure, Functions, and Mechanical Properties of Keratin. *JOM*, 64(4):449–468, April 2012.

[50] Andrew V. Suarez and Neil D. Tsutsui. The Value of Museum Collections for Research and Society. *BioScience*, 54(1):66, 2004.

[51] P Russell, D Batchelor, and J Thornton. SEM and AFM: Complementary Techniques for High Resolution Surface Investigations. *Veeco Instruments Inc., AN46, Rev A*, 1:2004, 2001.

[52] Marie-Christin G. Klein and Stanislav N. Gorb. Ultrastructure and wear patterns of the ventral epidermis of four snake species (Squamata, Serpentes). *Zoology*, 117(5):295–314, October 2014.

[53] Marcelle I. Arrigo, Luis M. De Oliveira Vilaca, Anamarija Fofonjka, Achyuthan N. Srikanthan, Adrien Debry, and Michel C. Milinkovitch. Phylogenetic mapping of scale nanostructure diversity in snakes. *BMC Evolutionary Biology*, 19(1):91, December 2019.

[54] Bruce C. Jayne, Steven J. Newman, Michele M. Zentkovich, and H. Matthew Berns. Why arboreal snakes should not be cylindrical: body shape, incline and surface roughness have interactive effects on locomotion. *Journal of Experimental Biology*, 218(24):3978–3986, December 2015.

[55] Henry C. Astley and Bruce C. Jayne. Arboreal habitat structure affects the performance and modes of locomotion of corn snakes (*Elaphe guttata*). *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology*, 311A(3):207–216, March 2009.

[56] Derek J. Jurestovsky, Logan R. Usher, and Henry C. Astley. Generation of propulsive force via vertical undulations in snakes. *Journal of Experimental Biology*, 224(13):jeb239020, July 2021.

[57] Rachel H. Mansfield and Bruce C. Jayne. Arboreal habitat structure affects route choice by rat snakes. *Journal of Comparative Physiology A*, 197(1):119–129, January 2011.

[58] Hamidreza Marvi and David L. Hu. Friction enhancement in concertina locomotion of snakes. *Journal of The Royal Society Interface*, 9(76):3067–3080, November 2012.

[59] D.G. Broadley. *FitzSimons' Snakes of Southern Africa*. Delta Books, 1983.

[60] Jessica L Tingle. Facultatively Sidewinding Snakes and the Origins of Locomotor Specialization. *Integrative and Comparative Biology*, 60(1):202–214, July 2020.

[61] Jessica L Tingle and Theodore Garland Jr. Morphological evolution in relationship to sidewinding, arboreality and precipitation in snakes of the family Viperidae. *Biological Journal of the Linnean Society*, page 18, 2021.

[62] H. Marvi, C. Gong, N. Gravish, H. Astley, M. Travers, R. L. Hatton, J. R. Mendelson, H. Choset, D. L. Hu, and D. I. Goldman. Sidewinding with minimal slip: Snake and robot ascent of sandy slopes. *Science*, 346(6206):224–229, October 2014.

[63] Marie-Christin G. Klein and Stanislav N. Gorb. Epidermis architecture and material properties of the skin of four snake species. *Journal of The Royal Society Interface*, 9(76):3140–3155, November 2012.

[64] Marie-Christin G. Klein, Julia K. Deuschle, and Stanislav N. Gorb. Material properties of the skin of the Kenyan sand boa Gongylophis colubrinus (Squamata, Boidae). *Journal of Comparative Physiology A*, 196(9):659–668, September 2010.

[65] Weibin Wu, Shudong Yu, Paul Schreiber, Antje Dollmann, Christian Lutz, Guillaume Gomard, Christian Greiner, and Hendrik Hölscher. Variation of the frictional anisotropy on ventral scales of snakes caused by nanoscale steps. *Bioinspiration & Biomimetics*, 15(5):056014, August 2020.

[66] Weibin Wu, Markus Guttmann, Marc Schneider, Richard Thelen, Matthias Worgull, Guillaume Gomard, and Hendrik Hölscher. Snake-Inspired, Nano-Stepped Surface with Tunable Frictional Anisotropy Made from a Shape-Memory Polymer for Unidirectional Transport of Microparticles. *Advanced Functional Materials*, 31(19):2009611, May 2021.

[67] Martina J Baum, Lars Heepe, Elena Fadeeva, and Stanislav N Gorb. Dry friction of microstructured polymer surfaces inspired by snake skin. *Beilstein Journal of Nanotechnology*, 5:1091–1103, July 2014.

[68] Laura Rodrigues Vieira de Alencar, Marcio Martins, Gustavo Burin, and Tiago Bosisio Quental. Arboreality constrains morphological evolution but not species diversification in vipers. *Proceedings of the Royal Society B: Biological Sciences*, 284(1869):20171775, December 2017.

[69] Richard Shine, Claire Goiran, Catherine Shilton, Shai Meiri, and Gregory P Brown. The life aquatic: an association between habitat type and skin thickness in snakes. *Biological Journal of the Linnean Society*, page blz136, October 2019.

[70] Marlene Spinner, Stanislav N. Gorb, Alexander Balmert, Horst Bleckmann, and Guido Westhoff. Non-Contaminating Camouflage: Multifunctional Skin Microornamentation in the West African Gaboon Viper (Bitis rhinoceros). *PLoS ONE*, 9(3):e91087, March 2014.

[71] C. Gans and D. Baic. Regional Specialization of Reptilian Scale Surfaces: Relation of Texture and Biologic Role. *Science*, 195(4284):1348–1350, March 1977.

[72] David J. Gower. Scale microornamentation of uropeltid snakes. *Journal of Morphology*, 258(2):249–268, November 2003.

[73] Rooban Thavarajah, Vidya Kazhiyur Mudimbaimannar, Joshua Elizabeth, Umadevi Krishnamohan Rao, and Kannan Ranganathan. Chemical and physical

basics of routine formaldehyde fixation. *Journal of oral and maxillofacial pathology,* 16(3):400–405, 2012.

[74] John E Simmons. *Things great and small: Collection management policies.* Rowman & Littlefield, 2024.