Article

Phenotypic plasticity as a function of genetic polymorphism: thermal dominance reversal

in *Drosophila* species with contrasting melanism

Jean R. David[1,2*], Béatrice Denis[1], Pascaline Francelle[3], Alicia Lemaire[1], Aparup Das[4], Sujata

Mohanty[5], Patricia Gibert[6], Amir Yassin[1,2**]

[1]Laboratoire Évolution, Génomes, Comportement et Écologie, CNRS, IRD, Université Paris-

Saclay – Institut Diversité, Écologie et Évolution (IDEEV), 91190 Gif-sur-Yvette, France.

[2]Institut Systématique Evolution Biodiversité (ISYEB) CNRS, MNHN, Sorbonne Université,

EPHE, Université des Antilles, 57 rue Cuvier, CP 50, 75005 Paris, France.

[3]School of Biological and Behavioural Sciences, Queen Mary University of London, Mile End

Road, London E1 4NS, United Kingdom.

[4]ICMR-Regional Medical Research Centre, Sri Vijaya Puram, Andaman and Nicobar Islands,

744103, India.

[5]Department of Biotechnology, Jaypee Institute of Information Technology, Sector 62, Noida,

Uttar Pradesh-201309, India.

[6]Université Claude Bernard Lyon 1, Laboratoire de Biométrie et Biologie Evolutive, CNRS,

UMR5558, Villeurbanne, Rhônes-Alpes, 69100, France.

Running title: Phenotypic plasticity and heterozygosity

*Deceased in 2021.

Corresponding author: Laboratoire Évolution, Génomes, Comportement et Écologie,

CNRS, IRD, Université Paris-Saclay – Institut Diversité, Écologie et Évolution (IDEEV), 12 route

128, 91190 Gif-sur-Yvette, France. Email : amir.yassin@universite-paris-saclay.fr

Authors contribution

J.R.D. – conceptualization, formal analysis, investigation, resources; B.D. – resources, writing review and editing; P.F. – data curation, formal analysis, visualization; A.L. – data curation, formal analysis, visualization; A.D. – investigation, resources; S.M. – investigation, resources; P.G. – writing review and editing; A.Y. – conceptualization, data curation, methodology, supervision, visualization, writing original draft.

Conflicts of interest

The authors declare no conflict of interest.

Data availability statement

All data associated with this study are provided in a Figshare Folder and will be released upon publication.

Acknowledgments

The authors thank Artyom Kopp for comments on the manuscript and for sharing with Yuichi Fukutomi the association between female-limited color dimorphism and polymorphism at the *pdm3* locus in *D. jambulina* and *D.* cf. *bocqueti*.

Abstract

Phenotypic plasticity is often seen as an alternative adaptive strategy to genetic polymorphism, especially in response to rapid environmental changes. Indeed, a link between plasticity and heterozygosity, *i.e.* the measure of polymorphism, has previously been dismissed. Here, we compare the thermal plasticity of abdominal pigmentation in eight *Drosophila* species, four belonging to the *melanogaster* species group and four to the *montium* group. Despite a conserved developmental pathway for melanin synthesis, the genetic architecture of its variation has significantly evolved, being polygenic in most species (like *D. melanogaster*) and Mendelian or invariable in others. By investigating the thermal plasticity of this trait in species with distinct architectures, we show the degree of plasticity to strongly associate with heterozygosity. Plasticity was resurrected in hybrids between species with no plastic responses but with contrasting melanism, and was higher in heterozygotes in species with simple Mendelian polymorphism. In plastic cases, pigmentation dominance is reversed depending on the developmental temperature. We propose simple genetic models with empirical molecular support to explain this link between phenotypic plasticity and genetic polymorphism. The relationship between these two phenomena, and the impact of each on the evolution of the other, may be more relevant than it is currently appreciated.

Keywords

Overdominance; genetic architecture; reaction norm; heterozygosity; dominance reversal.

Introduction

Phenotypic plasticity is the ability of a genotype to produce different phenotypes under distinct environmental conditions (Bradshaw 1965; Scheiner 1993; Pigliucci 2001; West-Eberhard 2003; Sultan 2021). When optimal phenotypic values are generated in the different environments, plasticity can facilitate adaptation even in the absence of genetic variation (Ghalambor et al. 2015). This becomes particularly important in cases where the population loses a significant portion of genetic polymorphism, such as after a founder bottleneck during colonization of a new environment, or when the environment change is faster than the rate of accumulation of new adaptive mutations, the so-called 'buying time' role of plasticity (Diamond and Martin 2021). Phenotypic plasticity and genetic polymorphism might be considered as distinct alternative adaptive strategies (e.g., Wennersten and Forsman 2012).

Despite the definition of plasticity as a property of a genotype, it is usually studied for complex phenotypic traits whose underlying genotypes are not fully known (e.g., flowering time in *Arabidopsis*, growth in yeasts, etc.) (Goldstein and Ehrenreich 2021; Kovuri et al. 2023). Genotypes do not only differ in the degree of their phenotypic plasticity (e.g., the slope of the reaction norm) but also in their degree of genetic variability, with heterozygous genotypes in non-haploid organisms having more alleles than hemi- or homozygotes. Among the early genetic models of phenotypic plasticity, the Overdominance model explicitly related plasticity to genotypic diversity (Scheiner 1993). The model postulated that plasticity is a function of homozygosity, with the heterozygous genotype considered to be the most advantageous because it is the most robust, *i.e.* the least plastic. This assumption reflected the view of plasticity at the mid-20[th] century as a developmental nuisance (Lerner 1954; Bradshaw 2006). Indeed, some of the earliest empirical examples of the overdominance model came from studies of wing morphology in *Drosophila melanogaster*, where two mutants showed the highest temperature-sensitivity when found in the homozygous state, but their heterozygote

recapitulated the wildtype thermal response, *i.e.* it was less plastic than either mutant (Harnly

and Harnly 1936; Schmalhausen 1949). Gillespie and Turelli (1989) extended this view to a

multi-locus quantitative genetics model and concluded that stabilizing selection on more

environmentally robust heterozygotes will maintain genetic variation in populations. Pigliucci

(1992) developed a one-locus, two-allele classical population genetics model for testing the

evolution of plasticity under overdominance and concurred with Gillespie and Turelli's (1989)

multilocus result. He concluded that plasticity becomes "increasingly disadvantageous when

the system approaches a perfect equilibrium between the probabilities of occurrence of the

two environments."

A direct link between heterozygosity and robustness was not always evident from the

empirical literature, and overdominance was disfavored relative to other genetic models such

as pleiotropy, *i.e.* plasticity is a function of the expression of a gene, and epistasis, *i.e.* plasticity

is a function of interactions between loci. Scheiner (1993) concluded that "plasticity is not a

function of heterozygosity", and Pigliucci (2005), a decade later, stated that "heterozygosity

ha[d] little, if anything, to do with the genetic basis of plasticity." A search using "heterozygo*"

in a recent 400-page long book on *Phenotypic Plasticity and Evolution* (Pfennig 2021) yielded

no result. However, almost none of the studies cited by Scheiner (1993) and Pigliucci (2005) to

reach their conclusions has addressed the link between heterozygosity and plasticity in the most

optimal setting similar to Pigliucci's (1992) theoretical model, *i.e.* comparing the reaction norms

of a phenotype with a simple, well-characterized genotypic basis. Conspicuous color

polymorphisms are ideally suited for these experiments, because they often involve a simple

genetic architecture and color synthesis in many species results from a well conserved gene

network (Orteu and Jiggins 2020). In several lineages of moths, simple Mendelian

polymorphisms have evolved in response to the industrial revolution, with the melanic wing

allele being dominant in some species and recessive in others (Ford 1945). The genetic basis of

this industrial melanism has recently been identified to be a conserved micro-RNA close to the

locus of the transcription factor *Cortex* (Tian et al. 2024). Remarkably, several butterfly species

also show a seasonal color polyphenism, with cold season morphs being darker. van der Burg

et al. (2020) showed that mutations at the same locus containing *Cortex*, and consequently the

linked micro-RNA, underlie this polyphenism in *Junonia caenia*. Plasticity in simple color

polymorphism systems was also observed in *Harmonia* ladybugs morphs (Michie et al. 2010).

Much of our knowledge about the genetic and developmental basis of melanin synthesis

comes from studies in *Drosophila*. Pigmentation differences between body segments, sexes,

geographical populations, seasons, and species are often due to *cis*- and *trans*-regulatory

modifications of melanin synthesis genes, such as *tan (t)*, *yellow (y)* or *ebony (e)*, and their

transcription factors, such as *bric-à-brac (bab)*, *POU-domain-motif-3 (pdm3)* or *Abdominal-B*

(Abd-B) (Massey and Wittkopp 2016; Hughes et al. 2023). Wild-caught flies from the same

locality usually differ in their degree of pigmentation (Figure 1 A; Bastide et al. 2013, 2016;

Dembeck et al. 2015; Endler et al. 2016; Castro et al. 2018), but flies collected from cold high

altitudes, latitudes, and seasons tend to be dark, even when their progeny is maintained under

standard thermal conditions (Telonis-Scott et al. 2011; Bastide et al. 2014, 2016; Takahashi et

al. 2014; Negoua et al. 2019; Berardi et al. 2024). Similarly, when full-sib flies issued from the

same isofemale line are grown under different temperatures, a negative reaction norm is

detected, with flies grown under low temperatures being darker (Figure 1; (Gibert et al. 1996,

1998, 2000; Chakir et al. 2002; Pétavy et al. 2002; Gibert et al. 2004b, 2009; Pétavy et al. 2018;

Lafuente et al. 2021, 2024). Thermal plasticity in *D. melanogaster* is due to the higher

expression of the *t* and *y* genes in colder temperatures (Gibert et al. 2016, 2017b). A natural

variation in the transcription factor *bab* underlying color differences also showed distinct

responses to developmental temperature through repression of *t* and *y* in hotter environments

(Castro et al. 2018). This parallelism between genetic polymorphism in the wild and phenotypic

plasticity in the lab is strongly suggestive for an adaptive role for darker pigmentation in thermoregulation. Indeed, drosophilids with darker cuticle have higher body temperature (Freoa et al. 2023).

Figure 1 – Pigmentation variation in *Drosophila melanogaster* and *D. kikkawai*. A) Variation in abdominal pigmentation in *D. melanogaster* can be due to genetic polymorphism (left) or thermal phenotypic plasticity (right). B) Contour plots showing pigmentation thermal plasticity in *D. melanogaster* females (left) and males (right) at each abdominal segment (A2-A7). C) Variation in abdominal pigmentation in *D. kikkawai* is caused by a single locus with two alleles dark (*D*) and light (*L*) whose effect is more pronounced on females (upper *DD* and middle *LL* photos) than males (lower photo). D) Contour plots showing pigmentation thermal plasticity in *D. kikkawai* females (above) and males (below) for the three genotypes *DD*, *DL*, and *LL*. For each abdominal segment, a reaction norm was inferred as a linear regression slope using mean values available published in Gibert et al. (1999, 2000). Photos are from Jean R. David's archive.

The genetic architecture of pigmentation variation has dramatically evolved between drosophilid species. Whereas female abdominal pigmentation variation is continuous and polygenic in *D. melanogaster* as well as most species, it is discrete and monogenic in multiple species (Payant 1986; Yassin et al. 2016a,b). Female-limited Mendelian melanism has evolved in *D. erecta* of the *melanogaster* group and in 22 species of the *montium* group belonging to five distinct subgroups (Yassin et al. 2016b; Yassin 2018). Of the 16 *montium* species where the genetics was studied, the dark allele was dominant in 13 species (Yassin et al. 2016b; Prigent et al. 2020). The phenotypic plasticity of abdominal pigmentation was investigated by Gibert et al. (1999) in one species, namely *D. kikkawai*, where females have two morphs and males have a light abdomen. The authors found major effect of genotype, segment, sex, and temperature (Figure 1B). In some other species, pigmentation is nearly fixed with very slight variability even between the sexes. This is the case of the two sister species, *D. santomea* and *D. yakuba*, where both males and females' posterior abdomens are invariably light or dark, respectively (Lachaise et al. 2000; Jeong et al. 2008; David et al. 2022).

In this paper, we compared the level of pigmentation phenotypic plasticity in eight species with distinct genetic architectures grown under five thermal regimes. To this end, we focused on the effect of heterozygosity, as it can be seen in heterozygous genotypes of five species with monogenic architectures as well as in species hybrids (*i.e.* the ultimate viable heterozygotes). Our results support a positive link between plasticity and the degree of genetic polymorphism that was not previously assumed.

Materials and Methods

Strains used and fly husbandry

We studied five cases where for each the thermal phenotypic plasticity can be compared between homozygous dark and light lines and their reciprocal hybrids. Four cases considered within-species monogenic Mendelian color polymorphism with one (*D. erecta*) from the *melanogaster* species group and three cases (*D. jambulina*, *D. burlai*, and *D.* cf. *bocqueti*) from its sister clade, the *montium* species group. For each of these species homozygous dark and light lines for the color causing allele were established through selection from a polymorphic mass culture (Table 1). The monogenic basis of female pigmentation in these species are known (Yassin et al. 2016a,b; Prigent et al. 2020). In two species (*D. erecta* and *D. burlai*), the dark allele is dominant, although the causative locus is different being the X-linked *tan* in *D. erecta* and the autosomal *pdm3* in *D. burlai* (Yassin et al. 2016a,b). In the two other species (*D. jambulina* and *D.* cf. *bocqueti*), the light allele is dominant (Watanabe et al. 1982; Prigent et al. 2020), but the causative locus is still not mapped although color variation associates with polymorphism in *pdm3* in the two species (Fukutomi et al. 2025). The fifth case involved hybridization between the two sister species *D. santomea* and *D. yakuba* of the *melanogaster* group, whose posterior segments are either monomorphically light or dark, respectively (Table 1). The pigmentation difference between these two species is controlled by mutations in at least six loci (Liu et al. 2019; David et al. 2022). All strains were kept in a rearing room at 21°C with 12:12 dark:light cycles on a food medium composed of 10 g agar, 101 g sugar, 25 g yeast, 100 g corn flour, 6.9 g nipagin, and 50 mL ethanol per 1 L. We also obtained published data of the thermal plasticity response of *D. melanogaster* (Gibert et al. 2000), the nomitypical species of the *melanogaster* group with polygenic basis of pigmentation variation, and the monogenic *D. kikkawai* (Gibert et al. 1999) for comparison purposes (Table 1).

208   Table 1. *Drosophila* strains analyzed in this study.

Group	Subgroup	Complex	Species	Color variation	Strain
melano-gaster	*melano-gaster*	*melano-gaster*	*melanogaster*	Complex	Bordeaux, France (Gibert et al. 2000)
melano-gaster	*melano-gaster*	*yakuba*	*yakuba mayottensis*	Nearly absent	Mayotte, France, 2013 (J. R. David, V. Debat, N. Gidaszewski)
melano-gaster	*melano-gaster*	*yakuba*	*santomea*	Nearly absent	Sao Tomé, Sao Tomé & Principe, 1998 (D. Lachaise)
melano-gaster	*melano-gaster*	*erecta*	*erecta*	Monogenic	La Lopé Forest, Gabon, 1998 (D. Lachaise)
montium	*kikkawai*	*kikkawai*	*kikkawai*	Monogenic	Chandigarh, India (Gibert et al. 1999)
montium	*seguyi*	*jambulina*	*jambulina*	Monogenic	New Delhi, India, 2014 (J. R. David, S. Mohanty)
montium	*seguyi*	*bocqueti*	*burlai*	Monogenic	Mt Oku, Cameroon, 2008 (M. Veuille)
montium	*seguyi*	*bocqueti*	cf. *bocqueti*	Monogenic	Sao Tomé, Sao Tomé & Principe, 2015 (J. R. David, M. Lang)

Pigmentation scoring

For monogenic species, we followed the nomenclature given in Gibert et al.'s (1999)

paper, referring to the light and dark alleles as *L* and *D*, respectively, and to the three genotypes

as *LL*, *LD* or *DL*, and *DD*. For heterozygotes, the allele from the mother is given first depending

on the cross. For each genotype, pigmentation was scored on >4 days old 10 males and 10

females as in Yassin et al. (2022) following the pigmentation score scale proposed by David et

al. (1990). This pigmentation score visually quantifies the proportion of the pigmented part of

an abdominal segment when viewed on the lateral side of a fly under a stereoscope and it goes

from 0 (no pigmentation) to 10 (fully pigmented). It is therefore different from other

pigmentation scores in *Drosophila* that depend on the intensity of melanism or the length of the

pigmented stripe quantified after numerical photographing (e.g., Bastide et al. 2014; Lafuente

et al. 2021). Scorings were made by the same observer (the first author) between 2013 and 2019

(but mostly between March 2016 and April 2017) to limit observation bias. Data were manually

entered on a printed sheet with a template table for morphometric data and then the means for

each segment were calculated in a separate sheet. All sheets are archived. Scorings were made

by the same observer to limit observation bias between 2013 and 2019 (but mostly between

March 2016 and April 2017). Abdomens of some flies at extreme temperatures (see below)

were dissected and mounted on microscopic slides and photomicrographed (cf. data associated

with this study). In each case, pigmentation was scored on six and five abdominal segments in

females and males respectively (denoted A2-7 with F and M for females and males,

respectively).

Flies were grown at five temperatures going from 12°C to 31°C with 3-4° increments.

For most cases, the five categories were 14, 17, 21, 25, and 28°C, but a few cases were scored

at 19 (*D. jambulina DD*), 29 (*D.* cf. *bocqueti DL*), and 30°C (*D. erecta DD*, *DL*, *LD*, and *LL*).

Mortality during development was not assessed. For *D. melanogaster* and *D. kikkawai*, only

mean values per genotype were retrieved from the literature for seven (12, 14, 17, 21, 25, 28,

and 31°C) and four (12, 17, 25, and 30°C) temperature degrees, respectively (Gibert et al. 1999,

2000). For three genotypes < 10 flies per sex were available at the time of observation (namely,

D. cf. *bocqueti LL* males at 21°C ($N = 2$), *D. jambulina DD* females and males at 25°C ($N = 5$).

Consequently, temperature was coded either as a degree or as a class depending on the statistical

test used (see below), with the five classes being: cold (12-14°C), mildly cold (17-19°C), mild

(21°C), mildly hot (25°C), and hot (28-31°C). In total, pigmentation was scored for nearly 8,703

abdominal segments over 1,582 flies for the four species with monogenic melanism. For the

hybridization experiment between *D. santomea* and *D. yakuba*, only the sheet recording the

means for each genotype per sex per temperature was retrieved for the five temperatures.

However, photomicrographs of abdominal segments for the four genotypes (parental strains

and reciprocal crosses) at three temperatures (14, 21, and 28°C) are provided in the data

associated with this study. Analyses on the whole set of temperatures for this experiment were

therefore limited to the mean values.

Statistical analyses

All statistical analyses were conducted using R (R Core Team 2023). As a serially

homologous trait, abdominal pigmentation can be tested either as in an integrative (*i.e.* the

overall pigment production) or a modular way (*i.e.* pigmentation patterns across segments).

Indeed, pigmentation in all segments is produced by the same melanin synthesis genes and their

variants that each individual carries. Previous works in *D. melanogaster* and *D. kikkawai* have

shown a substantial degree of correlation especially among adjacent segments (David et al.

1990; Gibert et al. 1999, 2000). However, pigmentation variation associates with different

genetic variants in different abdominal segment in *D. melanogaster* measured on the same

isogenic lines (Dembeck et al. 2015), and the reaction norms of the thermal plasticity are

different even among adjacent abdominal segments in *D. simulans* (Gibert et al. 1998). To test

for the independence between the abdominal segments we conducted correlation analyses on

the whole individual-based dataset using the Corrplot package in R (Wei and Simko 2024), and

found indeed substantial degrees of correlations which were stronger among adjacent segments

(see Results). Consequently, we conducted our further analyses either on each segment

separately or using the sum of all segments per an individual depending on the test.

We first conducted a multi-factor Analysis of Variance (ANOVA) on the individual-

based dataset for the four species with monogenic melanism to test of the effect of each factor

(species, genotype, sex, segment, and temperature) as well as of their interactions on

pigmentation variation. We estimated the effect of these factors as the proportion of variance

explained (η^2) using the effectsize package (Ben-Shachar et al. 2020) in R. To test for the effect

of reciprocal crosses, we conducted the same analysis only on the *DL* and *LD* genotypes. Because no significant difference was found between these genotypes, they were pooled into a single F_1 heterozygous category for subsequent analyses. For a balanced ANOVA, equal number of segments were used for each sex. To do so in *D. kikkawai*, Gibert et al. (1999) excluded the most posterior female abdominal segment, *i.e.* FA7. However, subsequent analyses showed that this segment is among the most variable within and between Drosophila species (Bastide et al. 2013; Yassin et al. 2016a,b). Consequently, we excluded here the most anterior female segment, *i.e.* A2, which usually shows less variation and strong correlation with other anterior segments (see Results). Similarly, temperature classes rather than temperature degrees were used to test the effect of temperature in a balanced way across species and genotypes.

Previous analyses have shown the shape of the thermal reaction norms to differ between segments of the same isofemale lines. For example, segments 5, 6, and 7 in *D. simulans* females have quadratic, linear, and sigmoid (logistic) norms, respectively, although all show the same response of higher melanism in colder environments (Gibert et al. 1998). Because our main focus in this paper is to compare the level of plasticity between heterozygotes and homozygotes rather than the diversity of the shape of the reaction norms, linear regressions of pigmentation scores over temperature degrees were conducted using the lm function in R for every segment as well as for the sum of all segments, and the slope of the regression line (*b*) was considered as a proxy for the degree of plasticity. Regression analyses were conducted on the whole individual-based dataset rather than on averages for the measurements generated by this study, although for the data retrieved from the literature for *D. melanogaster* and *D. kikkawai*, and for the hybridization experiment, only averages were available.

To visualize the patterns of segment-specific diversity for each genotype per sex per species we used a customized perl script to translate the linear regression equations generated

above into a pigmentation pattern. Each segment was divided into a grid of 100 centiles so that

a pigmentation score of 2.2 for a segment means that the 22 most posterior centiles of that

segment have a pigmentation score of 10, whereas the 78 most anterior centiles have a score of

0. A 2D contour map was then drawn using the Plotly package in R (Sievert 2020). For the

overall pigmentation score (*i.e.* sum of all segments), 2D plots were drawn for each sex per

species using the plot function in R. Genetic dominance was inferred as the difference between

the slope of pooled heterozygotes (*DL* and *LD*) and the slope of the pooled homozygotes (*DD*

and *LL*), with dominance reversal being witnessed at the intersection between the two slopes

within the range of vital temperatures between 10 and 30°C. To test if heterozygotes had

different plasticity than homozygotes, we ranked the slope of each of the three genotypes from

the highest to the lowest as 3, 2, and 1 for each species and conducted a non-parametric

Wilcoxon test on the relation between rank and genotype.

For the hybridization experiment between *D. santomea* and *D. yakuba*, only mean scores

for each genotype per sex per temperature were available (see above). We first conducted a

multi-factor ANOVA on these values, after excluding female A2 and considering the five

temperature classes as for the species with the monogenic melanism. We also conducted

ANOVA on only the hybrid genotypes to test for the effect of reciprocal crosses on the whole

dataset. To quantify and visualize plasticity at each segment as well as on the total pigmentation

scores we conducted linear regression analyses on the mean values as above. We note that for

species with monogenic melanism, regression slopes strongly correlated between the whole

datasets and the means ($r = 0.997$, $P < 2.2 \times 10^{-16}$; Student's t $P = 0.871$), although the standard

error of the slope value was significantly reduced in individual-based analyses ($r = 0.849$, $P <$

2.2×10^{-16}; Student's t $P = 1.671 \times 10^{21}$). Consequently, the slope reported for the mean values

of this experiment correctly reflects the plasticity response.

Results

Pigmentation thermal plasticity varies with species, sex, segment, and genotype in species with monogenic melanism

ANOVA on the whole dataset of the four species with monogenic melanism indicated a strong effect of all factors and their interactions (Table 2). However, the most significant effect was due to segment position (explaining 30% of the variance), followed by temperature and genotype (5%), then species (2%) and sex (1%). To test for the effect of reciprocal crosses, we conducted the same ANOVA on the data set of only heterozygous genotypes (*LD* and *DL*). We found that the genotype had no effect at all, whether individually or in interaction with other factors. For simplicity, we pooled data for heterozygous individuals issued from reciprocal crosses into a single genotypic group (F_1). When we conduct the analysis using the complete genotypic dataset (*i.e.* with pooled *LD* and *DL*), the ANOVA results showed the significant effect of all factors and their interactions.

Table 2. Analysis of variance (ANOVA) for pigmentation variation in four species with monogenic melanism. *D* and *L* refers to dark and light alleles, respectively. Df = degree of freedom, GEN = genotype, SEG = segment position, SPE = species, and TCL = temperature class. *** $P < 0.001$.

Genotypes	*DD, DL, LD, LL*			*DL, LD*			*DD, F₁, LL*		
Factor	Df	F	η^2	Df	F	Eta2	Df	F	η^2
SPE	3	222.438***	0.02	3	104.487***	0.02	3	222.897***	0.02
GEN	3	546.944***	0.05	1	0.433	<0.01	2	821.871***	0.05
TCL	1	1459.131***	0.05	1	1183.554***	0.08	1	1462.142***	0.05
SEX	1	337.125***	0.01	1	87.979***	<0.01	1	337.820***	0.01
SEG	1	9325.489***	0.30	1	5121.901***	0.36	1	9344.735***	0.30
SPE:GEN	9	21.791***	<0.01	3	0.328	<0.01	6	32.573***	<0.01
SPE:TCL	3	70.243***	<0.01	3	79.879***	0.02	3	70.388***	<0.01
GEN:TCL	3	55.130***	<0.01	1	3.112	<0.01	2	81.154***	<0.01
SPE:SEX	3	156.685***	0.02	3	115.028***	0.02	3	157.008***	0.02

	df	value	p	df	value	p	df	value	p
GEN:SEX	3	306.050***	0.03	1	0.107	<0.01	2	459.963***	0.03
TCL:SEX	1	96.736***	<0.01	1	178.999***	0.01	1	96.936***	<0.01
SPE:SEG	3	736.924***	0.07	3	503.505***	0.11	3	738.445***	0.07
GEN:SEG	3	518.281***	0.05	1	0.781	<0.01	2	778.596***	0.05
TCL:SEG	1	45.414***	<0.01	1	119.222***	<0.01	1	45.508***	<0.01
SEX:SEG	1	1118.545***	0.04	1	308.347***	0.02	1	1120.853***	0.04
SPE:GEN:TCL	9	11.695***	<0.01	3	1.447	<0.01	6	16.783***	<0.01
SPE:GEN:SEX	9	13.581***	<0.01	3	0.138	<0.01	6	20.337***	<0.01
SPE:TCL:SEX	3	20.594***	<0.01	3	32.869***	<0.01	3	20.636***	<0.01
GEN:TCL:SEX	3	34.231***	<0.01	1	0.056	<0.01	2	51.421***	<0.01
SPE:GEN:SEG	9	22.254***	<0.01	3	0.051	<0.01	6	33.421***	<0.01
SPE:TCL:SEG	3	10.443***	<0.01	3	26.690***	<0.01	3	10.464***	<0.01
GEN:TCL:SEG	3	30.370***	<0.01	1	0.079	<0.01	2	45.606***	<0.01
SPE:SEX:SEG	3	55.643***	<0.01	3	96.490***	0.02	3	55.758***	<0.01
GEN:SEX:SEG	3	347.385***	0.03	1	1.411	<0.01	2	521.377***	0.03
TCL:SEX:SEG	1	115.968***	<0.01	1	227.193***	0.02	1	116.208***	<0.01
SPE:GEN:TCL:SEX	9	6.391***	<0.01	3	0.085	<0.01	6	9.559***	<0.01
SPE:GEN:TCL:SEG	9	9.518***	<0.01	3	0.412	<0.01	6	14.079***	<0.01
SPE:GEN:SEX:SEG	9	20.399***	<0.01	3	0.212	<0.01	6	30.545***	<0.01
SPE:TCL:SEX:SEG	3	25.856***	<0.01	3	49.527***	0.01	3	25.910***	<0.01
GEN:TCL:SEX:SEG	3	45.620***	<0.01	1	0.207	<0.01	2	68.457***	<0.01
SPE:GEN:TCL:SEX:SEG	9	11.697***	<0.01	3	0.097	<0.01	6	17.529***	<0.01
Residuals	7780			3936			7812		

Higher plasticity of overall pigmentation in heterozygous genotypes in species with monogenic

melanism

Despite the strong effect of body segmentation on pigmentation diversity between species, sexes, and genotypes, previous studies have reported strong correlation in pigmentation between adjacent segments (David et al. 1990; Gibert et al. 1999, 2000). We tested this hypothesis using individual data for the four species and indeed found substantial degrees of correlations, especially among adjacent segments, indicating the non-independence of segment-

based effect (Figure 2A). Consequently, to exclude such non-independence, we summarized

pigmentation variation as sum scores across segments per individual. A multi-factor ANOVA

on the sum pigmentation scores showed a significant effect of all factors and their interactions,

with the strongest effect being of temperature (explaining 21% of the variance), followed by

genotype (18%) and species (11%). To visualize the effect of these three factors, we plotted

sum pigmentation scores of each genotype per species over temperature. In all cases, a

significant negative linear relation was found, with pigmentation increases in colder

environments (Figure 2B). Adjusted R^2 ranged from 0.339 to 0.899, indicating that some

relationships may better be explained by more complex non-linear regressions. However, for

the sake of comparisons across species and genotypes, we used the slope of the linear regression

of each genotype as a proxy for the degree of phenotypic plasticity. Species greatly differed in

these slopes (averaged over three genotypes and two sexes), being highest in *D. jambulina*

(mean *b*=-1.189±0.617), followed by *D.* cf. *bocqueti* (*b*=-0.781±0.329), *D. burlai* (*b*=-

0.470±0.010), and *D. erecta* (*b*=-0.441±0.080). For each species, we ranked the three genotypes

from the most plastic (*i.e.* strongest slope) to the least plastic, and found that heterozygosity

strongly correlated with higher plasticity (Wilcoxon's $W = 16$, $P = 0.002$).

Figure 2 – Correlation between abdominal segments and plasticity of total pigmentation in species with monogenic melanism. A) Correlation between pigmentation scores of abdominal segment in females (left) and males (right) showing stronger correlation between adjacent segments. B) Linear regression of total pigmentation (sum of segment scores) over developmental temperature in homozygous dark (*DD*) and light (*LL*) genotypes and pooled heterozygotes (*DD* and *LL*). Dominance can be inferred as the difference between the slope of the pooled heterozygotes (black dashed) and that of the pooled homozygotes (red solid).

 Higher plasticity of heterozygotes associates with dominance reversal

The stronger slope of the heterozygote relative to the homozygotes indicate that

dominance between the dark *D* and light *L* alleles likely changes with developmental

temperature. To evaluate this effect, we inferred the slope of pooled homozygous genotypes,

which is a proxy of the mid-parent values across temperatures. We found indeed that the slope

of the heterozygotes approaches this mid-parent slope in almost all cases at higher temperature,

indicating a reduction of the dominance of the *D* allele with increasing temperature. The

heterozygotes' slope intercrossed that of the mid-parent at 17°C in *D.* cf. *bocqueti* males, 19°C

in *D.* cf. *bocqueti* females, 20°C in *D. jambulina* females, 25°C in *D. jambulina* males and 30°C

in *D. burlai* males, indicating a reversal of dominance within the vital temperature range

between 10 and 30°C. In two cases, *D. erecta* and *D. burlai* females the intercrossing was

expected at >30°C, *i.e.* the *D* allele is dominant throughout the vital range although with

decreasing dominance. In *D. erecta* males, the slope superimposed that of the mid-parent

throughout the temperature range, indicating a non-plastic codominance of the *D* and *L* alleles.

Visual depiction of the regression slope at each segment showed differences between the

segments especially towards the posterior of the abdomen (Figures 3 and 4). The reversal of

dominance in *Drosophila jambulina* is interesting, since the dominance relationship in this

species is ambiguous. Parkash and Sharma (1978) found dark dominance for an Indian

population, whereas Watanabe et al. (1982) found light dominance also for Indian populations

and concluded that dominance in *D. jambulina* was "a great problem existing in a simple genetic

phenomenon". Our results suggest that Parkash and Sharma (1978) and Watanabe et al. (1982)

might have conducted their experiments at different temperatures, hence reaching contradictory

results. For *D.* cf. *bocqueti*, Prigent et al. (2020) reported that the light allele was dominant in

the Sao Tomé strains that we analyze here. Our results indicate that this statement is

temperature-dependent. However, because both *D. jambulina* and *D.* cf. *bocqueti*, are tropical,

predominantly living in environments with an average temperature ≥21°C, the light allele may

be considered the dominant one in nature.

Figure 3 – Thermal phenotypic plasticity of pigmentation on successive abdominal segments
in *D. erecta* (A) and *D. jambulina* (B). For each panel, contour plots show pigmentation thermal
plasticity (proportion of black stripe) in females (above) and males (below) for the three
genotypes *DD*, *DL*, and *LL*. For each abdominal segment, a reaction norm was inferred as a
linear regression slope using individual values.

Figure 4 – Thermal phenotypic plasticity of pigmentation on successive abdominal segments in *D. burlai* (A) and *D.* cf. *bocqueti* (B). For each panel, contour plots show pigmentation thermal plasticity (proportion of black stripe) in females (above) and males (below) for the three genotypes *DD*, *DL*, and *LL*. For each abdominal segment, a reaction norm was inferred as a linear regression slope using individual values.

 Plasticity increases in hybrids of D. santomea *and* D. yakuba

Multi-factor ANOVA showed all factors but sex to have a significant effect on

abdominal pigmentation in *D. santomea*, *D. yakuba*, and their hybrids, with the genotype being

the most important factor (accounting for nearly 39% of the variance), followed by the segment

position (25%), and temperature (4%) (Table 3). Indeed, the two parental species are known to

show very little sexual dimorphism, but very different pigmentation especially on the posterior

segments, with *D. yakuba* and *D. santomea* being nearly completely dark and light, respectively

(David et al. 2022; Yassin et al. 2022). Contrary to species with monogenic melanism, multi-

factor ANOVA on the heterozygotes from the reciprocal crosses showed a significant effect of

the genotype (Table 3).

Table 3. Analysis of variance (ANOVA) for pigmentation variation in *D. santomea* (*LL*), *D.*

yakuba (*DD*), and their hybrids. *D* and *L* refers to dark (*D. yakuba*) and light (*D. santomea*)

alleles, respectively. Df = degree of freedom, GEN = genotype, SEG = segment position, and

TCL = temperature class. ** $P < 0.01$; *** $P < 0.001$.

Experiment	DD, DL, LD, LL			DL, LD		
Factor	Df	F	η^2	Df	F	η^2
GEN	3	230.863***	0.39	1	43.272***	0.08
TCL	1	64.244***	0.04	1	65.130***	0.12
SEX	1	2.511	1.41e-03	1	3.352.	5.97e-03
SEG	1	450.899***	0.25	1	238.814***	0.43
GEN:TCL	3	7.945***	0.01	1	0.132	2.35e-04
GEN:SEX	3	10.601***	0.02	1	23.579***	0.04
TCL:SEX	1	0.001	3.04e-07	1	0.321	5.71e-04
GEN:SEG	3	73.875***	0.12	1	16.177***	0.03
TCL:SEG	1	22.129***	0.01	1	35.368***	0.06
SEX:SEG	1	13.212***	7.42e-03	1	14.030***	0.02
GEN:TCL:SEX	3	0.683	1.15e-03	1	0.970	1.73e-03
GEN:TCL:SEG	3	7.495***	0.01	1	0.835	1.49e-03
GEN:SEX:SEG	3	5.626**	9.48e-03	1	9.489**	0.02
TCL:SEX:SEG	1	2.662	1.50e-03	1	3.439.	6.13e-03
GEN:TCL :SEX:SEG	3	1.686	2.84e-03	1	2.550	4.54e-03
Residuals	208			104		

Total pigmentation (summed over all segments) showed significant thermal plasticity in

both sexes in *D. yakuba* (females: b = -0.365 ± 0.039, P < 0.001; males: b = -0.513 ± 0.057, P

< 0.001) but only and to a lesser degree in *D. santomea* females (females: b = -0.093 ± 0.020,

P < 0.01 males: b = -0.145 ± 0.057, P = 0.065) (Figure 5). The slope of the linear regression

was higher in hybrids, approaching -1 in females (*DL*: b = -1.277 ± 0.155, P < 0.01; *LD*: b = -

0.941 ± 0.092, P < 0.001) and males (*DL*: b = -0.866 ± 0.161, P < 0.01; *LD*: b = -1.020 ± 0.096,

P < 0.001) (Figure 5). As for species with monogenic melanism, the slope of the heterozygotes

approached or crossed that of the mid-parents indicating a reversal of dominance. In females of

reciprocal crosses, the intercrossing occurred at intermediated temperatures (17-20°C). In

males, on the other hand, the slope of the heterozygotes approached that of the pooled

homozygotes but in opposite sides. For the *san x yak (LD)* males, the *santomea* light phenotype

was dominant throughout the range, approaching codominance at cold temperatures. For the

yak x san (DL) males, the opposite was found with the *yakuba* dark phenotype was dominant

throughout the range approaching codominance at hot temperatures. This indicates a strong

effect of the X chromosome, which carries two melanin-synthesis genes with loss-of-function

mutations in *D. santomea* (Liu et al. 2019; David et al. 2022; see Discussion).

Figure 5 – Thermal phenotypic plasticity of abdominal pigmentation in *D. yakuba*, *D. santomea*, and their hybrids. A) Linear regression of total pigmentation (sum of segment scores) over developmental temperature in homozygous dark (*DD*) and light (*LL*) genotypes and pooled heterozygotes (*DD* and *LL*). Dominance can be inferred as the difference between the slope of the pooled heterozygotes (black dashed) and that of the pooled homozygotes (red solid). B) Plasticity patterns at each abdominal segment. For each panel, contour plots show pigmentation thermal plasticity (proportion of black stripe) in females (above) and males (below) for the three genotypes *DD*, *DL*, and *LL*. For each abdominal segment, a reaction norm was inferred as a linear regression slope using mean values (see text).

As indicated from the ANOVA analyses, segment position had very important effect on

pigmentation, especially in comparing reciprocal crosses where it explained 43% of the

variance (Table 3). This is perfectly seen in the visual depiction for each segment, where

differences in pigmentation plasticity were more pronounced in posterior segments (Figure 5).

Discussion

The molecular basis of the association between plasticity and heterozygosity in Drosophila

Although it has long been recognized that genetic variation of plasticity exists, the nature

of this variation remains elusive. Early works suggested three non-mutually exclusive

mechanisms, namely overdominance, pleiotropy, and epistasis (Scheiner 1993). Of the three

models, only overdominance made an explicit prediction about the relationship between

plasticity and heterozygosity. The lack of strong evidence that heterozygous genotypes were

more robust has led to the dismissal of the overdominance model, and current studies on the

molecular basis of plasticity either consider the epigenetic and transcriptomic landscapes (*i.e.*

the pleiotropy model) or the shape of regulatory networks (*i.e.* the epistasis model) (Goldstein

and Ehrenreich 2021; Chevin et al. 2022). Our analysis of abdominal pigmentation in multiple

Drosophila species supports an association between plasticity and heterozygosity but in a

positive rather than negative way, hence revisiting the classical overdominance model.

However, as indicated by Scheiner (1993), the three models are not mutually-exclusive, and we

will illustrate how our results can agree with the different models depending on the species and

the sex considered.

In its simplest form, a gene regulatory network consists of a transcription factor

activating or suppressing an effector gene. Of the five monogenic cases studied here, the

underlying locus involves an effector gene, *t*, in one species (*D. erecta*) (Yassin et al. 2016a),

and a transcription factor, *pdm3*, in four species (*D. kikkawai*, *D. jambulina*, *D. burlai* and *D.*

cf. *bocqueti*) (Yassin et al. 2016b; Fukutomi et al. 2025) (Figure 6). Loss-of-function regulatory

mutations of *t* reduce plasticity in *D. erecta* recessive light homozygotes (Figure 2B), and only

slightly affect the degree of dominance, *i.e.* plasticity of the functional allele is nearly

codominant near the upper thermal limit when both dark heterozygotes and homozygotes

become lighter (Figure 2B). Indeed, the *t* regulatory element which is involved in female color

variation in both *D. erecta* and *D. melanogaster* (Bastide et al. 2013; Endler et al. 2016; Gibert

et al. 2017a) partly affects pigmentation plasticity in the later species (Gibert et al. 2016). On

the other hand, mutations in the transcription factor *pdm3* show more complex relationships

(Figures 2B). First, the plasticity of both homozygotes shows almost similar slopes (but

different means). Second, dominance relationship differed between species, even among sister

species such as *D. burlai* and *D.* cf. *bocqueti*. In two species (*D. kikkawai* and *D. burlai*), the

dark allele is dominant throughout the temperature range, whereas in the two other species (*D.*

jambulina and *D.* cf. *bocqueti*), a strong dominance reversal was observed in females.

Polymorphism in *pdm3* associates with female abdominal pigmentation variation in European

and African populations of *D. melanogaster* (Endler et al. 2016). Mutations in this gene also

underlie pigmentation evolution between the sister species *D. yakuba* and *D. santomea* with

contrasting melanism (Liu et al. 2019; David et al. 2022). Whereas pigmentation plasticity in

these two species is low, their reciprocal hybrids, which are heterozygous for *pdm3* interspecific

differences along with other genes, are highly plastic (Figure 2A; Figure 5B). This may indicate

that the relation between heterozygosity and plasticity may differ according to the type of the

underlying gene.

	Phenotypic variation	Genetic polymorphism	Thermal plasticity
melanogaster group			
melanogaster	Continuous	E: *t, e* TF: *bab, pdm3*	High
simulans	Continuous	Unknown	High
yakuba	No variation	E: *y, t, e* TF: *Abd-B,* *pdm3, Gug*	Low Hybrid high
santomea	No variation		Low
erecta	Discrete, dark dominant	E: *t*	Low
suzukii	Continuous	Unknown	High
kikkawai	Discrete, dark dominant	E: *y* TF: *pdm3*	Low
jambulina	Discrete, light dominant	TF: *pdm3*	Heterozygous high Homozygous low
burlai	Discrete, dark dominant	TF: *pdm3*	Low
cf. *bocqueti*	Discrete, light dominant	TF: *pdm3*	Heterozygous high Homozygous low
montium group			

507 **Figure 6** – Evolution of the genetic architecture of pigmentation variation and its thermal
508 plasticity in the *melanogaster* and *montium* species groups. E = effector melanin-synthesis
509 gene; TF = transcription factor. Phylogenetic relationships are after Suvorov et al. (2022).
510

Besides, the types of the mutations in the alleles of the polymorphic gene may also affect

the degree of the plastic response. Indeed, plasticity was highest in the heterozygotes of the two

montium group species, *D. jambulina* and *D.* cf. *bocqueti*, where the light allele was dominant

under optimal developmental temperature regimes, *i.e.* ≥21°C. The reversal of dominance

according to the temperature, the main reason of the steepness of the reaction norm, is

associated with the evolution of dominance between the *pdm3* alleles in these species. In *D.*

melanogaster, the wildtype light allele is dominant (Tichy et al. 2008), but RNAi lines targeting

different parts of this gene yields contrasting results, either increasing pigmentation (mostly in

females) or decreasing it (mostly in males) (Rogers et al. 2014). Consequently, it is possible

that the *pdm3* mutations underlying the dark allele in *montium* species where it is dominant,

such as in *D. kikkawai* and *D. burlai*, are of the dominant negative type, which are common for

transcription factors (Gerasimavicius et al. 2022). Such transcription factors usually have

multiple isoforms, one of them has the DNA binding domain but lacks polymerase or other

protein binding domains that are needed to express a gene, whereas another isoform has both

DNA and protein binding domains. Consequently, the isoforms have opposite effects on the

trait. In the heterozygote, the isoform lacking the protein binding domain competes with the

isoform bearing this domain, hence altering its function. On the other hand, mutations in species

with the light allele being dominant, like *D. jambulina* and *D. cf. bocqueti*, may be of the

classical loss-of-function type that may reduce but do not completely eliminate the effect of the

wild-type light allele. Further molecular dissection of the underlying alleles is required to

understand the evolution of dominance reversal at *pdm3*.

532   The effect of mutations in both modifier and effector genes, *i.e.* the epistatic model of

plasticity, is best illustrated in *D. kikkawai* and the hybrids between *D. santomea* and *D. yakuba*.

In *D. kikkawai*, the effector *yellow (y)* gene lacks a conserved *cis*-regulatory element leading to

the loss of body pigmentation (Jeong et al. 2008). Because this gene is X-linked, a dosage effect

may occur between the sexes. Plasticity in males additively increased with the number of copies

of the dominant dark *pdm3* allele, which is an indirect regulator of *y* (Liu et al. 2019). Putatively

negative dominant mutations in this gene are expected to activate *y*, but the effect may depend

on the number of copies of *y*, being higher in females. Similarly, whereas hybrid males

mothered by *D. santomea*, *i.e.* those carrying defective X-linked melanin synthesis genes *t* and

y, had similar plasticity slopes to hybrid females. But this slope was halved in hybrid males

mothered by *D. yakuba*, with functional *y* and *t*. Therefore, plasticity is affected by the

functionality of the polymorphism at multiple genes, as predicted by the epistasis model,

although the degree of plasticity was higher in fully heterozygous background.

The association between plasticity and polymorphism may be more general than currently

appreciated

We have restricted our analyses to eight species belonging to two species groups of the

subgenus *Sophophora*. Despite the differences in their reaction norms, both groups contain

species with sexually dimorphic melanism, and a single gene with antagonistic sexual effect

may underlie the association between plasticity and heterozygosity, hence reducing the

independence of our observations. Future analyses should investigate pigmentation thermal

plasticity in non-*Sophophora* fly species lacking sexual dimorphism, with or without Mendelian

polymorphism (e.g., Brisson et al. 2005; Rocha et al. 2009; Wittkopp et al. 2009). Beyond

Drosophila, color polymorphism abounds in nature (Orteu and Jiggins 2020; Sapir et al. 2021).

Investigation of the diversity in reaction norms in species with heterozygous and homozygous

color genotypes, especially in these with a hierarchy of dominance (e.g., Durand et al. 2014; Le

Poul et al. 2014), should shed light on the generality of the relationship between heterozygosity

and the degree of plasticity.

Most quantitative traits for which phenotypic plasticity was studied in *Drosophila*, such

as body size, wing shape, bristle number, ovariole number or life-history traits, are highly

polygenic with often a redundant genetic basis (David et al. 2004; Gibert et al. 2004a; Mackay

and Lyman 2005; Lobell et al. 2017; Lafuente et al. 2018; Pitchers et al. 2019). Identifying the

underlying genotypes of these traits to test for the effect of heterozygosity may be difficult. An

alternative to the 'forward genetics' approach that we leveraged here, is to use a 'reverse

genetics' analysis, wherein the transcription level of polymorphic genes can be analyzed in

homozygous vs. heterozygous states in different environments, even in the lack of knowledge

about the phenotypic consequences of the genotypes. One such significant example comes from

a study by Chen et al. (2015) who showed that 62% of genes with thermal plastic response in

D. melanogaster (*i.e.* ~51% of total genes) swapped allelic dominance in heterozygotes

between cold and hot environments. Future similar transcriptomic analyses are therefore highly

needed to test the generality of these patterns, especially these using hybrids between species.

The evolutionary consequences of links between phenotypic plasticity and polymorphism

A link between phenotypic plasticity and polymorphism can provide simple

explanations on how plasticity evolves. In fact, plasticity can be lost through any mechanism

reducing genetic heterozygosity such as inbreeding or directional selection. Indeed, the 'flexible

stem hypothesis' postulates that from a plastic ancestor, descendants with fixed phenotypes

from the ancestral phenotypic spectrum can evolve (West-Eberhard 2003; Gibert 2017). A

likely example from our data is the case of *D. santomea* and *D. yakuba*. According to the

phylogeny given in Figure 6, it is likely that the ancestor of these species was highly plastic

given the high plasticity of *D. melanogaster* (Gibert et al. 2000) and *D. suzukii* (Colinet and

Kustre 2025). Pigmentation difference between *D. yakuba* and *D. santomea* is due to mutations

in at least six loci (Liu et al. 2019; David et al. 2022), but it is unlikely that all these mutations

have arisen simultaneously. This means that a substantial degree of heterozygosity segregated

in the ancestor of these species, and consequently this ancestor has gradually lost its plasticity

as more and more alleles became fixed in each lineage. Indeed, pigmentation plays a role in the

reproductive isolation between these two species (David et al. 2022), and a hybrid zone on the

African island of Sao Tomé between these two species exists (Lachaise et al. 2000; Llopart et

al. 2005; Turissini and Matute 2017). A plastic hybrid is disadvantaged and the 'flexible stem

hypothesis' may in essence reflects a simple model of 'underdominance' of the heterozygote.

Conversely, phenotypic plasticity can be gained by any mechanism that increases

genetic variability and heterozygosity in a population, such as mutations, outbreeding (e.g.,

gene flow and introgression), and balancing selection. Indeed, any new mutation in a di- or

polyploid organism first arises in the heterozygous state, and it has been argued that a

heterozygous advantage is essential for the spread of new mutations and counteracting the loss

by random genetic drift (Sellis et al. 2011). In cases where plasticity is adaptive, the higher

plasticity of the heterozygote can facilitate the spread of this mutation. Such adaptive plasticity

gain model is in essence an 'overdominance' model. An important consequence of the

overdominance model is that plasticity could be a mechanism to maintain genetic diversity, *i.e.*

rather than 'buying time' for genetic adaptation to occur, plasticity may be a major source for

'saving time' by maintaining standing genetic variation, such as in the case of introduced

species or multivoltine species inhabiting a heterogeneous spatial or seasonal environment

(Gulisija et al. 2016; Wittmann et al. 2017; Promy et al. 2023). Further theoretical and empirical

investigations of the link between heterozygosity and plasticity will definitively improve our

understanding of the evolution of both.

References

Bastide, H., A. Betancourt, V. Nolte, R. Tobler, P. Stöbe, A. Futschik, and C. Schlötterer.

2013. A Genome-Wide, Fine-Scale Map of Natural Pigmentation Variation in Drosophila

melanogaster. PLOS Genetics 9:e1003534. Public Library of Science.

Bastide, H., J. D. Lange, J. B. Lack, A. Yassin, and J. E. Pool. 2016. A Variable Genetic

Architecture of Melanic Evolution in Drosophila melanogaster. Genetics 204:1307–1319.

Bastide, H., A. Yassin, E. J. Johanning, and J. E. Pool. 2014. Pigmentation in Drosophila

melanogaster reaches its maximum in Ethiopia and correlates most strongly with ultra-violet

radiation in sub-Saharan Africa. BMC Evolutionary Biology 14:179.

Ben-Shachar, M. S., D. Lüdecke, and D. Makowski. 2020. effectsize: Estimation of Effect

Size Indices and Standardized Parameters. Journal of Open Source Software 5:2815.

Berardi, S., J. A. Rhodes, M. C. Berner, S. I. Greenblum, M. C. Bitter, E. L. Behrman, N. J.

Betancourt, A. O. Bergland, D. A. Petrov, S. Rajpurohit, and P. Schmidt. 2024. Drosophila

melanogaster pigmentation demonstrates adaptive phenotypic parallelism but genomic

unpredictability over multiple timescales. bioRxiv.

Bradshaw, A. D. 1965. Evolutionary Significance of Phenotypic Plasticity in Plants. Pp. 115–

155 *in* E. W. Caspari and J. M. Thoday, eds. Advances in Genetics. Academic Press.

Bradshaw, A. D. 2006. Unravelling phenotypic plasticity – why should we bother? New

Phytologist 170:644–648.

Brisson, J. A., D. C. D. Toni, I. Duncan, and A. R. Templeton. 2005. Abdominal

Pigmentation Variation in Drosophila Polymorpha: Geographic Variation in the Trait, and

Underlying Phylogeography. Evolution 59:1046–1059.

Castro, S. D., F. Peronnet, J.-F. Gilles, E. Mouchel-Vielh, and J.-M. Gibert. 2018. bric à brac

(bab), a central player in the gene regulatory network that mediates thermal plasticity of

pigmentation in Drosophila melanogaster. PLOS Genetics 14:e1007573. Public Library of

Science.

Chakir, M., A. Chafik, P. Gibert, and J. R. David. 2002. Phenotypic plasticity of adult size

and pigmentation in *Drosophila*: thermosensitive periods during development in two sibling

species. Journal of Thermal Biology 27:61–70.

Chen, J., V. Nolte, and C. Schlötterer. 2015. Temperature Stress Mediates Decanalization and

Dominance of Gene Expression in Drosophila melanogaster. PLOS Genetics 11:e1004883.

Public Library of Science.

Chevin, L.-M., C. Leung, A. Le Rouzic, and T. Uller. 2022. Using phenotypic plasticity to

understand the structure and evolution of the genotype–phenotype map. Genetica 150:209–

221.

Colinet, H., and A. Kustre. 2025. The apparent seasonal biphenism in stems in reality from

continuous reaction norms. Pest Management Science 81:507–517.

David, J. R., R. Allemand, P. Capy, M. Chakir, P. Gibert, G. Pétavy, and B. Moreteau. 2004.

Comparative life histories and ecophysiology of Drosophila melanogaster and D. simulans.

Genetica 120:151–163.

David, J. R., P. Capy, and J.-P. Gauthier. 1990. Abdominal pigmentation and growth

temperature in Drosophila melanogaster: Similarities and differences in the norms of reaction

of successive segments. Journal of Evolutionary Biology 3:429–445.

David, J. R., E. A. Ferreira, L. Jabaud, D. Ogereau, H. Bastide, and A. Yassin. 2022.

Evolution of assortative mating following selective introgression of pigmentation genes

between two Drosophila species. Ecology and Evolution 12:e8821.

Dembeck, L. M., W. Huang, M. M. Magwire, F. Lawrence, R. F. Lyman, and T. F. C.

Mackay. 2015. Genetic Architecture of Abdominal Pigmentation in Drosophila melanogaster.

PLOS Genetics 11:e1005163. Public Library of Science.

Diamond, S. E., and R. A. Martin. 2021. Buying Time: Plasticity and Population Persistence.

P. *in* Phenotypic Plasticity & Evolution. CRC Press.

Durand, E., R. Méheust, M. Soucaze, P. M. Goubet, S. Gallina, C. Poux, I. Fobis-Loisy, E.

Guillon, T. Gaude, A. Sarazin, M. Figeac, E. Prat, W. Marande, H. Bergès, X. Vekemans, S.

Billiard, and V. Castric. 2014. Dominance hierarchy arising from the evolution of a complex

small RNA regulatory network. Science 346:1200–1205. American Association for the

Advancement of Science.

Endler, L., A. J. Betancourt, V. Nolte, and C. Schlötterer. 2016. Reconciling differences in

pool-GWAS between populations: a case study of female abdominal pigmentation in

Drosophila melanogaster. Genetics 202:843–855. Oxford University Press.

Ford, E. B. 1945. Polymorphism. Biological Reviews 20:73–88.

Freoa, L., L.-M. Chevin, P. Christol, S. Méléard, M. Rera, A. Véber, and J.-M. Gibert. 2023.

Drosophilids with darker cuticle have higher body temperature under light. Sci Rep 13:3513.

Nature Publishing Group.

Fukutomi, Y., A. Phillips-Garcia, J. Liu, A. Chuang, M. Watada, S. Ramniwas, and A. Kopp.

2025. Evolution of dominance in a Mendelian trait is linked to the evolution of environmental

plasticity. bioRxiv, doi: https://doi.org/10.1101/2025.05.30.657093.

Gerasimavicius, L., B. J. Livesey, and J. A. Marsh. 2022. Loss-of-function, gain-of-function

and dominant-negative mutations have profoundly different effects on protein structure. Nat

Commun 13:3895. Nature Publishing Group.

Ghalambor, C. K., K. L. Hoke, E. W. Ruell, E. K. Fischer, D. N. Reznick, and K. A. Hughes.

2015. Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in

nature. Nature 525:372–375. Nature Publishing Group.

Gibert, J.-M. 2017. The flexible stem hypothesis: evidence from genetic data. Dev Genes

Evol 227:297–307.

Gibert, J.-M., J. Blanco, M. Dolezal, V. Nolte, F. Peronnet, and C. Schlötterer. 2017a. Strong

epistatic and additive effects of linked candidate SNPs for Drosophila pigmentation have

implications for analysis of genome-wide association studies results. Genome Biol 18:126.

Gibert, J.-M., E. Mouchel-Vielh, S. D. Castro, and F. Peronnet. 2016. Phenotypic Plasticity

through Transcriptional Regulation of the Evolutionary Hotspot Gene tan in Drosophila

melanogaster. PLOS Genetics 12:e1006218. Public Library of Science.

Gibert, J.-M., E. Mouchel-Vielh, and F. Peronnet. 2017b. Modulation of yellow expression

contributes to thermal plasticity of female abdominal pigmentation in Drosophila

melanogaster. Sci Rep 7:43370. Nature Publishing Group.

Gibert, P., P. Capy, A. Imasheva, B. Moreteau, J. P. Morin, G. Pétavy, and J. R. David.

2004a. Comparative Analysis of Morphological Traits Among Drosophila Melanogaster and

D. Simulans: Genetic Variability, Clines and Phenotypic Plasticity. Genetica 120:165–179.

Gibert, P., B. Moreteau, and J. R. David. 2000. Developmental constraints on an adaptive

plasticity: reaction norms of pigmentation in adult segments of Drosophila melanogaster.

Evolution & Development 2:249–260.

Gibert, P., B. Moreteau, and J. R. David. 2009. Phenotypic plasticity of abdomen

pigmentation in two geographic populations of Drosophila melanogaster: male–female

comparison and sexual dimorphism. Genetica 135:403–413.

Gibert, P., B. Moreteau, and J. R. David. 1999. Phenotypic plasticity of abdominal

pigmentation in Drosophila kikkawai: multiple interactions between a major gene, sex,

abdomen segment and growth temperature. Genetica 105:165–176.

Gibert, P., B. Moreteau, and J. R. David. 2004b. Phenotypic plasticity of body pigmentation

in Drosophila melanogaster: Genetic repeatability of quantitative parameters in two

successive generations. Heredity 92:499–507. Nature Publishing Group.

Gibert, P., B. Moreteau, J. R. David, and S. M. Scheiner. 1998. Describing the evolution of

reaction norm shape: body pigmentation in Drosophila. Evolution 52:1501–1506.

Gibert, P., B. Moreteau, J.-C. Moreteau, and J. R. David. 1996. Growth Temperature and

Adult Pigmentation in Two Drosophila Sibling Species: An Adaptive Convergence of

Reaction Norms in Sympatric Populations? Evolution 50:2346–2353.

Gillespie, J. H., and M. Turelli. 1989. Genotype-environment interactions and the

maintenance of polygenic variation. Genetics 121:129–138.

Goldstein, I., and I. M. Ehrenreich. 2021. Genetic Variation in Phenotypic Plasticity. P. *in*

Phenotypic Plasticity & Evolution. CRC Press.

Gulisija, D., Y. Kim, and J. B. Plotkin. 2016. Phenotypic Plasticity Promotes Balanced

Polymorphism in Periodic Environments by a Genomic Storage Effect. Genetics 202:1437–

1448.

Harnly, M. H., and M. L. Harnly. 1936. The effects of the gene on growth and differentiation

as shown by the temperature responses of pennant and its heterozygote in D. melanogaster.

Journal of Experimental Zoology 74:41–59.

Hughes, J., Williams ME, Rebeiz M, and Williams TM. 2023. Widespread cis- and trans-

regulatory evolution underlies the origin, diversification, and loss of a sexually dimorphic

fruit fly pigmentation trait. J Exp Zool B Mol Dev Evol 340:143–161. Wiley-Liss, United

States.

Jeong, S., M. Rebeiz, P. Andolfatto, T. Werner, J. True, and S. B. Carroll. 2008. The

Evolution of Gene Regulation Underlies a Morphological Difference between Two

Drosophila Sister Species. Cell 132:783–793. Elsevier.

Kovuri, P., A. Yadav, and H. Sinha. 2023. Role of genetic architecture in phenotypic

plasticity. Trends in Genetics 39:703–714. Elsevier.

Lachaise, D., M. Harry, M. Solignac, F. Lemeunier, V. Bénassi, and M.-L. Cariou. 2000.

Evolutionary novelties in islands: Drosophila santomea, a new melanogaster sister species

from São Tomé. Proceedings of the Royal Society of London. Series B: Biological Sciences

267:1487–1495. Royal Society.

Lafuente, E., F. Alves, J. G. King, C. M. Peralta, and P. Beldade. 2021. Many ways to make

darker flies: Intra- and interspecific variation in Drosophila body pigmentation components.

Ecology and Evolution 11:8136–8155.

Lafuente, E., D. Duneau, and P. Beldade. 2018. Genetic basis of thermal plasticity variation in

Drosophila melanogaster body size. PLOS Genetics 14:e1007686. Public Library of Science.

Lafuente, E., D. Duneau, and P. Beldade. 2024. Genetic basis of variation in thermal

developmental plasticity for Drosophila melanogaster body pigmentation. Molecular Ecology

33:e17294.

Le Poul, Y., A. Whibley, M. Chouteau, F. Prunier, V. Llaurens, and M. Joron. 2014.

Evolution of dominance mechanisms at a butterfly mimicry supergene. Nat Commun 5:5644.

Nature Publishing Group.

Lerner, I. M. 1954. Genetic homeostasis. Oliver and Boyd. London.

Liu, Y., M. Ramos-Womack, C. Han, P. Reilly, K. L. Brackett, W. Rogers, T. M. Williams, P.

Andolfatto, D. L. Stern, and M. Rebeiz. 2019. Changes throughout a Genetic Network Mask

the Contribution of Hox Gene Evolution. Current Biology 29:2157-2166.e6. Elsevier.

Llopart, A., D. Lachaise, and J. A. Coyne. 2005. An Anomalous Hybrid Zone in Drosophila.

Evolution 59:2602–2607.

Lobell, A. S., R. R. Kaspari, Y. L. Serrano Negron, and S. T. Harbison. 2017. The Genetic

Architecture of Ovariole Number in Drosophila melanogaster: Genes with Major,

Quantitative, and Pleiotropic Effects. G3 Genes|Genomes|Genetics 7:2391–2403.

Mackay, T. F. C., and R. F. Lyman. 2005. Drosophila bristles and the nature of quantitative

genetic variation. Philosophical Transactions of the Royal Society B: Biological Sciences

360:1513–1527. Royal Society.

Massey, J. H., and P. J. Wittkopp. 2016. The Genetic Basis of Pigmentation Differences

Within and Between *Drosophila* Species. Pp. 27–61 *in* V. Orgogozo, ed. Current Topics in

Developmental Biology. Academic Press.

Michie, L. J., F. Mallard, M. E. N. Majerus, and F. M. Jiggins. 2010. Melanic through nature

or nurture: genetic polymorphism and phenotypic plasticity in Harmonia axyridis. Journal of

Evolutionary Biology 23:1699–1707.

Negoua, H., M. Chakir, J. R. David, and P. Capy. 2019. Climatic adaptation in Drosophila:

phenotypic plasticity of morphological traits along a seasonal cycle. Annales de la Société

entomologique de France (N.S.) 55:48–60. Taylor & Francis.

Orteu, A., and C. D. Jiggins. 2020. The genomics of coloration provides insights into adaptive evolution. Nat Rev Genet 21:461–475. Nature Publishing Group.

Parkash, R., and A. K. Sharma. 1978. Colour dimorphism in Drosophila jambulina. Drosophila Information Service 166.

Payant, V. 1986. Le polymorphisme de la coloration abdominale dans le genre Drosophila. L'Annee biologique. Comptes Rendus Annuels des Travaux de Biologie Generale 25:167–184.

Pétavy, G., B. Moreteau, J. R. David, and P. Gibert. 2018. Influence of extreme heat or cold stresses on body pigmentation of *Drosophila melanogaster*. Journal of Thermal Biology 72:118–126.

Pétavy, G., B. Moreteau, P. Gibert, and J. R. David. 2002. Phenotypic plasticity of body pigmentation in Drosophila: influence of a developmental thermoperiodic regime in two sibling species. Physiological Entomology 27:124–135.

Pfennig, D. W. (ed). 2021. Phenotypic Plasticity & Evolution: Causes, Consequences, Controversies. CRC Press, Boca Raton.

Pigliucci, M. 2005. Evolution of phenotypic plasticity: where are we going now? Trends in Ecology & Evolution 20:481–486. Elsevier.

Pigliucci, M. 1992. Modelling phenotypic plasticity. I. Linear and higher-order effects of dominance, drift, environmental frequency and selection on a one-locus, two-allele model. J. Genet. 71:135–150.

Pigliucci, M. 2001. Phenotypic Plasticity: Beyond Nature and Nurture. JHU Press.

Pitchers, W., J. Nye, E. J. Márquez, A. Kowalski, I. Dworkin, and D. Houle. 2019. A Multivariate Genome-Wide Association Study of Wing Shape in Drosophila melanogaster. Genetics 211:1429–1447.

Prigent, S. R., Lang ,Michael, Nagy ,Olga, Acurio ,Andrea, Matamoro-Vidal ,Alexis,

Courtier-Orgogozo ,Virginie, and J. R. and David. 2020. Field collections reveal that São

Tomé is the Afrotropical island with the highest diversity of drosophilid flies (Diptera:

Drosophilidae). Annales de la Société entomologique de France (N.S.) 56:1–14. Taylor &

Francis.

Promy, N. T., M. Newberry, and D. Gulisija. 2023. Rapid evolution of phenotypic plasticity

in patchy habitats. Sci Rep 13:19158. Nature Publishing Group.

R Core Team. 2023. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria.

Rocha, F., H. F. Medeiros, and L. B. Klaczko. 2009. The reaction norm of abdominal

pigmentation and its curve in Drosophila mediopunctata depend on the mean phenotypic

value. Evolution 63:280–287.

Rogers, W. A., S. Grover, S. J. Stringer, J. Parks, M. Rebeiz, and T. M. Williams. 2014. A

survey of the trans-regulatory landscape for *Drosophila melanogaster* abdominal

pigmentation. Developmental Biology 385:417–432.

Sapir, Y., M. K. Gallagher, and E. Senden. 2021. What Maintains Flower Colour Variation

within Populations? Trends in Ecology & Evolution 36:507–519. Elsevier.

Scheiner, S. M. 1993. Genetics and Evolution of Phenotypic Plasticity. Annual Review of

Ecology, Evolution, and Systematics 24:35–68. Annual Reviews.

Schmalhausen, I. 1949. Factors of evolution: the theory of stabilizing selection. Blakiston.

Toronto, Canada.

Sellis, D., B. J. Callahan, D. A. Petrov, and P. W. Messer. 2011. Heterozygote advantage as a

natural consequence of adaptation in diploids. Proceedings of the National Academy of

Sciences 108:20666–20671. Proceedings of the National Academy of Sciences.

Sievert, C. 2020. Interactive Web-Based Data Visualization with R, plotly, and shiny.

Chapman and Hall/CRC, New York.

Sultan, S. E. 2021. Phenotypic Plasticity as an Intrinsic Property of Organisms. P. *in*

Phenotypic Plasticity & Evolution. CRC Press.

Suvorov, A., B. Y. Kim, J. Wang, E. E. Armstrong, D. Peede, E. R. R. D'Agostino, D. K.

Price, P. J. Waddell, M. Lang, V. Courtier-Orgogozo, J. R. David, D. Petrov, D. R. Matute, D.

R. Schrider, and A. A. Comeault. 2022. Widespread introgression across a phylogeny of 155

Drosophila genomes. Current Biology 32:111-123.e5.

Takahashi, Y., K. Kagawa, E. I. Svensson, and M. Kawata. 2014. Evolution of increased

phenotypic diversity enhances population performance by reducing sexual harassment in

damselflies. Nat Commun 5:4468. Nature Publishing Group.

Telonis-Scott, M., A. A. Hoffmann, and C. M. Sgrò. 2011. The molecular genetics of clinal

variation: a case study of ebony and thoracic trident pigmentation in Drosophila melanogaster

from eastern Australia. Molecular Ecology 20:2100–2110.

Tian, S., Y. Asano, T. Das Banerjee, S. Komata, J. L. Q. Wee, A. Lamb, Y. Wang, S. N.

Murugesan, H. Fujiwara, K. Ui-Tei, P. J. Wittkopp, and A. Monteiro. 2024. A microRNA is

the effector gene of a classic evolutionary hotspot locus. Science 386:1135–1141. American

Association for the Advancement of Science.

Tichy, A. L., A. Ray, and J. R. Carlson. 2008. A New Drosophila POU Gene, pdm3, Acts in

Odor Receptor Expression and Axon Targeting of Olfactory Neurons. J. Neurosci. 28:7121–

7129. Society for Neuroscience.

Turissini, D. A., and D. R. Matute. 2017. Fine scale mapping of genomic introgressions

within the Drosophila yakuba clade. PLOS Genetics 13:e1006971. Public Library of Science.

van der Burg, K. R. L., J. J. Lewis, B. J. Brack, R. A. Fandino, A. Mazo-Vargas, and R. D.

Reed. 2020. Genomic architecture of a genetically assimilated seasonal color pattern. Science

370:721–725. American Association for the Advancement of Science.

Watanabe, T. K., M. Matsuda, S. Ohnishi, and F. Hihara. 1982. Notes on the systematics of

Drosophila jambulina. The Japanese journal of genetics 57:561–567.

Wei, T., and V. Simko. 2024. R package "corrplot": Visualization of a Correlation Matrix.

Wennersten, L., and A. Forsman. 2012. Population-level consequences of polymorphism,

plasticity and randomized phenotype switching: a review of predictions. Biological Reviews

87:756–767.

West-Eberhard, M. J. 2003. Developmental Plasticity and Evolution. Oxford University Press.

Wittkopp, P. J., E. E. Stewart, L. L. Arnold, A. H. Neidert, B. K. Haerum, E. M. Thompson,

S. Akhras, G. Smith-Winberry, and L. Shefner. 2009. Intraspecific Polymorphism to

Interspecific Divergence: Genetics of Pigmentation in Drosophila. Science 326:540–544.

American Association for the Advancement of Science.

Wittmann, M. J., A. O. Bergland, M. W. Feldman, P. S. Schmidt, and D. A. Petrov. 2017.

Seasonally fluctuating selection can maintain polymorphism at many loci via segregation lift.

Proceedings of the National Academy of Sciences 114:E9932–E9941. Proceedings of the

National Academy of Sciences.

Yassin, A. 2018. Phylogenetic biogeography and classification of the *Drosophila montium*

species group (Diptera: Drosophilidae). Annales de la Société entomologique de France

(N.S.) 54:167–175.

Yassin, A., H. Bastide, H. Chung, M. Veuille, J. R. David, and J. E. Pool. 2016a. Ancient

balancing selection at tan underlies female colour dimorphism in Drosophila erecta. Nat

Commun 7:10400. Nature Publishing Group.

Yassin, A., E. K. Delaney, A. J. Reddiex, T. D. Seher, H. Bastide, N. C. Appleton, J. B. Lack,

J. R. David, S. F. Chenoweth, and J. E. Pool. 2016b. The pdm3 locus is a hotspot for recurrent

evolution of female-limited color dimorphism in Drosophila. Current Biology 26:2412–2422.

Elsevier.

Yassin, A., N. Gidaszewski, V. Debat, and J. R. David. 2022. Long-term evolution of

quantitative traits in the Drosophila melanogaster species subgroup. Genetica 150:343–353.