Article

Phenotypic plasticity as a function of genetic polymorphism: thermal dominance reversal in

***Drosophila* species with contrasting melanism**

Jean R. David[1,2*], Béatrice Denis[1], Pascaline Francelle[2], Alicia Lemaire[1], Aparup Das[3], Sujata

Mohanty[4], Patricia Gibert[5], Amir Yassin[1,2**]

[1]Laboratoire Évolution, Génomes, Comportement et Écologie, CNRS, IRD, Université Paris-

Saclay – Institut Diversité, Écologie et Évolution (IDEEV), 91190 Gif-sur-Yvette, France.

[2]Institut Systématique Evolution Biodiversité (ISYEB) CNRS, MNHN, Sorbonne Université,

EPHE, Université des Antilles, 57 rue Cuvier, CP 50, 75005 Paris, France.

[3]ICMR-National Institute of Research in Tribal Health, Jabalpur, Madhya Pradesh, India.

[4]Department of Biotechnology, Jaypee Institute of Information Technology, Sector 62, Noida,

Uttar Pradesh-201309, India.

[5]Université Claude Bernard Lyon 1, Laboratoire de Biométrie et Biologie Evolutive, CNRS,

UMR5558, Villeurbanne, Rhônes-Alpes, 69100, France.

Running title: Phenotypic plasticity and heterozygosity

*Deceased in 2021.

Corresponding author: amir.yassin@universite-paris-saclay.fr

Abstract

Phenotypic plasticity is often seen as an alternative adaptive strategy to genetic polymorphism, especially in response to rapid environmental changes. Indeed, a link between plasticity and heterozygosity, *i.e.* the basic measure of polymorphism, has previously been dismissed. Here, we investigate the thermal plasticity of abdominal pigmentation in eight *Drosophila* species, four belonging to the *melanogaster* species group and four to the *montium* group. Despite a conserved developmental pathway for melanin synthesis, the genetic architecture of its variation has significantly evolved, being polygenic in most species (like *D. melanogaster*) and Mendelian or invariable in others. By investigating the thermal plasticity of this trait in species with distinct architectures, we show the degree of plasticity to strongly associate with heterozygosity. Plasticity was resurrected in hybrids between species with no plastic responses but with contrasting melanism, and was higher in heterozygotes in species with simple Mendelian polymorphism. In plastic cases, pigmentation dominance is reversed depending on the developmental temperature. We propose simple genetic models with empirical molecular support to explain this link between phenotypic plasticity and genetic polymorphism. The relationship between these two phenomena, and the impact of each on the evolution of the other, may be more relevant than it is currently appreciated.

Keywords

Overdominance; genetic architecture; reaction norm; heterozygosity; dominance reversal.

Introduction

Phenotypic plasticity is the ability of a genotype to produce different phenotypes under

distinct environmental conditions (Bradshaw 1965; Scheiner 1993; Pigliucci 2001; West-

Eberhard 2003; Sultan 2021). Plasticity produces responses that are often summarized in

terms of a reaction norm (Woltereck 1909; Dupont et al. 2024). When such a response

generates the optimal phenotypic values in the different environments, plasticity can facilitate

adaptation even in the absence of genetic variation (Ghalambor et al. 2015). This becomes

particularly important in cases where the population loses a significant portion of genetic

polymorphism, such as after a founder bottleneck during colonization of a new environment,

or when the environment change is faster than the rate of accumulation of new adaptive

mutations, the so-called 'buying time' role of plasticity (Diamond and Martin 2021).

Consequently, several authors have considered phenotypic plasticity and genetic

polymorphism as distinct alternative adaptive strategies (e.g., Wennersten and Forsman

2012).

Despite the definition of plasticity as a property of a genotype, it is usually studied for

complex phenotypic traits whose underlying genotypes are not fully known (e.g., flowering

time in *Arabidopsis*, growth in yeasts, etc.) (Goldstein and Ehrenreich 2021; Kovuri et al. 2023).

Genotypes do not only differ in the degree of their phenotypic plasticity (e.g., the slope of the

reaction norm) but also in their degree of genetic variability, with heterozygous genotypes in

non-haploid organisms having more alleles than hemi- or homozygotes. Among the early

genetic models of phenotypic plasticity, the Overdominance model explicitly related plasticity

to genotypic diversity (Scheiner 1993). The model postulated that plasticity is a function of

homozygosity, with the heterozygous genotype considered to be the most advantageous

because it is the most robust, *i.e.* the least plastic. This assumption reflected the view of

plasticity at the mid-20[th] century as a developmental nuisance (Lerner 1954; Bradshaw 2006).

Indeed, some of the earliest empirical examples of the overdominance model came from

studies of wing morphology in *Drosophila melanogaster*, where two mutants showed the

highest temperature-sensitivity when found in the homozygous state, but their heterozygote

recapitulated the wildtype thermal response, *i.e.* it was less plastic than either mutant (Harnly

and Harnly 1936; Schmalhausen 1949). Gillespie and Turelli (1989) extended this view to a

multi-locus quantitative genetics model and concluded that stabilizing selection on more

environmentally robust heterozygotes will maintain genetic variation in populations. Pigliucci

(1992) developed a one-locus, two-allele classical population genetics model for testing the

evolution of plasticity under overdominance and concurred with Gillespie and Turelli's (1989)

multilocus result. He concluded that plasticity becomes "increasingly disadvantageous when

the system approaches a perfect equilibrium between the probabilities of occurrence of the

two environments."

A direct link between heterozygosity and robustness was not always evident from the

empirical literature, and overdominance was disfavored relative to other genetic models such

as pleiotropy, *i.e.* plasticity is a function of the expression of a gene, and epistasis, *i.e.* plasticity

is a function of interactions between loci. Scheiner (1993) concluded that "plasticity is not a

function of heterozygosity", and Pigliucci (2005), a decade later, stated that "heterozygosity

ha[d] little, if anything, to do with the genetic basis of plasticity." A search using "heterozygo*"

in a recent 400-page long book on *Phenotypic Plasticity and Evolution* (Pfennig 2021) yielded

no result. However, almost none of the studies cited by Scheiner (1993) and Pigliucci (2005) to

reach their conclusions has addressed the link between heterozygosity and plasticity in the

most optimal setting similar to Pigliucci's (1992) theoretical model, *i.e.* comparing the reaction

norms of a phenotype with a simple, well-characterized genotypic basis. Conspicuous color

polymorphisms are ideally suited for these experiments, because they often involve a simple

genetic architecture and color synthesis in many species results from a well conserved gene

network (Orteu and Jiggins 2020). This is particularly true for melanism. In several lineages of

butterflies, simple Mendelian polymorphisms have evolved in response to the industrial

revolution, with the melanic wing allele being dominant in some species and recessive in

others (Ford 1945). The genetic basis of this industrial melanism has recently been identified

to be a conserved micro-RNA close to the locus of the transcription factor *Cortex* (Tian et al.

2024). Remarkably, several butterfly species also show a seasonal melanic polyphenism, with

cold season morphs being darker. van der Burg et al. (2020) showed that mutations at the

same locus containing *Cortex*, and consequently the linked micro-RNA, underlie this

polyphenism in *Junonia caenia*. Plasticity in simple color polymorphism systems was also

observed in *Harmonia* ladybugs morphs (Michie et al. 2010).

Much of our knowledge about the genetic and developmental basis of melanin

synthesis comes from studies in *Drosophila*. Pigmentation differences between body

segments, sexes, geographical populations, seasons, and species are often due to *cis-* and

trans-regulatory modifications of melanin synthesis genes, such as *tan (t)*, *yellow (y)* or *ebony*

(e), and their transcription factors, such as *bric-à-brac (bab)*, *POU-domain-motif-3 (pdm3)* or

Abdominal-B (Abd-B) (Massey and Wittkopp 2016; Hughes et al. 2023). Wild-caught flies from

the same locality usually differ in their degree of pigmentation (Figure 1 A; Bastide et al. 2013,

2016; Dembeck et al. 2015; Endler et al. 2016; Castro et al. 2018), but flies collected from cold

high altitudes, latitudes and seasons tend to be dark, even when their progeny is maintained

under standard thermal conditions (Telonis-Scott et al. 2011; Bastide et al. 2014, 2016;

Takahashi et al. 2014; Negoua et al. 2019; Berardi et al. 2024). Similarly, when full-sib flies

issued from the same isofemale line are grown under different temperatures, a negative

reaction norm is detected, with flies grown under low temperatures being darker (Figure 1;

(Gibert et al. 1996, 1998, 2000, 2004b, 2009; Chakir et al. 2002; Pétavy et al. 2002, 2018;

Lafuente et al. 2021, 2024). Thermal plasticity in *D. melanogaster* is due to the higher

expression of the *t* and *y* genes in colder temperatures (Gibert et al. 2016, 2017b). A natural

variation in the transcription factor *bab* underlying color differences also showed distinct

responses to developmental temperature through repression of *t* and *y* in hotter environments

(Castro et al. 2018). This parallelism between genetic polymorphism in the wild and

phenotypic plasticity in the lab is strongly suggestive for an adaptive role for darker

pigmentation in thermoregulation. Indeed, drosophilids with darker cuticle have higher body

temperature (Freoa et al. 2023).

FA6	-0.40±0.03***
FA7	-0.58±0.06***

MA5	-0.03±0.00**
MA6	0.00±0.00

Figure 1 – Pigmentation variation in *Drosophila melanogaster*. A) Variation in abdominal pigmentation can be due to genetic polymorphism (left) or thermal phenotypic plasticity (right) (Photos from Jean R. David's archive). B) Three-dimensional analysis of pigmentation variation across growth temperatures and abdominal segments. C) Pigmentation thermal plasticity in females. D) Pigmentation thermal plasticity in males. Reaction norm slopes for the last two abdominal segments (A6 and A7 in females (F) and A5 and A6 in males (M)) are given for each sex. (Data from Gibert et al. 2000).

The genetic architecture of pigmentation variation has dramatically evolved between

drosophilid species. Whereas female abdominal pigmentation variation is continuous and

polygenic in *D. melanogaster* as well as most species, it is discrete and monogenic in multiple

species (Payant 1986; Yassin et al. 2016a,b). In some species, pigmentation is nearly fixed with very slight variability even between the sexes. This is the case of the two sister species, *D. santomea* and *D. yakuba*, where both males and females' posterior abdomens are invariably light or dark, respectively (Lachaise et al. 2000; Jeong et al. 2008; David et al. 2022). We aim here at quantifying the level of pigmentation phenotypic plasticity in eight species with distinct genetic architectures grown under six thermal regimes. To this end, we focused on the effect of heterozygosity, as it can be seen in species hybrids (*i.e.* the ultimate viable heterozygotes) and heterozygous genotypes of five species with monogenic architectures. Our results support a positive link between plasticity and the degree of genetic polymorphism that was not previously assumed.

Materials and Methods

We studied five cases where for each the thermal phenotypic plasticity can be compared between homozygous dark and light lines and their reciprocal hybrids. Four cases considered within-species monogenic Mendelian color polymorphism with one (*D. erecta*) from the *melanogaster* species group and three cases (*D. jambulina*, *D. burlai* and *D.* cf. *bocqueti*) from its sister clade, the *montium* species group. For each of these species homozygous dark and light lines were established through selection from a polymorphic mass culture (Table 1). The monogenic basis of female pigmentation in these species are known (Yassin et al. 2016a,b; Prigent et al. 2020). In two species (*D. erecta* and *D. burlai*), the dark allele is dominant (Yassin et al. 2016a,b), whereas in the two other species (*D. jambulina* and *D.* cf. *bocqueti*), the light allele is dominant (Watanabe et al. 1982; Prigent et al. 2020). The fifth case involved hybridization between the two sister species *D. santomea* and *D. yakuba* of the *melanogaster* group (Table 1). We also obtained published data of the thermal plasticity

162   response of *D. melanogaster* (Gibert et al. 2000), the nomitypical species of the *melanogaster*

163   group with polygenic basis of pigmentation variation, and the monogenic *D. kikkawai* (Gibert

164   et al. 1999) for comparison purposes (Table 1).

166   Table 1. *Drosophila* strains analyzed in this study.

Group	Subgroup	Complex	Species	Strain
melano-gaster	*melano-gaster*	*melano-gaster*	*melanogaster*	Bordeaux, France (Gibert et al. 2000)
melano-gaster	*melano-gaster*	*yakuba*	*yakuba*	Kunde, Cameroon, 1967 (L. Tsacas)
melano-gaster	*melano-gaster*	*yakuba*	*santomea*	Sao Tomé, Sao Tomé & Principe, 1998 (D. Lachaise)
melano-gaster	*melano-gaster*	*erecta*	*erecta*	La Lopé Forest, Gabon, 1998 (D. Lachaise)
montium	*kikkawai*	*kikkawai*	*kikkawai*	Chandigarh, India (Gibert et al. 1999)
montium	*seguyi*	*jambulina*	*jambulina*	New Delhi, India, 2014 (J. R. David, S. Mohanty)
montium	*seguyi*	*bocqueti*	*burlai*	Mt Oku, Cameroon, 2008 (M. Veuille)
montium	*seguyi*	*bocqueti*	cf. *bocqueti*	Sao Tomé, Sao Tomé & Principe, 2015 (J. R. David, M. Lang)

For each case, pigmentation was scored on >4 days old 10 males and 10 females as in

Yassin et al. (2022) following the pigmentation score scale proposed by David et al. (1990).

Flies were grown at five temperatures going from 14°C to 28°C with 3° increments. In each

case, pigmentation was scored on six and five abdominal segments in females and males

respectively (denoted A2-7 with F and M for females and males, respectively). All statistical

analyses were conducted using R (R Core Team 2023), with 3D plots obtained using the Plotly

package in R (Sievert 2020). For the sake of simplicity, the degree of phenotypic plasticity (*i.e.*

the shape of the reaction norm) was summarized in terms of the slope of a linear regression.

For monogenic species, we followed the nomenclature given in Gibert et al.'s (1999) paper,

referring to the light and dark alleles as *L* and *D*, respectively, and to the three genotypes as *L*

x L, *L x D* or *D x L*, and *D x D*. For heterozygotes, the allele from the mother is given first

depending on the cross.

Results

Ancestral plasticity resurrection in D. santomea *and* D. yakuba *hybrids*

Abdominal pigmentation is the most discriminating quantitative trait among the nine

species of the *melanogaster* subgroup (Yassin et al. 2022). The thermal plasticity of this trait

was examined only in two sibling species, *D. melanogaster* and *D. simulans*, and showed quasi-

parallel reaction norms (Pétavy et al. 2002). In females of both species, the phenotypic

plasticity between cold and warm environments increases towards most posterior abdominal

segments, with the pigmentation of A7 being the most plastic (Figure 1C). In males, on the

other hand, this pattern is restricted to the first three abdominal segments (A2, A3 and A4),

whereas the last two segments (A5 and A6) are completely dark with almost no plastic

response (Figure 1D). Both species belong to the *melanogaster* complex, but similar patterns

of pigmentation plasticity was observed in *D. suzukii* (Colinet and Kustre 2025), an outgroup

species belonging to another subgroup of the *melanogaster* group, indicating that plasticity is

likely ancestral.

Drosophila santomea and *D. yakuba* belong to the *yakuba* complex of the

melanogaster subgroup. The two sister species are monomorphically either light (*D.*

santomea) or black (*D. yakuba*) and reciprocal crosses produce hybrid fertile females and

sterile males with intermediate pigmentation phenotypes in both (Llopart et al. 2005; Liu et

al. 2019; David et al. 2022). At least two X-linked and four autosomal genes underlie the

pigmentation differences between these species (Liu et al. 2019; David et al. 2022). We found

the two species to lack plasticity (Figure 2A). However, in reciprocal crosses, plasticity was

generated in both sexes in a pattern similar to that of *D. melanogaster* females, *i.e.* darker

phenotypes in cold temperature (Figure 2A). The degree of plasticity, as measured by the slope

of the reaction norms of the two last abdominal segments, did not differ between reciprocal

hybrid females, since these females share the same heterozygous genotypes for both the X-

linked and autosomal genes. In reciprocal hybrid males, on the other hand, males mothered

by *D. santomea*, *i.e.* those carrying loss-of-function mutations in the X-linked pigmentation

genes *y* and *t*, had a steeper slope than males mothered by *D. yakuba* with functional copies

of these genes. In females and males mothered by *D. santomea*, dominance is completely

reversed with pigmentation of hybrid flies resembling *D. yakuba* in cold temperature and *D.*

santomea in hot temperature.

Loss-of-function mutations at t *leads to loss of plasticity in* D. erecta *recessive homozygotes*

Of the six loci underlying pigmentation difference between *D. yakuba* and *D. santomea*,

only one, the X-linked *t*, underlies female-limited Mendelian color dimorphism in *D. erecta*

(Yassin et al. 2016a), a species of the *erecta* complex of the *melanogaster* subgroup. We found

that the loss-of-function mutations in a likely female-specific suppressor of this gene likely led

to no thermal plasticity in the recessive homozygous light line (Figure 2B). However, these

mutations are completely recessive in heterozygous females, where similar thermal slopes to

that of the homozygous dark females were found. Regardless of the genotype, males showed

no plastic response. So, plasticity does not correlate with heterozygosity in *D. erecta*, but

rather, to a low extent, with a functional *t* gene.

A   *D. santomea*, *D. yakuba* and their hybrids

	san x san	san x yak	yak x san	yak x yak
FA6	-0.02±0.01	-0.24±0.07*	-0.32±0.09*	-0.04±0.08
FA7	0.00±0.00	-0.56±0.07**	-0.67±0.06***	-0.07±0.03
MA5	-0.02±0.01	-0.29±0.04**	-0.20±0.01***	-0.07±0.03
MA6	-0.06±0.02	-0.53±0.05***	-0.21±0.02***	-0.08±0.01**

B   *D. erecta*

	L x L	L x D	D x L	D x D
FA6	-0.04±0.03	-0.57±0.09**	-0.54±0.07**	-0.55±0.06**
FA7	-0.08±0.05	-0.11±0.05	-0.06±0.03	0.00±0.00
MA5	-0.09±0.05	-0.10±0.06	-0.03±0.07	-0.09±0.08
MA6	0.00±0.00	0.00±0.00	0.0±0.00	0.00±0.00

Figure 2 – Thermal phenotypic plasticity of pigmentation on successive abdominal segments in species of the *melanogaster* group. A) *Drosophila santomea (san)*, *D. yakuba (yak)* and their reciprocal hybrids. B) *D. erecta*. For each cross, the genotype of the mother is given first. *L* and *D* for light and dark alleles, respectively. Reaction norm slopes for the last two abdominal segments (A6 and A7 in females and A5 and A6 in males) are given for each sex and genotype. The 3D plot axes are as in Figure 1B.

Phenotypic plasticity correlates with heterozygosity in females of species of the montium *group*

with monogenic variation and dark dominant allele

Female-limited Mendelian melanism, similar to that found in *D. erecta*, has evolved in

22 *montium* group species belonging to five distinct subgroups (Yassin et al. 2016b; Yassin

2018). Of the 15 species where the genetics was studied, the dark allele was dominant in 13

species (Yassin et al. 2016b). In four species with dark-dominant allele namely *D. serrata*

(*serrata* subgroup), *D. kikkawai* and *D. leontia* (*kikkawai* subgroup) and *D. burlai* (*seguyi*

subgroup), the polymorphism maps to a single locus, the autosomal transcription factor *pdm3*

(Yassin et al. 2016b). In these species, males have a light abdomen in the three former species,

which is characteristic to their respective subgroups, whereas males have a dark abdomen in

D. burlai, which is common in the *seguyi* subgroup to which it belongs (Yassin et al. 2016b;

Yassin 2018). The loss of male pigmentation in *D. kikkawai* is, at least partially, due to *cis*-

regulatory mutations in the X-linked melanin synthesis gene *y* (Jeong et al. 2006).

Gibert et al. (1999) compared the thermal plasticity of abdominal pigmentation

between homozygous light and dark *D. kikkawai* lines and their progeny from reciprocal

crosses. For females, heterozygotes had nearly twice the slope of either homozygous line in

A6, whereas in males, plasticity increases with the number of copies of the dark *pdm3* allele

for both A5 and A6 (Figure 3A). In *D. burlai*, female heterozygotes had stronger plasticity than

homozygotes only for A7, whereas no difference was found at either A5 and A6 in males, with

most of the genotypes being plastic (except A6 in homozygous dark males) (Figure 3B).

A *D. kikkawai*

	L x L	L x D	D x L	D x D
Female				
FA5	-0.09±0.05	-0.18±0.04*	-0.28±0.09	-0.07±0.07
FA6	-0.07±0.03	-0.11±0.02*	-0.12±0.02*	-0.01±0.02
FA7	-0.39±0.11	-0.02±0.00	-0.03±0.02	-0.01±0.00
Male				
MA4	-0.07±0.01*	-0.04±0.01*	-0.01±0.02	-0.08±0.02*
MA5	-0.01±0.00*	-0.03±0.01	-0.03±0.01*	-0.08±0.02*
MA6	0.00±0.00	-0.27±0.11	-0.23±0.05*	-0.58±0.15

B *D. burlai*

	L x L	L x D	D x L	D x D
Female				
FA6	-0.11±0.02***	-0.08±0.02***	-0.01±0.00*	0.00±0.00
FA7	-0.01±0.01	-0.13±0.01***	-0.05±0.01***	0.00±0.00
Male				
MA5	-0.14±0.01***	-0.35±0.04***	-0.24±0.03***	-0.27±0.03***
MA6	-0.10±0.02***	-0.01±0.00**	-0.01±0.01*	0.00±0.00

Figure 3 – Thermal phenotypic plasticity of pigmentation on successive abdominal segments in monogenic species of the *montium* group where the dark allele is dominant. A) *Drosophila kikkawai* (data from (Gibert et al. 1999). B) *D. burlai*. For each cross, the genotype of the mother is given first. *L* and *D* for light and dark alleles, respectively. Reaction norm slopes for the last two abdominal segments (A6 and A7 in females and A5 and A6 in males) are given for each sex and genotype. The 3D plot axes are as in Figure 1B.

265   *Phenotypic plasticity reaches its maximum in female heterozygotes of species of the* montium

266   *group with monogenic variation and light dominant allele*

267   *Drosophila jambulina* is one of three species of the *montium* group with female

Mendelian melanism wherein the light allele was reported to be dominant (Yassin et al. 2016b;

Prigent et al. 2020). However, this dominance relationship is ambiguous since Parkash and

Sharma (1978) found dark dominance for an Indian population, whereas Watanabe et al.

(1982) found light dominance also for Indian populations and concluded that dominance in *D.*

jambulina was "a great problem existing in a simple genetic phenomenon". We analyzed the

phenotypic plasticity of homozygous light and dark female strains along with their reciprocal

heterozygotes. The female posterior abdominal pigmentation of both homozygous lines

completely lacked a plastic response (Figure 4A). The heterozygous females on the other hand

showed the steepest plastic response with a slope ≥0.80 for both A6 and A7. Indeed, at

temperatures ≥21°C, the light allele is dominant, but at temperatures <21°C the dark allele is

dominant. It is thus possible that Parkash and Sharma (1978) and Watanabe et al. (1982) have

conducted their experiments at different temperatures reaching contradictory results. *D.*

jambulina, like *D. burlai*, belongs to the *seguyi* subgroup, with males having dark posterior

abdomen (Yassin 2018). Like *D. burlai*, A5 was plastic for nearly all genotypes (except

homozygous light), whereas A6 showed weak plasticity.

A *D. jambulina*

	L x L	L x D	D x L	D x D
Female				

| FA6 | -0.02±0.04 | -0.80±0.06*** | -0.92±0.05*** | 0.00±0.00 |
| FA7 | -0.02±0.04 | -0.84±0.06*** | -0.95±0.05*** | 0.00±0.00 |

Male

| MA5 | -0.12±0.02 | -0.25±0.02*** | -0.18±0.02*** | -0.11±0.02*** |
| MA6 | -0.07±0.01*** | -0.01±0.00* | 0.00±0.00 | 0.00±0.00 |

B *D. cf. bocqueti*

	L x L	L x D	D x L	D x D
Female				

| FA6 | 0.00±0.00 | -0.73±0.04*** | -0.69±0.04*** | -0.03±0.01*** |
| FA7 | 0.00±0.00 | -0.79±0.06*** | -0.74±0.05*** | -0.06±0.02*** |

Male

| MA5 | -0.57±0.08*** | -0.23±0.03*** | -0.29±0.02*** | -0.13±0.02*** |
| MA6 | 0.00±0.00 | -0.01±0.00 | -0.02±0.01** | -0.01±0.01 |

Figure 4 – Thermal phenotypic plasticity of pigmentation on successive abdominal segments in monogenic species of the *montium* group where the light allele is dominant. A) *Drosophila jambulina*. B) *D.* cf. *bocqueti*. For each cross, the genotype of the mother is given first. *L* and *D* for light and dark alleles, respectively. Reaction norm slopes for the last two abdominal segments (A6 and A7 in females and A5 and A6 in males) are given for each sex and genotype. The 3D plot axes are as in Figure 1B.

Prigent et al. (2020) reported a third case of a *montium* species with dominant light

allele in the Sao Tomé strains that we analyze here as *D.* cf. *bocqueti*. Like *D. burlai* and *D.*

jambulina, *D.* cf. *bocqueti* belongs to the *seguyi* subgroup, but it is closely related to *D. burlai*,

with whom and along a third species, *D. chauvacae*, they constitute the Afrotropical *bocqueti*

species complex within the *seguyi* subgroup (Yassin 2018). We crossed *D.* cf. *bocqueti* and *D.*

burlai and reciprocal crosses yielded inviable progeny, hence supporting the close relationship

between them and their proper specific status despite their great morphological similarity.

Unlike *D. burlai*, where the dark allele is constantly dominant, we found strong dominance

reversal in heterozygotes similar to that found in the distantly-related *D. jambulina*; the light

allele was dominant at temperatures ≥21°C and the dark allele is dominant at <21°C (Figure

4B). Like *D. jambulina*, pigmentation of female heterozygotes has a very steep slope ≥0.70,

whereas the slope of the homozygotes never exceeded 0.06 for both A6 and A7. Similar to *D.*

jambulina also, the posterior abdomen of males showed a weak plasticity, with the strongest

slope found for the homozygous light genotype but for A5 rather than for A6. Because both

species are tropical, predominantly living in environments with an average temperature ≥21°C,

the light allele may be considered the dominant one.

Excess heterozygosity among the most plastic abdominal segments

The distribution of the total 272 reaction norms that we estimated for all segments,

sexes, strains and species has a near zero mean ~-0.13, meaning that most segments have low

plasticity and are developmentally robust (Figure 5A). However, the distribution is strongly

negatively skewed as expected from the general pattern of low pigmentation under high

temperatures. Nearly 10% of the reaction norms fall below -0.25, and these correspond to the

most plastic segments. This pool of strongly plastic segments was not enriched for species (chi-

squared P = 0.080) or sex (chi-squared P = 0.258). However, there was a bias for posterior

segments (A5-7) (chi-squared P = 4.2 x 10^{-5}). Most importantly, 80% of the most plastic

segments were found in heterozygous genotypes (chi-squared P = 6.4 x 10^{-4}), with the opposite

trend, *i.e.* excess homozygotes, in the pool of most robust segments with slopes above the 0.9

quantile (chi-squared P = 0.001) (Figure 5A).

All cases where a heterozygous genotype had a strong plasticity showed a reversal of

dominance, *i.e.* the phenotype of the heterozygote (F1, averaged over reciprocal crosses)

tended towards the dark homozygous genotype in cold temperatures (the blue curve of F1 –

D x D in Figure 5B), and towards the light homozygous genotype in hot temperatures (the red

curve of *L x L* – F1 in Figure 5B). The intersection between the two curves corresponds to the

temperature at which dominance is reversed, as well as the phenotypic value of the

codominance at that temperature, *i.e.* the point at which the heterozygote is at mid-distance

between both homozygotes. From these curves, showed for the last two abdominal segments

of each sex in Figure 5B, dominance reversal occurred for both males and females in the case

of the hybrids between *D. santomea* and *D. yakuba*, only in females in *D. erecta*, *D. jambulina*

and *D.* cf. *bocqueti*, and only in males in *D. burlai*. No intersection was found in *D. kikkawai*,

but an intersection at extreme temperature in the female at A5 was found. *D. kikkawai* is the

only species showing reversed slopes depending on the sex. The phenotype of the male

heterozygote approaches the dark genotype at high temperatures, where they both become

light, whereas the phenotype of the female heterozygote approaches the light genotype at

low temperatures, where they both become dark.

Figure 5 – Thermal phenotypic plasticity associates with heterozygosity. A) Histogram of the
reaction norm slopes of 272 abdominal segments analyzed across all species. The red line

indicates the 0.1 quantile (values below which plasticity is high). The pie charts reflect the proportions of the heterozygous and homozygous genotypes among the most plastic (≤0.1 quantile) and the most robust (≥0.9 quantile) abdominal segments. B) Patterns of temperature-dependent dominance reversal in the six analyzed cases. F1 is averaged across reciprocal crosses. *LL* and *DD* are the light and dark homozygous genotypes, respectively. For each case the penultimate and ultimate abdominal segments are represented for both sexes on the left and right, respectively. Female slopes are represented as solid lines and male slopes as dashed lines. The graphs demonstrate the pigmentation difference between the heterozygous genotype (F1), and both homozygotes. The value of the heterozygote is presented as 0, *i.e.* F1 − F1, whereas the differences from the homozygotes are presented as *LL* − F1 and *DD* − F1, which should produce negative values as long as dominance is incomplete. Note that in species with dominance reversal, the *LL* − F1 difference tends to approach the value of the heterozygote in high temperatures, whereas the F1 − *DD* difference tends to approach the value of the heterozygote in cold temperatures. The point of intersection between *LL* − F1 and *DD* − F1 indicates the temperature at which dominance is reversed.

Discussion

The molecular basis of the association between plasticity and heterozygosity in Drosophila

Although it has long been recognized that genetic variation of plasticity exists, the nature of this variation remains elusive. Early works suggested three non-mutually exclusive mechanisms, namely overdominance, pleiotropy, and epistasis (Scheiner 1993). Of the three models, only overdominance made an explicit prediction about the relationship between plasticity and heterozygosity. The lack of strong evidence that heterozygous genotypes were more robust has led to the dismissal of the overdominance model, and current studies on the molecular basis of plasticity either consider the epigenetic and transcriptomic landscapes (*i.e.* the pleiotropy model) or the shape of regulatory networks (*i.e.* the epistasis model) (Goldstein and Ehrenreich 2021; Chevin et al. 2022). Our analysis of 272 reaction norms of a serially homologous trait supports an association between plasticity and heterozygosity but in a positive rather than negative way, hence revisiting the classical overdominance model.

In its simplest form, a gene regulatory network consists of a transcription factor activating or suppressing an effector gene. Of the five monogenic cases studied here, the

underlying locus involves an effector gene, *t*, in one species (*D. erecta*) (Yassin et al. 2016a),

and a transcription factor, *pdm3*, in four species (*D. kikkawai*, *D. jambulina*, *D. burlai* and *D.* cf.

bocqueti) (Yassin et al. 2016b; Fukutomi et al. 2025) (Figure 6). Loss-of-function regulatory

mutations of *t* reduce plasticity in *D. erecta* recessive light homozygotes (Figure 2B), and only

slightly affect the degree of dominance, *i.e.* plasticity of the functional allele is nearly

completely dominant with a reversal of dominance near the upper thermal limit when both

dark heterozygotes and homozygotes become light (Figure 5B). Indeed, the *t* regulatory

element which is involved in female color variation in both *D. erecta* and *D. melanogaster*

(Bastide et al. 2013; Endler et al. 2016; Gibert et al. 2017a) partly affects pigmentation

plasticity in the later species (Gibert et al. 2016). On the other hand, mutations in the

transcription factor *pdm3* increases plasticity when present in a heterozygous state in the four

species of the *montium* group. Polymorphism in *pdm3* associates with female abdominal

pigmentation variation in European and African populations of *D. melanogaster* (Endler et al.

2016). Mutations in this gene also underlie pigmentation evolution between the sister species

D. yakuba and *D. santomea* with contrasting melanism (Liu et al. 2019; David et al. 2022).

Whereas pigmentation in these two species is not plastic, their reciprocal hybrids, which are

heterozygous for *pdm3* interspecific differences along with other genes, are highly plastic

(Figure 2A; Figure 5B). This may indicate that the relation between heterozygosity and

plasticity may mainly depend on the position of the polymorphic gene in the genetic network,

i.e. being more pronounced if the gene is a transcription factor rather than an effector gene.

	Genetic variation	Genes	Thermal plasticity
melanogaster group			
melanogaster	Continuous	E: *t, e* TF: *bab, pdm3*	High
simulans	Continuous	Unknown	High
yakuba	No variation	E: *y, t, e* TF: *Abd-B,* *pdm3, Gug*	No
santomea	No variation		Hybrid high No
erecta	Discrete, dark dominant	E: *t*	No
suzukii	Continuous	Unknown	High
kikkawai	Discrete, dark dominant	TF: *pdm3*	Low
jambulina	Discrete, light dominant	TF: *pdm3*	Heterozygous high Homozygous no
burlai	Discrete, dark dominant	TF: *pdm3*	No
cf. bocqueti	Discrete, light dominant	TF: *pdm3*	Heterozygous high Homozygous no
montium group			

Figure 6 – Evolution of the genetic architecture of pigmentation variation and its thermal plasticity in the *melanogaster* and *montium* species groups. E = effector melanin-synthesis gene; TF = transcription factor. Phylogenetic relationships are after Suvorov et al. (2022).

Besides, the types of the mutations in the alleles of the polymorphic gene may also affect the degree of the plastic response. Indeed, plasticity was highest in the heterozygotes of the two *montium* group species, *D. jambulina* and *D.* cf. *bocqueti*, where the light allele was dominant under optimal developmental temperature regimes, *i.e.* ≥21°C. The reversal of dominance according to the temperature, the main reason of the steepness of the reaction

norm, is associated with the evolution of dominance between the *pdm3* alleles in these

species. In *D. melanogaster*, the wildtype light allele is dominant (Tichy et al. 2008), but RNAi

lines targeting different parts of this gene yields contrasting results, either increasing

pigmentation (mostly in females) or decreasing it (mostly in males) (Rogers et al. 2014).

Consequently, it is possible that the *pdm3* mutations underlying the dark allele in *montium*

species where it is dominant, such as in *D. kikkawai* and *D. burlai*, are of the dominant negative

type, which are common for transcription factors (Gerasimavicius et al. 2022). *Pdm3* has

multiple isoforms in *D. melanogaster* and if one of them has the DNA binding domain but lacks

polymerase or other protein binding domains that are needed to express a gene, whereas

another isoform has both DNA and protein binding domains, the isoforms will have opposite

effects on pigmentation. In the heterozygote, the isoform lacking the protein binding domain

will compete with the isoform bearing this domain, hence altering its function. On the other

hand, mutations in species with the light allele being dominant, like *D. jambulina* and *D. cf.*

bocqueti, may be of the classical loss-of-function type that may reduce but do not completely

eliminate the effect of the wild-type light allele. Further molecular dissection of the underlying

alleles is required to understand the evolution of dominance reversal at *pdm3*.

The effect of mutations in both modifier and effector genes on plasticity is best

illustrated in *D. kikkawai* and the hybrids between *D. santomea* and *D. yakuba*. In *D. kikkawai*,

the effector *yellow (y)* gene lacks a conserved *cis*-regulatory element leading to the loss of

body pigmentation (Jeong et al. 2008). Because this gene is X-linked, a dosage effect may occur

between the sexes. Plasticity in males additively increased with the number of copies of the

dominant dark *pdm3* allele, which is an indirect regulator of *y* (Liu et al. 2019). Putatively

negative dominant mutations in this gene are expected to activate *y*, but the effect may

depend on the number of copies of *y*, being higher in females. Similarly, whereas hybrid males

mothered by *D. santomea*, *i.e.* those carrying defective X-linked melanin synthesis genes *t* and *y* had similar plasticity slopes to hybrid females, but this slope was halved in hybrid males mothered by *D. yakuba*, with functional *y* and *t*. Therefore, the degree of plasticity is affected by the functionality of the polymorphism at the underlying genes.

The association between plasticity and polymorphism may be more general than currently appreciated

We have restricted our analyses to eight species belonging to two species groups of the subgenus *Sophophora*. Despite the differences in their reaction norms, both groups contain species with sexually dimorphic melanism, and a single gene with antagonistic sexual effect may underlie the association between plasticity and heterozygosity, hence reducing the independence of our observations. Future analyses should investigate pigmentation thermal plasticity in non-*Sophophora* fly species lacking sexual dimorphism, with or without Mendelian polymorphism (e.g., Brisson et al. 2005; Rocha et al. 2009; Wittkopp et al. 2009). Beyond *Drosophila*, color polymorphism abounds in nature (Orteu and Jiggins 2020; Sapir et al. 2021). Investigation of the diversity in reaction norms in species with heterozygous and homozygous color genotypes, especially in these with a hierarchy of dominance (e.g., Durand et al. 2014; Le Poul et al. 2014), should shed light on the generality of the relationship between heterozygosity and the degree of plasticity.

Most quantitative traits for which phenotypic plasticity was studied in *Drosophila*, such as body size, wing shape, bristle number, ovariole number or life-history traits, are highly polygenic with often a redundant genetic basis (David et al. 2004; Gibert et al. 2004a; Mackay and Lyman 2005; Lobell et al. 2017; Lafuente et al. 2018; Pitchers et al. 2019). Identifying the underlying genotypes of these traits to test for the effect of heterozygosity may be difficult. An

alternative to the 'forward genetics' approach that we leveraged here, is to use a 'reverse

genetics' analysis, wherein the transcription level of polymorphic genes can be analyzed in

homozygous vs. heterozygous states in different environments, even in the lack of knowledge

about the phenotypic consequences of the genotypes. One such significant example comes

from a study by Chen et al. (2015) who showed that 62% of genes with thermal plastic

response in *D. melanogaster* (*i.e.* ~51% of total genes) swapped allelic dominance in

heterozygotes between cold and hot environments. Future similar transcriptomic analyses are

therefore highly needed to test the generality of these patterns, especially these using hybrids

between species.

Dominance reversal is a major mechanism for higher plasticity in heterozygotes

Major insight on the genome-wide ubiquity of dominance reversal was recently

demonstrated by Grieshop and Arnqvist (2018). These authors compared fitness, a polygenic

trait by nature, between the sexes in crosses between inbred lines of seed beetles and found

a strong dominance reversal between the sexes. Grieshop et al. (2024) reviewed other

examples including environment-, trait- and sex-specific dominance reversal. Such examples

are ideal models to test the link between plasticity and heterozygosity, especially in the context

when sexual antagonism, *i.e.* distinct fitness optima for each sex, may interfere with

environment-specific dominance reversal.

We note here that in our analysis, a sexual dimorphism in thermal plastic response was

observed. In *D. burlai*, females are less plastic for all genotypes, whereas the opposite is true

for *D. melanogaster*. In *D. jambulina* and *D. cf. bocqueti*, homozygous genotypes were more

plastic in males than in females, whereas heterozygous genotypes were more plastic in

females. The strongest example was in *D. kikkawai*, wherein the dominance of the dark *pdm3*

allele reversed between the sexes (in the heterozygotes) with higher plasticity in opposite

homozygous genotypes (Gibert et al. 1999) (Figure 3A). Indeed, the sexual dimorphism in *D.*

melanogaster is caused by the action of the transcription factor *bab* which suppresses melanin

synthesis genes in females (Kopp et al. 2000; Rogers et al. 2014; Hughes et al. 2023). This same

gene is also responsible for the further suppression of pigmentation in females grown at high

temperature (Castro et al. 2018). Because *bab* is repressed in dark *D. burlai* females and

expressed in light *D. bocqueti* females (Kopp et al. 2000), and *bab* is a direct regulator of *y*, it

is possible that *bab* sexual expression and plasticity are regulated by *pdm3*. These observations

highlight the complexity underlying the link between heterozygosity and plasticity, and the

necessity for future theoretical modeling of this link to account for possible interactions

between the environment and the sex.

To the best of our knowledge, the only simple population genetics model, *i.e.* a single

locus with two alleles, for the evolution of plasticity was developed by Pigliucci (1992). He

modeled fitness as a phenotype and considered the heterozygous genotype to have equal

fitness in two environments, *i.e.* a robust fitness, in contrast to homozygous genotypes that

had symmetrically opposite plastic fitness. However, one should distinguish the difference

between developmental plasticity, *i.e.* the genotype-phenotype map, from fitness plasticity. In

fact, Pigliucci's (1992) model of complete dominance agrees with a model of developmental

dominance reversal, which can occur if the heterozygote switches its phenotype to that of

each homozygous genotype in the environment where the phenotype is beneficial.

Overdominance will follow if the homozygotes cannot switch their phenotype between the

environments, *i.e.* a robust genotype has a plastic fitness, whereas a plastic genotype has a

robust fitness. This will lead to fitness relationships similar to the ones that we present in

Figure 5B (compare to Figure 1 in Pigliucci 1992). The realization of this distinction between

developmental and fitness plasticity leads us to the exact opposite interpretations of Pigliucci's

(1992) model: plasticity, not robustness, should prevail when the system approaches

equilibrium between the probabilities of occurrences of the different environments.

The evolutionary consequences of links between phenotypic plasticity and polymorphism

A link between phenotypic plasticity and polymorphism can provide simple

explanations on how plasticity evolves. In fact, plasticity can be lost through any mechanism

reducing genetic heterozygosity such as inbreeding or directional selection. Indeed, the

'flexible stem hypothesis' postulates that from a plastic ancestor, descendants with fixed

phenotypes from the ancestral phenotypic spectrum can evolve (West-Eberhard 2003; Gibert

2017). A likely example from our data is the case of *D. santomea* and *D. yakuba*. Pigmentation

difference between these two species is due to mutations in at least six loci (Liu et al. 2019;

David et al. 2022), but it is unlikely that all these mutations have arisen simultaneously. This

means that a substantial degree of heterozygosity segregated in the ancestor of these species,

and consequently this ancestor has gradually lost its plasticity as more and more alleles

became fixed in each lineage. Indeed, pigmentation plays a role in the reproductive isolation

between these two species (David et al. 2022), and a hybrid zone on the African island of Sao

Tomé between these two species exists (Lachaise et al. 2000; Llopart et al. 2005; Turissini and

Matute 2017). A plastic hybrid is disadvantaged and the 'flexible stem hypothesis' may in

essence reflects a simple model of 'underdominance' of the heterozygote.

Conversely, phenotypic plasticity can be gained by any mechanism that increases

genetic variability and heterozygosity in a population, such as mutations, outbreeding (e.g.,

gene flow and introgression), and balancing selection. Indeed, any new mutation in a di- or

polyploid organism first arises in the heterozygous state, and it has been argued that a

heterozygous advantage is essential for the spread of new mutations and counteracting the

loss by random genetic drift (Sellis et al. 2011). In cases where plasticity is adaptive, the higher

plasticity of the heterozygote can facilitate the spread of this mutation. Such adaptive plasticity

gain model is in essence an 'overdominance' model. An important consequence of the

overdominance model is that plasticity could be a mechanism to maintain genetic diversity,

i.e. rather than 'buying time' for genetic adaptation to occur, plasticity may be a major source

for 'saving time' by maintaining standing genetic variation, such as in the case of introduced

species or multivoltine species inhabiting a heterogeneous spatial or seasonal environment

(Gulisija et al. 2016; Wittmann et al. 2017; Promy et al. 2023). Further theoretical and empirical

investigations of the link between heterozygosity and plasticity will definitively improve our

understanding of the evolution of both.

Acknowledgments

The authors thank Artyom Kopp for comments on the manuscript and for sharing with Yuichi

Fukutomi the association between female-limited color dimorphism and polymorphism at the

pdm3 locus in *D. jambulina* and *D.* cf. *bocqueti*.

References

Bastide, H., A. Betancourt, V. Nolte, R. Tobler, P. Stöbe, A. Futschik, and C. Schlötterer.

2013. A Genome-Wide, Fine-Scale Map of Natural Pigmentation Variation in Drosophila

melanogaster. PLOS Genetics 9:e1003534. Public Library of Science.

Bastide, H., J. D. Lange, J. B. Lack, A. Yassin, and J. E. Pool. 2016. A Variable Genetic

Architecture of Melanic Evolution in Drosophila melanogaster. Genetics 204:1307–1319.

Bastide, H., A. Yassin, E. J. Johanning, and J. E. Pool. 2014. Pigmentation in Drosophila

melanogaster reaches its maximum in Ethiopia and correlates most strongly with ultra-violet

radiation in sub-Saharan Africa. BMC Evolutionary Biology 14:179.

Berardi, S., J. A. Rhodes, M. C. Berner, S. I. Greenblum, M. C. Bitter, E. L. Behrman, N. J.

Betancourt, A. O. Bergland, D. A. Petrov, S. Rajpurohit, and P. Schmidt. 2024. Drosophila

melanogaster pigmentation demonstrates adaptive phenotypic parallelism but genomic

unpredictability over multiple timescales. bioRxiv.

Bradshaw, A. D. 1965. Evolutionary Significance of Phenotypic Plasticity in Plants. Pp. 115–

155 *in* E. W. Caspari and J. M. Thoday, eds. Advances in Genetics. Academic Press.

Bradshaw, A. D. 2006. Unravelling phenotypic plasticity – why should we bother? New

Phytologist 170:644–648.

Brisson, J. A., D. C. D. Toni, I. Duncan, and A. R. Templeton. 2005. Abdominal

Pigmentation Variation in Drosophila Polymorpha: Geographic Variation in the Trait, and

Underlying Phylogeography. Evolution 59:1046–1059.

Castro, S. D., F. Peronnet, J.-F. Gilles, E. Mouchel-Vielh, and J.-M. Gibert. 2018. bric à brac

(bab), a central player in the gene regulatory network that mediates thermal plasticity of

pigmentation in Drosophila melanogaster. PLOS Genetics 14:e1007573. Public Library of

Science.

Chakir, M., A. Chafik, P. Gibert, and J. R. David. 2002. Phenotypic plasticity of adult size

and pigmentation in *Drosophila*: thermosensitive periods during development in two sibling

species. Journal of Thermal Biology 27:61–70.

Chen, J., V. Nolte, and C. Schlötterer. 2015. Temperature Stress Mediates Decanalization and

Dominance of Gene Expression in Drosophila melanogaster. PLOS Genetics 11:e1004883.

Public Library of Science.

Chevin, L.-M., C. Leung, A. Le Rouzic, and T. Uller. 2022. Using phenotypic plasticity to

understand the structure and evolution of the genotype–phenotype map. Genetica 150:209–

221.

Colinet, H., and A. Kustre. 2025. The apparent seasonal biphenism in stems in reality from continuous reaction norms. Pest Management Science 81:507–517.

David, J. R., R. Allemand, P. Capy, M. Chakir, P. Gibert, G. Pétavy, and B. Moreteau. 2004. Comparative life histories and ecophysiology of Drosophila melanogaster and D. simulans. Genetica 120:151–163.

David, J. R., P. Capy, and J.-P. Gauthier. 1990. Abdominal pigmentation and growth temperature in Drosophila melanogaster: Similarities and differences in the norms of reaction of successive segments. Journal of Evolutionary Biology 3:429–445.

David, J. R., E. A. Ferreira, L. Jabaud, D. Ogereau, H. Bastide, and A. Yassin. 2022. Evolution of assortative mating following selective introgression of pigmentation genes between two Drosophila species. Ecology and Evolution 12:e8821.

Dembeck, L. M., W. Huang, M. M. Magwire, F. Lawrence, R. F. Lyman, and T. F. C. Mackay. 2015. Genetic Architecture of Abdominal Pigmentation in Drosophila melanogaster. PLOS Genetics 11:e1005163. Public Library of Science.

Diamond, S. E., and R. A. Martin. 2021. Buying Time: Plasticity and Population Persistence. P. *in* Phenotypic Plasticity & Evolution. CRC Press.

Dupont, L., M. Thierry, L. Zinger, D. Legrand, and S. Jacob. 2024. Beyond reaction norms: the temporal dynamics of phenotypic plasticity. Trends in Ecology & Evolution 39:41–51. Elsevier.

Durand, E., R. Méheust, M. Soucaze, P. M. Goubet, S. Gallina, C. Poux, I. Fobis-Loisy, E. Guillon, T. Gaude, A. Sarazin, M. Figeac, E. Prat, W. Marande, H. Bergès, X. Vekemans, S. Billiard, and V. Castric. 2014. Dominance hierarchy arising from the evolution of a complex small RNA regulatory network. Science 346:1200–1205. American Association for the Advancement of Science.

Endler, L., A. J. Betancourt, V. Nolte, and C. Schlötterer. 2016. Reconciling differences in

pool-GWAS between populations: a case study of female abdominal pigmentation in

Drosophila melanogaster. Genetics 202:843–855. Oxford University Press.

Ford, E. B. 1945. Polymorphism. Biological Reviews 20:73–88.

Freoa, L., L.-M. Chevin, P. Christol, S. Méléard, M. Rera, A. Véber, and J.-M. Gibert. 2023.

Drosophilids with darker cuticle have higher body temperature under light. Sci Rep 13:3513.

Nature Publishing Group.

Fukutomi, Y., A. Phillips-Garcia, J. Liu, A. Chuang, M. Watada, S. Ramniwas, and A. Kopp.

2025. Evolution of dominance in a Mendelian trait is linked to the evolution of environmental

plasticity. submitted.

Gerasimavicius, L., B. J. Livesey, and J. A. Marsh. 2022. Loss-of-function, gain-of-function

and dominant-negative mutations have profoundly different effects on protein structure. Nat

Commun 13:3895. Nature Publishing Group.

Ghalambor, C. K., K. L. Hoke, E. W. Ruell, E. K. Fischer, D. N. Reznick, and K. A. Hughes.

2015. Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in

nature. Nature 525:372–375. Nature Publishing Group.

Gibert, J.-M. 2017. The flexible stem hypothesis: evidence from genetic data. Dev Genes

Evol 227:297–307.

Gibert, J.-M., J. Blanco, M. Dolezal, V. Nolte, F. Peronnet, and C. Schlötterer. 2017a. Strong

epistatic and additive effects of linked candidate SNPs for Drosophila pigmentation have

implications for analysis of genome-wide association studies results. Genome Biol 18:126.

Gibert, J.-M., E. Mouchel-Vielh, S. D. Castro, and F. Peronnet. 2016. Phenotypic Plasticity

through Transcriptional Regulation of the Evolutionary Hotspot Gene tan in Drosophila

melanogaster. PLOS Genetics 12:e1006218. Public Library of Science.

Gibert, J.-M., E. Mouchel-Vielh, and F. Peronnet. 2017b. Modulation of yellow expression

contributes to thermal plasticity of female abdominal pigmentation in Drosophila

melanogaster. Sci Rep 7:43370. Nature Publishing Group.

Gibert, P., P. Capy, A. Imasheva, B. Moreteau, J. P. Morin, G. Pétavy, and J. R. David.

2004a. Comparative Analysis of Morphological Traits Among Drosophila Melanogaster and

D. Simulans: Genetic Variability, Clines and Phenotypic Plasticity. Genetica 120:165–179.

Gibert, P., B. Moreteau, and J. R. David. 2000. Developmental constraints on an adaptive

plasticity: reaction norms of pigmentation in adult segments of Drosophila melanogaster.

Evolution & Development 2:249–260.

Gibert, P., B. Moreteau, and J. R. David. 2009. Phenotypic plasticity of abdomen

pigmentation in two geographic populations of Drosophila melanogaster: male–female

comparison and sexual dimorphism. Genetica 135:403–413.

Gibert, P., B. Moreteau, and J. R. David. 1999. Phenotypic plasticity of abdominal

pigmentation in Drosophila kikkawai: multiple interactions between a major gene, sex,

abdomen segment and growth temperature. Genetica 105:165–176.

Gibert, P., B. Moreteau, and J. R. David. 2004b. Phenotypic plasticity of body pigmentation

in Drosophila melanogaster: Genetic repeatability of quantitative parameters in two

successive generations. Heredity 92:499–507. Nature Publishing Group.

Gibert, P., B. Moreteau, J. R. David, and S. M. Scheiner. 1998. Describing the evolution of

reaction norm shape: body pigmentation in Drosophila. Evolution 52:1501–1506.

Gibert, P., B. Moreteau, J.-C. Moreteau, and J. R. David. 1996. Growth Temperature and

Adult Pigmentation in Two Drosophila Sibling Species: An Adaptive Convergence of

Reaction Norms in Sympatric Populations? Evolution 50:2346–2353.

Gillespie, J. H., and M. Turelli. 1989. Genotype-environment interactions and the

maintenance of polygenic variation. Genetics 121:129–138.

Goldstein, I., and I. M. Ehrenreich. 2021. Genetic Variation in Phenotypic Plasticity. P. *in*

Phenotypic Plasticity & Evolution. CRC Press.

Grieshop, K., and G. Arnqvist. 2018. Sex-specific dominance reversal of genetic variation for fitness. PLOS Biology 16:e2006810. Public Library of Science.

Grieshop, K., E. K. H. Ho, and K. R. Kasimatis. 2024. Dominance reversals: the resolution of genetic conflict and maintenance of genetic variation. Proceedings of the Royal Society B: Biological Sciences 291:20232816. Royal Society.

Gulisija, D., Y. Kim, and J. B. Plotkin. 2016. Phenotypic Plasticity Promotes Balanced Polymorphism in Periodic Environments by a Genomic Storage Effect. Genetics 202:1437–1448.

Harnly, M. H., and M. L. Harnly. 1936. The effects of the gene on growth and differentiation as shown by the temperature responses of pennant and its heterozygote in D. melanogaster. Journal of Experimental Zoology 74:41–59.

Hughes, J., Williams ME, Rebeiz M, and Williams TM. 2023. Widespread cis- and trans-regulatory evolution underlies the origin, diversification, and loss of a sexually dimorphic fruit fly pigmentation trait. J Exp Zool B Mol Dev Evol 340:143–161. Wiley-Liss, United States.

Jeong, S., M. Rebeiz, P. Andolfatto, T. Werner, J. True, and S. B. Carroll. 2008. The Evolution of Gene Regulation Underlies a Morphological Difference between Two Drosophila Sister Species. Cell 132:783–793. Elsevier.

Jeong, S., A. Rokas, and S. B. Carroll. 2006. Regulation of Body Pigmentation by the Abdominal-B Hox Protein and Its Gain and Loss in Drosophila Evolution. Cell 125:1387–1399. Elsevier.

Kopp, A., I. Duncan, and S. B. Carroll. 2000. Genetic control and evolution of sexually dimorphic characters in Drosophila. Nature 408:553–559. Nature Publishing Group.

Kovuri, P., A. Yadav, and H. Sinha. 2023. Role of genetic architecture in phenotypic

plasticity. Trends in Genetics 39:703–714. Elsevier.

Lachaise, D., M. Harry, M. Solignac, F. Lemeunier, V. Bénassi, and M.-L. Cariou. 2000.

Evolutionary novelties in islands: Drosophila santomea, a new melanogaster sister species

from São Tomé. Proceedings of the Royal Society of London. Series B: Biological Sciences

267:1487–1495. Royal Society.

Lafuente, E., F. Alves, J. G. King, C. M. Peralta, and P. Beldade. 2021. Many ways to make

darker flies: Intra- and interspecific variation in Drosophila body pigmentation components.

Ecology and Evolution 11:8136–8155.

Lafuente, E., D. Duneau, and P. Beldade. 2018. Genetic basis of thermal plasticity variation in

Drosophila melanogaster body size. PLOS Genetics 14:e1007686. Public Library of Science.

Lafuente, E., D. Duneau, and P. Beldade. 2024. Genetic basis of variation in thermal

developmental plasticity for Drosophila melanogaster body pigmentation. Molecular Ecology

33:e17294.

Le Poul, Y., A. Whibley, M. Chouteau, F. Prunier, V. Llaurens, and M. Joron. 2014.

Evolution of dominance mechanisms at a butterfly mimicry supergene. Nat Commun 5:5644.

Nature Publishing Group.

Lerner, I. M. 1954. Genetic homeostasis. Oliver and Boyd. London.

Liu, Y., M. Ramos-Womack, C. Han, P. Reilly, K. L. Brackett, W. Rogers, T. M. Williams, P.

Andolfatto, D. L. Stern, and M. Rebeiz. 2019. Changes throughout a Genetic Network Mask

the Contribution of Hox Gene Evolution. Current Biology 29:2157-2166.e6. Elsevier.

Llopart, A., D. Lachaise, and J. A. Coyne. 2005. An Anomalous Hybrid Zone in Drosophila.

Evolution 59:2602–2607.

Lobell, A. S., R. R. Kaspari, Y. L. Serrano Negron, and S. T. Harbison. 2017. The Genetic

Architecture of Ovariole Number in Drosophila melanogaster: Genes with Major,

Quantitative, and Pleiotropic Effects. G3 Genes|Genomes|Genetics 7:2391–2403.

Mackay, T. F. C., and R. F. Lyman. 2005. Drosophila bristles and the nature of quantitative

genetic variation. Philosophical Transactions of the Royal Society B: Biological Sciences

360:1513–1527. Royal Society.

Massey, J. H., and P. J. Wittkopp. 2016. The Genetic Basis of Pigmentation Differences

Within and Between *Drosophila* Species. Pp. 27–61 *in* V. Orgogozo, ed. Current Topics in

Developmental Biology. Academic Press.

Michie, L. J., F. Mallard, M. E. N. Majerus, and F. M. Jiggins. 2010. Melanic through nature

or nurture: genetic polymorphism and phenotypic plasticity in Harmonia axyridis. Journal of

Evolutionary Biology 23:1699–1707.

Negoua, H., M. Chakir, J. R. David, and P. Capy. 2019. Climatic adaptation in Drosophila:

phenotypic plasticity of morphological traits along a seasonal cycle. Annales de la Société

entomologique de France (N.S.) 55:48–60. Taylor & Francis.

Orteu, A., and C. D. Jiggins. 2020. The genomics of coloration provides insights into adaptive

evolution. Nat Rev Genet 21:461–475. Nature Publishing Group.

Parkash, R., and A. K. Sharma. 1978. Colour dimorphism in Drosophila jambulina.

Drosophila Information Service 166.

Payant, V. 1986. Le polymorphisme de la coloration abdominale dans le genre Drosophila.

L'Annee biologique. Comptes Rendus Annuels des Travaux de Biologie Generale 25:167–

184.

Pétavy, G., B. Moreteau, J. R. David, and P. Gibert. 2018. Influence of extreme heat or cold

stresses on body pigmentation of *Drosophila melanogaster*. Journal of Thermal Biology

72:118–126.

Pétavy, G., B. Moreteau, P. Gibert, and J. R. David. 2002. Phenotypic plasticity of body

pigmentation in Drosophila: influence of a developmental thermoperiodic regime in two

sibling species. Physiological Entomology 27:124–135.

Pfennig, D. W. (ed). 2021. Phenotypic Plasticity & Evolution: Causes, Consequences,

Controversies. CRC Press, Boca Raton.

Pigliucci, M. 2005. Evolution of phenotypic plasticity: where are we going now? Trends in

Ecology & Evolution 20:481–486. Elsevier.

Pigliucci, M. 1992. Modelling phenotypic plasticity. I. Linear and higher-order effects of

dominance, drift, environmental frequency and selection on a one-locus, two-allele model. J.

Genet. 71:135–150.

Pigliucci, M. 2001. Phenotypic Plasticity: Beyond Nature and Nurture. JHU Press.

Pitchers, W., J. Nye, E. J. Márquez, A. Kowalski, I. Dworkin, and D. Houle. 2019. A

Multivariate Genome-Wide Association Study of Wing Shape in Drosophila melanogaster.

Genetics 211:1429–1447.

Prigent, S. R., Lang ,Michael, Nagy ,Olga, Acurio ,Andrea, Matamoro-Vidal ,Alexis,

Courtier-Orgogozo ,Virginie, and J. R. and David. 2020. Field collections reveal that São

Tomé is the Afrotropical island with the highest diversity of drosophilid flies (Diptera:

Drosophilidae). Annales de la Société entomologique de France (N.S.) 56:1–14. Taylor &

Francis.

Promy, N. T., M. Newberry, and D. Gulisija. 2023. Rapid evolution of phenotypic plasticity

in patchy habitats. Sci Rep 13:19158. Nature Publishing Group.

R Core Team. 2023. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria.

Rocha, F., H. F. Medeiros, and L. B. Klaczko. 2009. The reaction norm of abdominal

pigmentation and its curve in Drosophila mediopunctata depend on the mean phenotypic

value. Evolution 63:280–287.

Rogers, W. A., S. Grover, S. J. Stringer, J. Parks, M. Rebeiz, and T. M. Williams. 2014. A

survey of the trans-regulatory landscape for *Drosophila melanogaster* abdominal

pigmentation. Developmental Biology 385:417–432.

Sapir, Y., M. K. Gallagher, and E. Senden. 2021. What Maintains Flower Colour Variation

within Populations? Trends in Ecology & Evolution 36:507–519. Elsevier.

Scheiner, S. M. 1993. Genetics and Evolution of Phenotypic Plasticity. Annual Review of

Ecology, Evolution, and Systematics 24:35–68. Annual Reviews.

Schmalhausen, I. 1949. Factors of evolution: the theory of stabilizing selection. Blakiston.

Toronto, Canada.

Sellis, D., B. J. Callahan, D. A. Petrov, and P. W. Messer. 2011. Heterozygote advantage as a

natural consequence of adaptation in diploids. Proceedings of the National Academy of

Sciences 108:20666–20671. Proceedings of the National Academy of Sciences.

Sievert, C. 2020. Interactive Web-Based Data Visualization with R, plotly, and shiny.

Chapman and Hall/CRC, New York.

Sultan, S. E. 2021. Phenotypic Plasticity as an Intrinsic Property of Organisms. P. *in*

Phenotypic Plasticity & Evolution. CRC Press.

Suvorov, A., B. Y. Kim, J. Wang, E. E. Armstrong, D. Peede, E. R. R. D'Agostino, D. K.

Price, P. J. Waddell, M. Lang, V. Courtier-Orgogozo, J. R. David, D. Petrov, D. R. Matute, D.

R. Schrider, and A. A. Comeault. 2022. Widespread introgression across a phylogeny of 155

Drosophila genomes. Current Biology 32:111-123.e5.

Takahashi, Y., K. Kagawa, E. I. Svensson, and M. Kawata. 2014. Evolution of increased

phenotypic diversity enhances population performance by reducing sexual harassment in

damselflies. Nat Commun 5:4468. Nature Publishing Group.

Telonis-Scott, M., A. A. Hoffmann, and C. M. Sgrò. 2011. The molecular genetics of clinal

variation: a case study of ebony and thoracic trident pigmentation in Drosophila melanogaster

from eastern Australia. Molecular Ecology 20:2100–2110.

Tian, S., Y. Asano, T. Das Banerjee, S. Komata, J. L. Q. Wee, A. Lamb, Y. Wang, S. N.

Murugesan, H. Fujiwara, K. Ui-Tei, P. J. Wittkopp, and A. Monteiro. 2024. A microRNA is the effector gene of a classic evolutionary hotspot locus. Science 386:1135–1141. American Association for the Advancement of Science.

Tichy, A. L., A. Ray, and J. R. Carlson. 2008. A New Drosophila POU Gene, pdm3, Acts in Odor Receptor Expression and Axon Targeting of Olfactory Neurons. J. Neurosci. 28:7121–7129. Society for Neuroscience.

Turissini, D. A., and D. R. Matute. 2017. Fine scale mapping of genomic introgressions within the Drosophila yakuba clade. PLOS Genetics 13:e1006971. Public Library of Science.

van der Burg, K. R. L., J. J. Lewis, B. J. Brack, R. A. Fandino, A. Mazo-Vargas, and R. D. Reed. 2020. Genomic architecture of a genetically assimilated seasonal color pattern. Science 370:721–725. American Association for the Advancement of Science.

Watanabe, T. K., M. Matsuda, S. Ohnishi, and F. Hihara. 1982. Notes on the systematics of Drosophila jambulina. The Japanese journal of genetics 57:561–567.

Wennersten, L., and A. Forsman. 2012. Population-level consequences of polymorphism, plasticity and randomized phenotype switching: a review of predictions. Biological Reviews 87:756–767.

West-Eberhard, M. J. 2003. Developmental Plasticity and Evolution. Oxford University Press.

Wittkopp, P. J., E. E. Stewart, L. L. Arnold, A. H. Neidert, B. K. Haerum, E. M. Thompson, S. Akhras, G. Smith-Winberry, and L. Shefner. 2009. Intraspecific Polymorphism to Interspecific Divergence: Genetics of Pigmentation in Drosophila. Science 326:540–544. American Association for the Advancement of Science.

Wittmann, M. J., A. O. Bergland, M. W. Feldman, P. S. Schmidt, and D. A. Petrov. 2017. Seasonally fluctuating selection can maintain polymorphism at many loci via segregation lift. Proceedings of the National Academy of Sciences 114:E9932–E9941. Proceedings of the National Academy of Sciences.

Woltereck, R. 1909. Weitere experimentelle Untersuchungen über Artveränderung, speziel

über das Wesen quantitativer Artunterschiede bei Daphniden. erhandlungen der deutschen

zoologischen Gesellschaft 19:110–173.

Yassin, A. 2018. Phylogenetic biogeography and classification of the *Drosophila montium*

species group (Diptera: Drosophilidae). Annales de la Société entomologique de France

(N.S.) 54:167–175.

Yassin, A., H. Bastide, H. Chung, M. Veuille, J. R. David, and J. E. Pool. 2016a. Ancient

balancing selection at tan underlies female colour dimorphism in Drosophila erecta. Nat

Commun 7:10400. Nature Publishing Group.

Yassin, A., E. K. Delaney, A. J. Reddiex, T. D. Seher, H. Bastide, N. C. Appleton, J. B. Lack,

J. R. David, S. F. Chenoweth, and J. E. Pool. 2016b. The pdm3 locus is a hotspot for recurrent

evolution of female-limited color dimorphism in Drosophila. Current Biology 26:2412–2422.

Elsevier.

Yassin, A., N. Gidaszewski, V. Debat, and J. R. David. 2022. Long-term evolution of

quantitative traits in the Drosophila melanogaster species subgroup. Genetica 150:343–353.