Social experience alters behaviors by reprogramming the Fruitless pathway and circadian state in *Drosophila*

Chengcheng Du [1], Sumie Okuwa [1], Lanling Jia [1], Marta Rozados Barreiro [1], Shayna Scott [1], Jesús

Emiliano Sotelo Fonseca [1], Yuta Mabuchi [3,5], Shania Appadoo [1], Luis Garcia [1], Stephanie Rohrbach [1],

Sare Koruk [1], Efe Balkanli [1], Josh Lin [1], Nilay Yapici [3], Corbin D. Jones [4], and Pelin C. Volkan [1, 2†]

[1]Department of Biology, Duke University, Durham, NC 27708, USA

[2]Department of Neurobiology, Duke University Medical Center, Durham, NC 27710, USA

[3]Department of Neurobiology and Behavior, Cornell University, Ithaca, NY 14853, UAS

[4]Department of Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA

[5]Current Address: Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY,

10027-2325, USA

[†]Corresponding author: pelin.volkan@duke.edu

4341 French Family Science Center

124 Science Drive

Campus Box 90338, Duke University

Durham, NC 27708, USA

Abstract

From flies to humans, social experience affects various cognitive and behavioral processes. Previous studies have shown that group housing suppresses many behaviors like courtship, aggression, and feeding in *Drosophila melanogaster,* in addition to resetting the circadian state. Here, we focus on group housing-induced courtship suppression. To determine the mechanisms by which social experience modulates courtship behaviors, we performed bulk tissue RNAseq and single-cell RNAseq from cells expressing Fruitless^M (Fru^M) and Doublesex^M (Dsx^M), two transcription factors that label interconnected neural circuits for socially driven behaviors, from grouped or isolated male brains. These revealed that social isolation alters *fru* and *dsx* levels throughout the brain. Knocking down *fru^M* in different *fru^M*-positive neuron subpopulations in the brains has diverse effects on social experience-dependent changes in courtship. Furthermore, group housing increases the expression of *stripe* (*sr*) and *Hormone receptor-like in 38* (*Hr38*) genes encoding neural activity-induced transcription factors in most neurons within social circuits. We found that knocking down *sr* in *fru^M*-positive neurons effectively eliminates the impact of social experience by increasing courtship in group-housed males. Importantly, social experience also alters the expression of Fru^M/Dsx^M putative target genes regulating circadian states throughout the brain. Disrupting the function of multiple circadian genes diminishes the effect of group housing on courtship. Our findings suggest that group housing/social enrichment suppresses courtship by reprogramming the circadian arousal state, whereas courtship-elevating effects of social experience rely on unique influences of Fru^M expression and function in different neurons within social and clock circuits. These results are significant as they point to modulation of circadian arousal state as a possible central strategy for mediating the pleiotropic effects of social experience on organismal responses.

Introduction

Social experience, particularly social isolation or loneliness, is known to have long-lasting effects on behavior and physiology in animals ranging from mice to flies (Anders, 1978; Cacioppo et al., 2006; de Melo Sene, 1977; Hoffman et al., 1975; Sánchez et al., 1998). Over the past decade, growing awareness of these effects—exacerbated by global health crises—has led to the recognition of social isolation and loneliness as a new epidemic, as declared by the Surgeon General (General, 2023; Jeste et al., 2020; Lin, 2023). Adverse social experience, including isolation, disrupts various cognitive processes, including sleep, learning, memory, and attention, while also increasing behaviors such as aggression and feeding (Brown et al., 2017; Cacioppo and Hawkley, 2009; Cacioppo et al., 2006; Cohen, 2004; Liu et al., 2025; Zelikowsky et al., 2018). Consequently, in humans, isolation is linked to a rise in neuropsychiatric and neurodegenerative diseases (Cacioppo et al., 2015; Cacioppo et al., 2006; Cohen, 2004). The molecular and circuit-based mechanisms of social experience-dependent behavioral and physiological effects remain unclear.

Social experience similarly shapes physiology and behavior across animal species. Animals modulate their behaviors by signals from the social environment and prior social experience (Anders, 1978; de Melo Sene, 1977; Eban-Rothschild and Bloch, 2015; Luciano and Lore, 1975; Matsumoto et al., 2005; Ueda and Kidokoro, 2002; Wang et al., 2022). Emerging evidence from both vertebrate and invertebrate models demonstrates that social environment exerts profound effects on the regulation of genes critical for maintaining neuronal homeostasis, physiological function, and synaptic transmission, ultimately shaping neural circuit activity and behavioral outputs (Cushing and Kramer, 2005; Deanhardt et al., 2023; Flavell and Greenberg, 2008; West and Greenberg, 2011; Zhao et al., 2020). Despite this progress, how the detection of social cues by peripheral sensory systems alters gene expression throughout central circuits to ultimately modify circuit function and behaviors remains poorly understood.

Exploring molecular and circuit mechanisms by which social experience modulates behaviors requires a well-defined model system with causal links between genes, circuits, and behavioral function. The genetically tractable fruit fly is an excellent model because of its adaptable social behaviors, simple nervous system with a widely mapped connectome, and known master regulatory genes for circuit structure and function. Beyond the general advantages of *Drosophila* as a model organism, its pronounced sensitivity to social experience makes it particularly valuable for identifying genes and circuits driving responses to social experience. Previous studies reported that social isolation elevates many fundamental behaviors, like feeding, aggression, and courtship, while disrupting cognitive processes such as sleep, learning, and memory in fruit flies (Agrawal et al., 2020; Dankert et al., 2009; Ganguly-Fitzgerald et al., 2006; Goncharova et al., 2016; Kacsoh et al., 2018; Li et al., 2021; Ueda and Kidokoro, 2002; Wang et al., 2008). Social experience also resets the circadian state and alters the activity of the clock neurons in *Drosophila* (Kim et al., 2013; Levine et al., 2002), which might drive the pleiotropic effects on diverse behavioral and physiological responses. Socially isolated flies sleep less than grouped flies, and this is mediated by contributions from dopaminergic modulation, cAMP signaling, and memory-related genes (Ganguly-Fitzgerald et al., 2006; Li et al., 2021). The influence of social experience extends to alterations in social behaviors such as courtship and aggression. *Drosophila melanogaster* males raised in groups display less intense courtship behavior toward female targets than isolated males (Dankert et al., 2009; Goncharova et al., 2016). In addition, *Drosophila melanogaster* males with prior exposure to mated females learn to reject mated females compared to males without such experience (Dukas, 2005). Social isolation has been shown to consistently increase aggression in various *Drosophila* species, both in males and females, including *D. melanogaster* (Agrawal et al., 2020; Liu et al., 2011; Ueda and Kidokoro, 2002; Ueda and Wu, 2009; Wang et al., 2008) and *D. suzukii* (Belenioti and Chaniotakis, 2020). The shared

effects of social experience, combined with an unmatched set of genetic tools for labeling and
manipulating neurons, make *Drosophila* an excellent model to explore molecular and circuit mechanisms
by which social experience alters gene expression, circuit function, and behaviors.

Genes and circuits regulating *Drosophila* social behaviors, like courtship and aggression, have been well
established. Two transcription factors, Fruitless (Fru) and Doublesex (Dsx), play crucial roles in
governing the development and function of social circuits in *Drosophila*. Fru establishes most of the
circuits mediating courtship and aggression (Anand et al., 2001; Billeter et al., 2006; Dickson, 2008;
Koganezawa et al., 2016; Neville et al., 2014; Vrontou et al., 2006; Yamamoto and Koganezawa, 2013).
The *fruitless* locus generates multiple alternative splice isoforms for RNAs transcribed from seven
promoters (Billeter et al., 2006; Dalton et al., 2013; Demir and Dickson, 2005; Goodwin et al., 2000; Lee
et al., 2000; Neville et al., 2014; von Philipsborn et al., 2014). The transcripts from *fru* P1 promoter are
alternatively spliced between males and females by Transformer (Tra), where the male isoforms (fru^M)
encode functional proteins, whereas female isoforms (fru^F) do not produce any protein product (Goodwin
et al., 2000; Lee et al., 2000). Other *fru* promoters produce common isoforms in both male and female
circuitry (Goodwin et al., 2000; Lee et al., 2000). Fru^M is expressed in an interconnected neural network
throughout both the peripheral and central nervous system, and its expression is required for the
development, function, and plasticity of the circuit that drives male-specific behaviors (Billeter et al.,
2006; Dalton et al., 2013; Demir and Dickson, 2005; Goodwin et al., 2000; Lee et al., 2000; Neville et al.,
2014; von Philipsborn et al., 2014). Owing to the function of Fru^M in establishing the structural and
functional dimorphism of social circuits, it is unsurprising that Fru^M is both necessary and sufficient for
sex-specific behaviors like male courtship.

A second transcription factor, Dsx, is essential for sex specification and differentiation in various tissues
(Burtis and Baker, 1989; Dauwalder, 2011; Hildreth, 1965; Rideout et al., 2010; Villella and Hall, 1996).
Unlike *fru,* which exhibits a high degree of evolutionary divergence, *dsx* is molecularly and functionally
conserved across invertebrates and vertebrates (Clough et al., 2014; Kopp, 2012; Murphy et al., 2007;
Rideout et al., 2010). Dsx belongs to the DMRT (Doublesex and Mab-3 Related Transcription factor)
family, which is involved in regulating sex-specific differentiation (Kopp, 2012; Murphy et al., 2007).
Doublesex is also differentially spliced in males (dsx^M) and females (dsx^F) in *Drosophila*, and shows
sexually dimorphic expression in the central brain (Burtis and Baker, 1989; Dauwalder, 2011; Hildreth,
1965; Rideout et al., 2010; Villella and Hall, 1996; Waterbury et al., 1999), albeit in fewer neurons
compared to fru^M. Disrupting Dsx function or blocking the activity of *dsx*-positive neurons alters
courtship in both males and females (Han et al., 2022; Pan and Baker, 2014; Rideout et al., 2010; Zhou et
al., 2014). Dsx^M also mediates social experience-dependent courtship learning exhibited by fru^M mutant
males, suggesting Fru^M drives innate courtship behaviors, while Dsx^M has a more modulatory effect that
can be influenced by social experience in the absence of Fru^M (Pan and Baker, 2014). This is likely driven
by the redundant function of Dsx^M and Fru^M both of which are expressed in a set of central courtship
command neurons (P1 neurons) in males, which are causally linked to initiation of courtship (Inagaki et
al., 2014; Kimura et al., 2008; Pan et al., 2012; Von Philipsborn et al., 2011).

Previous studies have shown that social context reprograms the expression and chromatin state of *fru* and
dsx in pheromone-sensing neurons and alters the responses of these neurons to modify behaviors in a
fru^M-dependent manner (Deanhardt et al., 2023; Hueston et al., 2016; Sethi et al., 2019; Zhao et al., 2020).
These findings highlight an intimate connection between social experience, chromatin remodeling,
transcriptional modulation, and neuronal activity in the peripheral nervous system. Yet, it remains unclear
how gene expression and neuronal activity are regulated in the central brain in response to social
experience, and how these changes contribute to behavioral plasticity, particularly in male courtship.

Here, we investigated the impact of social experience on gene expression and neuronal activity in the
central nervous system and how it reprograms male courtship behaviors in *Drosophila melanogaster.* We
showed that group housing suppresses both evoked neural responses in courtship circuits and male
courtship vigor, in agreement with previous research causally linking the activity of P1 neurons and
courtship behaviors (Inagaki et al., 2014; Kimura et al., 2008; Pan et al., 2012). To identify differentially
expressed genes responding to social experience, we carried out bulk RNAseq from the whole brain and
single-cell RNAseq from fru^M- and dsx^M-positive neurons in the brains of grouped and isolated males,
which revealed three major findings: 1) Social isolation increases fru^M and dsx^M reporter expression, the
expression of *fru* in clock neurons, and *dsx* throughout the brain. Manipulating Fru^M function in different
neuronal populations within social circuits has differential effects on social experience dependent
modulation of courtship. 2) Group housing increases the expression of immediate early genes like *sr* and
Hr38 throughout the neurons within social circuits. The knockdown of *sr* in *fru*-positive neurons
derepresses courtship in group-housed males to levels seen in isolated males. 3) Social experience alters
the expression of Fru^M/Dsx^M target genes encoding circadian regulators (*i.e. Clk*, *cry*) and regulators of
male mating rhythm (*Sik3*) in the brain, which causally drive courtship responses in group-housed males.
Overall, our results suggest that social experience reprograms Fru^M transcriptional cascades and their
downstream target circadian genes within the social circuits to modulate courtship vigor.

Results

Social experience induces plastic changes in male fruit fly courtship behaviors

To assess how social experience influences male courtship behaviors, we examined the courtship activity
of wild-type *Canton-S* (*CS*) male flies that were either group-housed (GH) or single-housed (SH) for
seven days starting right after eclosion (Fig. 1). Males kept in groups exhibited significantly lower
courtship intensity toward virgin females and exhibited suppressed P1 courtship command neuron activity
compared to those raised in isolation (Fig. 1A-D; Supplementary Fig. 1B-D). Notably, the reduced
courtship vigor in GH males was primarily due to an increased latency of the courtship onset (Fig. 1D).
Once aroused, both GH and SH males courted females actively and persistently (Fig. 1A-D;
Supplementary Fig. 1A).

To determine whether these socially induced behavioral changes are reversible, we tested the courtship
behaviors in four different conditions: continuous GH, GH followed by SH (GH > SH), SH followed by
GH (SH > GH), and continuous SH, across 7- and 14-day durations, with the switch occurring at 3.5 days
and 7 days, respectively (Fig. 1A-D; Supplementary Fig. 1A). Across both time frames, males housed
entirely in groups consistently courted less vigorously than those housed entirely in isolation (Fig. 1A-D;
Supplementary Fig. 1A). When males were transferred from GH to SH (GH > SH), their courtship
behavior shifted to resemble that of SH males in both 7 and 14 days (Fig. 1A-D; Supplementary Fig. 1A),
suggesting a restoration of courtship drive upon social isolation. Conversely, the effects were more
complex when flies were first isolated and then grouped (SH > GH). We found that courtship is
suppressed in males housed in groups compared to isolation for either 7 or 14 days. Switching from SH to
GH halfway through the 7-day housing duration led to a slight decline in courtship activity (Fig. 1A-D;
Supplementary Fig. 1A). Yet, switching to GH on day 7 led to courtship levels resembling GH levels by
the end of 14 days (Fig. 1A-D; Supplementary Fig. 1A). This indicates that the effect of isolation is more
dominant and that the suppressive effect of group housing accumulates over time. These results overall
point to the reversible impact of social experience on courtship behavior, with isolation exerting a
stronger enhancing effect and prolonged grouping exerting a more substantial suppressive effect.

To further examine the effect of population density on courtship behavior, we compared the courtship
index of 7-day-old isolated males with that of males raised in groups of 5, 10, 15, 20, or 30
(Supplementary Fig. 1E). The results showed that even the smallest group size (5 males) was sufficient to
suppress courtship behavior, and increasing group size beyond that did not produce significantly stronger
suppression (Supplementary Fig. 1E). This suggests that social cues from a relatively small number of
conspecifics are enough to modulate male courtship behavior.

190 Fru^M, but not Dsx^M, mediates social experience-dependent changes in courtship
191   vigor

Fru^M and Dsx^M are transcription factors in the sex-determination pathway and regulate the development
and function of circuits for male courtship behaviors (Dauwalder, 2011; Rideout et al., 2010; Ryner et al.,
1996). In the central brain, fru^M is expressed throughout interconnected neurons, including the P1
courtship command neurons that co-express both fru^M and dsx^M (Kimura et al., 2008; Lee et al., 2000; Pan
et al., 2012). Given the important function of Fru^M and Dsx^M in regulating courtship, we first investigated
whether fru^M or dsx^M is required in P1 neurons for the social experience-dependent modulation of
courtship. We performed RNAi knockdown of fru^M or dsx^M in P1 neurons using an intersectional
stringent driver (*P1-spGAL4*, Fig. 1E, see Methods for genotypes) and a broader P1 driver (*R71G01-*
GAL4, Fig. 1F). While fru^M knockdown with the stringent driver did not alter courtship responses to
social experience, knockdown with the broader P1 driver increased courtship in GH males to SH levels
(Fig. 1E and F). Furthermore, knocking down fru^M in *fru*-positive neurons in the central brain using an
intersectional strategy (*fru* brain-GAL4, see Methods for genotypes) also eliminated the courtship
difference, this time due to reduced courtship in SH males (Fig. 1G; Supplementary Fig. 2A). This is
consistent with previous studies demonstrating that fru^M is essential for male courtship behavior (Anand
et al., 2001; Demir and Dickson, 2005; Pan and Baker, 2014; Ryner et al., 1996). Both fru^M mutants and
fru^M knockdown in all *fru+* cells result in significantly reduced courtship behaviors in both GH and SH
conditions (Supplementary Fig. 2C and D). These findings suggest that, besides its canonical roles in the
specification of male courtship behaviors, fru^M might have diverse roles in different neuronal populations
in reprogramming courtship behaviors in response to social experience.

RNAi-based knock down of dsx^M in P1 neurons—using either the stringent or the broad P1 drivers (Fig.
1E and F)—and *dsx*-positive brain neurons (Fig. 1H; Supplementary Fig. 2B) did not alter the effect of
social experience on courtship behavior. Although $dsx^{0683/1649}$ mutant males consistently courted less than
wild-type controls in agreement with previous reports (Rideout et al., 2010; Villella and Hall, 1996), GH
$dsx^{0683/1649}$ mutants still courted females less than their SH counterparts (Supplementary Fig. 2E). Thus,
while dsx^M affects normal male courtship and is necessary for keeping the high level of courtship, it might
not mediate the modulation of courtship behaviors with social experience. Alternatively, the function of
Dsx in mediating responses to social experience might reside in cells outside the brain, like head fat body
cells (Clough et al., 2014). These results suggest that the majority of the effects of social experience on
courtship are largely mediated by Fru^M but not Dsx^M.

221   Bulk and cell-type-specific single-cell RNA sequencing revealed transcriptional
222   responses to social experience

Social experience-dependent changes in circuit function and courtship behaviors may be driven by
alterations in gene expression in response to neural activity/inactivity during chronic social enrichment or
isolation. Previous reports showed that, in the peripheral olfactory system, social experience and
pheromone signaling alter fru^M and dsx^M expression, and the expression of many Fru^M target genes,
including modulators of membrane potential (Deanhardt et al., 2023; Hueston et al., 2016; Sethi et al.,

2019; Zhao et al., 2020). These transcriptional responses to social experience are accompanied by FruM-
dependent alterations in the pheromone responses of olfactory neurons (Sethi et al., 2019).

To identify differentially expressed genes in response to social experience, we performed bulk RNA
sequencing on brain tissues from 7-day-old wild-type *Canton-S* (*CS*) males raised in grouped or isolated
conditions (Fig. 2). Principal component analysis (PCA) revealed a clear separation between the two
housing conditions across three biological replicates (Fig. 2A), indicating consistent transcriptomic
differences. The expression of housekeeping genes remained stable between the two groups
(Supplementary Fig. 3). Approximately 240 genes were differentially expressed (Fig. 2B, C), with
roughly half upregulated in grouped males and the other half downregulated. Based on Gene Ontology
terms (GO terms), these differentially expressed genes (DEGs) are involved in biosynthesis, metabolism,
circadian rhythm, behavior, and defense responses (Fig. 2E-G). Notably, *dsx* showed a near-significant
trend for reduction in transcript levels in GH males (Fig. 2D). In contrast, *fru* transcript levels did not
show a significant difference between GH and SH males (Fig. 2D).

While bulk brain RNA sequencing provided insights into overall gene expression changes, it averages
across all cells, masking small but meaningful changes within specific cell populations. To explore how
social experience modulates gene expression at the cellular level, we performed single-cell RNA
sequencing (scRNA-seq) on a combined pool of fru^M+ brain cells and dsx^M+ brain cells sorted from 7-
day-old males raised under either GH or SH conditions (Fig. 3A). Using GFP reporters and fluorescence-
activated cell sorting (FACS), we selectively sorted GFP-labeled fru^M+ cells and dsx^M+ cells from male
brains (Fig. 3A). Notably, the percentages of $fru^M::GFP$ and $dsx^M::GFP$ positive cells in FACS were
consistently higher in SH males compared to GH males (Fig. 3B, Supplementary Fig.4A). When analyzed
separately, both fru^M+ and dsx^M+ populations showed an increased GFP+ cell ratio in SH male brains
(Fig. 3B, Supplementary Fig.4A).

We then sought to determine whether this increase was restricted to particular cell types or distributed
across many. At resolution 1, cells were grouped into 83 small clusters from six major clusters (Fig. 3C
and D). All clusters expressed fru^M, and some subsets expressed dsx^M (Fig. 3E, Supplementary Fig.6A
and B). Based on previously known gene markers (Davie et al., 2018; Park et al., 2022), we manually
annotated 21 clusters (Fig. 3D). Overall, the proportions of these annotated cell types were similar
between GH and SH samples, with clusters 6 and 18 showing the most significant variability across
replicates, likely due to low RNA counts (Supplementary Fig. 4). Thus, they were excluded from further
comparative analysis (Supplementary Fig. 5). Side-by-side comparisons of replicates 1 and 2 showed
mostly consistent or near clustering proportions (Supplementary Fig. 5A and C). However, comparing
proportions alone can obscure actual differences in cell abundance, as each sample's proportions are
normalized to add up to 1. To capture changes in cell numbers more accurately, we adjusted for the
percentages of GFP+ cells from FACS data and applied these scaling factors to the cluster counts
(Supplementary Fig. 5B and D). With this adjustment, many clusters from SH brains exhibited increased
counts compared to GH brains (Supplementary Fig. 5B and D). This indicates that the rise in
fru^M+/dsx^M+ cells in SH male brains results from small increases, likely in transcriptional reporter
activity, across many clusters, rather than expansion of a specific fru^M+ or dsx^M+ cell population.

Next, we conducted pseudobulk differential expression analysis across clusters to identify genes altered
by social experience. Two immediate early genes (IEGs), *Hr38* (*Hormone receptor-like in 38*) and *sr*
(*stripe*), known to respond to neural activity (Chen et al., 2016; Fujita et al., 2013; Takayanagi-Kiya et al.,
2023), showed robust upregulation in most clusters (excluding glia) in GH brains compared to SH brains
(Fig. 3E, Supplementary Fig. 6D and E). A third activity-responding gene, *cbt*, exhibited differential
expression in only a few clusters (Supplementary Fig. 6E). These trends were largely consistent with the

bulk RNA-seq data, where *Hr38* and *sr* also showed higher expression in GH brains, though *Hr38* did not
reach statistical significance (Supplementary Fig. 6C). Given that *Hr38*, *sr,* and *cbt* showed strong
upregulation in GH brains across multiple cell clusters, we asked if this contributes to courtship
suppression using RNAi-mediated knockdowns in *fru+* or *dsx+* neurons under GH conditions. We found
that *fru^{GAL4}* or *dsx^{GAL4}* driven expression of most RNAi transgenes against *Hr38* and *sr* in the brain
increased courtship compared to controls in the first round of screening (Supplementary Fig. 9A and B).
We next tested whether this increase is sufficient to eliminate the courtship difference between grouped
and isolated males by analyzing *Hr38* and *sr* RNAi knockdown phenotypes in both GH and SH
conditions with additional controls (Fig. 4). Knocking down *sr,* but not *Hr38,* in *fru^M+* neurons in the
central brain (*fru* brain-GAL4) elevated courtship in grouped males, eliminating the courtship difference
between GH and SH males (Fig. 4A). In contrast, knocking down *Hr38* or *sr* in *dsx^M+* neurons in the
brain (*dsx* brain-GAL4) did not eliminate the effect of social experience on courtship difference between
GH and SH conditions (Fig. 4B). In addition to the role of *fru^M* in social modulation of courtship, these
results also implicate transcriptional changes in immediate early genes like *sr* in *fru^M* neuronal circuits in
mediating courtship suppression in response to group housing.

As shown previously, P1 command neurons exhibit reduced evoked activity in GH males (Supplementary
Fig. 1B-D). To determine transcriptional changes in P1 neurons that might underlie this reduction in
activity, we also examined the DEGs in P1 neurons (Supplementary Fig. 10K). *Hr38* was again
upregulated in GH P1 neurons. However, knocking down *Hr38* specifically in P1 neurons using *P1-*
spGAL4 did not affect the courtship response to group housing (Supplementary Fig. 10L), suggesting that
P1 activity modulation could be driven by upstream circuit inputs rather than intrinsic transcriptional
changes. Taken together, these results suggest that upregulation of immediate early genes within social
circuits is associated with mediating the effects of group housing on courtship suppression.

RNAi screening identifies candidate genes mediating social experience-dependent courtship plasticity

Single-cell RNAseq also revealed additional DEGs in various clusters, which are listed in Supplementary
Figure 7. These DEGs are enriched in diverse biological processes, including neuronal signaling,
neuromodulation, circadian regulation, and behavior (Supplementary Fig. 7B). To determine whether the
DEGs identified in single-cell RNA-seq are causally involved in mediating social experience-dependent
changes in courtship behavior, we conducted an RNAi screen. Even though some DEGs were in
previously annotated clusters, like dopaminergic neurons or glia cells, there were still many DEGs in
unannotated cells (Fig. 3D). Therefore, we used the FindMarkers function to identify marker genes for the
four biggest clusters—excluding Cluster 1, which corresponded to mushroom body γ-lobe Kenyon cells
with few DEGs (Fig. 3, Supplementary Fig. 8A).

Based on available GAL4 lines, we selected *Kr* and *spab* from Cluster 0, *dsf* and *Fer1* from Cluster 2, and
Sox102F and *SoxN* from Cluster 3 as marker genes (Supplementary Fig. 8A). While none of these genes
are exclusive to a single cluster, they served as valid entry points. We then performed pseudobulk DEG
analysis in subsets expressing these markers, as well as in the *fru+* and *dsx+* subsets (Fig. 3F,
Supplementary Fig. 8B). DEGs in these marker-defined populations were enriched in metabolism,
signaling, and circadian processes, as well as behavior modulation (Supplementary Fig. 8C).

We used the cluster-specific marker GAL4 lines to knock down many of the DEGs and tested their effect
on social modulation of courtship (Supplementary Fig. 9 and 10). Many DEGs within the *dsx+* subset
were from *dsx+* glial cells (Supplementary Fig. 9C), whereas *dsx* brain-GAL4 does not target glia. These
genes were thus tested genetically using *repo-GAL4* driven RNAi in glial cells (Supplementary Fig. 9 and

10). While many other RNAi lines showed no effect, a few genes, such as *Sik3* and *Dh31*, emerged as
potent modulators of courtship plasticity, consistently showing elevated courtship in group-housed males
when knocked down with multiple drivers and RNAi lines (Supplementary Figs. 9 and 10). Both genes
are involved in circadian modulation of behaviors and are upregulated in specific cell clusters in the brain
in response to group housing (Fig. 3F, Supplementary Fig. 7). *Sik3* knockdown in central brain *fru+*
neurons increased courtship in GH males and dramatically decreased the behavioral difference between
GH and SH conditions (Fig. 4D). Similarly, knocking down *Sik3* in *spab+* cells also diminished the GH–
SH difference in courtship vigor by increasing courtship in grouped males (Fig. 4C). Meanwhile, even
though knocking down *Dh31* in *spab+* cells eliminated the difference between GH and SH, the
confidence was decreased due to the effects seen in controls (Fig. 4C). These findings implicate the
circadian gene *Sik3* as an essential regulator of social experience-dependent modulation of courtship
behaviors.

Circadian genes mediate the effects of group housing on courtship behaviors

Previous studies have implicated *Sik3* in regulating the circadian male sex drive rhythm (Fujii et al.,
2017), suggesting that social experience may influence internal arousal states by reprogramming the
circadian system with downstream effects on the brain and behaviors. Thus, we revisited the DEGs in the
bulk tissue RNAseq, where the GO terms centered on circadian rhythm-related processes (Fig. 2).

In the bulk brain RNAseq, three key circadian genes—*cryptochrome* (*cry*), *timeless* (*tim*), and *Clock*
(*Clk*)—exhibited significant differential expression between GH and SH males (Fig. 2C). Interestingly,
these four circadian genes (*Sik3, cry, tim* and *Clk*) are indicated as potential target genes transcriptionally
regulated by Fru^M/Dsx^M (Clough et al., 2014; Neville et al., 2014). The brain expression patterns of
circadian genes *cry*, *Clk*, and *tim* in the clock network overlap, but are not always coexpressed (Fig. 5A
and D). To confirm social experience-dependent transcriptional changes in *Clk, cry,* and *tim,* we analyzed
their expression *in vivo* using existing transcriptional reporters and protein-GFP fusions in the brain (Fig.
5A and B, also see (Du et al., 2025)). Among the three circadian genes, we found that *Clk* transcript levels
were upregulated in response to single housing (Fig. 5A and B). In contrast, we found Cry protein levels
exhibited an increase in response to group housing (Fig. 5A and B). *tim,* on the other hand, did not show
an overall difference in transcript levels (Fig. 5A and B). Overall, these results suggest that some
circadian genes are modulated by social experience.

To investigate whether changes in circadian gene expression with social experience contribute to
experience-dependent changes in courtship behavior, we examined the courtship in *cry, tim,* and *Clk*
mutants. We also investigated *Pdf* mutants, because of its coexpresssion and interaction with *cry* in a
subset of clock neurons (Cusumano et al., 2009; Zhang et al., 2009), as well as its transcriptional
downregulation in response to group housing (Supplementary Fig. 7A). Compared to wild-type *CS* males,
Clk mutants completely eliminated the difference in courtship by increasing courtship in GH males (Fig.
5C; Supplementary Fig. 11B). *tim* mutants also showed a reduced difference in courtship between GH
and SH conditions due to increased courtship in GH males (Fig. 5C; Supplementary Fig. 11B). In
contrast, *cry* and *Pdf* mutants showed an accentuated courtship difference between GH and SH males,
mostly due to further courtship suppression in GH males (Fig. 5C; Supplementary Fig. 11B). We also
manipulated neuronal activity within clock neuron populations. Silencing or stimulating neural activity
using *Clk-GAL4*-driven *UAS-Kir2.1* or *UAS-dTrpA1* expression did not alter social experience-dependent
changes in courtship behaviors (Fig. 5D, F and G). *tim-GAL4* driven *UAS-Kir2.1* or *UAS-TrpA1*
expression disrupted viability (Fig. 5D). While *cry-GAL4* driven *UAS-Kir2.1* did not yield viable flies,
interestingly, activating neuronal function using *cry-GAL4* driven *UAS-dTrpA1* mimicked *cry* and *Pdf*
mutants in courtship responses to grouping (Fig. 5C-E; Supplementary Fig. 11C).

As *cry*, *tim*, and *Clk* are partially coexpressed with fru^M in subsets of neurons in the brain that label the
clock neural network, we further explored how gene expression is altered in these overlapping
populations. Given the strong behavioral phenotype of *Clk* mutants, we focused on the *Clk+* cell
population and analyzed DEGs within this subset using single-cell RNA-seq data (Fig. 5H,
Supplementary Fig. 11A). We observed that in group-housed males, *Hr38* and *sr* were upregulated in
Clk+ cells, while *fru* was significantly downregulated in this population (Fig. 5H). We validated this
expression level by quantifying Fru^M (FruV5) in *Clk*-expressing cells *in vivo* using imaging (Fig. 5I). To
test whether these changes contribute to courtship modulation, we manipulated the expression of *Hr38*,
sr, and fru^M in *Clk+* cells, followed by courtship assays. Specifically, we knocked down *Hr38*, *sr*
(Supplementary Fig. 11D), and fru^M, and overexpressed fru^M in *Clk+* neurons (Fig. 5J-K). However, none
of these manipulations affected the courtship difference between GH and SH males compared to controls.
Together, these results suggest that while circadian genes like *Clk*, *tim, pdf, cry Sik3*, are crucial for
mediating the effect of group housing on courtship suppression, fru^M and IEG function in *Clk*-positive
neurons and their neuronal activity are not implicated. Rather, Fru^M function in other clock neurons (i.e,
Pdf or *cry*-expressing neurons included in broad P1 drivers in Fig.1) might modulate courtship
suppression in group-housed males. Yet, given the *Clk* mutant phenotype, these results raise the
possibility that non-neuronal *Clk*-positive cells around the brain, like fat body cells or glia cells (Patop et
al., 2023), might still have an influence on social modulation of behaviors. Overall, our results suggest
that group housing might impinge upon the circadian networks serving as a general hub for
reprogramming the arousal state or responsiveness of the core courtship circuits in the brain.

Discussion

Social experiences, from isolation to social enrichment, exert profound and enduring influences on
cognition, physiology, and behavior (Cacioppo and Hawkley, 2009; Dow-Edwards et al., 2019; Jeste et
al., 2020). Loneliness is associated with an increased incidence and severity of numerous neuropsychiatric
and neurodegenerative disorders, highlighting its emergence as a contemporary epidemic, particularly
detrimental to elderly populations who are more susceptible to social isolation epidemic (Cacioppo and
Hawkley, 2009; Dow-Edwards et al., 2019; Jeste et al., 2020). Despite these widely known effects of
social isolation on brains and behaviors, the molecular and circuit-based mechanisms through which
social experiences shape health and behavior remain largely unexplored. In this study, we
used *Drosophila melanogaster* to investigate the neural and molecular pathways by which social
experiences influence male courtship behavior.

Group housing suppresses male courtship activity and reduces the evoked responses in P1 courtship
command neurons. The transcription factor Fru^M plays an essential role in this modulation, acting within
distinct neuronal subtypes to integrate social signals and modulate behavioral output. Furthermore, the
immediate early gene *sr*, functioning in *fru*-expressing neural circuits, are critical for maintaining the
behavioral distinctions between grouped and isolated males. Beyond fru^M, circadian genes (*Sik3*, *tim, cry,*
Clk and *Pdf*) - some are putative Fru^M targets (Neville et al., 2014) - also contribute significantly to
sustaining the social context-dependent variation in courtship behavior. Given that social experience
resets the circadian rhythm and alters the activity within central clock neurons (Kim et al., 2013; Levine
et al., 2002), our results highlights modulation of circadian state as a causal mechanism through which
social experience influences courtship behaviors.

Previous studies have shown that monosexual group housing suppresses male courtship behaviors, such
as wing extension and courtship song (Dankert et al., 2009; Goncharova et al., 2016). Goncharova et al.

further proposed that pre-grouped males avoid contact during courtship tests, possibly due to a
conditioned fear response arising from male-male aggression during group housing. In the present study,
we confirmed that group housing suppresses courtship vigor; however, this suppression is reversible—
isolating previously grouped males restores their courtship activity. Notably, the degree of suppression
increases with prolonged group housing, suggesting a time- and exposure-dependent mechanism
underlying this behavioral plasticity.

At the neural circuit level, P1 neurons are key command neurons for male courtship behavior, capable of
sustaining courtship-associated states for minutes (Inagaki et al., 2014; Jung et al., 2020; Kimura et al.,
2008; Pan et al., 2012; Von Philipsborn et al., 2011). Previous studies found that isolated males were
more sensitive to the same level of optogenetic stimulation of P1 neurons and showed an increase in wing
extension compared to grouped males in response to the stimulation (Inagaki et al., 2014). *Ex vivo*
imaging of P1 neurons in grouped male brains showed lower evoked responses than isolated male brains
(Inagaki et al., 2014; Zhao et al., 2024). These results suggest that P1 neuronal activity less sensitized in
GH male brains. Consistent with these findings, in this study, we show that P1 neurons in live-behaving
grouped males exhibited reduced evoked responses to female cues. Together, these data support the idea
that group housing diminishes P1 neuron sensitivity, thereby increasing courtship latency and reducing
overall courtship vigor.

The question then arises: how does social experience regulate the activity of P1 neurons? As an
integrative hub for multiple sensory inputs, P1 neurons are modulated by upstream circuit components
(Clowney et al., 2015; Kohatsu et al., 2011; Nässel and Wu, 2022). Social conditions likely alter the
activity or connectivity of these upstream inputs. For instance, Drosulfakinin (DSK) has been shown to
inhibit male courtship through its receptor CCKLR-17D3 on P1 neurons, and group housing increases
synaptic connections between DSK-expressing neurons and P1 neurons in male brains (Wu et al., 2019).
This enhanced inhibition may underlie the reduced courtship vigor observed in grouped males. In the
future, monitoring responses of individual neurons upstream of P1 in the courtship circuit in grouped or
isolated male brains will reveal neuronal paths implicated in the modulation of behaviors with social
experience.

Given that P1 neurons express both *fruitless^M* and *doublesex^M*, it is important to determine whether these
courtship regulators contribute to the social experience-dependent modulation of courtship. Single-cell
and bulk RNAseq experiments revealed that social isolation increases *fru^M* and *dsx^M* reporter expression,
the levels of *fru^M* transcripts in *Clk*+ cells, and levels of *dsx^M* transcripts throughout the brain. Even
though bulk RNAseq experiments showed that *dsx^M* expression increases in isolated male brains, it is not
required in neurons for driving the courtship difference between grouped and isolated males. This
suggests that *dsx^M* may influence other behaviors or structural changes in the brain rather than courtship
per se, or its function in this process is located in non-neuronal cells around the brain. In contrast,
knocking down *fru^M* in *fru^M*+ neurons in the brain abolished the behavioral differences between grouped
and isolated males, by decreasing courtship in isolated males. This result is in agreement with a role for
Fru^M in positively regulating courtship. Interestingly, knocking down *fru^M* in certain neurons increases
courtship in grouped males (Fig. 1F), suggesting Fru^M might play different roles in various neuronal
populations with opposing influences on courtship circuits. Regardless, these results demonstrate that
Fru^M function is essential for encoding the behavioral impact of social experience.

Beyond *fru^M*, this study also identified circadian-related genes, such as *Clk*, *tim*, *Pdf, cry,* and *Sik3*, as key
modulators of courtship behavior under different social contexts. *Sik3* expression, which we found to be
upregulated in group-housed brains, has been shown to influence male sexual circadian rhythms (Fujii et
al., 2017). Mutants for *tim* and *Clk* exhibited disrupted or eliminated differences in courtship between

grouped and isolated males, indicating that circadian reprogramming likely contributes to social
modulation of courtship. Compared to *Clk* and *tim* mutants, *cry* mutants further decrease courtship
responses in grouped males, suggesting the effects of circadian genes on reprogramming behaviors are
restricted to group-housed conditions. It is interesting that some of these circadian genes have binding
sites for Fru^M or Dsx^M in their upstream regulatory elements (Clough et al., 2014; Neville et al., 2014),
suggesting potential transcriptional cascades in response to social experience. Future experiments will
reveal the cells where these regulatory interactions occur and how activity/inactivity within courtship
circuits induced through social experience affects their expression.

Within the *Clk+* cells, *sr* and *Hr38* expression levels were upregulated in grouped male brains compared
to isolated brains, whereas *fru* expression was decreased. Yet, manipulating *fru^M*, *sr*, or *Hr38* expression
in *Clk+* neurons did not impact courtship behavior, suggesting that they do not function in *Clk+* cells for
grouping-induced suppression of courtship. A recent study showed that neurons where *Clk* and *fru^M*
overlap influence courtship, and that *fru^M+* neurons receive input from *Clk+* neurons (Deluca et al., 2025).
These findings suggest that while *Clk+* neurons may not be part of the core courtship circuit, they can
indirectly influence the function of neurons within courtship circuits. *Clk* mutants exhibited elevated
courtship behavior in grouped males compared to wild-type, yet manipulating *Clk*-positive neural activity
does not have an effect. One possible explanation is that in group-housed male brains, Clk protein
functions in non-neuronal cells (i.e. fat body or glia) to suppress the function of the courtship circuits. In
contrast, both manipulating the function of *cry* and *Pdf,* normally co-expressed in a subset of clock
neurons, or the function of *cry*-positive neurons, resulted in further suppression of courtship in group
housed males. These results suggest opposing function of different circadian genes and the activity of a
subset of clock neurons mediate the effects of social experience on courtship responses to group housing.

Even though grouping-induced courtship suppression is disrupted in *Clk* and *tim* mutants, their mode of
transcriptional modulation by social experience and their impact on courtship are not straightforward.
Among the three core circadian genes, *Clk* expression consistently shows a downregulation in response to
group housing using multiple approaches. In contrast, the expression of *tim* and *cry* shows some variation
based on the methodology used. Given the complex transcriptional and translational regulation of the core
circadian molecular oscillator, a deeper investigation into the social modulation of the circadian clock will
help resolve these discrepancies.

One surprising finding we observed is that for both *Clk* and *cry* genes, the phenotypic trends tend to
manifest in opposition to predicted outcomes based on transcriptional responses to social experience. For
example, the *Clk* gene is upregulated in isolated male brains, while *Clk* mutants increase courtship in
group-housed males. Similarly, Cry is upregulated in group-housed male brains, while *cry* mutants further
decrease courtship in response to group housing. These are possibly due to the existence of double-
negative regulatory loops regulating the function of clock neural networks by the circadian protein. Given
that the phenotypic effect of circadian mutants is restricted to GH males, it is also possible that they
contribute much less to the modulation of behavior in response to socially isolated conditions. These
results support the hypothesis that circadian genes play essential roles in mediating the effects of social
experience on male courtship behavior, agreeing with previous reports on reprogrammed circadian state
with social experience (Levine et al., 2002). They also point to opposing effects of Cry and Pdf versus
Tim and Clk in mediating the effects of group housing on courtship behaviors, and that there might be
more dominant programs mediating behavioral response to social isolation that rely on Fru^M function.

Within both *Clk+*, *fru^M+*, and *dsx^M+* neuronal subsets, the immediate early genes *sr* and *Hr38* showed
elevated expression in grouped male courtship circuits. Functional experiments revealed that upregulation
of *sr* within *fru^M+* neurons in response to group housing is required to suppress courtship. While it does

not appear to act within P1 or *Clk+* neurons directly, it may influence specific, yet unidentified, subsets of
fru+ neurons or other clock neuron subsets. Future studies are needed to map the precise cellular
substrates through which *sr* operate to mediate socially driven behavioral changes.

Beyond courtship, social experience modulates a wide array of behaviors in *Drosophila*, including sleep,
feeding, locomotion, and aggression (Agrawal et al., 2020; Dankert et al., 2009; Ganguly-Fitzgerald et al.,
2006; Goncharova et al., 2016; Kacsoh et al., 2018; Li et al., 2021; Ueda and Kidokoro, 2002; Wang et
al., 2008; Zhao et al., 2024). The social isolation-induced rise in aggression is suggested to be regulated
by genes involved in reactive oxygen species (ROS) metabolism, *Hyperkinetic* (*Hk*) and *glutathione S-*
transferase-S1 (*gsts1*) (Ueda and Wu, 2009), as well as the neuropeptide Drosulfakinin (Dsk) in the
central brain (Agrawal et al., 2020), and cytochrome gene *Cyp6a20* in the peripheral olfactory systems
(Wang et al., 2008). Furthermore, Or65a olfactory receptor neurons (ORNs), which respond to the male
pheromone 11-*cis*-vaccenyl acetate (cis-VA), mediate the suppression of aggression in group housed
male (Liu et al., 2011). Group-housed *Drosophila* males also decrease their locomotion compared to
single-housed males, and this is mediated by P1 neurons and other dsx^M-positive neurons, where
neuropeptides for rhythmic locomotion and aggression, DH44 and TK, oppositely regulate locomotion
through feedback loops (Zhao et al., 2024).

Whether a shared molecular or circuit-level mechanism underlies the effects of social experience across
these diverse behaviors remains an open and compelling question. Given the pleiotropic suppressive
effects of social experience on a variety of behaviors, there is potential for central arousal-related
networks in mediating the effects. The finding that group housing induced courtship suppression requires
circadian reprogramming is significant, as it might also underlie modulation of many of the behaviors in
response to social experience, particularly in light of previous reports suggesting social experience resets
circadian state (Levine et al., 2002). For example, many behaviors are suppressed during nighttime due to
the modulation of arousal through molecular and circuit-level oscillations of the circadian clock (Duhart
et al., 2023). Changes happening in the circadian system in response to social experience can provide a
central modulatory hub that mediates many of the effects of social experience on organismal behaviors
and physiology. It is likely that any change in circadian state would result in state changes in downstream
neurons driving diverse behaviors. These can include P1 command neurons and dopaminergic neurons, as
activity in both is affected by social experience and was shown to modulate courtship, aggression,
feeding, and sleep (Agrawal et al., 2019; Alekseyenko et al., 2013; Ganguly-Fitzgerald et al., 2006;
Karam et al., 2020; Zhang et al., 2018). A systematic examination of transcriptional and circuit
mechanisms by which social signals engage arousal systems, like the circadian system, will be crucial for
understanding the central strategies mediating the brain's adaptive responses to social environments.
Social isolation has been increasingly recognized as a risk factor not only for mental health conditions
such as depression and anxiety, but also for neurodegenerative diseases like Alzheimer's (Cacioppo et al.,
2015; Cacioppo et al., 2006; Cohen, 2004). By uncovering the mechanisms through which the social
environment influences neural circuits, this study may help inform strategies to mitigate the negative
consequences of social isolation on brain health across species, including humans. Thus, our findings
contribute to a deeper understanding of how social experience shapes brain function and behavior.

Methods

Drosophila strains and husbandry

Flies were raised on standard fly food from Archon Scientific company at 25 °C (except the courtship assays with dTrpA1, labeled in figures) under a 12-h light: 12-h dark (LD) cycle. All experiments used male flies except targets of courtship behavior assays. Male flies used in courtship behavior assays, bulk tissue RNA sequencing, single-cell RNA sequencing, two-photon calcium live imaging and quantification of immunofluorescence and confocal microscope imaging were unmated and 7 days old unless mentioned otherwise. For male-female single-pair courtship behavior assays, unmated males were used as the test fly, and 4 to 6-day-old w^{1118} virgin females were used as the target fly. Male flies used for courtship behavior assays were w^+ background (except *dsx* mutant) and did not have balancers. Fly stocks were obtained from the Bloomington Drosophila Stock Center (BDSC), the Vienna Drosophila Resource Center (VDRC), or kindly provided by other labs. Specific genotypes of males used in each figure are listed together with the sources for each fly stock in Table 1 (same sources are listed only once, at their first appearance). Split GAL4 lines *R15A01-AD* and *R71G01-DBD* were crossed together for intersectional labeling of P1 neurons (*P1-spGAL4*). *Otd-FLP/tub>GAL80>; fru^{GAL4}* or *Otd-FLP; fru^{GAL4}/tub>GAL80>* restricts expression from *fru^{GAL4}* to neurons in the central brain (*fru brain-GAL4*) while *Otd-FLP/tub>GAL80>; dsx^{GAL4}* restricts expression from *dsx^{GAL4}* to neurons in the central brain (*dsx brain-GAL4*).

Table 1. Genotypes and sources

Figure number	Genotypes	Sources
Figure 1A-D, 2, 5B, Supplementary Figure 1A and E, 3, 6C, 11A	• *Canton-S (CS)*	• *Canton-S (CS):* Gift of the Yapici lab from Cornell University
Supplementary Figure 1B - D	• *R15A01-p65.AD; R71G01-GAL4.DBD/20XUAS-IVS-jGCaMP7s*	• *R15A01-p65.AD*: BDSC #68837 • *R71G01-GAL4.DBD*: BDSC #69507 • *20XUAS-IVS-jGCaMP7s*: BDSC #79032
Figure 1E1	• *R15A01-p65.AD/10XUAS-IVS-mCD8GFP; R71G01-GAL4.DBD/+*	• *10XUAS-IVS-mCD8GFP*: BDSC #32186
Figure 1E2	• *w+; R15A01-p65.AD/+; R71G01-GAL4.DBD/+* • *w+; +/+; UAS-fru^M-miRNA/+* • *w+; R15A01-p65.AD/+; R71G01-GAL4.DBD/UAS-fru^M-miRNA*	• *UAS-fru^M-miRNA* (Meissner et al., 2016): Gift of the Ding lab from University of Pennsylvania.
Figure 1E3	• *w+; R15A01-p65.AD/+; R71G01-GAL4.DBD/+* • *w+; +/+; UAS-dsx^M-RNAi/+* • *w+; R15A01-p65.AD/+; R71G01-GAL4.DBD/UAS-dsx^M-RNAi*	• *UAS-dsx^M-RNAi* (Peng et al., 2022): Gift of the Pan lab from Southeast University.
Figure 1F1	• *10XUAS-IVS-mCD8GFP/+; R71G01-GAL4/+*	• *R71G01-GAL4*: BDSC #39599

Figure 1F2	• $w+; +/+; R71G01\text{-}GAL4/+$ • $w+; +/+; UAS\text{-}fru^M\text{-}miRNA/+$ • $w+; +/+; R71G01\text{-}GAL4/UAS\text{-}fru^M\text{-}miRNA$	
Figure 1F3	• $w+; +/+; R71G01\text{-}GAL4/+$ • $w+; +/+; UAS\text{-}dsx^M\text{-}RNAi/+$ • $w+; +/+; R71G01\text{-}GAL4/UAS\text{-}dsx^M\text{-}RNAi$	
Figure 1G1, Supplementary Figure 2A	• $w+; Otd\text{-}FLP/tub>GAL80>; fru^{GAL4}/40XUAS\text{-}IVS\text{-}mCD8GFP$	• $Otd\text{-}FLP$ (Asahina et al., 2014): Gift of the Asahina lab from the Salk Institute for Biological Studies • $tub>GAL80>$ (Chr2): Gift of the Asahina lab from the Salk Institute for Biological Studies • fru^{GAL4} (Stockinger et al., 2005): Lab stock • $40XUAS\text{-}IVS\text{-}mCD8GFP$: BDSC #32195
Figure 1G2	• $w+; Otd\text{-}FLP/tub>GAL80>; fru^{GAL4}/+$ • $w+; tub>GAL80>/+; UAS\text{-}fru^M\text{-}miRNA/+$ • $w+; Otd\text{-}FLP/tub>GAL80>; fru^{GAL4}/UAS\text{-}fru^M\text{-}miRNA$	
Figure 1H1, Supplementary Figure 2B	• $w+; Otd\text{-}FLP/tub>GAL80>; dsx^{GAL4}/40XUAS\text{-}IVS\text{-}mCD8GFP$	• dsx^{GAL4} (Rideout et al., 2010): Lab stock
Figure H2	• $w+; Otd\text{-}FLP/tub>GAL80>; dsx^{GAL4}/+$ • $w+; tub>GAL80>/+; UAS\text{-}dsx^M\text{-}RNAi/+$ • $w+; Otd\text{-}FLP/tub>GAL80>; dsx^{GAL4}/UAS\text{-}dsx^M\text{-}RNAi$	
Supplementary Figure 2C	• $Canton\text{-}S (CS)$ • $w+; +/+; fru^{LexA/4\text{–}40}$	• fru^{LexA}: Lab stock (Anand et al., 2001; Mellert et al., 2010). • $fru^{4\text{–}40}$: Lab stock (Anand et al., 2001; Mellert et al., 2010).
Supplementary Figure 2D	• $w+; +/+; fru^{GAL4}/+$ • $w+; +/+; UAS\text{-}fru^M\text{-}miRNA/+$ • $w+; +/+; fru^{GAL4}/UAS\text{-}fru^M\text{-}miRNA$	
Supplementary Figure 2E	• $Canton\text{-}S (CS)$ • $w\text{-}; +/+; dsx^{f00683\text{-}d07058/f01649\text{-}d09625}$	• $dsx^{f00683\text{-}d07058}$: Lab stock (Chatterjee et al., 2011). • $dsx^{f01649\text{-}d09625}$: Lab stock (Chatterjee et al., 2011).
Figure 3, 5H, Supplementary	• $w+; +/+; fru^{GAL4} 40XUAS\text{-}IVS\text{-}mCD8GFP/ dsx^{LexA} 26XLexAop2\text{-}mCD8GFP$	• dsx^{LexA} (Zhou et al., 2015): Gift of the Yang lab from Department of

Figure 4, 5, 6, 7, 8, 9C, 10K, 11A		Neurobiology at Duke University. • *26XLexAop2-mCD8GFP*: BDSC #32207
Figure 3B, Supplementary Figure 4A	• *w+; +/+; fru^{GAL4} 40XUAS-IVS-mCD8GFP* • *w+; +/+; dsx^{LexA} 26XLexAop2-mCD8GFP*	
Figure 4A	• *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/+* • *w+; tub>GAL80>/+; UAS-Hr38-RNAi JF02541/+* • *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-Hr38-RNAi JF02541* • *w+; tub>GAL80>/+; UAS-sr-RNAi HMC05017/+* • *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-sr-RNAi HMC05017*	• *UAS-Hr38-RNAi JF02541*: BDSC #29377 • *UAS-sr-RNAi HMC05017*: BDSC #60024
Figure 4B	• *w+; Otd-FLP/tub>GAL80>; dsx^{GAL4}/+* • *w+; tub>GAL80>/+; UAS-Hr38-RNAi JF02541/+* • *w+; Otd-FLP/tub>GAL80>; dsx^{GAL4}/UAS-Hr38-RNAi JF02541* • *w+; tub>GAL80>/+; UAS-sr-RNAi HMC05017/+* • *w+; Otd-FLP/tub>GAL80>; dsx^{GAL4}/UAS-sr-RNAi HMC05017*	
Figure 4C	• *w+; spab^{GAL4}/+; +/+* • *w+; +/+; UAS-Dh31-RNAi HMS02354/+* • *w+; spab^{GAL4}/+; UAS-Dh31-RNAi HMS02354/+* • *w+; +/+; UAS-Sik3-RNAi HMC04380/+* • *w+; spab^{GAL4}/+; UAS-Sik3-RNAi HMC04380/+*	• *spab^{GAL4}*: BDSC #84617 • *UAS-Dh31-RNAi HMS02354*: BDSC #41957 • *UAS-Sik3-RNAi HMC04380:* BDSC #56941
Figure 4D	• *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/+* • *w+; tub>GAL80>/+; UAS-Sik3-RNAi HMC04380/+* • *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-Sik3-RNAi HMC04380*	

Supplementary Figure 9A	• *w+; Otd-FLP/+; fru^{GAL4}/+*
	• *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-cbt-RNAi 43136*
	• *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-Gk2-RNAi 35164*
	• *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-Hr38-RNAi 29376*
	• *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-Hr38-RNAi 29377*
		• *tub>GAL80>* (Chr3): BDSC #38881
		• BDSC stock numbers are included in line names:
		  ○ *UAS-cbt-RNAi 43136*
		  ○ *UAS-Gk2-RNAi 35164*
		  ○ *UAS-Hr38-RNAi 29376*

	• *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-Sik3-RNAi 28366* • *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-Sik3-RNAi 35378* • *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-Sik3-RNAi 56941* • *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-Sik3-RNAi 39005* • *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-sr-RNAi 27701* • *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-sr-RNAi 60024* • *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-Pdf-RNAi 25802* • *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-Pdf-RNAi v4380* • *w+; Otd-FLP/UAS-Pdf-RNAi v50749; fru^{GAL4}/tub>GAL80>* • *w+; Otd-FLP/tub>GAL80>; fru^{GAL4}/UAS-Pdf-RNAi v50750*	○ *UAS-Hr38-RNAi 29377* ○ *UAS-Sik3-RNAi 28366* ○ *UAS-Sik3-RNAi 35378* ○ *UAS-Sik3-RNAi 56941* ○ *UAS-Sik3-RNAi 39005* ○ *UAS-sr-RNAi 27701* ○ *UAS-sr-RNAi 60024* ○ *UAS-Pdf-RNAi 25802* • VDRC stock numbers are included in line names: ○ *UAS-Pdf-RNAi v4380* ○ *UAS-Pdf-RNAi v50749* ○ *UAS-Pdf-RNAi v50750*
Supplementary Figure 9B	• *w+; Otd-FLP/+; dsx^{GAL4}/+* • *w+; Otd-FLP/tub>GAL80>; dsx^{GAL4}/UAS-cbt-RNAi 43136* • *w+; Otd-FLP/tub>GAL80>; dsx^{GAL4}/UAS-Hr38-RNAi 29376* • *w+; Otd-FLP/tub>GAL80>; dsx^{GAL4}/UAS-Hr38-RNAi 29377* • *w+; Otd-FLP/tub>GAL80>; dsx^{GAL4}/UAS-sr-RNAi 27701* • *w+; Otd-FLP/tub>GAL80>; dsx^{GAL4}/UAS-sr-RNAi 60024* • *w+; Otd-FLP/tub>GAL80>; dsx^{GAL4}/UAS-ct-RNAi 29625* • *w+; Otd-FLP/tub>GAL80>; dsx^{GAL4}/UAS-ct-RNAi 33967* • *w+; Otd-FLP/tub>GAL80>; dsx^{GAL4}/UAS-LpR2-RNAi 31150* • *w+; Otd-FLP/tub>GAL80>; dsx^{GAL4}/UAS-LpR2-RNAi 54461*	• BDSC stock numbers are included in line names: ○ *UAS-ct-RNAi 29625* ○ *UAS-ct-RNAi 33967* ○ *UAS-LpR2-RNAi 31150* ○ *UAS-LpR2-RNAi 54461*
Supplementary Figure 10A	• *w+; 10XUAS-IVS-mCD8GFP/+; repo-GAL4/+* • *w+; +/+; repo-GAL4/+* • *w+; UAS-Ae2-RNAi 51827/+; repoGAL4/+* • *w+; UAS-Ae2-RNAi 63577/+; repoGAL4/+* • *w+; +/+; repoGAL4/UAS-Arc1-RNAi 25954* • *w+; +/+; repoGAL4/UAS-Hsp68-RNAi 50637* • *w+; UAS-Jhbp14-RNAi 60061/+; repoGAL4/+* • *w+; +/+; repoGAL4/UAS-Obp19d-RNAi 29432* • *w+; +/+; repoGAL4/UAS-aay-RNAi 28577* • *w+; +/+; repoGAL4/UAS-apolpp-RNAi 28946*	• *repo-GAL4:* BDSC #7415 • BDSC stock numbers are included in line names: ○ *UAS-Ae2-RNAi 51827* ○ *UAS-Ae2-RNAi 63577* ○ *UAS-Arc1-RNAi 25954*

	• *w+; +/+; repoGAL4/UAS-apolpp-RNAi 33388* • *w+; UAS-axo-RNAi 65911/+; repoGAL4/+* • *w+; UAS-Gs2-RNAi 40949/+; repoGAL4/+* • *w+; UAS-mfas-RNAi 52905/+; repoGAL4/+* • *w+; UAS-mfas-RNAi 58256/+; repoGAL4/+* • *w+; UAS-Msr-110-RNAi 51034/ +; repoGAL4/+* • *w+; +/+; repoGAL4/UAS-Rh50-RNAi 50666* • *w+; UAS-Tsf1-RNAi 55936/+; repoGAL4/+* *w+; UAS-Tsf1-RNAi 62968/+; repoGAL4/+* • *w+; +/+; repoGAL4/UAS-wdp-RNAi 28907* • *w+; UAS-wdp-RNAi 66004/+; repoGAL4/+*	○ *UAS-Hsp68-RNAi 50637* ○ *UAS-Jhbp14-RNAi 60061* ○ *UAS-Obp19d-RNAi 29432* ○ *UAS-aay-RNAi 28577* ○ *UAS-apolpp-RNAi 28946* ○ *UAS-apolpp-RNAi 33388* ○ *UAS-axo-RNAi 65911* ○ *UAS-Gs2-RNAi 40949* ○ *UAS-mfas-RNAi 52905* ○ *UAS-mfas-RNAi 58256* ○ *UAS-Msr-110-RNAi 51034* ○ *UAS-Rh50-RNAi 50666* ○ *UAS-Tsf1-RNAi 55936* ○ *UAS-Tsf1-RNAi 62968* ○ *UAS-wdp-RNAi 28907* ○ *UAS-wdp-RNAi 66004*
Supplementary Figure 10B	• *TH-GAL4* refers to a GAL4 insertion in the *ple* locus (*ple^{GAL4}*). • *w+; 10XUAS-IVS-mCD8GFP/+; TH-GAL4/+* • *w+; +/+; TH-GAL4/+* • *w+; UAS-cbt-RNAi 38276/+; TH-GAL4/+* • *w+; +/+; TH-GAL4/UAS-cbt-RNAi 43136* • *w+; +/+; TH-GAL4/UAS-CCAP-RNAi 29569* • *w+; UAS-CCAP-RNAi 62470/+; TH-GAL4/+* • *w+; +/+; TH-GAL4/UAS-Ms-RNAi 26245* • *w+; +/+; TH-GAL4/UAS-sNPF-RNAi 25867*	• *ple^{GAL4}*: BDSC #86289 • BDSC stock numbers are included in line names: ○ *UAS-cbt-RNAi 38276* ○ *UAS-CCAP-RNAi 29569* ○ *UAS-CCAP-RNAi 62470* ○ *UAS-Ms-RNAi 26245* ○ *UAS-sNPF-RNAi 25867*
Supplementary Figure 10C	• *w+; 10XUAS-IVS-mCD8GFP/+; Kr-GAL4/+* • *w+; +/+; Kr-GAL4/+* • *w+; +/+; Kr-GAL4/UAS-CCAP-RNAi 29569* • *w+; UAS-CCAP-RNAi 62470/+; Kr-GAL4/+* • *w+; +/+; Kr-GAL4/UAS-Gk2-RNAi 35164* • *w+; UAS-Gk2-RNAi 57522/+; Kr-GAL4/+* • *w+; +/+; Kr-GAL4/UAS-Capa-RNAi 28345* • *w+; +/+; Kr-GAL4/UAS-Ms-RNAi 26245* • *w+; +/+; Kr-GAL4/UAS-sNPF-RNAi 25867* • *w+; +/+; Kr-GAL4/UAS-spab-RNAi 27069*	• *Kr-GAL4:* BDCS #58800 • BDSC stock numbers are included in line names: ○ *UAS-Gk2-RNAi 57522* ○ *UAS-Capa-RNAi 28345* ○ *UAS-spab-RNAi 27069*

Supplementary Figure 10D	• $w+; spab^{GAL4}/10XUAS\text{-}IVS\text{-}mCD8GFP; +/+$ • $w+; spab^{GAL4}/+; +/+$ • $w+; spab^{GAL4}/UAS\text{-}CCHa1\text{-}RNAi\ 57562; +/+$ • $w+; spab^{GAL4}/+; UAS\text{-}Dh31\text{-}RNAi\ 41957/+$ • $w+; spab^{GAL4}/+; UAS\text{-}Gk2\text{-}RNAi\ 35164/+$ • $w+; spab^{GAL4}/UAS\text{-}Gk2\text{-}RNAi\ 57522; +/+$ • $w+; spab^{GAL4}/+; UAS\text{-}Sik3\text{-}RNAi\ 28366/+$ • $w+; spab^{GAL4}/+; UAS\text{-}Sik3\text{-}RNAi\ 35378/+$ • $w+; spab^{GAL4}/+; UAS\text{-}Sik3\text{-}RNAi\ 39005/+$ • $w+; spab^{GAL4}/+; UAS\text{-}Sik3\text{-}RNAi\ 56941/+$ • $w+; spab^{GAL4}/UAS\text{-}TwdlW\text{-}RNAi\ 57387; +/+$ • $w+; spab^{GAL4}/+; UAS\text{-}dve\text{-}RNAi\ 26225/+$ • $w+; spab^{GAL4}/+; UAS\text{-}FMRFa\text{-}RNAi\ 25870/+$ • $w+; spab^{GAL4}/+; UAS\text{-}FMRFa\text{-}RNAi\ 58197/+$ • $w+; spab^{GAL4}/+; UAS\text{-}MFS9\text{-}RNAi\ 29305/+$ • $w+; spab^{GAL4}/+; UAS\text{-}Ms\text{-}RNAi\ 26245/+$	• BDSC stock numbers are included in line names: ○ *UAS-CCHa1-RNAi 57562:* ○ *UAS-TwdlW-RNAi 57387* ○ *UAS-dve-RNAi 26225* ○ *UAS-FMRFa-RNAi 25870* ○ *UAS-FMRFa-RNAi 58197* ○ *UAS-MFS9-RNAi 29305*
Supplementary Figure 10E	• *w+; 10XUAS-IVS-mCD8GFP/+; dsf R29G10-GAL4/+* • *w+; +/+; dsf R29G10-GAL4/+* • *w+; +/+; dsf R29G10-GAL4/UAS-Sik3 RNAi 28366* • *w+; +/+; dsf R29G10-GAL4/UAS-Sik3 RNAi 35378* • *w+; +/+; dsf R29G10-GAL4/UAS-Sik3 RNAi 39005* • *w+; +/+; dsf R29G10-GAL4/UAS-Sik3 RNAi 56941*	• *dsf R29G10-GAL4*: BDSC #49501
Supplementary Figure 10F	• *w+; 10XUAS-IVS-mCD8GFP/+; Fer1 R57G11-GAL4/+* • *w+; +/+; Fer1 R57G11-GAL4/+* • *w+; +/+; Fer1 R57G11-GAL4/UAS-Dh31 RNAi 41957* • *w+; +/+; Fer1 R57G11-GAL4/UAS-Tk RNAi 25800* • *w+; +/+; Fer1 R57G11-GAL4/UAS-Sik3 RNAi 28366* • *w+; +/+; Fer1 R57G11-GAL4/UAS-Sik3 RNAi 35378* • *w+; +/+; Fer1 R57G11-GAL4/UAS-Sik3 RNAi 39005* • *w+; +/+; Fer1 R57G11-GAL4/UAS-Sik3 RNAi 56941*	• *Fer1 R57G11-GAL4*: BDSC #47674 • BDSC stock numbers are included in line names: ○ *UAS-Tk RNAi 25800*
Supplementary Figure 10G	• *w+; 10XUAS-IVS-mCD8GFP/+; Fer1 R11D04-GAL4/+* • *w+; +/+; Fer1 R11D04-GAL4/+* • *w+; +/+; Fer1 R11D04-GAL4/UAS-Dh31 RNAi 41957*	• *Fer1 R11D04-GAL4*: BDSC #48454

	• *w+; +/+; Fer1 R11D04-GAL4/UAS-Tk RNAi 25800* • *w+; +/+; Fer1 R11D04-GAL4/UAS-Sik3 RNAi 28366* • *w+; +/+; Fer1 R11D04-GAL4/UAS-Sik3 RNAi 35378* • *w+; +/+; Fer1 R11D04-GAL4/UAS-Sik3 RNAi 39005* • *w+; +/+; Fer1 R11D04-GAL4/UAS-Sik3 RNAi 56941*	
Supplementary Figure 10H	• *w+; 10XUAS-IVS-mCD8GFP/+; Sox102F R40A01-GAL4/+* • *w+; +/+; Sox102F R40A01-GAL4/+* • *w+; +/+; Sox102F R40A01-GAL4/UAS-ITP-RNAi 25799*	• *Sox102F R40A01-GAL4:* BDSC #50072 • BDSC stock numbers are included in line names: o *UAS-ITP-RNAi 25799*
Supplementary Figure 10I	• *w+; 10XUAS-IVS-mCD8GFP/+; Sox102F R42H07-GAL4/+* • *w+; +/+; Sox102F R42H07-GAL4/+* • *w+; +/+; Sox102F R42H07-GAL4/UAS-ITP-RNAi 25799*	• *Sox102F R42H07-GAL4:* BDSC #50172
Supplementary Figure 10J	• *w+; SoxN^{GAL4}/10XUAS-IVS-mCD8GFP; +/+* • *w+; SoxN^{GAL4}/+; +/+* • *w+; SoxN^{GAL4}/UAS-TwdlW-RNAi 57387; +/+* • *w+; SoxN^{GAL4}/UAS-Gs2-RNAi 40949; +/+*	• *SoxN^{GAL4}*: BDSC #95015 • BDSC stock numbers are included in line names: o *UAS-TwdlW-RNAi 57387*
Supplementary Figure 10L	• *w+; R15A01-p65.AD/+; R71G01-GAL4.DBD/+* • *w+; +/+; UAS-Hr38-RNAi JF02541/+* • *w+; R15A01-p65.AD/+; R71G01-GAL4.DBD/UAS-Hr38-RNAi JF02541*	
Figure 5A, B	• *w+; tim-GAL4/Cry-GFP; UAS-mCherry.NLS/+* • *w+; Clk856-GAL4/Cry-GFP; UAS-mCherry.NLS/+*	• *Cry-GFP*: BDSC #76317 • *tim-GAL4*: BDSC #7126 • *Clk856-GAL4*: BDSC #93198 • *UAS-mCherry.NLS*: BDSC #38424
Figure 5C, Supplementary Figure 11B	• *Canton-S (CS)* • *w+; +/+; cry$^{02/02}$* • *w+; tim$^{01/01}$; +/+* • *w+; +/+; Clk$^{out/out}$* • *w+; +/+; Pdf$^{01/01}$*	• *cry^{02}*: BDSC #86267 • *tim^{01}*: BDSC #80930 • *Clk^{out}*: BDSC #56754 • *Pdf$^{01/01}$* BDSC #26654
Figure 5D	• *10XUAS-IVS-mCD8GFP/+; cry-GAL4/+* • *tim-GAL4/10XUAS-IVS-mCD8GFP; +/+* • *Clk856-GAL4/10XUAS-IVS-mCD8GFP; +/+*	• *cry-GAL4:* BDSC #24514

Figure 5E, Supplementary Figure 11C	• *w+; +; cry-GAL4/+* • *w+; UAS-dTrpA1/+; +* • *w+; UAS-dTrpA1/+; cry-GAL4/+*	• *UAS-dTrpA1:* Gift of the Yang lab from Duke University.
Figure 5F	• *w+; Clk856-GAL4+; +* • *w+; UAS-dTrpA1/+; +* • *w+; Clk856-GAL4/UAS-dTrpA1; +*	
Figure 5G	• *w+; Clk856-GAL4+; +* • *w+; UAS-Kir2.1/+; +* • *w+; Clk856-GAL4/UAS-Kir2.1; +*	*UAS-Kir2.1:* Gift of the Yang lab from Duke University.
Figure 5I	• *w+; Clk856-GAL4/+; UAS-mCherry.NLS/FruV5*	• *FruV5* (Chen et al., 2021)*:* Gift of the Pan lab from Southeast University.
Figure 5J, K	• *w+; Clk856-GAL4/+; +/+* • *w+; +/+; UAS-fru^M-miRNA/+* • *w+; Clk856-GAL4/+; UAS-fru^M-miRNA/+* • *w+; U6-fru-sgRNA/+; UAS-3xFLAG.dCas9.VPR/+* • *w+; Clk856-GAL4/U6-fru-sgRNA; UAS-3xFLAG.dCas9.VPR/+*	• *U6-fru-sgRNA:* BDSC #80225 • *UAS-3xFLAG.dCas9.VPR*: BDSC #66562
Supplementary Figure 11D	• *w+; Clk856-GAL4/+; +/+* • *w+; +/+; UAS-Hr38-RNAi JF02541/+* • *w+; Clk856-GAL4/+; UAS-Hr38-RNAi JF02541/+* • *w+; +/+; UAS-sr-RNAi HMC05017/+* • *w+; Clk856-GAL4/+; UAS-sr-RNAi HMC05017/+*	

Social isolation and group housing setup

Dark pupae (80–100 hours old) were picked out and separated by sex. For social isolation (single housing, SH) condition, each pupa was placed into individual vials, allowed to eclose alone and aged to 7 days. For the group housing (GH) condition, more than 30 pupae with the same sex were collected and placed into food vials, and 30 newly eclosed and unmated flies with correct genotypes were collected and put in a fresh new vial the next day of the collection day (day 0). All *w^{1118}* virgin females used for courtship assay were raised under GH condition.

Male-female single-pair courtship behavior assay setup and analysis

Courtship behavior assays were performed during ZT 7 – ZT 10. Two types of chambers were used for courtship assays. The plastic chambers were used for manual scoring blindly. Agarose chambers were designed for automated scoring to improve the efficiency of scoring courtship behavior assays.

Plastic chamber and manual annotation: Each plastic chamber has two layers and 10 arenas, with each arena measuring 10 mm in diameter and 2.5 mm in height in each layer. To set up the experiment, one unmated male and one virgin *w^{1118}* female were gently aspirated to each arena in separate layers. When all the flies were loaded, the chamber was placed under a camera. Two layers were rotated to allow two flies to meet each other in one connected arena, and video recording was started immediately. After the recording, videos were processed by SLEAP (Pereira et al., 2022), and each arena was cropped out

separately. Copulation behavior and other courtship behaviors (including chasing, orientation, tapping,
single wing extension, licking, and attempted copulation) of flies in each arena were manually annotated
using the Boris software (Friard and Gamba, 2016).

Agarose chamber and automated annotation: Each agarose chamber has 12 round arenas, with each
arena measuring 25 mm in diameter and 3mm in height. To set up the experiment, one unmated male and
one virgin w^{1118} female were gently aspirated to each arena of an unused agarose chamber, separated by
metal barriers before the start of the recording. When all flies were loaded, the chamber was placed at a
fixed position under a camera with stable consistent light source. Barries were removed to allow two flies
to meet each other, and video recording was started immediately. The trajectory of each fly was tracked
by FlyTracker (Eyjolfsdottir et al., 2014) on MATLAB. Fly identity and the start time of copulation were
determined manually. JAABA was trained to automatically annotate other courtship behaviors (Kabra et
al., 2013).

Quantification: For each courtship behavior video, only the first 10 min were used for the scoring of the
courtship index (CI):

CI = (the time a male fly spends on courtship within 10 min)/(10 min)

The time spent on copulation was also included in the CI. The results from two kinds of chambers were
consistent by multiple different experiments. The housing switch (Fig. 1A-D; Supplementary Fig. 1A)
and *fru* mutant (Supplementary Fig. 2C) and *dsx* mutant (Supplementary Fig. 2E) experiments presented
above were performed on plastic chambers. All other courtship behavior experiments presented above
were performed on agarose chambers.

Bulk tissue RNA sequencing

Brain dissection and RNA extraction: Male brains were dissected in cold RNase-free 1X PBS and
transferred immediately in RNAlater (Sigma-Aldrich, R0901) on ice. Each biological replicate contained
30 male fly brains. After the dissection, dilute the RNAlater using 1X RNase-free PBS with 2 times the
volume of the RNAlater. Centrifuge for 1 min at max speed and remove the supernatant. Brain samples
were washed with 1 ml 1X RNase-free PBS and centrifuged for 1 min at max speed. The supernatant was
removed and replaced with 300 ul of TRIzol reagent (Invitrogen, 15596026). Briefly spin down to make
sure all tissues are in TRIzol. Brain samples were either processed immediately or stored at -80 ˚C for
long term. To extract RNA, samples were ground sufficiently and processed with QIAGEN Shredder kit
(QIAGEN, 79654) and then RNeasy Mini Kit (QIAGEN, 74104). DNase I (TURBO DNA-free Kit,
Invitrogen, Thermo Fisher Scientific AM1907) was applied to remove genome DNA after the RNA
extraction.

Library preparation and sequencing: Around 500 ng total RNA in each sample was prepared for
library preparation. KAPA Stranded mRNA-Seq Kit was used to generate mRNA libraries. Libraries were
sequenced on the NextSeq 2000 sequencer, generating 75-bp or 100-bp paired-end reads.

Bulk RNA sequencing data analysis: Raw fastq files were processed with the standard nf-core/rnaseq
pipeline (Ewels et al., 2020). Briefly, reads were mapped to the *Drosophila melanogaster* BDGP6.46
genome with STAR, and read counts were generated using the Salmon method. The following analysis
was performed in R (R Core Team, 2013; Wickham and Sievert, 2009). Differential expression analysis

was performed with DESeq2 (Love et al., 2014). Gene ontology (GO) analyses was performed using the clusterProfiler library (Yu et al., 2012).

Single-cell RNA sequencing

Single-cell suspension preparation and GFP positive cells sorting: Male fly brains were dissected in dissection buffer (Schneider′s Insect Medium, Sigma-Aldrich S0146 with 1% BSA) and transferred into dissection buffer in 1.5 ml tubes on ice. The old dissection buffer was replaced with the fresh dissection buffer. Collagenase Type I (Gibco, 17018029) stock solution was added directly into samples for a final 2mg/mL concentration. Samples were incubated at 37 °C for 20 minutes. During the incubation, samples were dissociated by gentle pipetting 20-30 times every 7 min with P200. The collagenase solution was removed after centrifugation at 300 xg for 5 min at 4°C. Cells were resuspended with 200 ul cold elution buffer (1X PBS, Gibco, 70011-044 with 0.1% BSA). Cell suspensions were passed through 30 um cell strainer caps into 5 ml collection tubes on ice. Cell strainer caps were rinsed with 800 ul cold elution buffer. Cells were processed by flow cytometry (Sony SH800 Cell Sorter) with wild-type *CS* brain cells as the negative control. GFP-positive single cells were sorted into 1.5 ml tubes and immediately transferred on ice to Duke Molecular Genomics Core (MGC) for 10X single-cell RNA seq library preparation and sequencing. Each biological replicate used thirty male fly brains, and each housing condition has two independent biological replicates. Each batch includes one GH and one SH sample processed simultaneously.

10X Genomics library preparation and sequence: Cell viability and concentration were assessed by Cellometer K2 (Revvity PN CMT-K2-MX-150). Single-cell libraries were generated using Chromium X (PN 1000326), Chromium Controller (PN 110203), and 10x Genomics v3 Next GEM 3' Single Cell chemistry (PN 1000269) following the manufacturer's protocol, with around 20K single cells in each sample. Libraries were sequenced by the NovaSeq sequencer, and 36K -50K reads/cell were sequenced. For quality control, cDNA QCs were performed using Agilent High Sensitivity D5000 ScreenTape (Cat #5067-5592) with High Sensitivity D5000 Reagents (Cat #5067-5593). Final library QCs were performed using the Agilent High Sensitivity D1000 ScreenTape (Cat #5067-5584) with High Sensitivity D1000 Reagents (Cat #5067-5585).

10X data processing: The basic analysis for the single-cell RNA sequencing followed the recommended protocols from 10X Genomics. Briefly, Cell Ranger v7.1.0 was used to demultiplex raw FASTQ files, align the reads to FlyBase release dmel-all-chromosome-r6.50 reference genome (Tweedie et al., 2009), and generate gene expression matrices for all single cells in each sample. Around 10K cells were recovered in each sample. Seurat objectives were generated for the following analysis (Satija et al., 2015). Doublets were removed by scDblFinder (Germain et al., 2022). Low-quality cells were removed (nUMI lower than 1000 or higher than 20000, gene numbers lower than 200 or higher than 4500, mitochondria genes higher than 15 percent). To do the clustering, two replicates of each housing condition were integrated first, and four datasets were merged. Clustering was performed using a standard Seurat pipeline with top 40 PCAs. Under resolution 1, the *fru+* or *dsx+* cells were sorted into 83 clusters in UMAPs (McInnes et al., 2018). Cell types were annotated manually based on existing cell markers (Davie et al., 2018; Park et al., 2022). Pseudo-bulk DEGs analysis were performed using DESeq2 (Love et al., 2014). Gene ontology (GO) analyses was performed using the clusterProfiler library (Yu et al., 2012).

Immunofluorescence and confocal microscope imaging

Fly brains were dissected in cold PBT buffer (1XPBS with 10% Triton X-100) and were fixed in 4% paraformaldehyde (PFA) on nutators at room temperature for 30 minutes. Fixed brains were washed three times with fresh PBT on nutators, and every wash was 10 min at room temperature. Brain samples were stained with primary antibodies on nutators at 4 °C overnight and washed three times with fresh PBT on

nutators, and every wash was 20 min at room temperature. Then, brains were stained with secondary
antibodies on nutators at 4 °C overnight and washed three times with fresh PBT on nutators, and every
wash was 20 min at room temperature. Antibodies used are: Rat-anti-NCad (monoclonal, DSHB-Ex#8-s,
1:20), Rabbit-anti-GFP (polyclonal, Invitrogen A11122, 1:1000), Mouse-anti-V5 (Bio-Rad,
MCA1360GA, 1:200), Goat-anti-rat 647 (polyclonal, Invitrogen A21247, 1:200), Goat-anti-rabbit 488
(polyclonal, Invitrogen A11008, 1:1000), Goat-anti-mouse 488 (polyclonal, Invitrogen A11029, 1:200).
Brains were mounted using Fluoromount-G Slide Mounting Medium (SouthernBiotech). Images were
taken by Olympus FluoView FV1000 confocal microscope.

Fluorescence intensity quantification was performed using Python. Background-subtracted fluorescence
signals were obtained by subtracting the background from the raw fluorescence intensity in each image
layer, and total intensity was calculated by summing across all layers. For quantification of FruV5 signal
in *Clk*-positive cells, the overlapping area between FruV5 and *Clk* signals was defined first, and FruV5
intensity within this region was calculated as described above. Statistical analyses were performed in R.

Two-photon calcium live imaging

The two-photon calcium live imaging was performed as described previously by the Yapici lab (Mabuchi
et al., 2023). Briefly, male flies were mounted on dissection holders and put in dark for one hour before
the experiment. Right before imagining, the cuticle on the posterior of the head between the eyes was
removed in saline. Flies were then mounted on the two-photon microscope under the objective. The P1
area was located under the scope. During the living imaging, a virgin female was delivered to a reachable
position by fly forelegs to stimulate the test male fly. Neuronal responses reflected by GCaMP7s signals
were recorded by the two-photon microscopy.

Statistics

682 Control and experimental flies were raised and tested in parallel under identical conditions. Statistical
683 analyses were performed using R or GraphPad. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$. All
684 experiments were repeated in different batches. Specific animal numbers were shown in each figure or in
685 each method part.

Acknowledgements

We would like to thank members of the Volkan lab for discussions and help with the manuscript. We
thank Dr. Yun Ding (University of Pennsylvania), Dr. Yufeng Pan (Southeast University), Dr. Kenta
Asahina (Salk Institute for Biological Studies) and Dr. Rebecca Chung-Hui Yang (Duke University) for
kindly sharing fly lines. We would like to acknowledge the Duke Molecular Physiology Institute
Molecular Genomics Core, especially Karen Abramson, Ellora Haukenfrers and Michael Aksu for their
help on the generation of single cell RNA sequencing data. We thank the Duke High Throughput
Sequencing Facility, Bloomington Stock Center, Developmental Studies Hybridoma Bank, Vienna
Drosophila Resource Center, and Drosophila Genomics Resource Center for their services.

Funding

698 This study was supported by the National Institute of Health grant R01 GM146010 and the National
699 Science Foundation award 2006471 to PCV.

Author contributions

701 Conceptualization: CD and PCV. Investigation: CD, SO, LJ, MRB and SS with help from YM, SA, LG,
702 SR, SK, EB, and JL. Analysis/interpretation of data: CD, JESF, CDJ, NY, and PCV. Manuscript—
703 original draft: CD and PCV.

Figures

Figure 1. Social experience induces plastic changes in male fruit fly courtship behaviors and P1 neuronal activity. (A) Experimental paradigm for the setup of four different housing conditions. **(B)** Pie-Donut charts showing the percentage of flies at each courtship stage. **(C-D)** Courtship Index **(C)** and Courtship Latency **(D)** in male-female single pair assays with male flies raised under four different housing conditions. The Kruskal–Wallis rank sum test, followed by Dunn's multiple comparisons test, was applied for the comparison among different housing conditions grouped by age. Significance of adjusted P-value (P_adj): *P_adj < 0.05, **P_adj < 0.01, ***P_adj < 0.001, ****P_adj < 0.0001. **(E1)** P1 neurons labelled by *P1-spGAL4> mCD8::GFP* in the brain, stained for GFP (green) and Ncad (magenta). See Methods for genotypes. **(E2-E3)** Courtship Index from male–female single-pair assays in control and experimental groups with *fru* **(E2)** or *dsx* **(E3)** knockdown in P1 neurons labeled by the stricter P1 driver. **(F1)** Neurons labelled by *R71G01GAL4> mCD8::GFP* in the brain, stained for GFP (green) and Ncad (magenta). **(F2-F3)** Courtship Index from male–female single-pair assays in control and experimental groups with *fru* **(F2)** or *dsx* **(F3)** knockdown in P1 neurons labeled by the broader P1 driver. **(G1)** Neurons labelled by *fru brain-GAL4> mCD8::GFP* in the brain, stained for GFP (green) and Ncad (magenta). **(G2)** Courtship Index from male–female single-pair assays in control and experimental groups with *fru* knockdown in *fru* positive neurons in the brain labeled by the intersection of *fru* and *Otd*. **(H1)**

Neurons labelled by *dsx brain-GAL4> mCD8::GFP* in the brain, stained for GFP (green) and Ncad
(magenta). **(H2)** Courtship Index from male–female single-pair assays in control and experimental groups
with *dsx* knockdown in *dsx* positive neurons in the brain labeled by the intersection of *dsx* and *Otd*. The
Wilcox test was applied to the comparison between GH and SH conditions of the same genotype.
Significance of adjusted P-value (P_adj): *P_adj < 0.05, **P_adj < 0.01, ***P_adj < 0.001, ****P_adj <
0.0001.

Figure 2. Bulk tissue RNA sequencing revealed diverse transcriptional changes in male brains to social experience. (A) PCA analysis of transcriptional profiles among biological replicates from brains of group-housed and single-housed wild-type *CS* male flies. **(B)** DEG numbers of GH/SH comparisons (adjusted P-value < 0.001, |log₂FC| > 0.1). Red indicates genes upregulated in the GH condition, and blue indicates genes downregulated in the GH condition. **(C)** Volcano plots displaying Log2FoldChange (GH/SH) and -Log10(P-adj) of genes. Red indicates genes upregulated in the GH condition, and blue indicates genes downregulated in the GH condition. **(D)** Transcript levels for *fru* and *dsx* among all male brain samples. **(E)** Summary of GO terms enriched in DEGs from the GH CS vs. SH CS comparison in male brains. Colors of branches indicate categories of GO terms. Colors of the pie indicate clusters of DEGs. The size of the pie indicates gene numbers. **(F)** Top 20 most significantly enriched GO terms for DEGs of GH/SH comparisons (q-value < 0.05). **(G)** Connection plots of the top 20 most significantly enriched GO terms for DEGs from GH/SH comparisons (q-value < 0.05). Big pies are GO terms, and small dots indicate different genes. Colors indicate clusters of DEGs. The size of the pie indicates gene numbers.

Supplementary Figure 3. Transcript levels for representative housekeeping genes among all male brain samples in the bulk tissue RNAseq.

776
777

Figure 3. Cell-type-specific single-cell RNA sequencing revealed diverse transcriptional changes in male brains to social experience. (A) Workflow of the cell type-specific single-cell RNA sequencing of combined *fru+* and *dsx+* cells *(fru^GAL4 40XUAS-mCD8GFP; dsx-LexA 26XLexAop-mCD8GFP)* from brains of GH and SH males with the 10X Genomics platform. FACS: fluorescence-activated cell sorting. Created with BioRender. **(B)** Proportion of GFP-labeled combined *fru^+* cells and *dsx^+* cells, *fru^+* only cells, and *dsx^+* only cells in FACS. **(C-D)** Cluster annotation of cell-type-specific single-cell RNA sequencing. **(C)** UMAP plot illustrating six major cell classes, distinguishing glial cells from neurons characterized by their neurotransmitter profiles. **(D)** UMAP plot illustrating 83 specific cell clusters integrated across all brain samples. The annotations (bottom) were determined using marker gene expression. **(E)** Visualization of expression pattern of *fru*, *dsx*, *Hr38,* and *sr* in GH and SH male brains. Each dot is one cell colored by the expression levels of genes. **(F)** Pseudobulk analysis results showing Log2 fold changes (GH/SH) and -Log10 adjusted P-values of DEGs across cell subsets defined by specific marker genes, as indicated on the x-axis.

Supplementary Figure 4. The proportions of different cell clusters in GH and SH male brains from
single-cell RNAseq. (A) Proportion of GFP-labeled combined *fru*+ cells and *dsx*+ cells, *fru*+ only cells,
and *dsx*+ only cells. The X-axes indicate GFP fluorescence intensity. Gray-shaded regions (GFP+) are
magnified in the right panels and quantified in Figure 3B. Red and blue represent GH and SH conditions,
respectively; gray lines indicate negative controls from *CS* brain cells. **(B)** Comparison of cluster
proportions in GH and SH conditions in two biological replicates. **(C)** UMAP plot illustrating the overall
level of transcript counts.

Supplementary Figure 5. The increase in the number of *fru-* and *dsx*-positive cells originate from many clusters in the SH condition. (A) Comparison of cluster proportions in GH and SH conditions in two biological replicates, excluding clusters 6 and 18. **(B)** Comparison of processed counts in GH and SH conditions, adjusted for the actual percentage of GFP+ cells measured by FACS, with clusters 6 and 18 excluded. **(C)** Differences in cluster proportions between GH and SH conditions across two biological replicates in (A). Red indicates clusters with a higher proportion in GH, while blue indicates clusters with a higher proportion in SH. **(D)** Adjusted differences in cluster counts between GH and SH conditions across two biological replicates in (B). Red indicates clusters with more cells in GH, while blue indicates clusters with more cells in SH.

Supplementary Figure 6. The expression pattern of *fru, dsx, Hr38,* and *sr* in each cell cluster in single-cell RNA sequencing. (A) Violin plots comparing the distribution of *fru* and *dsx* expression levels across different cell clusters in GH and SH conditions in single-cell RNA sequencing. **(B)** Violin plots comparing the distribution of overall *fru* and *dsx* expression levels in cells with detectable *fru* level or *dsx* level, respectively, in GH and SH conditions in single-cell RNA sequencing. **(C)** Transcript levels for *Hr38* and *sr* among all male brain samples in the bulk tissue RNAseq. **(D)** Violin plots comparing the

821 distribution of *Hr38* and *sr* expression levels across different cell clusters in GH and SH conditions in
822 single-cell RNA sequencing. **(E)** Results of pseudobulk analysis showing Log2FoldChange (GH/SH) and
823 -Log10(P-adj) of *sr, Hr38* and *cbt* in each cell cluster.

A

B

Gene Ontology Top 20

Supplementary Figure 7. DEGs of each cell cluster from the GH vs SH comparison in single-cell
RNA sequencing. (A) Results of pseudobulk analysis showing Log2FoldChange (GH/SH) and -
Log10(P-adj) of DEGs in each cell cluster. **(B)** Top 20 most significantly enriched GO terms for DEGs of
GH/SH comparisons in different cell clusters from (A) (q-value < 0.01).

Supplementary Figure 8. DEGs in different subsets defined by marker genes in single-cell RNA
sequencing. (A) Marker genes for cell cluster 0, cluster 2, and cluster 3 from the UMAP clusters in
Figure 3D. **(B)** Volcano plots displaying Log2FoldChange (GH/SH) and -Log10(P-value) of genes in
different subsets defined by marker genes. Red indicates DEGs from pseudobulk analysis. **(C)** Top 10
most significantly enriched GO terms for DEGs of GH/SH comparisons in different cell subsets from
Figure 3F (q-value < 0.05).

Figure 4. Knocking down *sr* or *Sik3* eliminated the courtship difference between GH and SH males.
(A-B) Courtship Index from male–female single-pair assays in control and experimental groups with
Hr38 or *sr* knockdown in *fru* brain neurons **(A)** or *dsx* brain neurons **(B)**. **(C)** Courtship Index from male–
female single-pair assays in control and experimental groups with *Dh31* or *Sik3* knockdown in *spab+*
cells. **(D)** Courtship Index from male–female single-pair assays in control and experimental groups with
Sik3 knockdown in *fru* brain neurons. The Wilcox test was applied to the comparison between GH and
SH conditions of the same genotype. Significance of adjusted P-value (P_adj): *P_adj < 0.05, **P_adj <
0.01, ***P_adj < 0.001, ****P_adj < 0.0001.

Supplementary Figure 9. RNAi screening of DEGs in *fru* or *dsx* brain neurons. (A-B) Courtship
index of male flies with RNAi-mediated knockdown of selected DEGs in *fru*- **(A)** or *dsx*- **(B)** expressing
brain neurons. In each plot, the first column (+) represents the control group, with red indicating GH
conditions and blue indicating SH conditions. The dashed lines indicate the mean courtship index for
control groups. Each dot represents an individual male fly, with light red dots indicating that the
knockdown group exhibited significantly higher courtship levels compared to the control, and light blue
dots indicating that the knockdown group exhibited significantly lower courtship levels compared to the
control. The Wilcox test was applied to the comparison between each RNAi group and the control group.
Significance of adjusted P-value (P_adj): *P_adj < 0.05, **P_adj < 0.01, ***P_adj < 0.001, ****P_adj <
0.0001. **(C)** Visualization of the expression pattern of selected genes. Each dot is one cell colored by the
expression levels of each gene.

Supplementary Figure 10. RNAi screening identifies candidate genes mediating social experience-dependent courtship plasticity. (A-J) RNAi-mediated knockdown screening of selected DEGs in *repo-* **(A)**, *ple-* (TH) **(B)**, *Kr-* **(C)**, *spab-* **(D)**, *dsf-* **(E)**, *Fer1-* **(F-G)**, *Sox102F-* **(H-I)**, and *SoxN-* **(J)** expressing cells. Left: Cells labelled by different drivers in the brain, stained for GFP (green) and Ncad (magenta). Right: Courtship index of male-female courtship assays in each screening group. In each plot, the first column (+) represents the control group, with red indicating GH conditions and blue indicating SH

conditions. The dashed lines indicate the mean courtship index for control groups. Each dot represents an
individual male fly, with light red dots indicating that the knockdown group exhibited significantly higher
courtship levels compared to the control, and light blue dots indicating that the knockdown group
exhibited significantly lower courtship levels compared to the control. The Wilcox test was applied to the
comparison between each RNAi group and the control group. Significance of adjusted P-value (P_adj):
*P_adj < 0.05, **P_adj < 0.01, ***P_adj < 0.001, ****P_adj < 0.0001. **(K)** Volcano plots displaying
Log2FoldChange (GH/SH) and -Log10(P-value) of genes in P1 neurons. Red indicates DEGs from
pseudobulk analysis. **(L)** Courtship Index from male–female single-pair assays in control and
experimental groups with *Hr38* knockdown in P1 neurons with the P1-spGAL4 driver. The Wilcox test
was applied to the comparison between GH and SH conditions of the same genotype. Significance of
adjusted P-value (P_adj): *P_adj < 0.05, **P_adj < 0.01, ***P_adj < 0.001, ****P_adj < 0.0001.

Figure 5. Circadian genes mediate the effects of group housing on courtship behaviors. (A) Cry+ cells labelled by GFP, with *tim+* or *Clk+* cells labelled by mCherryNLS in male brains, stained for GFP (green). **(B)** Quantification of fluorescence intensity of Cry labelled by GFP, and *tim+* or *Clk+* labelled by mCherryNLS in **(A)**. The sum of net fluorescence signals (fluorescence signals – background signals) was compared between GH and SH male brains. Unpaired t-test was applied. **(C)** Courtship Index from male–female single-pair assays in *CS*, *cry* mutants, *tim* mutants, *Clk* mutants and *Pdf* mutants. **(D)** The GFP expression pattern driven by *cry-GAL4*, *tim-GAL4* and *Clk856-GAL4* used for activation/silencing of neuronal activity, stained for GFP (green) and Ncad (magenta). **(E-F)** Courtship Index from male–female single-pair assays with activation of *cry+* **(E)** or *Clk+* **(F)** neurons using dTrpA1. **(G)** Courtship Index from male–female single-pair assays with silencing of *Clk+* neurons using Kir2.1. **(H)** Volcano plots displaying Log2FoldChange (GH/SH) and -Log10(P-value) of genes in *Clk+* cells from the single-cell RNAseq. Red indicates DEGs from pseudobulk analysis. **(I)** Left: Fru+ cells labelled by V5, with *Clk+* cells labelled by mCherryNLS in male brains, stained for V5 (green). Right: Quantification of fluorescence intensity of Fru labelled by V5 in *Clk+* cells. The sum of net fluorescence signals (fluorescence signals – background signals) was compared between GH and SH male brains. Unpaired t-test was applied. **(J-K)** Courtship Index from male–female single-pair assays comparing control and experimental groups of *fru* knockdown **(J)** or *fru* overexpression **(K)** in *Clk*-expressing cells. For all courtship behaviors, the Wilcox test was applied to the comparison between GH and SH conditions of the

899　same genotype. Significance of adjusted P-value (P_adj): *P_adj < 0.05, **P_adj < 0.01, ***P_adj <
900　0.001, ****P_adj < 0.0001.

Supplementary Figure 11. The effect of the clock system and immediate early genes in social experience-dependent modulation of courtship. (A) Visualization of the expression pattern of *Clk* in the single-cell RNAseq. **(B)** Courtship Index from male–female single-pair assays in *CS*, and mutants in *cry*, *tim*, *Clk* and *Pdf*. **(C)** Courtship Index from male–female single-pair courtship assays with *cry-GAL4* driving dTrpA1 under 22°C and 29°C. **(B-C)** The Kruskal–Wallis rank sum test, followed by Dunn's multiple comparisons test, was applied for the comparison among different genotypes grouped by housing. **(D)** Courtship Index from male–female single-pair assays comparing control and experimental groups of *Hr38* and *sr* knockdown in *Clk* expressing cells. The Wilcox test was applied to the comparison between GH and SH conditions of the same genotype. Significance of adjusted P-value (P_adj): *P_adj < 0.05, **P_adj < 0.01, ***P_adj < 0.001, ****P_adj < 0.0001.

Figure 6. A model for social experience-dependent modulation of courtship. Group housing
suppresses the courtship vigor of male flies. The transcription factor Fru^M plays an essential role in this
modulation, acting within distinct neuronal subtypes to integrate social signals and modulate behavioral
output. The increase in fru^M+ cells in the central circuits likely occurs as an increase in fru^M expression,
which drives an increase in courtship in response to social isolation. In group-housed conditions,
upregulation of the immediate early gene, *sr,* in *fru+* neural circuits mediates male courtship suppression.
Circadian genes (*Sik3*, *tim*, *Clk*, *cry, Pdf*), potentially regulated by Fru^M/Dsx^M transcription factors, also
contribute significantly to social experience-dependent modulation in courtship behaviors. *cry* and *Pdf*
function oppositely to *Sik3*, *tim* and *Clk* to modulate courtship behaviors in the GH condition. In contrast,
the effects of these circadian genes are limited in socially isolated males, which might dominate the
behavioral elevation by regulating additional factors. These changes all together lead to suppressed
evoked P1 courtship command neuron responses and male courtship behaviors under GH condition
relative to the SH condition.

References

Agrawal, P., Chung, P., Heberlein, U., and Kent, C. (2019). Enabling cell-type-specific behavioral epigenetics in Drosophila: a modified high-yield INTACT method reveals the impact of social environment on the epigenetic landscape in dopaminergic neurons. BMC biology *17*, 1-19.

Agrawal, P., Kao, D., Chung, P., and Looger, L.L. (2020). The neuropeptide Drosulfakinin regulates social isolation-induced aggression in Drosophila. Journal of Experimental Biology *223*, jeb207407.

Alekseyenko, O.V., Chan, Y.-B., Li, R., and Kravitz, E.A. (2013). Single dopaminergic neurons that modulate aggression in Drosophila. Proceedings of the National Academy of Sciences *110*, 6151-6156.

Anand, A., Villella, A., Ryner, L.C., Carlo, T., Goodwin, S.F., Song, H.-J., Gailey, D.A., Morales, A., Hall, J.C., and Baker, B.S. (2001). Molecular genetic dissection of the sex-specific and vital functions of the Drosophila melanogaster sex determination gene fruitless. Genetics *158*, 1569-1595.

Anders, F. (1978). The effect of social isolation on the aggressive and sexual behaviour in a cichlid fish, Haplochromis burtoni. Behaviour *65*, 43-61.

Asahina, K., Watanabe, K., Duistermars, B.J., Hoopfer, E., González, C.R., Eyjólfsdóttir, E.A., Perona, P., and Anderson, D.J. (2014). Tachykinin-expressing neurons control male-specific aggressive arousal in Drosophila. Cell *156*, 221-235.

Belenioti, M., and Chaniotakis, N. (2020). Aggressive behaviour of Drosophila suzukii in relation to environmental and social factors. Scientific Reports *10*, 7898.

Billeter, J.-C., Villella, A., Allendorfer, J.B., Dornan, A.J., Richardson, M., Gailey, D.A., and Goodwin, S.F. (2006). Isoform-specific control of male neuronal differentiation and behavior in Drosophila by the fruitless gene. Current Biology *16*, 1063-1076.

Brown, M.K., Strus, E., and Naidoo, N. (2017). Reduced sleep during social isolation leads to cellular stress and induction of the unfolded protein response. Sleep *40*, zsx095.

Burtis, K.C., and Baker, B.S. (1989). Drosophila doublesex gene controls somatic sexual differentiation by producing alternatively spliced mRNAs encoding related sex-specific polypeptides. Cell *56*, 997-1010.

Cacioppo, J.T., Cacioppo, S., Capitanio, J.P., and Cole, S.W. (2015). The neuroendocrinology of social isolation. Annual review of psychology *66*, 733-767.

Cacioppo, J.T., and Hawkley, L.C. (2009). Perceived social isolation and cognition. Trends in cognitive sciences *13*, 447-454.

Cacioppo, J.T., Hughes, M.E., Waite, L.J., Hawkley, L.C., and Thisted, R.A. (2006). Loneliness as a specific risk factor for depressive symptoms: cross-sectional and longitudinal analyses. Psychology and aging *21*, 140.

Chatterjee, S.S., Uppendahl, L.D., Chowdhury, M.A., Ip, P.-L., and Siegal, M.L. (2011). The female-specific doublesex isoform regulates pleiotropic transcription factors to pattern genital development in Drosophila. Development *138*, 1099-1109.

Chen, J., Jin, S., Chen, D., Cao, J., Ji, X., Peng, Q., and Pan, Y. (2021). Fruitless tunes functional flexibility of courtship circuitry during development. Elife *10*, e59224.

Chen, X., Rahman, R., Guo, F., and Rosbash, M. (2016). Genome-wide identification of neuronal activity-regulated genes in Drosophila. elife *5*, e19942.

Clough, E., Jimenez, E., Kim, Y.-A., Whitworth, C., Neville, M.C., Hempel, L.U., Pavlou, H.J., Chen, Z.-X., Sturgill, D., and Dale, R.K. (2014). Sex-and tissue-specific functions of Drosophila doublesex transcription factor target genes. Developmental cell *31*, 761-773.

Clowney, E.J., Iguchi, S., Bussell, J.J., Scheer, E., and Ruta, V. (2015). Multimodal chemosensory
circuits controlling male courtship in Drosophila. Neuron *87*, 1036-1049.

Cohen, S. (2004). Social relationships and health. American psychologist *59*, 676.

Cushing, B.S., and Kramer, K.M. (2005). Mechanisms underlying epigenetic effects of early social
experience: the role of neuropeptides and steroids. Neuroscience & Biobehavioral Reviews *29*, 1089-
1105.

Cusumano, P., Klarsfeld, A., Chélot, E., Picot, M., Richier, B., and Rouyer, F. (2009). modulated visual
inputs and cryptochrome define diurnal behavior in Drosophila. Nature neuroscience *12*, 1431-1437.

Dalton, J.E., Fear, J.M., Knott, S., Baker, B.S., McIntyre, L.M., and Arbeitman, M.N. (2013). Male-
specific Fruitless isoforms have different regulatory roles conferred by distinct zinc finger DNA binding
domains. Bmc Genomics *14*, 1-15.

Dankert, H., Wang, L., Hoopfer, E.D., Anderson, D.J., and Perona, P. (2009). Automated monitoring and
analysis of social behavior in Drosophila. Nature methods *6*, 297-303.

Dauwalder, B. (2011). The roles of fruitless and doublesex in the control of male courtship. International
review of neurobiology *99*, 87-105.

Davie, K., Janssens, J., Koldere, D., De Waegeneer, M., Pech, U., Kreft, Ł., Aibar, S., Makhzami, S.,
Christiaens, V., and González-Blas, C.B. (2018). A single-cell transcriptome atlas of the aging Drosophila
brain. Cell *174*, 982-998. e920.

de Melo Sene, F. (1977). Effect of social isolation on behavior of Drosophila silvestris from Hawaii.

Deanhardt, B., Duan, Q., Du, C., Soeder, C., Morlote, A., Garg, D., Saha, A., Jones, C.D., and Volkan,
P.C. (2023). Social experience and pheromone receptor activity reprogram gene expression in sensory
neurons. G3: Genes, Genomes, Genetics *13*, jkad072.

Deluca, A., Bascom, B., Planas, D.A.K., Kocher, M.A., Torres, M., and Arbeitman, M.N. (2025).
Contribution of neurons that express fruitless and Clock transcription factors to behavioral rhythms and
courtship. iScience *28*.

Demir, E., and Dickson, B.J. (2005). fruitless splicing specifies male courtship behavior in Drosophila.
Cell *121*, 785-794.

Dickson, B.J. (2008). Wired for sex: the neurobiology of Drosophila mating decisions. Science *322*, 904-
909.

Dow-Edwards, D., MacMaster, F.P., Peterson, B.S., Niesink, R., Andersen, S., and Braams, B.R. (2019).
Experience during adolescence shapes brain development: From synapses and networks to normal and
pathological behavior. Neurotoxicology and teratology *76*, 106834.

Du, C., Sotelo Fonseca, J.E., Jia, L., Scott, S., Okuwa, S., Koruk, S., Jones, C., and Volkan, P.C. (2025).
Pheromone circuits and transcriptional cascades modulating transcriptional and chromatin states in the
Drosophila central brain with social experience. bioRxiv, 2025.2009. 2012.675922.

Duhart, J.M., Inami, S., and Koh, K. (2023). Many faces of sleep regulation: beyond the time of day and
prior wake time. The FEBS journal *290*, 931-950.

Dukas, R. (2005). Experience improves courtship in male fruit flies. Animal Behaviour *69*, 1203-1209.

Eban-Rothschild, A., and Bloch, G. (2015). The colony environment modulates sleep in honey bee
workers. Journal of Experimental Biology *218*, 404-411.

Ewels, P.A., Peltzer, A., Fillinger, S., Patel, H., Alneberg, J., Wilm, A., Garcia, M.U., Di Tommaso, P.,
and Nahnsen, S. (2020). The nf-core framework for community-curated bioinformatics pipelines. Nature
biotechnology *38*, 276-278.

Eyjolfsdottir, E., Branson, S., Burgos-Artizzu, X.P., Hoopfer, E.D., Schor, J., Anderson, D.J., and Perona,
P. (2014). Detecting social actions of fruit flies. Paper presented at: Computer Vision–ECCV 2014: 13th
European Conference, Zurich, Switzerland, September 6-12, 2014, Proceedings, Part II 13 (Springer).

Flavell, S.W., and Greenberg, M.E. (2008). Signaling mechanisms linking neuronal activity to gene
expression and plasticity of the nervous system. Annu Rev Neurosci *31*, 563-590.

Friard, O., and Gamba, M. (2016). BORIS: a free, versatile open-source event-logging software for
video/audio coding and live observations. Methods in ecology and evolution *7*, 1325-1330.

Fujii, S., Emery, P., and Amrein, H. (2017). SIK3–HDAC4 signaling regulates Drosophila circadian male
sex drive rhythm via modulating the DN1 clock neurons. Proceedings of the National Academy of
Sciences *114*, E6669-E6677.

Fujita, N., Nagata, Y., Nishiuchi, T., Sato, M., Iwami, M., and Kiya, T. (2013). Visualization of neural
activity in insect brains using a conserved immediate early gene, Hr38. Current Biology *23*, 2063-2070.

Ganguly-Fitzgerald, I., Donlea, J., and Shaw, P.J. (2006). Waking experience affects sleep need in
Drosophila. Science *313*, 1775-1781.

General, O.o.t.S. (2023). Our epidemic of loneliness and isolation: The US Surgeon General's Advisory
on the healing effects of social connection and community [Internet].

Germain, P.-L., Lun, A., Meixide, C.G., Macnair, W., and Robinson, M.D. (2022). Doublet identification
in single-cell sequencing data using scDblFinder. f1000research *10*, 979.

Goncharova, A., Bragina, Y.V., Fedotov, S., and Kamyshev, N. (2016). Influence of group rearing on
sexual behavior of Drosophila melanogaster males. Journal of Evolutionary Biochemistry and Physiology
52, 454-462.

Goodwin, S.F., Taylor, B.J., Villella, A., Foss, M., Ryner, L.C., Baker, B.S., and Hall, J.C. (2000).
Aberrant splicing and altered spatial expression patterns in fruitless mutants of Drosophila melanogaster.
Genetics *154*, 725-745.

Han, C., Peng, Q., Su, X., Xing, L., Ji, X., and Pan, Y. (2022). A male-specific doublesex isoform reveals
an evolutionary pathway of sexual development via distinct alternative splicing mechanisms.
Communications Biology *5*, 728.

Hildreth, P.E. (1965). Doublesex, a recessive gene that tranforms both males and females of Drosophila
into intersexes. Genetics *51*, 659.

Hoffman, H.S., Boskoff, K.J., Eiserer, L.A., and Klein, S.H. (1975). Isolation-induced aggression in
newly hatched ducklings. Journal of Comparative and Physiological Psychology *89*, 447.

Hueston, C.E., Olsen, D., Li, Q., Okuwa, S., Peng, B., Wu, J., and Volkan, P.C. (2016). Chromatin
modulatory proteins and olfactory receptor signaling in the refinement and maintenance of fruitless
expression in olfactory receptor neurons. PLoS biology *14*, e1002443.

Inagaki, H.K., Jung, Y., Hoopfer, E.D., Wong, A.M., Mishra, N., Lin, J.Y., Tsien, R.Y., and Anderson,
D. (2014). Optogenetic control of Drosophila using a red-shifted channelrhodopsin reveals experience-
dependent influences on courtship. Nature methods *11*, 325-332.

Jeste, D.V., Lee, E.E., and Cacioppo, S. (2020). Battling the modern behavioral epidemic of loneliness:
suggestions for research and interventions. JAMA psychiatry *77*, 553-554.

Jung, Y., Kennedy, A., Chiu, H., Mohammad, F., Claridge-Chang, A., and Anderson, D.J. (2020). Neurons that function within an integrator to promote a persistent behavioral state in Drosophila. Neuron *105*, 322-333. e325.

Kabra, M., Robie, A.A., Rivera-Alba, M., Branson, S., and Branson, K. (2013). JAABA: interactive machine learning for automatic annotation of animal behavior. Nature methods *10*, 64-67.

Kacsoh, B.Z., Bozler, J., and Bosco, G. (2018). Drosophila species learn dialects through communal living. PLoS genetics *14*, e1007430.

Karam, C.S., Jones, S.K., and Javitch, J.A. (2020). Come Fly with Me: An overview of dopamine receptors in Drosophila melanogaster. Basic & clinical pharmacology & toxicology *126*, 56-65.

Kim, W.J., Jan, L.Y., and Jan, Y.N. (2013). A PDF/NPF neuropeptide signaling circuitry of male Drosophila melanogaster controls rival-induced prolonged mating. Neuron *80*, 1190-1205.

Kimura, K.-i., Hachiya, T., Koganezawa, M., Tazawa, T., and Yamamoto, D. (2008). Fruitless and doublesex coordinate to generate male-specific neurons that can initiate courtship. Neuron *59*, 759-769.

Koganezawa, M., Kimura, K.-i., and Yamamoto, D. (2016). The neural circuitry that functions as a switch for courtship versus aggression in Drosophila males. Current Biology *26*, 1395-1403.

Kohatsu, S., Koganezawa, M., and Yamamoto, D. (2011). Female contact activates male-specific interneurons that trigger stereotypic courtship behavior in Drosophila. Neuron *69*, 498-508.

Kopp, A. (2012). Dmrt genes in the development and evolution of sexual dimorphism. Trends in Genetics *28*, 175-184.

Lee, G., Foss, M., Goodwin, S.F., Carlo, T., Taylor, B.J., and Hall, J.C. (2000). Spatial, temporal, and sexually dimorphic expression patterns of the fruitless gene in the Drosophila central nervous system. Journal of neurobiology *43*, 404-426.

Levine, J.D., Funes, P., Dowse, H.B., and Hall, J.C. (2002). Resetting the circadian clock by social experience in Drosophila melanogaster. Science *298*, 2010-2012.

Li, Q., Barish, S., Okuwa, S., Maciejewski, A., Brandt, A.T., Reinhold, D., Jones, C.D., and Volkan, P.C. (2016). A functionally conserved gene regulatory network module governing olfactory neuron diversity. PLoS genetics *12*, e1005780.

Li, W., Wang, Z., Syed, S., Lyu, C., Lincoln, S., O'Neil, J., Nguyen, A.D., Feng, I., and Young, M.W. (2021). Chronic social isolation signals starvation and reduces sleep in Drosophila. Nature *597*, 239-244.

Lin, S.L. (2023). The "loneliness epidemic", intersecting risk factors and relations to mental health help-seeking: A population-based study during COVID-19 lockdown in Canada. Journal of affective disorders *320*, 7-17.

Liu, D., Rahman, M., Johnson, A., Amo, R., Tsutsui-Kimura, I., Sullivan, Z.A., Pena, N., Talay, M., Logeman, B.L., and Finkbeiner, S. (2025). A hypothalamic circuit underlying the dynamic control of social homeostasis. Nature, 1-11.

Liu, W., Liang, X., Gong, J., Yang, Z., Zhang, Y.-H., Zhang, J.-X., and Rao, Y. (2011). Social regulation of aggression by pheromonal activation of Or65a olfactory neurons in Drosophila. Nature neuroscience *14*, 896-902.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome biology *15*, 1-21.

Luciano, D., and Lore, R. (1975). Aggression and social experience in domesticated rats. Journal of Comparative and Physiological Psychology *88*, 917.

Mabuchi, Y., Cui, X., Xie, L., Kim, H., Jiang, T., and Yapici, N. (2023). Visual feedback neurons fine-tune Drosophila male courtship via GABA-mediated inhibition. Current Biology *33*, 3896-3910. e3897.

Matsumoto, K., Pinna, G., Puia, G., Guidotti, A., and Costa, E. (2005). Social isolation stress-induced aggression in mice: a model to study the pharmacology of neurosteroidogenesis. Stress *8*, 85-93.

McInnes, L., Healy, J., and Melville, J. (2018). Umap: Uniform manifold approximation and projection for dimension reduction. arXiv preprint arXiv:180203426.

Meissner, G.W., Luo, S.D., Dias, B.G., Texada, M.J., and Baker, B.S. (2016). Sex-specific regulation of Lgr3 in Drosophila neurons. Proceedings of the National Academy of Sciences *113*, E1256-E1265.

Mellert, D.J., Knapp, J.-M., Manoli, D.S., Meissner, G.W., and Baker, B.S. (2010). Midline crossing by gustatory receptor neuron axons is regulated by fruitless, doublesex and the Roundabout receptors. Development *137*, 323-332.

Murphy, M.W., Zarkower, D., and Bardwell, V.J. (2007). Vertebrate DM domain proteins bind similar DNA sequences and can heterodimerize on DNA. BMC molecular biology *8*, 1-14.

Nässel, D.R., and Wu, S.-F. (2022). Cholecystokinin/sulfakinin peptide signaling: conserved roles at the intersection between feeding, mating and aggression. Cellular and molecular life sciences *79*, 188.

Neville, M.C., Nojima, T., Ashley, E., Parker, D.J., Walker, J., Southall, T., Van de Sande, B., Marques, A.C., Fischer, B., and Brand, A.H. (2014). Male-specific fruitless isoforms target neurodevelopmental genes to specify a sexually dimorphic nervous system. Current Biology *24*, 229-241.

Pan, Y., and Baker, B.S. (2014). Genetic identification and separation of innate and experience-dependent courtship behaviors in Drosophila. Cell *156*, 236-248.

Pan, Y., Meissner, G.W., and Baker, B.S. (2012). Joint control of Drosophila male courtship behavior by motion cues and activation of male-specific P1 neurons. Proceedings of the National Academy of Sciences *109*, 10065-10070.

Park, A., Croset, V., Otto, N., Agarwal, D., Treiber, C.D., Meschi, E., Sims, D., and Waddell, S. (2022). Gliotransmission of D-serine promotes thirst-directed behaviors in Drosophila. Current Biology *32*, 3952-3970. e3958.

Patop, I.L., Anduaga, A.M., Bussi, I.L., Ceriani, M.F., and Kadener, S. (2023). Organismal landscape of clock cells and circadian gene expression in Drosophila. bioRxiv.

Peng, Q., Chen, J., Su, X., Wang, R., Han, C., and Pan, Y. (2022). The sex determination gene doublesex is required during adulthood to maintain sexual orientation. Journal of genetics and genomics= Yi chuan xue bao *49*, 165-168.

Pereira, T.D., Tabris, N., Matsliah, A., Turner, D.M., Li, J., Ravindranath, S., Papadoyannis, E.S., Normand, E., Deutsch, D.S., and Wang, Z.Y. (2022). SLEAP: A deep learning system for multi-animal pose tracking. Nature methods *19*, 486-495.

R Core Team, R. (2013). R: A language and environment for statistical computing.

Rideout, E.J., Dornan, A.J., Neville, M.C., Eadie, S., and Goodwin, S.F. (2010). Control of sexual differentiation and behavior by the doublesex gene in Drosophila melanogaster. Nature neuroscience *13*, 458-466.

Ryner, L.C., Goodwin, S.F., Castrillon, D.H., Anand, A., Villella, A., Baker, B.S., Hall, J.C., Taylor, B.J., and Wasserman, S.A. (1996). Control of male sexual behavior and sexual orientation in Drosophila by the fruitless gene. Cell *87*, 1079-1089.

Sánchez, M.M., Hearn, E.F., Do, D., Rilling, J.K., and Herndon, J.G. (1998). Differential rearing affects corpus callosum size and cognitive function of rhesus monkeys. Brain research *812*, 38-49.

Satija, R., Farrell, J.A., Gennert, D., Schier, A.F., and Regev, A. (2015). Spatial reconstruction of single-
cell gene expression data. Nature biotechnology *33*, 495-502.

Sethi, S., Lin, H.-H., Shepherd, A.K., Volkan, P.C., Su, C.-Y., and Wang, J.W. (2019). Social context
enhances hormonal modulation of pheromone detection in Drosophila. Current Biology *29*, 3887-3898.
e3884.

Stockinger, P., Kvitsiani, D., Rotkopf, S., Tirián, L., and Dickson, B.J. (2005). Neural circuitry that
governs Drosophila male courtship behavior. Cell *121*, 795-807.

Takayanagi-Kiya, S., Shioya, N., Nishiuchi, T., Iwami, M., and Kiya, T. (2023). Cell assembly analysis
of neural circuits for innate behavior in Drosophila melanogaster using an immediate early gene
stripe/egr-1. Proceedings of the National Academy of Sciences *120*, e2303318120.

Tweedie, S., Ashburner, M., Falls, K., Leyland, P., McQuilton, P., Marygold, S., Millburn, G., Osumi-
Sutherland, D., Schroeder, A., and Seal, R. (2009). FlyBase: enhancing Drosophila gene ontology
annotations. Nucleic acids research *37*, D555-D559.

Ueda, A., and Kidokoro, Y. (2002). Aggressive behaviours of female Drosophila melanogaster are
influenced by their social experience and food resources. Physiological Entomology *27*, 21-28.

Ueda, A., and Wu, C.-F. (2009). Effects of social isolation on neuromuscular excitability and aggressive
behaviors in Drosophila: altered responses by Hk and gsts1, two mutations implicated in redox regulation.
Journal of neurogenetics *23*, 378-394.

Villella, A., and Hall, J.C. (1996). Courtship anomalies caused by doublesex mutations in Drosophila
melanogaster. Genetics *143*, 331-344.

von Philipsborn, A.C., Jörchel, S., Tirian, L., Demir, E., Morita, T., Stern, D.L., and Dickson, B.J. (2014).
Cellular and behavioral functions of fruitless isoforms in Drosophila courtship. Current Biology *24*, 242-
251.

Von Philipsborn, A.C., Liu, T., Jai, Y.Y., Masser, C., Bidaye, S.S., and Dickson, B.J. (2011). Neuronal
control of Drosophila courtship song. Neuron *69*, 509-522.

Vrontou, E., Nilsen, S.P., Demir, E., Kravitz, E.A., and Dickson, B.J. (2006). fruitless regulates
aggression and dominance in Drosophila. Nature neuroscience *9*, 1469-1471.

Wang, L., Dankert, H., Perona, P., and Anderson, D.J. (2008). A common genetic target for
environmental and heritable influences on aggressiveness in Drosophila. Proceedings of the National
Academy of Sciences *105*, 5657-5663.

Wang, Z.Y., McKenzie-Smith, G.C., Liu, W., Cho, H.J., Pereira, T., Dhanerawala, Z., Shaevitz, J.W., and
Kocher, S.D. (2022). Isolation disrupts social interactions and destabilizes brain development in
bumblebees. Current Biology *32*, 2754-2764. e2755.

Waterbury, J.A., Jackson, L.L., and Schedl, P. (1999). Analysis of the doublesex female protein in
Drosophila melanogaster: role in sexual differentiation and behavior and dependence on intersex.
Genetics *152*, 1653-1667.

West, A.E., and Greenberg, M.E. (2011). Neuronal activity–regulated gene transcription in synapse
development and cognitive function. Cold Spring Harbor perspectives in biology *3*, a005744.

Wickham, H., and Sievert, C. (2009). ggplot2: elegant graphics for data analysis, Vol 10 (springer New
York).

Wu, S., Guo, C., Zhao, H., Sun, M., Chen, J., Han, C., Peng, Q., Qiao, H., Peng, P., and Liu, Y. (2019).
Drosulfakinin signaling in fruitless circuitry antagonizes p1 neurons to regulate sexual arousal in
Drosophila. Nat Commun 10: 4770.

Yamamoto, D., and Koganezawa, M. (2013). Genes and circuits of courtship behaviour in Drosophila
males. Nature Reviews Neuroscience *14*, 681-692.

Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R package for comparing
biological themes among gene clusters. Omics: a journal of integrative biology *16*, 284-287.

Zelikowsky, M., Hui, M., Karigo, T., Choe, A., Yang, B., Blanco, M.R., Beadle, K., Gradinaru, V.,
Deverman, B.E., and Anderson, D.J. (2018). The neuropeptide Tac2 controls a distributed brain state
induced by chronic social isolation stress. Cell *173*, 1265-1279. e1219.

Zhang, L., Lear, B.C., Seluzicki, A., and Allada, R. (2009). The CRYPTOCHROME photoreceptor gates
PDF neuropeptide signaling to set circadian network hierarchy in Drosophila. Current Biology *19*, 2050-
2055.

Zhang, W., Guo, C., Chen, D., Peng, Q., and Pan, Y. (2018). Hierarchical control of Drosophila sleep,
courtship, and feeding behaviors by male-specific P1 neurons. Neuroscience bulletin *34*, 1105-1110.

Zhao, H., Jiang, X., Ma, M., Xing, L., Ji, X., and Pan, Y. (2024). A neural pathway for social modulation
of spontaneous locomotor activity (SoMo-SLA) in Drosophila. Proceedings of the National Academy of
Sciences *121*, e2314393121.

Zhao, S., Deanhardt, B., Barlow, G.T., Schleske, P.G., Rossi, A.M., and Volkan, P.C. (2020). Chromatin-
based reprogramming of a courtship regulator by concurrent pheromone perception and hormone
signaling. Science advances *6*, eaba6913.

Zhou, C., Franconville, R., Vaughan, A.G., Robinett, C.C., Jayaraman, V., and Baker, B.S. (2015).
Central neural circuitry mediating courtship song perception in male Drosophila. Elife *4*, e08477.

Zhou, C., Pan, Y., Robinett, C.C., Meissner, G.W., and Baker, B.S. (2014). Central brain neurons
expressing doublesex regulate female receptivity in Drosophila. Neuron *83*, 149-163.

1210