Species interactions determine plasmid persistence in a 3-member bacterial community

Kaitlin A. Schaal[1]*, Ying-Jie Wang[2], Johannes Nauta[3], Manlio De Domenico[3], Shai Pilosof[2], James P. J. Hall[1]*

Correspondence to kschaal@liverpool.ac.uk; j.p.j.hall@liverpool.ac.uk

Affiliations

1. Department of Evolution, Ecology and Behaviour, Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool UK
2. Department of Life Sciences, Ben-Gurion University of the Negev, Be'er-Sheva, Israel
3. Department of Physics & Astronomy 'Galileo Galilei', University of Padua, Padua, Italy

Author contributions

Conceptualisation: all authors. Data curation: KS. Formal analysis: KS, JPJH. Funding acquisition: MdeD, SP, JPJH. Investigation: KS. Methodology: KS. Project administration: MdeD, SP, JPJH. Supervision: MdeD, SP, JPJH. Visualisation: KS. Writing — original draft: all authors. Writing — review & editing: all authors.

Abstract. Microbial communities are shaped by complex forces, including interspecies interactions and the effects of mobile genetic elements such as plasmids. How these forces interact to affect community responses to environmental perturbations remains unclear, particularly considering the qualitatively different natures of competitive and plasmid transfer interactions. We investigated the role of bacterial and plasmid interaction networks in community responses to single and multiple environmental perturbations, using a model community of three bacteria and two plasmids grown in rich media over five 2-day transfers. Bacterial interactions were the primary driver of community response, to the extent that plasmids were not always retained in the community even when they carried relevant resistance genes. Overall, our results indicate that while bacterial and plasmid interactions both shape community responses to environmental perturbations, bacterial interactions may be the primary drivers of community dynamics.

Introduction

Microbial communities are governed by diverse interactions. Prominent amongst these are the interactions between different species and strains. Organisms can interact antagonistically, competing for the limited resources necessary for growth and replication, or producing factors that kill or impede the growth of competitors (Granato et al. 2019). Organisms can also act mutualistically, facilitating one another's growth by cross-feeding and cooperation to utilise resources (Lawrence et al. 2012; Evans et al. 2020; van den Berg et al. 2022; Bisesi et al. 2025). However, interactions in microbiomes are not restricted to interactions between cellular organisms. Superimposed on the various species- and strain-level interactions are interactions between cells and their cognate mobile genetic elements (MGEs), which include bacteriophage and plasmids (Hall et al. 2022). Through such interactions, MGEs can exert significant effects on a community. MGEs can act as parasites, burdening community members in a manner that can prevent a species or strain from dominating (as is the case with so-called Kill-The-Winner dynamics in bacteriophage) (Thingstad 2000; Castledine and Buckling 2024). Conversely, MGEs can confer context-dependent beneficial traits on the microbes that host them, including resistance to environmental perturbations like antibiotics or heavy metals, the ability to metabolise novel resources, or traits allowing colonisation of new niches (Eberhard 1990; Scanlan et al. 2015).

Central to MGE-microbe interactions is the concept of host range: the ability of an MGE to enter and replicate within a given recipient cell type (its 'host'). In the context of beneficial MGE-host interactions, strains that are susceptible to acquiring MGEs can rapidly gain new traits which might accelerate adaptation to changing environments. Hosts of MGEs act as reservoirs of adaptive traits that can be disseminated to others in the community (Hall

et al. 2016), enhancing community-wide adaptation, and strains which can acquire multiple MGEs ('hub' or 'bridge' strains) constitute a location where recombination between MGEs could enable new combinations of functionality (Lam et al. 2019; Wyres et al. 2019). However, bridge strains may suffer excessively from metabolic and other burdens associated with carriage of multiple MGEs, ultimately resulting in their out-competition by other organisms, and local extinction (San Millan et al. 2014; Igler et al. 2022), with potential consequences for MGE persistence.

   To experimentally address how the presence of a bridge strain affects community dynamics and the persistence of MGEs, we developed a simple microbial community consisting of three bacterial strains (*Pseudomonas fluorescens* SBW25, *Escherichia coli* MG1655, and *P. putida* KT2440) and two plasmids (pQBR57, which confers mercury resistance, and pKJK5, which confers kanamycin resistance) (Figure 1A). Under our experimental conditions, each plasmid can be hosted by two strains: pQBR57 by *P. fluorescens* SBW25 and *P. putida* KT2440, and pKJK5 by *E. coli* MG1655 and *P. putida* KT2440. As *P. putida* KT2440 can host both plasmids (and can harbour both at the same time) it can be considered a bridge strain, whereas the other two strains are considered 'peripheral' strains. We tracked bacterial and plasmid populations, under non-selective conditions, selective levels of mercury, selective levels of kanamycin, and both stressors in combination, for ~38 generations. We compared the results with communities lacking plasmids, lacking the bridge strain, or lacking both, to ascertain the contribution of each partner to the outcome (Figure 1B). We found that the presence of plasmids increased community resilience in the face of environmental stresses, but species interactions played a predominant role in the dynamics, occasionally driving plasmid extinction even under positive selection.

Methods

Bacterial strains and plasmids. We used *Pseudomonas fluorescens* SBW25, *Pseudomonas putida* KT2440, and *Escherichia coli* MG1655. SBW25 (Rainey et al. 1994; Bailey et al. 1995) and the plasmid pQBR57 (Lilley et al. 1994, 1996; Lilley and Bailey 1997) were isolated independently from a sugar beet field in Oxfordshire, UK. KT2440 is a derivative of *P. putida* mt-2 (Bagdasarian et al. 1981). MG1655 is a derivative of *E. coli* K-12 (Blattner et al. 1997). Interactions between SBW25 and KT2440 have been previously studied, as well as dynamics between the two strains involving pQBR57, which carries the *merA* gene for mercury resistance (Hall et al. 2015, 2016). pQBR57 is considered a narrow host range plasmid, as transconjugants have only been isolated using Pseudomonads (Hall et al. 2015). pKJK5 is a broad host range plasmid known to infect both KT2440 and various *E. coli* strains

96	(Sengeløv et al. 2001; Bahl et al. 2007) including MG1655 (Klümper et al. 2015). Though
97	pKJK5 has been reported to conjugate into SBW25 (Sünderhauf et al. 2023), we have been
98	unable to retrieve SBW25(pKJK5) transconjugants either under the experimental conditions
99	outlined in this study, or an extended range of conditions including broth and filter mating,
100	and selection for pKJK5-encoded traits. We therefore consider SBW25 as incapable of
101	hosting pKJK5 in our experimental system. Both plasmids are relevant for soil microbial
102	communities (Hall et al. 2015; Klümper et al. 2015).
103	We used derivatives of each strain to enable enumeration of different populations:
104	SBW25-Sm^R*lacZ* (Hall et al. 2015), KT2440-Gm^R (Hall et al. 2016), and wild-type MG1655
105	(which carries the *lacZ* gene). We used pQBR57::tdTomato (Wright et al. 2024) and
106	pKJK5::GFP, which has the GFP gene inserted into the tetracycline resistance gene and is
107	resistant to kanamycin but not tetracycline (Klümper et al. 2015). This allowed us to track the
108	three bacterial strains and plasmid carriage throughout the experiment: the plasmids by
109	transfluorescence on a blue light box, and the strains by plating on agar with 250 µg/ml
110	streptomycin and incubating at 28°C (for SBW25) and on agar with 50 µg/ml X-Gal and
111	incubating at 37°C (for KT2440 and MG1655).
112	
113	*Sub-population construction.* For brevity, we named our different community treatments SMK
114	(for SBW25 + MG1655 + KT2440) or SM (for SBW25 + MG1655) for communities containing
115	or lacking the bridge strain respectively, appended with '/p' for communities in which
116	plasmids were initially present. Our experimental design (Figure 1) used a link-balanced
117	initial community composition (Table 1), where all populations in treatments including
118	plasmids start 50% plasmid-free, with the remaining 50% evenly comprised of plasmid-
119	carrying sub-populations (one each for SBW25 and MG1655, three for KT2440). We
120	therefore constructed all possible donor strains (Table 2) prior to starting the experiment,
121	using the patch conjugation method from the Barrick lab (Leonard and Perreau 2024). To
122	test effects of plasmid carriage on host growth rate, we grew overnight cultures of each
123	plasmid-free and plasmid-carrying strain, transferred them by pin replicator (~1.5 µl volume)
124	into 200 µl LB broth in a 96-well plate, and performed a 48-hr growth curve at 28 °C in a
125	Tecan Infinite F Nano+ plate reader, reading OD$_{600}$ every 15 minutes.
126	

Table 1. Initial community compositions

Community	Population	Fraction	Sub-population	Fraction
SM	SBW25	0.5		
	MG1655	0.5		
SMK	SBW25	0.33		
	MG1655	0.33		
	KT2440	0.33		
SM/p	SBW25	0.5	SBW25	0.25
			SBW25(pQBR57)	0.25
	MG1655	0.5	MG1655	0.25
			MG1655(pKJK5)	0.25
SMK/p	SBW25	0.33	SBW25	0.165
			SBW25(pQBR57)	0.165
	MG1655	0.33	MG1655	0.165
			MG1655(pKJK5)	0.165
	KT2440	0.33	KT2440	0.165
			KT2440(pQBR57)	0.055
			KT2440(pKJKJ5)	0.055
			KT2440(pKJK5, pQBR57)	0.055

127
128

Table 2. Plasmid-hosting bacterial sub-populations

Strain	Plasmid(s)
SBW25-Sm^R*lacZ*	pQBR57::tdTomato
MG1655	pKJK5::GFP
KT2440-Gm^R	pQBR57::tdTomato
KT2440-Gm^R	pKJK5::GFP
KT2440-Gm^R	pQBR57::tdTomato and pKJK5::GFP

130	
131	*Experiment.* We streaked out all strains on LB agar plates and incubated overnight at 28°C.
132	We selected 6 independent colonies of each and grew them overnight in LB liquid at 28°C
133	and 180 RPM in 50-ml Falcon tubes in an Innova 42 incubator shaker. We standardized
134	overnight cultures to OD_{600} 1 in LB then diluted 10x, and assembled communities by
135	combining strains in the ratios shown in Table 2 in a total volume of 600 µl. We prepared a
136	96-well plate with 180 µl LB per well and inoculated each replicate community four times
137	(once per environmental treatment), to a total of 200 µl. Communities were started with 5 x
138	10^7 CFU. The plating pattern was randomized. No environmental stresses were added
139	during the first cycle. We incubated the experimental plate for 2 days at 28°C and 640 RPM
140	on a Stuart Microtitre SSL-5 orbital shaker then transferred to the next cycle by Boekel
141	Scientific pin replicator (transfer volume ~1 µl). All subsequent 96-well plates were prepared
142	with 200 µl LB per well, with ¼ of wells containing 10 µM HgCl2 [Hg(II)], ¼ containing 12.5
143	µg/ml kanamycin sulfate, and ¼ containing 10 µM HgCl2 [Hg(II)] and 12.5 µg/ml kanamycin
144	sulfate; the remaining ¼ received no perturbation. Concentrations of mercury and kanamycin
145	were the minimum inhibitory concentrations for the non-plasmid-hosting strains. We
146	passaged the cultures 4 times over a total of 10 days. We quantified population and sub-
147	populations of the initial communities and after cycles 1, 3, and 5 (days 2, 6, and 10), with a
148	lower limit of detection of 5 x 10^5, or >0.5% of community carrying capacity. With a transfer
149	volume of 1% of the community, we expected sub-populations below our limit of detection to
150	be removed from the community across transfers.
151	
152	*Statistical analysis.* We summarized overall dynamics of bacteria and plasmids for statistical
153	analysis by calculating trapezoidal area under the curve ('AUC', Figures S2-3), to avoid
154	analytical complications that would have been introduced in timecourse analysis by
155	autocorrelation between timepoints. To assess the effects of community structure and
156	environmental stresses, we used general linear models followed by Tukey HSD tests. To
157	assess extinction probability of SBW25, we ranked extinction as a binary event and
158	performed a Fisher's exact test. We analysed the data using R version 4.3.3 (R Core Team
159	2024) and RStudio version 2024.12.1+563 (RStudio Team 2020). We organized the data
160	using the *tidyr* (Wickham et al. 2024) and *dplyr* (Wickham et al. 2023) packages, and
161	visualized them using *ggplot2* (Wickham 2016) and *viridis* (Garnier et al. 2024).
162	
163	
164	

Results

Plasmids rescue total community population size from stress. First, we considered communities lacking the bridge strain. In the absence of plasmids, exogenous stresses (mercury, kanamycin, or both) decreased cumulative community size (Figure S4; ANOVA main effect of environment on area under the curve, $F_{[3,80]} = 72.54$, *p*-value < 0.0001), an effect principally driven by the impact of kanamycin (post-hoc Tukey tests effect of kanamycin *p*-values < 0.0002; effect of mercury *p*-value = 0.055). As expected, adding resistance-encoding plasmids to the community ameliorated these effects (ANOVA main effect of plasmids, $F_{[1,80]} = 173.4$, *p*-value < 0.0001), effectively restoring pre-stress community population size (post-hoc Tukey tests, *p*-values > 0.89). Plasmid addition had no significant effect on total community size in the absence of stress (post-hoc Tukey test, *p*-value = 1). At the community level, therefore, plasmids provided robustness to stress.

Plasmids stabilized the two-member community. Under non-stressful conditions, plasmid addition did not significantly alter the dynamics of either of the two community members (Figure 2; area under the curve analysis, post-hoc Tukey HSD tests, *p*-value > 0.9 for all comparisons), possibly because the presence of plasmid-free subpopulations for each strain could mask any negative effect of plasmid burden on the overall population counts. Stresses particularly impacted SBW25, which performed worse under all stress treatments (post-hoc Tukey HSD tests, Hg *p* = 0.03, Kn and Hg + Kn *p*-values < 0.001). As predicted given the host ranges of the plasmids, for SBW25, plasmid addition mitigated only the effect of mercury stress (SM community vs SM/p community, *p*-value = 0.0006), and could not rescue the population when faced with kanamycin stress (SM community vs SM/p community, *p*-values = 1), where SBW25's population fell below the detection limit of the experiment in each case.

By contrast, MG1655 was primarily affected only by kanamycin stress (Hg *p* = 0.25, Kn and Hg + Kn *p*-values < 0.003), and though kanamycin significantly reduced MG1655's population size (post-hoc Tukey tests, *p*-value < 0.0001), the strain remained at appreciable levels even when plasmids were absent from the community (AUC difference from non-stressful conditions of -7.34 for Kn vs. -21.41 for Hg + Kn), possibly by emergence of a *de novo* mutant with increased kanamycin resistance. Plasmid addition improved MG1655's growth under kanamycin conditions, even rescuing MG1655 from dual stress (*p*-values < 0.0001).

In a two-member community with no between-species HGT, plasmids were therefore able to rescue their hosts from relevant stresses. Despite the inability to host the plasmid conferring mercury resistance, MG1655 demonstrated an ability to survive under mercury

conditions regardless of plasmid presence, or under dual stress when it carries the plasmid
conferring kanamycin resistance (Figure 2).

Adding the bridge strain destabilized the community. We predicted that community members
present in the host ranges of multiple plasmids (i.e. bridge strains, Figure 1) may experience
higher plasmid burden, altering their dynamics and influence on the overall community. We
therefore investigated the dynamics of communities which included a third member, *P. putida*
KT2440, which can host both pQBR57 and pKJK5 and thus acts as a bridge strain when the
plasmids are present.

First, we considered the effect of the bridge strain on plasmid-free communities. The
presence of the bridge strain did not significantly influence total population size (Figure S4;
ANOVA, $F_{[1,80]}$ = 2.2, *p*-value = 0.13), and MG1655 was unaffected by KT2440's presence
across environmental treatments (Figures 2 & 3, ANOVA, $F_{[1,80]}$ = 0.095, *p*-value = 0.75).
However, SBW25 was significantly impacted, experiencing population decline over the
course of the experiment when the bridge strain was present (Figures 2 & 3, ANOVA, main
effect of bridge strain, $F_{[1,80]}$ = 150.7 *p*-value < 0.0001). This may have been due to niche
overlap between the two *Pseudomonas*, resulting in nutrient competition (Zhang et al. 2012).
The effect of KT2440 on SBW25 was observed both in non-stressful and mercury conditions
(post-hoc Tukey HSD tests, *p*-values < 0.002) and drove SBW25 below the detection limit
under non-stressful conditions (Figures 2 & 3, first column; Fisher's Exact test, *p*-value =
0.004). (We did not detect an effect in treatments containing kanamycin (*p*-values > 0.99)
because SBW25 quickly fell below the detection limit regardless of KT2440 presence (Figure
2)). The presence of the bridge strain therefore shaped the overall community principally by
influencing the dynamics and persistence of specific community members, in this case
SBW25.

Stress type determines plasmid effects on community stability. Next, we investigated how
the presence of plasmids and bridge strain interacted to affect community members. Hosting
pKJK5 imposes a cost relative to hosting pQBR57 (Figure S1; ANOVA for differences in
growth dynamics of KT2440 sub-populations, $F_{[3,23]}$ = 3.86, *p*-value = 0.023; post-hoc Tukey
HSD test for pKJK5 vs pQBR57, *p*-value = 0.034). However, we expected plasmids to
facilitate growth of strains under conditions that selected for the cognate resistance traits.

Under mercury conditions, in contrast to the two-species communities, the presence
of plasmids in the three-species communities did not improve survival of the pQBR57-
hosting strains, SBW25 and KT2440, . In fact, while there was no significant effect on
SBW25 survival (Figure 3; post-hoc Tukey HSD tests for SMK vs SMK/p, *p*-value >0.99),
plasmid presence exerted a negative effect on KT2440, with loss from the community more

| likely when plasmids were present than when they were absent (Figure S5; Fisher's Exact
| test, *p*-value = 0.015). This outcome suggests that the interaction of plasmids and
| environmental conditions can influence strain interactions in non-intuitive ways.
| By contrast, kanamycin stress did enhance the growth of strains capable of hosting
| the plasmid carrying the cognate kanamycin resistance gene, MG1655 and KT2440.
| Plasmids improved MG1655 survival under both conditions containing kanamycin (Figures 2
| & 3 "Kn" and "Hg + Kn"; post-hoc Tukey HSD tests, *p*-value < 0.0001), while plasmids
| rescued KT2440 under kanamycin stress (Figure 3 "Kn"; post-hoc Tukey HSD test, *p* <
| 0.001), but not dual stress, even though KT2440 is capable of hosting both plasmids and
| thus expressing resistances to both stresses (post-hoc Tukey HSD test, *p*-value > 0.99).
| Overall, the bridge strain appears to have destabilized the community via its effect on
| SBW25, and the plasmids could in some cases exacerbate this effect, resulting in loss of
| species diversity from the community.
|
| *Plasmids were not always rescued by the corresponding stress*. Communities which
| contained plasmids had increased robustness to environmental stressors to which those
| plasmids conferred resistance. We therefore expected plasmids to increase in frequency
| within host populations experiencing a stress for the plasmid carries a resistance gene (i.e.
| plasmid selection (Hall et al. 2020)). However, we did not always find this to be the case. For
| pQBR57, which encodes resistance to mercury, dynamics were determined by the hosting
| strain, presence of the bridge strain in the community, and environmental conditions (Figure
| 4; ANOVA, $F_{[1,63]}$ = 32.41 strain, $F_{[1,63]}$ = 23.32 bridge, $F_{[3,63]}$ = 25.28 stress, $F_{[3,63]}$ = 21.98
| bridge x stress interaction, *p*-values < 0.0001). As expected, plasmid frequency declined in
| the absence of selection, likely owing to fitness cost. But while mercury selection was able to
| prevent decline of pQBR57 (Figure 4, post-hoc Tukey HSD tests, *p*-values < 0.0001), this
| was true only for the 2-member community, where SBW25 was the only available host (post-
| hoc Tukey HSD tests, *p*-values < 0.0001). The presence of the bridge strain removed any
| benefit provided by environmental selection (post-hoc Tukey HSD tests, comparison
| between SM/p and SMK/p for Hg, *p*-value < 0.0001), and pQBR57 declined similarly across
| all environmental treatments (post-hoc Tukey HSD tests, *p*-values > 0.3).
| By contrast, while host strain had a significant impact on pKJK5 (Figure 4; ANOVA,
| $F_{[1,60]}$ = 285.03, *p*-value < 0.0001), the overall presence of the bridge strain in the community
| did not (ANOVA, $F_{[1,60]}$ = 0.072, *p*-value = 0.82). Kanamycin selection had a significant effect
| on pKJK5 (post-hoc Tukey HSD tests comparing kanamycin and non-kanamycin
| environments, *p*-values < 0.0002), and though there was a quantitative difference between
| the kanamycin and dual selection in the three-member (but not the two-member) community,
| thanks to the contribution by the bridge strain (*p*-values < 0.0002), the overall dynamics were

276	qualitatively similar. In most cases pKJK5 dynamics follow those of MG1655, its predominant
277	host, with little contribution from the bridge strain.
278	

Discussion

280	Here, we explored the interaction of several forces — inter-species interactions, mobile
281	genetic elements, and environmental change — to assess their interacting effects on
282	microbial community composition and dynamics. We found that interactions between strains,
283	known to have a predominant role in community dynamics, fundamentally impacted the
284	success of their resident MGEs. Such effects could even overwhelm any benefits provided
285	by the MGEs in changing, stressful environments. For example, we found that the mercury
286	resistance plasmid pQBR57 was lost even from environments containing mercury because
287	its resident hosts were outcompeted, while *E. coli* MG1655 persisted (allowing maintenance
288	of its plasmid pKJK5) regardless of environmental selection pressures. Such patterns
289	challenge our intuition as to drivers of MGEs in communities, indicating that MGE dynamics
290	are fundamentally determined by internal community interactions, and not only by extrinsic
291	selective pressures.
292	We manipulated community composition by the addition of a bridge strain, *P. putida*
293	KT2440, that could harbour both possible plasmids. The main effect of the bridge strain on
294	community dynamics was through competition with *P. fluorescens* SBW25, which suffered
295	from reduced abundances across environmental and plasmid conditions. Previous work
296	found that competitive interactions between these species exerted only a mild effect on
297	abundances in a structured soil environment (Hall et al. 2018), or in heterogenous liquid
298	culture, and though the effects were intensified in well-mixed rich media ((Zhang et al. 2012),
299	as used in the current study) they tended not to result in local extinction of SBW25.
300	Additional competitive pressure from the other community member, MG1655, may have
301	been sufficient to exclude SBW25 from stressful or bridge-strain-containing conditions in our
302	experiments; such competition may be further intensified in minimal media, as opposed to
303	the rich medium used here (Goldford et al. 2018). Natural communities, evolving to
304	specialize on a single labile carbon source, tend to converge on a conserved composition at
305	the Family level, consisting principally of Enterobacteriaceae and Pseudomonadaceae owing
306	to the metabolic capabilities of with each family (Goldford et al. 2018). Similar species-
307	sorting interactions may be at work in our simpler communities, resulting in the exclusion of
308	the less competitive Pseudomonad.
309	While we expected that the relative success of the bridge strain ought to promote the
310	persistence of both plasmids, plasmid maintenance was generally reduced in communities
311	containing the bridge strain. Previous work found that *P. putida* KT2440 was a less
312	favourable host than *P. fluorescens* SBW25 for plasmid pQBR57, even if the fitness effects

in each strain are similar ((Hall et al. 2016; Kottara et al. 2018), Fig. S2). The alternative plasmid pKJK5 was likewise more costly to the bridge strain than to its peripheral strain *E. coli* MG1655 (Fig. S2). Competitive interactions between organisms, and variance in MGE stability across species and strains (possibly depending on co-infection status of the host (San Millan et al. 2014; Igler et al. 2022)), thus interact to have consequential effects for maintenance and spread of MGEs that could not be predicted from MGE host ranges alone (Alonso-del Valle et al. 2021). Similarly, we expected that selection for plasmid-borne resistances would have promoted the corresponding plasmid, and the organisms able to host that plasmid. However, our results suggest that selection for plasmid-borne resistances was insufficient to rescue plasmids, or hosts, in a community context — most strikingly in communities treated with mercury, where both hosts of the mercury resistance plasmid went extinct. Higher concentrations of mercury may have rescued pQBR57 (Hall et al. 2020), suggesting that local plasmid fate may vary across heterogeneous environments.

Embedding bacterial-plasmid interactions in a community has been shown to impact plasmid fate. Walker-Sünderhauf et al. (2023) showed that maintenance of pKJK5 in two different microbes isolated from potting soil (species of *Stenotrophomonas* and *Variovorax*) was disrupted in the context of a five-species community, leading to plasmid loss from these organisms. Kottara et al. (2021) similarly showed that plasmid prevalence in a focal species could be reduced by the epidemiological dilution effect of living in a multi-species community, as conjugation events were more likely to result in plasmid transfer to non-focal organisms. Besides shaping microbial community composition, strain competition (i.e. features of the bacterial interaction matrix 'A', Fig. 1) can exert consequential influences on subsequent opportunities for HGT, by driving the dynamics of the vectors of gene transfer, sometimes to extinction. The influence of strain competition on plasmid survival might explain why some plasmids may have evolved to manipulate expression of secreted competitive molecules (Di Venanzio et al. 2019; Billane et al. 2022).

In a reciprocal manner, microbial community dynamics can be profoundly altered in the context of horizontal gene transfer. HGT can have a positive effect on community diversity and microbiome stability by enabling beneficial resistances to transfer across community members, facilitating their survival (Coyte et al. 2022). The transfer of costly genes can also maintain community stability in a process of 'dynamic neutrality', reducing variance in growth rates between species and mitigating competition (Zhu et al. 2024), although this process may be influenced by changes in species interactions. Our work suggests that structure in the HGT network can have a critical effect modulating the impact of HGT on communities, allowing – or preventing – mobile genes to hitch-hike on competitive community members.

The non-*Pseudomonas* species in our experiment, *E. coli* MG1655, offers a contrast. MG1655 was successful at competition regardless of the presence or absence of the bridge strain, and was able to maintain pKJK5 successfully. Moreover, MG1655 was successful at surviving in the face of single stresses even when plasmids were absent. Our concentrations of selective agents were determined based on single-species MIC measurements; growth in a multi-species community — together with the large population size and standing genetic diversity gained during the first growth cycle (from which stresses were absent) — likely allowed survival beyond these minimal concentrations. Moreover, mercury resistance is achieved by environmental detoxification (Barkay et al. 2010), a 'public good' which could benefit other, nearby community members lacking specific mercury resistance genes. Interestingly, MG1655 could survive dual stresses only in communities with plasmids, even though it couldn't maintain both plasmids itself, suggesting that resistance to one stress improved the capacity of this strain to cope with a heterologous stress, possibly by enabling sufficient population size for *de novo* mutants to emerge.

Overall, our work demonstrates the importance of considering multiple factors to understand community dynamics. Further, we suggest that plasmids are at risk of being lost from a community due to impacts on their hosts, even when environmental conditions would be expected to select for them. This idea suggests additional complexity that may be at play in plasmid evolution.

Acknowledgements

This work was supported by a Human Frontiers Science Programme Project Grant to SP, MdeD, JPJH (RGY0064/2022). JPJH is supported by an MRC Career Development Award (MR/W02666X/1). We thank Uli Klümper (TU Dresden) for providing pKJK5::GFP, and Mike Bottery (University of Manchester) for *E. coli* MG1655.

References

Alonso-del Valle, A., R. León-Sampedro, J. Rodríguez-Beltrán, J. DelaFuente, M. Hernández-García, P. Ruiz-Garbajosa, R. Cantón, R. Peña-Miller, and A. San Millán. 2021. Variability of plasmid fitness effects contributes to plasmid persistence in bacterial communities. Nat Commun 12:2653.

Bagdasarian, M., R. Lurz, B. Rückert, F. C. H. Franklin, M. M. Bagdasarian, J. Frey, and K. N. Timmis. 1981. Specific-purpose plasmid cloning vectors II. Broad host range, high

copy number, RSF 1010-derived vectors, and a host-vector system for gene cloning
in Pseudomonas. Gene 16:237–247.

Bahl, M. I., L. H. Hansen, A. Goesmann, and S. J. Sørensen. 2007. The multiple antibiotic
resistance IncP-1 plasmid pKJK5 isolated from a soil environment is phylogenetically
divergent from members of the previously established α, β and δ sub-groups.
Plasmid 58:31–43.

Bailey, M. J., A. K. Lilley, I. P. Thompson, P. B. Rainey, and R. J. Ellis. 1995. Site directed
chromosomal marking of a fluorescent pseudomonad isolated from the phytosphere
of sugar beet; stability and potential for marker gene transfer. Molecular Ecology
4:755–764.

Barkay, T., K. Kritee, E. Boyd, and G. Geesey. 2010. A thermophilic bacterial origin and
subsequent constraints by redox, light and salinity on the evolution of the microbial
mercuric reductase. Environmental Microbiology 12:2904–2917.

Billane, K., E. Harrison, D. Cameron, and M. A. Brockhurst. 2022. Why do plasmids
manipulate the expression of bacterial phenotypes? Philosophical Transactions of the
Royal Society B: Biological Sciences 377.

Bisesi, A. T., R. P. Carlson, L. Cotner, and W. R. Harcombe. 2025. Metabolic remodeling of
microorganisms by obligate intracellular parasites alters mutualistic community
composition. BioRxiv.

Blattner, F. R., G. Plunkett, C. A. Bloch, N. T. Perna, V. Burland, M. Riley, J. Collado-Vides, J.
D. Glasner, C. K. Rode, G. F. Mayhew, J. Gregor, N. W. Davis, H. A. Kirkpatrick, M. A.
Goeden, D. J. Rose, B. Mau, and Y. Shao. 1997. The Complete Genome Sequence
of *Escherichia coli* K-12. Science 277:1453–1462.

Castledine, M., and A. Buckling. 2024. Critically evaluating the relative importance of phage
in shaping microbial community composition. Trends in Microbiology 32:957–969.

Coyte, K. Z., C. Stevenson, C. G. Knight, E. Harrison, J. P. J. Hall, and M. A. Brockhurst.
2022. Horizontal gene transfer and ecological interactions jointly control microbiome
stability. PLoS Biology 20:e3001847.

Di Venanzio, G., A. L. Flores-Mireles, J. J. Calix, M. F. Haurat, N. E. Scott, L. D. Palmer, R. F.
Potter, M. E. Hibbing, L. Friedman, B. Wang, G. Dantas, E. P. Skaar, S. J. Hultgren,
and M. F. Feldman. 2019. Urinary tract colonization is enhanced by a plasmid that
regulates uropathogenic Acinetobacter baumannii chromosomal genes. Nat Commun
10:2763.

Eberhard, W. G. 1990. Evolution in Bacterial Plasmids and Levels of Selection. The
Quarterly Review of Biology 65:3–22.

Evans, R., A. P. Beckerman, R. C. T. Wright, S. McQueen-Mason, N. C. Bruce, and M. A.
Brockhurst. 2020. Eco-evolutionary Dynamics Set the Tempo and Trajectory of

Metabolic Evolution in Multispecies Communities. Current Biology 30:4984-4988.e4.

Elsevier Ltd.

Garnier, S., N. Ross, R. Rudis, A. P. Camargo, M. Sciaini, and C. Scherer. 2024. viridis(Lite)

- Colorblind-Friendly Color Maps for R.

Goldford, J. E., N. Lu, D. Bajić, S. Estrela, M. Tikhonov, A. Sanchez-Gorostiaga, D. Segrè, P.

Mehta, and A. Sanchez. 2018. Emergent simplicity in microbial community assembly.

Science 361:469–474.

Granato, E. T., T. A. Meiller-Legrand, and K. R. Foster. 2019. The Evolution and Ecology of

Bacterial Warfare. Current Biology 29:R521–R537. Elsevier Ltd.

Hall, J. P. J., E. Harrison, and D. A. Baltrus. 2022. Introduction: The secret lives of microbial

mobile genetic elements. Philosophical Transactions of the Royal Society B:

Biological Sciences 377.

Hall, J. P. J., E. Harrison, and M. A. Brockhurst. 2018. Competitive species interactions

constrain abiotic adaptation in a bacterial soil community. Evolution Letters 2:580–

589.

Hall, J. P. J., E. Harrison, A. K. Lilley, S. Paterson, A. J. Spiers, and M. A. Brockhurst. 2015.

Environmentally co-occurring mercury resistance plasmids are genetically and

phenotypically diverse and confer variable context-dependent fitness effects.

Environmental Microbiology 17:5008–5022.

Hall, J. P. J., E. Harrison, K. Pärnänen, M. Virta, and M. A. Brockhurst. 2020. The Impact of

Mercury Selection and Conjugative Genetic Elements on Community Structure and

Resistance Gene Transfer. Frontiers in Microbiology 11:1–14.

Hall, J. P. J., A. J. Wood, E. Harrison, and M. A. Brockhurst. 2016. Source-sink plasmid

transfer dynamics maintain gene mobility in soil bacterial communities. Proceedings

of the National Academy of Sciences of the United States of America 113:8260–

8265.

Igler, C., J. S. Huisman, B. Siedentop, S. Bonhoeffer, and S. Lehtinen. 2022. Plasmid co-

infection: linking biological mechanisms to ecological and evolutionary dynamics.

Phil. Trans. R. Soc. B 377:20200478.

Klümper, U., L. Riber, A. Dechesne, A. Sannazzarro, L. H. Hansen, S. J. Sørensen, and B. F.

Smets. 2015. Broad host range plasmids can invade an unexpectedly diverse

fraction of a soil bacterial community. ISME Journal 9:934–945.

Kottara, A., L. Carrilero, E. Harrison, J. P. J. Hall, and M. A. Brockhurst. 2021. The dilution

effect limits plasmid horizontal transmission in multispecies bacterial communities.

Microbiology 167:1–7.

Kottara, A., J. P. J. Hall, E. Harrison, and M. A. Brockhurst. 2018. Variable plasmid fitness
effects and mobile genetic element dynamics across Pseudomonas species. FEMS
Microbiology Ecology 94:1–7.
Lam, M. M. C., K. L. Wyres, R. R. Wick, L. M. Judd, A. Fostervold, K. E. Holt, and I. H. Löhr.
2019. Convergence of virulence and MDR in a single plasmid vector in MDR
Klebsiella pneumoniae ST15. Journal of Antimicrobial Chemotherapy 74:1218–1222.
Lawrence, D., F. Fiegna, V. Behrends, J. G. Bundy, A. B. Phillimore, T. Bell, and T. G.
Barraclough. 2012. Species interactions alter evolutionary responses to a novel
environment. PLOS biology 10:e1001330.
Leonard, S., and J. Perreau. 2024. General Conjugation Protocol.
Lilley, A. K., and M. J. Bailey. 1997. The acquisition of indigenous plasmids by a genetically
marked pseudomonad population colonizing the sugar beet phytosphere is related to
local environmental conditions. Appl Environ Microbiol 63:1577–1583.
Lilley, A. K., M. J. Bailey, M. J. Day, and J. C. Fry. 1996. Diversity of mercury resistance
plasmids obtained by exogenous isolation from the bacteria of sugar beet in three
successive years. FEMS Microbiology Ecology 20:211–227.
Lilley, A. K., J. C. Fry, M. J. Day, and M. J. Bailey. 1994. In situ transfer of an exogenously
isolated plasmid between *Pseudomonas* spp. in sugar beet rhizosphere.
Microbiology 140:27–33.
R Core Team. 2024. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria.
Rainey, P. B., M. J. Bailey, and I. P. Thompson. 1994. Phenotypic and genotypic diversity of
fluorescent pseudomonads isolated from field-grown sugar beet. Microbiology
140:2315–2331.
RStudio Team. 2020. RStudio: Integrated Development for R. RStudio, PBC, Boston, MA.
San Millan, A., K. Heilbron, and R. C. MacLean. 2014. Positive epistasis between co-
infecting plasmids promotes plasmid survival in bacterial populations. The ISME
Journal 8:601–612.
Scanlan, P. D., A. R. Hall, G. Blackshields, V. P. Friman, M. R. Davis, J. B. Goldberg, and A.
Buckling. 2015. Coevolution with bacteriophages drives genome-wide host evolution
and constrains the acquisition of abiotic-beneficial mutations. Molecular Biology and
Evolution 32:1425–1435.
Sengeløv, G., K. J. Kristensen, A. H. Sørensen, N. Kroer, and S. J. Sørensen. 2001. Effect of
genomic location on horizontal transfer of a recombinant gene cassette between
Pseudomonas strains in the rhizosphere and spermosphere of barley seedlings.
Current Microbiology 42:160–167.

Sünderhauf, D., U. Klümper, E. Pursey, E. R. Westra, W. H. Gaze, and S. Van Houte. 2023.

Removal of AMR plasmids using a mobile, broad host-range CRISPR-Cas9 delivery

tool. Microbiology 169.

Thingstad, T. F. 2000. Elements of a theory for the mechanisms controlling abundance,

diversity, and biogeochemical role of lytic bacterial viruses in aquatic systems.

Limnology & Oceanography 45:1320–1328.

van den Berg, N. I., D. Machado, S. Santos, I. Rocha, J. Chacón, W. Harcombe, S. Mitri, and

K. R. Patil. 2022. Ecological modelling approaches for predicting emergent properties

in microbial communities. Nature Ecology and Evolution 6:855–865. Springer US.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

Wickham, H., R. François, L. Henry, K. Müller, and V. Davis. 2023. dplyr: A Grammar of Data

Manipulation.

Wickham, H., D. Vaughan, and M. Girlich. 2024. tidyr: Tidy Messy Data.

Wright, R. C. T., A. J. Wood, M. J. Bottery, K. J. Muddiman, S. Paterson, E. Harrison, M. A.

Brockhurst, and J. P. J. Hall. 2024. A chromosomal mutation is superior to a plasmid-

encoded mutation for plasmid fitness cost compensation. PLoS Biol 22:e3002926.

Wyres, K. L., R. R. Wick, L. M. Judd, R. Froumine, A. Tokolyi, C. L. Gorrie, M. M. C. Lam, S.

Duchêne, A. Jenney, and K. E. Holt. 2019. Distinct evolutionary dynamics of

horizontal gene transfer in drug resistant and virulent clones of Klebsiella

pneumoniae. PLoS Genet 15:e1008114.

Zhang, Q.-G., R. J. Ellis, and H. C. J. Godfray. 2012. THE EFFECT OF A COMPETITOR ON

A MODEL ADAPTIVE RADIATION. Evolution 66:1985–1990.

Zhu, S., J. Hong, and T. Wang. 2024. Horizontal gene transfer is predicted to overcome the

diversity limit of competing microbial species. Nat Commun 15:800.

518
519
520
Figure 1. Experimental design. The community interaction network (A) consists of three bacterial
strains (lower layer) and two plasmids (upper layer). The bacteria are *P. fluorescens* SBW25 (green
oval), *P. putida* KT2440 (blue oval), and *E. coli* MG1655 (purple oval). The plasmids are pQBR57 (red
dotted ring) and pKJK5 (green dashed ring). KT2440 is able to host both plasmids, making it a bridge
strain within the community. We analysed dynamics of four communities under four environmental
conditions, including three stress treatments (B).

527
528
Figure 2. Bacterial dynamics in the 2-member community. Population sizes of SBW25 and
MG1655 over 10 days under different environmental conditions (environmental perturbations were
included from day 2 onward). The upper row of panels shows dynamics when plasmids were absent
from the community, and the lower row shows bacterial dynamics when plasmids were present. Small
dots represent replicate values, large dots represent means of 6 biological replicates, and vertical
lines represent 95% confidence intervals of the means. Stresses were added after day 2.

Figure 3. Bacterial dynamics in the 3-member community. Population sizes of SBW25, KT2440,

and MG1655 over 10 days under different environmental conditions (environmental perturbations

were included from day 2 onward). The upper row of panels shows dynamics when plasmids were

absent from the community, and the lower row shows bacterial dynamics when plasmids were

present. Small dots represent replicate values, large dots represent means of 6 biological replicates,

and vertical lines represent 95% confidence intervals of the means. Stresses were added after day 2.

Figure 4. Plasmid frequency dynamics. Frequencies of plasmid-carrying subpopulations of the host

bacteria over 10 days under different environmental conditions, where pQBR57 confers Hg resistance

and pKJK5 confers Kn resistance. A frequency of 1 indicates that all cells of that host contain the

plasmid. Colors represent plasmid status (hosting one plasmid, the other, or both plasmids) and

shapes represent host species. The upper row of panels shows plasmid frequency dynamics in the 2-

member community, and the lower row shows dynamics in the 3-member community. Small dots

represent replicate values, large dots represent means of 6 biological replicates, and vertical lines

represent 95% confidence intervals of the means

553
554
Figure S1. Plasmid fitness costs across hosts. Growth profiles of all possible subpopulations
(each bacterial strain with each possible plasmid status) over 48 hours, showing the effect of the
plasmids on host growth dynamics. Each time step represents 15 minutes. There are 3 biological
replicates of each subpopulation.

Figure S2. Areas under the curve (AUCs) for bacterial populations. (A) AUCs for strain dynamics

across communities and environmental conditions. (B) AUCs for total community dynamics across

environmental conditions. Bars are averages of six biological replicates; error bars show 95%

confidence intervals.

Figure S3. Areas under the curve (AUCs) for plasmids. AUCs for plasmids by host strain, across communities and environmental stresses. Bars are averages of six biological replicates; error bars show 95% confidence intervals.

Figure S4. Total community size across environments. The total size of all bacterial populations in each community over 10 days, as affected by the environmental conditions. Small dots represent replicate values, large dots represent means of 6 biological replicates, and vertical lines represent 95% confidence intervals of the means

578 **Figure S5. Strain survival.** Fraction of replicates, out of six total replicates, in which each strain was

579 detected at the final timepoint, across communities and environmental conditions. Limit of detection

580 was 5×10^5 CFUs.