Species interactions determine plasmid persistence in a 3-member bacterial community

Kaitlin A. Schaal[1]*, Ying-Jie Wang[2], Johannes Nauta[3], Manlio De Domenico[3], Shai Pilosof[2], James P. J. Hall[1]*

Correspondence to kschaal@liverpool.ac.uk; j.p.j.hall@liverpool.ac.uk

Affiliations
1. Department of Evolution, Ecology and Behaviour, Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, UK
2. Department of Life Sciences, Ben-Gurion University of the Negev, Be'er-Sheva, Israel
3. Department of Physics & Astronomy 'Galileo Galilei', University of Padua, Padua, Italy

Author contributions

Conceptualization: all authors. Data curation: KS. Formal analysis: KS, JPJH. Funding acquisition: MdeD, SP, JPJH. Investigation: KS. Methodology: KS. Project administration: MdeD, SP, JPJH. Supervision: MdeD, SP, JPJH. Visualization: KS. Writing — original draft: all authors. Writing — review & editing: all authors.

Acknowledgements

This work was supported by a Human Frontiers Science Programme Project Grant to SP, MdeD, JPJH (RGY0064/2022). YJW is supported by the Kreitman School of Advanced Graduate Studies, Ben-Gurion University of the Negev. JPJH is supported by an MRC Career Development Award (MR/W02666X/1). We thank Uli Klümper (TU Dresden) for providing pKJK5::GFP, and Mike Bottery (University of Manchester) for E. coli MG1655.

Species interactions determine plasmid persistence in a 3-member bacterial community

Abstract. Microbial communities are shaped by multiple forces, including interspecies interactions and the effects of mobile genetic elements such as plasmids. How these forces interact to affect community responses to environmental perturbations remains unclear, particularly considering the qualitatively different natures and the complexity of competitive and plasmid transfer interactions. We investigated the role of bacterial and plasmid interaction networks in community responses to single and multiple environmental perturbations, using a model community of three bacteria and two plasmids grown in rich media over five 2-day transfers. Bacterial interactions were the primary driver of community response, to the extent that plasmids were not always retained in the community even when they carried relevant resistance genes. Overall, our results indicate that while bacterial and plasmid interactions both shape community responses to environmental perturbations, bacteria-bacteria interactions may be the primary driver of community dynamics and their response to perturbations.

Introduction

Microbial communities are governed by diverse interactions. Microbes can interact antagonistically, competing for the limited resources necessary for growth and replication, or producing factors that kill or impede the growth of competitors (1). They can also interact mutualistically, facilitating one another's growth by cross-feeding and cooperation to use resources (2–5). However, interactions in microbiomes are not restricted to interactions between cellular organisms. Superimposed on the various species- and strain-level interactions are interactions between cells and their cognate mobile genetic elements (MGEs), which include bacteriophages and plasmids (6). Through such interactions, MGEs can exert significant effects on a community. MGEs can act as parasites, burdening community members in a manner that can prevent a species or strain from dominating (e.g., Kill-The-Winner dynamics in bacteriophages) (7,8). Conversely, MGEs can confer context-dependent beneficial traits on the microbes that host them, including resistance to environmental perturbations like antibiotics or heavy metals, the ability to metabolize novel resources, or traits allowing colonization of new niches (9,10).

Central to MGE-microbe interactions is the concept of host range: the ability of an MGE to enter and replicate within a given recipient cell type (its 'host'). In the context of beneficial MGE-host interactions, strains that are susceptible to acquiring MGEs can rapidly gain new traits which might accelerate adaptation to changing environments. Hosts of MGEs act as reservoirs of adaptive traits that can be disseminated to others in the community (11), enhancing community-wide adaptation, and strains which can acquire multiple MGEs ('bridge' strains) and thereby enable recombination between MGEs could enable new combinations of functionality (12,13). However, bridge strains may suffer excessively from metabolic and other burdens associated with carriage of multiple MGEs, ultimately resulting in their out-competition by other organisms and local extinction (14,15), with potential consequences for MGE persistence. Bridge strains are therefore expected to face a trade-off shaped by the costs of multiple plasmids versus the environment-dependent benefits of plasmids' genetic cargo.

To experimentally address how the presence of a bridge strain affects community dynamics and the persistence of MGEs, we developed a simple microbial community consisting of three bacterial strains (*Pseudomonas fluorescens* SBW25, *Escherichia coli* MG1655, and *P. putida* KT2440) and two plasmids (pQBR57, which confers mercury resistance, and pKJK5, which confers kanamycin resistance) (Figure 1A). Under our experimental conditions, each plasmid can be hosted by two strains: pQBR57 by *P. fluorescens* SBW25 and *P. putida* KT2440, and pKJK5 by *E. coli* MG1655 and *P. putida* KT2440. As *P. putida* KT2440 can host both plasmids (and can harbour both at the same time) we consider it a 'bridge' strain, and the other two as 'peripheral' strains. We tracked bacterial and plasmid populations, under non-selective conditions, selective levels of mercury, selective levels of kanamycin, and under combined selection from both, for ~38 generations. We compared the results with communities lacking plasmids, lacking the bridge strain, or lacking both, to ascertain the contribution of each member to community dynamics (Figure 1B). We found that the presence of plasmids increased community resilience in the face of environmental stresses. However, bacterial species interactions played a predominant role in the dynamics, occasionally driving plasmid extinction even under positive selection.

Figure 1. Experimental design. The community interaction network (A) consists of three bacterial strains (lower layer) and two plasmids (upper layer). The bacteria are *P. fluorescens* SBW25 (green oval), *P. putida* KT2440 (blue oval), and *E. coli* MG1655 (purple oval). The plasmids are pQBR57 (red dotted ring) and pKJK5 (green dashed ring). KT2440 can host both plasmids, making it a bridge strain within the community. We analysed dynamics of four communities under four environmental conditions, including three stress treatments (B).

Methods

Bacterial strains and plasmids. We used *Pseudomonas fluorescens* SBW25, *Pseudomonas putida* KT2440, and *Escherichia coli* MG1655. SBW25 (16,17) and the plasmid pQBR57 (18–20) were isolated independently from a sugar beet field in Oxfordshire, UK. KT2440 is a derivative of *P. putida* mt-2 (21). MG1655 is a derivative of *E. coli* K-12 (22). Interactions between SBW25 and KT2440 have been previously studied, as well as dynamics between the two strains involving pQBR57, which carries the *merA* gene for mercury resistance (11,23). pQBR57 is considered a narrow host range plasmid, as transconjugants have only been isolated using Pseudomonads (23). pKJK5 is a broad host range plasmid known to infect both KT2440 and various *E. coli* strains (24,25) including MG1655 (26). Though pKJK5 has been reported to conjugate into SBW25 (27), we have been unable to retrieve SBW25 (pKJK5) transconjugants either under the experimental conditions outlined in this study, or an extended range of conditions including broth and filter mating, and selection for

pKJK5-encoded traits. We therefore consider SBW25 as incapable of hosting pKJK5 in the
context of our experimental system. Both plasmids are relevant for soil microbial
communities (23,26). Strains were cultured in LB (10 g/L peptone from casein, 5 g/L yeast
extract, 10 g/L NaCl, with agar at 1.5% where required) and KB (King's B; 20 g/L proteose
peptone, 1.5 g/L MgSO$_4$, 1.15 g/L K$_2$HPO$_4$, 10 g/L glycerol, with agar at 1.2% as required) as
described.

We used derivatives of each strain to enable enumeration of different populations:
SBW25-Sm^R*lacZ* (23), KT2440-Gm^R (11), and wild-type MG1655 (which carries the *lacZ*
gene). We used pQBR57::tdTomato (28) and pKJK5::GFP, which has the GFP gene inserted
into the tetracycline resistance gene and is resistant to kanamycin but not tetracycline (26).
This allowed us to track the three bacterial strains and plasmid carriage throughout the
experiment: the plasmids by transfluorescence on a blue light box, and the strains by plating
on agar with 250 µg/ml streptomycin and incubating at 28°C (for SBW25) and on agar with
50 µg/ml X-Gal and incubating at 37°C (for KT2440 and MG1655).

Sub-population construction. For brevity, we named our different community treatments SMK
(for SBW25 + MG1655 + KT2440) or SM (for SBW25 + MG1655) for communities containing
or lacking the bridge strain respectively, appended with '/p' for communities in which
plasmids were initially present. Our experimental design (Figure 1) used a 'link-balanced'
initial community composition (Table 1), whereby all populations in treatments including
plasmids are initiated with 50% plasmid-free cells, with the remaining 50% evenly comprised
of plasmid-carrying sub-populations (one each for SBW25 and MG1655, three for KT2440).
We therefore constructed all possible donor strains (Table S1) prior to starting the
experiment, using the patch conjugation method (29). To test effects of plasmid carriage on
bacterial growth rate, we grew overnight cultures of each plasmid-free and plasmid-carrying
strain, transferred them by pin replicator (~1.5 µl volume) into 200 µl LB broth in a 96-well
plate, and performed a 48-hr growth curve at 28 °C shaking in a Tecan Infinite F Nano+ plate
reader (Tecan Group Ltd., Männedorf, Switzerland), reading OD$_{600}$ every 15 minutes. To test
the effect of environmental stressors, growth curves were also established under a range of
mercury conditions (0, 2.5, 5, 10, 20 µM).

Table 1. Initial community compositions. Fraction refers to the initial proportion of cells from each population.

Community	Population	Fraction	Sub-population	Fraction
SM	SBW25	0.5		
	MG1655	0.5		
SMK	SBW25	0.33		
	MG1655	0.33		
	KT2440	0.33		
SM/p	SBW25	0.5	SBW25	0.25
			SBW25(pQBR57)	0.25
	MG1655	0.5	MG1655	0.25
			MG1655(pKJK5)	0.25
SMK/p	SBW25	0.33	SBW25	0.165
			SBW25(pQBR57)	0.165
	MG1655	0.33	MG1655	0.165
			MG1655(pKJK5)	0.165
	KT2440	0.33	KT2440	0.165
			KT2440(pQBR57)	0.055
			KT2440(pKJKJ5)	0.055
			KT2440(pKJK5, pQBR57)	0.055

Experiment. We streaked out all strains on LB agar plates and incubated overnight at 28°C.
We selected 6 independent colonies of each and grew them overnight in LB liquid at 28°C
and 180 RPM in 50-ml Falcon tubes in an Innova 42 incubator shaker. We standardized
overnight cultures to OD_{600} 1 in LB then diluted 10x, and assembled communities by
combining strains in the ratios shown in Table S1 in a total volume of 600 µl. We prepared a
96-well plate with 180 µl LB per well and inoculated each replicate community four times
(once per environmental treatment), to a total of 200 µl. Communities were started with 5 x
10^7 CFU. The plating pattern was randomized by sorting the list of samples by the rand()
function in Microsoft Excel. No environmental stresses were added during the first growth
cycle. We incubated the experimental plate for 2 days at 28°C and 640 RPM on a Stuart
Microtiter SSL-5 orbital shaker then transferred to the next cycle by Boekel Scientific pin
replicator (transfer volume ~1 µl). All subsequent 96-well plates were prepared with 200 µl
LB per well, with ¼ of wells containing 10 µM HgCl2 [Hg(II)], ¼ containing 12.5 µg/ml

kanamycin sulfate, and ¼ containing 10 μM HgCl2 [Hg(II)] and 12.5 μg/ml kanamycin sulfate; the remaining ¼ received no perturbation. Concentrations of mercury and kanamycin were the lowest relevant stress levels for the non-plasmid-hosting strains. We passaged the cultures 4 times over a total of 10 days. We quantified population and sub-populations of the initial communities and after cycles 1, 3, and 5 (days 2, 6, and 10), with a lower limit of detection of 5 x 10^5, or >0.5% of community carrying capacity. With a transfer volume of 1% of the community, we expected sub-populations below our limit of detection to be ultimately removed from the community owing to dilution during transfers.

Statistical analysis. We summarized overall dynamics of bacteria and plasmids for statistical analysis by calculating trapezoidal area under the curve (AUC, Figures 2 & 3), to avoid analytical complications that would have been introduced in time-series analysis by autocorrelation between timepoints (30–32). To quantify the effects of community structure and environmental stresses on bacterial and plasmid dynamics, we used general linear models followed by Tukey HSD tests. To quantify the extinction probability of SBW25, we ranked extinction as a binary event and performed a Fisher's exact test. We analysed the data using R version 4.5.2 (33) and RStudio version 2026.01.1+403 (34). We organized the data using the "tidyr" (35) and "dplyr" (36) packages, analysed growth curve data with "gcplyr" (37), and visualized them using "ggplot2" (38) and "viridis" (39). All data and analysis files can be found at [public DOI after review].

Results

Plasmids rescue total community population size from stress. To investigate the effect that plasmid presence has on the overall community, we first considered communities lacking the bridge strain. In the absence of plasmids, exogenous stresses (mercury, kanamycin, or both) decreased cumulative community size (Figure S1; ANOVA main effect of environment on AUC, $F_{[3,80]}$ = 72.54, *p*-value < 0.0001), an effect principally driven by the impact of kanamycin (post-hoc Tukey tests effect of kanamycin *p*-values < 0.0002; effect of mercury *p*-value = 0.055). As expected, adding resistance-encoding plasmids to the communities ameliorated these effects (ANOVA main effect of plasmids, $F_{[1,80]}$ = 173.4, *p*-value < 0.0001), effectively restoring pre-stress community population size (post-hoc Tukey tests, *p*-values > 0.89). Plasmid addition had no significant effect on total community size in the absence of stress (post-hoc Tukey test, *p*-value = 1). At the community level, therefore, plasmids provided robustness to stress.

Figure 2. Bacterial dynamics. Population sizes of SBW25, KT2440, and MG1655 over 10 days
under different environmental conditions (environmental perturbations were included from day 2
onward). The upper row of panels shows dynamics when plasmids were absent from the community,
and the lower row shows bacterial dynamics when plasmids were present. Small dots represent
replicate values, large dots represent means of 6 biological replicates, and vertical lines represent
95% confidence intervals of the means. Stresses were added after day 2.

Plasmids stabilized the two-member community. Under non-stressful conditions, plasmid
addition did not significantly alter the dynamics of either of the two community members
(Figure 2A&E; AUC post-hoc Tukey HSD tests, *p*-value > 0.9 for all comparisons), possibly
because the presence of plasmid-free subpopulations in each strain masked the negative
effect of plasmid burden on the overall population counts. Stresses particularly impacted
SBW25, which performed worse under all stress treatments (Figure 2A-D; post-hoc Tukey
HSD tests, Hg *p* = 0.03, Kn and Hg + Kn *p*-values < 0.001). As predicted given the host
ranges of the plasmids, for SBW25 plasmid addition mitigated only the effect of mercury
stress (Figure 2B; SM community vs SM/p community, *p*-value = 0.0006), and could not
rescue the population when faced with kanamycin stress (Figure 2C; SM community vs
SM/p community, *p*-values = 1), where SBW25's population fell below the detection limit of
the experiment in each case.

In contrast, MG1655 was only significantly affected by kanamycin stress (Figure 2E-
H, Hg p = 0.25, Kn and Hg + Kn p-values < 0.003), and though kanamycin significantly
reduced MG1655's population size (Figure 2G&H; post-hoc Tukey tests, p-value < 0.0001),
the strain remained at appreciable levels even when plasmids were absent from the
community (Figure 2G&H; AUC difference from non-stressful conditions of -7.34 for Kn vs. -
21.41 for Hg + Kn), possibly by emergence of a *de novo* mutant with increased kanamycin
resistance. MG1655 clones isolated from plasmid-free populations show higher resistance to
increased kanamycin concentrations than the ancestor, but lower than what is conferred by
pKJK5 (Figure S2). Plasmid addition improved MG1655's growth under kanamycin
conditions, even rescuing MG1655 from dual stress (p-values < 0.0001).
In a two-member community with no between-species HGT, plasmids were therefore
able to rescue their hosts from relevant stresses. Despite its inability to host the plasmid
conferring mercury resistance, MG1655 demonstrated an ability to survive under mercury
conditions regardless of plasmid presence, or under dual stress when it carries the plasmid
conferring kanamycin resistance (Figure 2F&H). This may be due to mercury absorption by
the growth medium and a higher overall tolerance to mercury than SBW25 and KT2440
(Supplemental results).
202
Adding the bridge strain destabilized the community. We predicted that a bridge strain
(Figure 1) will experience higher plasmid burden, altering its dynamics compared to when
grown with no plasmids. We therefore investigated the dynamics of communities including a
third member, *P. putida* KT2440, which can host both pQBR57 and pKJK5 and thus acts as
a bridge strain when the plasmids are present.
First, we considered the effect of the bridge strain on plasmid-free communities. The
presence of the bridge strain did not significantly influence total population size (Figure S1;
ANOVA, $F_{[1,80]}$ = 2.2, p-value = 0.13), and MG1655 was unaffected by KT2440's presence
across environmental treatments (Figure 2E-H; ANOVA, $F_{[1,80]}$ = 0.095, p-value = 0.75).
However, SBW25 was significantly impacted, experiencing population decline over the
course of the experiment when the bridge strain was present (Figure 2A&B; ANOVA, main
effect of bridge strain, $F_{[1,80]}$ = 150.7 p-value < 0.0001). This difference in dynamics between
the bacterial strains may have occurred due to niche overlap between the two *Pseudomonas*
strains, resulting in nutrient competition (40). The effect of KT2440 on SBW25 was observed
both in non-stressful and mercury conditions (Figure 2A&B; post-hoc Tukey HSD tests, p-
values < 0.002) and drove SBW25 below the detection limit under non-stressful conditions
(Figure 2A; Fisher's Exact test, p-value = 0.004). We did not detect an effect in treatments
containing kanamycin (p-values > 0.99) because SBW25 quickly fell below the detection limit
regardless of KT2440 presence (Figure 2C&D). The presence of the bridge strain therefore

| shaped the overall community principally by influencing the dynamics and persistence of
| specific community members, in this case SBW25.
|
| *Stress type determines plasmid effects on community stability*. Given that plasmids
| stabilized the 2-member community and the bridge strain destabilized the plasmid-free
| community, we investigated how the presence of plasmids and bridge strain interacted to
| affect community members. Hosting pKJK5 imposes a cost on the bridge strain relative to
| hosting pQBR57 (Figure S3; ANOVA for differences in growth dynamics of KT2440 sub-
| populations, $F_{[3,23]}$ = 3.86, *p*-value = 0.023; post-hoc Tukey HSD test for pKJK5 vs pQBR57,
| *p*-value = 0.034). However, we expected plasmids to facilitate growth of their hosts under
| conditions that selected for the cognate resistance traits.
| Under mercury conditions, in contrast to the two-species communities, the presence
| of plasmids in the three-species communities did not improve survival of the pQBR57-
| hosting strains, SBW25 and KT2440. In fact, while there was no significant effect on SBW25
| survival (Figure 2B; post-hoc Tukey HSD tests for SMK vs SMK/p, *p*-value >0.99), plasmid
| presence exerted a negative effect on KT2440, whose loss from the community was more
| likely when plasmids were present than when they were absent (Figure S4; Fisher's Exact
| test, *p*-value = 0.015). This outcome suggests that the interaction between plasmids and
| environmental conditions can influence strain interactions in non-intuitive ways.
| By contrast, pKJK5 hosts MG1655 and KT2440 grew better under kanamycin stress
| when plasmids were present. Plasmids improved MG1655 survival under both conditions
| containing kanamycin (Figure 2G&H; post-hoc Tukey HSD tests, *p*-value < 0.0001), while
| plasmids rescued KT2440 under kanamycin stress (Figure 2L; post-hoc Tukey HSD test, *p* <
| 0.001), but not dual stress, even though KT2440 is capable of hosting both plasmids and
| thus expressing resistances to both stresses (Figure 2M; post-hoc Tukey HSD test, *p*-value
| > 0.99). Overall, the bridge strain appears to have destabilized the community via its effect
| on SBW25, and the plasmids could in some cases exacerbate this effect, resulting in loss of
| species diversity from the community.

Figure 3. Plasmid frequency dynamics. Frequencies of plasmid-carrying subpopulations of the host bacteria over 10 days under different environmental conditions. pQBR57 confers Hg resistance and pKJK5 confers Kn resistance. A frequency of 1 indicates that all cells of that host contain the plasmid. Colours represent plasmid status (hosting one plasmid, the other, or both plasmids) and shapes represent host species. The upper row of panels shows plasmid frequency dynamics in the 2-member community, and the lower row shows dynamics in the 3-member community. Small dots represent replicate values, large dots represent means of 6 biological replicates, and vertical lines represent 95% confidence intervals of the means. Missing data indicates where hosts were no longer detected in the community, to differentiate from plasmid frequency of 0 within the host.

Plasmids were not always rescued by the corresponding stress. Communities which contained plasmids had increased robustness to environmental stressors to which those plasmids conferred resistance. We therefore expected plasmids to increase in frequency within host populations exposed to a stressor against which the plasmid confers resistance (*i.e.*, positive plasmid selection (41)). However, we did not consistently observe this. For pQBR57, which encodes resistance to mercury, dynamics were determined by the hosting strain, presence of the bridge strain in the community, and environmental conditions (Figure 3A-D; ANOVA, $F_{[1,63]}$ = 32.41 strain, $F_{[1,63]}$ = 23.32 bridge, $F_{[3,63]}$ = 25.28 stress, $F_{[3,63]}$ = 21.98 bridge x stress interaction, *p*-values < 0.0001). As expected, plasmid frequency declined in the absence of selection, likely owing to fitness cost. While mercury selection was able to

prevent decline of pQBR57 (Figure 3B; post-hoc Tukey HSD tests, *p*-values < 0.0001), this

was true only for the 2-member community, where SBW25 was the only available host

(Figure 3B, circles; post-hoc Tukey HSD tests, *p*-values < 0.0001). The presence of the

bridge strain removed any benefit provided by environmental selection (Figure 3B; post-hoc

Tukey HSD tests, comparison between SM/p and SMK/p for Hg, *p*-value < 0.0001), and

pQBR57 declined similarly across all environmental treatments (Figure 3A-D, triangles; post-

hoc Tukey HSD tests, *p*-values > 0.3). Loss of pQBR57 from the community under mercury

conditions after day 6 (Figure 3B, triangles) suggests an interplay between bacteria-plasmid

and bacteria-bacteria interactions, which we investigated further (next section).

While host strain had a significant impact on pKJK5 (Figure 3E-H; ANOVA, $F_{[1,60]}$ =

285.03, *p*-value < 0.0001), the overall presence of the bridge strain in the community did not

(ANOVA, $F_{[1,60]}$ = 0.072, *p*-value = 0.82). Kanamycin selection had a significant effect on

pKJK5 (Figure3G&H; post-hoc Tukey HSD tests comparing kanamycin and non-kanamycin

environments, *p*-values < 0.0002). Although there was a quantitative difference in pKJK5

dynamics between the kanamycin and dual selection environments in the three-member (but

not the two-member) community, thanks to the contribution by the bridge strain (Figure

3G&H, triangles; *p*-values < 0.0002), the overall dynamics were qualitatively similar. In most

cases pKJK5 dynamics followed those of MG1655, its predominant host, with little

contribution from the bridge strain.

We expected that dual stress conditions (Figure 3D,H,M) would select for KT2440

hosting both plasmids. Indeed, this is the only condition under which we observed any

notable presence of co-hosting cells within the community (Figure 3M). However, their

increase in frequency was transient, and by day 10 pQBR57 had apparently been lost from

the community, while pKJK5 had reached fixation in both of its hosts (Figure 3H). This may

once again be due to the interplay of differing plasmid fitness costs for MG1655 and

KT2440, different impacts of mercury on the two strains, and competition between them.

Figure 4. Interplay among strain competition, plasmid hosting, and environment under mercury conditions. We tested the effect of mercury on growth of the three plasmid-free strains and the two pQBR57-hosting strains. (A) Kinetic growth curves of the five strains across a range of mercury concentrations. Solid lines indicate averages of 4 replicate freezer-stock inoculations and dashed lines with shaded areas indicate standard error. (B) Area under the curve of the growth curves in panel A. Boxes show the median (middle line) and the first and third quartiles, whiskers show the smallest and largest value no farther than 1.5 x the inter-quartile range, and dots show outliers). (C) Frequency of pQBR57 within SBW25 and KT2440 under non-stressful and mercury conditions, showing the effect of mercury on plasmid prevalence within each strain. Difference from Figure 3: here KT2440 cells carrying both plasmids are included when quantifying pQBR57 frequency.

pQBR57 loss under mercury conditions indicates interplay between bacteria-plasmid and bacteria-bacteria interactions. The unexpected loss, only under mercury conditions, of both strains able to carry the plasmid which conferred resistance to mercury (Figure 2B&K) led us to hypothesize that presence of mercury increased the cost of the plasmid (42), exacerbating the competition between SBW25 and KT2440 and driving strain extinction. To test this, we

conducted growth curves of SBW25, SBW25 + pQBR57, KT2440, KT2440 + pQBR57, and MG1655 across a range of mercury concentrations (Figure 4A). A comparison of the AUC for the plasmid-free strains shows that increasing mercury concentration has a greater impact on growth of SBW25 and KT2440 than on MG1655, indicating that MG1655 is better able to tolerate mercury stress up to and including 10 µM mercury (Figure 4B), which explains MG1655's survival under mercury stress. Next, by examining the total frequency of pQBR57 across both hosts (including KT2440 also hosting pKJK5) we found a decline in the non-stressful environment, consistent with pQBR57's known fitness cost on these strains, and increases in the mercury environment (Figure 4C). Therefore, we propose that in the mercury environment, the toxicity to SBW25 and KT2440 means that only cells harbouring the burdensome plasmid will survive, while they continue to face competition from MG1655, which is able to tolerate the stress without carrying a costly plasmid. This combined effect may also explain the predominance of MG1655 in the 3-member community under dual stress conditions (Figure 2H) over KT2440 (Figure 2M), whose population was ultimately dominated by cells hosting pKJK5 alone (Figure 3H&M). Overall, these results indicate that the combination of host-plasmid-environment interactions and inter-strain competition together shape community outcomes.

Discussion

While interactions among members shape community diversity, function, and stability, our understanding of the relative roles and cumulative impacts of different types of interactions (*e.g.*, between bacterial species or between a bacterium and a plasmid) remains limited. This gap hinders our ability to predict the responses of microbial communities to environmental change. Here, we explored the interplay of several forces — inter-species interactions, mobile genetic elements, and environmental change — on microbial community composition and dynamics. We found that interactions between organisms, known to have a predominant role in community dynamics, fundamentally impacted the success of their resident MGEs. Such effects could even overwhelm any benefits provided by the MGEs in relevant stressful environments. For example, we found that the mercury resistance plasmid pQBR57 was lost even from environments containing mercury because its resident hosts were outcompeted, while *E. coli* MG1655 persisted (allowing maintenance of its plasmid pKJK5) regardless of environmental selection pressures. Such patterns challenge our intuition as to drivers of MGEs in communities, indicating that MGE dynamics are fundamentally determined by internal community interactions, and not only by extrinsic selective pressures.

We manipulated community composition by the addition of a bridge strain, *P. putida*
KT2440, that could harbour both plasmids. The main effect of the bridge strain on
community dynamics was through competition with *P. fluorescens* SBW25, which suffered
from reduced abundances across environmental and plasmid conditions (Figure 2A-D).
Previous work found that competitive interactions between these species exerted only a mild
effect on abundances in a structured soil environment (43) or in heterogenous liquid culture,
and though the effects were intensified in well-mixed rich media ((40), as used in the current
study), they tended not to result in local extinction of SBW25. Additional competitive
pressure from the other community member, MG1655, may have been sufficient to exclude
SBW25 from stressful or bridge-strain-containing conditions in our experiments; such
competition may be further intensified in minimal media, as opposed to the rich medium
used here (44). Natural communities, evolving to specialize on a single labile carbon source,
tend to converge on a conserved composition at the Family level, consisting principally of
Enterobacteriaceae and Pseudomonadaceae owing to the metabolic capabilities of with
each family (44). Similar species-sorting interactions may be at work in our simpler
communities, resulting in the exclusion of the less competitive Pseudomonad. As we
performed our assays in well-mixed rich media, we encourage future work to investigate
community outcomes under resource limitation or in structured, soil-like environments which
are expected to better approximate natural conditions.

While we expected that the relative success of the bridge strain would promote the
persistence of both plasmids, plasmid maintenance was generally reduced in communities
containing the bridge strain. Previous work found that *P. putida* KT2440 was a less
favourable host than *P. fluorescens* SBW25 for plasmid pQBR57, even if the fitness effects
in each strain are similar ((11,45), Figure S3). The plasmid pKJK5 was more costly to the
bridge strain than to its peripheral strain *E. coli* MG1655 (Figure S3) and likewise less likely
to be maintained by KT2440. Competitive interactions between organisms, and variance in
MGE stability across species and strains (possibly depending on co-infection status of the
host (14,15)), thus interact to have consequential effects for maintenance and spread of
MGEs that could not be predicted from MGE host ranges alone (46). The plasmid dynamics
we report here reflect the net balance of transfer, cost, and loss alongside host competition;
future work should additionally quantify specific host-plasmid interactions such as
conjugation and segregational loss to further explore drivers of MGEs in communities.

We expected that selection for plasmid-borne resistances would have promoted the
corresponding plasmid, and the organisms able to host that plasmid. However, our results
suggest that selection for plasmid-borne resistances was insufficient to rescue plasmids, or
hosts, in a community context — most strikingly in communities treated with mercury, where
both hosts of the mercury resistance plasmid went extinct. The toxicity of mercury to SBW25

and KT2440 (Figure 4) selected for cells harbouring the burdensome plasmid while also allowing ongoing competition from MG1655. The combination of host-plasmid-environment interactions and inter-strain competition may therefore counteract environmental selection and determine plasmid maintenance. Higher concentrations of mercury may have rescued pQBR57 (41), suggesting that local plasmid fate may vary across heterogeneous environments.

Embedding bacterial-plasmid interactions in a community has been shown to impact plasmid fate. Walker-Sünderhauf et al. (47) showed that maintenance of pKJK5 in two different microbes isolated from potting soil (species of *Stenotrophomonas* and *Variovorax*) was disrupted in the context of a five-species community, leading to plasmid loss from these organisms. Kottara et al. (32) similarly showed that plasmid prevalence in a focal species could be reduced by the epidemiological dilution effect of living in a multi-species community, as conjugation events were more likely to result in plasmid transfer to non-focal organisms. Besides shaping microbial community composition, strain competition (*i.e.*, features of the bacterial interaction matrix 'A'; Figure 1A) can influence subsequent opportunities for HGT, by driving the dynamics of the vectors of gene transfer, sometimes to extinction. The influence of strain competition on plasmid survival might explain why some plasmids may have evolved to manipulate expression of secreted competitive molecules (48,49).

In a reciprocal manner, microbial community dynamics can be profoundly altered in the context of horizontal gene transfer. HGT can have a positive effect on community diversity and microbiome stability by enabling beneficial resistances to transfer across community members, facilitating their survival (50). The transfer of costly genes can also maintain community stability in a process of 'dynamic neutrality', reducing variance in growth rates between species and mitigating competition (51), although this process may be influenced by changes in species interactions. Our work suggests that structure in the HGT network (e.g. presence of a bridge strain, creating partly overlapping host ranges) can have a critical effect modulating the impact of HGT on communities, allowing – or preventing – mobile genes to hitch-hike on competitive community members.

The non-*Pseudomonas* species in our experiment, *E. coli* MG1655, offers a contrast. MG1655 was successful at competition regardless of the presence or absence of the bridge strain and was able to maintain pKJK5 successfully. Moreover, MG1655 was successful at surviving in the face of single stresses even when plasmids were absent. Our concentrations of selective agents were determined based on single-species MIC measurements; growth in a multi-species community, together with the large population size and standing genetic diversity gained during the first growth cycle (from which stresses were absent), likely allowed survival beyond these minimal concentrations. Moreover, mercury resistance is achieved by environmental detoxification (52), a 'public good' which could benefit other,

nearby community members lacking specific mercury resistance genes. Interestingly,

MG1655 could survive dual stresses only in communities with plasmids, even though it

couldn't maintain both plasmids itself, suggesting that resistance to one stress improved the

capacity of this strain to cope with a heterologous stress, possibly by enabling sufficient

population size for *de novo* mutants to emerge.

Our work demonstrates the importance of considering the interplay of key factors in

understanding community dynamics, not just the effects of those factors in isolation.

Bacterial community members face effects of biotic and abiotic drivers, such as competition

and cooperation, plasmid hosting, and environmental stresses, which can interact in ways

that impact their survival differently than when each driver is considered alone. Further, we

suggest that plasmids are at risk of being lost from a community due to impacts on their

hosts, even when environmental conditions would be expected to select for them. This idea

suggests additional complexity that may be at play in plasmid evolution.

References

1. Granato ET, Meiller-Legrand TA, Foster KR. The Evolution and Ecology of Bacterial Warfare. Current Biology. 2019;29(11):R521–37. doi:10.1016/j.cub.2019.04.024 PubMed PMID: 31163166.

2. Lawrence D, Fiegna F, Behrends V, Bundy JG, Phillimore AB, Bell T, et al. Species interactions alter evolutionary responses to a novel environment. PLOS biology. 2012;10(5):e1001330. doi:10.1371/journal.pbio.1001330 PubMed PMID: 22615541.

3. Evans R, Beckerman AP, Wright RCT, McQueen-Mason S, Bruce NC, Brockhurst MA. Eco-evolutionary Dynamics Set the Tempo and Trajectory of Metabolic Evolution in Multispecies Communities. Current Biology. 2020;30(24):4984-4988.e4. doi:10.1016/j.cub.2020.09.028 PubMed PMID: 33035481.

4. van den Berg NI, Machado D, Santos S, Rocha I, Chacón J, Harcombe W, et al. Ecological modelling approaches for predicting emergent properties in microbial communities. Nature Ecology and Evolution. 2022;6(JulY):855–65. doi:10.1038/s41559-022-01746-7

5. Bisesi AT, Carlson RP, Cotner L, Harcombe WR. Metabolic remodeling of microorganisms by obligate intracellular parasites alters mutualistic community composition [Internet]. BioRxiv; 2025 [cited 2025 Apr 11]. Available from: http://biorxiv.org/lookup/doi/10.1101/2025.01.29.635536 doi:10.1101/2025.01.29.635536

6. Hall JPJ, Harrison E, Baltrus DA. Introduction: The secret lives of microbial mobile genetic elements. Philosophical Transactions of the Royal Society B: Biological Sciences. 2022;377(1842). doi:10.1098/rstb.2020.0460 PubMed PMID: 34839706.

7. Thingstad TF. Elements of a theory for the mechanisms controlling abundance, diversity,
and biogeochemical role of lytic bacterial viruses in aquatic systems. Limnology &
Oceanography. 2000 Sep;45(6):1320–8. doi:10.4319/lo.2000.45.6.1320

8. Castledine M, Buckling A. Critically evaluating the relative importance of phage in
shaping microbial community composition. Trends in Microbiology. 2024 Oct;32(10):957–
69. doi:10.1016/j.tim.2024.02.014

9. Eberhard WG. Evolution in Bacterial Plasmids and Levels of Selection. The Quarterly
Review of Biology. 1990 Mar;65(1):3–22. doi:10.1086/416582

10. Scanlan PD, Hall AR, Blackshields G, Friman VP, Davis MR, Goldberg JB, et al.
Coevolution with bacteriophages drives genome-wide host evolution and constrains the
acquisition of abiotic-beneficial mutations. Molecular Biology and Evolution.
2015;32(6):1425–35. doi:10.1093/molbev/msv032 PubMed PMID: 25681383.

11. Hall JPJ, Wood AJ, Harrison E, Brockhurst MA. Source-sink plasmid transfer dynamics
maintain gene mobility in soil bacterial communities. Proceedings of the National
Academy of Sciences. 2016;113(29):8260–5. doi:10.1073/pnas.1600974113 PubMed
PMID: 27385827.

12. Lam MMC, Wyres KL, Wick RR, Judd LM, Fostervold A, Holt KE, et al. Convergence of
virulence and MDR in a single plasmid vector in MDR Klebsiella pneumoniae ST15.
Journal of Antimicrobial Chemotherapy. 2019 May 1;74(5):1218–22.
doi:10.1093/jac/dkz028

13. Wyres KL, Wick RR, Judd LM, Froumine R, Tokolyi A, Gorrie CL, et al. Distinct
evolutionary dynamics of horizontal gene transfer in drug resistant and virulent clones of
Klebsiella pneumoniae. Hughes D, editor. PLoS Genet. 2019 Apr 15;15(4):e1008114.
doi:10.1371/journal.pgen.1008114

14. San Millan A, Heilbron K, MacLean RC. Positive epistasis between co-infecting plasmids
promotes plasmid survival in bacterial populations. The ISME Journal. 2014 Mar
1;8(3):601–12. doi:10.1038/ismej.2013.182

15. Igler C, Huisman JS, Siedentop B, Bonhoeffer S, Lehtinen S. Plasmid co-infection:
linking biological mechanisms to ecological and evolutionary dynamics. Phil Trans R Soc
B. 2022 Jan 17;377(1842):20200478. doi:10.1098/rstb.2020.0478

16. Rainey PB, Bailey MJ, Thompson IP. Phenotypic and genotypic diversity of fluorescent
pseudomonads isolated from field-grown sugar beet. Microbiology. 1994 Sep
1;140(9):2315–31. doi:10.1099/13500872-140-9-2315

17. Bailey MJ, Lilley AK, Thompson IP, Rainey PB, Ellis RJ. Site directed chromosomal
marking of a fluorescent pseudomonad isolated from the phytosphere of sugar beet;
stability and potential for marker gene transfer. Molecular Ecology. 1995 Dec;4(6):755–
64. doi:10.1111/j.1365-294X.1995.tb00276.x

18. Lilley AK, Fry JC, Day MJ, Bailey MJ. In situ transfer of an exogenously isolated plasmid
between *Pseudomonas* spp. in sugar beet rhizosphere. Microbiology. 1994;140(1):27–
33. doi:10.1099/13500872-140-1-27

19. Lilley AK, Bailey MJ, Day MJ, Fry JC. Diversity of mercury resistance plasmids obtained
by exogenous isolation from the bacteria of sugar beet in three successive years. FEMS
Microbiology Ecology. 1996 Aug;20(4):211–27. doi:10.1111/j.1574-6941.1996.tb00320.x

20. Lilley AK, Bailey MJ. The acquisition of indigenous plasmids by a genetically marked
pseudomonad population colonizing the sugar beet phytosphere is related to local
environmental conditions. Appl Environ Microbiol. 1997 Apr;63(4):1577–83.
doi:10.1128/aem.63.4.1577-1583.1997

21. Bagdasarian M, Lurz R, Rückert B, Franklin FCH, Bagdasarian MM, Frey J, et al.
Specific-purpose plasmid cloning vectors II. Broad host range, high copy number, RSF
1010-derived vectors, and a host-vector system for gene cloning in Pseudomonas.
Gene. 1981 Dec;16(1–3):237–47. doi:10.1016/0378-1119(81)90080-9

22. Blattner FR, Plunkett G, Bloch CA, Perna NT, Burland V, Riley M, et al. The Complete
Genome Sequence of *Escherichia coli* K-12. Science. 1997 Sep 5;277(5331):1453–62.
doi:10.1126/science.277.5331.1453

23. Hall JPJ, Harrison E, Lilley AK, Paterson S, Spiers AJ, Brockhurst MA. Environmentally
co-occurring mercury resistance plasmids are genetically and phenotypically diverse and
confer variable context-dependent fitness effects. Environmental Microbiology.
2015;17(12):5008–22. doi:10.1111/1462-2920.12901 PubMed PMID: 25969927.

24. Sengeløv G, Kristensen KJ, Sørensen AH, Kroer N, Sørensen SJ. Effect of genomic
location on horizontal transfer of a recombinant gene cassette between *Pseudomonas*
strains in the rhizosphere and spermosphere of barley seedlings. Current Microbiology.
2001;42(3):160–7. doi:10.1007/s002840010197 PubMed PMID: 11270648.

25. Bahl MI, Hansen LH, Goesmann A, Sørensen SJ. The multiple antibiotic resistance IncP-
1 plasmid pKJK5 isolated from a soil environment is phylogenetically divergent from
members of the previously established α, β and δ sub-groups. Plasmid. 2007
Jul;58(1):31–43. doi:10.1016/j.plasmid.2006.11.007

26. Klümper U, Riber L, Dechesne A, Sannazzarro A, Hansen LH, Sørensen SJ, et al. Broad
host range plasmids can invade an unexpectedly diverse fraction of a soil bacterial
community. ISME Journal. 2015;9:934–45. doi:10.1038/ismej.2014.191 PubMed PMID:
25333461.

27. Sünderhauf D, Klümper U, Pursey E, Westra ER, Gaze WH, Van Houte S. Removal of
AMR plasmids using a mobile, broad host-range CRISPR-Cas9 delivery tool.
Microbiology. 2023 May 25;169(5). doi:10.1099/mic.0.001334

28. Wright RCT, Wood AJ, Bottery MJ, Muddiman KJ, Paterson S, Harrison E, et al. A
chromosomal mutation is superior to a plasmid-encoded mutation for plasmid fitness
cost compensation. De Visser JAGM, editor. PLoS Biol. 2024 Dec 2;22(12):e3002926.
doi:10.1371/journal.pbio.3002926

29. Leonard S, Perreau J. Barrick Lab [Internet]. 2024 [cited 2025 Mar 21]. General
Conjugation Protocol. Available from:
https://barricklab.org/twiki/bin/view/Lab/ProtocolsConjugation

30. Carrilero L, Kottara A, Guymer D, Harrison E, Hall JPJ, Brockhurst MA. Positive
selection inhibits plasmid coexistence in bacterial genomes. mBio. 2021;12(3).
doi:10.1128/mBio.00558-21 PubMed PMID: 33975933.

31. Carrilero L, Dunn SJ, Moran RA, McNally A, Brockhurst MA. Evolutionary Responses to
Acquiring a Multidrug Resistance Plasmid Are Dominated by Metabolic Functions across
Diverse Escherichia coli Lineages. Rodríguez-Verdugo A, editor. mSystems. 2023 Feb
23;8(1):e00713-22. doi:10.1128/msystems.00713-22

32. Kottara A, Carrilero L, Harrison E, Hall JPJ, Brockhurst MA. The dilution effect limits
plasmid horizontal transmission in multispecies bacterial communities. Microbiology.
2021 Sep;167(9):1–7. doi:10.1099/mic.0.001086 PubMed PMID: 34494951.

33. R Core Team. R: A language and environment for statistical computing [Internet]. Vienna,
Austria: R Foundation for Statistical Computing; 2024. Available from: https://www.r-
project.org/

34. RStudio Team. RStudio: Integrated Development for R. Boston, MA: RStudio, PBC;
2020.

35. Wickham H, Vaughan D, Girlich M. tidyr: Tidy Messy Data [Internet]. 2024. Available
from: https://cran.r-project.org/package=tidyr

36. Wickham H, François R, Henry L, Müller K, Davis V. dplyr: A Grammar of Data
Manipulation [Internet]. 2023. Available from: https://cran.r-project.org/package=dplyr

37. Blazanin M. gcplyr: an R package for microbial growth curve data analysis. BMC
Bioinformatics. 2024 Jul 9;25(1):232. doi:10.1186/s12859-024-05817-3

38. Wickham H. ggplot2: Elegant Graphics for Data Analysis [Internet]. Springer-Verlag New
York; 2016. Available from: https://ggplot2.tidyverse.org

39. Garnier S, Ross N, Rudis R, Camargo AP, Sciaini M, Scherer C. viridis(Lite) - Colorblind-
Friendly Color Maps for R. 2024.

40. Zhang QG, Ellis RJ, Godfray HCJ. THE EFFECT OF A COMPETITOR ON A MODEL
ADAPTIVE RADIATION. Evolution. 2012 Jun;66(6):1985–90. doi:10.1111/j.1558-
5646.2011.01559.x

41. Hall JPJ, Harrison E, Pärnänen K, Virta M, Brockhurst MA. The Impact of Mercury
Selection and Conjugative Genetic Elements on Community Structure and Resistance
Gene Transfer. Frontiers in Microbiology. 2020;11(August):1–14.
doi:10.3389/fmicb.2020.01846

42. Boyd ES, Barkay T. The Mercury Resistance Operon: From an Origin in a Geothermal
Environment to an Efficient Detoxification Machine. Front Microbio. 2012;3.
doi:10.3389/fmicb.2012.00349

43. Hall JPJ, Harrison E, Brockhurst MA. Competitive species interactions constrain abiotic
adaptation in a bacterial soil community. Evolution Letters. 2018;2(6):580–9.
doi:10.1002/evl3.83

44. Goldford JE, Lu N, Bajić D, Estrela S, Tikhonov M, Sanchez-Gorostiaga A, et al.
Emergent simplicity in microbial community assembly. Science. 2018;361(6401):469–74.
doi:10.1126/science.aat1168 PubMed PMID: 30072533.

45. Kottara A, Hall JPJ, Harrison E, Brockhurst MA. Variable plasmid fitness effects and
mobile genetic element dynamics across Pseudomonas species. FEMS Microbiology
Ecology. 2018;94(1):1–7. doi:10.1093/femsec/fix172 PubMed PMID: 29228229.

46. Alonso-del Valle A, León-Sampedro R, Rodríguez-Beltrán J, DelaFuente J, Hernández-
García M, Ruiz-Garbajosa P, et al. Variability of plasmid fitness effects contributes to
plasmid persistence in bacterial communities. Nat Commun. 2021 May 11;12(1):2653.
doi:10.1038/s41467-021-22849-y

47. Sünderhauf D, Klümper U, Gaze WH, Westra ER, Van Houte S. Interspecific competition
can drive plasmid loss from a focal species in a microbial community. The ISME Journal.
2023 Oct 1;17(10):1765–73. doi:10.1038/s41396-023-01487-w

48. Di Venanzio G, Flores-Mireles AL, Calix JJ, Haurat MF, Scott NE, Palmer LD, et al.
Urinary tract colonization is enhanced by a plasmid that regulates uropathogenic
Acinetobacter baumannii chromosomal genes. Nat Commun. 2019 Jun 24;10(1):2763.
doi:10.1038/s41467-019-10706-y

49. Billane K, Harrison E, Cameron D, Brockhurst MA. Why do plasmids manipulate the
expression of bacterial phenotypes? Philosophical Transactions of the Royal Society B:
Biological Sciences. 2022;377(1842). doi:10.1098/rstb.2020.0461 PubMed PMID:
34839708.

50. Coyte KZ, Stevenson C, Knight CG, Harrison E, Hall JPJ, Brockhurst MA. Horizontal
gene transfer and ecological interactions jointly control microbiome stability. PLoS
Biology. 2022;20(11):e3001847. doi:10.1371/journal.pbio.3001847 PubMed PMID:
36350849.

51. Zhu S, Hong J, Wang T. Horizontal gene transfer is predicted to overcome the diversity
limit of competing microbial species. Nat Commun. 2024 Jan 27;15(1):800.
doi:10.1038/s41467-024-45154-w

52. Barkay T, Kritee K, Boyd E, Geesey G. A thermophilic bacterial origin and subsequent
constraints by redox, light and salinity on the evolution of the microbial mercuric
reductase. Environmental Microbiology. 2010 Nov;12(11):2904–17. doi:10.1111/j.1462-
2920.2010.02260.x

613