Inter-individual variability in immune response to AAV ocular gene delivery impedes immunomonitoring

Duohao REN[1,2,3], Gaelle CHAUVEAU[1,2,3], Julie VENDOMELE[1,2], Emilie CABON[1,2,3], Audrey PINEIRO[1,2], Catherine VIGNAL-CLERMONT[4,5], Hanadi SALIBA[1,2], Giuseppe RONZITTI[1,2], Anne GALY[1,2], Deniz DALKARA[3], Juliette PULMAN[3], Divya AIL[1,3], Sylvain FISSON[1,2,3,6,*]

[1]Université Paris-Saclay, Univ Evry, Inserm, Genethon, Integrare research unit UMR_S951, F-91000 Evry, France
[2]Genethon, F-91000 Evry, France
[3]Sorbonne Université, CNRS, Inserm, Institut de la Vision, F-75012 Paris, France
[4]Department of Neuro Ophthalmology and Emergencies, Rothschild Foundation Hospital, Paris, France
[5]Centre Hospitalier National d'Ophtalmologie des Quinze Vingts, Paris, France
[6]Lead contact
*Correspondence: sylvain.fisson@univ-evry.fr

ABSTRACT

Adeno-associated viruses (AAVs) have been used in gene therapy, especially for inherited retinal diseases. Despite their effectiveness in gene transduction, immune responses to the AAV capsid and transgene products have been reported, which can compromise both the efficacy and safety of AAV-mediated therapies. The eye is regarded as an immune-privileged organ where immune activity is constitutively suppressed. Here, we highlighted that immunomonitoring in an ocular gene therapy clinical trial and a non-human primate experiment revealed variable immune responses, whatever the site of injection, the cassette, the dose. We further explored factors contributing to this variability, investigating the correlation among immune parameters in a controlled experimental setting. In a syngeneic murine model after an intraocular injection of AAV, our results highlight an inter-individual variability of immune parameters, emphasizing the importance of considering inherent variability among individuals while designing personalized therapies.

INTRODUCTION

Adeno-associated virus (AAV) vectors have become one of the most promising tools in gene therapy, largely due to their ability to deliver genetic material efficiently and safely into a variety of cell types. Unlike other viral vectors, AAV is non-pathogenic and exhibits minimal toxicity, making it suitable for treating genetic disorders such as hemophilia, muscular dystrophy, and retinal diseases(Sahel & Dalkara, 2019; Jauze *et al*, 2024; Yamaguti-Hayakawa & Ozelo, 2022; Saad *et al*, 2024). Since the first attempt of gene transfer in the retina with AAV in 1996 proving the efficiency of transduction(Ali *et al*, 1996), preclinical research in the application of AAV-mediated gene therapy for retinal diseases is on the rise. In 2007, the first clinical trial using AAV to treat Leber congenital amaurosis was initiated(Hauswirth *et al*, 2008) which eventually led to the first FDA-approved AAV-based ocular gene therapy, Luxturna (voretigene neparvovec), approved in 2017 for RPE65 mutation-associated retinal dystrophy(LUXTURNA® (voretigene neparvovec-rzyl) - Inherited Retinal Disease). And by the end of 2023, 94 clinical trials have been conducted or are ongoing targeting 12 retinal diseases with 6 AAV serotypes (clinicaltrials.gov, Key word: AAV, retina). These outputs have provided convincing proof for the application of AAV-based gene therapies to treat retinal genetic diseases. However, an often overlooked or under-reported aspect of AAV-based therapies, especially in the ocular field, is the immune responses induced by the AAV capsid and transgene product. The therapeutic success of AAV-based gene therapy is significantly influenced by the host's immune response, which can on one hand limit the effectiveness of the treatment(Emami *et al*, 2023), and on the other cause adverse secondary effects such as inflammation.

Although the eye is considered as an immune-privileged organ(Streilein, 1999), safety of ocular gene therapy mediated by AAV is not guaranteed. Indeed, studies have shown that microglial cells can be activated after subretinal injection of AAVs in murine models(Khabou *et al*, 2018). Systemic humoral and cellular immune responses can also be elicited by AAV and transgene products that are delivered by subretinal injections(Ail *et al*, 2022; Vendomèle *et al*, 2024). A study evaluating the immune responses to AAV delivery in the retina of non-human primates (NHP) reported elevated levels of anti-AAV antibodies

in the serum (systemic immune response) as well as ocular inflammation (local immune response)(Ail *et al*, 2022; Hakim *et al*, 2021). Another study, evaluating the cellular immune response by immunostaining in NHPs after AAV subretinal injections in the retina, reported CD8[+] T-cell infiltration(Reichel *et al*, 2017).

Immune responses have not only been observed and reported in animal models used in gene therapy research, but also in clinical trials. Ocular inflammation was reported in patients undergoing ocular AAV gene therapy, and both anti-capsid humoral and cellular immune responses were reported, indicating the potential strong side effects of the AAV gene therapy(Bainbridge *et al*, 2008, 2015). Often, patients in these trials are provided with immunosuppression regimens that aim at managing the immune response-related side effects(Newman *et al*, 2021; Russell *et al*, 2017) Despite this, some patients developed ocular inflammation and systemic immune responses(Duke *et al*, 2022; Cehajic-Kapetanovic *et al*, 2020). An intriguing point is that non-consistent immune responses were observed whatever the condition in clinical trial. For example, in clinical trial NCT00749957, 3 out of 12 patients developed ocular inflammation and 5 out of 12 patients developed anti-capsid antibodies while no patients developed anti-capsid T cell response(Weleber *et al*, 2016). This has been supposedly attributed to differences in disease stage, treatment prior to gene therapy, genetics (Tong & Fairfax, 2020; Klein & Flanagan, 2016), lifestyle and environment(Huang *et al*, 2023) which are confounding factors that can influence the correlations among the immunomonitoring parameters in the patients and the efficiency of the gene therapy. Thus, we would like to test that in controlled animal models. It's not necessarily pertinent to use the same vector in the human and animal as the vector used in the human may not be the functional in the animals and the injection may not be the same either. What is important is to know if there is still an inter-individual variability despite vectors.

In the present study, we first highlighted the inter-individual differences in the anti-AAV immune responses in an ocular gene therapy clinical trial and in an NHP experiment. Next, we conducted analyses of local and systemic immune responses induced by AAV injection, maintaining consistent biological and experimental parameters to identify potential correlations among immune parameters. To that end, we administered AAV

vectors through subretinal delivery in syngeneic mice and evaluated various immune parameters, including antibody production, T cell response, local inflammation, and cytokine secretion. Despite the controlled conditions, we observed significant inter-individual variability in immune responses, with only limited correlations identified among the assessed parameters.

RESULTS

Inter-individual variability of immune response is observed in the human clinical trial and NHP model after intraocular AAV gene transfer

Immunomonitoring after intraocular AAV2 gene transfer showed immune responses in both a human clinical trial(Bouquet *et al*, 2019) and NHPs(Ail *et al*, 2022). Since different strategies yielded distinct outcomes, we investigated the potential relations between the different immunomonitoring data from a human clinical trial described previously (NCT02064569)(Bouquet *et al*, 2019). In this clinical trial, 15 patients diagnosed with ND4 Leber Hereditary Optic Neuropathy (LHON), were distributed into four cohorts and treated with an intravitreal injection of a recombinant AAV2 vector carrying the ND4 gene at 4 different doses. A composite global ocular inflammation score (OIS) was determined using 4 separate grades according to Standardization of Uveitis Nomenclature (SUN) classification (Figure 1A). In addition, total antibody (TAb) and neutralizing antibody (NAb) levels against the capsid were also measured by ELISA and NAb assays respectively in patients. TAb and NAb levels were generally increased in patients' post-injection serum samples, with no clear dose effect (Figures 1B, 1B'). Immune profiles were generated for each patient, incorporating fold-change of TAb and Nab against AAV2, maximal OIS and anti-capsid cellular immune responses measured by IFNγ ELISpot in peripheral blood mononuclear cells (PBMCs) isolated from patients. The immunomonitoring revealed distinct patterns in patients who received the same dose, exhibiting different levels of ocular inflammation scores (OIS), cellular and humoral responses regardless of the AAV dose received (Figures 1C-F, Supplementary table 1).

In order to further investigate immune responses under more controlled environmental conditions, immunomonitoring data from a study on NHPs were analyzed(Ail *et al*, 2022). All the 8 NHPs received an intravitreal injection of 5×10^{11} vector genomes (vg) of AAV2.7m8 encoding ChrimsonR in both eyes (Figure 2A). Ocular inflammation was assessed by slit lamp 1-month post-injection and scores were determined by SUN classification on anterior chamber cells (ACC), anterior chamber flare (ACF), vitreous haze (VH), and vitreous cells (VC). ELISA and NAb assays were used to measure capsid-specific

TAb and NAb levels 2 to 3 months post-injection (PI). Both TAb and NAb against capsid in NHP sera increased after AAV administration (Figures 2B, 2B'). Normalized immune profiles of NHPs highlighted varying contributions of humoral immunity and inflammation by slit lamp, including ACC, ACF, VH, and VC to the overall response (Figure 2C, Supplementary table 2). Both species exhibited variability in immunomonitoring outputs following intraocular AAV injection.

Systemic adaptive immune responses both against capsid and transgene product are induced after subretinal injections of AAV8-GFP-HY in mice

Due to the diversity in the disease stage, genome and environmental factors in humans and NHPs, syngeneic murine models seem to be pertinent to explore the consistency of the immune responses following AAV-mediated AAV gene transfer. Previous studies have demonstrated that AAV intraocular injections can induce both humoral and cellular immune responses against capsids and transgene products that can be detected systemically in blood or in lymphoid organs in murine models(Li *et al*, 2024; Ertl, 2021; Calcedo *et al*, 2013; Bucher *et al*, 2021); however, these studies typically focused on limited immune parameters like antibody production and local inflammation. To comprehensively assess the humoral and cellular immune responses against both capsids and transgene products, a male peptide named HY which was known to induce systemic T cell responses(Vendomèle *et al*, 2018, 2024) in female mice was packaged into AAV8 capsid along with GFP. The AAV vectors ($5x10^{10}$ vg/eye) were administered into the eye of female C57Bl/6 mice via subretinal injection (Figure 3A). Sera and spleens were collected 21 days post-injection as previously described(Vendomèle *et al*, 2018, 2024) to analyze the cellular and humoral immune response against the capsid and transgene product (Figure 3A). IFNγ ELISpot revealed a significant increase in activated T cells against AAV (p-value = 0.0001) and transgene product (p-value < 0.0001) in AAV-injected mice compared to PBS-injected control mice (Figures 3B, 3C and Figure S1A). HY peptides (pHY) are composed of DBY, activating CD4[+] T cells and UTY, activating CD8[+] T cells(Carpentier *et al*, 2015). ELISpot assay showed an IFNγ secretion by CD4[+] and CD8[+] T cell specific to transgene product (Figures S1B, S1C). ELISA measurements showed that anti-AAV8 and anti-GFP antibodies significantly increased (p-value < 0.0001) after AAV subretinal injection compared to PBS-injected control mice (Figures 3D, 3E). The systemic cytokine profile contributing to the inflammation and immune response in mice was evaluated following AAV subretinal injection. Spleen cells were isolated from mice that received either PBS or AAV. These spleen cells were then stimulated *in vitro* with pHY or AAV. A cytometric bead array (CBA) was used to measure cytokines in the culture supernatant to determine the cell polarizations: Th1/Tc1 (IL-2, IFNγ, TNFα,

GM-CSF), Th2/Tc2 (IL-4, IL-10, IL-13), Th17/Tc17 (IL-17), and those involved in inflammation and migration (IL-1β, IL-6, RANTES, MCP-1) in the culture supernatant to determine the cell polarizations. Radar charts were generated to visualize the proportional production of these cytokines. Spleen cells from PBS-injected control mice mainly did not secrete cytokines (Figures 4A, 4A'). In contrast, spleen cells from AAV-injected mice produced production of multiple cytokines (Figures 4B, 4B'). Among cytokines involving inflammation and migration, only RANTES (p-value = 0.01) showed a significant increase (Figure S2A). The Th1/Tc1 cytokines, IL-2 (p-value = 0.0046), IFNγ (p-value = 0.0001), TNFα (p-value = 0.0388), were significantly upregulated (Figure S2B), in response to *in vitro* AAV stimulation. Other cytokines remained unchanged (Figures S2A, S2B, S2C, and S2D). Upon *in vitro* stimulation with pHY, cytokines such as RANTES (p-value < 0.0001), IFNγ (p-value = 0.0001), TNFα (p-value = 0.0007), and IL-10 (p-value = 0.0024) were significantly upregulated (Figure S3).

Local transgene expression in the retina is correlated with cytotoxicity against transgene-expressing cells post-injection of AAV-GFP-HY

Further, the level of transgene expression was measured to assess the efficacy of AAV transduction and its relationship with cytotoxic effects using Droplet Digital PCR (ddPCR). In AAV-injected mice, significant expressions of the HY (p-value = 0.0014) and GFP (p-value = 0.0002) transgenes were observed 21 days after injection, and a correlation between the expression levels of both transgenes was identified ($R^2 = 0.818$) (Figures 5A, 5B). To evaluate the clearance ability of specific antigen-expressing cells in these mice, an *in vivo* cytotoxicity assay was performed using male spleen cells expressing the HY antigen as target cells which means the more target cells survive, the less cytotoxicity there is. We found a significant reduction (p-value = 0.0004) in the number of male target cells in AAV-injected mice, indicating the development of HY-specific cytotoxicity (Figure 5C). Direct correlations between target cell survival in the blood and retinal transgene expression ($R^2 = 0.7702$ for GFP; $R^2 = 0.7935$ for HY) were demonstrated suggesting an inverse correlation between cytotoxicity effect and transgene expression in the mice (Figures 5D, 5E).

Local inflammation and correlated expressions between MHC II molecules and CYBB are found in the retina post-injection of AAV-GFP-HY

The direct correlation between target cell survival and retinal transgene expression suggested a potential antigen presenting cell (APC) contribution to present transgene product peptides, associated with local inflammation. It was assessed by evaluating the transcript expression of MHC II molecules(Lipski *et al*, 2017) (H2-Eb1 and H2-Ab1) and the type 1 macrophage marker CYBB(Frazão *et al*, 2015) in the retina, using ddPCR 21 days after the injection. We observed a significant increase in the expression of both MHC II molecules (p-value = 0.0002) 21 days after the subretinal injection (Figure 6A). Since H2-Eb1 and H2-Ab1 are co-dominant molecules, their correlated expressions were confirmed (R^2 = 0.8686) (Figure 6B). Interestingly, the expression of CYBB increased significantly in AAV-injected mice (p-value = 0.0002) (Figure 6C), and correlations were found between CYBB and MHC II molecules expression (R^2 = 0.9659 for H2-Eb1; R^2 = 0.8165 for H2-Ab1), suggesting a link between antigen presentation and macrophage activation (Figures 6D, 6E).

Diversity in immune responses is noticed in syngeneic mice 21d post-injection of AAV-GFP-HY

After analysis of all immune parameters typically collected for immunomonitoring in clinical trials, variability in the immune parameters was observed even in this syngeneic model. To figure out the main factors driving this variability, principal component analysis (PCA) was performed, and angle sectors were then used to show the immune profiles and cytokine profiles of each mouse. In angle sectors, all analyzed transgene expression, immune parameters and significantly regulated cytokines were normalized to their corresponding highest values to provide a more direct visualization. The immune profile included transgene expression (HY-Tg and GFP-Tg), local inflammation (H2-Eb1, H2-Ab1 and CYBB), humoral immune response (AAV-Ab and GFP-Ab) and cellular immune response (%male, ELISpot-HY, ELISpot-DBY, ELISpot-UTY, ELISpot-AAV-PGK-Luc2 and ELISpot-AAV-PGK-GFP-HY). The cytokine profile was composed of cytokines significantly affected by AAV *in vitro* stimulation (IL-2, IFNγ, TNFα and RANTES) or HY *in vitro* stimulation (IFNγ, TNFα, IL-10 and RANTES). No similar immune profiles were found in AAV-injected mice suggesting that the immune responses against AAV and its transgene product may vary among mice (Figures 7A, 7B, Supplementary table 3). The Principal Component Analysis (PCA) results confirmed that AAV-injected mice could be clearly separated from PBS-injected mice as expected and that the AAV-injected group exhibited a broad distribution of the individuals which was consistent with immune and cytokine profiles, highlighting the variability of immune parameters after AAV subretinal injection (Figure 7C).

Absence of correlation between immune parameters in syngeneic mice 21 days following AAV-GFP-HY delivery

To explore relationships among immune parameters, a correlation analysis was conducted. With a correlation matrix, R^2 values for each parameter pair was displayed and was further visualized using a network analysis to illustrate the connections among these parameters. Distinct correlations emerged between certain parameters; for instance, anti-transgene product antibody levels showed a clear correlation with transgene expression, which in turn was linked inversely to male cell death, indicating the cytotoxic effect. However, most parameters displayed no significant correlations, particularly across the four primary domains: local transgene expression, local inflammation, systemic immune response, and cytokine profile (Figures 8A, 8B). The absence of correlation between immune markers suggests that subretinal AAV administration leads to individualized immune responses, with significant variation between subjects. This indicates that each subject may mount a distinct immunological reaction to the treatment.

DISCUSSION

Gene therapy offers a promising approach for treating genetic diseases, with AAV vectors employed to deliver the therapy (Niemeyer *et al*, 2009; Bainbridge *et al*, 2003; Miyoshi *et al*, 1997). Nevertheless, immune responses triggered by AAV capsids and transgene products remain a concern, as they are observed both peripherally and locally in preclinical and clinical studies, and they are generally considered as factors that can negatively affect the effectiveness of gene therapy(Ertl, 2022; Shen *et al*, 2022; Ren *et al*, 2022). We highlight here that immunomonitoring results in clinics vary among patients. The immune response variations observed in patients is usually attributed to the stage of disease, treatment prior to gene therapy, genetics, environment and lifestyle. However, in our study despite the use of a syngeneic murine model treated with AAV vectors in highly controlled conditions (same sex, same age, kept in the same environment, subjected to the same treatment), a considerable inter-individual variability was still observed, similar to the high inter-individual variability noticed both in human clinical and NHP data.

Inter-individual variability of immune response after intraocular AAV injections was evident in humans and NHPs. None of the individuals, whether human or NHP, who received the same dose and vector injection exhibited identical immune responses. This observation confirms that there is a widespread variability in heterogeneous individuals which has not been well understood and emphasizes the critical need to identify the factors contributing to this inter-individual variability. In order to confirm if the inter-individual variability persists despite the route of administration, transgene and serotypes, another AAV vector was applied in murine experiments. Thus, systemic immune response induced by AAV subretinal injection was characterized in the syngeneic murine model. HY-GFP were packaged into AAV which was previously shown to induce systemic T cell immune responses(Vendomèle *et al*, 2018, 2024) against transgene products. All the mice developed adaptive systemic immune responses and surprisingly innate immune responses persisted 3 weeks after the injection. In previous studies aiming to explore the factors impacting the AAV-induced immune responses, a dose-dependency was repetitively identified(Ronzitti *et al*, 2020; Hamilton & Wright, 2021) which can also be influenced by serotype (Gehrke *et al*, 2022; Bentler *et al*, 2023; Chan *et al*,

2021), administration route(Ge *et al*, 2001), transgene(Boisgerault *et al*, 2013; Xiong *et al*, 2019; Khabou *et al*, 2018; Pfromm *et al*, 2022) and the sex of the recipient(Clare *et al*, 2025). In the context of this study, optimization of vector dose and transgenes could help reduce the risk of inducing a potential immune response. Furthermore, developing novel AAV vectors with lower immunogenicity can be another alternative option to evade part of the immune responses directed against the capsid. In addition to directed evolution, capsid shuffling and rational design, modeling in silico and machine learning are increasingly being used for AAV capsid development. In silico strategies can predict which mutations are likely to enhance AAV functionality and narrow down potential candidates to test in experimental setting(Wec *et al*, 2021).

Correlation data analyses revealed a novel finding: the lack of significant relationships between immune parameters, including local transgene expression, local inflammation, and systemic immune responses. This independence of immune factors has not been previously documented. Among the studied parameters, transgene expression level appears to be inversely correlated with the cytotoxicity against transgene expressing cells, which is worthy of further exploration. In contrast, the absence of correlation found with the commonly used clinical immunomonitoring methods, such as ELISA and ELISpot(Bainbridge *et al*, 2015; VIGNAL-CLERMONT *et al*, 2023; Britten *et al*, 2013), which investigate peripheral adaptive immune responses, suggests that they may not fully capture the complexities of immune reactions in ocular gene therapy patients. Thus, the use of these immunomonitoring techniques may not reflect the actual immune status and cannot be used to determine the application of immunosuppression regimens in patients. Analysis of systemic cytokine profiles and putative inflammatory biomarkers(Calcedo *et al*, 2018) proved to be poor predictors of immune responses following AAV gene delivery to the eye. However, a more extensive assessment of local cytokine expression patterns could provide additional insights. Besides, patients with retinal diseases express local inflammation cytokines in the retina that have the potential to leak into the periphery which further complicates the sensitivity of cytokines as biomarkers(Tao *et al*, 2024). Therefore, novel immunomonitoring strategies need to be explored to provide a complete and exhaustive

view of immunological status which can further aid in the standardization of the application of immunosuppression strategies in clinics.

Inter-individual variability in immune responses has been observed in human clinical trials as well as in NHPs receiving ocular gene therapy. This is expected since even in human monozygotic twins, the immune responses can differ due to T and B cell repertoire(HAVERKORN *et al*, 1975; Pogorelyy *et al*, 2018). However, with our study we demonstrated that even in a syngeneic murine model where most of the factors were kept the same, high inter-individual variability in immune responses is observed. Even though factors like the surgery and operation of the experiments cannot be exactly the same in all the mice, this kind of artificial variation also exists in human and other animal models(Puranen *et al*, 2023). Similar inter-individual variability of immune response has been reported in vaccine and anti-tumor research within syngeneic murine models(Audebert *et al*, 2021; Rotnemer-Golinkin & Ilan, 2022; Leete *et al*, 2022; Sivan *et al*, 2015). The factors and mechanisms impacting the inter-individual variability are still not well understood. One study applied vaccinia virus to activate the T cell response in C57BL/6 mice to follow CD8[+] T cell dynamics and inter-individual variability was observed in priming efficiency, effector expansion and memory cell generation(Audebert *et al*, 2021). Additionally, another study reported a varied immune response in C57BL/6 mice administered with concanavalin A (ConA), which can induce immune-mediated hepatitis, and measured kinetics of alanine aminotransferase (ALT) levels (a marker of liver damage), emphasizing immune responses vary significantly in timing and magnitude, even among genetically identical subjects(Rotnemer-Golinkin & Ilan, 2022). Besides, a study tracked CD8[+] cell in CT26 syngeneic murine tumor model to assess anti-PD-1 therapy response and high variability in anti-tumor responses was observed, indicating contribution of the rate of CD8[+] T cell activation to the inter-individual variability(Leete *et al*, 2022). Another study showed that different intestinal microbiota could impact the immune response in syngeneic mice. Mice that were raised in two different facilities showed differences in CD8[+] cell priming and accumulation during anti-tumor therapy(Sivan *et al*, 2015). In our study however, all the mice were raised in the same facility, under the same diet and housing conditions, and are expected

to have similar intestinal microbiota. Nonetheless, this can be a factor influencing the variability in human and other animal models and merits consideration. Such studies reinforce the idea that inter-individual immune variability is a common phenomenon in immunological contexts beyond gene therapy and further exploration is needed to elucidate the underlying mechanisms.

Animal models provide crucial inputs for gene therapy development and fundamental immunology research. In our study, despite using a syngeneic murine model to eliminate most internal and external variability factors, an inter-individual variability in immune responses was observed. Alternatively, research on cell and organoid levels, which are used in gene therapy transduction (Ramamurthy *et al*, 2022), may assist in understanding more about the factors impacting the variability by focusing for example on innate immune reactivity. Regardless of the systemic collaboration of immune systems, cell lines and organoid models can still provide the insight of the immune sensitivity to specific antigens where attempts to use cell culture to predict immunogenicity have been initiated(Fischer *et al*, 2011; Ozawa *et al*, 2021).

Our findings underscore the need for individualized patient care strategies, highlighting the value of a personalized medicine approach(Zheng *et al*, 2015; Ong *et al*, 2013). The understanding and surveillance of individual immune response variations could inform personalized decisions about dosing, treatment selection, duration, and immunosuppressive regimens. The contribution of artificial intelligence (AI) in biology has been expanded widely during the last decade, which allows meta-data analysis and predictions. Multiple studies have used AI in cancer immunotherapy and different AI models have been trained to predict not only various immune signatures but also direct immunotherapy responses(Yang *et al*, 2022). Despite the existence of predictive tools of the immune response in specific cells or signaling pathways(Penhaskashi *et al*, 2023; Chandler *et al*, 2022), those have not been adapted to the context of gene therapy, and challenges in predicting immune responses in this field still largely persist. Current limitations in our understanding of immune system complexity may prevent AI from generating accurate predictions.

In humans and complex animal models like NHPs, inter-individual differences in immune responses are expected and attempts are made to minimize their impact on safety and efficacy (immunosuppression strategies) or to resolve symptoms (anti-inflammation strategies), with little to no information on the underlying mechanisms. Our goal with the present study was to identify the most pertinent parameter or a combination of parameters that can be reliably used to predict, follow-up and manage immune responses post-therapy. However, even in syngeneic mice receiving the same treatments we observed high variability in immune responses. Few correlations were observed among local transgene expression, local inflammation, and systemic immune responses revealing the limitation of the current immunomonitoring strategies in ocular gene therapy clinical trials. Our study reinforces two crucial points – the first one being that the current immunomonitoring strategies can hardly be used to infer on the ongoing immune response and adapt the immunosuppressive regimens. Efforts should be focused on the understanding of the underlying mechanisms leading to the individual differences observed in the immune response. Second, in the therapeutic context patients may benefit from 'personalized therapies' that are designed taking into consideration their own unique immune profiles.

Limitations of the study

In this study, a limited range of immune parameters were evaluated concerning local inflammation, systemic cytokines and systemic anti-transgene product and anti-capsid humoral and cellular immune response. These outputs which correspond to those mostly used in clinical immunomonitoring may not reflect the complex individual immune responses after AAV subretinal injection. Therefore, a more thorough analysis of local inflammation or the systemic immune response using alternative techniques can strengthen the conclusions regarding immune response variability while also identifying new potential biomarkers and immunomonitoring strategies. Besides, all the experiments were performed based on a controlled syngeneic murine model with the same age and gender. Our results should be confirmed in other syngeneic murine models to exclude the potential impact of the model, and possibly with additional conditions such as the animals' age, more AAV serotypes.

Methods

Reagents and Tools table

REAGENT OR RESOURCE	SOURCE	IDENTIFIER
Antibodies		
IFNγ Monoclonal Antibody	eBiosciences	Cat#14-7318-81; RRID:AB_468475
PE Mouse Anti-Mouse CD45.1	BD Biosciences	Cat#553776; RRID:AB_395044
Anti-AAV8 antibody	Humimmu	Cat#A20016-100ug; RRID:AB_3675841
Anti-GFP antibody	Abcam	Cat#ab1218; RRID:AB_298911
Goat Anti-Mouse IgG, Human ads-HRP	Southern Biotech	Cat#1030-05; RRID:AB_2619742
Bacterial and virus strains		
AAV8-GFP-HY	Généthon	N/A
AAV8-Luc2-HY	Généthon	N/A
Chemicals, peptides, and recombinant proteins		
Ketamine	Virbac	Cat#AMM FR/V/5646687 8/1986
Xylazine	Bayer	Cat#AMM FR/V/8146715 2/1980
Isoflurane	Laboratorios Karizoo	N/A
Phosphate-buffered saline (PBS)	Gibco	Cat#14190-094
NH_4Cl	Sigma	Cat#A9434
EDTA	Sigma	Cat#27285
$KHCO_3$	Sigma	Cat#P-9144
Concanavalin A (Con A)	Sigma	Cat#234567M
β-mercaptoethanol	Sigma	Cat#M6250
$NaHCO_3$	Sigma	Cat#55761-1kg
Na_2CO_3	Sigma	Cat#223530-1kg
Bovine serum albumin (BSA)	VWR	Cat#422361V
TMB reagent	BD Biosciences	Cat#555214
Sulfuric acid	Sigma	Cat#S-1526
The DEAD Box polypeptide 3Y-linked (DBY)	Genepep	NAGFNSNRANSSRSS
UTY peptide	Genepep	WMHHNMDLI
Complete Freund's adjuvant	Sigma	Cat#F5881-10ML
BCIP/NBT	Mabtech	Cat#3650-10
Streptavidin-Alkaline Phosphatase	Roche Diagnostics	Cat#03137031103
GFP	Chromotek	Cat#EGFP-250
Critical commercial assays		
Mouse IL-1β Flex Set	BD Biosciences	Cat#560232
Mouse IL-2 Flex Set	BD Biosciences	Cat#558297

Mouse IL-4 Flex Set	BD Biosciences	Cat#558298
Mouse IL-6 Flex Set	BD Biosciences	Cat#558301
Mouse IL-10 Flex Set	BD Biosciences	Cat#558300
Mouse IL-13 Flex Set	BD Biosciences	Cat#558349
Mouse IL-17A Flex Set	BD Biosciences	Cat#560283
Mouse IFN-γ Flex Set	BD Biosciences	Cat#558296
Mouse GM-CSF Flex Set	BD Biosciences	Cat#558347
Mouse TNF Flex Set	BD Biosciences	Cat#558299
Mouse RANTES Flex Set	BD Biosciences	Cat#558345
Mouse MCP-1 Flex Set	BD Biosciences	Cat#558342
BD Cytometric Bead Array (CBA) Mouse/Rat Soluble Protein Master Buffer Kit	BD Biosciences	Cat#558267
Cell trace violet (CTV) cell proliferation kit	Invitrogen	Cat#C34557
Rneasy kit	Qiagen	Cat#74136
Rnase free Dnase Set	Qiagen	Cat#79254
Verso cDNA Synthesis Kit	ThermoFisher Scientific	Cat#AB1453B
ddPCR Supermix for probes (no dUTP)	BIO-RAD	Cat#1863024
Experimental models: Organisms/strains		
C57BL/6J	Charles River Laboratories	Strain#027
Oligonucleotides		
GAPDH forward primer	Eurofins genomics	CATGGCCTTCCGTGTTCCTA
GAPDH reverse primer	Eurofins genomics	GCGGCACGTCAGATCCA
GAPDH probe (HEX)	Eurofins genomics	CCCCAATGTGTCCGTC
eGFP forward primer	Eurofins genomics	ACGTCTATATCATGGCCGAC
eGFP reverse primer	Eurofins genomics	GTGCTCAGGTAGTGGTTGTC
eGFP probe (FAM)	Eurofins genomics	ACGGCCCCGTGCTGCTGCCC
HY forward primer	Eurofins genomics	CCAATAGCAGCCGAAGTAGT
HY reverse primer	Eurofins genomics	GGTGCATCCAACCTAACTG
HY probe (FAM)	Eurofins genomics	TGGTGGAGGTGGCTATGGAGG
H2-Eb1 forward primer	Eurofins genomics	CTGGTCCGAAATGGAGACTG
H2-Eb1 reverse primer	Eurofins genomics	TGTGCTTTCCACTCGACC
H2-Eb1 probe (FAM)	Eurofins genomics	ACCTGCCAGGTGGAGCATCCCAGCC
H2-Ab1 forward primer	Eurofins genomics	TGTGCAGACACAACTACGAG
H2-Ab1 reverse primer	Eurofins genomics	GACATTGGGCTGTTCAAGC

H2-Ab1 probe (FAM)	Eurofins genomics	ACCCACACCTCCCTGCG GCG
CYBB forward primer	Eurofins genomics	TCGAAAACTCCTTGGGT CAG
CYBB reverse primer	Eurofins genomics	TCTTCGAATCCTTGTCG AGC
CYBB probe (FAM)	Eurofins genomics	CGGGCACCTGCAGCCTG CCTGAATT
Software and algorithms		
AID ELISpot Reader v6.0 software	AID	https://www.aid-diagnostika.com/en
FCAP software	BD Biosciences	Cat#652099
Cytoscape v3.10.2	Cytoscape	https://cytoscape.org/
GraphPad Prism V10.0	GraphPad	https://www.graphpad.com/
CytExpert software	Beckman Coulter	https://www.beckman.fr/flo w-cytometry/research-flow-cytometers/cytoflex/software
QXManager	BIO-RAD	Cat#12018108
Other		
CytoFLEX LX flow cytometer	Beckman Coulter	Cat#C00445
AID ELISpot Reader system ILF05	AID	https://www.aid-diagnostika.com/en
96-well Maxisorp plates	ThermoFisher scientific	Cat#439454
IFN-γ Enzyme-Linked Immunospot plates	Millipore	Cat#MAHAS4510
Automated Droplet Generator	BIO-RAD	Cat#1864101
QX 200 Droplet Reader	BIO-RAD	Cat#1864003
Non-fat milk	Régilait	REGILAIT-Skimmed Powdered Milk

Methods and protocols

Murine model

Wild-type 6- to 8-week-old female C57BL/6J mice (H-2^b) were purchased from Charles River Laboratories (L'Arbresle, France). Animals were anesthetized either by intraperitoneal injection of 120 mg/kg ketamine (Virbac, Carros, France) and 6 mg/kg xylazine (Bayer, Lyon, France) or by inhalation of isoflurane (Baxter, Guyancourt, France). They were euthanized by cervical elongation. All mice were housed, cared for, and handled in accordance with the European Union guidelines and with the approval of the local research ethics committee (CEEA-51 Ethics Committee in Animal Experimentation, Evry, France; authorization number 2015102117539948).

Methods and Protocols

Clinical data and NHP data analysis

Immunomonitoring data from the clinical trial (NCT02064569) that includes 15 patients with LHON carrying the G11778A-ND4 mutation was analyzed. These patients were divided into 4 dose cohorts (9×10^9, 3×10^{10}, 9×10^{10}, and 1.8×10^{11} vg per eye) and each cohort received an intravitreal injection. Ocular inflammation, anti-AAV2 TAb, anti-AAV2 NAb and anti-AAV2 cellular immune response were quantified, and details were described previously(Bouquet *et al*, 2019). In brief, a composite global OIS was determined according to SUN classification. Anti-AAV2 TAb were determined by ELISA from human serum samples prediluted at 1:50. NAb assay was used to measure anti-AAV2 NAb with HEK293 cells cultured with an rAAV2 expressing luciferase under the control of the cytomegalovirus promoter (8×10^7vg per well) and either alone or serial fold dilutions of human serum samples. The half maximal inhibitory concentration was determined using the intercept at 50% of the regression curve and expressed as a dilution factor. Peripheral blood mononuclear cells (PBMCs) were isolated and an IFNγ ELISpot assay was performed to measure cellular immune response against AAV2 antigens. Angle sector diagrams for patients were generated based on the FC of anti-AAV2 TAb and NAb, maximal OIS and anti-AAV2 cellular immune response which had been normalized with the highest value corresponding to each aspect.

Immunomonitoring data from 8 NHPs was analyzed. NHPs received an intravitreal injection of $1x10^9$ vg AAV2.7m8 in both eye. Ocular inflammation, anti-AAV2 TAb and Nab were evaluated and details were described previously(Ail *et al*, 2022). In brief, anti-AAV2 TAb were determined by ELISA from NHP serum samples prediluted at 1:100. NAb assay for AAV2 was performed using HEK293T cells cultured with an rAAV2 expressing luciferase under the control of the cytomegalovirus promoter (at a multiplicity of infection (MOI) of 6,400 per well) and either alone or serial fold dilutions of NHP serum samples. The half maximal inhibitory concentration was determined using the intercept at 50% of the regression curve and expressed as a dilution factor. A Spectralis HRA + OCT system was used to acquire OCT images. SUN classification was applied for grading the "Anterior chamber cells". For grading the "Vitreous cells", the NIH grading system is used. The British Medical Journal (BMJ) grading system is used for grading the

"Posterior Uveitis".

AAV vectors

AAV8-GFP-HY and AAV8-Luc2-HY vectors were produced by Généthon in Evry (France) using the tritransfection technique in 293T cells cultured in roller bottles(Liu *et al*, 2003). Transgenes were under the ubiquitous Phosphoglycerate kinase (PGK) promoter. HY is a male antigen which is immunogenic in female mice. AAV vectors were purified by cesium chloride gradients centrifugation, and vector titters were determined by qPCR. Endotoxin levels were below 6 E.U/mL.

Peptides

The DEAD Box polypeptide 3Y-linked (DBY) and Ubiquitously Transcribed tetratricopeptide repeat gene Y-linked (UTY) peptides, NAGFNSNRANSSRSS and WMHHNMDLI, respectively, were synthesized by Genepep (Montpellier, France) and shown to be more than 95% pure. UTY and DBY are immunodominant peptides of the HY antigen, restricted to MHC-I and MHC-II, respectively.

Injections

Subretinal injections were performed as described previously(Terrada *et al*, 2006, 2008). Briefly, the right eye was protruded under microscopic visualization, and the sclera was perforated with a 27G beveled needle. A blunt 32G needle set on a 10 μL Hamilton syringe was inserted in the hole and the same volume (2 μL) of PBS, or AAV (5E10vg), was injected into the subretinal space via the vitreous. The quality of the injection was verified by checking the detachment of the retina and the absence of reflux outside the eye.

Cell extraction from murine spleen

After euthanasia, cells were extracted from spleen as described previously(Vendomèle *et al*, 2018). Briefly, spleens were removed and crushed with a syringe plunger on a 70-μm filter in 2 mL of RPMI medium. Red cells were lysed by adding ACK buffer (8.29 g/L NH_4Cl, 0.037 g/L EDTA, and 1 g/L $KHCO_3$)

for 1 min. Lysis was stopped by addition of complete RPMI medium (10% FBS, 1% penicillin/streptomycin, 1% glutamine, and 50 µM β-mercaptoethanol). After centrifugation, cells were counted and the concentration was adjusted in complete RPMI medium.

Murine IFNγ ELISpot assay

IFNγ ELISpot assay was performed as described previously(Vendomèle *et al*, 2018). IFN-γ Enzyme-Linked Immunospot plates (MAHAS4510, Millipore, Molsheim, France) were coated with anti-IFNγ antibody (eBiosciences, San Diego, CA, USA) overnight at +4°C. Stimulation media (complete RPMI), AAV (1×10^{11} vg/mL), UTY (2 µg/mL), DBY (2 µg/mL), UTY + DBY (2 µg/mL), or Concanavalin A (Sigma, Lyon, France) (5 µg/mL) were plated and 5×10^5 spleen cells/well were added. After 24 h of culture at +37°C, plates were washed and the secretion of IFNγ was revealed with a biotinylated anti-IFNγ antibody (eBiosciences), Streptavidin-Alcalin Phosphatase (Roche Diagnostics, Mannheim, Germany), and BCIP/NBT (Mabtech, Les Ulis, France). Spots were counted with an AID ELISpot Reader system ILF05 and the AID ELISpot Reader v6.0 software. Results are expressed in index where IFNγ secretion of the positive control was set to 100, based on the positive control for anti-HY immune response.

Cytokine titration by multiplex cytometric bead array

Cytokine titration by multiplex cytometric bead array was performed as described previously(Vendomèle *et al*, 2018). Stimulation media (complete RPMI), AAV (1×10^{11} vg/mL), UTY (2 µg/mL), DBY (2 µg/mL), UTY + DBY (2 µg/mL), or Concanavalin A (Sigma, Lyon, France) (5 µg/mL) were plated and 1×10^6 spleen cells/well were added. After 36 h of culture at +37°C, supernatants from triplicates were pooled and frozen at −80°C until the titration. Cytometric bead arrays were performed with BD Biosciences flex kits (IL-2, IL-4, IL-6, IL-10, IL-13, IL-17A, IFN-γ, GM-CSF, TNFα, and MCP-1). Briefly, capturing bead populations with distinct fluorescence intensities and coated with cytokine-specific capture antibodies were mixed together. Next, 25 µL of the bead mix of beads was distributed and 25 µL of each sample (supernatants) was added. After 1 h of incubation at room temperature, cytokine-specific PE-antibodies were mixed and 25 µL of this mix was added. After 1 h of incubation at room temperature, beads were washed with 1 mL of Wash

buffer and data were acquired with an LSRII flow cytometer (BD Biosciences). FCAP software (BD Biosciences) was used for the analysis.

Radar diagrams represent the percentage of cytokine secretion in the different groups based on the maximum of cytokine secretion and were performed with Excel software. The 100% radar scale fits the maximum value of cytokine secretion, and the black area values correspond to the means of these cytokine secretions

In vivo cell cytotoxicity assay

IVC assay was performed as previously described(Vendomèle *et al*, 2024).

Spleen cells from CD45.1$^+$ CD45.2$^-$ male (expressing HY antigen) and CD45.1$^-$ CD45.2$^+$ female (not expressing HY antigen) C57BL/6 wild-type mice were harvested as described above and stained with cell trace violet (CTV) cell proliferation kit (Molecular Probes) in PBS at different concentrations: 2 μM for male and 20 μM for female cells, according to the protocol of the kit. CTV staining level allows tracking separately male, and female transferred cells. A mixture of $3x10^6$ male cells (CTV^{low}) and $3x10^6$ female cells (CTV^{high}) in 200 μL was injected intravenously in the experimented female C57BL/6 mice (CD45.1$^-$ CD45.2$^+$ and not expressing HY antigen) at day 17 of the protocol. CD45.1$^-$ CD45.2$^+$ CTV^{high} female cells are used as a control of cell survival, as they are not targeted by anti-HY immune responses. Three days after injection, blood was harvested, red blood cells were lysed by adding ACK buffer, washed in PBS 1X, and leukocytes were stained for flow cytometry with an anti-CD45.1-PE antibody to analyze the male cell survival *in vivo* (Pharmingen, BD Biosciences). Data were acquired on a CytoFLEX LX flow cytometer (Beckman Coulter) and analyzed with the CytExpert software (Beckman Coulter).

Enzyme-linked immunosorbent assay (ELISA)

96-well Maxisorp plates (ThermoFisher scientific) were coated with full AAV8 capsid ($5x10^8$ vg/well) or GFP (0.5 μg/well, Chromotek) diluted with coating buffer (0.84% NaHCO$_3$, 0.356%Na$_2$CO$_3$, pH9.5) overnight at +4°C. The wells were emptied and washed with blocking buffer (PBS 1X-6% milk) before

incubating the plate with blocking buffer at room temperature for 2 hours. Serial dilutions of primary antibodies (Anti-AAV8 antibody, Hum Immu; Anti-GFP antibody, Abcam) or serum were prepared during the incubation. Primary antibodies were diluted with dilution buffer (PBS 1X-1% BSA) to ensure a gradient on the plate (100ng, 50ng, 25ng, 12.5ng, 6.25ng, 3.2ng, 1.56ng, 0.781ng, 0.391ng, 0.195ng, 0.0975ng for AAV8 and 25ng, 12.5ng, 6.25ng, 3.2ng, 1.56ng, 0.781ng, 0.391ng, 0.195ng for GFP) to calculate the standard curve for the experiment. Murine sera were diluted to 1/1000 with dilution buffer for AAV ELISA and 1/2500 for GFP ELISA. After the incubation, primary antibodies dilution and serum dilution were added in the wells and the plate were incubated for 1 hour in the incubator at 37°C after 3 times washing with washing buffer (PBS 1X-0.05% Tween 20). 1/4000 dilution of secondary antibodies (Goat Anti-Mouse IgG, Southern Biotech) was prepared with dilution buffer. Secondary antibodies were added into each well after washing 3 times and the plate was incubated for 1 hour in 37°C incubator. The TMB reagent (BD Biosciences) and stop solution (1M sulfuric acid) were placed at room temperature 30 minutes before the end of the previous incubation. When the last incubation ended, the plate was washed 3 times with washing buffer. TMB reagent was added and a blue color appeared. Stop solution was added to stop the reaction after 10 minutes. The plate was read at 450nm to get the optical density of each sample.

RNA extraction from mouse models retinas and reverse transcription

Total RNA was isolated using RNeasy kit, QIAGEN (ref: 74106). The procedure was performed according to the manufacturer's specification. The purification included a DNase treatment using the RNase free DNase Set (Qiagen). The yield and purity of the RNA was measured with NanoDrop 8000 spectrometer. Verso cDNA Synthesis Kit (ThermoFisher Scientific) was applied for reverse transcription following the protocol provided.

Droplet Digital PCR (ddPCR)

Preparation of the primers and probes for the genes of interests. For each gene, a mix of primer and probe was prepared with 18µL of primers forward, 18µL of primers reverse, 5µL of probe (Eurofin genomics,

initial concentration of 100mM) and 59µL of H_2O MilliQ.11µL of ddPCR Supermix for probes (no dUTP) (Biorad), 1µL from the primers/probes mix prepared previously (the ratio of housekeeping gene and target gene was 1:1), 6µL of water, and 2µL of the cDNA sample from mice (diluted with water to obtain quantities of either 5ng or 2.5ng in the wells) were added in one well. The droplets were generated by the Automated Droplet Generator from BIO-RAD. A PCR was proceeded by the C1000 Touch Thermal Cycler from BIO-RAD, with the program as 10min at 95°C, repetition of 30sec at 94°C and 1min at 58.1°C 40 times, then 10min at 98°C to finalize at 12°C infinite hold. The results were obtained by QX 200 Droplet Reader from BIO-RAD with the software QXManager for the analysis. Copy numbers of the genes were obtained. Copy number of gene of interest was normalized with copy number of housekeeping gene and the ratio was multiplied by 100.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analyses were performed with GraphPad Prism V10.0. Mann-Whitney tests, correlation matrix and principal component analysis were performed. p value <0.05: *, <0.01: **, <0.001: ***, <0.0001: ****. Radar diagrams and angle sector diagrams made by Excel software represent the percentage of measured value in the different groups normalized with the maximal value in the corresponding aspect. Network analysis was performed with Cytoscape v3.10.2. The computation of half maximal inhibitory concentration of NAb (IC_{50}) has been performed using R.

ACKNOWLEDGMENTS

D.R. was supported by CSC (China Scholarship Council, No. 202108070132). This work was supported by Fondation de France (Berthe Fouassier), LabEx LIFESENSES (ANR-10-LABX-65) and IHU FOReSIGHT (ANR-18-IAHU-01), DIM C-BRAINS funded by the Conseil Régional d'Ile-de-France Région Ile-de-France. The authors would like to thank Peggy Sanatine for flow cytometry processing, Jérémie Cosette for advice on data analysis. The authors would like to thank Elena Brazhnikova, Céline Nouvel-Jaillard and the MirCEN NHP facility for assistance with NHP experiments. We would like to thank Melissa Desrosiers and Camille Robert for their technical assistance with the production of plasmids and viral vectors for the NHP experiments. The schematic illustrations were created on BioRender.com.

AUTHOR CONTRIBUTIONS

Conceptualization, S.F., D.A. and D.R.; methodology, D.R., G.C., J.V., E.C. and A.P.; Investigation, S.F., D.A., and D.R.; writing—original draft, D.A. and D.R.; writing—review & editing, C.V., J.P., D.R., D.A., A.G., D.D., H.S. and S.F.; funding acquisition, A.G., D.D. and G.R.; resources, C.V.; supervision, D.A., and S.F.

DECLARATION OF INTERESTS

C.V was the Principal Investigator of the clinical Phase ½ study (NCT02064569).

DATA AVAILABILITY

This study includes no data deposited in external repositories.

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Sylvain Fisson (sylvain.fisson@univ-evry.fr)

Any additional information required to reanalyze the data reported in this work paper is available from the lead contact upon request, Sylvain Fisson (sylvain.fisson@univ-evry.fr)

REFERENCES

Ail D, Ren D, Brazhnikova E, Nouvel-Jaillard C, Bertin S, Mirashrafi SB, Fisson S & Dalkara D (2022) Systemic and local immune responses to intraocular AAV vector administration in non-human primates. *Mol Ther Methods Clin Dev* 24: 306–316

Ali RR, Reichel MB, Thrasher AJ, Levinsky RJ, Kinnon C, Kanuga N, Hunt DM & Bhattacharya SS (1996) Gene transfer into the mouse retina mediated by an adeno-associated viral vector. *Hum Mol Genet* 5: 591–594

Audebert C, Laubreton D, Arpin C, Gandrillon O, Marvel J & Crauste F (2021) Modeling and characterization of inter-individual variability in CD8 T cell responses in mice. *In Silico Biol* 14: 13–39

Bainbridge JWB, Mehat MS, Sundaram V, Robbie SJ, Barker SE, Ripamonti C, Georgiadis A, Mowat FM, Beattie SG, Gardner PJ, *et al* (2015) Long-Term Effect of Gene Therapy on Leber's Congenital Amaurosis. *N Engl J Med* 372: 1887

Bainbridge JWB, Mistry A, Schlichtenbrede FC, Smith A, Broderick C, De Alwis M, Georgiadis A, Taylor PM, Squires M, Sethi C, *et al* (2003) Stable rAAV-mediated transduction of rod and cone photoreceptors in the canine retina. *Gene Ther* 10: 1336–1344

Bainbridge JWB, Smith AJ, Barker SS, Robbie S, Henderson R, Balaggan K, Viswanathan A, Holder GE, Stockman A, Tyler N, *et al* (2008) Effect of gene therapy on visual function in Leber's congenital amaurosis. *N Engl J Med* 358: 2231–2239

Bentler M, Hardet R, Ertelt M, Rudolf D, Kaniowska D, Schneider A, Vondran FWR, Schoeder CT, Delphin M, Lucifora J, *et al* (2023) Modifying immune responses to adeno-associated virus vectors by capsid engineering. *Mol Ther Methods Clin Dev* 30: 576–592

Boisgerault F, Gross DA, Ferrand M, Poupiot J, Darocha S, Richard I & Galy A (2013) Prolonged gene expression in muscle is achieved without active immune tolerance using microrRNA 142.3p-regulated rAAV gene transfer. *Hum Gene Ther* 24: 393–405

Bouquet C, Vignal Clermont C, Galy A, Fitoussi S, Blouin L, Munk MR, Valero S, Meunier S, Katz B, Sahel JA, *et al* (2019) Immune Response and Intraocular Inflammation in Patients With Leber Hereditary Optic Neuropathy Treated With Intravitreal Injection of Recombinant Adeno-Associated Virus 2 Carrying the ND4 Gene: A Secondary Analysis of a Phase 1/2 Clinical Trial. *JAMA Ophthalmol* 137: 399–406

Britten CM, Walter S & Janetzki S (2013) Immunological Monitoring to Rationally Guide AAV Gene Therapy. *Front Immunol* 4

Bucher K, Rodríguez-Bocanegra E, Dauletbekov D & Fischer MD (2021) Immune responses to retinal gene therapy using adeno-associated viral vectors - Implications for treatment success and safety. *Prog Retin Eye Res* 83

Calcedo R, Chichester JA & Wilson JM (2018) Assessment of Humoral, Innate, and T-Cell Immune Responses to Adeno-Associated Virus Vectors. *Hum Gene Ther Methods* 29: 86–95

Calcedo R, Wilson JM, Weber T & Therapy G (2013) Humoral immune response to AAV.

Carpentier M, Lorain S, Chappert P, Lalfer M, Hardet R, Urbain D, Peccate C, Adriouch S, Garcia L, Davoust J, *et al* (2015) Intrinsic Transgene Immunogenicity Gears CD8+ T-cell Priming After rAAV-Mediated Muscle Gene Transfer. *Molecular Therapy* 23: 697

Cehajic-Kapetanovic J, Xue K, Martinez-Fernandez de la Camara C, Nanda A, Davies A, Wood LJ, Salvetti AP, Fischer MD, Aylward JW, Barnard AR, *et al* (2020) Retinal gene therapy in X-linked retinitis pigmentosa caused by mutations in RPGR: Results at 6 months in a first in human clinical trial. *Nat Med* 26: 354

Chan YK, Wang SK, Chu CJ, Copland DA, Letizia AJ, Verdera HC, Chiang JJ, Sethi M, Wang MK, Neidermyer WJ, *et al* (2021) Engineering adeno-associated viral vectors to evade innate immune and inflammatory responses. *Sci Transl Med* 13

Chandler M, Jain S, Halman J, Hong E, Dobrovolskaia MA, Zakharov A V. & Afonin KA (2022) Artificial Immune Cell, AI-cell, a New Tool to Predict Interferon Production by Peripheral Blood Monocytes in Response to Nucleic Acid Nanoparticles. *Small* 18: 2204941

Clare AJ, Langer PM, Ward A, Chan YK, Dick AD & Copland DA (2025) Characterization of the ocular inflammatory response to AAV reveals divergence by sex and age. *Molecular Therapy* 33

Duke BR, El-Husayni S, Jaskolka MC, Erlwein A, Myers R, Pennesi ME, Pierce E, Shearman MS, Zhang K & Mukherjee S (2022) Exploratory Immuno-Safety Profile of EDIT-101, a First-in-Human In Vivo CRISPR

Gene Editing Therapy for CEP290-Related Retinal Degeneration.

Emami MR, Espinoza A, Young CS, Ma F, Farahat PK, Felgner PL, Chamberlain JS, Xu X, Pyle AD, Pellegrini M, *et al* (2023) Innate and adaptive AAV-mediated immune responses in a mouse model of Duchenne muscular dystrophy. *Mol Ther Methods Clin Dev* 30: 90–102

Ertl HCJ (2021) T Cell-Mediated Immune Responses to AAV and AAV Vectors. *Front Immunol* 12

Ertl HCJ (2022) Immunogenicity and toxicity of AAV gene therapy. *Front Immunol* 13

Fischer A, Koeper LM & Vohr HW (2011) Specific antibody responses of primary cells from different cell sources are able to predict immunotoxicity in vitro. *Toxicol In Vitro* 25: 1966–1973

Frazão JB, Thain A, Zhu Z, Luengo M, Condino-Neto A & Newburger PE (2015) Regulation of CYBB Gene Expression in Human Phagocytes by a Distant Upstream NF-κB Binding Site. *J Cell Biochem* 116: 2008–2017

Ge Y, Powell S, Van Roey M & McArthur JG (2001) Factors influencing the development of an anti-factor IX (FIX) immune response following administration of adeno-associated virus-FIX. *Blood* 97: 3733–3737

Gehrke M, Diedrichs-Möhring M, Bogedein J, Büning H, Michalakis S & Wildner G (2022) Immunogenicity of Novel AAV Capsids for Retinal Gene Therapy. *Cells* 11

Hakim CH, Kumar SRP, Pérez-López DO, Wasala NB, Zhang D, Yue Y, Teixeira J, Pan X, Zhang K, Million ED, *et al* (2021) Cas9-specific immune responses compromise local and systemic AAV CRISPR therapy in multiple dystrophic canine models. *Nat Commun* 12

Hamilton BA & Wright JF (2021) Challenges Posed by Immune Responses to AAV Vectors: Addressing Root Causes. *Front Immunol* 12

Hauswirth WW, Aleman TS, Kaushal S, Cideciyan A V., Schwartz SB, Wang L, Conlon TJ, Boye SL, Flotte TR, Byrne BJ, *et al* (2008) Treatment of leber congenital amaurosis due to RPE65 mutations by ocular subretinal injection of adeno-associated virus gene vector: short-term results of a phase I trial. *Hum Gene Ther* 19: 979–990

HAVERKORN MJ, HOFMAN B, Masurel N & van Rood JJ (1975) HL-A linked genetic control of immune response in man. *Transplant Rev* 22: 120–124

Huang D, Taha MS, Nocera AL, Workman AD, Amiji MM & Bleier BS (2023) Cold exposure impairs extracellular vesicle swarm-mediated nasal antiviral immunity. *J Allergy Clin Immunol* 151: 509-525.e8

Jauze L, Vie M, Miagoux Q, Rossiaud L, Vidal P, Montalvo-Romeral V, Saliba H, Jarrige M, Polveche H, Nozi J, *et al* (2024) Synergism of dual AAV gene therapy and rapamycin rescues GSDIII phenotype in muscle and liver. *JCI Insight* 9

Khabou H, Cordeau C, Pacot L, Fisson S & Dalkara D (2018) Dosage Thresholds and Influence of Transgene Cassette in Adeno-Associated Virus-Related Toxicity. *Hum Gene Ther* 29: 1235–1241

Klein SL & Flanagan KL (2016) Sex differences in immune responses.

Leete JC, Zager MG, Musante CJ, Shtylla B & Qiao W (2022) Sources of inter-individual variability leading to significant changes in anti-PD-1 and anti-PD-L1 efficacy identified in mouse tumor models using a QSP framework. *Front Pharmacol* 13: 1056365

Li N, Kumar SRP, Cao D, Munoz-Melero M, Arisa S, Brian BA, Greenwood CM, Yamada K, Duan D & Herzog RW (2024) Redundancy in Innate Immune Pathways That Promote CD8+ T-Cell Responses in AAV1 Muscle Gene Transfer. *Viruses* 16

Lipski DA, Dewispelaere R, Foucart V, Caspers LE, Defrance M, Bruyns C & Willermain F (2017) MHC class II expression and potential antigen-presenting cells in the retina during experimental autoimmune uveitis. *J Neuroinflammation* 14

Liu YL, Wagner K, Robinson N, Sabatino D, Margaritis P, Xiao W & Herzog RW (2003) Optimized production of high-titer recombinant adeno-associated virus in roller bottles. *Biotechniques* 34: 184–189

LUXTURNA® (voretigene neparvovec-rzyl) - Inherited Retinal Disease

Miyoshi H, Takahashi M, Gage FH & Verma IM (1997) Stable and efficient gene transfer into the retina using an HIV-based lentiviral vector

Newman NJ, Yu-Wai-Man P, Carelli V, Moster ML, Biousse V, Vignal-Clermont C, Sergott RC, Klopstock T, Sadun AA, Barboni P, *et al* (2021) Efficacy and Safety of Intravitreal Gene Therapy for Leber Hereditary Optic Neuropathy Treated within 6 Months of Disease Onset. *Ophthalmology* 128: 649–660

Niemeyer GP, Herzog RW, Mount J, Arruda VR, Tillson DM, Hathcock J, Van Ginkel FW, High KA & Lothrop CD (2009) Long-term correction of inhibitor-prone hemophilia B dogs treated with liver-directed AAV2-

mediated factor IX gene therapy. *Blood* 113: 797–806

Ong FS, Kuo JZ, Wu WC, Cheng CY, Blackwell WLB, Taylor BL, Grody WW, Rotter JI, Lai CC & Wong TY (2013) Personalized Medicine in Ophthalmology: From Pharmacogenetic Biomarkers to Therapeutic and Dosage Optimization. *J Pers Med* 3: 40–69

Ozawa S, Miura T, Terashima J, Habano W & Ishida S (2021) Recent Progress in Prediction Systems for Drug-induced Liver Injury Using In vitro Cell Culture. *Drug Metab Lett* 14: 25–40

Penhaskashi J, Sekimoto O & Chiappelli F (2023) Permafrost viremia and immune tweening. *Bioinformation* 19: 685

Pfromm JK, Bonillo M, Dauletbekov D, Bucher K & Fischer MD (2022) Plasmid-mediated gene transfer of Cas9 induces vector-related but not SpCas9-related immune responses in human retinal pigment epithelial cells. *Sci Rep* 12: 13202

Pogorelyy M V., Minervina AA, Touzel MP, Sycheva AL, Komech EA, Kovalenko EI, Karganova GG, Egorov ES, Komkov AY, Chudakov DM, *et al* (2018) Precise tracking of vaccine-responding T cell clones reveals convergent and personalized response in identical twins. *Proc Natl Acad Sci U S A* 115: 12704–12709

Puranen J, Ranta VP, Ruponen M, Urtti A & Sadeghi A (2023) Quantitative intravitreal pharmacokinetics in mouse as a step towards inter-species translation. *Exp Eye Res* 235: 109638

Ramamurthy RM, Atala A, Porada CD & Almeida-Porada G (2022) Organoids and microphysiological systems: Promising models for accelerating AAV gene therapy studies. *Front Immunol* 13

Reichel FF, Dauletbekov DL, Klein R, Peters T, Ochakovski GA, Seitz IP, Wilhelm B, Ueffing M, Biel M, Wissinger B, *et al* (2017) AAV8 Can Induce Innate and Adaptive Immune Response in the Primate Eye. *Molecular Therapy* 25: 2648–2660

Ren D, Fisson S, Dalkara D & Ail D (2022) Immune Responses to Gene Editing by Viral and Non-Viral Delivery Vectors Used in Retinal Gene Therapy. *Pharmaceutics* 14

Ronzitti G, Gross DA & Mingozzi F (2020) Human Immune Responses to Adeno-Associated Virus (AAV) Vectors. *Front Immunol* 11

Rotnemer-Golinkin D & Ilan Y (2022) Personalized-Inherent Variability in a Time-Dependent Immune Response: A Look into the Fifth Dimension in Biology. *Pharmacology* 107: 417–422

Russell S, Bennett J, Wellman JA, Chung DC, Yu Z-F, Tillman A, Wittes J, Pappas J, Elci O, Mccague S, *et al* (2017) Efficacy and safety of voretigene neparvovec (AAV2-hRPE65v2) in patients with RPE65-mediated inherited retinal dystrophy: a randomised, controlled, open-label, phase 3 trial. *www.thelancet.com* 390: 849

Saad FA, Saad JF, Siciliano G, Merlini L & Angelini C (2024) Duchenne Muscular Dystrophy Gene Therapy. *Curr Gene Ther* 24: 17–28

Sahel JA & Dalkara D (2019) Gene therapy for retinal dystrophy. *Nat Med* 25: 198–199

Shen W, Liu S & Ou L (2022) rAAV immunogenicity, toxicity, and durability in 255 clinical trials: A meta-analysis. *Front Immunol* 13: 1001263

Sivan A, Corrales L, Hubert N, Williams JB, Aquino-Michaels K, Earley ZM, Benyamin FW, Lei YM, Jabri B, Alegre ML, *et al* (2015) Commensal Bifidobacterium promotes antitumor immunity and facilitates anti-PD-L1 efficacy. *Science* 350: 1084–1089

Streilein JW (1999) Immunologic privilege of the eye. *Springer Semin Immunopathol* 21: 95–111

Tao Y, Fukushima M, Shimokawa S, Zhao H, Okita A, Fujiwara K, Takeda A, Mukai S, Sonoda KH & Murakami Y (2024) Ocular and Serum Profiles of Inflammatory Molecules Associated With Retinitis Pigmentosa. *Transl Vis Sci Technol* 13: 18

Terrada C, Fisson S, De Kozak Y, Kaddouri M, Lehoang P, Klatzmann D, Salomon BL & Bodaghi B (2006) Regulatory T cells control uveoretinitis induced by pathogenic Th1 cells reacting to a specific retinal neoantigen. *J Immunol* 176: 7171–7179

Terrada C, Pâques M, Fisson S, De Kozak Y, Klatzmann D, Salomon B, LeHoang P & Bodaghi B (2008) [Neoretinal antigen expression: a comparison of anatomical and clinical features of a murine uveoretinitis model]. *J Fr Ophtalmol* 31: 147–154

Tong O & Fairfax BP (2020) Dissecting genetic determinants of variation in human immune responses. *Curr Opin Immunol* 65: 74–78

Vendomèle J, Chauveau GA, Dalkara D, Galy A & Fisson S (2024) Peripheral Cellular Immune Responses Induced by Subretinal Adeno-Associated Virus Gene Transfer Can Be Restrained by the Subretinal-

Associated Immune Inhibition Mechanism. *Hum Gene Ther* 35: 464–476

Vendomèle J, Dehmani S, Khebizi Q, Galy A & Fisson S (2018) Subretinal injection of HY peptides induces systemic antigen-specific inhibition of effector CD4+ and CD8+ T-cell responses. *Front Immunol* 9

VIGNAL-CLERMONT C, YU-WAI-MAN P, NEWMAN NJ, CARELLI V, MOSTER ML, BIOUSSE V, SUBRAMANIAN PS, WANG ANGUOR, DONAHUE SP, LEROY BP, *et al* (2023) Safety of Lenadogene Nolparvovec Gene Therapy Over 5 Years in 189 Patients With Leber Hereditary Optic Neuropathy. *Am J Ophthalmol* 249: 108–125

Wec AZ, Lin KS, Kwasnieski JC, Sinai S, Gerold J & Kelsic ED (2021) Overcoming Immunological Challenges Limiting Capsid-Mediated Gene Therapy With Machine Learning. *Front Immunol* 12

Weleber RG, Pennesi ME, Wilson DJ, Kaushal S, Erker LR, Jensen L, McBride MT, Flotte TR, Humphries M, Calcedo R, *et al* (2016) Results at 2 Years after Gene Therapy for RPE65-Deficient Leber Congenital Amaurosis and Severe Early-Childhood–Onset Retinal Dystrophy. *Ophthalmology* 123: 1606–1620

Xiong W, Wu DM, Xue Y, Wang SK, Chung MJ, Ji X, Rana P, Zhao SR, Mai S & Cepko CL (2019) AAV cis-regulatory sequences are correlated with ocular toxicity. *Proc Natl Acad Sci U S A* 116: 5785–5794

Yamaguti-Hayakawa GG & Ozelo MC (2022) Gene therapy for hemophilia: looking beyond factor expression. *Exp Biol Med (Maywood)* 247: 2223–2232

Yang Y, Zhao Y, Liu X & Huang J (2022) Artificial intelligence for prediction of response to cancer immunotherapy. *Semin Cancer Biol* 87: 137–147

Zheng A, Li Y & Tsang SH (2015) Personalized therapeutic strategies for patients with retinitis pigmentosa. *Expert Opin Biol Ther* 15: 391–402

Figure 1. Inter-individual variability of immune response is observed in the human clinical trial (NCT02064569) after ocular AAV gene transfer.

A Ocular inflammation score (OIS) in human patients who received intraocular AAV injections at baseline and maximal value (obtained within 20 weeks after injection, except patient 003 for 40 weeks).

B, B' Total antibody (TAb) **(B)** and neutralizing antibody (NAb) **(B')** levels measured by ELISA and NAb assay against AAV2 in human patients who received intraocular AAV injections at baseline and maximal value (obtained within 20 weeks after injection). IC_{50}: half maximal inhibitory concentration.

C-F Immune profiles of individual patients, who received dose 1 – $9x10^9$ vg **(C)**, dose 2– $3x10^{10}$ vg **(D)**, dose 3– $9x10^{10}$ vg **(E)**, dose 4– $1.8x10^{11}$ vg **(F)** of AAV, showing maximum ocular OIS, foldchange (FC) of anti-AAV2 TAb and NAb, anti-AAV2 cellular immune response. Each slice of the pie corresponds to one immune parameter which is normalized against the highest value for the parameter. The code of each patient is shown on the above each chart. vg: vector genome.

Figure 2. Inter-individual variability of immune response is observed in the NHP model after ocular AAV gene transfer.

A Schematic of ocular gene injections in NHPs.

B, B' Total antibody (TAb) **(B)** and neutralizing antibody (NAb) **(B')** levels measured by ELISA and NAb assay against AAV2 in NHPs who received intraocular AAV injections with dose $1 – 5x10^{11}$ vg before (BI) and 2-3 month post injection (PI). IC$_{50}$: half maximal inhibitory concentration.

C Immune profiles of individual NHPs showing FC of anti-AAV2 TAb and NAb, grading score at month 1 for Anterior Chamber Cells (ACC), Anterior Chamber Flare (ACF), Vitreous Haze (VH) and Vitreous Cells (VC). Each slice of the pie corresponds to one immune parameter which is normalized against the highest value for the parameter. The code of each NHP is shown on the above each chart. Vg: vector genome.

Figure 3. Systemic adaptive immune responses are induced after subretinal injections of AAV8-GFP-HY in mice.

A Schematic representation of the experimental procedure showing the subretinal injection of AAV8-GFP-HY in mouse, followed by harvest of the spleen and serum 21d post-injection (PI) to test cellular and humoral responses against the capsid and transgene.

B, C T cell activation measured by ELISpot assay against **(B)** the AAV8 capsid and **(C)** the transgene-HY.

D, E Antibody levels measured by ELISA against **(D)** the AAV8 capsid and **(E)** the transgene-GFP.

Data information: Results obtained from 2 independent experiments (n=10 per group). Bars correspond to mean + SEM. *P < 0.05, **P < 0.01, ***P < 0.001, and ****P < 0.0001 with unpaired Mann-Whitney test.

Figure 4. Global view of the systemic pro-inflammatory cytokine profile in mouse 21d post-injection (PI) of AAV-GFP-HY.

A, A' Cytokine profiles of PBS-injected group with **(A)** AAV *in vitro* stimulation and **(A')** peptide HY (pHY) stimulation.

B, B' Cytokine profiles of AAV-injected group with **(B)** AAV *in vitro* stimulation and **(B')** pHY stimulation. The 100% radar scale fits to the maximum value of cytokine secretion, and the black area values correspond to the mean of these cytokine secretions.

Cytokines tested are ILs: Interleukin, TNFα: Tumor Necrosis Factor alpha, IFNγ: Interferon gamma, GM-CSF: Granulocyte-Macrophage Colony-Stimulating Factor, RANTES: Regulated upon Activation, Normal T cell Expressed and Secreted (also known as CCL5), MCP-1: Monocyte Chemoattractant Protein 1 (also known as CCL2). The concentration of each cytokine in each condition can be found in Supplementary figure 2, 3 and Supplementary table 3.

Figure 5. Local transgene expressions in the retina are correlated with cytotoxicity against transgene[+] cells post-injection (PI) of AAV-GFP-HY.

A HY and GFP transgene expression levels in female mouse retina 21d PI measured by ddPCR. AU: Arbitrary Units.

B Correlation between HY and GFP expressions.

C Activation of cytotoxic cells specific to the HY peptide 21 PI evaluated by *in vivo* cytotoxicity assay.

D, E Correlation between the survival of male cells and local transgene expression in AAV-injected mice for **(D)** GFP and **(E)** HY.

Data information: Results obtained from 2 independent experiments (n=8 per group). Bars correspond to mean + SEM. *P < 0.05, **P < 0.001, and ***P < 0.0001 with unpaired Mann-Whitney test. R^2 is calculated via linear regression.

Figure 6. Local inflammation and correlated expressions between MHC II molecules and CYBB are found in the retina post-injection (PI) of AAV-GFP-HY.
A Gene expression of H2-Eb1 and H2-Ab1 measured by ddPCR in the mouse retina 21d PI. AU: Arbitrary Units.
B Correlation between H2-Eb1 and H2-Ab1 expression levels.
C Gene expression of CYBB measured by ddPCR in the mouse retina 21d PI.
D, E Correlation between (D) CYBB and H2-Eb1 expression; and **(E)** CYBB and H2-Ab1 expression.
Data information: Results obtained from 2 independent experiments (n=8 per group). Bars correspond to mean + SEM. *P < 0.05, **P < 0.001, and ***P < 0.0001 with unpaired Mann-Whitney test. R^2 is calculated via linear regression.

Figure 7. Diversity in immune responses is noticed in syngeneic mice 21d post-injection (PI) of AAV-GFP-HY.

A Immune profiles of each individual mouse showing transgene expression (HY-Tg and GFP-Tg), local inflammation (H2-Eb1, H2-Ab1 and CYBB) and humoral immune response (AAV-Ab and GFP-Ab) and cellular immune response (%male, ELISpot-HY, ELISpot-DBY, ELISpot-UTY, ELISpot-AAV-PGK-Luc2 and ELISpot-AAV-PGK-GFP-HY) 21d PI of PBS (control) and AAV8-GFP-HY (AAV).

B Cytokine profiles of individual mouse showing cytokine levels (AAV *in vitro* stimulation: IL-2, IFNγ, TNFα and RANTES; HY *in vitro* stimulation: IFNγ, TNFα, IL-10 and RANTES) 21d PI of PBS (control) and AAV8-GFP-HY (AAV). Each slice of the pie corresponds to one immune parameter which is normalized against the highest value into percentage for the parameter.

C Principal component analysis of PBS-injected mice (blue dots) compared to AAV-injected mice (red dots). Data information: The mouse code is shown related to the dot. The code of mouse is shown on the above each chart, Results obtained from 2 independent experiments (n=8 per group).

Figure 8. Immune parameters show no correlation in syngeneic mice 21 days following AAV-GFP-HY delivery.

A Correlation matrix showing the coefficient of determination of each pair of immune parameters tested experimentally. Positive correlations are marked in shades of red with R^2 above 0 and negative correlations in shades of blue with minus label (-) before R^2. The intensity of shade corresponds to the strength of correlation.

B Network analysis map generated based on the correlation matrix. The color of the lines connecting two parameters (edges) indicate positive (red), negative (blue) or no (grey) correlation and the thickness of the edges indicate the strength of correlation with thicker lines indicating higher correlation and dotted lines showing no correlation.

Supplementary Figure 1. CD4[+] T cell and CD8[+] T cell activation by peptides.
A T cell activation specific to AAV capsid measured by ELISpot assay.
B CD8[+] T cell activation specific to UTY peptides.
C CD4[+] T cell activation specific to DBY peptides.
Data information: Results obtained from 2 independent experiments (n=10 per group). Bars correspond to mean + SEM. *P < 0.05, **P < 0.01, ***P < 0.001, and ****P < 0.0001 with unpaired Mann-Whitney test.

Supplementary Figure 2. Cytokine secretion in mouse spleen cells 21d post-injection (PI) of AAV-GFP-HY stimulated with AAV *in vitro*.

A-D Expression of cytokines participating in (A) Inflammation and cellular migration (IL-1b, IL-6, MCP-1, RANTES), **(B)** Th1 function (IL-2, IFNγ, TNFα, GM-CSF), **(C)** Th2 function (IL-4, IL-10, IL-13) and **(D)** Th17 function (IL-17).

Data information: Results obtained from 2 independent experiments (n=10 per group). Bars correspond to mean + SEM. *P < 0.05, **P < 0.01, ***P < 0.001, and ****P < 0.0001 with unpaired Mann-Whitney test.

Supplementary Figure 3. Cytokine secretion in mouse spleen cells 21d post-injection (PI) of AAV-GFP-HY stimulated with HY peptides (pHY) *in vitro*.
A-D Expression of cytokines participating in (A) Inflammation and cellular migration (IL-1b, IL-6, MCP-1, RANTES), **(B)** Th1 function (IL-2, IFNγ, TNFα, GM-CSF), **(C)** Th2 function (IL-4, IL-10, IL-13) and **(D)** Th17 function (IL-17).
Data information: Results obtained from 2 independent experiments (n=10 per group). Bars correspond to mean + SEM. *P < 0.05, **P < 0.01, ***P < 0.001, and ****P < 0.0001 with unpaired Mann-Whitney test.

Supplementary Table 1. Characteristics of individual immune responses in patients

Dose Level, Patient No.	Maximum, OIS	IgG (×10^{-3} AU/mL)		NAb (IC$_{50}$)		Cellular Response
		Baseline	Maximum Value	Baseline	Maximum Value	
9E9 vg						
001	0	7115	7565	1 440	3050	Negative
003	0.5	3520	14895	520	3650	Negative
005	0	1903	5355	360	3400	Positive
3E10 vg						
006	1	155	10665	23	2200	Negative
007	1.5	33	336	0	40	Negative
008	0.5	536	2585	48	488	Negative
9E10 vg						
009	5	2110	14240	420	3200	Negative
011	1.5	13	1148	0	120	Negative
012	1	32	177	0	60	Negative
017	0.5	918	1215	228	396	Negative
018	9.5	10295	67240	2850	40 000	Positive
019	1.5	26	78	0	24	Negative
1.8E11 vg						
013	1.5	46	760	0	68	Negative
014	1	4952	18010	630	8400	Negative
015	0.5	155	1229	0	220	Negative

AU, arbitrary unit; IC$_{50}$, half maximal inhibitory concentration; NAb, neutralizing antibody; OI, ocular inflammation; OIS, ocular inflammation score; vg, viral genome.

Supplementary Table 2. Characteristics of individual immune responses in NHPs

No.	Eye	Anti-AAV2 TAb (%)		Anti-AAV2 NAb (IC$_{50}$)		Inflammation at Month 1			
		T0	T2	T0	T2	ACC	ACF	VH	VC
NHP1	LE	100	453.86	23.64	443.83	3	1	0	1
	RE					1	0	0	1
NHP2	LE	100	307.03	3.35	487.05	0.5	0	0	1
	RE					0.5	0	0	1
NHP3	LE	100	537.08	3.56	159.55	0	0	0	1
	RE					0.5	0	0	1
NHP4	LE	100	408.2	1.57	309.11	1	0	0	1
	RE					1	0	0	1
NHP5	LE	100	142.86	3.16	17.7	1	0	0	1
	RE					1	0	0	1
NHP6	LE	100	383.36	7.94	247.3	0.5	0	0	1
	RE					1	0	0	1
NHP7	LE	100	366.88	8.74	432.64	0.5	0	0	1
	RE					0.5	0	0	1
NHP8	LE	100	230	100	181.75	1	0	0	1
	RE					1	0	0	1

TAb, total antibody; NAb, neeutralizing antibody; IC$_{50}$, half maximal inhibitory concentration;
ACC, anterior chamber cell; ACF, anterior chamber flare; VH, vitreous haze; VC, vitreous cell; LE, left eye; RE, right eye.

Supplementary Table 3. Characteristics of individual immune responses in mice

Mouse No.	Local transgene and inflammation					Systemic cellular immune response					IVC
	ddPCR					ELISpot					
	GFP	HY	H2-Ab1	H2-Eb1	CYBB	UTY	DBY	HY	AAV-GFP-HY	AAV-Luc2	% male
1-1	0.01	0.00	0.36	1.04	0.19	62.96	8.93	8.38	0.00	1.64	35.71
1-2	0.04	0.00	2.72	3.07	0.48	0.00	142.86	36.31	0.41	0.00	26.67
1-3	0.02	0.00	0.00	3.17	0.26	0.00	0.00	0.00	0.00	0.00	50.00
1-4	0.03	0.01	5.64	5.32	0.60	0.00	0.00	2.79	0.41	0.00	34.69
1-5	0.07	0.00	4.47	7.75	0.78	0.00	35.71	0.00	0.00	0.00	50.00
2-1	0.00	0.00	6.30	12.11	0.66	0.70	18.68	18.96	1.39	1.11	67.59
2-2	0.00	0.00	4.58	7.86	1.30	2.11	42.02	18.96	0.56	1.67	54.55
2-3	0.00	0.00	5.46	8.27	1.08	0.00	77.82	23.24	1.11	0.56	55.68
1-17	47.83	0.00	146.78	199.45	14.99	266.67	857.14	865.92	71.31	75.41	0.00
1-18	1.72	6.03	60.10	68.57	6.06	118.52	321.43	226.26	13.11	2.05	0.00
1-19	86.96	132.01	29.15	30.37	1.49	114.81	294.64	212.29	0.41	4.92	20.74
1-20	3.28	4.89	42.95	45.05	5.11	459.26	1607.14	631.28	22.95	18.85	0.00
2-21	47.30	96.70	86.38	132.65	10.15	262.92	227.24	263.00	18.64	9.18	9.52
2-23	68.53	235.40	234.72	497.38	46.56	395.08	373.54	444.04	24.77	12.24	16.88
2-24	174.13	285.04	296.05	379.32	34.32	136.38	244.36	222.63	6.68	5.57	29.17
2-25	167.71	319.50	138.01	204.76	10.77	72.41	197.67	169.42	26.71	10.30	19.05

IVC, in vivo cytotoxicity; CBA, cytometric bead array.

Humoral immune response		Cytokine profile									
---	---	---	---	---	---	---	---	---	---	---	
ELISA		CBA, AAV *in vitro* stimulation				CBA, HY *in vitro* stimulation					
Anti-AAV TAb	Anti-GFP TAb	IL-2 (AAV)	IFN-γ (AAV)	TNF-α (AAV)	RANTES (AAV)	IL-2 (HY)	IFN-γ (HY)	TNF-α (HY)	IL-10 (HY)	RANTES (HY)	
1.66	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
1.95	0.00	0.00	0.00	0.00	0.00	1.85	6.21	0.00	23.43	0.00	
1.66	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
2.25	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
2.83	0.00	0.00	0.00	0.00	0.00	6.74	8.06	0.45	2.31	23.79	
22.64	0.00	0.92	0.00	45.69	8.10	7.00	0.00	5.80	2.64	0.00	
25.33	0.00	0.75	0.00	43.52	28.28	5.04	2.03	0.00	0.33	0.00	
22.52	0.00	0.00	0.39	38.48	5.88	3.54	4.30	0.00	0.00	0.00	
7866.66	267205.17	10.63	162.53	33.69	124.50	3.77	308.17	21.50	88.62	79.45	
4473.36	58669.90	3.08	3.71	2.06	3.71	2.41	19.65	1.79	13.62	5.90	
3166.20	19897.68	0.00	0.00	2.61	42.08	0.00	2.33	2.39	11.59	50.94	
1478.52	36187.40	5.10	9.01	1.70	0.00	4.35	90.91	23.07	69.44	137.00	
5077.02	439856.61	9.00	89.96	99.32	42.01	0.00	129.36	24.97	12.23	57.40	
5780.30	444835.37	14.28	19.68	57.61	79.23	2.54	102.78	25.17	32.01	238.93	
5030.14	580257.57	10.10	6.37	43.72	18.61	2.30	7.34	0.00	0.00	11.48	
4127.59	788037.71	8.08	9.73	106.33	24.88	4.42	56.76	18.15	12.12	41.35	