TGF-beta coordinates changes in Keratin gene expression during complex tissue regeneration.

Dipak D Meshram*[§], Yuchang Liu*[§], Molly Worth*, Changqing Zhang[b], Tom J. Carney[b] and Henry H. Roehl*

*School of Bioscience
The University of Sheffield
Firth Court, Western Bank
Sheffield S10 2TN
United Kingdom
[b]Lee Kong Chian School of Medicine,
Nanyang Technological University,
Singapore, Singapore.

[§] These authors contributed equally
Corresponding author: h.roehl@sheffield.ac.uk

Abstract:

Zebrafish larval tail regeneration can be broadly broken down into three phases: wound repair, initiation of regeneration and redevelopment of the tail. We and others have shown that activation of the Hedgehog and TGF-beta signalling pathways is required to initiate tail regeneration. In this study we initially focus on these two pathways to better understand their role regeneration. We have found that Hedgehog signalling acts upstream of the TGF-beta pathway by activating transcription of the TGF-beta ligand *tgfb1a*. We have identified differentially expressed genes that are regulated after tail excision as well as those that are dependent upon TGF-beta signalling. From this analysis we found that Keratin genes fall into categories of co-expression: one set that is upregulated by TGF-beta signalling after injury (krt_{up}) and a second set that is downregulated (krt_{down}). To see whether these two sets show co-expression more broadly, we analysed single cell RNA sequencing data and found that krts fall into the same categories during zebrafish development. We have mapped the expression of a representative member of krt_{up} (*krt97*) and krt_{down} (*krt5*) sets during development and regeneration and show that *krt97* is found in epithelia that is moving and *krt5* is expressed in epithelia that is stationary. Based upon our results we propose that the krt_{up} set allows for collective epithelial migration while the krt_{down} set inhibits mobility. These results reveal one mechanism by which TGF-beta may initiate regeneration and could help to explain how changes in cytoskeletal composition influence cell movement during development and repair.

Introduction:

Although mammals are able to heal wounds and regrow many tissues such as skin and muscle, the regeneration of complex structures is limited. Aquatic vertebrates on the other hand, are able to regenerate large portions of organs including limb, tail, spinal cord, retina and heart(Poss and Tanaka 2024). When comparable damage occurs in mammals, tissue usually fails to regrow and scarring occurs. One possible explanation for the poor regenerative potential observed in mammals may be a failure to recruit competent cells to the site of injury in sufficient numbers to restore the missing tissue. Thus, the characterisation of signalling pathways that are able to induce regenerative cell

migration in model organisms may aid in the development of novel regenerative medicine strategies.

Figure 1. Signalling events during tail regeneration. (a) Larval tail excision removes the end of the zebrafish tail and includes different cell types. **(b)** Within 0.5 hours post excision (hpe) early wound signals are released and the wound is closed. **(c)** The body axis shortens and the notochord bead is extruded. **(d)** The notochord bead becomes a HH signalling centre that activates regeneration. **(e)** Regenerative cells migrate towards the stump. **(f)** The wound epithelium and blastema form. **(g)** The tail reforms. **(h and i)** Models presented in this paper.

Zebrafish larval tail excision offers a rapid and simple model to understand the basic principles of regeneration(Fig. 1a)(Rojas-Muñoz et al. 2009; Romero et al. 2018). Excision is typically performed between 48 and 72 hours post fertilisation (hpf) and involves removal of some of the larval fin (finfold), muscle, neural tube, notochord and other tissue. Following excision, early wound signals (ATP, Calcium and ROS) are released and the wound rapidly closes (Fig. 1b)(Niethammer 2016). The body axis shortens and a ball of notochord cells forms at the stump (called the notochord bead)(Fig. 1c-d). The notochord bead expresses Hedgehog (HH) ligands which induce migration of regenerative cells

to the stump (Fig. 1d-e). This results in the formation of the blastema and wound epithelium (Fig. 1f). The blastema and wound epithelium express signalling pathways which orchestrate the redevelopment of the tail (Fig. 1g). Regeneration of the tail completes by around 72 hours post excision (hpe).

The TGF-beta pathway plays crucial roles in wound healing in mammalian systems and is involved with wound closure, fibroblast activation, inflammation and fibrosis(Massagué and Sheppard 2023). It has also been shown to act during the regeneration of different zebrafish tissues including heart, spinal cord, lateral line and adult skin(Richardson et al. 2016; Bensimon-Brito et al. 2020; Keatinge et al. 2021; Hsu et al. 2024). Recent analysis has shown that TGF-beta signalling acts during larval tail regeneration and is linked closely with pro-regenerative metabolic reprogramming(Sinclair et al. 2021). How the TGF-beta pathway relates to the existing signalling network for tail regeneration (Fig. 1a-g) is not clear.

Keratins (Krts) are intermediate filament proteins that are found mainly in the cytoskeleton of epithelial cells and are associated with desmosomes and hemidesmosomes. They give strength to many structures including skin, hair and nails. In addition to their structural roles they have effects on diverse cellular processes such as cell signalling, metabolism and mechanotransduction(Redmond and Coulombe 2021). Studies in zebrafish have primarily used *krts* as markers of epithelia and transgenic fish carrying fluorescent proteins driven by *krt* promoters are widely used(Love, Huber, and Anders 2014; Gong et al. 2002; Lee, Asharani, and Carney 2014).

In this study we have propose that the TGF-beta pathway plays a central role during tail regeneration. We find that it is activated by notochord-derived HH signalling through the upregulation of transcription of the ligand *tgfb1a* in the notochord bead (Fig. 1h). Notochordal TGF-beta signalling then inhibits HH signalling by repressing transcription of *indian hedgehog a* (*ihha*) in the notochord and further activates *tgfb1a* in the notochord and surrounding tissue. To better understand the downstream effects of TGF-beta signalling we use bulk RNA-sequencing (RNA-seq) to identify differentially expressed genes (DEGs) after tail excision and after pharmacological inhibition of TGF-beta signalling. We identify members of the *krt* gene family as targets of the TGF-beta pathway after excision, and find that they fall into two categories of co-expression (synexpression): one set that is upregulated by TGF-beta signalling after injury (krt_{up}) and a second set that is downregulated by TGF-beta signalling after injury (krt_{down}). Next we focus on two *krt* genes, *krt97* and *krt5* as representative members of krt_{up} and krt_{down} to examine their expression during development and regeneration. We find that whereas *krt97* is expressed in epithelia that are actively moving, *krt5* is expressed in epithelium that is stationary. Using developmental single cell RNA sequencing (scRNA-seq) datasets we show that genes in krt_{up} and krt_{down} tend not to be co-expressed suggesting that these two groups have distinct functions during development. Collectively these data

suggest a model where TGF-beta allows for collective cell movement in the epithelia by modulating expression of different members of the *krt* gene family.

Results:

TGF-beta signalling acts downstream of HH signalling.

Pharmacological inhibition of the HH and TGF-beta signalling pathways have striking similarities: In both cases the wound closes and the notochord bead forms, but there is no recruitment of regenerative cells to the stump(Sinclair et al. 2021; Romero et al. 2018). We have previously shown that HH ligands *ihhb* and *shha* are expressed in the notochord both before and after tail excision. To begin to understand the relationship between the HH and TGF-beta pathways we used standard RNA in situ methods(Thisse and Thisse 2014) to identify *tgfb1a* as a candidate ligand gene that is expressed in the notochord bead immediately after wounding (Fig. 2a). Expression is initially just in the notochord and then spreads to the surrounding tissue. To determine whether the two pathways regulate each other we treated fish with either a HH pathway inhibitor (cyclopamine) or a TGF-beta pathway inhibitor (SB431542). We found that cyclopamine treatment blocks *tgfb1a* transcriptional activation (Fig. 2b) and SB431542 treatment increases *ihha* expression (Fig. 2c). These two findings indicate that TGF-beta signalling is likely to act downstream of the HH pathway and forms a negative feedback loop to modulate HH activity. In addition we found that TGF-beta activates its own transcription especially in the tissue surrounding the notochord bead (Fig. 2d). We have previously found that while the standard whole-mount RNA in situ methods detect expression in the notochord bead, it does not detect genes expressed in the intact notochord(Romero et al. 2018). To investigate whether *tgfb1a* is expressed in notochord cells prior to injury, we switched to the hybridization chain reaction fluorescent (HCR) in situ method which utilises much smaller probes(Choi, Schwarzkopf, and Pierce 2020). Whereas, *ihhb* is expressed through-out the notochord before and after injury, *tgfb1a* is only activated in the notochord bead upon injury (Fig. 2e). Together these findings suggest a model whereby the formation of the notochord bead causes the induction of *tgfb1a* expression by HH signalling.

Figure 2. *tgfb1a* **expression is activated by Hedgehog signalling in the notochord bead. (a)** Expression of *tgfb1a* at three time points after tail excision. Expression is initially confined to the notochord, but spreads to surrounding tissue over time. Scale bar 100um **(b)** Cyclopamine treatment from -2 to 24hpe results in a significant reduction in *tgfb1a* expression. **(c)** SB431542 treatment from -2 to 24hpe results in a significant increase in *ihhb* expression. **(d)** SB431542 treatment from 4 to 24hpe results in a significant decrease in *tgfb1a* expression in the tissue that surrounds the notochord bead (arrow heads). **(e)** HCR in situs of an uninjured fish at 72hpf (top row) and injured fish at 24hpe (bottom row). In uninjured fish *ihhb* is expressed throughout the notochord prior to excision and *tgfb1a* is absent. After excision *tgfb1a* expression is activated only in the notochord bead (yellow arrows). Units for the graphs in b-d are thresholded pixels (see methods for details). Asterisks indicate significance (* = 0.05; ** = 0.01; *** = 0.001; **** = 0.0001)

results using the unpaired t test and ns is not shown to be significant.SB431542 concentration was 50uM and cyclopamine concentration was 20uM.

Transcriptional analysis of TGF-beta signalling during regeneration

To begin to understand the role of TGF-beta signalling in regenerative cell recruitment, we performed bulk RNA-seq to identify DEGs after tail excision (Fig. 3a). To focus on genes that are regulated at the start of regeneration, fish were processed at 18 hours post excision (hpe). RNA was collected using four treatments: DMSO (vehicle control) unoperated, DMSO operated, SB431542 unoperated and SB431542 operated (Fig. 3b). Three replicates were done for each treatment and the data was processed using the DESeq2 package in R(Choi, Schwarzkopf, and Pierce 2020; Love, Huber, and Anders 2014). One conclusion that can be drawn from this analysis is that DEGs

Figure 3. **Transcriptome changes due to injury and TGF-beta activation. (a)** Preparation of samples. Operated samples have approximately 200um of posterior tail tissue removed and are incubated until 18hpe. A second 200um tissue is then removed and processed for sequencing. The unoperated samples take a similar region of the tail for processing. **(b)** Four-way comparisons are made between the samples. **(c)** Comparison of potential TGF-beta pathway targets after injury to potential TGF-beta pathway targets without injury. The X-axis is SB431542 operated compared to DMSO operated (2 in panel b) and the Y-axis is SB431542 unoperated compared to DMSO unoperated (3 in panel b). Values are in Log2. Only DEGs that have an adjusted P-value (Padj) value <0.05 in the SB431542 operated compared to DMSO operated set are plotted. DEGs in the upper right quadrant are genes that are potentially suppressed by TGF-beta

signalling in both operated and unoperated samples. DEGs in the lower left quadrant are those that are potentially activated by TGF-beta signalling in both operated and unoperated samples. The lines indicate the regression with 95% confidence. The regression slope is 0.62 indicating that regulation is stronger in the injured dataset. **(d)** Comparison of potential DEGs after injury to potential TGF-beta pathway targets after injury. The X-axis is DMSO operated compared to unoperated (1 in panel b) and the Y-axis is SB431542 operated compared to DMSO operated (2 in panel b). Values are in Log2. Only DEGs that have a Padj value <0.05 in the operated compared to unoperated set are plotted. DEGs in the upper left quadrant are genes that are potentially suppressed by TGF-beta signalling following injury. DEGs in the lower right quadrant are those that are potentially activated by TGF-beta signalling after injury. The lines indicate the regression with 95% confidence. **(e)** Comparison of potential targets after injury to potential TGF-beta pathway targets after injury. The X-axis is DMSO operated compared to unoperated (1 in panel b) and the Y-axis is SB431542 operated compared to SB431542 unoperated (4 in panel b). Values are in Log2. Only DEGs that have a Padj value <0.05 in the SB431542 operated compared to SB431542 unoperated set are plotted. The solid line represents a slope of one, and the dotted line represents the linear regression.

that are found in the SB431542 operated versus DMSO operated comparisons tend to also be DEGs in the SB431542 unoperated versus DMSO unoperated set (Fig. 3c). This suggests that genes that are targets of TGF-beta signalling during regeneration are also responding to TGF-beta activity in uninjured fish. This is unexpected as treatment with SB431542 in unoperated fish confers no visible changes. There are however some DEGs that are regulated by SB431542 in operated fish that do not show significant regulation by SB431542 in unoperated fish. These genes are potential wound/regeneration specific targets of the TGF-beta pathway (see Supplemental Table 1).

A second conclusion from the RNA-seq comparisons is that genes that are regulated during regeneration are usually influenced by SB431542 treatment (Fig. 3d). For example, DEGs that are upregulated following injury are likely to also be downregulated when injured fish are treated with SB431542. This supports the model that TGF-beta is pro-regenerative and is required during the onset of regeneration to recruit cells. However, there are some DEGs that do not fit this model (see upper right and lower left quadrants in Fig.3d). For example DEGs in the upper right quadrant of Fig3d are upregulated following injury and are upregulated when injured fish are treated with SB431542. It is not immediately clear why this is, but one possible explanation is that there is negative feedback affecting these genes and that testing at earlier time points would find that they are initially downregulated by TGF-beta. The most significant DEGs that are identified by this comparison are presented in Supplemental Table 2.

Further evidence that TGF-beta has a pro-regenerative role comes from comparison of DEGs after injury in DMSO to DEGs after injury in fish treated with SB431542 (Fig. 3e). In this graph, DEGs that are unaffected by the presence of SB431542 after injury fall on a slope of 1. The regression slope is closer to 0.9 indicating that there is a bias towards DEGs that are injury specific and are responding to TGF-beta. For example, genes that fall below the slope of 1 line in the upper right quadrant are predicted to be activated by TGF-beta whilst those that fall above the slope of 1 line in the lower left quadrant are predicted to be repressed by TGF-beta during regeneration. A subset of these genes are presented in Supplemental Table 3.

To find genes that are injury-specific but not TGF-beta targets we can take DEGs that fall close to the slope of 1 in Fig3e. We can also identify this type of gene by selecting DEGs that lie close to the X-axis in Fig3d. By applying both criteria we identified a subset of DEGs that are potential TGF-beta independent regeneration genes (Supplemental Table 4).

Figure 4. Zebrafish *krt* gene regulation and relatedness. (a) A table with zebrafish *krt* genes, their ensemble identifiers, the fold level of change in all four comparisons, the adjusted counts for each sample and their allocation into *krt*_{up} or *krt*_{down} groups. **(b)** Molecular phylogeny dendrogram with zebrafish (zKrt) and human (hKRT) representatives of the Keratin family. The arrows indicate upregulation or downregulation as shown in panel a. Black arrows indicate significance (P < 0.05) and grey arrows indicate a trend that was not significant. Phylogram tree and alignments were done with "one click" settings www.phylogeny.fr(Dereeper et al. 2008) **(c)** The genomic locations of the zebrafish *krts* with an arrowhead showing their chromosomal orientation and colour showing upregulation (pink) or downregulation (blue) by TGF-beta. The genome assembly for this analysis was GRCz11 (www.ensemble.org).

Members of the Keratin family show TGF-beta-dependent responses to injury

To learn more about coordinated gene expression during regeneration we searched for families of genes in the RNA-seq data (Supplemental Table 5). Of these, the Krt family stands out because most members are regulated by TGF-beta and these members are either activated during regeneration (krt_{up}) or inhibited (krt_{down}) (Fig. 4a). We grouped the *krts* as being in these synexpression groups based upon being significant ($P < 0.05$) in at least one of the comparisons. There are nine *krts* that do not fall into these groupings. Two of these, *krt18a.2* and *krt95*, were not identified in our analysis. Of the remaining seven, six are expressed at low levels at the time of our assay which perhaps affects our detection, and the seventh (*krt222*) is only distantly related to other members of the *krt* family.

The zebrafish *krt* family is smaller than the mammalian family and is less well studied(Krushna Padhi, Akimenko, and Ekker 2006; Ho et al. 2022). To begin to understand the significance of these two groupings, we first performed phylogenetic analysis to see whether members of Krt$_{up}$ or Krt$_{down}$ are more closely related to each other based upon their predicted protein sequence (Fig. 4 b). We have included members of the human Krt (hKRT) family in this analysis for comparison. There are three type II Krts with one Krt$_{up}$ and two Krt$_{down}$ members. These fall next to the human type II Krts in one cluster. The type I branch has greater variation with one of the human Krts (hKRT18) showing more similarity to zebrafish genes than to other human Krts. Within the type I branch there is some correlation between expression and relatedness. Krt$_{up}$ and Krt$_{down}$ appear to separate between Krt15 and Krt94. Intriguingly, the members that show regulation but a P value that is greater than 0.5 do still show a correlation in their relatedness. For example, although *krt1-c5* does not show significance in pairwise comparisons, it does show the expected changes in expression for Krt$_{up}$ genes (Fig 4a) and falls into the Krt$_{up}$ phylogenetic branch. Likewise, *krt1-19d* has the expected changes in expression for Krt$_{down}$ and falls into the Krt$_{down}$ branch. The correlation in the type I group between expression and relatedness may indicate that there are structural differences between these Krt$_{up}$ and Krt$_{down}$ members and that the two groups were formed by one evolutionarily event.

To see whether this correlation extends to their chromosomal organisation, we looked at the genomic organisation of *krts*. *krts* are found clustered on six chromosomes. The cluster on chromosome 19 contains up to three members of *krt$_{down}$* and the cluster on chromosome 22 has two members of *krt$_{up}$*. Three of the other chromosomes are mixed synexpression genes and chromosome 5 has a single *krt* gene. Overall the genomic organisation does not show a strong correlation with the type of synexpression.

***krts* have distinct domains of expression during regeneration and development**

To understand the relevance of *krt* regulation at the tissue level we decided to do HCR in situ analysis. For this we have focused on *krt5* as a representative of *krt*$_{down}$ and *krt97* as a representative of *krt*$_{up}$. We first tested the response of these at three stages, uninjured (96hpf), 24hpe(96hpf) and 48hpe(120hpf) (Fig. 5). In fish at these stages we found that *krt97* and *krt5* are expressed in distinct domains. *krt97* is primarily in epithelial cells of the finfold in a crescent shape around the end of the tail (Fig. 5a). This region of expression is removed following tail excision. *krt5* on the other hand is found in the epithelia that sits on the trunk. At 24hpe epithelial cells have gathered on the end of the stump to form the wound epithelium. At this time, *krt97* expression is upregulated in wound epithelium and *krt5* expression is reduced in the trunk epithelia nearest to the stump (Fig. 5b). By 48hpe the fin fold is reforming and krt97 is expressed in the new area of growth (Fig. 5d). On the other hand *krt5* is strongly downregulated near to the regrowing tail. Inhibition of TGF-beta with SB431542 during regeneration results in a strong increase in *krt5* and a reduction in *krt97* (Fig. 5c and e) consistent with the RNA-seq data. To further confirm our results we probed scRNA-seq data for fish at 24hpe and 48hpe(Sinclair et al. 2021). We see that *krt97*[+] and *krt5*[+] cells are in distinct domains with *krt97*[+] cells increasing after excision and *krt5*[+] cells decreasing. Pseudo bulk RNA-seq analysis of this scRNA-seq data suggests an increase of 1.5 fold for *krt97* at 24hpe and a decrease of *krt5* expression to 0.5 at 24hpe and 0.38 at 48hpe. Together these data suggest that *krts* respond to injury in a TGF-beta dependent fashion and are expressed in distinct compartments of the skin.

Figure 5. *krt97* and *krt5* are inversely regulated during tail regeneration. (a) In situ analysis of *krt97* and krt5 in uninjured and injured fish treated with SB431542. Larvae were treated with DMSO or 50 μM SB431542 for 2 hours prior to tail excision and kept in the chemical solution until 24 or 48 hpe. Yellow dotted lines indicate the excision plane. Yellow stars indicate non-specific fluorescence in the vasculature. Scale bar is 50um **(b)** Quantification of fluorescent in situs as shown in (a). Asterisks indicate significance as in Fig.2 resulting from the unpaired t test. **(c)** Expression of *krt97* and *krt5* in scRNA-seq cell clusters representing periderm (P) and epithelia (E) during regeneration.

Given that regeneration of the fin fold is likely to recapitulate aspects of fin fold development, we decided to look at tail developmental stages to compare development to regeneration. The fin fold emerges between 20hpf and 48hpf (Figure 6a). At 20hpf it can be seen as a ridge of epidermal cells that sits on the

Figure 6. Dynamic expression of *krt97* and *krt5* during finfold development. (a) HCR in situ analysis of *krt97* and *krt5* between 20hpf and 48hpf. Yellow stars indicate non-specific fluorescence in the vasculature. Scale bar for 20hpf is 50um and 100um for 24hpf to 48hpf. **(b)** Expression of *krt97* and *krt5* in scRNA-seq cell clusters. The outline shows all cells in the dataset (3hpf to 120hpf) Cells in the 36-46hpf data set are shown in grey, yellow or red according to their level of expression. Clusters representing periderm (P), fin epithelium (FE) and Epithelium (E) are labelled. Data taken from Daniocell scRNA-seq dataset(Farrell et al. 2018; Sur et al. 2023).

midline of the left and right sides of the tail. At this stage *krt5* is in scattered cells throughout the tail epidermis and at 24hpf and 32hpf *krt5* is expressed throughout the tail epidermis. Between 36hpf and 48hpf the fin fold goes through morphogenesis and *krt5* expression is limited to the trunk of the tail. At 20hpf *krt97* expression is in a layer of cells at the base of the epidermal ridge in a crescent of cells around the end of the tail. This expression expands and as the finfold expands it becomes more restricted to the caudal end of the finfold. Intriguingly *krt97* is not expressed in the most distal layer of cells of the finfold. To summarise, *krt5* and *krt97* are expressed in different areas of the tail epidermis with *krt97* being expressed in the fin fold as it undergoes morphogenesis.

To further understand the compartmentalisation of *krt5* and *krt97* expression we analysed the Daniocell scRNA-seq dataset which contains cells from 3hpf to 120hpf(Farrell et al. 2018; Sur et al. 2023). We found that *krt97* and *krt5* show separate domains of expression primarily between 30hpf and 58hpf and at other time points the expression domains are not distinct (data not shown). Figure 6g shows pooled cells from 36-46hpf and *krt5* is expressed in a cluster annotated as periderm and *krt97* is in fin epithelium and epithelium clusters.

Treatment with the inhibitor SB431542 during finfold development does not confer a morphological phenotype suggesting that TGF-beta signalling does not strongly affect finfold development (data not shown). However, considering our result that most genes which are regulated by TGF-beta during regeneration also show some regulation in uninjured fish, we decided to see whether SB431542 affects *krt* gene expression during development. We found that treatment between 16.5hpf to 24hpf and 40hpf to 48hpf results in a strong reduction in *krt97* expression (Fig. 7). On the other hand, *krt5* did not show a change in expression. This suggests that the regulation of *krts* during development may be related to that seen during regeneration, at least for *krt97*.

Figure 7. Inhibition of TGF-beta signalling affects krt97 expression during development. (a) HCR in situ staining for *krt97* and *krt5* with and without SB43154. Orthogonal views are shown to the right and below each image. The planes used for the orthogonal views are indicated by the yellow lines. Scale bar is 20um. **(b)** Quantification of fluorescence in (a). Asterisks indicate significance as in Fig.2 resulting from the unpaired t test. For the 24hpf timepoint, embryos at the 15 somites stages (16.5 hpf) were treated with DMSO or 100 uM SB431542 continuously. For the 48 hpf stage, larvae at 40 hpf were treated with DMSO or 100 uM SB431542.

This compartmentalised expression of *krt5* and *krt97* during finfold development and regeneration raises the possibility that other members of *krt$_{up}$* and *krt$_{down}$* show segregated expression patterns during development. We have tested other keratins and find that there is limited overlap between *krt$_{up}$* and *krt$_{down}$* gene expression (Supplemental Figure 1 and data not shown). To see whether this is the case more broadly we pooled periderm, fin epithelium and epithelium clusters from the Daniocell scRNA-seq dataset and assessed the degree of co-expression between *krt$_{up}$* and *krt$_{down}$* using Pearson's correlation coefficient analysis (Fig. 8). This data shows that there is a strong degree of coexpression within each group. For example, *krt1-19e* expression is correlated to other members of *krt$_{down}$* especially in the epidermis with *krt5* and *krt91* showing the highest association with *krt1-19e*. *krt1-19e* shows a negative correlation to members of *krt$_{up}$* particularly with *krt18a.1*, *krt97* and *krt8* in the periderm. This

data shows the co-regulation of *krts* seen during regeneration and with manipulation of TGF-beta signalling is also found more broadly in epithelia during developmental stages.

Figure 8. Coexpression of *krts* in skin clusters. scRNA-seq data from the Daniocell analysis (Farrell et al. 2018; Sur et al. 2023) was subsetted based upon periderm, fin epithelium and epidermis identities using all available timepoints. Each of these subsets was then tested for correlation (using Pearson Correlation in Seurat) for the krt_{down} and krt_{up} genes. Red colour indicates a positive correlation and blue colour indicates a negative correlation.

The expression *krt97* during finfold regeneration and development correlates to regions that are undergoing morphogenesis. These epithelia need to engage in collective cell migration and are likely to alter their cytoskeleton to allow this to occur. On the other hand, *krt5* is reduced during finfold morphogenesis and regeneration suggesting that it is inhibitory to epithelial movement. To investigate this hypothesis, we decided to analyse epithelial cell movement during development and regeneration to see whether different epithelial regions have different mobility. Our predictions were that regions that express *krt97* (epithelia of the fin fold) move more quickly than regions that express *krt5* (epithelia on the trunk). To test cell movement during regeneration we used photo-conversion of a nuclear localised dendra transgene (Fig 9a, b, g and Supplemental Fig. 2). We found that cells in the proximal finfold move the furthest averaging approximately 70um per day relative to cells on the trunk. In unoperated animals there was very little movement. To assess cell movement during development, we made timelapse recordings of a nuclear-localised fluorescent reporter and measured movement relative to the underlying somites (Movie 1). As during regeneration we found that during development cells on the finfold move faster than cells on the trunk (by approximately 15um/hour) (Fig. 9c-f, h). These results show that epithelia move at dramatically different rates during regeneration and development, and that the differences in movement may correlate to differences in *krt* expression.

Figure 9. Collective migration of epithelial cells is faster in the finfold than on the trunk. **(a)** Cell nuclei were followed over a period of three days after tail excision using Dendra photoconversion (red nuclei). The asterisks mark the border of converted nuclei on the trunk and the arrows indicate the distance travelled by nuclei on the fin epithelia relative to the trunk. Scale bar = 50um **(b)** Unoperated control fish show very little movement over these four days. Scale bar = 100um **(c)** Overview image (z-projection) showing nuclei labelled red fluorescence *(actb2:h2b-mCherry)* at 24hpf. Scale bar = 100um. **(d)** Close-up of the box in (c) showing three time points during a time-lapse recording. The red arrowheads point to two finfold epithelium nuclei at the different time points. The yellow arrows indicate the distance travelled. **(e)** Sections showing somite boundary positions at three timepoints with arrows marking the same position. **(f)** Sections following epithelium nuclei on the trunk at three timepoints with arrows marking their positions. **(g)** Quantification of movement of finfold cells after excision as in panels (a) and (b). Lines indicate non-linear fit 95% confidence interval. Individual unpaired t test for each timepoint are < 0.0001. **(h)** Quantification of movement of finfold cells during development as in panels (c-f). Note that the midline somite boundary is used to standardise the epithelial cell movements. Asterisks indicate significance as in Fig.2 resulting from the unpaired t test.

Discussion

In this study we focus on how regenerative cells are recruited to the site of injury after tail excision. We present a model whereby HH signalling activates regenerative cell recruitment by transcriptionally activating *tgfb1a* expression in the notochord. The TGF-beta pathway then induces more expression of *tgfb1a* whilst inhibiting expression of *ihhb* in the notochord. After establishing that TGF-beta signalling acts downstream of the HH pathway, we then focus on the identification of genes regulated by TGF-beta during regeneration to investigate how cell recruitment takes place. These data reveal sets of genes that are important in the regeneration and indicate that TGF-beta plays a central role in this process. We propose that members of the *krt* gene family as potential effectors of regenerative TGF-beta signalling and that TGF-beta signalling enables collective cell migration by modulating the expression of different *krts*.

TGF-beta has been linked to wound healing by mammalian studies(Lichtman, Otero-Vinas, and Falanga 2016). The initial activation of the pathway during wounding is by serum protease activity which releases TGF-beta1 from platelets. The activated pathway then has many roles including keratinocyte recruitment. Our results suggest that during tail regeneration the role in keratinocyte recruitment is similar, but the initial activation of the TGF-beta pathway involves transcriptional activation of *tgfb1a*. Another similarity is that there is also some evidence for TGF-beta regulation of mammalian *Krts* during wounding. For example, one study has found that *Krt15* is repressed during wound healing and is also repressed by TGF-beta signalling suggesting that it would be in the *krt_{off}* synexpression group(Werner and Munz 2000). Furthermore, links between keratin and TGFbeta have been characterised for different tumour types and keratins are used as biomarkers for cancer prognosis(Li et al. 2024; Moll, Divo, and Langbein 2008).

Mammalian studies that primarily use scratch assays have suggested that Krt's usually have negative roles on cell movement, but can also increase migration (Sanghvi-Shah and Weber 2017; Chung, Rotty, and Coulombe 2013); (Beil et al. 2003; Yoon and Leube 2019). For example, Keratin 6 and 16 are activated after wounding and are thought to play a role in retraction of intermediate filaments from the cell periphery(Paladini et al. 1996) and cause a reduction in cell adhesion(Wawersik and Coulombe 2000); (Wang et al. 2018); (Wawersik and Coulombe 2000). In addition, Krts may affect cell migration by altering cellular elasticity(Beil et al. 2003) and myosin activation(Nanes et al. 2024). In Xenopus keratins are involved with adhesion and cell contraction during gastrulation(Sonavane et al. 2017). Understanding the precise molecular role of zebrafish Krts during collective epithelial cell migration will take further investigation.

Regulated expression of *krts* during regeneration has been observed in adult zebrafish tail regeneration(Martorana et al. 2001), xenopus larval tail regeneration(Tazawa, Shimizu-Nishikawa, and Yoshizato 2006) and newt limbs(Corcoran and Ferretti 1997). This suggests that dynamic expression of *krts* is conserved during organ regeneration. Our finding that *krts* have similar regional expression domains during development and regeneration may indicate that embryonic expression patterns are re-established during regeneration. The significance of regionalised expression domains of *krts* during development has yet to be explored.

Movie 1 Description. The movie shows the animal imaged in Figure 9c-f. Panel A shows the maximum intensity projection with yellow arrows following two finfold epithelium nuclei. Panel B shows sections through the somites with a somite boundary position marked with an arrow. Panel C shows superficial sections of the trunk epithelium with arrows following two nuclei.

Methods

General methods

Experimental procedures and fish maintenance were performed using standard methods49. All animal husbandry and experimentation was carried out under the supervision and approval of the Home Office (UK) and the University of Sheffield Ethics Board. Adult zebrafish were maintained with a 14 h light/10 h dark cycle at 28 °C according to standard protocols and were mated using pair mating in individual cross tanks. Fish were anesthetised in 40 µg/ml Tricaine (3-amino benzoic acidethylester) in E3. For more information on how individual experiments were performed, please refer to the figure legends and the sections below.

Bulk RNA-seq analysis

Zebrafish embryos were raised until 72hpf. A scalpel was used to remove the end of the tail from anaesthetised larvae using the pigment gap as a reference (Romero et al). The four conditions were: Unoperated DMSO, Operated DMSO, Unoperated SB431542 and Operated SB431542. There were 3 replicates for each condition and these each contained 100 tails pieces. Each replicate was processed in 1ml of TRI Reagent (Merck) using a plastic pestle to dissociate the tissue. Following the TRI Reagent procedure the RNA was further purified by LiCL precipitation. The RNA quality was checked by Qubit (ThermoFisher) and TapeStation (Agilent) protocols. The libraries were made using the Lexogen QuantSeq 3' FWD library kit and using NextSeq500 (Illumina) to generate approximately 5 million 75SE reads for each library. The resulting sequence data was demultiplexed using the BlueBee tool (Illumina), aligned to the ensemble zebrafish reference genome and analysed in DESeq2(Love, Huber, and Anders

2014). Plots were generated using Prism 10 (Graphpad). Data sets available at GEO Series GSE296469.

RNA in situ hybridization

Standard RNA in situ hybridisation was done using the Thisse protocol (Thisse and Thisse). HCR in situ hybridization was performed using zebrafish whole mount protocol(Choi, Schwarzkopf, and Pierce 2020) and reagents purchased from Molecular Instruments (Los Angeles, California). To report in situ expression patterns, a representative animal is shown in the figure panel. For quantification, graphs were generated and analysed in Prism 10 software. Error bars indicate the 95% confidence interval and the centre bar represents the mean. Individual circles represent individual animals tested. The figure legends indicate the type of statistical test applied.

PhotoConversion

ubi:h2b-dendra(Gurskaya et al. 2006) transgenic fish were used for photoconversion experiments. Larvae at 48hpf were placed in a holding chamber in a glass bottom dish. Rectangular ROI was used to draw the region on the tail and photoconverted exposing to 405nm with 100% laser power for 2 min. Unoperated and Operated fish were monitored over a period of time, and images were captured at 543nm and 488nm wavelength using Nikon A1 confocal microscope. For quantification cells at the centre of the trunk were used as a reference point, and the maximum distance travelled by fin fold cells was measured. The analysis was done on six operated animals and three unoperated animals.

Time-lapse

actb2:h2b-mCherry transgenic fish were used for these experiments. Zebrafish Larvae were anaesthetised and imaged by light sheet microscopy (Bruker TrueLive3D) for 280 minutes. For quantification, three fish were analysed, (two started at 22hpf and one at 24hpf). Three epithelial cell nuclei were measured in the proximal fin fold and three epithelial cell nuclei in the centre of the trunk. The boundary between two somites was used as a reference point to standardise movement of the epithelial cells.

scRNA-seq analysis

Data sets GSE145497 and GSE223922 were downloaded from the NCBI/Geo server. Analysis was done using Seurat version 5.1.0, R (version 4.4.1 2024-06-14) and R Studio (version 2024.04.2+764). A complete list of installed packages and versions is available in Supplemental Table 6.

Acknowledgements

We would like to thank: Gideon Hughes for technical assistance during collection of the tissue for RNA-seq analysis and for providing the *ubi:h2b-dendra* line; Dimitris Rovanias for initial experiments with TGF-beta expression. Fish lines were maintained by the University of Sheffield Biological Services Aquarium under Home Office project licence 3627554. Imaging services were provided by the Wolfson Light Microscopy Facility at the University of Sheffield (Biotechnology and Biological Sciences Research Council Grant BB/W019450/1). Funding provided to HHR by the Biotechnology and Biological Sciences Research Council (Grant BB/S015086/1) and the Leverhulme Trust (Grant RPG-2021-073). For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising.

Author Contributions Statement

Experiments were conceived and designed by T.J.C. and H.H.R. and were performed and analysed by D.D.M., Y.L., M.W. H.H.R. and C.Z. The manuscript was written by T.J.C and H.R. All authors reviewed the manuscript.

References

Beil, Michael, Alexandre Micoulet, Götz von Wichert, Stephan Paschke, Paul Walther, M. Bishr Omary, Paul P. Van Veldhoven, et al. 2003. "Sphingosylphosphorylcholine Regulates Keratin Network Architecture and Visco-Elastic Properties of Human Cancer Cells." *Nature Cell Biology* 5 (9): 803–11.

Bensimon-Brito, Anabela, Srinath Ramkumar, Giulia L. M. Boezio, Stefan Guenther, Carsten Kuenne, Christian S. M. Helker, Héctor Sánchez-Iranzo, et al. 2020. "TGF-β Signaling Promotes Tissue Formation during Cardiac Valve Regeneration in Adult Zebrafish." *Developmental Cell* 52 (1): 9–20.e7.

Choi, Harry M. T., Maayan Schwarzkopf, and Niles A. Pierce. 2020. "Multiplexed Quantitative In Situ Hybridization with Subcellular or Single-Molecule Resolution Within Whole-Mount Vertebrate Embryos: qHCR and dHCR Imaging (v3.0)." *Methods in Molecular Biology (Clifton, N.J.)* 2148:159–78.

Chung, Byung-Min, Jeremy D. Rotty, and Pierre A. Coulombe. 2013. "Networking Galore: Intermediate Filaments and Cell Migration." *Current Opinion in Cell Biology* 25 (5): 600–612.

Corcoran, J. P., and P. Ferretti. 1997. "Keratin 8 and 18 Expression in Mesenchymal Progenitor Cells of Regenerating Limbs Is Associated with Cell Proliferation and Differentiation." *Developmental Dynamics : An Official Publication of the American Association of Anatomists* 210 (4): 355–70.

Dereeper, A., V. Guignon, G. Blanc, S. Audic, S. Buffet, F. Chevenet, J-F Dufayard, et al. 2008. "Phylogeny.fr: Robust Phylogenetic Analysis for the Non-Specialist." *Nucleic Acids Research* 36 (Web Server issue): W465–69.

Farrell, Jeffrey A., Yiqun Wang, Samantha J. Riesenfeld, Karthik Shekhar, Aviv Regev, and Alexander F. Schier. 2018. "Single-Cell Reconstruction of Developmental Trajectories during Zebrafish Embryogenesis." *Science (New York, N.Y.)* 360 (6392). https://doi.org/10.1126/science.aar3131.

Gong, Zhiyuan, Bensheng Ju, Xukun Wang, Jiangyan He, Haiyan Wan, Putter Mudumana Sudha, and Tie Yan. 2002. "Green Fluorescent Protein Expression in

Germ-Line Transmitted Transgenic Zebrafish under a Stratified Epithelial Promoter from keratin8." *Developmental Dynamics : An Official Publication of the American Association of Anatomists* 223 (2): 204–15.

Gurskaya, Nadya G., Vladislav V. Verkhusha, Alexander S. Shcheglov, Dmitry B. Staroverov, Tatyana V. Chepurnykh, Arkady F. Fradkov, Sergey Lukyanov, and Konstantin A. Lukyanov. 2006. "Engineering of a Monomeric Green-to-Red Photoactivatable Fluorescent Protein Induced by Blue Light." *Nature Biotechnology* 24 (4): 461–65.

Ho, Minh, Brian Thompson, Jeffrey Nicholas Fisk, Daniel W. Nebert, Elspeth A. Bruford, Vasilis Vasiliou, and Christopher G. Bunick. 2022. "Update of the Keratin Gene Family: Evolution, Tissue-Specific Expression Patterns, and Relevance to Clinical Disorders." *Human Genomics* 16 (1): 1.

Hsu, Wei-Lin, Yu-Chi Lin, Meng-Ju Lin, Yi-Wen Wang, and Shyh-Jye Lee. 2024. "Macrophages Enhance Regeneration of Lateral Line Neuromast Derived from Interneuromast Cells through TGF-β in Zebrafish." *Development, Growth & Differentiation* 66 (2): 133–44.

Keatinge, Marcus, Themistoklis M. Tsarouchas, Tahimina Munir, Nicola J. Porter, Juan Larraz, Davide Gianni, Hui-Hsin Tsai, Catherina G. Becker, David A. Lyons, and Thomas Becker. 2021. "CRISPR gRNA Phenotypic Screening in Zebrafish Reveals pro-Regenerative Genes in Spinal Cord Injury." *PLoS Genetics* 17 (4): e1009515.

Krushna Padhi, Bhaja, Marie-Andrée Akimenko, and Marc Ekker. 2006. "Independent Expansion of the Keratin Gene Family in Teleostean Fish and Mammals: An Insight from Phylogenetic Analysis and Radiation Hybrid Mapping of Keratin Genes in Zebrafish." *Gene* 368 (March):37–45.

Lee, Raymond T. H., P. V. Asharani, and Thomas J. Carney. 2014. "Basal Keratinocytes Contribute to All Strata of the Adult Zebrafish Epidermis." *PLoS One* 9 (1): e84858.

Lichtman, Michael K., Marta Otero-Vinas, and Vincent Falanga. 2016. "Transforming Growth Factor Beta (TGF-β) Isoforms in Wound Healing and Fibrosis." *Wound Repair and Regeneration : Official Publication of the Wound Healing Society [and] the European Tissue Repair Society* 24 (2): 215–22.

Li, Gang, Jinbao Guo, Yunfei Mou, Qingsong Luo, Xuehai Wang, Wei Xue, Ting Hou, Tianyang Zeng, and Yi Yang. 2024. "Keratin Gene Signature Expression Drives Epithelial-Mesenchymal Transition through Enhanced TGF-β Signaling Pathway Activation and Correlates with Adverse Prognosis in Lung Adenocarcinoma." *Heliyon* 10 (3): e24549.

Love, Michael I., Wolfgang Huber, and Simon Anders. 2014. "Moderated Estimation of Fold Change and Dispersion for RNA-Seq Data with DESeq2." *Genome Biology* 15 (12): 550.

Martorana, M. L., M. Tawk, T. Lapointe, N. Barre, M. Imboden, C. Joulie, J. Géraudie, and S. Vriz. 2001. "Zebrafish Keratin 8 Is Expressed at High Levels in the Epidermis of Regenerating Caudal Fin." *The International Journal of Developmental Biology* 45 (2): 449–52.

Massagué, Joan, and Dean Sheppard. 2023. "TGF-β Signaling in Health and Disease." *Cell* 186 (19): 4007–37.

Moll, Roland, Markus Divo, and Lutz Langbein. 2008. "The Human Keratins: Biology and Pathology." *Histochemistry and Cell Biology* 129 (6): 705–33.

Nanes, Benjamin A., Kushal Bhatt, Evgenia Azarova, Divya Rajendran, Sabahat Munawar, Tadamoto Isogai, Kevin M. Dean, and Gaudenz Danuser. 2024. "Shifts in Keratin Isoform Expression Activate Motility Signals during Wound Healing." *Developmental Cell* 59 (20): 2759–71.e11.

Niethammer, Philipp. 2016. "The Early Wound Signals." *Current Opinion in Genetics &*

Development 40 (October):17–22.

Paladini, R. D., K. Takahashi, N. S. Bravo, and P. A. Coulombe. 1996. "Onset of Re-Epithelialization after Skin Injury Correlates with a Reorganization of Keratin Filaments in Wound Edge Keratinocytes: Defining a Potential Role for Keratin 16." *The Journal of Cell Biology* 132 (3): 381–97.

Poss, Kenneth D., and Elly M. Tanaka. 2024. "Hallmarks of Regeneration." *Cell Stem Cell* 31 (9): 1244–61.

Redmond, Catherine J., and Pierre A. Coulombe. 2021. "Intermediate Filaments as Effectors of Differentiation." *Current Opinion in Cell Biology* 68 (February):155–62.

Richardson, Rebecca, Manuel Metzger, Philipp Knyphausen, Thomas Ramezani, Krasimir Slanchev, Christopher Kraus, Elmon Schmelzer, and Matthias Hammerschmidt. 2016. "Re-Epithelialization of Cutaneous Wounds in Adult Zebrafish Combines Mechanisms of Wound Closure in Embryonic and Adult Mammals." *Development (Cambridge, England)* 143 (12): 2077–88.

Rojas-Muñoz, Agustin, Shibani Rajadhyksha, Darren Gilmour, Frauke van Bebber, Christopher Antos, Concepción Rodríguez Esteban, Christiane Nüsslein-Volhard, and Juan Carlos Izpisúa Belmonte. 2009. "ErbB2 and ErbB3 Regulate Amputation-Induced Proliferation and Migration during Vertebrate Regeneration." *Developmental Biology* 327 (1): 177–90.

Romero, Maria Montserrat Garcia, Gareth McCathie, Philip Jankun, and Henry Hamilton Roehl. 2018. "Damage-Induced Reactive Oxygen Species Enable Zebrafish Tail Regeneration by Repositioning of Hedgehog Expressing Cells." *Nature Communications* 9 (1): 4010.

Sanghvi-Shah, Rucha, and Gregory F. Weber. 2017. "Intermediate Filaments at the Junction of Mechanotransduction, Migration, and Development." *Frontiers in Cell and Developmental Biology* 5 (September):81.

Sinclair, Jason W., David R. Hoying, Erica Bresciani, Damian Dalle Nogare, Carli D. Needle, Alexandra Berger, Weiwei Wu, et al. 2021. "The Warburg Effect Is Necessary to Promote Glycosylation in the Blastema during Zebrafish Tail Regeneration." *NPJ Regenerative Medicine* 6 (1): 55.

Sonavane, Pooja R., Chong Wang, Bette Dzamba, Gregory F. Weber, Ammasi Periasamy, and Douglas W. DeSimone. 2017. "Mechanical and Signaling Roles for Keratin Intermediate Filaments in the Assembly and Morphogenesis of Mesendoderm Tissue at Gastrulation." *Development (Cambridge, England)* 144 (23): 4363–76.

Sur, Abhinav, Yiqun Wang, Paulina Capar, Gennady Margolin, Morgan Kathleen Prochaska, and Jeffrey A. Farrell. 2023. "Single-Cell Analysis of Shared Signatures and Transcriptional Diversity during Zebrafish Development." *Developmental Cell* 58 (24): 3028–47.e12.

Tazawa, Ichiro, Keiko Shimizu-Nishikawa, and Katsutoshi Yoshizato. 2006. "A Novel Xenopus Laevis Larval Keratin Gene, xlk2: Its Gene Structure and Expression during Regeneration and Metamorphosis of Limb and Tail." *Biochimica et Biophysica Acta* 1759 (5): 216–24.

Thisse, Bernard, and Christine Thisse. 2014. "In Situ Hybridization on Whole-Mount Zebrafish Embryos and Young Larvae." *Methods in Molecular Biology (Clifton, N.J.)* 1211:53–67.

Wang, Fengrong, Song Chen, Hans B. Liu, Carole A. Parent, and Pierre A. Coulombe. 2018. "Keratin 6 Regulates Collective Keratinocyte Migration by Altering Cell-Cell and Cell-Matrix Adhesion." *The Journal of Cell Biology* 217 (12): 4314–30.

Wawersik, M., and P. A. Coulombe. 2000. "Forced Expression of Keratin 16 Alters the Adhesion, Differentiation, and Migration of Mouse Skin Keratinocytes." *Molecular*

Biology of the Cell 11 (10): 3315–27.

Werner, S., and B. Munz. 2000. "Suppression of Keratin 15 Expression by Transforming Growth Factor Beta in Vitro and by Cutaneous Injury in Vivo." *Experimental Cell Research* 254 (1): 80–90.

Yoon, Sungjun, and Rudolf E. Leube. 2019. "Keratin Intermediate Filaments: Intermediaries of Epithelial Cell Migration." *Essays in Biochemistry* 63 (5): 521–33.

Supplemental Data

Supplemental Figures 1-2

Supplemental Tables 1-6

Supplemental Figure 1: *krt1-19e* is expressed in the trunk region and does not overlap with *krt97* at 48hpf. (a-c) Low magnification and (d-f) high magnification images of *krt1-19e* expression is shown in green, DAPI staining of nuclei in blue and *krt97* in red. *krt1-19e* is expressed in the trunk and does not overlap with *krt97* at this stage.

Supplemental Figure 2: Examples of photoconversion experiments during regeneration.

Gene Symbol	Gene Title	ensemble	operated vs unoperated DMSO	operated vs unoperated SB43	SB43 vs DMSO operated	SB43 vs DMSO unoperated
lama3	laminin subunit alpha-3	ENSDARG00000022615	7.41	1.23	0.18	1.11
manf	mesencephalic astrocyte-derived neurotrophic	ENSDARG00000063177	3.22	1.87	0.59	1.02
tram1	translocation associated membrane protein 1	ENSDARG00000019137	2.99	1.36	0.49	1.08
efemp2a	EGF containing fibulin-like extracellular matrix	ENSDARG00000094324	2.92	1.98	0.61	0.90
krt96	keratin 96	ENSDARG00000095147	2.86	0.59	0.19	0.92
si:dkey	protein coding gene	ENSDARG00000090552	2.83	1.82	0.67	1.05
hpcal4	hippocalcin like 4	ENSDARG00000070491	2.75	0.85	0.33	1.09
slc39a7	solute carrier family 39 (zinc transporter),	ENSDARG00000104451	2.46	1.53	0.59	0.95
si:dkey	protein coding gene	ENSDARG00000096678	2.13	0.77	0.39	1.07
BX511082.1	protein coding gene	ENSDARG00000113678	1.82	0.99	0.52	0.97
serpinh1a	serpin peptidase inhibitor, clade H (heat shock	ENSDARG00000075954	1.79	1.00	0.51	0.91
arhgdig	Rho GDP dissociation inhibitor (GDI) gamma	ENSDARG00000004034	1.75	1.17	0.72	1.07
cpm	carboxypeptidase M	ENSDARG00000011769	1.66	1.01	0.56	0.91
sox4b	SRY (sex determining region Y)-box 4b	ENSDARG00000098834	1.56	0.82	0.51	0.96
sec23a	Sec23 homolog A, COPII coat complex	ENSDARG00000104230	1.56	1.09	0.65	0.94
virma	si:ch211-79I20.4	ENSDARG00000075824	1.50	0.92	0.57	0.94
dlx5a	distal-less homeobox 5a	ENSDARG00000042296	1.47	0.81	0.53	0.96
bhlha9	basic helix-loop-helix family, member a9	ENSDARG00000103981	1.47	0.77	0.50	0.95
psat1	phosphoserine aminotransferase 1	ENSDARG00000016733	1.41	0.75	0.50	0.94
cct5	chaperonin containing TCP1, subunit 5	ENSDARG00000045399	1.40	0.97	0.63	0.91
acot7		ENSDARG00000018653	1.39	0.77	0.51	0.92
psmd6	proteasome 26S subunit, non-ATPase 6	ENSDARG00000070674	1.37	0.94	0.62	0.90
fstl1b	follistatin-like 1b	ENSDARG00000039576	1.28	0.87	0.66	0.97
ppp1r3b	protein phosphatase 1, regulatory subunit 3B	ENSDARG00000044691	1.27	0.57	0.44	0.98
gps1	G protein pathway suppressor 1	ENSDARG00000040650	1.26	0.98	0.73	0.94
ccdc6a	coiled-coil domain containing 6a	ENSDARG00000043334	1.21	0.87	0.70	0.97
zgc:92744	zgc:92744	ENSDARG00000076223	1.06	0.76	0.69	0.96
immt	inner membrane protein, mitochondrial	ENSDARG00000102874	1.03	0.79	0.69	0.90
ACADSB		ENSDARG00000098174	1.03	0.71	0.61	0.88
btr12	bloodthirsty-related gene family, member 12	ENSDARG00000051809	0.97	3.43	3.88	1.10
mfn2	mitofusin 2	ENSDARG00000079504	0.94	0.53	0.63	1.11
tmem131	transmembrane protein 131	ENSDARG00000056259	0.93	1.40	1.53	1.02
got1	glutamic-oxaloacetic transaminase 1, soluble	ENSDARG00000039093	0.85	1.46	1.58	0.93
galnt6	UDP-N-acetyl-alpha-D-galactosamine:	ENSDARG00000014386	0.73	2.05	2.51	0.90
rfx2	regulatory factor X, 2 (influences HLA class II	ENSDARG00000013575	0.72	1.33	1.91	1.03
pmp22a	peripheral myelin protein 22a	ENSDARG00000105223	0.58	0.88	1.57	1.03
matn3b		ENSDARG00000069265	0.57	0.39	0.66	0.97
evpla	envoplakin a	ENSDARG00000019808	0.53	1.04	1.91	0.97
fh	fumarate hydratase	ENSDARG00000075132	0.52	0.84	1.63	1.01
lipg	lipase, endothelial	ENSDARG00000031044	0.42	0.74	1.89	1.07

Fold change
>5.0 = red
>3.0 = light red
>1.2 = pale red
<0.85 = pale blue
<0.65 = light blue
<0.5 = blue
p<0.05 p<0.1

Gene counts
>1
>10
>100
>1000

Count columns: DMSO unoperated COUNT, DMSO operated COUNT, SB43 unoperated COUNT, SB43 operated COUNT

Supplemental Table 1. Potential injury-specific TGF-beta target DEGs. DEGs with SB431542 operated versus DMSO operated (with P_{adj} value < 0.05) and a lack of differential expression in SB431542 unoperated versus DMSO unoperated are presented in this table. The columns show the ensemble identifiers, the fold level of change and the adjusted counts for each sample. These are the DEGs that lie in the boxed area of the graph (as in Fig 3c) and have a Log_2 score of between 0.2 and -0.2 in SB431542 unoperated versus DMSO unoperated comparison (i.e. genes that are between 0.87 and 1.15 fold change). Note that the graph is in Log_2 scale and the table shows fold change.

Fold change
- >5.0 = red
- >3.0 = light red
- >1.2 = pale red
- <0.85 = pale blue
- <0.65 = light blue
- <0.5 = blue

Gene counts
- >1
- >10
- >100
- >1000

■ p<0.05   p<0.1

Gene Symbol	Gene Title	ensemble	operated vs unoperated DMSO	operated vs unoperated SB43	SB43 vs DMSO operated	SB43 vs DMSO unoperated	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
mmp9	matrix metallopeptidase 9	ENSDARG00000042816	10.35	7.44	2.42	3.37				
lama3	laminin subunit alpha-3	ENSDARG00000022615	7.41	1.23	0.18	1.11				
ctss2.1	cathepsin Sb, tandem duplicate 1	ENSDARG00000074656	6.91	6.61	2.06	2.15				
f13a1b	coagulation factor XIII, A1 polypeptide b	ENSDARG00000036893	6.28	1.73	0.41	1.49				
sp6	*Sp6 transcription factor*	ENSDARG00000099880	5.89	1.95	0.22	0.67				
si:ch211	Protein coding	ENSDARG00000060927	5.61	2.53	0.34	0.75				
lcp1	*lymphocyte cytosolic protein 1 (L-plastin)*	ENSDARG00000023188	4.46	3.25	0.48	0.66				
f5	coagulation factor V	ENSDARG00000055705	4.40	2.30	0.41	0.79				
apoeb	apolipoprotein Eb	ENSDARG00000040295	4.15	1.79	0.56	1.26				
c1qtnf5	C1q and tumor necrosis factor related protein 5	ENSDARG00000056134	3.88	2.42	0.40	0.64				
snorc	secondary ossification center associated	ENSDARG00000092383	3.71	1.17	0.19	0.62				
txn	thioredoxin	ENSDARG00000044125	3.36	4.22	1.81	1.44				
manf	mesencephalic astrocyte-derived neurotrophic	ENSDARG00000063177	3.22	1.87	0.59	1.02				
loxl5b	*lysyl oxidase-like 5b*	ENSDARG00000076904	3.16	1.33	0.26	0.63				
tram1	translocation associated membrane protein 1	ENSDARG00000019137	2.99	1.36	0.49	1.08				
efemp2a	EGF containing fibulin-like extracellular matrix	ENSDARG00000094324	2.92	1.98	0.61	0.90				
cmpk	cytidylate kinase	ENSDARG00000019924	2.90	2.11	0.52	0.72				
si:dkey	protein coding gene	ENSDARG00000090552	2.83	1.82	0.67	1.05				
hpcal4	hippocalcin like 4	ENSDARG00000070491	2.75	0.85	0.33	1.09				
krt97	keratin 97	ENSDARG00000000212	2.66	1.08	0.30	0.74				
slc39a7	solute carrier family 39 (zinc transporter),	ENSDARG00000104451	2.46	1.53	0.59	0.95				
mcm10	minichromosome maintenance 10 replication	ENSDARG00000045815	2.38	1.71	0.39	0.54				
s100a10b	S100 calcium binding protein A10b	ENSDARG00000025254	2.08	2.58	1.59	1.28				
prps1b	phosphoribosyl pyrophosphate synthetase 1B	ENSDARG00000037506	1.97	1.35	0.56	0.82				
rpa3	replication protein A3	ENSDARG00000002613	1.86	1.47	0.60	0.76				
ywhaqb	tyrosine 3-monooxygenase/tryptophan 5-	ENSDARG00000023323	1.85	2.27	1.54	1.26				
BX511082.1	protein coding gene	ENSDARG00000113678	1.82	0.99	0.52	0.97				
arhgdig	Rho GDP dissociation inhibitor (GDI) gamma	ENSDARG00000004034	1.75	1.17	0.72	1.07				
ap1m1		ENSDARG00000096249	1.67	1.21	0.63	0.86				
cpm	carboxypeptidase M	ENSDARG00000011769	1.66	1.01	0.56	0.91				
ccn1l2		ENSDARG00000099985	1.65	1.15	0.60	0.86				
clta	clathrin, light chain A	ENSDARG00000045618	1.64	1.27	0.56	0.73				
arpp19b	cAMP-regulated phosphoprotein 19b /// cAMP-	ENSDARG00000039880	1.59	1.16	0.58	0.80				
pqbp1	polyglutamine binding protein 1	ENSDARG00000029724	1.58	1.20	0.54	0.72				
nars1	asparaginyl-tRNA synthetase	ENSDARG00000061100	1.57	1.06	0.57	0.84				
sec23a	Sec23 homolog A, COPII coat complex	ENSDARG00000104230	1.56	1.09	0.65	0.94				
dnajc9	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000031293	1.56	1.33	0.62	0.73				
cd81a	CD81 molecule a	ENSDARG00000036080	1.40	1.17	0.71	0.85				
abcb5	ATP-binding cassette, sub-family B	ENSDARG00000021787	0.61	0.84	1.84	1.33				
pmp22a	peripheral myelin protein 22a	ENSDARG00000105223	0.58	0.88	1.57	1.03				
sdhaf4	succinate dehydrogenase complex assembly	ENSDARG00000039390	0.58	0.94	1.94	1.19				
PPM1K		ENSDARG00000076011	0.58	0.66	1.52	1.33				
matn3b		ENSDARG00000069265	0.57	0.39	0.66	0.97				
tmem41ab	transmembrane protein 41ab	ENSDARG00000026771	0.53	0.67	1.70	1.35				
evpla	envoplakin a	ENSDARG00000019808	0.53	1.04	1.91	0.97				
fh	fumarate hydratase	ENSDARG00000075132	0.52	0.84	1.63	1.01				
ucp1	uncoupling protein 1	ENSDARG00000023151	0.49	0.77	1.85	1.17				
si:dkey	protein coding gene	ENSDARG00000101128	0.48	0.66	2.33	1.70				
nmrk2	nicotinamide riboside kinase 2	ENSDARG00000067848	0.45	0.61	1.96	1.43				
lipg	lipase, endothelial	ENSDARG00000031044	0.42	0.74	1.89	1.07				
ankrd9	ankyrin repeat domain 9	ENSDARG00000028804	0.36	0.53	4.21	2.86				
col7a1	collagen, type VII, alpha 1	ENSDARG00000021720	0.21	0.60	2.36	0.84				

Supplemental Table 2. Potential TGF-beta dependent DEGs. These are DEGs that have a P_{adj} value < 0.05 for both DMSO operated versus DMSO unoperated and SB431542 operated versus DMSO operated. The columns show the ensemble identifiers, the fold level of change and the adjusted counts for each sample. The graph is the same as in Fig. 3d but only showings DEGs that are significant for both comparisons. Note that the graph is in Log_2 scale and the table shows fold change.

Fold change
- ■ >5.0 = red
- ■ >3.0 = light red
- ■ >1.2 = pale red
- ■ <0.85 = pale blue
- ■ <0.65 = light blue
- ■ <0.5 = blue
- ■ p<0.05   ■ p<0.1

Gene counts
- ■ >1
- ■ >10
- ■ >100
- ■ >1000

Gene Symbol	Gene Title	ensemble	operated vs unoperated DMSO	operated vs unoperated SB43	SB43 vs DMSO operated	SB43 vs DMSO unoperated	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
fn1b	fibronectin 1b	ENSDARG00000006526	32.47	21.34	0.67	1.02				
cart3		ENSDARG00000035852	18.77	12.90	0.71	1.03				
timp2b	TIMP metallopeptidase inhibitor 2b	ENSDARG00000075261	16.15	13.34	0.74	0.90				
grn1	granulin 1	ENSDARG00000089362	16.13	10.23	0.80	1.26				
si:ch211		ENSDARG00000109648	9.76	5.54	0.75	1.32				
ms4a17a.4	membrane-spanning 4-domains	ENSDARG00000014024	9.71	6.57	0.70	1.04				
slc7a7	solute carrier family 7 (amino acid transporter	ENSDARG00000055226	7.70	5.15	0.67	1.00				
tgm2l	transglutaminase 2, like	ENSDARG00000093381	7.27	2.70	0.78	2.09				
adam8a		ENSDARG00000001452	6.52	4.10	0.83	1.32				
clu	clusterin	ENSDARG00000010434	6.43	5.29	0.75	0.91				
ms4a17a.5	membrane-spanning 4-domains	ENSDARG00000092204	6.24	3.21	0.67	1.30				
cbx7a		ENSDARG00000038025	6.16	3.66	0.69	1.16				
c7a		ENSDARG00000042172	5.95	4.73	0.60	0.75				
ms4a17a.7	membrane-spanning 4-domains	ENSDARG00000043796	5.36	2.64	0.81	1.65				
krt18a.1	keratin 18	ENSDARG00000018042	5.05	3.20	0.75	1.18				
FO704661.1		ENSDARG00000102758	4.98	3.40	0.81	1.18				
ctsk	cathepsin K	ENSDARG00000040251	4.58	2.58	0.82	1.46				
lcp1	lymphocyte cytosolic protein 1 (L-plastin)	ENSDARG00000023188	4.46	3.25	0.48	0.66				
mpeg1.1		ENSDARG00000055290	4.46	2.80	0.70	1.11				
kif23	kinesin family member 23	ENSDARG00000014943	4.37	2.50	0.71	1.24				
snap23.2		ENSDARG00000055252	4.35	3.77	0.70	0.81				
apoeb	apolipoprotein Eb	ENSDARG00000040295	4.15	1.79	0.56	1.26				
si:rp71	dispanin subfamily A member 2b-like	ENSDARG00000097746	4.09	2.90	0.75	1.06				
ccdc88b		ENSDARG00000076189	3.98	3.15	0.86	1.08				
tagln	transgelin	ENSDARG00000045408	3.96	2.87	0.66	0.92				
tnnt2c	troponin T2c, cardiac	ENSDARG00000032242	3.90	2.82	0.73	1.01				
c1qtnf5	C1q and tumor necrosis factor related protein 5	ENSDARG00000056134	3.88	2.42	0.40	0.64				
fcer1gl		ENSDARG00000104077	3.68	2.33	0.68	1.07				
spi1b	Spi-1 proto-oncogene b	ENSDARG00000000767	3.65	2.49	0.72	1.06				
manf	mesencephalic astrocyte-derived neurotrophic	ENSDARG00000063177	3.22	1.87	0.59	1.02				
si:ch211	si:ch211-1a19.3	ENSDARG00000100968	3.10	2.39	0.76	0.99				
cdca8	cell division cycle associated 8	ENSDARG00000043137	3.00	2.18	0.73	1.01				
efemp2a	EGF containing fibulin-like extracellular matrix	ENSDARG00000094324	2.92	1.98	0.61	0.90				
cmpk	cytidylate kinase	ENSDARG00000019924	2.90	2.11	0.52	0.72				
si:dkey	protein coding gene	ENSDARG00000090552	2.83	1.82	0.67	1.05				
hyou1	hypoxia up-regulated 1	ENSDARG00000013670	2.78	1.84	0.83	1.26				
ms4a17a.12	membrane-spanning 4-domains	ENSDARG00000053563	2.74	2.27	0.86	1.04				
clic2		ENSDARG00000010625	2.62	1.92	0.82	1.12				
pitpnaa	phosphatidylinositol transfer protein, alpha a	ENSDARG00000039490	2.61	2.26	0.76	0.88				
il6st	interleukin 6 signal transducer ///	ENSDARG00000104693	2.47	2.04	0.77	0.93				
capgb	capping protein (actin filament), gelsolin-like b	ENSDARG00000099672	2.41	1.80	0.83	1.12				
ptges3b	prostaglandin E synthase 3b (cytosolic)	ENSDARG00000089626	2.21	1.78	0.85	1.06				
rrbp1a	ribosome binding protein 1 homolog a (dog)	ENSDARG00000013763	2.19	1.72	0.73	0.94				
dad1	defender against cell death 1	ENSDARG00000102105	2.01	1.54	0.69	0.90				
thop1	zgc:92139	ENSDARG00000013776	1.96	1.66	0.79	0.93				
tmsb4x	thymosin, beta 4 x	ENSDARG00000077777	1.72	1.47	0.83	0.96				
atp2a1l	ATPase, Ca++ transporting, cardiac muscle,	ENSDARG00000035458	0.52	0.69	1.36	1.03				
cs	citrate synthase	ENSDARG00000103364	0.50	0.62	1.15	0.93				
krt4	keratin 4	ENSDARG00000017624	0.49	0.66	1.33	0.99				
casq1a		ENSDARG00000038716	0.48	0.57	1.48	1.24				
dnajc18	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000056005	0.47	0.58	1.19	0.97				
si:ch211		ENSDARG00000096616	0.46	0.62	1.33	0.98				
tuba8l2	tubulin, alpha 8 like 2	ENSDARG00000031164	0.45	0.56	1.17	0.94				
fndc1		ENSDARG00000002847	0.40	0.50	1.28	1.03				
nrap		ENSDARG00000009341	0.35	0.41	1.27	1.08				
col2a1a	collagen, type II, alpha 1a	ENSDARG00000096093	0.32	0.37	1.77	1.51				
col11a2	collagen, type XI, alpha 2	ENSDARG00000012422	0.30	0.38	1.16	0.92				
epyc		ENSDARG00000056950	0.27	0.42	1.21	0.77				

Supplemental Table 3. Potential regeneration DEGs. To generate this table DEGs must meet all of the following criteria (1) a gene count of >10 for all conditions (2) a P_{adj} value < 0.05 for DMSO operated versus unoperated and for SB431542 operated versus unoperated. (3) Either a $\log_2$ score above 0 for SB431542 operated versus unoperated and a ratio below 0.87 for SB431542 operated versus unoperated/ DMSO operated versus unoperated (coloured DEGs in the top right quadrant of graph) OR a $\log_2$ score below 0 for SB431542 operated versus unoperated and a ratio above 1.15 for SB431542 operated versus unoperated/ DMSO operated versus unoperated (coloured DEGs in the lower left quadrant of graph). The columns show the ensemble identifiers, the fold level of change and the adjusted counts for each sample. Note that the graph is the same data as in Fig 3e with the line indicating a slope of one. The graph is in $\log_2$ scale and the table shows fold change.

graph 1

graph 2

Supplemental Table 4. Potential regeneration DEGs that are not strongly regulated by TGF-beta. To generate this table DEGs must meet all of the following criteria (1) a gene count of >10 for all conditions (2) a P_{adj} value < 0.05 for DMSO operated versus DMSO unoperated and a Log_2 score of between 0.2 and -0.2 in the SB431542 operated versus DMSO operated comparison (box in graph 1) (3) a P_{adj} value < 0.05 for SB431542 operated versus unoperated (4) a similar expression change between DMSO operated versus DMSO unoperated and SB431542 operated versus SB431542 unoperated (ratio between 0.87 and 1.15) (box in graph 2). The columns show the ensemble identifiers, the fold level of change and the adjusted counts for each sample. Note that the graph is in Log_2 scale and the table shows fold change. Graph 1 is data from Figure 3d and graph 2 is Figure 3e.

Supplemental Table 5. Comparison of families of genes within RNA-seq data. This table shows families of genes with five or more members that were detected in our RNA-seq experiments. The columns show the ensemble identifiers, the fold level of change and the adjusted counts for each sample.

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
abcb4	ATP-binding cassette, sub-family B	ENSDARG00000010936	7.14	0.47	1.15	2.92				
abcc10	ATP-binding cassette, sub-family C	ENSDARG00000077988	2.16	1.90	2.18	1.88				
abcf2a	ATP-binding cassette, sub-family F (GCN20),	ENSDARG00000038785	1.73	0.93	1.10	1.47				
abca1a	ATP-binding cassette, sub-family A (ABC1),	ENSDARG00000074635	1.52	0.85	1.15	1.12				
abce1	ATP-binding cassette, sub-family E (OABP),	ENSDARG00000007216	1.48	0.69	0.85	1.20				
abca1b	ATP-binding cassette, sub-family A (ABC1),	ENSDARG00000079009	1.27	0.90	0.66	1.73				
abcf1	ATP-binding cassette, sub-family F (GCN20),	ENSDARG00000031795	1.20	0.82	0.73	1.34				
abca5	ATP-binding cassette, sub-family A (ABC1),	ENSDARG00000074041	1.03	1.03	0.78	1.35				
abca3b	ATP-binding cassette, sub-family A (ABC1),	ENSDARG00000100524	1.02	1.13	0.94	1.23				
abcb7	ATP-binding cassette, sub-family B	ENSDARG00000062795	0.96	0.87	1.07	0.78				
abcd3a	ATP-binding cassette, sub-family D (ALD),	ENSDARG00000104085	0.93	0.74	0.83	0.83				
abcc5	ATP-binding cassette, sub-family C	ENSDARG00000061233	0.88	2.13	0.60	3.10				
abcb8	ATP-binding cassette, sub-family B	ENSDARG00000056672	0.64	2.31	1.88	0.78				
abcb5	ATP-binding cassette, sub-family B	ENSDARG00000021787	0.61	1.84	1.33	0.84				
abcd4	ATP-binding cassette, sub-family D (ALD),	ENSDARG00000061770	0.57	4.75	2.48	1.10				
actr2a	ARP2 actin-related protein 2a homolog (yeast)	ENSDARG00000052438	2.98	1.08	1.58	2.04				
actb1	actin, beta 1	ENSDARG00000037746	1.72	0.99	0.83	2.04				
actb2	actin, beta 2	ENSDARG00000037870	1.24	1.02	1.10	1.15				
actl6a	actin-like 6A	ENSDARG00000070828	1.06	1.21	0.72	1.77				
actr6	ARP6 actin-related protein 6 homolog (yeast)	ENSDARG00000021370	1.05	2.37	1.84	1.36				
actr1	ARP1 actin-related protein 1, centractin (yeast)	ENSDARG00000011611	1.05	1.21	1.10	1.16				
actr8	ARP8 actin-related protein 8 homolog (yeast)	ENSDARG00000103610	1.05	0.64	1.01	0.66				
actr10	actin-related protein 10 homolog (S. cerevisiae)	ENSDARG00000038432	0.94	0.84	0.72	1.10				
actc1b	actin, alpha, cardiac muscle 1b	ENSDARG00000099197	0.69	0.83	0.97	0.59				
acta2	actin, alpha 2, smooth muscle, aorta	ENSDARG00000045180	0.64	0.84	0.69	0.78				
actc1	actin, alpha, cardiac muscle 2	ENSDARG00000057911	0.64	0.79	1.23	0.41				
actc1a	actin alpha cardiac muscle 1a	ENSDARG00000042535	0.59	0.83	0.93	0.53				
acta1b	actin alpha 1, skeletal muscle b	ENSDARG00000055618	0.50	0.98	0.84	0.58				
acta1a	actin, alpha 1a, skeletal muscle	ENSDARG00000036371	0.35	1.01	0.81	0.44				
actc1c	actin alpha cardiac muscle 1c	ENSDARG00000079111	0.21	1.00	0.84	0.26				
actn1	actinin, alpha 1	ENSDARG00000007219	2.58	3.11	4.51	1.78				
actn4	actinin, alpha 4	ENSDARG00000099786	1.95	1.02	0.97	2.05				
actn3b	actinin alpha 3b	ENSDARG00000001431	0.78	1.15	1.35	0.67				
actn2b	actinin, alpha 2b	ENSDARG00000071090	0.73	0.70	0.74	0.69				
actn3a	actinin alpha 3a	ENSDARG00000013755	0.61	1.03	1.09	0.58				
ankrd10a	ankyrin repeat domain 10a	ENSDARG00000037100	1.81	0.79	0.96	1.50				
ankrd49	ankyrin repeat domain 49	ENSDARG00000023508	1.53	0.90	1.33	1.04				
ankmy2a	ankyrin repeat and MYND domain containing	ENSDARG00000005948	1.48	0.50	0.96	0.77				
ankfy1	ankyrin repeat and FYVE domain containing 1	ENSDARG00000061013	1.22	0.78	0.89	1.07				
ankrd28b	ankyrin repeat domain 28b	ENSDARG00000009023	1.12	1.13	1.27	1.00				
ankar	ankyrin and armadillo repeat containing	ENSDARG00000000516	1.00	2.31	1.00	1.70				
ankhd1	ankyrin repeat and KH domain containing 1	ENSDARG00000077860	0.97	1.05	0.86	1.18				
ankra2	ankyrin repeat, family A (RFXANK-like), 2	ENSDARG00000035399	0.95	0.97	1.33	0.69				
ankrd24	ankyrin repeat domain 24	ENSDARG00000062103	0.95	0.40	0.69	0.55				
ankmy1	ankyrin repeat and MYND domain containing 1	ENSDARG00000062702	0.94	2.15	0.78	2.61				
ankrd13c	ankyrin repeat domain 13C	ENSDARG00000103831	0.93	0.89	0.91	0.91				
anks1b	ankyrin repeat and sterile alpha motif domain	ENSDARG00000003512	0.79	0.79	0.90	0.69				
ankrd22	ankyrin repeat domain 22	ENSDARG00000002298	0.76	1.48	1.41	0.79				
ankrd46b	ankyrin repeat domain 46b	ENSDARG00000015780	0.71	1.38	1.56	0.63				

Fold change
- >5.0 = red
- >3.0 = light red
- >1.2 = pale red
- <0.85 = pale blue
- <0.65 = light blue
- <0.5 = blue

Gene counts
- >1
- >10
- >100
- >1000

- p<0.05
- p<0.1

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
ankrd12	ankyrin repeat domain 12	ENSDARG00000052419	0.68	1.00	0.67	1.02				
ank2b	ankyrin 2b, neuronal	ENSDARG00000043313	0.59	1.65	0.89	1.10				
ankrd1b	ankyrin repeat domain 1b (cardiac muscle)	ENSDARG00000076192	0.53	1.23	1.43	0.46				
ankrd11	ankyrin repeat domain 11	ENSDARG00000051886	0.52	1.86	0.83	1.16				
ank1a	ankyrin 1, erythrocytic a	ENSDARG00000092143	0.40	1.18	0.75	0.63				
ankrd9	ankyrin repeat domain 9	ENSDARG00000028804	0.36	4.21	2.86	0.53				
ankrd1a	ankyrin repeat domain 1a (cardiac muscle)	ENSDARG00000075263	0.28	1.99	1.15	0.48				
apoda.2	apolipoprotein Da, duplicate 2	ENSDARG00000060350	14.99	0.38	7.41	0.77				
apoba	apolipoprotein Ba	ENSDARG00000042780	11.88	0.07	3.53	0.24				
apobb.1	apolipoprotein Bb, tandem duplicate 1	ENSDARG00000022767	8.78	0.12	3.74	0.28				
apoea	apolipoprotein Ea	ENSDARG00000102004	7.70	0.38	4.57	0.63				
apoa4b.2	apolipoprotein A-IV b, tandem duplicate 2	ENSDARG00000020866	7.45	0.06	2.89	0.15				
apoa1a	apolipoprotein A-Ia	ENSDARG00000012076	7.29	0.75	3.44	1.59				
apoc1	apolipoprotein C-I like	ENSDARG00000092170	5.77	0.83	2.29	2.10				
apobb.2	apolipoprotein B-100	ENSDARG00000075016	5.56	0.15	1.00	1.00				
apoa2	apolipoprotein A-II	ENSDARG00000015866	4.40	0.43	1.55	1.23				
apoeb	apolipoprotein Eb	ENSDARG00000040295	4.15	0.56	1.26	1.79				
apoa4b.1	apolipoprotein A-IV b, tandem duplicate 1	ENSDARG00000040298	3.26	0.46	0.93	1.62				
apoa1b	apolipoprotein A-Ib	ENSDARG00000101324	1.61	1.06	1.54	1.11				
apodb	apolipoprotein Db	ENSDARG00000057437	1.34	1.74	2.01	1.16				
apoa4a	apolipoprotein A-IV a	ENSDARG00000101160	1.00	6.86	3.04	1.63				
apoob	apolipoprotein O, b	ENSDARG00000026444	0.99	0.76	0.90	0.84				
apol1	apolipoprotein L, 1	ENSDARG00000007425	0.96	1.24	1.68	0.71				
apool	apolipoprotein O-like	ENSDARG00000039374	0.94	1.05	1.42	0.69				
apoc2	apolipoprotein C-II	ENSDARG00000092155	0.84	0.83	0.33	2.07				
apooa	apolipoprotein O, a	ENSDARG00000104696	0.74	1.15	1.11	0.76				
apof	apolipoprotein F	ENSDARG00000090980	0.68	2.83	3.01	0.64				
arpc5a	actin related protein 2/3 complex, subunit 5A	ENSDARG00000039142	2.95	0.78	0.76	3.01				
arpc5b	actin related protein 2/3 complex, subunit 5B	ENSDARG00000019062	2.61	0.95	2.39	1.04				
arpc1b	actin related protein 2/3 complex, subunit 1B	ENSDARG00000027063	2.24	1.01	1.31	1.73				
arpc3	actin related protein 2/3 complex, subunit 3	ENSDARG00000057882	1.64	0.95	1.04	1.50				
arpc4	actin related protein 2/3 complex, subunit 4	ENSDARG00000054063	1.61	0.94	1.17	1.29				
arpc2	actin related protein 2/3 complex, subunit 2	ENSDARG00000075989	1.61	0.78	0.93	1.35				
arpc4l	actin related protein 2/3 complex, subunit 4,	ENSDARG00000058225	1.32	1.41	1.34	1.39				
arpc1a	actin related protein 2/3 complex, subunit 1A	ENSDARG00000008383	1.18	0.83	0.76	1.29				
atp9b	ATPase, class II, type 9B	ENSDARG00000062521	20.41	0.47	16.43	0.58				
atp2a2b	ATPase, Ca++ transporting, cardiac muscle,	ENSDARG00000005122	2.82	0.89	1.61	1.57				
atp6v0b	ATPase H+ transporting V0 subunit b	ENSDARG00000031681	2.75	1.38	2.14	1.77				
atp6v1ba	ATPase, H+ transporting, lysosomal, V1	ENSDARG00000013443	2.07	0.75	1.10	1.41				
atp6ap1b	ATPase, H+ transporting, lysosomal accessory	ENSDARG00000037153	2.07	0.73	1.17	1.30				
atp6v0ca	ATPase, H+ transporting, lysosomal, V0	ENSDARG00000057853	1.96	0.65	0.80	1.58				
atp6v1ab	ATPase, H+ transporting, lysosomal, V1	ENSDARG00000076318	1.75	0.86	1.33	1.13				
atp6v1f	ATPase, H+ transporting, lysosomal, V1	ENSDARG00000045543	1.61	0.63	0.97	1.05				
atp6v1h	ATPase, H+ transporting, lysosomal V1 subunit	ENSDARG00000006370	1.59	0.75	1.04	1.15				
atp6v1aa	ATPase H+ transporting V1 subunit Aa	ENSDARG00000034534	1.43	0.80	0.64	1.80				
atp6v1d	ATPase, H+ transporting, lysosomal V1 subunit	ENSDARG00000011175	1.36	0.69	0.84	1.11				
atp6v1e1b	ATPase, H+ transporting, lysosomal, V1	ENSDARG00000030694	1.33	0.85	0.94	1.20				
atp1a1a.1	ATPase, Na+/K+ transporting, alpha 1a	ENSDARG00000002791	1.30	0.96	0.88	1.42				
atp6v1g1	ATPase, H+ transporting, lysosomal, V1	ENSDARG00000022315	1.29	0.95	1.13	1.08				
atp6ap2	ATPase, H+ transporting, lysosomal accessory	ENSDARG00000008735	1.29	0.80	1.04	0.99				

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
atp6v1c1b	ATPase, H+ transporting, lysosomal, V1	ENSDARG00000035880	1.28	0.79	0.74	1.36				
atp1b1b	ATPase, Na+/K+ transporting, beta 1b	ENSDARG00000076833	1.24	1.06	1.49	0.88				
atp6v0d1	ATPase, H+ transporting, lysosomal V0 subunit	ENSDARG00000069090	1.23	1.03	1.21	1.05				
atp6v1c1a	ATPase, H+ transporting, lysosomal, V1	ENSDARG00000023967	1.13	1.40	1.46	1.09				
atp13a1	ATPase type 13A1	ENSDARG00000029931	1.13	1.00	0.60	1.88				
atp1b2b	ATPase, Na+/K+ transporting, beta 2b	ENSDARG00000034424	1.12	0.59	0.52	1.26				
atp1b1a	ATPase, Na+/K+ transporting, beta 1a	ENSDARG00000013144	1.12	0.79	0.81	1.09				
atp1b2a	ATPase, Na+/K+ transporting, beta 2a	ENSDARG00000099203	1.08	0.96	1.36	0.77				
atp1a1a.2	ATPase, Na+/K+ transporting, alpha 1a	ENSDARG00000007739	1.03	0.88	1.15	0.79				
atp5if1b	ATPase inhibitory factor 1b	ENSDARG00000044092	0.97	1.15	1.50	0.75				
atp1a1b	ATPase, Na+/K+ transporting, alpha 1b	ENSDARG00000019856	0.96	0.83	0.79	1.01				
atpv0e2	ATPase, H+ transporting V0 subunit e2	ENSDARG00000059057	0.91	1.00	1.64	0.56				
atp10b	ATPase, class V, type 10B	ENSDARG00000076230	0.86	1.12	1.10	0.88				
atp1b3a	ATPase, Na+/K+ transporting, beta 3a	ENSDARG00000015790	0.84	1.08	1.07	0.84				
atp6v1b2	ATPase, H+ transporting, lysosomal, V1	ENSDARG00000043465	0.82	1.17	1.00	0.96				
atp2b4	ATPase, Ca++ transporting, plasma membrane	ENSDARG00000044902	0.80	1.01	0.76	1.07				
atp2a1	ATPase, Ca++ transporting, cardiac muscle,	ENSDARG00000020574	0.80	1.12	1.13	0.79				
atp2b3b	ATPase plasma membrane Ca2+ transporting	ENSDARG00000023445	0.76	0.94	1.16	0.62				
atp2b2	ATPase, Ca++ transporting, plasma membrane	ENSDARG00000063433	0.76	0.96	0.82	0.89				
atp1a3b	ATPase, Na+/K+ transporting, alpha 3b	ENSDARG00000104139	0.76	1.03	0.77	1.01				
atp2b3a	ATPase, Ca++ transporting, plasma membrane	ENSDARG00000043474	0.76	1.06	0.75	1.08				
atp1a2a	ATPase, Na+/K+ transporting, alpha 2a	ENSDARG00000010472	0.74	1.72	1.63	0.79				
atp1a3a	ATPase, Na+/K+ transporting, alpha 3a	ENSDARG00000018259	0.74	1.13	0.93	0.89				
atp2b1a	ATPase, Ca++ transporting, plasma membrane	ENSDARG00000012684	0.71	0.83	0.75	0.79				
atp5if1a	ATPase inhibitory factor 1a	ENSDARG00000067975	0.70	0.76	0.70	0.76				
atp6ap1la	ATPase, H+ transporting, lysosomal accessory	ENSDARG00000091509	0.63	0.76	0.61	0.79				
atp1b3b	ATPase, Na+/K+ transporting, beta 3b	ENSDARG00000042837	0.58	0.98	0.86	0.66				
atp2a2a	ATPase, Ca++ transporting, cardiac muscle,	ENSDARG00000029439	0.56	0.98	0.73	0.75				
atp6v0cb	ATPase, H+ transporting, lysosomal, V0	ENSDARG00000036577	0.54	1.07	0.80	0.73				
atp2a1l	ATPase, Ca++ transporting, cardiac muscle,	ENSDARG00000035458	0.52	1.36	1.03	0.69				
atp1a1a.4	ATPase, Na+/K+ transporting, alpha 1a	ENSDARG00000001870	0.52	1.70	0.85	1.04				
chd4a	chromodomain helicase DNA binding protein	ENSDARG00000063535	1.32	0.98	0.76	1.70				
chd1	chromodomain helicase DNA binding protein 1	ENSDARG00000103787	1.24	0.93	0.94	1.23				
chd1l	chromodomain helicase DNA binding protein 1-	ENSDARG00000015471	1.18	0.85	1.00	1.01				
chd4b	chromodomain helicase DNA binding protein	ENSDARG00000025789	1.05	1.05	0.94	1.17				
chd7	chromodomain helicase DNA binding protein 7	ENSDARG00000075211	0.88	0.98	0.78	1.11				
chd2	chromodomain helicase DNA binding protein 2	ENSDARG00000060687	0.81	1.25	0.91	1.11				
chd9	chromodomain helicase DNA binding protein 9	ENSDARG00000074498	0.77	1.03	0.86	0.92				
chd6	chromodomain helicase DNA binding protein 6	ENSDARG00000017244	0.75	0.87	0.65	1.00				
chd5	chromodomain helicase DNA binding protein 5	ENSDARG00000105083	0.57	0.74	0.45	0.93				
chd3	chromodomain helicase DNA binding protein 3	ENSDARG00000021405	0.53	0.98	0.72	0.72				
cldn5a	claudin 5a	ENSDARG00000043716	1.52	2.35	0.93	3.85				
cldni	claudin i	ENSDARG00000054616	1.42	1.01	1.04	1.38				
cldne	claudin e	ENSDARG00000043128	1.23	1.37	1.54	1.10				
cldn7b	claudin 7b	ENSDARG00000014047	1.19	1.45	1.52	1.13				
cldn15la	claudin 15-like a	ENSDARG00000016081	1.15	1.63	2.68	0.70				
cldn12	claudin 12	ENSDARG00000003927	1.10	0.77	0.88	0.96				
cldna	claudin a	ENSDARG00000069888	0.91	3.36	1.21	2.51				
cldnb	claudin b	ENSDARG00000009544	0.88	1.84	1.57	1.03				
cldnh	claudin h	ENSDARG00000069503	0.79	1.42	1.11	1.02				
cldn11b	claudin 11b	ENSDARG00000030723	0.78	0.91	0.92	0.76				

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
cldng	claudin g	ENSDARG00000003701	0.74	0.87	0.92	0.71				
cldnk	claudin k	ENSDARG00000042357	0.74	1.05	1.33	0.58				
cldn8.3	claudin 8.3	ENSDARG00000099950	0.72	1.60	4.43	0.26				
cldnd1a	claudin domain containing 1a	ENSDARG00000104439	0.71	0.74	0.45	1.16				
cldnc	claudin c	ENSDARG00000015955	0.70	2.99	1.86	1.13				
cldn19	claudin 19	ENSDARG00000044569	0.61	1.87	0.86	1.34				
cldnd	claudin d	ENSDARG00000006580	0.59	0.85	1.00	0.50				
cldn11a	claudin 11a	ENSDARG00000020031	0.50	0.91	0.86	0.53				
col17a1b	collagen, type XVII, alpha 1b	ENSDARG00000079011	1.46	0.87	1.04	1.23				
col12a1a	collagen, type XII, alpha 1a	ENSDARG00000078322	1.45	0.81	0.94	1.26				
col5a3a	collagen type V alpha-3a	ENSDARG00000098294	1.33	0.58	0.20	3.92				
col5a2a	collagen, type V, alpha 2a	ENSDARG00000031678	1.13	0.96	1.09	1.00				
col8a1a	collagen, type VIII, alpha 1a	ENSDARG00000077403	1.00	19.40	1.00	14.29				
col4a1	collagen, type IV, alpha 1	ENSDARG00000055009	1.00	1.04	0.83	1.25				
col5a1	procollagen, type V, alpha 1	ENSDARG00000012593	0.91	0.84	0.85	0.90				
col28a2a	collagen type XXVIII alpha 1 a	ENSDARG00000076321	0.88	0.77	1.04	0.65				
col15a1b	collagen, type XV, alpha 1b	ENSDARG00000061848	0.84	0.58	1.29	0.38				
col1a2	collagen, type I, alpha 2	ENSDARG00000020007	0.77	0.89	0.90	0.76				
col1a1b	collagen, type I, alpha 1b	ENSDARG00000035809	0.76	1.17	1.22	0.73				
col14a1a	collagen, type XIV, alpha 1a	ENSDARG00000005762	0.74	1.17	1.08	0.80				
col4a6	collagen, type IV, alpha 6	ENSDARG00000052061	0.72	1.49	1.35	0.80				
col1a1a	collagen, type I, alpha 1a	ENSDARG00000012405	0.72	1.01	1.07	0.68				
col4a5	collagen, type IV, alpha 5 (Alport syndrome)	ENSDARG00000052063	0.68	1.29	1.06	0.83				
col10a1a	collagen, type X, alpha 1a	ENSDARG00000054753	0.50	0.57	0.54	0.53				
col9a2	procollagen, type IX, alpha 2	ENSDARG00000024492	0.40	0.84	0.90	0.37				
col9a3	collagen, type IX, alpha 3	ENSDARG00000037845	0.40	0.97	1.17	0.33				
col9a1b	collagen, type IX, alpha 1b	ENSDARG00000031483	0.35	1.13	1.00	0.40				
col11a1a	collagen, type XI, alpha 1a	ENSDARG00000026165	0.33	1.05	0.76	0.46				
col2a1a	collagen, type II, alpha 1a	ENSDARG00000069093	0.32	1.77	1.51	0.37				
col11a2	collagen, type XI, alpha 2	ENSDARG00000012422	0.30	1.16	0.92	0.38				
col7a1	collagen, type VII, alpha 1	ENSDARG00000021720	0.21	2.36	0.84	0.60				
ctsl.1	cathepsin L.1	ENSDARG00000003902	11.91	0.91	1.28	8.44				
ctss2.1	cathepsin Sb, tandem duplicate 1	ENSDARG00000074656	6.91	2.06	2.15	6.61				
ctsk	cathepsin K	ENSDARG00000040251	4.58	0.82	1.46	2.58				
ctss2.2	cathepsin Sb, tandem duplicate 2	ENSDARG00000013771	3.58	1.46	3.70	1.42				
ctsh	cathepsin H	ENSDARG00000041108	2.99	0.86	1.41	1.83				
ctsba	cathepsin Ba	ENSDARG00000055120	2.29	1.16	2.44	1.09				
ctsc	cathepsin C	ENSDARG00000101334	2.27	0.90	1.01	2.01				
ctsz	cathepsin Z	ENSDARG00000043081	2.02	0.79	1.08	1.48				
ctsla	cathepsin La	ENSDARG00000007836	1.94	1.12	1.43	1.51				
ctsd	cathepsin D	ENSDARG00000057698	1.67	1.09	1.24	1.46				
ctss1	cathepsin Sa	ENSDARG00000036940	1.07	1.04	1.24	0.89				
ctsll	cathepsin L, like	ENSDARG00000011701	0.73	1.17	5.20	0.16				
ctsf	cathepsin F	ENSDARG00000063095	0.70	1.06	1.05	0.72				
cyba	cytochrome b-245, alpha polypeptide	ENSDARG00000018283	3.59	1.17	1.66	2.54				
cybb	cytochrome b-245, beta polypeptide (chronic	ENSDARG00000056615	3.51	0.58	1.17	1.74				
cyb5b	cytochrome b5 type B	ENSDARG00000099774	2.36	0.75	0.83	2.14				
cyb5d1	cytochrome b5 domain containing 1	ENSDARG00000056007	1.66	1.03	0.37	4.66				
cyb5r3	cytochrome b5 reductase 3	ENSDARG00000005891	1.65	1.11	1.74	1.05				
cyb5a	cytochrome b5 type A (microsomal)	ENSDARG00000098589	1.51	1.30	1.59	1.23				

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
cyb5r1	cytochrome b5 reductase 1	ENSDARG00000018966	1.06	1.11	1.10	1.07				
dnajb9b		ENSDARG00000016886	3.27	1.16	0.50	7.50				
dnajc25		ENSDARG00000067613	2.75	4.37	1.00	12.25				
dnajc3a	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000041110	2.05	0.80	0.98	1.67				
dnajb11	DnaJ heat shock protein family (Hsp40)	ENSDARG00000015088	1.98	0.74	0.79	1.85				
dnajc12	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000086691	1.72	1.44	2.93	0.84				
dnajc5b		ENSDARG00000058147	1.65	1.14	1.64	1.14				
dnajb6b	DnaJ (Hsp40) homolog, subfamily B, member	ENSDARG00000020953	1.61	0.86	1.17	1.19				
dnajc9	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000031293	1.56	0.62	0.73	1.33				
dnajc16		ENSDARG00000059699	1.40	0.68	0.17	5.79				
dnaja1	DnaJ (Hsp40) homolog, subfamily A, member 1	ENSDARG00000030972	1.34	0.93	0.98	1.28				
dnajb12a	DnaJ (Hsp40) homolog, subfamily B, member	ENSDARG00000039363	1.33	1.03	1.73	0.80				
dnajc3b	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000017874	1.30	1.00	0.96	1.35				
dnajb1a	DnaJ (Hsp40) homolog, subfamily B, member	ENSDARG00000099383	1.28	0.96	1.03	1.19				
dnajc10	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000074727	1.26	0.76	0.76	1.26				
dnaja2b	DnaJ (Hsp40) homolog, subfamily A, member	ENSDARG00000010745	1.26	0.76	1.02	0.94				
dnajc8	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000059373	1.26	0.60	0.68	1.11				
dnajb1b	DnaJ (Hsp40) homolog, subfamily B, member	ENSDARG00000041394	1.24	6.57	4.66	1.75				
dnajc5gb	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000017687	1.22	2.53	2.48	1.25				
dnajc17	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000104959	1.16	0.55	0.63	1.01				
dnajb12b		ENSDARG00000087473	1.15	1.26	1.09	1.33				
dnajc21	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000105195	1.14	0.72	0.77	1.06				
dnajc16l		ENSDARG00000060725	1.12	1.12	1.47	0.86				
dnaja3a	DnaJ (Hsp40) homolog, subfamily A, member	ENSDARG00000058494	1.11	0.90	1.15	0.86				
dnaja2a	DnaJ (Hsp40) homolog, subfamily A, member 2	ENSDARG00000104066	1.09	0.82	0.86	1.04				
dnajc11a	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000011196	1.04	0.94	0.97	1.00				
dnajc11b		ENSDARG00000102696	1.02	1.08	1.01	1.09				
dnajb14		ENSDARG00000069996	1.00	0.88	1.31	0.68				
dnajc1	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000001940	0.99	0.94	0.91	1.02				
dnajb2		ENSDARG00000058644	0.96	1.69	2.52	0.65				
dnajc15		ENSDARG00000038309	0.92	1.38	1.39	0.92				
dnajc7	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000058148	0.89	0.93	0.89	0.93				
dnajc5ga	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000041896	0.89	0.71	0.87	0.72				
dnajc6		ENSDARG00000079891	0.88	1.02	1.14	0.79				
dnajc27	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000070916	0.84	0.95	1.23	0.65				
dnajc19	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000044420	0.83	0.78	0.89	0.73				
dnajc5ab	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000004836	0.83	0.82	0.80	0.85				
dnajc5aa		ENSDARG00000042948	0.82	1.15	1.14	0.82				
dnajb4		ENSDARG00000038978	0.80	1.35	1.16	0.93				
dnajc24	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000023927	0.79	0.68	0.59	0.89				
dnajc28		ENSDARG00000018181	0.78	1.30	1.19	0.86				
dnajc4		ENSDARG00000024090	0.78	1.44	1.38	0.81				
dnaja3b	DnaJ (Hsp40) homolog, subfamily A, member	ENSDARG00000102295	0.78	2.28	2.25	0.79				
dnajc2		ENSDARG00000070477	0.71	0.94	0.74	0.90				
dnajc14		ENSDARG00000105398	0.67	2.08	0.35	4.02				
dnajb6a	DnaJ (Hsp40) homolog, subfamily B, member	ENSDARG00000004680	0.67	2.48	2.72	0.61				
dnajb13		ENSDARG00000043157	0.66	1.38	0.81	1.14				
dnajc22		ENSDARG00000037067	0.57	2.10	0.49	2.44				
dnajc30b	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000079415	0.56	1.17	2.49	0.26				
dnajc18	DnaJ (Hsp40) homolog, subfamily C, member	ENSDARG00000056005	0.47	1.19	0.97	0.58				
dnajb1	DnaJ (Hsp40) homolog, subfamily B, member 1	ENSDARG00000015831								

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
hoxc5a	homeobox C5a	ENSDARG00000070340	8.15	1.47	1.00	12.17				
hoxa1a	homeobox A1a	ENSDARG00000104307	7.96	1.06	1.00	8.62				
hoxa13b	homeobox A13b	ENSDARG00000036254	1.80	0.99	1.35	1.32				
hoxd13a	homeobox D13a	ENSDARG00000059256	1.70	0.84	1.72	0.82				
hoxa11a	homeobox A11a	ENSDARG00000104162	1.58	1.69	1.99	1.34				
hoxb7a	homeobox B7a	ENSDARG00000056030	1.53	1.24	2.61	0.73				
hoxd12a	homeobox D12a	ENSDARG00000059263	1.52	1.57	2.63	0.91				
hoxc13a	homeobox C13a	ENSDARG00000070353	1.37	0.88	0.73	1.66				
hoxb8b	homeobox B8b	ENSDARG00000054025	1.31	0.72	2.58	0.37				
hoxb2a	homeobox B2a	ENSDARG00000000175	1.31	1.02	1.37	0.97				
hoxb2a	homeobox B2a	ENSDARG00000000175	1.31	1.02	1.37	0.97				
hoxb6b	homeobox B6b	ENSDARG00000026513	1.26	1.13	1.28	1.11				
hoxa9b	homeobox A9b	ENSDARG00000056819	1.15	1.04	1.59	0.76				
hoxa2b	homeobox A2b	ENSDARG00000023031	1.13	1.14	1.14	1.13				
hoxa9a	homeobox A9a	ENSDARG00000105013	1.09	1.15	1.51	0.83				
hoxb4a	homeobox B4a	ENSDARG00000013533	1.00	2.31	1.00	1.70				
hoxc4a	homeobox C4a	ENSDARG00000070338	1.00	1.00	1.91	0.45				
hoxa10b	homeobox A10b	ENSDARG00000031337	1.00	0.87	1.07	0.81				
hoxd10a	homeobox D10a	ENSDARG00000057859	0.94	1.04	1.08	0.90				
hoxc12b	homeobox C12b	ENSDARG00000103133	0.93	0.89	0.88	0.93				
hoxc9a	homeobox C9a	ENSDARG00000092809	0.91	1.40	0.70	1.81				
hoxb8a	homeobox B8a	ENSDARG00000056027	0.90	0.83	0.98	0.77				
hoxb5b	homeobox B5b	ENSDARG00000054030	0.88	0.71	0.63	0.99				
hoxb5a	homeobox B5a	ENSDARG00000013057	0.87	0.97	0.97	0.86				
hoxc11b	homeobox C11b	ENSDARG00000102631	0.86	1.14	1.19	0.82				
hoxd9a	homeobox D9a	ENSDARG00000059274	0.84	0.63	0.75	0.72				
hoxb3a	homeobox B3a	ENSDARG00000029263	0.82	0.86	1.01	0.70				
hoxd3a	homeobox D3a	ENSDARG00000059280	0.82	1.56	0.86	1.47				
hoxb6a	homeobox B6a	ENSDARG00000010630	0.79	0.73	0.59	0.98				
hoxa11b	homeobox A11b	ENSDARG00000007009	0.77	0.97	0.86	0.86				
hoxa4a	homeobox A3a /// homeobox A4a	ENSDARG00000103862	0.72	0.94	0.88	0.76				
hoxd11a	homeobox D11a	ENSDARG00000059267	0.70	0.95	1.28	0.52				
hoxc6b	homeobox C6b	ENSDARG00000101954	0.67	1.23	0.86	0.97				
hoxd4a	homeobox D4a	ENSDARG00000059276	0.67	1.18	0.94	0.84				
hoxc10a	homeobox C10a	ENSDARG00000070348	0.65	0.94	0.83	0.74				
hoxc6a	homeobox C6a	ENSDARG00000070343	0.59	1.20	0.59	1.19				
hoxc8a	homeobox C8a	ENSDARG00000070346	0.49	1.16	0.46	1.24				
hsp90b1	heat shock protein 90, beta (grp94), member 1	ENSDARG00000114206	11.57	0.87	1.04	9.60				
hsp90b1	heat shock protein 90, beta (grp94), member 1	ENSDARG00000003570	2.00	0.95	1.02	1.86				
hspa4a	heat shock protein 4a	ENSDARG00000004754	1.69	0.87	0.93	1.57				
hspd1	heat shock 60 protein 1	ENSDARG00000056160	1.55	1.06	1.24	1.33				
hspa4b	heat shock protein 4b	ENSDARG00000018989	1.50	0.69	0.87	1.18				
hspa5	heat shock protein 5	ENSDARG00000103846	1.45	0.86	0.99	1.25				
hspa5	heat shock protein 5	ENSDARG00000103846	1.45	0.86	0.99	1.25				
hspb1	heat shock protein, alpha-crystallin-related, 1	ENSDARG00000041065	1.34	0.78	0.90	1.16				
hsp90aa1.2	heat shock protein 90, alpha (cytosolic), class	ENSDARG00000024746	1.33	0.87	0.84	1.38				
hspa14	heat shock protein 14	ENSDARG00000058030	1.23	0.75	0.88	1.05				
hspe1	heat shock 10 protein 1	ENSDARG00000056167	1.17	0.91	0.99	1.08				
hspa9	heat shock protein 9	ENSDARG00000003035	1.14	1.29	1.60	0.92				
hspa8	heat shock protein 8	ENSDARG00000068992	1.11	0.94	0.93	1.13				
hsp90ab1	heat shock protein 90, alpha (cytosolic), class	ENSDARG00000029150	0.91	0.72	0.66	0.99				

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
hspb6	heat shock protein, alpha-crystallin-related, b6	ENSDARG00000077236	0.91	1.55	1.44	0.97				
hspbp1	HSPA (heat shock 70kDa) binding protein,	ENSDARG00000102937	0.84	1.02	1.28	0.67				
hspb8	heat shock protein b8	ENSDARG00000058365	0.78	1.14	1.09	0.82				
hsp90aa1.1	heat shock protein 90, alpha (cytosolic), class	ENSDARG00000010478	0.74	0.91	1.02	0.66				
hspa4l	heat shock protein 4 like	ENSDARG00000053544	0.70	1.11	0.90	0.86				
hspa12a	heat shock protein 12A	ENSDARG00000070603	0.65	0.98	1.06	0.60				
hspb15	heat shock protein, alpha-crystallin-related, b15	ENSDARG00000078411	0.57	1.20	0.87	0.79				
igfbp5a		ENSDARG00000039264	2.75	0.12	1.52	0.22				
igfbp1a	insulin-like growth factor binding protein 1a	ENSDARG00000099351	2.01	1.59	1.27	2.53				
igfbp1b	insulin-like growth factor binding protein 1b	ENSDARG00000038666	1.36	5.17	1.21	5.85				
igfbp6a		ENSDARG00000070941	1.15	0.41	1.60	0.29				
igfbp7		ENSDARG00000104138	1.00	2.31	1.91	0.87				
igfbp6b	insulin-like growth factor binding protein 6b	ENSDARG00000090833	0.90	0.50	0.63	0.71				
igfbp5b	insulin-like growth factor binding protein 5b	ENSDARG00000025348	0.87	1.89	1.72	0.96				
igfbp2b		ENSDARG00000031422	0.85	0.68	0.13	4.62				
igfbp3	insulin-like growth factor binding protein 3	ENSDARG00000099144	0.82	3.38	3.79	0.73				
igfbp2a	insulin-like growth factor binding protein 2a	ENSDARG00000052470	0.75	0.80	1.09	0.55				
krt93	xkeratin 93	ENSDARG00000044976	5.95	0.14	1.00	1.00				
krt18a.1	keratin 18	ENSDARG00000018404	5.05	0.75	1.18	3.20				
krt96	keratin 96	ENSDARG00000095147	2.86	0.19	0.92	0.59				
krt97	keratin 97	ENSDARG00000000212	2.66	0.30	0.74	1.08				
krt94	keratin 94	ENSDARG00000044975	1.93	0.64	0.54	2.31				
krt18b	keratin zgc:77517 krtt1c6	ENSDARG00000028618	1.91	0.86	0.93	1.76				
krt8	keratin 8	ENSDARG00000058358	1.67	0.75	0.95	1.32				
krt1-c5	keratin 1	ENSDARG00000026979	1.22	0.71	0.74	1.17				
krt222	keratin 222	ENSDARG00000071518	1.18	0.90	1.25	0.85				
krt99	keratin 99 zgc:110712	ENSDARG00000019365	0.94	0.58	1.41	0.39				
krt98	keratin 98	ENSDARG00000044973	0.71	0.31	0.66	0.33				
krt5	keratin 5	ENSDARG00000058371	0.56	1.44	1.30	0.62				
krt92	keratin 96?	ENSDARG00000036834	0.53	2.05	1.01	1.08				
krt4	keratin 4	ENSDARG00000017624	0.49	1.33	0.99	0.66				
krtt1c19e	keratin type 1 c19e	ENSDARG00000090268	0.47	0.96	0.90	0.51				
krt91	keratin 91	ENSDARG00000036830	0.47	1.19	0.83	0.67				
krt17	keratin 17	ENSDARG00000094041	0.46	1.24	0.85	0.68				
krt15	keratin 15	ENSDARG00000036840	0.36	1.35	0.65	0.74				
krt1-19d	keratin, type 1, gene 19d	ENSDARG00000023082	0.23	1.97	0.82	0.55				
map1lc3c	microtubule-associated protein 1 light chain 3	ENSDARG00000100528	6.03	0.14	0.25	3.45				
mapre1b	microtubule-associated protein, RP/EB family,	ENSDARG00000002659	1.67	0.67	0.91	1.24				
map1lc3a	microtubule-associated protein 1 light chain 3	ENSDARG00000033609	1.37	1.30	1.89	0.94				
mapre1a	microtubule-associated protein, RP/EB family,	ENSDARG00000042927	1.14	1.11	1.11	1.14				
map1lc3b	microtubule-associated protein 1 light chain 3	ENSDARG00000101127	1.02	1.01	1.00	1.03				
map9	microtubule-associated protein 9	ENSDARG00000037276	1.02	1.08	0.65	1.68				
map1sa	microtubule-associated protein 1Sa	ENSDARG00000060805	0.79	0.86	0.84	0.81				
maptb	microtubule-associated protein tau b	ENSDARG00000087616	0.73	0.58	0.73	0.58				
map1ab	microtubule-associated protein 1Ab	ENSDARG00000022045	0.62	0.94	0.84	0.69				
mapre3b	microtubule-associated protein, RP/EB family,	ENSDARG00000102878	0.58	1.01	0.93	0.64				
map1lc3cl	microtubule-associated protein 1 light chain 3	ENSDARG00000075727	0.29	0.82	1.08	0.22				
mmp13b	matrix metallopeptidase 13b	ENSDARG00000100794	78.68	0.94	6.93	10.68				
mmp9	matrix metallopeptidase 9	ENSDARG00000042816	10.35	2.42	3.37	7.44				

Fold change
- >5.0 = red
- >3.0 = light red
- >1.2 = pale red
- <0.85 = pale blue
- <0.65 = light blue
- <0.5 = blue

Gene counts
- >1
- >10
- >100
- >1000

- p<0.05
- p<0.1

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
mmp13a	matrix metallopeptidase 13a	ENSDARG00000012395	7.49	1.01	1.20	6.30				
mmp13a	matrix metallopeptidase 13a	ENSDARG00000114451	4.34	1.91	1.53	5.42				
mmp14b	matrix metallopeptidase 14b (membrane-	ENSDARG00000008388	2.01	0.94	1.88	1.01				
mmp2	matrix metallopeptidase 2	ENSDARG00000017676	1.64	1.61	3.05	0.87				
mmp14a	matrix metallopeptidase 14a (membrane-	ENSDARG00000002235	1.27	0.96	1.17	1.04				
MMP23B	matrix metallopeptidase 23B	ENSDARG00000043079	1.00	3.92	1.00	2.89				
mmp17b	matrix metallopeptidase 17b	ENSDARG00000102956	0.88	1.28	1.18	0.96				
mmp30	matrix metallopeptidase 30	ENSDARG00000045887	0.62	2.44	1.22	1.25				
ms4a17a.4	membrane-spanning 4-domains	ENSDARG00000014024	9.71	0.70	1.04	6.57				
ms4a17a.5	membrane-spanning 4-domains	ENSDARG00000092204	6.24	0.67	1.30	3.21				
ms4a17a.2	membrane-spanning 4-domains	ENSDARG00000105674	5.82	0.70	0.96	4.23				
ms4a17a.7	membrane-spanning 4-domains	ENSDARG00000043796	5.36	0.81	1.65	2.64				
ms4a17a.1	membrane-spanning 4-domains, subfamily A,	ENSDARG00000043798	4.40	0.57	0.48	5.14				
ms4a17a.12	membrane-spanning 4-domains	ENSDARG00000053563	2.74	0.86	1.04	2.27				
ms4a17a.11	membrane-spanning 4-domains	ENSDARG00000094809	2.71	0.88	0.86	2.77				
ms4a17a.9	membrane-spanning 4-domains	ENSDARG00000094854	2.32	0.87	1.23	1.64				
mybphb	myosin binding protein Hb	ENSDARG00000003081	0.99	2.04	2.21	0.92				
mybpc2b	myosin binding protein C, fast type b	ENSDARG00000021265	0.88	0.66	0.77	0.75				
mybpc3	myosin binding protein C, cardiac	ENSDARG00000011615	0.63	0.98	1.02	0.61				
mybpc1	myosin binding protein C1	ENSDARG00000045560	0.52	0.71	0.82	0.45				
mybpha	myosin binding protein Ha	ENSDARG00000058799	0.33	1.33	0.99	0.44				
myl12.1	myosin, light chain 12, genome duplicate 1	ENSDARG00000099766	1.56	1.18	1.55	1.19				
mylipa	myosin regulatory light chain interacting protein	ENSDARG00000008859	1.01	1.45	1.45	1.01				
myl7	myosin, light chain 7, regulatory	ENSDARG00000019096	1.00	2.31	1.91	0.87				
myl12.2	myosin, light chain 12, genome duplicate 2	ENSDARG00000025326	0.99	0.84	0.95	0.87				
myl6	myosin, light chain 6, alkali, smooth muscle	ENSDARG00000008494	0.86	0.92	0.87	0.91				
myl10	myosin, light chain 10, regulatory	ENSDARG00000062592	0.77	0.75	0.91	0.64				
myl1	myosin, light chain 1, alkali; skeletal, fast	ENSDARG00000014196	0.73	0.80	0.99	0.59				
mylpfb	myosin light chain, phosphorylatable, fast	ENSDARG00000002589	0.72	1.06	1.25	0.61				
myl13	myosin, light chain 13	ENSDARG00000042245	0.69	1.07	1.02	0.72				
mylpfa	myosin light chain, phosphorylatable, fast	ENSDARG00000053254	0.62	1.22	1.06	0.72				
mylz3	myosin, light polypeptide 3, skeletal muscle	ENSDARG00000017441	0.61	0.99	0.94	0.64				
mylk4a	myosin light chain kinase 3-like	ENSDARG00000091260	0.54	2.73	0.76	1.93				
ndufs1	NADH dehydrogenase (ubiquinone) Fe-S	ENSDARG00000028546	4.06	0.21	1.00	1.00				
ndufa4	NADH dehydrogenase (ubiquinone) 1 alpha	ENSDARG00000056108	1.03	2.49	3.37	0.76				
ndufaf4	NADH dehydrogenase (ubiquinone) complex I,	ENSDARG00000077859	0.97	1.16	1.29	0.88				
ndufs8a	NADH dehydrogenase (ubiquinone) Fe-S	ENSDARG00000051986	0.93	1.03	1.14	0.84				
ndufs5	NADH dehydrogenase (ubiquinone) Fe-S	ENSDARG00000006290	0.90	0.80	0.88	0.82				
ndufb4	NADH dehydrogenase (ubiquinone) 1 beta	ENSDARG00000019332	0.86	1.36	1.36	0.87				
ndufb8	NADH dehydrogenase (ubiquinone) 1 beta	ENSDARG00000010113	0.86	0.81	0.98	0.71				
ndufb9	NADH dehydrogenase (ubiquinone) 1 beta	ENSDARG00000041314	0.85	0.82	0.91	0.76				
ndufv2	NADH dehydrogenase (ubiquinone)	ENSDARG00000013044	0.84	0.90	1.31	0.58				
ndufc2	NADH dehydrogenase (ubiquinone) 1,	ENSDARG00000102115	0.84	0.94	1.05	0.75				
ndufa6	NADH dehydrogenase (ubiquinone) 1 alpha	ENSDARG00000038028	0.82	0.73	0.75	0.80				
ndufb10	NADH dehydrogenase (ubiquinone) 1 beta	ENSDARG00000028889	0.82	0.96	0.99	0.79				
ndufaf1	NADH dehydrogenase (ubiquinone) complex I,	ENSDARG00000025549	0.80	1.66	1.47	0.90				
ndufab1a	NADH dehydrogenase (ubiquinone) 1,	ENSDARG00000058463	0.79	1.19	1.10	0.85				
ndufv1	NADH dehydrogenase (ubiquinone)	ENSDARG00000036438	0.78	1.12	1.07	0.82				
ndufs2	NADH dehydrogenase (ubiquinone) Fe-S	ENSDARG00000007526	0.78	1.03	1.04	0.77				

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
ndufb5	NADH dehydrogenase (ubiquinone) 1 beta	ENSDARG00000070824	0.76	1.13	1.26	0.69				
ndufaf6	NADH dehydrogenase (ubiquinone) complex I,	ENSDARG00000053652	0.75	0.85	0.69	0.92				
ndufb6	NADH dehydrogenase (ubiquinone) 1 beta	ENSDARG00000037259	0.75	1.12	1.10	0.76				
ndufs6	NADH dehydrogenase (ubiquinone) Fe-S	ENSDARG00000056583	0.75	1.46	2.00	0.54				
ndufa8	NADH dehydrogenase (ubiquinone) 1 alpha	ENSDARG00000058041	0.73	1.19	1.22	0.71				
ndufb3	NADH dehydrogenase (ubiquinone) 1 beta	ENSDARG00000075709	0.72	0.86	0.89	0.70				
ndufb7	NADH dehydrogenase (ubiquinone) 1 beta	ENSDARG00000033789	0.71	0.85	0.88	0.69				
ndufs4	NADH dehydrogenase (ubiquinone) Fe-S	ENSDARG00000052840	0.71	0.89	0.85	0.74				
ndufa10	NADH dehydrogenase (ubiquinone) 1 alpha	ENSDARG00000013333	0.69	0.94	0.82	0.80				
ndufa4l	NADH dehydrogenase (ubiquinone) 1 alpha	ENSDARG00000099499	0.68	1.13	1.03	0.74				
NDUFB1	NADH dehydrogenase (ubiquinone) 1 beta	ENSDARG00000087456	0.67	1.05	0.87	0.81				
ndufab1b	NADH dehydrogenase (ubiquinone) 1,	ENSDARG00000014915	0.66	0.99	0.95	0.69				
ndufa5	NADH dehydrogenase (ubiquinone) 1 alpha	ENSDARG00000039346	0.65	1.16	0.96	0.79				
ndufb2	NADH dehydrogenase (ubiquinone) 1 beta	ENSDARG00000045490	0.65	1.01	0.89	0.74				
ndufv3	NADH:ubiquinone oxidoreductase subunit V3	ENSDARG00000090389	0.64	0.89	0.84	0.68				
ndufaf5	NADH dehydrogenase (ubiquinone) complex I,	ENSDARG00000061629	0.64	0.90	0.77	0.75				
ndufa2	NADH dehydrogenase (ubiquinone) 1 alpha	ENSDARG00000021984	0.62	0.90	0.75	0.75				
ndufa11	NADH dehydrogenase (ubiquinone) 1 alpha	ENSDARG00000042777	0.62	1.01	0.89	0.70				
ndufs7	NADH dehydrogenase (ubiquinone) Fe-S	ENSDARG00000074552	0.54	1.06	0.77	0.75				
ndufa1	NADH dehydrogenase (ubiquinone) 1 alpha	ENSDARG00000036329	0.52	1.10	0.80	0.71				
nr5a5	nuclear receptor subfamily 5, group A, member	ENSDARG00000039116	1.60	6.65	2.67	3.98				
nr5a1b	nuclear receptor subfamily 5, group A, member	ENSDARG00000023362	1.41	2.32	4.27	0.76				
nr1h3	nuclear receptor subfamily 1, group H, member	ENSDARG00000098439	1.34	1.06	1.26	1.13				
nrip1b	nuclear receptor interacting protein 1b	ENSDARG00000068894	1.30	1.47	2.00	0.95				
nrf1	nuclear respiratory factor 1	ENSDARG00000000018	1.22	0.89	1.18	0.93				
nr2f6a	nuclear receptor subfamily 2, group F, member	ENSDARG00000003607	1.19	0.69	0.96	0.85				
nr2f1a	nuclear receptor subfamily 2, group F, member	ENSDARG00000052695	1.11	0.76	1.03	0.83				
nrbf2b	nuclear receptor binding factor 2b	ENSDARG00000023591	1.05	1.10	1.14	1.01				
nr2f5	nuclear receptor subfamily 2, group F, member	ENSDARG00000033172	1.01	1.61	1.34	1.22				
nr2e1	nuclear receptor subfamily 2, group E, member	ENSDARG00000017107	1.00	1.00	3.04	0.28				
nr2f6b	nuclear receptor subfamily 2, group F, member	ENSDARG00000003165	0.98	1.04	0.98	1.04				
nr1h5	nuclear receptor subfamily 1, group H, member	ENSDARG00000031046	0.91	2.98	3.21	0.84				
nr3c1	nuclear receptor subfamily 3, group C, member	ENSDARG00000025032	0.85	0.87	0.85	0.86				
nr2f2	nuclear receptor subfamily 2, group F, member	ENSDARG00000040926	0.78	1.08	1.08	0.78				
nr4a2b	nuclear receptor subfamily 4, group A, member	ENSDARG00000044532	0.67	1.63	1.13	0.97				
nr2f1b	nuclear receptor subfamily 2, group F, member	ENSDARG00000017168	0.67	1.11	0.67	1.10				
nr1d2b	nuclear receptor subfamily 1, group D, member	ENSDARG00000009594	0.64	1.23	1.94	0.41				
nr1d2a	nuclear receptor subfamily 1, group D, member	ENSDARG00000003820	0.62	1.39	1.15	0.75				
nr0b2a	nuclear receptor subfamily 0, group B, member	ENSDARG00000044685	0.61	2.92	2.29	0.77				
nr2c2	nuclear receptor subfamily 2, group C, member	ENSDARG00000042477	0.53	0.97	0.93	0.55				
nr4a1	nuclear receptor subfamily 4, group A, member	ENSDARG00000000796	0.52	3.44	2.55	0.70				
nr1d1	nuclear receptor subfamily 1, group d, member	ENSDARG00000033160	0.49	0.92	1.41	0.31				
nr6a1a	nuclear receptor subfamily 6, group A, member	ENSDARG00000101508	0.47	0.61	0.33	1.00				
nr5a2	nuclear receptor subfamily 5, group A, member	ENSDARG00000100940	0.42	1.11	0.50	0.95				
nr5a1a	nuclear receptor subfamily 5, group A, member	ENSDARG00000103176	0.24	3.76	0.77	1.16				
ppp1r18	protein phosphatase 1, regulatory subunit 18	ENSDARG00000071251	4.57	0.55	1.16	2.15				
ppp1r11	protein phosphatase 1, regulatory (inhibitor)	ENSDARG00000036063	4.50	0.72	4.57	0.71				
ppp2r2ab	protein phosphatase 2, regulatory subunit B,	ENSDARG00000006624	2.48	1.81	1.56	2.88				
ppp2r2ab	protein phosphatase 2, regulatory subunit B,	ENSDARG00000006624	2.48	1.81	1.56	2.88				
ppp4r2b	protein phosphatase 4, regulatory subunit 2b	ENSDARG00000053447	2.15	0.79	1.56	1.08				
ppp1r3ca	protein phosphatase 1, regulatory subunit 3Ca	ENSDARG00000071005	2.05	1.35	0.80	3.46				

Legend:

Fold change
>5.0 = red
>3.0 = light red
>1.2 = pale red
<0.85 = pale blue
<0.65 = light blue
<0.5 = blue

Gene counts
>1
>10
>100
>1000

p<0.05   p<0.1

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
ppp1r3da	protein phosphatase 1, regulatory subunit 3Da	ENSDARG00000077513	2.00	0.95	1.69	1.12				
ppp4ca	protein phosphatase 4, catalytic subunit a	ENSDARG00000070570	1.86	0.91	1.54	1.10				
ppp4r2a	protein phosphatase 4, regulatory subunit 2a	ENSDARG00000026540	1.75	0.77	1.23	1.10				
ppp1caa	protein phosphatase 1, catalytic subunit, alpha	ENSDARG00000003486	1.48	1.43	1.94	1.09				
ppp1r15a	protein phosphatase 1, regulatory subunit 15A	ENSDARG00000069135	1.42	2.59	1.58	2.33				
ppp1cab	protein phosphatase 1, catalytic subunit, alpha	ENSDARG00000071566	1.41	0.82	1.02	1.13				
ppp1r8b	protein phosphatase 1, regulatory subunit 8b	ENSDARG00000022430	1.37	0.80	0.87	1.25				
ppp1r14bb	protein phosphatase 1, regulatory (inhibitor)	ENSDARG00000030161	1.36	1.06	1.67	0.87				
ppp1r3b	protein phosphatase 1, regulatory subunit 3B	ENSDARG00000044691	1.27	0.44	0.98	0.57				
ppp2cb	protein phosphatase 2, catalytic subunit, beta	ENSDARG00000099241	1.18	0.85	0.80	1.25				
ppp2r1ba	protein phosphatase 2, regulatory subunit A,	ENSDARG00000007791	1.16	0.70	0.77	1.05				
ppp4r3b	protein phosphatase 4, regulatory subunit 3B	ENSDARG00000103415	1.16	0.63	0.50	1.45				
ppp1r37	protein phosphatase 1, regulatory subunit 37	ENSDARG00000078458	1.13	1.57	1.34	1.32				
ppp1r13ba	protein phosphatase 1, regulatory subunit 13Ba	ENSDARG00000004377	1.11	0.99	1.33	0.82				
ppp1r13l	protein phosphatase 1, regulatory subunit 13	ENSDARG00000013777	1.09	0.82	0.74	1.21				
ppp6r3	protein phosphatase 6, regulatory subunit 3	ENSDARG00000013379	1.06	1.06	1.02	1.11				
ppp1r9bb	neurabin-2-like /// protein phosphatase 1,	ENSDARG00000071709	1.04	0.65	0.99	0.69				
ppp2r3c	protein phosphatase 2, regulatory subunit B",	ENSDARG00000043972	1.04	1.38	1.53	0.93				
ppp6r2a	protein phosphatase 6, regulatory subunit 2a	ENSDARG00000045540	1.03	1.14	1.12	1.04				
ppp2r5d	protein phosphatase 2, regulatory subunit B',	ENSDARG00000014428	1.03	0.96	0.98	1.01				
ppp5c	protein phosphatase 5, catalytic subunit	ENSDARG00000034313	1.02	0.76	0.80	0.96				
ppp1r2	protein phosphatase 1, regulatory (inhibitor)	ENSDARG00000054007	0.98	1.71	1.40	1.20				
ppp1cb	protein phosphatase 1, catalytic subunit, beta	ENSDARG00000044153	0.98	1.05	1.16	0.88				
ppp1r12a	protein phosphatase 1, regulatory subunit 12A	ENSDARG00000010784	0.98	1.46	1.27	1.13				
ppp2r1bb	protein phosphatase 2, regulatory subunit A,	ENSDARG00000032430	0.97	0.87	0.85	0.99				
ppp2r5eb	protein phosphatase 2, regulatory subunit B',	ENSDARG00000069118	0.96	1.45	1.78	0.78				
ppp2r3b	protein phosphatase 2, regulatory subunit B",	ENSDARG00000076004	0.89	1.27	1.09	1.04				
ppp1r7	protein phosphatase 1, regulatory (inhibitor)	ENSDARG00000009740	0.89	0.84	0.84	0.89				
ppp1r10	protein phosphatase 1, regulatory subunit 10	ENSDARG00000032651	0.89	1.20	1.01	1.06				
ppp3r1b	protein phosphatase 3 (formerly 2B), regulatory	ENSDARG00000069360	0.88	0.81	1.00	0.71				
ppp1r14ab	protein phosphatase 1, regulatory (inhibitor)	ENSDARG00000017710	0.86	1.02	0.96	0.91				
ppp1r14aa	protein phosphatase 1, regulatory (inhibitor)	ENSDARG00000011239	0.82	0.94	1.17	0.66				
ppp1r42	protein phosphatase 1, regulatory subunit 42	ENSDARG00000057632	0.80	0.70	1.03	0.54				
ppp3cb	protein phosphatase 3, catalytic subunit, beta	ENSDARG00000025106	0.79	1.00	1.09	0.72				
ppp6c	protein phosphatase 6, catalytic subunit	ENSDARG00000002949	0.78	2.39	1.78	1.05				
ppp2r5ea	protein phosphatase 2, regulatory subunit B',	ENSDARG00000015474	0.77	1.13	0.98	0.88				
ppp4r4	protein phosphatase 4, regulatory subunit 4	ENSDARG00000010407	0.74	2.05	1.46	1.04				
ppp1r13bb	protein phosphatase 1, regulatory subunit 13Bb	ENSDARG00000009142	0.71	0.92	0.70	0.94				
ppp2r2ca	protein phosphatase 2, regulatory subunit B,	ENSDARG00000056797	0.71	0.87	0.55	1.12				
ppp1r15b	protein phosphatase 1, regulatory subunit 15B	ENSDARG00000068128	0.69	2.17	0.90	1.67				
ppp3cca	protein phosphatase 3, catalytic subunit,	ENSDARG00000014962	0.66	0.90	0.64	0.94				
ppp2r5b	protein phosphatase 2, regulatory subunit B',	ENSDARG00000054931	0.61	1.79	0.98	1.11				
ppp3ca	protein phosphatase 3, catalytic subunit, alpha	ENSDARG00000004988	0.55	1.40	1.01	0.76				
ppp1r3ab	protein phosphatase 1, regulatory subunit 3Ab	ENSDARG00000088813	0.54	1.52	0.86	0.96				
ppp2r5ca	protein phosphatase 2, regulatory subunit B',	ENSDARG00000059083	0.50	2.93	0.50	2.90				
ppp1r1b	protein phosphatase 1, regulatory (inhibitor)	ENSDARG00000076280	0.46	0.93	0.58	0.73				
ppp1r3cb	protein phosphatase 1, regulatory subunit 3Cb	ENSDARG00000014554	0.42	1.18	0.72	0.69				
pvalb8	parvalbumin 8	ENSDARG00000037790	0.94	0.82	1.00	0.78				
pvalb2	parvalbumin 2	ENSDARG00000002768	0.69	1.41	1.40	0.69				
pvalb4	parvalbumin 4	ENSDARG00000024433	0.66	0.99	0.99	0.66				
pvalb5	parvalbumin 5	ENSDARG00000032836	0.66	1.01	0.91	0.73				
pvalb9	parvalbumin 9	ENSDARG00000071601	0.59	0.54	0.82	0.39				

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
pvalb6	parvalbumin 6	ENSDARG00000009311	0.59	1.12	0.99	0.67				
pvalb3	parvalbumin 3	ENSDARG00000022817	0.54	1.10	0.97	0.61				
pvalb7	parvalbumin 7	ENSDARG00000034705	0.52	0.86	0.88	0.51				
pvalb1	parvalbumin 1	ENSDARG00000037789	0.49	1.08	0.79	0.67				
slc7a7	solute carrier family 7 (amino acid transporter	ENSDARG00000055226	7.70	0.67	1.00	5.15				
slc46a1	solute carrier family 46 (folate transporter),	ENSDARG00000026149	6.06	0.87	7.83	0.67				
slc22a7b.1	solute carrier family 22 (organic anion	ENSDARG00000056643	2.85	0.15	1.57	0.28				
slc39a7	solute carrier family 39 (zinc transporter),	ENSDARG00000104451	2.46	0.59	0.95	1.53				
slc12a8	solute carrier family 12, member 8	ENSDARG00000074384	2.16	1.15	1.58	1.57				
slc30a1a	solute carrier family 30 (zinc transporter),	ENSDARG00000005463	2.13	0.83	1.17	1.52				
slc51a	solute carrier family 51 member A	ENSDARG00000045306	2.05	0.82	0.80	2.10				
slc13a2	solute carrier family 13 (sodium-dependent	ENSDARG00000053853	1.98	2.55	2.32	2.17				
slc52a3	solute carrier family 52 (riboflavin transporter),	ENSDARG00000042737	1.97	1.51	1.39	2.15				
slc46a3	solute carrier family 46, member 3	ENSDARG00000077313	1.96	1.48	2.10	1.39				
slco1d1	solute carrier organic anion transporter family,	ENSDARG00000104108	1.95	0.68	1.50	0.88				
slco2b1	solute carrier organic anion transporter family,	ENSDARG00000054609	1.92	1.29	1.20	2.07				
slc25a19	solute carrier family 25 (mitochondrial thiamine	ENSDARG00000100385	1.89	1.76	4.65	0.71				
slc10a2	solute carrier family 10 (sodium/bile acid	ENSDARG00000014916	1.87	1.86	2.63	1.32				
slc2a10	solute carrier family 2 (facilitated glucose	ENSDARG00000090820	1.82	0.54	0.69	1.43				
slc24a2	solute carrier family 24	ENSDARG00000042988	1.79	1.71	1.58	1.95				
slc29a3	solute carrier family 29 (equilibrative nucleoside	ENSDARG00000077828	1.77	0.92	1.05	1.55				
slc10a7	solute carrier family 10, member 7	ENSDARG00000104508	1.74	0.69	0.95	1.27				
slc12a9	solute carrier family 12, member 9	ENSDARG00000060366	1.74	0.90	1.48	1.05				
slc20a1a	solute carrier family 20, member 1a	ENSDARG00000020114	1.73	1.10	1.80	1.06				
slc14a2	solute carrier family 14 member 2	ENSDARG00000051914	1.70	2.45	5.38	0.78				
slc9a3r1a	solute carrier family 9, subfamily A (NHE3,	ENSDARG00000000068	1.67	0.57	1.04	0.92				
slc35f6	solute carrier family 35, member F6	ENSDARG00000016745	1.66	0.85	0.94	1.52				
slc26a1	solute carrier family 26 (anion exchanger),	ENSDARG00000029832	1.65	0.47	1.07	0.73				
slc12a4	solute carrier family 12 (potassium/chloride	ENSDARG00000014378	1.64	1.52	1.94	1.29				
slc43a3b	solute carrier family 43 member 3b	ENSDARG00000057949	1.61	1.40	1.12	2.02				
slc52a2	solute carrier family 52 (riboflavin transporter),	ENSDARG00000102035	1.61	0.07	0.76	0.14				
slc4a1ap	solute carrier family 4 (anion exchanger),	ENSDARG00000034160	1.60	0.81	1.02	1.26				
slc2a11b	solute carrier family 2 (facilitated glucose	ENSDARG00000063288	1.59	0.79	1.59	0.79				
slc4a4b	solute carrier family 4 (sodium bicarbonate	ENSDARG00000044808	1.54	0.33	0.33	1.53				
slc26a11	solute carrier family 26 (anion exchanger),	ENSDARG00000043021	1.48	0.71	0.78	1.34				
slc12a3	solute carrier family 12 (sodium/chloride	ENSDARG00000013855	1.46	1.83	1.45	1.84				
slc39a9	solute carrier family 39, member 9	ENSDARG00000070447	1.46	0.87	0.59	2.14				
slc35f2	solute carrier family 35, member F2	ENSDARG00000069745	1.45	0.79	1.00	1.14				
slc31a1	solute carrier family 31 (copper transporter),	ENSDARG00000013961	1.41	0.65	0.78	1.18				
slc35b2	solute carrier family 35 (adenosine 3'-phospho	ENSDARG00000007886	1.41	1.47	2.30	0.90				
slc22a31	solute carrier family 22, member 31	ENSDARG00000078882	1.40	0.77	1.12	0.97				
slc30a7	solute carrier family 30 (zinc transporter),	ENSDARG00000019998	1.39	0.80	0.97	1.14				
slc35d1a	solute carrier family 35 (UDP-GlcA/UDP-	ENSDARG00000011973	1.36	0.89	1.44	0.84				
slc38a2	solute carrier family 38, member 2	ENSDARG00000045886	1.34	0.57	0.87	0.89				
slc17a9b	solute carrier family 17 (vesicular nucleotide	ENSDARG00000011049	1.33	0.85	0.61	1.84				
slc38a7	solute carrier family 38, member 7	ENSDARG00000012002	1.26	0.87	0.86	1.27				
slc25a33	solute carrier family 25 (pyrimidine nucleotide	ENSDARG00000039931	1.23	1.34	2.08	0.79				
slc16a9a	solute carrier family 16, member 9a	ENSDARG00000013926	1.22	2.42	0.93	3.17				
slc39a6	solute carrier family 39 (zinc transporter),	ENSDARG00000068143	1.20	0.76	0.73	1.25				
slc15a4	solute carrier family 15 (oligopeptide	ENSDARG00000102612	1.19	0.78	1.06	0.87				
slc13a5a	info solute carrier family 13 (sodium-dependent	ENSDARG00000077691	1.17	0.16	1.02	0.18				
slc40a1	solute carrier family 40 (iron-regulated	ENSDARG00000000241	1.16	1.10	1.34	0.95				

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)
slc20a1b	solute carrier family 20 (phosphate transporter),	ENSDARG00000010641	1.16	1.37	1.29	1.24
slc9a7	solute carrier family 9, subfamily A (NHE7,	ENSDARG00000076754	1.15	0.57	0.82	0.80
slc25a40	solute carrier family 25 member 40	ENSDARG00000015856	1.15	2.26	2.27	1.14
slc4a1a	solute carrier family 4 (anion exchanger),	ENSDARG00000012881	1.14	0.75	1.05	0.81
slc12a7a	solute carrier family 12 (potassium/chloride	ENSDARG00000073756	1.13	1.21	1.18	1.15
slc48a1b	solute carrier family 48 (heme transporter),	ENSDARG00000026109	1.11	1.29	2.08	0.69
slc16a9b	solute carrier family 16, member 9b	ENSDARG00000104687	1.11	1.64	2.21	0.82
slc25a20	solute carrier family 25 (carnitine/acylcarnitine	ENSDARG00000040401	1.09	0.74	0.84	0.97
slc12a7b	solute carrier family 12 (potassium/chloride	ENSDARG00000062058	1.09	0.97	1.19	0.89
slc16a6b	solute carrier family 16, member 6b	ENSDARG00000060246	1.07	1.48	1.30	1.22
slc5a6b	solute carrier family 5 (sodium/multivitamin and	ENSDARG00000014599	1.06	1.24	1.37	0.96
slc6a9	solute carrier family 6 (neurotransmitter	ENSDARG00000018534	1.06	1.03	1.12	0.97
slc43a2b	solute carrier family 43 (amino acid system L	ENSDARG00000061120	1.04	1.04	1.28	0.84
slc30a5	solute carrier family 30 (zinc transporter),	ENSDARG00000051921	1.03	1.20	1.31	0.95
slc2a15b	solute carrier family 2 (facilitated glucose	ENSDARG00000053269	1.03	1.60	3.19	0.52
slc35b4	solute carrier family 35, member B4	ENSDARG00000071087	1.03	1.00	0.93	1.12
slc16a1b	solute carrier family 16 (monocarboxylate	ENSDARG00000068572	1.02	0.56	0.68	0.84
slc35c2	solute carrier family 35 (GDP-fucose	ENSDARG00000037517	1.02	1.14	1.27	0.92
slc25a48	solute carrier family 25, member 48	ENSDARG00000021250	1.01	0.56	0.83	0.68
slc27a1a	solute carrier family 27 (fatty acid transporter),	ENSDARG00000006240	1.01	1.15	1.58	0.73
slc25a43	solute carrier family 25, member 43	ENSDARG00000102048	1.00	1.62	1.76	0.92
slc39a5	solute carrier family 39 (zinc transporter),	ENSDARG00000079525	1.00	2.31	1.00	1.70
slc25a26	solute carrier family 25 (S-adenosylmethionine	ENSDARG00000058208	1.00	0.92	0.93	0.99
slc25a55a	solute carrier family 25 (mitochondrial carrier:	ENSDARG00000020893	0.99	0.86	0.88	0.97
slc4a2a	solute carrier family 4 (anion exchanger),	ENSDARG00000028173	0.99	0.74	1.05	0.69
slc25a17	solute carrier family 25 (mitochondrial carrier;	ENSDARG00000061684	0.98	0.82	0.61	1.31
slc3a2a	solute carrier family 3 (amino acid transporter	ENSDARG00000036427	0.97	1.10	1.26	0.85
slc30a9	solute carrier family 30 (zinc transporter),	ENSDARG00000057272	0.96	1.20	1.28	0.90
slc37a4a	solute carrier family 37 (glucose-6-phosphate	ENSDARG00000038106	0.96	0.82	0.96	0.82
slc26a5	solute carrier family 26 (anion exchanger),	ENSDARG00000022424	0.95	1.21	1.62	0.71
slc16a10	solute carrier family 16 member 10	ENSDARG00000020984	0.95	0.63	0.85	0.70
slc25a44b	solute carrier family 25, member 44 b	ENSDARG00000035905	0.94	0.95	0.92	0.98
slc25a1a	solute carrier family 25 (mitochondrial carrier;	ENSDARG00000057110	0.94	1.10	1.23	0.84
slc6a1b	solute carrier family 6 (neurotransmitter	ENSDARG00000039647	0.93	0.78	0.74	0.98
slc1a4	solute carrier family 1 (glutamate/neutral amino	ENSDARG00000000551	0.90	0.68	0.98	0.63
slc22a2	solute carrier family 22 (organic cation	ENSDARG00000030530	0.89	1.56	1.66	0.84
slc25a38b	solute carrier family 25, member 38b	ENSDARG00000074533	0.88	0.59	0.73	0.71
slc6a19b	solute carrier family 6 (neutral amino acid	ENSDARG00000056719	0.87	3.33	0.32	9.20
slc1a2b	solute carrier family 1 (glial high affinity	ENSDARG00000102453	0.87	0.80	0.94	0.75
slc43a1a	solute carrier family 43 (amino acid system L	ENSDARG00000037393	0.85	0.70	0.76	0.79
slc35e1	solute carrier family 35, member E1	ENSDARG00000011945	0.85	0.99	0.86	0.98
slc29a2	solute carrier family 29 (equilibrative nucleoside	ENSDARG00000001767	0.85	1.24	1.21	0.87
slc7a6os	solute carrier family 7, member 6 opposite	ENSDARG00000010596	0.83	1.06	0.81	1.09
slc25a32a	solute carrier family 25 (mitochondrial folate	ENSDARG00000089791	0.82	1.09	1.01	0.88
slc24a4a	solute carrier family 24	ENSDARG00000015425	0.82	1.55	1.14	1.11
slc25a14	solute carrier family 25 (mitochondrial carrier,	ENSDARG00000026680	0.81	1.13	0.95	0.97
slc25a28	solute carrier family 25 (mitochondrial iron	ENSDARG00000052994	0.80	1.22	1.09	0.90
slc25a39	solute carrier family 25, member 39	ENSDARG00000007449	0.80	1.03	1.45	0.57
slc39a14	solute carrier family 39 (zinc transporter),	ENSDARG00000102387	0.79	1.03	0.35	2.36
slc6a17	solute carrier family 6 (neutral amino acid	ENSDARG00000068787	0.79	1.01	0.90	0.88
slc37a2	solute carrier family 37 (glucose-6-phosphate	ENSDARG00000023394	0.79	0.97	0.76	1.01
slc32a1	solute carrier family 32 (GABA vesicular	ENSDARG00000059775	0.79	1.00	0.81	0.97
slc20a2	solute carrier family 20 (phosphate transporter),	ENSDARG00000060796	0.79	1.27	1.02	0.98

Count columns (heat-map squares): DMSO unoperated COUNT, DMSO operated COUNT, SB43 unoperated COUNT, SB43 operated COUNT

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)	DMSO unoperated COUNT / DMSO operated COUNT / SB43 unoperated COUNT / SB43 operated COUNT
slc22a7b.2	solute carrier family 22 member 7-like	ENSDARG00000091252	0.78	0.68	0.24	2.27	
slc27a2a	solute carrier family 27 (fatty acid transporter),	ENSDARG00000036237	0.78	0.53	1.11	0.37	
slc7a6	solute carrier family 7 member 6	ENSDARG00000054423	0.78	1.20	1.60	0.59	
slc7a8a	solute carrier family 7 (cationic amino acid	ENSDARG00000075831	0.78	1.33	1.45	0.71	
slc25a12	solute carrier family 25 (aspartate/glutamate	ENSDARG00000102362	0.77	0.97	1.26	0.59	
slc25a11	solute carrier family 25 (mitochondrial carrier;	ENSDARG00000035741	0.77	0.86	0.90	0.73	
slc48a1a	solute carrier family 48 (heme transporter),	ENSDARG00000026907	0.75	2.03	1.54	1.00	
slc25a27	solute carrier family 25, member 27	ENSDARG00000042873	0.75	1.28	1.32	0.73	
slc25a5	solute carrier family 25 (mitochondrial carrier;	ENSDARG00000092553	0.75	1.30	1.16	0.84	
slc7a3a	solute carrier family 7 (cationic amino acid	ENSDARG00000020645	0.75	2.39	2.02	0.88	
slc2a3a	solute carrier family 2 (facilitated glucose	ENSDARG00000013295	0.75	1.39	1.56	0.67	
slc2a2	solute carrier family 2 (facilitated glucose	ENSDARG00000056196	0.72	1.19	0.98	0.87	
slc6a11a	solute carrier family 6 (neurotransmitter	ENSDARG00000074002	0.72	9.84	7.75	0.92	
slc1a3a	solute carrier family 1 (glial high affinity	ENSDARG00000104431	0.71	1.09	1.06	0.73	
slc38a5b	solute carrier family 38, member 5b	ENSDARG00000014587	0.71	0.81	0.78	0.74	
slc16a3	solute carrier family 16 (monocarboxylate	ENSDARG00000045051	0.71	1.15	1.15	0.71	
slc25a3b	solute carrier family 25 (mitochondrial carrier;	ENSDARG00000025566	0.70	0.86	0.79	0.76	
slc25a25a	solute carrier family 25 (mitochondrial carrier;	ENSDARG00000010572	0.70	2.26	1.46	1.08	
slc18a3a	solute carrier family 18 (vesicular acetylcholine	ENSDARG00000006356	0.69	0.92	0.83	0.76	
slc25a55b	solute carrier family 25 member 55b	ENSDARG00000013749	0.68	0.74	0.83	0.60	
slc39a13	solute carrier family 39 (zinc transporter),	ENSDARG00000000442	0.67	0.75	1.01	0.50	
slc16a12b	solute carrier family 16, member 12b	ENSDARG00000089885	0.65	1.14	0.94	0.78	
slc38a4	solute carrier family 38, member 4	ENSDARG00000018149	0.65	0.94	0.94	0.64	
slc43a2a	solute carrier family 43 (amino acid system L	ENSDARG00000036848	0.64	1.97	0.55	2.30	
slc35a4	solute carrier family 35, member A4	ENSDARG00000062379	0.63	1.17	0.85	0.87	
slc38a6	solute carrier family 38, member 6	ENSDARG00000054312	0.63	3.23	1.87	1.08	
slc3a2b	solute carrier family 3 (amino acid transporter	ENSDARG00000037012	0.62	1.75	0.71	1.53	
slc9a8	solute carrier family 9, subfamily A (NHE8,	ENSDARG00000020699	0.62	1.19	0.98	0.76	
slc37a4b	solute carrier family 37 (glucose-6-phosphate	ENSDARG00000077180	0.59	0.99	0.80	0.74	
slc6a19a.1	solute carrier family 6 (neutral amino acid	ENSDARG00000018621	0.59	2.23	1.79	0.74	
slc44a5b	solute carrier family 44 member 5b	ENSDARG00000057419	0.58	1.89	0.72	1.53	
slc25a51b	solute carrier family 25 member 51-like	ENSDARG00000040463	0.58	1.05	0.82	0.75	
slc25a36b	solute carrier family 25 member 36b	ENSDARG00000015915	0.57	1.18	0.86	0.78	
slc25a4	solute carrier family 25 (mitochondrial carrier;	ENSDARG00000027355	0.54	0.97	0.85	0.61	
slc16a4	solute carrier family 16 member 4	ENSDARG00000042807	0.52	1.29	0.75	0.89	
slc8a4b	solute carrier family 8 (sodium/calcium	ENSDARG00000037145	0.49	1.49	0.63	1.16	
slc2a11l	solute carrier family 2 (facilitated glucose	ENSDARG00000062873	0.44	0.21	0.33	0.28	
slc13a1	solute carrier family 13 (sodium/sulphate	ENSDARG00000045638	0.40	1.53	0.36	1.68	
slc5a12	solute carrier family 5	ENSDARG00000005004	0.40	6.00	1.29	1.85	
slc25a3a	solute carrier family 25 (mitochondrial carrier;	ENSDARG00000027424	0.35	2.76	0.69	1.38	
slc25a44a	solute carrier family 25, member 44 a	ENSDARG00000045927	0.16	3.92	0.62	1.03	
slc25a37	solute carrier family 25 (mitochondrial iron	ENSDARG00000073743	0.12	2.31	0.50	0.54	
tgm2l	transglutaminase 2, like	ENSDARG00000093381	7.27	0.78	2.09	2.70	
tgm2b	transglutaminase 2b	ENSDARG00000074094	4.20	0.66	3.23	0.86	
tgm2a	transglutaminase 2, C polypeptide A	ENSDARG00000070157	0.34	0.79	1.19	0.22	
tgm1l4	transglutaminase 1 like 4	ENSDARG00000101407	0.27	0.68	0.25	0.74	
tnnt2a	troponin T type 2a (cardiac)	ENSDARG00000020610	4.17	0.21	1.00	1.00	
tnnt2c	troponin T2c, cardiac	ENSDARG00000032242	3.90	0.73	1.01	2.82	
tnni2b.1	troponin I type 2b (skeletal, fast), tandem	ENSDARG00000035958	2.34	0.97	1.35	1.68	
tnni1b	troponin I type 1b (skeletal, slow)	ENSDARG00000052708	1.94	0.59	0.62	1.86	
tnnc1a	troponin C type 1a (slow)	ENSDARG00000011400	0.70	1.16	0.12	6.67	

Legend:

Fold change
>5.0 = red
>3.0 = light red
>1.2 = pale red
<0.85 = pale blue
<0.65 = light blue
<0.5 = blue

Gene counts
>1
>10
>100
>1000

p<0.05   p<0.1

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
tnnc1b	troponin C type 1b (slow)	ENSDARG00000037539	0.65	0.90	0.84	0.69				
tnni4b.2	troponin I4b, tandem duplicate 2	ENSDARG00000036671	0.65	1.14	1.01	0.74				
tnni1d	troponin I, skeletal, slow d	ENSDARG00000073766	0.62	0.89	0.97	0.57				
tnnt3a	troponin T type 3a (skeletal, fast)	ENSDARG00000030270	0.60	0.91	0.77	0.71				
tnnc2	troponin C type 2 (fast)	ENSDARG00000070835	0.56	1.22	0.99	0.69				
tnni2a.4	troponin I type 2a (skeletal, fast), tandem	ENSDARG00000029069	0.56	1.12	0.91	0.68				
tnnt3b	troponin T type 3b (skeletal, fast)	ENSDARG00000068457	0.55	1.02	0.92	0.61				
tnni2a.3	troponin I type 2a (skeletal, fast), tandem	ENSDARG00000013752	0.48	1.29	0.70	0.89				
tnnt2d	troponin T2d, cardiac	ENSDARG00000002988	0.43	0.98	0.84	0.51				
tnni1c	troponin I, skeletal, slow c	ENSDARG00000042559	0.32	0.74	0.60	0.39				
tnni2b.2	troponin I type 2b (skeletal, fast), tandem	ENSDARG00000029995	0.31	1.63	0.89	0.58				
tnni4b.1	troponin I4b, tandem duplicate 1	ENSDARG00000092999	0.25	2.11	0.60	0.87				
tubb6	tubulin, beta 6 class V	ENSDARG00000104801	5.22	0.64	1.20	2.78				
tuba1b	tubulin, alpha 1b	ENSDARG00000045367	2.64	0.73	0.71	2.68				
tubd1	tubulin, delta 1	ENSDARG00000058219	2.30	0.68	1.07	1.45				
tuba8l	tubulin, alpha 8 like	ENSDARG00000042708	1.86	0.67	0.69	1.81				
tuba8l4	tubulin, alpha 8 like 4	ENSDARG00000006260	1.58	0.75	0.83	1.43				
tubb4b	tubulin, beta 4B class IVb	ENSDARG00000002344	1.37	0.97	1.24	1.08				
tubb4b	tubulin, beta 4B class IVb	ENSDARG00000002344	1.37	0.97	1.24	1.08				
tubb2b	tubulin, beta 2b	ENSDARG00000098591	1.37	0.81	0.90	1.23				
tubg1	tubulin, gamma 1	ENSDARG00000015610	1.31	0.61	0.97	0.82				
tubgcp4	tubulin, gamma complex associated protein 4	ENSDARG00000005374	1.30	0.82	0.92	1.15				
tubgcp5	tubulin, gamma complex associated protein 5	ENSDARG00000077442	1.28	0.88	1.21	0.93				
tubgcp2	tubulin, gamma complex associated protein 2	ENSDARG00000013079	1.25	1.14	1.43	1.00				
tuba4l	tubulin, alpha 4 like	ENSDARG00000074289	1.22	1.09	0.93	1.43				
tuba7l	tubulin, alpha 7 like	ENSDARG00000104643	1.00	1.00	3.04	0.28				
tuba8l3	tubulin, alpha 8 like 3	ENSDARG00000070155	0.93	0.78	0.83	0.87				
tubb5	tubulin, beta 5	ENSDARG00000037997	0.91	1.67	1.31	1.15				
tubgcp3	tubulin, gamma complex associated protein 3	ENSDARG00000029133	0.89	2.13	2.08	0.91				
tuba2	tubulin, alpha 2	ENSDARG00000045014	0.79	1.11	1.29	0.68				
tubb2	tubulin, beta 2A class IIa	ENSDARG00000039522	0.70	1.20	1.09	0.77				
tuba1a	tubulin, alpha 1a /// zgc:123298	ENSDARG00000001889	0.67	0.89	0.79	0.75				
tuba1c	tubulin, alpha 1c	ENSDARG00000055216	0.59	1.00	0.92	0.64				
tuba8l2	tubulin, alpha 8 like 2	ENSDARG00000031164	0.45	1.17	0.94	0.56				
ube2c	ubiquitin-conjugating enzyme E2C	ENSDARG00000114670	2.33	0.88	0.97	2.11				
ube2d1b	ubiquitin-conjugating enzyme E2D 1b	ENSDARG00000038576	2.10	1.45	1.93	1.58				
ube2d2l	ubiquitin-conjugating enzyme E2D 2 (UBC4/5	ENSDARG00000099749	1.72	0.70	0.95	1.27				
ube3d	ubiquitin protein ligase E3D	ENSDARG00000026178	1.69	0.88	0.94	1.58				
ube2q2	ubiquitin-conjugating enzyme E2Q family	ENSDARG00000013990	1.60	2.18	2.16	1.62				
ube2d1a	ubiquitin-conjugating enzyme E2D 1a	ENSDARG00000029107	1.56	1.86	2.85	1.02				
ube2s	ubiquitin-conjugating enzyme E2S	ENSDARG00000031775	1.46	0.93	0.98	1.38				
ube2v2	ubiquitin-conjugating enzyme E2 variant 2	ENSDARG00000028198	1.36	1.02	0.84	1.64				
ube2l3b	ubiquitin-conjugating enzyme E2L 3b	ENSDARG00000027141	1.36	0.86	1.01	1.15				
ube4b	ubiquitination factor E4B, UFD2 homolog (S.	ENSDARG00000037017	1.29	0.76	0.90	1.09				
ube2na	ubiquitin-conjugating enzyme E2Na	ENSDARG00000008748	1.28	1.12	1.16	1.23				
ube2ia	ubiquitin-conjugating enzyme E2Ia	ENSDARG00000052649	1.25	1.17	1.47	0.99				
ube2r2	ubiquitin-conjugating enzyme E2R 2	ENSDARG00000058740	1.24	1.19	1.28	1.16				
ube2d2	ubiquitin-conjugating enzyme E2D 2 (UBC4/5	ENSDARG00000043484	1.24	0.96	0.91	1.30				
ube2g2	ubiquitin-conjugating enzyme E2G 2 (UBC7	ENSDARG00000025404	1.18	1.68	2.18	0.91				
ube2ib	ubiquitin-conjugating enzyme E2Ib	ENSDARG00000007438	1.15	0.72	0.78	1.06				
ube2kb	ubiquitin-conjugating enzyme E2Kb (UBC1	ENSDARG00000013505	1.11	0.87	0.76	1.28				

Gene Symbol	Gene Title	ensemble	(1) DMSO (operated vs unoperated)	(2) Operated (SB431542 vs DMSO)	(3) Unoperated (SB431542 vs DMSO)	(4) SB431542 (operated vs unoperated)	DMSO unoperated COUNT	DMSO operated COUNT	SB43 unoperated COUNT	SB43 operated COUNT
ube2g1a	ubiquitin-conjugating enzyme E2G 1a (UBC7	ENSDARG00000015292	1.09	1.01	1.23	0.89				
ube2e3	ubiquitin-conjugating enzyme E2E 3 (UBC4/5	ENSDARG00000012244	1.07	1.02	1.13	0.97				
ube2ql1	ubiquitin-conjugating enzyme E2Q family-like 1	ENSDARG00000079276	1.05	1.56	0.82	2.00				
ube2e2	ubiquitin-conjugating enzyme E2E 2	ENSDARG00000034670	1.04	1.11	1.27	0.91				
ube2q1	ubiquitin-conjugating enzyme E2Q family	ENSDARG00000100766	0.98	0.83	0.75	1.08				
ube2a	ubiquitin-conjugating enzyme E2A (RAD6	ENSDARG00000098466	0.95	0.97	1.15	0.80				
ube2nb	ubiquitin-conjugating enzyme E2Nb	ENSDARG00000045877	0.87	0.91	0.91	0.87				
ube2h	ubiquitin-conjugating enzyme E2H (UBC8	ENSDARG00000000019	0.80	0.86	0.98	0.71				
ube2d3	ubiquitin-conjugating enzyme E2D 3	ENSDARG00000038473	0.79	1.18	0.92	1.00				
ube2g1b	ubiquitin-conjugating enzyme E2G 1b (UBC7	ENSDARG00000069527	0.69	1.15	0.95	0.83				
ube2d4	ubiquitin-conjugating enzyme E2D 4	ENSDARG00000015057	0.38	0.95	0.37	0.98				
ywhabb	tyrosine 3-monooxygenase/tryptophan 5-	ENSDARG00000075758	2.08	1.13	1.36	1.72				
ywhaqb	tyrosine 3-monooxygenase/tryptophan 5-	ENSDARG00000023323	1.85	1.54	1.26	2.27				
ywhaba	tyrosine 3-monooxygenase/tryptophan 5-	ENSDARG00000013078	1.57	0.98	1.15	1.33				
ywhaqa	tyrosine 3-monooxygenase/tryptophan 5-	ENSDARG00000042539	1.50	0.81	0.94	1.30				
ywhaz	tyrosine 3-monooxygenase/tryptophan 5-	ENSDARG00000032575	1.49	0.95	0.71	2.00				
ywhae2	tyrosine 3-monooxygenase/tryptophan 5-	ENSDARG00000017014	1.25	0.92	0.93	1.24				
ywhah	tyrosine 3-monooxygenase/tryptophan 5-	ENSDARG00000005560	0.93	0.75	0.81	0.87				
ywhae1	tyrosine 3-monooxygenase/tryptophan 5-	ENSDARG00000006399	0.82	0.87	0.67	1.07				
ywhabl	tyrosine 3-monooxygenase/tryptophan 5-	ENSDARG00000040287	0.81	1.07	0.80	1.09				

Fold change
>5.0 = red
>3.0 = light red
>1.2 = pale red
<0.85 = pale blue
<0.65 = light blue
<0.5 = blue
p<0.05   p<0.1

Gene counts
>1
>10
>100
>1000

Supplemental Table 6. R packages. This table lists the packages and versions used for the scRNA-seq analysis presented in the study.

Package	Version
abind	1.4-8
askpass	1.2.0
assertthat	0.2.1
assorthead	0.99.8
backports	1.5.0
base64enc	0.1-3
beachmat	2.21.6
beeswarm	0.4.0
BH	1.84.0-0
Biobase	2.65.1
BiocGenerics	0.51.1
BiocManager	1.30.25
BiocParallel	1.39.0
BiocSingular	1.21.3
BiocVersion	3.20.0
bit	4.5.0
bit64	4.5.2
bitops	1.0-8
blob	1.2.4
brew	1.0-10
brio	1.1.5
broom	1.0.6
bslib	0.8.0
cachem	1.1.0
Cairo	1.6-2
callr	3.7.6
car	3.1-2
carData	3.0-5
caTools	1.18.3
cellranger	1.1.0
circlize	0.4.16
cli	3.6.3
clipr	0.8.0
clock	0.7.1
clue	0.3-65
colorspace	2.1-1
commonmark	1.9.1
ComplexHeatmap	2.21.0
conflicted	1.2.0
corrplot	0.94
COTAN	2.5.7
cowplot	1.1.3

cpp11	0.5.0
crayon	1.5.3
credentials	2.0.1
crosstalk	1.2.1
curl	5.2.3
data.table	1.16.0
DBI	1.2.3
dbplyr	2.5.0
DelayedArray	0.31.11
DelayedMatrixStats	1.27.3
deldir	2.0-4
dendextend	1.17.1
Deriv	4.1.6
desc	1.4.3
DESeq2	1.45.3
devtools	2.4.5
diagram	1.6.5
dials	1.3.0
DiceDesign	1.1
diffobj	0.3.5
digest	0.6.37
doBy	4.6.22
doFuture	1.0.1
doParallel	1.0.17
dotCall64	1.1-1
downlit	0.4.4
dplyr	1.1.4
dqrng	0.4.1
DropletUtils	1.25.2
dtplyr	1.3.1
edgeR	4.3.15
ellipsis	0.3.2
evaluate	1.0.0
fansi	1.0.6
farver	2.1.2
fastDummies	1.7.4
fastmap	1.2.0
fitdistrplus	1.2-1
FNN	1.1.4.1
fontawesome	0.5.2
forcats	1.0.0
foreach	1.5.2
formatR	1.14
fs	1.6.4
furrr	0.3.1
futile.logger	1.4.3
futile.options	1.0.1
future	1.34.0

future.apply	1.11.2
gargle	1.5.2
generics	0.1.3
GenomeInfoDb	1.41.1
GenomeInfoDbData	1.2.12
GenomicRanges	1.57.1
gert	2.1.2
GetoptLong	1.0.5
ggbeeswarm	0.7.2
ggplot2	3.4.4
ggplotify	0.1.2
ggprism	1.0.5
ggpubr	0.6.0
ggrastr	1.0.2
ggrepel	0.9.6
ggridges	0.5.6
ggsci	3.2.0
ggsignif	0.6.4
ggthemes	5.1.0
gh	1.4.1
gitcreds	0.1.2
GlobalOptions	0.1.2
globals	0.16.3
glue	1.7.0
goftest	1.2-3
googledrive	2.1.1
googlesheets4	1.1.1
gower	1.0.1
GPfit	1.0-8
gplots	3.1.3.1
gridExtra	2.3
gridGraphics	0.5-1
gtable	0.3.5
gtools	3.9.5
hardhat	1.4.0
haven	2.5.4
HDF5Array	1.33.6
hdf5r	1.3.11
here	1.0.1
highr	0.11
hms	1.1.3
htmltools	0.5.8.1
htmlwidgets	1.6.4
httpuv	1.6.15
httr	1.4.7
httr2	1.0.4
ica	1.0-3
ids	1.0.1

igraph	2.0.3
ini	0.3.1
ipred	0.9-15
IRanges	2.39.2
irlba	2.3.5.1
isoband	0.2.7
iterators	1.0.14
janitor	2.2.0
jquerylib	0.1.4
jsonlite	1.8.9
kernlab	0.9-33
knitr	1.48
ks	1.14.3
labeling	0.4.3
lambda.r	1.2.4
later	1.3.2
latex2exp	0.9.6
lava	1.8.0
lazyeval	0.2.2
leiden	0.4.3.1
lhs	1.2.0
lifecycle	1.0.4
limma	3.61.9
listenv	0.9.1
lme4	1.1-35.5
lmtest	0.9-40
locfit	1.5-9.10
lubridate	1.9.3
magrittr	2.0.3
markdown	1.13
MatrixGenerics	1.17.0
MatrixModels	0.5-3
matrixStats	1.4.1
mclust	6.1.1
memoise	2.0.1
microbenchmark	1.5.0
mime	0.12
miniUI	0.1.1.1
minqa	1.2.8
modelenv	0.1.1
modelr	0.1.11
multicool	1.0.1
munsell	0.5.1
mvtnorm	1.3-1
Nebulosa	1.15.0
nloptr	2.1.1
numDeriv	2016.8-1.1
openssl	2.2.2

openxlsx	4.2.7.1
paletteer	1.6.0
parallelDist	0.2.6
parallelly	1.38.0
parsnip	1.2.1
patchwork	1.3.0
pbapply	1.7-2
pbkrtest	0.5.3
PCAtools	2.17.0
pillar	1.9.0
pkgbuild	1.4.4
pkgconfig	2.0.3
pkgdown	2.1.1
pkgload	1.4.0
plotly	4.10.4
plyr	1.8.9
png	0.1-8
polyclip	1.10-7
polynom	1.4-1
pracma	2.4.4
praise	1.0.0
presto	1.0.0
prettyunits	1.2.0
prismatic	1.1.2
processx	3.8.4
prodlim	2024.06.25
profvis	0.4.0
progress	1.2.3
progressr	0.14.0
promises	1.3.0
ps	1.8.0
purrr	1.0.2
quantreg	5.98
R.methodsS3	1.8.2
R.oo	1.26.0
R.utils	2.12.3
R6	2.5.1
ragg	1.3.3
RANN	2.6.2
rappdirs	0.3.3
rcmdcheck	1.4.0
RColorBrewer	1.1-3
Rcpp	1.0.13
RcppAnnoy	0.0.22
RcppArmadillo	14.0.2-1
RcppEigen	0.3.4.0.2
RcppGSL	0.3.13
RcppHNSW	0.6.0

RcppParallel	5.1.9
RcppProgress	0.4.2
RcppTOML	0.2.2
RcppZiggurat	0.1.6
RCurl	1.98-1.16
readr	2.1.5
readxl	1.4.3
recipes	1.1.0
rematch	2.0.0
rematch2	2.1.2
remotes	2.5.0
reprex	2.1.1
reshape2	1.4.4
reticulate	1.39.0
Rfast	2.1.0
rhdf5	2.49.0
rhdf5filters	1.17.0
Rhdf5lib	1.27.0
rjson	0.2.23
rlang	1.1.4
rmarkdown	2.28
ROCR	1.0-11
roxygen2	7.3.2
rprojroot	2.0.4
rsample	1.2.1
RSpectra	0.16-2
rstatix	0.7.2
rstudioapi	0.16.0
rsvd	1.0.5
Rtsne	0.17
rversions	2.1.2
rvest	1.0.4
S4Arrays	1.5.7
S4Vectors	0.43.2
sass	0.4.9
ScaledMatrix	1.13.0
scales	1.3.0
scattermore	1.2
scCustomize	2.1.2
SCpubr	2.0.2
sctransform	0.4.1
scuttle	1.15.4
selectr	0.4-2
sessioninfo	1.2.2
Seurat	5.1.0
SeuratData	0.2.2.9001
SeuratObject	5.0.2
SeuratWrappers	0.3.2

sfd	0.1.0
shape	1.4.6.1
shiny	1.9.1
SingleCellExperiment	1.27.2
sitmo	2.0.2
slider	0.3.1
snakecase	0.11.1
snow	0.4-4
sourcetools	0.1.7-1
sp	2.1-4
spam	2.10-0
SparseArray	1.5.37
SparseM	1.84-2
sparseMatrixStats	1.17.2
spatstat.data	3.1-2
spatstat.explore	3.3-2
spatstat.geom	3.3-3
spatstat.random	3.3-2
spatstat.sparse	3.1-0
spatstat.univar	3.0-1
spatstat.utils	3.1-0
SQUAREM	2021.1
statmod	1.5.0
stringi	1.8.4
stringr	1.5.1
SummarizedExperimer	1.35.1
sys	3.4.2
systemfonts	1.1.0
tensor	1.5
testthat	3.2.1.1
textshaping	0.4.0
tibble	3.2.1
tidyr	1.3.1
tidyselect	1.2.1
tidyverse	2.0.0
timechange	0.3.0
timeDate	4041.11
tinytex	0.53
tune	1.2.1
tzdb	0.4.0
UCSC.utils	1.1.0
umap	0.2.10.0
urlchecker	1.0.1
usethis	3.0.0
utf8	1.2.4
uuid	1.2-1
uwot	0.2.2
vctrs	0.6.5

VennDiagram	1.7.3
vipor	0.4.7
viridis	0.6.5
viridisLite	0.4.2
vroom	1.6.5
waldo	0.5.3
warp	0.2.1
whisker	0.4.1
withr	3.0.1
workflows	1.1.4
xfun	0.47
xml2	1.3.6
xopen	1.0.1
xtable	1.8-4
XVector	0.45.0
yaml	2.3.10
yardstick	1.3.1
yulab.utils	0.1.7
zeallot	0.1.0
zip	2.3.1
zlibbioc	1.51.1
zoo	1.8-12