Genomic factors contributing to the resilience of *Salmonella enterica* on ready-to-eat muskmelon

Irene Esteban-Cuesta[1]*, Laura Führer[1], Steffen Porwollik[2], Weiping Chu[2], Steven R. Fiddaman[3,4], Irmak Sah[1], Michael McClelland[2], Claudia Guldimann[1]

[1] Competence Center for Food Safety, Chair of Food Safety and Analytics, Veterinary Faculty, LMU Munich, Oberschleissheim, Germany

[2] Department of Microbiology and Molecular Genetics, School of Medicine, University of California, Irvine, California, United States of America

[3] The Pirbright Institute, Ash Rd, Pirbright, Woking, UK

[4] Department of Biology, University of Oxford, Oxford, UK

* Corresponding author: Irene.Esteban@ls.vetmed.uni-muenchen.de

Funding. This work was supported by the German Research Foundation (Deutsche Forschungsgemeinschaft, DFG). Grant No. 494875429.

Keywords: transposon insertion sequencing, salmonellosis, microbial contamination, transposon library, food safety

Short Title: Interaction mechanisms of *S. enterica* with ready-to-eat muskmelon

Abstract

Salmonella outbreaks have repeatedly been associated with muskmelons. This fruit poses a food safety risk due to its growth on the soil surface and because it is eaten raw. To identify genes under selection in *S. enterica* growing in this food matrix, barcoded transposon mutant libraries in three serovars of *Salmonella,* Typhimurium, Enteritidis, and Newport, were screened for survival and growth on muskmelon. The role of selected genes was subsequently verified in competition assays. Genes whose disruption caused a fitness effect in all three serovars were enriched for functions related to RNA degradation and ribosome biogenesis. The polyribonucleotide nucleotidyltransferase gene, *pnp,* and the D-3-phosphoglycerate dehydrogenase gene, *serA*, conferred an advantage when growing in muskmelon but not in a nutrition-rich control environment. This study provides a rare insight into genome-wide adaptation mechanisms of multiple *Salmonella* serovars to growth on food matrices, exemplified by muskmelons.

Introduction

Non-typhoidal salmonellosis is a zoonotic disease that has a high impact on the economy and human and animal health. It is among the most relevant foodborne pathogens in the European Union (EU) and worldwide [1].

Ready-to-eat (RTE) melons pose a high infection risk to consumers since pathogen inactivation measures are rarely applied before consumption, and handling may enable the spread of surface contaminants to the inside of the fruit. *S. enterica* foodborne outbreaks associated with the consumption of muskmelons have been reported since 1990 [2, 3], with recent cases in the EU (2023)[4] and the United States (2024)[5]. Especially, netted rind melon varieties offer a favorable niche for bacterial attachment and persistence [6].

To establish infection, *S. enterica* must be able to survive and grow on the food matrix. Quantitative data for the growth of *S. enterica* on muskmelons show that *S. enterica* can grow on muskmelons at 8°C and sustain its population at temperatures as low as 4°C [7-9].

S. enterica responses to food-related stresses have been studied in the past [10, 11] and previous studies revealed fitness effects of gene disruptions vary significantly by food matrix, identifying key survival mechanisms on tomatoes [12, 13], alfalfa sprouts [14], and low-water-activity foods, such as pistachios[15], black peppercorns, almonds, and hazelnuts [16]. Transposon insertion sequencing (TIS) showed matrix-specific effects for *Salmonella* colonization and persistence by the inactivation of genes linked to the alternative sigma factor σ^S (RpoS), DNA recombination, the biogenesis of lipopolysaccharides (LPS)[15], glycolysis, amino acid and LPS production, fatty acid degradation, and purine and pyrimidine biosynthesis pathways [13].

Based on the existing data for other food matrices, we hypothesized that distinct genetic determinants may exist for *S. enterica* fitness on muskmelon. In this study, we used barcoded transposon libraries of *S.* Typhimurium 14028s, *S.* Enteritidis PT4 strain P125109, and *S.* Newport

C4.2, to identify genome-wide fitness effects of specific genes and the corresponding metabolic

pathways involved in the interaction of these *S. enterica* serovars with RTE muskmelon.

Results

Growth data

Transposon libraries on the food matrix grew 1.0 to 2.0 $\log_{10}$ CFU/g, while *no matrix*

samples remained stable at 22°C and decreased slightly at 8°C (**Figure S1**). SEN libraries showed

reduced growth at 8°C compared to SNP and STM. The transposon libraries and their WTs

exhibited comparable growth on RTE muskmelons at 8°C and 22°C (**Figure S2**), except for the

SNP library, which showed a minor reduction compared to the WT (-0.3 $\log_{10}$ CFU/g, *p*-adj≤0.01)

at d_1, 8°C.

Mesophilic aerobic bacteria (MAB) and Enterobacteria on *Salmonella*-free RTE

muskmelon increased by approximately 4.0 $\log_{10}$ CFU/g at 8°C and 5.0-6.0 $\log_{10}$ CFU/g at 22°C,

with high variability among samples (**Figure S3**).

Transposon Insertion Sequencing data analysis

TIS screening of STM, SEN, and SNP libraries on RTE muskmelon identified functionally

grouped mutants with significant fitness effects. All fitness data of the transposon assay after

DESeq2 analysis are presented in **Table S2**, while the subset of genes that fulfilled the strict dual

requirement of significance, as detailed in Materials and Methods, are displayed in **Table 2.** The

dynamics of mutant representation during incubation are represented in **Figure S4**, while

exemplary volcano plots for inoculum/end of storage comparison are shown in **Figure 1**.

Table 1. Genes with a significant fitness effect in *Salmonella enterica* serovars when grown on muskmelon, classified by function.

Function	Gene Number	Gene	STM 8°C	STM 22°C	SEN 8°C	SEN 22°C	SNP 8°C	SNP 22°C	Comment
RNA Degradation	STM14_3962	*deaD*	-	-	-	-	(-)	(-)	
	STM14_0013	*dnaK*	n.d.	n.d.	+ / -				
	STM14_0218	*pcnB*						-	
	STM14_3964.RJ	*pnp*	-	-	-	-	-	-	
	STM14_3066	*ppk*	-		(-)		-		
	STM14_3965	SraG	(-)	-	(-)				
	STM14_5250	*vacB*	-				-		
Ribosome biogenesis	STM14_2082	*rluB*			(-)	-	-	(-)	
	STM14_1359	*rluC*			(-)		-		
	STM14_3263	*rluD*	(-)	(-)	(-)	(-)		-	
	STM14_3240	*srmB*	-	-	-	-	-	-	
	STM14_4639.P	STnc400	-	-	n.d.	n.d.	n.d.	n.d.	
	STM14_4822	*typA*	-	-	-	-	-	-	in SEN: *typA* represented by 50537.33.peg.4013STM14_4822
	STM14_5328+550537.33.peg.4414+108619.9.peg.3873	*yjgA*	-	-	-	-	-	-	in SEN and SNP: STM14_5328 unchanged
Reactive Oxygen Species (ROS) stress response	STM14_4959	*oxyR*	- / +	- / +	- / +	- / +	- / +		
	STM14_4878	*sodA*			- / +	(-)	n.d.	n.d.	
	STM14_4713	*trxA*	-	(-)	(-)		(-)	(-)	
	STM14_1080.J	*trxB*			(-)	(-)	-	(-)	
Lipopolysaccharides, O-antigen and enterobacterial common antigen biosynthesis	STM14_4308	*dcrB*	-				-		
	STM14_4479.J	*rfaI*			(-)		(-)	-	
	STM14_4483	*rfaG*					-	(-)	
	STM14_4484.J	*rfaQ*			(-)			-	
	STM14_4477	*rfaY*					(-)	-	
	STM14_4722	*wecD*	-	(+)	(-)				
	STM14_4723	*wecE*	+ / -	(+)	(-)		(-)	(-)	
	STM14_4725	*wecF*	-	(+)					
	STM14_4004	*yrbI*		(-)	(-)	(-)	(-)	-	
Envelope, membrane transport, and antimicrobial resistance	STM14_2841	*arnF*			(-)	-			
	STM14_5093	*dgkA*				-	(-)	(-)	
	STM14_0267	*hlpA*	-	-	(-)	-		-	
	STM14_0111	*surA*			-		(-)		
Two-component system	STM14_4216	*envZ*					+ / -	(-)	
	STM14_3140	*glnB*			-		n.d.	n.d.	
	STM14_4817	*glnG*	- / +	- / +	(-)	(-)		(-)	
	STM14_4818	*glnL*	(+)	(- / +)	(-)	- / +		(- / +)	
	STM14_2797	*ompC*					+ / -	(+)	
	STM14_4217	*ompR*					+ / -	(-)	
	STM14_1408	*phoQ*	(+)		- / +				
Protein folding, degradation, and secretion system	STM14_3114	*hscA*			(-)	-		(-)	
	STM14_3162	*lepB*					-	-	
	STM14_0162	*secA*	(-)		-		n.d.	n.d.	
	STM14_4461	*secB*			-	-	(-)	-	
	STM14_4780+108619.9.peg.3424	*tatB*			(-)		-	(-)	in SNP: STM14_4780 nearly unchanged
Amino acid metabolism	STM14_1106	*aroA*					-	(-)	
	STM14_2232	*prc*				-			
	STM14_3699	*serA*		(-)		-		-	
	STM14_5499	*serB*		(-)		(-)		-	
	STM14_1105	*serC*		-	(-)	(-)		-	
Translation factors	STM14_2145	*hemK*			(-)	-			

	Locus tag	Gene							Notes
	STM14_5478	*prfC*	-		-	(-)	(-)		
tRNA Biogenesis	STM14_3116	*iscA*			(-)	-	(-)	(-)	
	STM14_3604	*metZ*	(-)				-		
	STM14_2196	*rnd*	-		(-)				
	STM14_0583	*ybaK*		(-)	(-)	-	(-)	(-)	
Mismatch repair	STM14_5372.J	*holC*	-						
	STM14_5475	*holD*	-	-			n.d.	n.d.	
	STM14_3077	*xseA*	-	-					
DNA replication and recombination	STM14_3608	*recD*					(-)	-	
	STM14_0566	*priC*					-		
Cytoskeleton and cell division	STM14_0159	*ftsZ*					-	(-)	
	STM14_3095	*yfgA*			(-)		-	-	
Transcription and transcriptional regulators	STM14_4399	*cspA*					-	(-)	in SNP: *cspA* represented by STM14_4399STM14_4399.J
	STM14_4991	*rpoC*	(-)			-	(-)		
	STM14_2310	*yebC*				-			
Purine and pyrimidine metabolism	STM14_3076.LR	*guaB*				-			
	STM14_5248	*purA*	(-)	(-)	(-)	-		(-)	
	STM14_3050	*purC*		(-)		-			
	STM14_5017	*purH*		(-)		-		(-)	
	STM14_0623	*purK*				-			
	STM14_1200	*pyrD*				(-)		-	
Phosphotransferase system	STM14_2990	*ptsI*			(-)	(-)	-	(-)	
Carbohydrate metabolism and Pentose phosphate pathways	STM14_2289	*eda*		-	(-)	-	n.d.	n.d.	
	STM14_2118	*galU*	n.d.	n.d.			-	(-)	
	STM14_4444	*mtlD*				- / +		(-)	
	STM14_3124	*suhB*	n.d.	n.d.	n.d.	n.d.	-	-	in SNP: *suhB* represented by STM14_3124STM14_3123.J
Bacterial defense and phage	STM14_3172	Phage genes					+ / -	(-)	
	STM14_1150.JSTM14_3224	Phage genes			n.d.	n.d.	+ / -	+ / -	
	STM14_1193	*sseI*					+ / -	(+ / -)	
	STM14_0340.J	putative RHS-like protein	(-)	-	n.d.	n.d.	n.d.	n.d.	
Uncharacterized proteins	STM14_2205	*yoaH*	-				(-)		
	STM14_4069	hypothetical protein					-		
	STM14_3096.J	hypothetical protein					-		
	STM14_2704	hypothetical protein						-	
	STM14_1197	hypothetical protein				(-)	+ / -	(+ / -)	
	108619.9.peg.3873	FIG138315 - putative alpha-helix protein	n.d.	n.d.	n.d.	n.d.	-	-	
	STM14_1530	putative cytoplasmatic protein			n.d.	n.d.	-		
	STM14_1076	putative inner membrane protein	(-)		(-)		-		
	STM14_1194	Pseudogene					+ / -	(-)	

Mutants listed met two criteria: (i) a two-fold change (i.e., $\log_2$-fc>|1.0|) with a p_{adj}≤0.01 for the matrix

versus *no matrix* comparison, AND (ii) a two-fold change with a p_{adj}≤0.01 for the inoculum versus matrix

comparison at the same sampling time point.

-, conditions for which a statistically significant negative fitness effect was observed.

+/-, positive fitness effect at the first sampling time point, followed by a negative fitness effect later.

-/+, negative fitness effect at the first sampling time point, followed by a positive fitness effect later.

(), mutants with a two-fold fitness effect at one OR the other criterion of the dual requirement, at p_{adj}≤0.1.

n.d., absence of mutants in the transposon library.

RHS - rearrangement hotspot.

Figure 1. Change in the mutant abundances after growing the random transposon libraries on RTE muskmelons. Volcano plots depict the fold change of relative abundance of mutants and their adjusted p-values between inoculum and growth on muskmelon for 24 h at 22˚C for S. Typhimurium 14028s (**A**), S. Enteritidis PT4 strain P125109 (**B**), and S. Newport C4.2 (**C**); and the growth on muskmelon for 4 days at 8˚C for S. Typhimurium 14028s (**D**), S. Enteritidis PT4 strain P125109 (**E**), and S. Newport C4.2 (**F**). Genes with significantly changed representation (at least 2-fold, p ≤ 0.01) are labelled in blue. Intergenic regions (IRs) with a significant fitness effect were excluded from the graph. Plots showing changes for all other sampled time points are shown in **Supplementary Figure S4**.

Three core genes (*pnp, typA,* and *srmB*) contributed to fitness in all three serovars at both

temperatures, while *hlpA* and *oxyR* mutants were found to be underrepresented in all three serovars

at 22°C. For STM, mutants in 24 genes showed a significant fitness effect on RTE muskmelon,

including 21 genes at 8°C and 15 genes at 22°C, with 11 overlapping. Most mutations had a

negative fitness effect, except for *oxyR* and *glnG,* that switched from early negative to late positive

fitness effects, and *wecE,* which showed the opposite trend. In SEN, mutants in 32 genes exhibited

a significant fitness on RTE muskmelon. At 8°C, 14 genes were identified, while at 22°C, mutants

in 26 genes showed fitness effects. Mutants in seven of these genes were significantly affected in

both temperatures (*pnp, deaD, yjgA, srmB, typA, secB, oxyR)*. In SNP, mutations in 24 genes

affected fitness on RTE muskmelon at 22°C, while 35 genes were implicated at 8°C. Temperature-

independent fitness effects were observed for eight of these genes (*pnp, srmB, typA, yfgA, yjgA,*

suhB, lepB, and FIG138315; **Table 2**).

In all three serovars, mutants in RNA degradation genes, ribosome biogenesis, envelope

and membrane transport, and tRNA biosynthesis showed a statistically significant difference after

growth on muskmelon. For some categories displayed in **Table 2**, statistical significance, as

defined in our criteria, was only observed for genes in one or two serovars. As an example, mutants

in genes of the purine and pyrimidine pathways were statistically significantly affected in growth

in SEN, but not in STM or SNP, and cytoskeleton and cell division gene mutants cleared the

significance threshold in SNP but not in the other two serovars (**Table 2**). However, the number

of transposon mutants in each gene and their abundance in the library differ among strains, so

these differences should not be misinterpreted as confirmation of a difference in the role of that

pathway between strains.

We performed KEGG enrichment analyses to gain a better insight into pathways significant

for the growth of *Salmonella* on muskmelon. Significantly enriched KEGG pathways ($q \leq 0.01$) and

associated GO terms ($p \leq 0.01$) are summarized for all three serovars in **Table 3**, with full results

in **Table S3.**

Table 2. Significantly enriched KEGG pathways and associated GO terms for *S.* Typhimurium 14028,

S. Enteritidis P125109, and *S.* Newport C4.2 on RTE muskmelon.

KEGG Pathway	Serovar	Temp.	*q*-value	Selected Genes	Associated GO Terms
Cellular Processes Cell motility	STM	8°C	2.5 * E-10	*flgBDFGI, flhAB, fliCFHIQR*	GO:0044780, GO:0040011, GO:0051674, GO:0071978, GO:0044781, GO:0071973
Flagellar assembly	SNP		0.005	*flgABCDEK*	GO:0040011, GO:0071978, GO:0051674
Genetic Information Processing	STM	8°C	0.004	***deaD, pnp, ppk, vacB***	GO:0006402, GO:0010501
Folding, sorting, and degradation	SNP		0.001	***deaD, pcnB, pnp, ppk, vacB***	GO:0010501, GO:0000455
RNA degradation	STM	22°C	0.007	***deaD, pcnB, pnp, vacB***	GO:0010501, GO:0006402
Genetic Information Processing Translation Ribosome	STM	22°C	0.007	***rrlABCDEGH, rrsG***	-
Genetic information processing Folding, sorting and degradation Protein export	SNP	8°C	0.001	***lepB, secB, tatABC***	GO:0043953, GO:0065002
Metabolism Glycan biosynthesis and metabolism Biosynthesis of nucleotide sugars	STM	8°C	0.004	***rfbK,*** <u>*wecBC,*</u> ***wecDE***	GO:0009246
Metabolism Nucleotide metabolism Purine metabolism	SEN	22°C	0.001	***guaB, purACDEGHK***	GO:0006189, GO:0046040

Gene lists were based on $p_{adj}<0.05$ in at least one matrix/*no matrix* and inoculum/timepoint comparison. Genes in **bold** were under negative selection between sampling time points, while <u>underlined</u> genes were positively selected.

Competition assays confirm genes with a fitness effect

A total of 25 genes with a fitness effect in STM, SEN, or SNP on RTE muskmelon were selected from the *S.* Typhimurium 14028 SGD mutant collection [17] for competition assays (**Table 4)**. Four of these genes (*serA*, *serB, ybaK*, and *purA*) displayed changes in STM that were not as significant as in SEN or SNP but were tested to verify whether the results of the transposon assays were, in fact, a representation of the true diversity between *Salmonella* serovars. Two genes that exhibited temporal abundance changes in STM, *oxyR* and *glnG*, were also included in these competition assays, along with *rfbI_2,* which showed no significant changes in any of the three strains analyzed. A final gene, *wecF [D],* showed opposite effects in the two different incubation temperatures in STM. However, significance ($p_{adj}<0.01$) was only achieved at 8˚C, while at 22°C only the inoculum/t_{24} comparison was significant.

For 18 of these 25 genes, we had two single-gene deletion mutants available for each gene, one where a kanamycin resistance (Kan^R) cassette replaced the gene in the sense orientation and one where a chloramphenicol cassette (Cm^R) replaced the gene in the antisense orientation. For

five genes (*yjgA, hlpA, holC, purA,* and *holD*), we only had a Kan^R mutant available, while for

wecE, our collection only harbored a Cm^R mutant. Competition assays were performed at the

temperature where transposon data suggested a stronger fitness effect. Assays were performed on

muskmelon and rich LB media, to verify whether any fitness effects occurred in any rich nutrient

environment or were potentially specific to growth on muskmelon. For selected genes, the fitness

effect observed in muskmelon or LB was also tested in PBS, where the nutrient-restricted

conditions only confer a growth difference due to differential death. The results are summarized

in **Table 4**. To confirm a negative fitness effect, both SGD mutants were required to display a

competition index where CI+STDEV<1. In contrast, for a confirmed positive fitness effect, both

mutants needed to adhere to the CI–STDEV>1 threshold. If only one SGD mutant was available,

an expected single mutant behavior as outlined above was sufficient to confirm the phenotype

observed in the transposon assay.

As shown in **Table 4**, the observed fitness effects in the transposon assays were confirmed

for 15/24 genes. Notably, *serA* and *pnp* deletions showed a negative fitness effect during the

competition with the WT on muskmelon that was not detected in LB. Similarly, the positive fitness

effect of transposon insertions in *glnG*, observed in STM at 8°C on d$_5$, was confirmed in

competition assays on muskmelon and not on LB, although with high standard deviations. Further

replicates would be needed to illuminate this difference further. Mutants in *xseA* exhibited a strong

polar effect in LB and RTE muskmelon, where one single-gene deletion mutant confirmed the

expected debilitated phenotype on muskmelon, while the other suggested a growth advantage. This

may be due to the different orientations of the constitutive promoter introduced into the gene

during the generation of the two mutant constructs. For four genes (*wecD, trxA, typA*, and *serB*),

one SGD mutant confirmed the expected phenotype while the other showed no significant change.

Discussion

This study investigated *S. enterica* fitness determinants on RTE muskmelon using TIS in three serovars: *S.* Typhimurium ATCC 14028, *S.* Enteritidis PT4 strain P125109, and *S.* Newport C4.2. The growth potential was comparable to previous studies with *S. enterica* on muskmelons [7, 8, 18], with minor variations probably related to strain differences, processing conditions, and time-temperature combinations. The slight reduction in SNP growth at d_1 (8°C) may be due to phage lysis under stress, as phage genes were negatively selected at this temperature.

Limited data is available on the background microbiota in muskmelons and their growth during storage. Compared to our data, higher initial MAB counts (3.5-4.0 $\log_{10}$ CFU/g total aerobic bacteria compared to ~2.0 $\log_{10}$ CFU/g in our study) and coliforms (slightly below 2.0 $\log_{10}$ CFU/g compared to 0.5-1.0 $\log_{10}$ CFU/g *Enterobacteriaceae* in our study) were reported in previous studies [7, 19].

TIS revealed shared fitness determinants across all three serovars, particularly genes involved in RNA metabolism, ribosome and amino acid biogenesis, oxidative stress response, and cell integrity and membrane transport. Serovar-specific fitness effects were observed for mutations in genes related to protein folding and transport, purine and pyrimidine metabolism, DNA replication, recombination, and mismatch repair. Although these differences have not been further analyzed and may be due to differences in the number of transposon mutants in each gene and their abundance in the library, this approach allowed us to spread a wider net, capturing genes under selection that would have been missed with only one TIS library. Future work may analyze single-gene mutant orthologs to elucidate whether candidate differences between serovars are indeed differences.

Mutations in genes involved in **RNA degradation** (*pnp*, *deaD*, *vacB,* SraG*, ppk, dnaK, pcnB)* **and ribosomal homeostasis** (*typA, srmB, yigA, rluBCD, STnc400*) had significant fitness effects across serovars.

PNPase (exoribonuclease polynucleotide phosphorylase (*pnp*)), a conserved RNA

degradosome component, regulates mRNA degradation, tRNA processing, and small RNA

turnover [20]. PNPase is critical for bacterial cold-shock responses, although *S. enterica pnp*

mutants exhibit milder cold sensitivity, consistent with its temperature-independent fitness effect

on muskmelon. Its relevant role during the interaction on muskmelon was confirmed with a

muskmelon-specific negative fitness effect absent in nutrient-rich media in the competition

experiments against the WT.

DeaD and SrmB are DEAD-box RNA helicases [21, 22]. Both *srmB* and *deaD* mutants

exhibit cold-sensitive growth defects in *E. coli* [21]*, Pseudomonas syringae* [23], and *Yersinia*

pseudotuberculosis [24], consistent with our findings, although the effect was observed at both

temperatures. Beyond cold stress, DeaD and SrmB participate in post-transcriptional regulation,

including the carbon storage regulatory system in *E. coli* [22], which may explain their

temperature-independent fitness effects. Co-transcription of *deaD* and *pnp* occurs in certain

bacterial species but is absent in others [25, 26]. This is mirrored in our data, where *deaD*

disruption affected fitness only in STM and SEN but not SNP. This phenotype was confirmed on

both muskmelon and nutrient-rich media in the competition experiments, and is therefore not

melon-specific.

PPK (Polyphosphate kinase) is highly conserved across bacteria. It catalyzes inorganic

polyphosphate (polyP) synthesis, regulates RNA degradation and mRNA biogenesis [27], and

contributes to *S. enterica* growth, survival, and virulence [28]. PolyP, a high-energy reservoir,

plays a key role in STM and SNP muskmelon interactions but shows no fitness effect in SEN.

Serovar-specific differences in *ppk* function were previously observed [28], with *ppk* mutants

exhibiting a greater growth defect in *S.* Typhimurium than *S.* Gallinarum, suggesting variations in

energy homeostasis or alternative energy use. PolyP synthesis occurs mainly in stationary phase

[28], likely balancing metabolism by regenerating ATP through the removal of inhibitory polyP

and nucleotide diphosphates [29]. On muskmelon, *ppk* mutants displayed a fitness effect only at

8°C and later time points, indicating polyP accumulation is critical under prolonged cold stress, possibly to maintain mRNA degradation homeostasis, a phenotype confirmed for muskmelon and nutrient-rich media. Nevertheless, polyP is also known to be essential for *S. enterica* long-term survival in LB, even at 37°C [30]. Additionally, *vacB* mutants (exoribonuclease R), deficient in RNA degradation and processing as well as in ribosome biogenesis [31], exhibited a fitness effect under the same conditions as Δ*ppk* mutants. Taken together, these findings indicate the importance of RNA and ribosomal homeostasis in *S. enterica* environmental adaptation.

In the presence of **reactive oxygen species (ROS)**, OxyR acts as key regulator of oxidative stress. Δ*oxyR* mutants exhibited a dynamic fitness effect that was initially negative but shifted to a strong positive selection over time in SEN and STM at both temperatures and in SNP at 22°C, suggesting *oxyR* disruption initially impairs bacterial adaptation but later provides a selective advantage. Members of the *oxyR*-regulon (e.g., *trxAB*)[32] also showed a negative fitness effect in STM and SNP at 8°C, reinforcing the role of oxidative stress responses in muskmelon adaptation. The competition assays mirrored the transposon results, albeit with high variability, further emphasizing the complex environmental regulation of *oxyR*. While plants generally generate ROS as a defense response [33], cut muskmelon pieces are unlikely to retain this response. Thus, the *oxyR*-effect may relate to alternative metabolic adaptations [34, 35] that shall be further investigated in future studies.

Cell wall integrity and **membrane transport** components were also identified. Several genes involved in LPS and membrane biosynthesis (*rfaG, dcrB*) and enterobacterial common antigen (ECA) biosynthesis (*rfaG, wecDEF, dcrB*) exhibited fitness effects, particularly at 8°C, although for SNP, the negative fitness effect was prevalent at 22°C. While the fitness effect at 8°C may represent an adaptation to altered membrane fluidity at cold temperatures [36], a special need for the membrane LPS or specifically for the ECA in SNP during its interaction with muskmelon at room temperature may be of importance. LPS biosynthesis was shown to be important for *S.*

enterica survival on pistachios [15], tomatoes, and mouse colonization [12], reinforcing their

broad role in environmental adaptation.

Disruptions in membrane transport (*secAB, lepB, hlpA, hscA, surA, tatB*) impaired bacterial

fitness on muskmelon, indicating that efficient protein translocation is critical for survival. The

chaperone SecB, required for protein targeting to the SecYEG transport system, was particularly

important in SEN and SNP. The outer membrane protein HlpA also exhibited a fitness effect at

22°C, although competition assays did not confirm this phenotype. Nevertheless, the enrichment

of membrane transport genes suggests that maintaining protein export balance is essential; either

the accumulation of metabolites in the cytoplasm or the depletion of functional proteins in the

periplasm may be detrimental for *S. enterica* growth on muskmelons.

Two-component systems (TCSs) and **motility** genes were also overrepresented in the TIS

data. Mutants in various TCSs showed fitness effects at 8°C, suggesting a role under cold stress.

Nitrogen assimilation TCS mutants (*glnG* in STM, *glnBL* in SEN) showed significant fitness

effects, while *phoQ* mutants (sensing antimicrobial substances and Mg^{2+} starvation) appeared to

be affected only in SEN. In SNP, osmolarity regulators (*envZ, ompR, ompC*) exhibited fitness

effects. Positive selection for *glnG, phoQ, envZ,* and *ompR* mutants during part of the interaction

period suggests a fitness advantage.

Positive selection was also observed for mutations in flagellar assembly genes, likely

aiding adaptation to the plant environment [37]. This correlates with the results of a previous study,

where *S.* Typhimurium 14028 largely downregulated flagellin production, while a subset of cells

exhibited high flagellin expression inside the plant [38].

Protein synthesis was relevant for the interaction of *S. enterica* with muskmelons, as

inactivation of genes involved in the serine (*serABC, prc)*, aromatic amino acids (*aroA),* and

glutamine metabolism (*glnB)* resulted in serovar- and temperature-dependent fitness effects.

Additionally, genes related to tRNA (*rnd, iscA, ybaK, metZ*) and translational regulation (*prfC,*

hemK) were also identified. Both *serA* and *glnB* were previously identified as necessary for tomato

colonization [12], suggesting a critical role for scavenging serine and glutamine from the food

matrix. The positive fitness effect observed at later time points for *glnG* mutants, a gene coding

for the nitrogen regulation protein NR(I), was confirmed in the competition assays to be

muskmelon-specific, suggesting that a switch to alternative nutrient sources or alternative

regulation mechanisms of the nitrogen metabolism may be beneficial for growth on muskmelon.

Perhaps nitrogen assimilation is unnecessary for the fitness of *S. enterica* on muskmelons.

ΔaroA mutants have demonstrated a fitness effect in SNP and SEN under desiccation stress

on low-moisture foods [15]. Since *aroA* mutants only showed an effect at 8°C, a role in cold stress

could be hypothesized, with a potential link to TypA as a tyrosine-based enzyme with a role in

cold-shock response [39]. A growth defect for *E. coli met*-mutants was previously demonstrated

[40, 41]. While the *de novo* biosynthesis of amino acids was also found to be critical for the

persistence of *S. enterica* on tomatoes, *S. enterica* appeared to prioritize amino acid scavenging

during animal infections [12, 13].

Mutants in genes with a role in **purine metabolism** were strongly repressed in SEN at

22°C. Notably, *ΔpurA* exhibited nearly four $\log_2$-fc reductions, suggesting *purA* may be essential

for growth. The purine biosynthesis was previously implicated in *S.* Typhimurium colonization of

tomatoes [12, 13], and competition experiments confirmed a negative fitness effect for STM

Δ*purA* mutants on ready-to-eat muskmelon and nutrient-rich media.

The *pyrD* transposon mutants, deficient for this gene involved in the *de novo* pyrimidine

metabolism and linked to both purine and pyrimidine *de novo* synthesis [42, 43], exhibited a fitness

effect in SNP. Pyrimidine metabolism is required by *S.* Typhimurium infection in animals [12,

44], and proliferation in tomatoes [12]. However, our study found fitness effects only in SNP,

although associated GO terms were also enriched for SEN and SNP. This might suggest an ability

to scavenge pyrimidines from the environment for certain *S. enterica* strains [13].

Mutants in genes related to **Fe-S-cluster biosynthesis** showed fitness defects in SEN and

SNP. These clusters play essential roles in electron transfer, substrate binding, Fe-S storage, gene

expression, and enzyme activity [45]. Disruption in *hscA* and *iscA*, involved in Fe-S cluster

assembly [46, 47], as well as *aroA,* required for siderophore synthesis and iron uptake, were

detrimental during muskmelon interaction. Similar iron uptake genes were previously linked to *S.*

enterica proliferation in tomatoes [48], and *iscA* was relevant during alfalfa sprouts' colonization

[14], suggesting a shared iron requirement for fresh produce colonization.

Our analyses during the competition assays revealed that the fitness effects observed during

the interaction of several STM mutants with muskmelon were similar to those observed in LB

medium, suggesting that muskmelon provides conditions resembling a rich nutrient medium for

S. enterica.

An important limitation of our study was the high inoculation loads required to maintain

the complexity of the transposon library, which was mitigated to some extent by conducting direct

competition assays between individual candidate mutants and the WT. Future studies could focus

on individual genes, allowing lower inoculation levels and a more precise assessment of growth

potential. A further limitation is that several genes identified in the transposon screen were either

unavailable as single mutants in the collection (*serC, secAB, vacB)* or not confirmed in competition

assays (*hlpA, serB, ybaK)*. The latter may be due to sensitivity differences between the two

methods: TIS detects subtle fitness effects within a mutant library, whereas competition assays

measure fitness relative to the WT, potentially overlooking small but biologically relevant

differences. Moreover, hitchhiking mutations within the mutant pool may obscure individual gene

contributions, as complete mutant purity cannot always be ensured.

This study identified key fitness determinants enabling *S. enterica* survival on muskmelon,

revealing both conserved and serovar-specific adaptation mechanisms. RNA metabolism, *de novo*

amino acid and ribosome biogenesis, oxidative stress response, and cell wall integrity emerged as

central survival factors, with additional metabolic adaptations influencing fitness under specific

conditions. Many of these mechanisms align with known plant-endophyte interaction pathways,

including ROS detoxification, nitrogen fixation, flagella, and cell wall components such as LPS,

surface-associated proteins, or transporters [49]. Future studies should investigate the metabolic interplay between *S. enterica* and plant-derived nutrients to further elucidate bacterial survival strategies in fresh-produce environments.

Material and Methods

Bacterial strains and culture media

S. enterica wild type (WT) strains used in this study and their sources are listed in **Table 1**. All strains were preserved at −80°C in Luria-Bertani broth (LB; Carl Roth GmbH & Co. KG, Karlsruhe, Germany) with 20% glycerol (Th. Geyer GmbH & Co. KG, Renningen, Germany). Strains were grown in LB broth or 1.5% LB agar (Oxoid Deutschland GmbH, Wesel, Germany). Where necessary, culture media were supplemented with 60 µg/ml kanamycin (LB^{kan}; Carl Roth), 20 µg/ml chloramphenicol (LB^{cm}, Carl Roth) or 15 µg/ml tetracycline (LB^{tet}; Carl Roth).

Table 3. *Salmonella enterica* wild type strains used for the barcoded transposon libraries.

Strain	Abbreviation	Source
Salmonella Typhimurium strain ATCC 14028	STM	Isolated from 4-week-old chickens in 1960 [50] NCBI accession no. CP001363.1/ CP001362.2
Salmonella Enteritidis PT4 strain P125109	SEN	Clinical strain associated with poultry outbreak in the United Kingdom [51] NCBI accession no. CP063700.1/ CP063701.1
Salmonella Newport C4.2	SNP	Isolated from a tomato field in Virginia, United States [13] NCBI accession no. GCA_006821125.1

Library construction

The construction of the barcoded Tn*5*-based mutant libraries has been previously described [12, 13]. Briefly, a library of Tn*5* insertion mutants was constructed for each strain with a mini-Tn*5* derivative into which an N$_{18}$ random barcode was inserted via PCR. This construct was randomly introduced into the genome with the EZ-Tn*5* <T7/KAN-2> promoter insertion kit

(Epicentre Biotechnologies, Madison, WI, United States). Transformed cells were recovered on

LB^{kan} agar after overnight culture at 37°C.

The methods for mapping the barcoded transposons to specific locations in the genome

have also been previously described [12]. The *S.* Typhimurium ATCC 14028 transposon mutant

library consisted of about 230,000 insertion mutants [12], the *S.* Enteritidis PT4 strain P125109

transposon mutant library consisted of about 140,000 insertion mutants [16], and the *S.* Newport

C4.2 transposon mutant library consisted of about 80,000 insertion mutants [13].

Comparative growth analysis

353 Growth curves on RTE muskmelon for all three barcoded transposon libraries were

354 assessed (eight replicates) and compared to those of the WT (five replicates) at 8°C and 22°C.

355 Inoculation and sampling were performed as described for the screenings below. The growth data

356 was subjected to an ANOVA test, where an adjusted *p*-value below 0.05 was considered

357 significant.

Library screening on RTE muskmelon

For screenings on RTE muskmelon, 300 µl frozen stocks of the *S. enterica* barcoded mutant

libraries were thawed and grown in LB^{kan} (Carl Roth; 60 µg/ml) at 37 °C and 200 rpm until OD$_{600}$

1.0 was reached. One ml of this culture was diluted 1:10 in phosphate-buffered saline (PBS; Carl

Roth) to reach approximately 8.0 log$_{10}$ CFU/ml, centrifuged at 4.500 rcf for 5 min, the supernatant

discarded, and the pellet resuspended in 5 ml PBS to create the inoculum.

Galia muskmelons (*Cucumis melo* L. *reticulatus*) were purchased at local grocery stores.

For inoculation, muskmelon cubes were cut with sterile knives, and 10 g were weighed into sterile

filter Stomacher® bags (0.4 L, Avantor VWR International GmbH, Darmstadt, Germany). *No*

matrix samples containing only the inoculum without the muskmelon were included as negative

controls. RTE muskmelon samples and *no matrix* samples were inoculated with 5 ml of the

inoculum to reach an initial concentration of approximately 7.0 $\log_{10}$ CFU/g or ml to ensure that

the complexity of the library was maintained. The inoculum was evenly distributed over the

muskmelon samples by manually massaging the bags, ensuring the muskmelon cubes were

completely covered. Samples were incubated at 22°C and 8°C for 24 h and 5 days, respectively.

No macroscopic changes were observed in the appearance of the muskmelon cubes during the

incubation period.

Additionally, uninoculated RTE muskmelon samples were examined for total mesophilic

aerobic bacteria (MAB; ISO 4833-2:2013-12) and *Enterobacteriaceae* (ISO 21528-2:2004;

anaerobic incubation: O_2 <0.1%, CO_2 7.0-15.0%) at every sampling time point. The absence of

pre-existing *Salmonella* spp. in the purchased melons was confirmed by qualitative

microbiological analysis according to ISO 6579-1:2017.

Mutant libraries were recovered after 1 h (t_1), 7 h (t_7) and 24 h (t_{24}) for experiments at 22°C,

and after 1 h (d_1), 48 h (d_3) and 96 h (d_5). For this, the full RTE muskmelon and *no matrix* samples

were transferred into 90 ml LB^{kan} (Carl Roth) in 100 ml-Erlenmeyer flasks and incubated at 37°C

and 200 rpm for 30 min to detach the *Salmonella* from the matrix. After this, *Salmonella* spp.

populations were quantitatively assessed by preparing serial dilutions in PBS and plating on XLT-

4 agar (Oxoid) that was then incubated at 37°C for 24 h. The RTE muskmelon samples in 90 ml

LB^{kan} were then centrifuged at 4.500 rcf for 5 min to eliminate muskmelon juice residues, and the

pellet was resuspended in 90 ml LB^{kan}. Samples were then incubated for a further 7.5 h at 37°C

and 200 rpm before genomic DNA extraction for subsequent sequencing. Five biological

replicates were performed.

Sequencing library preparation and data analysis

Aliquots of 40 µl from each sample were washed three times in sterile water, pelleted and

then lysed with proteinase K (Sigma-Aldrich; 100 µg/ml) in 20 µl 10 mM Tris [pH 8.0] with 1 mM

EDTA (Carl Roth) and 0.1% Triton X-100 (Carl Roth) for 2 h at 55°C. After enzyme inactivation

at 95°C for 10 min, 5 µl of extracts were subjected to PCR with indexing primers targeted to the regions directly adjacent to the barcode on the transposon (0.4 µM Illumina-barcoded Primers L and V, listed in **Table S1**). The PCR was carried out using the following thermal cycling conditions: an initial denaturation step at 98°C for 2 min, followed by 10 cycles of denaturation at 98°C for 10 seconds, annealing at 65°C for 10 seconds, and extension at 72°C for 20 seconds. This was followed by an additional 20 cycles consisting of denaturation at 98°C for 10 seconds and extension at 72°C for 20 seconds. A final extension was performed at 72°C for 2 minutes. The reaction was subsequently held at 20°C. The Q5™ Hot Start High-Fidelity 2x Master Mix (Item number M0494L; New England Biolabs, Frankfurt am Main, Germany) was added at a final concentration of 1x. Equal volumes of PCR products were pooled and purified using the QIAquick PCR Purification Kit (Qiagen, Item number 28106), following the manufacturer's instructions. Illumina sequencing was performed as a dual-indexed paired-end 50, 100 or 150-base run on a NovaSeq 6000 with at least one million reads/sample.

Raw reads were binned based on their indexes and analyzed using custom Python scripts to identify and enumerate barcodes bordered by the expected invariable DNA sequence. Barcode insertions were compiled into an aggregated count for each disrupted gene. The differences in the aggregated abundances of the different mutants between time points, as well as between the matrix and *no matrix* samples for each sampling time point, were statistically analyzed using DESeq2 [52] and $\log_2$-fold changes (fc) were reported (**Table S2**).

Mutations were considered to have a significant fitness effect for *S. enterica* on muskmelons if they met a dual requirement for at least one sampling time point: (i) they showed a two-fold change (i.e., $\log_2$-fc $>|1.0|$) with an adjusted *p*-value ≤ 0.01 in the comparison of the matrix versus *no matrix* samples, and (ii) fulfilled the same criteria for a comparison between the inoculum and the matrix sample at the same sampling time point. If criterion (i) was achieved at the final timepoint (t_{24} or d_5), criterion (ii) was also deemed fulfilled if in comparisons the first timepoint (t_1 or d_1) was used instead of the inoculum (i.e. t_1 matrix versus t_{24} matrix; d_1 matrix

versus d_5 matrix), reflecting interaction mechanisms *after* the adaptation to the matrix. This strict

set of criteria was implemented to identify the genes with the most significant role during bacterial

growth on muskmelon.

To generate gene lists for the KEGG [53]- and GO [54-56] analysis input, the above p-

value criterion of $p_{adj} \leq 0.01$ was relaxed to $p_{adj} < 0.05$. GO enrichment analysis was carried out using

TopGO [57] and KEGG enrichment analysis used enrichKEGG from ClusterProfiler [58] in R

(v.4.4.1). For KEGG results, pathways with q-values≤ 0.01 were considered significant. Resulting

GO terms with classic Fisher p-values≤ 0.01 were considered significant.

Competition assays

To confirm the phenotype of candidate genes identified using TIS, single-gene deletion

(SGD) mutants in *S*. Typhimurium 14028s were used in direct competition with the WT strain.

These mutants were obtained from an existing SGD library [17] in STM that includes two mutants

for a majority of all genes, one mutant harboring a kanamycin resistance gene in the sense direction

of the deleted gene (SGD-Kan^R) and one mutant harboring a chloramphenicol resistance gene in

the antisense direction (SGD-Cm^R). The WT (*S*. Typhimurium 14028s) used in the competition

experiments was isogenic except for a tetracycline resistance cassette inserted in the neutral *malXY*

genome location (WT$_{STM}$-Tet^R).

SGD mutants and the WT$_{STM}$-Tet^R were grown in 5 ml LB^{kan}, LB^{cm}, and LB^{tet}, respectively,

at 37°C with 200 rpm shaking until approximately OD$_{600}$ 0.9 was reached. For each selected gene,

the different SGD cultures were mixed in a 1:1:1 ratio (or 1:1 if only one SGD mutant was available

in the collection). The SGD-WT$_{STM}$-Tet^R mix was then pelleted, resuspended in PBS, and 5 ml

were inoculated onto 10 g RTE muskmelon at a concentration of approximately 7.0 log$_{10}$ CFU/g

and screened as previously described. Post-screening, samples were plated on LB^{kan}, LB^{cm}, and

LB^{tet} agar and incubated at 37°C for 24 h to assess changes in population ratios. All mutants were

similarly tested in competition assays in LB broth to investigate whether any genes played a role

specific to growth on muskmelon or whether their role was universal in a nutrient-rich environment. Furthermore, some mutants were also tested in competition assays in PBS to confirm lack of differential death in the absence of nutrients. Competition assays in PBS and LB were inoculated at approximately 7.0 $\log_{10}$ CFU/ml. All cultures were tested for cross-resistance before being combined into a mixed inoculum.

The mutant phenotype was confirmed by comparing the changes in colony counts between samplings in the respective sense and antisense mutants and the WT_{STM}-Tet^R. If the competitive index (the ratio of mutant/WT ratios of timepoint/inoculum) plus standard deviation from at least three experiments remained below 1, a negative fitness effect was deemed confirmed. If the competitive index minus standard deviation remained higher than 1, a positive fitness effect was deemed confirmed. Competition assays were performed in triplicate, except when no fitness effect was observed in the first two replicates.

Table 4. Fitness effect in selected *S.* Typhimurium 14028s gene deletion mutants in competition with the wild type.

| Condition | STM Gene Number | Gene | Gene Function | Transposon assays# | | | Competition assays | | | | | | | | | | | | | | |
| | | | | | | | muskmelon | | | | | LB | | | | | PBS | | | | |
				STM	SEN	SNP	CI sense	stdev sense	CI antisense	stdev antisense	outcome*	CI sense	stdev sense	CI antisense	stdev antisense	outcome*	CI sense	stdev sense	CI antisense	stdev antisense	outcome*
22°C, 24h	STM14_3962	*deaD*	ATP-dependent RNA helicase DeaD	-	-	(-)	0.28	0.45	0.03	0.02	-	0.34	0.54	0.04	0.02	-	1.01	0.30	1.34	0.14	O
	STM14_3964	*pnp*	Polynucleotide phosphorylase/polyadenylase	-	-	-	0.04	0.03	0.04	0.03	-	1.58	2.67	1.53	2.60	O	0.77	0.45	0.52	0.20	O
	STM14_5328	*yjgA*	Ribosome associated protein	-	-	-	0.08	0.08			-	0.11	0.12			-	1.27	0.46			O
	STM14_4959	*oxyR*	DNA-binding transcriptional regulator OxyR	(-/+)	(-/+)	(-/+)	0.55	0.33	0.23	0.35	-	0.19	0.24	0.01	0.01	-	0.74	0.29	0.38	0.24	O
	STM14_3699	*serA*	Phosphoserine phosphatase	(-)	-	-	0.43	0.17	0.35	0.07	-	1.21	0.51	1.29	0.77	O	1.25	0.67	1.20	0.97	O
	STM14_5499	*serB*	Phosphoserine phosphatase	(-)	(-)	-	0.86	0.47	0.57	0.24	O	0.76	0.38	0.67	0.29	O	1.29		0.59		O
	STM14_0583	*ybaK*	Putative cytoplasmic protein	(-)	-	(-)	1.30	1.88	1.06	1.68	O	0.97	0.31	0.69	0.14	O	2.16	1.13	1.50	1.23	O
	STM14_2587	*rfbl_2*	CDP-6-deoxy-delta-3,4-glucose reductase				1.58	0.30	0.96	0.17	O	1.36	0.24	0.96	0.15	O					
8°C, 5d	STM14_3066	*ppk*	Polyphosphate kinase	-	(-)	-	0.05	0.03	0.10	0.10	-	0.07	0.00	0.11	0.04	-	2.56	2.54	1.87	1.44	O
	STM14_3240	*srmB*	ATP-dependent RNA helicase SrmB	-	-	-	0.20	0.08	0.17	0.10	-	0.22	0.09	0.19	0.13	-	1.37	0.27	0.93	0.05	O
	STM14_4639	**STnc400**	sRNA	-	n.d.	n.d.			0.15	0.06	-			0.18	0.11	-			1.15	1.13	O
	STM14_4822	*typA*	GTP-binding protein	-	-	-	0.18	0.28	0.40	0.64	O	0.07	0.10	0.12	0.19	-	0.93		0.92	0.08	O
	STM14_4713	*trxA*	Thioredoxin	-	(-)	(-)	0.52	0.20	0.95	0.33	O	0.35	0.01	0.63	0.18	-					
	STM14_4722	*wecD*	TDP-fucosamine acetyltransferase	-	(-)		1.16	0.60	0.55	0.21	O	0.72	0.26	0.74	0.43	O					
	STM14_4723	*wecE*	TDP-4-oxo-6-deoxy-D-glucose transaminase	+/-	(-)	(-)			2.61	1.42	+			1.58	0.23	+					
	STM14_4725	*wecF [D]*	4-alpha-L-fucosyltransferase	(-)			0.34	0.29	0.32	0.25	-	0.31	0.12	0.29	0.10	-	1.54	0.75	1.05	0.49	O
	STM14_0267	*hlpA*	Periplasmic chaperone	-	(-)		1.94	1.18			O	1.26	0.81			O					
	STM14_4817	*glnG*	Nitrogen regulation protein NR(I)	(-/+)	(-)		2.39	1.38	1.96	0.55	+	1.58	1.15	1.28	0.50	O					
	STM14_5478	*prfC*	Peptide chain release factor 3	-	-	(-)	0.32	0.11	0.40	0.23	-	0.25	0.16	0.30	0.17	-	2.09	1.64	2.23	1.08	O
	STM14_2196	*rnd*	Ribonuclease D	-	(-)		0.42	0.17	0.32	0.10	-	0.23	0.06	0.22	0.06	-	0.74	0.01	1.05	0.28	O
	STM14_5372.J	*holC*	DNA polymerase III subunit chi	-			0.05	0.05			-	0.01	0.01			-	1.05	0.12			O

STM14_5475	**holD**	DNA polymerase III subunit psi	-		n.d.	0.01	0.00			-	0.01	0.01			-	0.90	0.43			O
STM14_3077	*xseA*	Exodeoxyribonuclease VII large subunit	-			1.37	0.11	0.02	0.02	O	0.87	0.70	0.02	0.02	O	1.68	1.28	0.64	0.19	O
STM14_5248	**purA**	Adenylosuccinate synthetase	(-)	(-)		0.41	0.25			-	0.61	0.12			-					
STM14_2205	**yoaH**	Hypothetical protein in *fadD-pabB* IR	-	(-)		0.16	0.05	0.15	0.03	-	0.08	0.01	0.05	0.04	-	0.77	0.49	1.42	1.16	O

From two to four separate experiments. Genes in **bold** are where the STM transposon assay was confirmed in the competition assays. Genes in **bold** and underlined were important for the growth of *Salmonella* on muskmelon but not in LB.

CI = (mutant/wild type)$_{condition}$ / (mutant/wild type)$_{inoculum}$.

#, Fitness effects, as described in **Table 2**.

* Fitness effects: -, single-gene deletion mutants with a significant growth disadvantage in competition assays compared to wild type (CI + stdev < 1); +, advantage in competition assays compared to wild type (CI - stdev > 1); ○, no effect or contradicting Tn data.

Sense – single-gene deletion mutant with a kanamycin resistance cassette inserted in the sense orientation of the gene; antisense – single-gene deletion mutant with a chloramphenicol resistance cassette inserted in the antisense orientation of the gene.

Acknowledgements

We would like to express our gratitude to Verena Hohenester for her invaluable technical support during this study. Additionally, we extend our thanks to Barbara Fritz, Johanna Dietz, Erika Altgenug, René Mamet, and Larissa Klose for their contributions and assistance with the competition experiments.

Supporting Information

Supplementary Figure S1. Growth of the *Salmonella enterica* barcoded transposon mutant libraries (lib) and the corresponding wild type strains (wt) on ready-to-eat muskmelon at 22°C (A) and 8°C (B).

Supplementary Figure S2. Quantitative assessment of the *Salmonella enterica* barcoded transposon mutant libraries.

Supplementary Figure S3. Quantitative assessment of the background microbiota on the ready-to-eat muskmelon samples.

Supplementary Figure S4. Changes in the mutant abundances after growing the random transposon libraries on ready-to-eat muskmelons.

Supplementary Table S1. Primers used for library preparation prior Illumina sequencing.

Supplementary Table S2. DESeq2 statistical analysis for the transposon insertion sequencing experiments of the three *Salmonella enterica* libraries on muskmelon.

Supplementary Table S3. Results of the KEGG- and GO- enrichment analysis.

References

1. EFSA and ECDC, European Food Safety Authority European Centre for Disease Prevention Control. The European Union One Health 2023 Zoonoses report. EFSA Journal, 2024. 22(12): e9106.DOI: https://doi.org/10.2903/j.efsa.2024.9106.

2. Ries, A. A multistate outbreak of Salmonella chester linked to imported cantaloupe, 30th Intersci. in Conf. Antimicrob. Agents Chemother. 1990.

3. Bowen, A., A. Fry, G. Richards, and L. Beuchat, Infections associated with cantaloupe consumption: a public health concern. Epidemiol Infect, 2006. 134(4): 675-685.DOI: 10.1017/s0950268805005480.

4. McGeoch, L.J., A. Hoban, C. Sawyer, H. Rabie, A. Painset, L. Browning, D. Brown, C. McCarthy, A. Nelson, A. Firme, Â. Pista, J. Moreno, J.V. Martins, L. Silveira, J. Machado, P. Vasconcelos, O. Olufon, C. Inzoungou-Massanga, A. Douglas, J. McCormick, L. Larkin, and S. Balasegaram, Salmonella Saintpaul outbreak associated with cantaloupe consumption, the United Kingdom and Portugal, September to November 2023. Epidemiology and Infection, 2024. 152: e78.DOI: 10.1017/S0950268824000670.

5. CDC. Centers for Disease Control and Prevention. Salmonella Outbreak Linked to Cantaloupes. 2024 15.07.2024]; Available from: https://www.cdc.gov/salmonella/sundsvall-11-23/index.html.

6. Fu, Y., A.K. Bhunia, and Y. Yao, Abrasive brushing reduces pathogen biofilms at cantaloupe rind surface. International Journal of Food Microbiology, 2020. 329: 108685.DOI: https://doi.org/10.1016/j.ijfoodmicro.2020.108685.

7. Ukuku, D.O. and G.M. Sapers, Effect of time before storage and storage temperature on survival of Salmonella inoculated on fresh-cut melons. Food Microbiol, 2007. 24(3): 288-95.DOI: 10.1016/j.fm.2006.04.007.

8. Huang, J., Y. Luo, and X. Nou, Growth of Salmonella enterica and Listeria monocytogenes on Fresh-Cut Cantaloupe under Different Temperature Abuse Scenarios. Journal of Food Protection, 2015. 78(6): 1125-1131.DOI: https://doi.org/10.4315/0362-028X.JFP-14-468.

9. BfR. Melons: Health Hazard through Contamination with Pathogenic Bacteria. 2013 [cited 2024 11.10.2024]; Available from: https://www.bfr.bund.de/cm/349/melons-health-hazard-through-contamination-with-pathogenic-bacteria.pdf.

10.   Spector, M.P. and W.J. Kenyon, Resistance and survival strategies of Salmonella enterica
to environmental stresses. Food Research International, 2012. 45(2): 455-481.DOI:
https://doi.org/10.1016/j.foodres.2011.06.056.

11.   Pradhan, D. and V. Devi Negi, Stress-induced adaptations in Salmonella: A ground for
shaping its pathogenesis. Microbiological Research, 2019. 229: 126311.DOI:
https://doi.org/10.1016/j.micres.2019.126311.

12.   de Moraes, M.H., P. Desai, S. Porwollik, R. Canals, D.R. Perez, W. Chu, M. McClelland,
and M. Teplitski, Salmonella persistence in tomatoes requires a distinct set of metabolic
functions identified by transposon insertion sequencing. Applied and Environmental
Microbiology, 2017. 83(5): e03028-16.DOI: 10.1128/aem.03028-16.

13.   de Moraes, M.H., E.B. Soto, I. Salas González, P. Desai, W. Chu, S. Porwollik, M.
McClelland, and M. Teplitski, Genome-wide comparative functional analyses reveal
adaptations of Salmonella sv. Newport to a plant colonization lifestyle. Front Microbiol,
2018. 9: 877.DOI: 10.3389/fmicb.2018.00877.

14.   Holden, E.R., J. Abi Assaf, H. Al-Khanaq, N. Vimont, M.A. Webber, and E. Trampari,
Identification of pathways required for Salmonella to colonize alfalfa using TraDIS-
Xpress. Appl Environ Microbiol, 2024. 90(7): e0013924.DOI: 10.1128/aem.00139-24.

15.   Jayeola, V., M. McClelland, S. Porwollik, W. Chu, J. Farber, and S. Kathariou,
Identification of Novel Genes Mediating Survival of Salmonella on Low-Moisture Foods
via Transposon Sequencing Analysis. Front Microbiol, 2020. 11: 726.DOI:
10.3389/fmicb.2020.00726.

16.   Li, Y., J.K. Salazar, Y. He, P. Desai, S. Porwollik, W. Chu, P.-S.S. Paola, M.L.
Tortorello, O. Juarez, H. Feng, M. McClelland, and W. Zhang, Mechanisms of
Salmonella Attachment and Survival on In-Shell Black Peppercorns, Almonds, and
Hazelnuts. Frontiers in Microbiology, 2020. 11(2311).DOI: 10.3389/fmicb.2020.582202.

17.   Porwollik, S., C.A. Santiviago, P. Cheng, F. Long, P. Desai, J. Fredlund, S. Srikumar,
C.A. Silva, W. Chu, X. Chen, R. Canals, M.M. Reynolds, L. Bogomolnaya, C. Shields, P.
Cui, J. Guo, Y. Zheng, T. Endicott-Yazdani, H.J. Yang, A. Maple, Y. Ragoza, C.J.
Blondel, C. Valenzuela, H. Andrews-Polymenis, and M. McClelland, Defined single-
gene and multi-gene deletion mutant collections in Salmonella enterica sv Typhimurium.
PLoS One, 2014. 9(7): e99820.DOI: 10.1371/journal.pone.0099820.

18.   Feng, K., W. Hu, A. Jiang, G. Saren, Y. Xu, Y. Ji, and W. Shao, Growth of Salmonella
spp. and Escherichia coli O157:H7 on fresh-cut fruits stored at different temperatures.
Foodborne Pathogens and Disease, 2017. 14(9): 510-517.DOI: 10.1089/fpd.2016.2255.

19. Song, H., A.R. Jang, S. Lee, and S.-Y. Lee, Application of sodium alginate-based edible coating with citric acid to improve the safety and quality of fresh-cut melon (Cucumis melo L.) during cold storage. Food Science and Biotechnology, 2024. 33(7): 1741-1750.DOI: 10.1007/s10068-023-01475-y.

20. Clements, M.O., S. Eriksson, A. Thompson, S. Lucchini, J.C. Hinton, S. Normark, and M. Rhen, Polynucleotide phosphorylase is a global regulator of virulence and persistency in Salmonella enterica. Proc Natl Acad Sci U S A, 2002. 99(13): 8784-9.DOI: 10.1073/pnas.132047099.

21. Charollais, J., D. Pflieger, J. Vinh, M. Dreyfus, and I. Iost, The DEAD-box RNA helicase SrmB is involved in the assembly of 50S ribosomal subunits in Escherichia coli. Mol Microbiol, 2003. 48(5): 1253-65.DOI: 10.1046/j.1365-2958.2003.03513.x.

22. Vakulskas, C.A., A. Pannuri, D. Cortés-Selva, T.R. Zere, B.M. Ahmer, P. Babitzke, and T. Romeo, Global effects of the DEAD-box RNA helicase DeaD (CsdA) on gene expression over a broad range of temperatures. Molecular Microbiology, 2014. 92(5): 945-958.DOI: https://doi.org/10.1111/mmi.12606.

23. Hussain, A. and M.K. Ray, Role of DEAD-box RNA helicases in low-temperature adapted growth of Antarctic Pseudomonas syringae Lz4W. Microbiol Spectr, 2024. 12(1): e0433522.DOI: 10.1128/spectrum.04335-22.

24. Jiang, X., R. Keto-Timonen, M. Skurnik, and H. Korkeala, Role of DEAD-box RNA helicase genes in the growth of Yersinia pseudotuberculosis IP32953 under cold, pH, osmotic, ethanol and oxidative stresses. PLoS One, 2019. 14(7): e0219422.DOI: 10.1371/journal.pone.0219422.

25. Favaro, R. and G. Dehò, Polynucleotide phosphorylase-deficient mutants of Pseudomonas putida. J Bacteriol, 2003. 185(17): 5279-86.DOI: 10.1128/jb.185.17.5279-5286.2003.

26. Jones, P.G., M. Mitta, Y. Kim, W. Jiang, and M. Inouye, Cold shock induces a major ribosomal-associated protein that unwinds double-stranded RNA in Escherichia coli. Proceedings of the National Academy of Sciences, 1996. 93(1): 76-80.DOI: doi:10.1073/pnas.93.1.76.

27. Zhu, Y., W. Huang, S.S. Lee, and W. Xu, Crystal structure of a polyphosphate kinase and its implications for polyphosphate synthesis. EMBO reports, 2005. 6(7): 681-687.

28. McMeechan, A., M.A. Lovell, T.A. Cogan, K.L. Marston, T.J. Humphrey, and P.A. Barrow, Inactivation of ppk differentially affects virulence and disrupts ATP homeostasis

587  in Salmonella enterica serovars Typhimurium and Gallinarum. Research in

588  Microbiology, 2007. 158(1): 79-85.DOI: https://doi.org/10.1016/j.resmic.2006.10.008.

589 29. Blum, E., B. Py, A.J. Carpousis, and C.F. Higgins, Polyphosphate kinase is a component

590  of the Escherichia coli RNA degradosome. Molecular Microbiology, 1997. 26(2): 387-

591  398.DOI: https://doi.org/10.1046/j.1365-2958.1997.5901947.x.

592 30. Kim, K.-S., N.N. Rao, C.D. Fraley, and A. Kornberg, Inorganic polyphosphate is

593  essential for long-term survival and virulence factors in Shigella and Salmonella spp.

594  Proceedings of the National Academy of Sciences, 2002. 99(11): 7675-7680.DOI:

595  doi:10.1073/pnas.112210499.

596 31. Hammarlöf, D.L., J.M. Bergman, E. Garmendia, and D. Hughes, Turnover of mRNAs is

597  one of the essential functions of RNase E. Mol Microbiol, 2015. 98(1): 34-45.DOI:

598  10.1111/mmi.13100.

599 32. Prieto-Alamo, M.J., J. Jurado, R. Gallardo-Madueno, F. Monje-Casas, A. Holmgren, and

600  C. Pueyo, Transcriptional regulation of glutaredoxin and thioredoxin pathways and

601  related enzymes in response to oxidative stress. J Biol Chem, 2000. 275(18): 13398-

602  405.DOI: 10.1074/jbc.275.18.13398.

603 33. Kuźniak, E. and T. Kopczewski, The chloroplast reactive oxygen species-redox system in

604  plant immunity and disease. Front Plant Sci, 2020. 11: 572686.DOI:

605  10.3389/fpls.2020.572686.

606 34. Cota, I., A.B. Blanc-Potard, and J. Casadesús, STM2209-STM2208 (opvAB): a phase

607  variation locus of Salmonella enterica involved in control of O-antigen chain length.

608  PLoS One, 2012. 7(5): e36863.DOI: 10.1371/journal.pone.0036863.

609 35. Cota, I., M.A. Sánchez-Romero, S.B. Hernández, M.G. Pucciarelli, F. García-Del

610  Portillo, and J. Casadesús, Epigenetic control of Salmonella enterica O-Antigen chain

611  length: a tradeoff between virulence and bacteriophage resistance. PLoS Genet, 2015.

612  11(11): e1005667.DOI: 10.1371/journal.pgen.1005667.

613 36. Ricke, S.C., T.M. Dawoud, S.A. Kim, S.H. Park, and Y.M. Kwon, Salmonella Cold

614  Stress Response: Mechanisms and Occurrence in Foods. Adv Appl Microbiol, 2018. 104:

615  1-38.DOI: 10.1016/bs.aambs.2018.03.001.

616 37. Sanguankiattichai, N., P. Buscaill, and G.M. Preston, How bacteria overcome flagellin

617  pattern recognition in plants. Current Opinion in Plant Biology, 2022. 67: 102224.DOI:

618  https://doi.org/10.1016/j.pbi.2022.102224.

619 38. Zarkani, A.A., N. López-Pagán, M. Grimm, M.A. Sánchez-Romero, J. Ruiz-Albert, C.R.

620  Beuzón, and A. Schikora, Salmonella Heterogeneously Expresses Flagellin during

Colonization of Plants. Microorganisms, 2020. 8(6).DOI:
10.3390/microorganisms8060815.

39. Fan, H., J. Hahm, S. Diggs, J.J.P. Perry, and G. Blaha, Structural and Functional Analysis of BipA, a Regulator of Virulence in Enteropathogenic Escherichia coli. Journal of Biological Chemistry, 2015. 290(34): 20856-20864.DOI: 10.1074/jbc.M115.659136.

40. Tiefenbacher, S., V. Pezo, P. Marlière, T.M. Roberts, and S. Panke, Systematic analysis of tRNA transcription unit deletions in E. coli reveals insights into tRNA gene essentiality and cellular adaptation. Scientific Reports, 2024. 14(1): 24102.DOI: 10.1038/s41598-024-73407-7.

41. Kenri, T., F. Imamoto, and Y. Kano, Construction and characterization of an Escherichia coli mutant deficient in the metY gene encoding tRNAf2Met: either tRNAf1Met or tRNAf2Met is required for cell growth. Gene, 1992. 114(1): 109-114.DOI: https://doi.org/10.1016/0378-1119(92)90715-2.

42. Vial, T.C., K.E. Baker, and R.A. Kelln, Dual control by purines and pyrimidines of the expression of the pyrD gene of Salmonella typhimurium. FEMS Microbiology Letters, 1993. 111(2-3): 309-314.DOI: 10.1111/j.1574-6968.1993.tb06403.x.

43. Wilson, H.R. and C.L. Turnbough, Jr., Role of the purine repressor in the regulation of pyrimidine gene expression in Escherichia coli K-12. J Bacteriol, 1990. 172(6): 3208-13.DOI: 10.1128/jb.172.6.3208-3213.1990.

44. Yang, H.J., L. Bogomolnaya, M. McClelland, and H. Andrews-Polymenis, De novo pyrimidine synthesis is necessary for intestinal colonization of Salmonella Typhimurium in chicks. PLoS One, 2017. 12(10): e0183751.DOI: 10.1371/journal.pone.0183751.

45. Johnson, D.C., D.R. Dean, A.D. Smith, and M.K. Johnson, Structure, function, and formation of biological iron-sulfur clusters. Annual Review of Biochemistry, 2005. 74(Volume 74, 2005): 247-281.DOI: https://doi.org/10.1146/annurev.biochem.74.082803.133518.

46. Vickery, L.E. and J.R. Cupp-Vickery, Molecular Chaperones HscA/Ssq1 and HscB/Jac1 and Their Roles in Iron-Sulfur Protein Maturation. Critical Reviews in Biochemistry and Molecular Biology, 2007. 42(2): 95-111.DOI: 10.1080/10409230701322298.

47. Yang, J., J.P. Bitoun, and H. Ding, Interplay of IscA and IscU in biogenesis of iron-sulfur clusters. J Biol Chem, 2006. 281(38): 27956-63.DOI: 10.1074/jbc.M601356200.

48. Nugent, S.L., F. Meng, G.B. Martin, and C. Altier, Acquisition of Iron Is Required for Growth of Salmonella spp. in Tomato Fruit. Applied and Environmental Microbiology, 2015. 81(11): 3663-3670.DOI: doi:10.1128/AEM.04257-14.

655   49.   Pinski, A., A. Betekhtin, K. Hupert-Kocurek, L.A.J. Mur, and R. Hasterok, Defining the
656   Genetic Basis of Plant‾Endophytic Bacteria Interactions. Int J Mol Sci, 2019. 20(8).DOI:
657   10.3390/ijms20081947.

658   50.   Jarvik, T., C. Smillie, E.A. Groisman, and H. Ochman, Short-term signatures of
659   evolutionary change in the Salmonella enterica serovar Typhimurium 14028 genome. J
660   Bacteriol, 2010. 192(2): 560-7.DOI: 10.1128/jb.01233-09.

661   51.   Toro, M., P. Retamal, S. Ayers, M. Barreto, M. Allard, E.W. Brown, and N. Gonzalez-
662   Escalona, Whole-genome sequencing analysis of Salmonella enterica serovar Enteritidis
663   isolates in Chile provides insights into possible transmission between gulls, poultry, and
664   humans. Appl Environ Microbiol, 2016. 82(20): 6223-6232.DOI: 10.1128/aem.01760-16.

665   52.   Love, M.I., W. Huber, and S. Anders, Moderated estimation of fold change and
666   dispersion for RNA-seq data with DESeq2. Genome Biol, 2014. 15(12): 550.DOI:
667   10.1186/s13059-014-0550-8.

668   53.   Kanehisa, M., M. Furumichi, Y. Sato, M. Kawashima, and M. Ishiguro-Watanabe,
669   KEGG for taxonomy-based analysis of pathways and genomes. Nucleic Acids Res, 2023.
670   51(D1): D587-d592.DOI: 10.1093/nar/gkac963.

671   54.   Thomas, P.D., D. Ebert, A. Muruganujan, T. Mushayahama, L.-P. Albou, and H. Mi,
672   PANTHER: Making genome-scale phylogenetics accessible to all. Protein Science, 2022.
673   31(1): 8-22.DOI: https://doi.org/10.1002/pro.4218.

674   55.   Ashburner, M., C.A. Ball, J.A. Blake, D. Botstein, H. Butler, J.M. Cherry, A.P. Davis, K.
675   Dolinski, S.S. Dwight, J.T. Eppig, M.A. Harris, D.P. Hill, L. Issel-Tarver, A. Kasarskis,
676   S. Lewis, J.C. Matese, J.E. Richardson, M. Ringwald, G.M. Rubin, and G. Sherlock,
677   Gene Ontology: tool for the unification of biology. Nature Genetics, 2000. 25(1): 25-
678   29.DOI: 10.1038/75556.

679   56.   Aleksander, S.A., J. Balhoff, S. Carbon, J.M. Cherry, H.J. Drabkin, D. Ebert, M.
680   Feuermann, P. Gaudet, N.L. Harris, D.P. Hill, R. Lee, H. Mi, S. Moxon, C.J. Mungall, A.
681   Muruganugan, T. Mushayahama, P.W. Sternberg, P.D. Thomas, K. Van Auken, J.
682   Ramsey, D.A. Siegele, R.L. Chisholm, P. Fey, M.C. Aspromonte, M.V. Nugnes, F.
683   Quaglia, S. Tosatto, M. Giglio, S. Nadendla, G. Antonazzo, H. Attrill, G. dos Santos, S.
684   Marygold, V. Strelets, C.J. Tabone, J. Thurmond, P. Zhou, S.H. Ahmed, P. Asanitthong,
685   D. Luna Buitrago, M.N. Erdol, M.C. Gage, M. Ali Kadhum, K.Y.C. Li, M. Long, A.
686   Michalak, A. Pesala, A. Pritazahra, S.C.C. Saverimuttu, R. Su, K.E. Thurlow, R.C.
687   Lovering, C. Logie, S. Oliferenko, J. Blake, K. Christie, L. Corbani, M.E. Dolan, H.J.
688   Drabkin, D.P. Hill, L. Ni, D. Sitnikov, C. Smith, A. Cuzick, J. Seager, L. Cooper, J.

Elser, P. Jaiswal, P. Gupta, P. Jaiswal, S. Naithani, M. Lera-Ramirez, K. Rutherford, V.

Wood, J.L. De Pons, M.R. Dwinell, G.T. Hayman, M.L. Kaldunski, A.E. Kwitek, S.J.F.

Laulederkind, M.A. Tutaj, M. Vedi, S.-J. Wang, P. D'Eustachio, L. Aimo, K. Axelsen, A.

Bridge, N. Hyka-Nouspikel, A. Morgat, S.A. Aleksander, J.M. Cherry, S.R. Engel, K.

Karra, S.R. Miyasato, R.S. Nash, M.S. Skrzypek, S. Weng, E.D. Wong, E. Bakker, T.Z.

Berardini, L. Reiser, A. Auchincloss, K. Axelsen, G. Argoud-Puy, M.-C. Blatter, E.

Boutet, L. Breuza, A. Bridge, C. Casals-Casas, E. Coudert, A. Estreicher, M. Livia

Famiglietti, M. Feuermann, A. Gos, N. Gruaz-Gumowski, C. Hulo, N. Hyka-Nouspikel,

F. Jungo, P. Le Mercier, D. Lieberherr, P. Masson, A. Morgat, I. Pedruzzi, L. Pourcel, S.

Poux, C. Rivoire, S. Sundaram, A. Bateman, E. Bowler-Barnett, H. Bye-A-Jee, P. Denny,

A. Ignatchenko, R. Ishtiaq, A. Lock, Y. Lussi, M. Magrane, M.J. Martin, S. Orchard, P.

Raposo, E. Speretta, N. Tyagi, K. Warner, R. Zaru, A.D. Diehl, R. Lee, J. Chan, S.

Diamantakis, D. Raciti, M. Zarowiecki, M. Fisher, C. James-Zorn, V. Ponferrada, A.

Zorn, S. Ramachandran, L. Ruzicka and M. Westerfield, The Gene Ontology

Consortium. The Gene Ontology knowledgebase in 2023. Genetics, 2023. 224(1).DOI:

10.1093/genetics/iyad031.

57.   Alexa, A. and J. Rahnenfuhrer. topGO: Enrichment Analysis for Gene Ontology. R

package version 2.59.0. 2024 12.05.2025]; Available from:

https://bioconductor.org/packages/topGO.

58.   Wu, T., E. Hu, S. Xu, M. Chen, P. Guo, Z. Dai, T. Feng, L. Zhou, W. Tang, L. Zhan, X.

Fu, S. Liu, X. Bo, and G. Yu, clusterProfiler 4.0: A universal enrichment tool for

interpreting omics data. The Innovation, 2021. 2(3).DOI: 10.1016/j.xinn.2021.100141.