Dissecting surveying behavior of reactive microglia under chronic neurodegeneration

Sunitha Subhramanian[1], Olga Bocharova[1], Natallia Makarava[1], Tarek Safadi[1], Ilia V. Baskakov[1*]

[1] Department of Neurobiology, University of Maryland School of Medicine, Baltimore, Maryland, USA

[*]Correspondence: 111 Penn St., Baltimore, MD, 21201, USA, Phone: 410-706-4562,
baskakov@som.umaryland.edu

Abstract
In the healthy brain, microglia maintain homeostasis by continuously surveying neuronal health through highly dynamic processes that form puringergic junctions with neuronal somas. These mechanisms are finely tuned for the rapid detection of acute injuries. However, during the transition to a chronically reactive state in neurodegenerative diseases, microglial ramification decreases even as the need for neuronal monitoring escalates. How reactive microglia adapt their surveillance strategies under these conditions remains poorly understood. Using time-lapse imaging of acute brain slices from prion-infected mice, we identified a previously unrecognized mode of neuronal surveillance employed by reactive microglia. Unlike homeostatic microglia, which exhibit low somatic mobility and high process motility, enabling broad, simultaneous monitoring, reactive microglia display high somatic mobility. These cells actively migrate through the brain parenchyma, pausing to form direct and extensive body-to-body contacts with individual neurons. Contact durations ranged from minutes to several hours, often involving partial or full somatic envelopment, with transitions between these states being both frequent and reversible. Notably, reactive microglia exhibited sustained intracellular calcium bursts correlated with their increased mobility. Pharmacological inhibition of the P2Y6 receptor partially reduced microglial migration without disrupting their ability to form neuronal contacts. Furthermore, this highly mobile behavior persisted in acutely isolated reactive microglia *in vitro*, even in the absence of external stimuli, indicating that dynamic mobility is an intrinsic feature of the reactive phenotype. These findings reveal a fundamental shift in microglial surveillance architecture during chronic neurodegeneration - transforming from static, multi-neuron monitoring to dynamic, neuron-by-neuron engagement. This work uncovers a novel, adaptive strategy of microglial behavior with critical implications for understanding microglia-neuron interaction under chronic neurodegeneration.

Introduction
Under homeostatic conditions, microglia continuously monitor neuronal health through specialized structures known as somatic purinergic junctions - sites where microglial processes make direct contact with neuronal cell bodies [1]. A key molecular component of these junctions is the P2Y12 receptor, which plays a critical role in mediating microglia-neuron communication [1,2]. Homeostatic, ramified microglia exhibit highly dynamic behavior, extending and retracting their processes in a phenomenon termed *microglial motility* [2-5]. These transient process-to-soma contacts constitute the primary mode of neuronal surveillance in the healthy brain.

Under chronic neurodegenerative conditions, microglia acquire a sustained reactive phenotype [6-9]. Reactive microglia differ markedly in morphology from their homeostatic counterparts, displaying an amoeboid shape with reduced process complexity. Both microglial ramification and P2Y12 receptor expression are significantly diminished as microglia transition to the reactive state [7,8,10-13]. Paradoxically, this occurs at a time when the demand for neuronal surveillance intensifies due to increased neuronal stress and disfunction. How microglia maintain effective surveillance of neurons under conditions of chronic neuroinflammation remains poorly understood.

Prion diseases, or transmissible spongiform encephalopathies, are fatal and infectious neurodegenerative disorders that affect both humans and animals [14]. The hallmark pathological event in

these diseases is the accumulation and spread of the misfolded, β-sheet-rich isoform of the prion protein
(PrP[Sc]) throughout the central nervous system (CNS). This process involves the templated conversion of
the host's normal cellular prion protein (PrP[C]) into the disease-associated PrP[Sc] conformation [15,16].
Chronic neuroinflammation is a prominent neuropathological feature of prion diseases, as well as other
neurodegenerative diseases such as Alzheimer's and Parkinson's diseases [6-9,17]. Notably, unlike animal
models of other neurodegenerative diseases, prion-infected animals including mice develop authentic
ultimately lethal neurodegenerative disease, that recapitulates key features of the human prion disease
[18,19].
Previously, we demonstrated that in prion-infected mice, reactive microglia engage in a distinct
form of microglia-neuron interaction characterized by partial envelopment of neuronal cell bodies [20].
These extensive body-to-body contacts differ from typical process-based surveillance. The enveloped
neurons appear structurally intact, lack classical apoptotic markers, but exhibit functional impairments.
This phenomenon is consistently observed across multiple brain regions exhibiting neuroinflammation in
both prion-infected mice and human cases of sporadic Creutzfeldt-Jakob disease [20]. Strikingly, genetic
ablation of P2Y12 increases the frequency of neuronal envelopment and accelerates disease progression
in prion-infected mice [21]. In non-infected mice, P2Y12 deletion also increase the prevalence of microglia-
neuron body-to-body interactions at the expense of process-to-body contacts [21]. These findings suggest
that neuronal envelopment represents an alternative, P2Y12-independent mode of microglial surveillance,
employed by reactive microglia under pathological conditions. To date, quantification of neuronal
envelopment has relied on fixed brain tissue [20,21], leaving key questions about the temporal dynamics of
these interactions unresolved. Although the frequency of envelopment increases with disease progression,
it remains unclear whether individual interactions are sustained over time or whether reactive microglia
sequentially engage multiple neurons through transient contacts.
The current study employed *ex vivo* time-lapse imaging of acute organotypic brain slices prepared
from prion-infected mice to investigate the surveillance behavior of reactive microglia under chronic
neurodegenerative conditions. Unlike homeostatic microglia, reactive myeloid cells loose territorial
confinement. In contrast to homeostatic microglia - which exhibit low somatic mobility but high process
motility, enabling simultaneous monitoring of multiple neurons - reactive microglia display high somatic
mobility. Reactive microglia move dynamically through tissue, engaging in sequential body-to-body
contacts with individual neurons. This shift in behavior represents a fundamental reorganization of
microglial surveillance strategies in the diseased brain. Surprisingly, acutely isolated reactive microglia
retained high mobility *in vitro* in the absence of external stimuli, indicating that this dynamic behavior is
an intrinsic feature of the reactive phenotype. The current studies reveal that in the reactive state,
microglia adopt fundamentally different surveillance strategies to respond to the altered demands of the
chronically inflamed and degenerating brain.
85
Results
Reactive microglia envelop neuron
To investigate microglial dynamics, we performed ex vivo time-lapse imaging using acute
organotypic brain slices prepared from Cx3cr1/EGFP mice (fractalkine receptor knockout/EGFP knock-
in). These mice express enhanced green fluorescent protein (EGFP) under the control of the
endogenous *Cx3cr1* promoter, restricting expression to myeloid cells. Previous studies have demonstrated
that *Cx3cr1* deficiency does not influence disease pathogenesis, survival, or microglial activation in mice
infected with the 22L or RML prion strains [22]. In agreement with these findings, we observed
that *Cx3cr1* deficiency and EGFP knock-in in Cx3cr1/EGFP mice did not alter the incubation time to
disease following infection with mouse-adapted SSLOW or 22L prion strains, compared to wild-type
(WT; C57BL/6J) controls (Figure S1a–c). Moreover, PrP[Sc] accumulation levels were comparable between
SSLOW-infected Cx3cr1/EGFP and WT mice (Figure S1c,d).
For subsequent experiments, we selected the SSLOW strain due to its strong induction of
neuroinflammation and the shortest incubation time among mouse-adapted prion strains [23,24]. Examination
of fixed brain slices from SSLOW-infected Cx3cr1/EGFP mice at the clinical stage revealed extensive

envelopment of neurons by Iba1[+] reactive microglia, characterized by prominent body-to-body contacts
(Figure 1a), consistent with our previous report in WT mice [20]. Notably, the cortices of SSLOW-
infected Cx3cr1/EGFP mice exhibited a high prevalence of neuronal envelopment events (Figure S1f). In
contrast, brain slices from age-matched, non-infected Cx3cr1/EGFP mice displayed microglia with a
highly ramified morphology, primarily engaging in process-to-cell body interactions with neurons (Figure
1a, Figure S1g), typical of homeostatic microglia.

Reactive myeloid cells exhibit high mobility

To investigate microglial dynamics in prion-affected brains, *ex vivo* time-lapse imaging was
performed on Cx3cr1/EGFP mice infected with the SSLOW prion strain via intraperitoneal (ip) injection.
Imaging was conducted at three stages of disease progression: late subclinical (111–113 days post-
inoculation, dpi), early clinical (125–128 dpi), and advanced (162–169 dpi). In mice inoculated with
SSLOW via ip route, clinical onset typically occurs around 122 dpi [20]. Age-matched non-
infected Cx3cr1/EGFP mice (160–164 days old) served as healthy controls.

To be consistent with previously established terminology [25], the term "motility" will be used to
refer to the movement of microglia processes, whereas the movement of cell bodies to new position will
be referred to by the term "mobility". Initial time-lapse recordings of acute brain slices from non-infected
adult Cx3cr1/EGFP mice, acquired at a rate of one frame every five minutes, demonstrated that microglial
morphology remained ramified - indicative of a homeostatic state - for at least six hours of continuous
imaging (Video S1). During this period, microglia largely remained within defined territories, displaying
limited somatic mobility but sustained dynamic process activity, confirming cellular viability (Video S1).
However, noticeable photobleaching occurred after approximately three hours of imaging, with
accelerated photobleaching observed at higher imaging frequencies. Accordingly, all subsequent time-
lapse videos were captured at a frame rate of one frame every five minutes.

Recent studies have shown that transcriptomic changes in slice cultures are most pronounced one
day post-preparation [26], though morphological, functional, and gene expression alterations can begin as
early as four hours post-slicing [27]. In contrast to the stable process motility observed in homeostatic
microglia, EGFP[+] cells in slices from SSLOW-infected animals exhibited both high process activity and
active somatic translocation (Video S2). To define the optimal imaging window for capturing dynamic
behavior, we tracked individual cell movements across six consecutive one-hour intervals following slice
preparation. Microglial activity progressively declined after 3–4 hours in both SSLOW-infected and
control slices (Figure S2a–c), prompting us to restrict all subsequent imaging experiments to a three-hour
window.

At all examined disease stages, EGFP[+] cells in prion-infected brains displayed significantly
increased mobility compared to controls, as reflected by elevated mean speed and total soma
displacement (Figure 2a–c). A comprehensive heatmap analysis of mobility parameters, including soma
displacement, speed, and directionality, demonstrated a progressive enhancement of microglial activity
with advancing disease (Figure 2d).

'Kiss-and-ride' surveying behavior of reactive myeloid cells

Analysis of SSLOW-infected brain slices based on the distance traveled by EGFP[+] cell somas
revealed two distinct behavioral patterns: high mobility and low mobility. Low-mobility cells remained
confined to a limited area, displaying minimal displacement and exhibiting localized, jiggling movements
(Figure 3a). In contrast, high-mobility cells traversed significantly longer distances across the tissue
(Figures 3a, 3c, 3d). These two phenotypes also differed in their mean directional change rate, a metric
quantifying the average shift in movement direction over time (Figure 3e). Low-mobility cells exhibited
higher directional change rates, consistent with their confined, wobbling behavior, whereas high-mobility
cells demonstrated more linear and directed trajectories.

Detailed tracking of high-mobility cells uncovered a distinctive, intermittent surveying pattern.
EGFP[+] cells extended processes in multiple directions, often simultaneously reaching out to several
neurons. Subsequently, the cell somas translocated along some of these processes while others retracted
(Figures 4a, 4d; Videos S3–S6, S12). While migrating through the extracellular matrix, these cells paused
to establish transient, soma-to-soma contact with neurons before resuming movement to engage with

subsequent neurons. We refer to this dynamic pattern as 'kiss-and-ride' behavior. The duration of these
transient contacts ranged from several minutes to over an hour (Videos S3–S5). Notably, this behavior
was consistently observed across all three stages of disease progression: subclinical, early clinical, and
advanced (Figures 4a, 4d; Videos S3–S6).

During a three-hour imaging window, some EGFP[+] cells were observed to contact up to six
different neurons (Figure 4c; Videos S4, S7), while others engaged with only one or two neurons,
maintaining prolonged interactions with each (Figures 4c, 4d; Videos S3, S5, S8). Occasionally, a single
EGFP[+] cell simultaneously contacted or partially enveloped two or three neuronal somas (Figures 4a, 5a,
5b; Videos S8, S9). Interestingly, some neurons were sequentially surveyed by two different EGFP[+] cells
within the same three-hour period (Figure 4b; Video S10). Collectively, the behaviors of high-mobility
EGFP[+] cells underscore a lack of territorial restriction, contrasting with the spatially constrained
dynamics typically seen in ramified microglia under homeostatic conditions.

Enveloping behavior of reactive myeloid cells

Low-mobility EGFP[+] cells typically maintained contact with a single neuronal soma throughout
the entire duration of time-lapse imaging (Videos S8, S11). Occasionally, these EGFP[+] cells were
observed slowly transitioning between neuronal somas or oscillating between two or three somas (Figure
5a,b; Videos S3, S8, S9). Body-to-body contact between EGFP[+] myeloid cells and neurons was often
sustained for extended periods and could progress to full envelopment of the neuronal soma (Figure 4c,
5c; Videos S12–S15). Notably, full envelopment did not necessarily culminate in phagocytosis; EGFP[+]
cells frequently reversed the process, returning from full envelopment to partial envelopment or contact
(Figure 4c; Video S12).

Previous analyses of fixed brain sections have shown that reactive myeloid cells selectively
envelop neuronal cells, avoiding other cell types such as astrocytes or oligodendrocytes [20,21]. In the
current study, we attempted to visualize neuronal somas during time-lapse imaging using the calcium-
sensitive dye Calbryte-590, following protocols from earlier research [2]. However, this proved challenging.
Unlike neurons in non-infected animals, neurons in prion-infected slices exhibited markedly reduced
Calbryte-590 fluorescence, suggesting dysfunction in calcium signaling. Despite the weak signal, some
neuronal somas undergoing envelopment still showed detectable Calbryte-590 fluorescence in SSLOW-
infected slices (Video S16). Interestingly, brighter calcium puncta were frequently observed in association
with reactive myeloid cells (Videos S14, S18).

The same behavioral patterns observed in SSLOW-infected brains were also evident in brain
slices from Cx3cr1/EGFP mice infected with the 22L mouse-adapted prion strain. These included both
high-mobility EGFP[+] cells exhibiting 'kiss-and-ride' interactions and low-mobility cells engaged in
neuronal envelopment (Video S17).

The interaction patterns between reactive myeloid cells and neurons that are seen in time-laps
imaging in dynamics, could be captured at a much higher resolution using confocal microscopy imaging
of fixed brain slices. Confocal imaging confirmed extensive body-to-body contact between Iba1[+] cells
and neurons, ranging from partial to full envelopment (Figure 5d,e). Some Iba1[+] cells enveloped one
neuron while simultaneously extending processes to contact or even envelop additional neurons. Cases
were also observed where two Iba1[+] cells partially enveloped a single neuron simultaneously. These
patterns emphasize the profound body-to-body interactions between reactive myeloid cells and neurons in
prion-infected brains, contrasting sharply with the predominantly process-mediated interactions seen
under homeostatic conditions.

All behavioral patterns - 'kiss-and-ride', partial envelopment, and full envelopment - were
consistently observed across all three stages of disease progression: preclinical, early clinical, and
advanced. Notably, the proportion of high-mobility myeloid cells increased from the preclinical to early
clinical stages (Figure 3b), possibly reflecting heightened surveillance demands in response to emerging
neuronal dysfunction. At advanced stages, however, this proportion declined (Figure 3b), which may
indicate a shift toward prolonged interactions with severely impaired neurons and/or the onset of intrinsic
dysfunction within the reactive myeloid population.

Morphology of both high- and low-mobility myeloid cells is consistent with reactive phenotype

To examine the morphological features associated with different cell mobilities, measurements of
EGFP⁺ cells were averaged across all time frames acquired during the three-hour imaging period. This
statistical averaging approach accounts for the dynamic and continuously changing shapes of individual
cells over time. Notable morphological differences were observed between cells from healthy animals and
those from SSLOW-infected animals. Based on parameters such as cell radius, area, and perimeter,
EGFP⁺ cells from SSLOW-infected animals exhibited a hypertrophic, amoeboid morphology - consistent
with a reactive phenotype (Fig. 5a–c). These morphological changes were evident in both high- and low-
mobility cell populations, suggesting that regardless of mobility behavior, the cells adopted a reactive
phenotype. A similar pattern was observed for the shape index, a dimensionless metric that quantifies
overall cell shape (Fig. 5d).
Interestingly, in comparison to low-mobility cells, the high-mobility cells from SSLOW-infected
animals appeared to be closer to the cells from healthy controls compared (Fig. 5d). This may reflect the
elongated morphology adopted by mobile cells as they navigate the extracellular matrix, influencing the
overall statistical representation of their shape. A trend toward lower values in cell radius, area, perimeter,
and shape index with disease progression aligned with confocal imaging data, which illustrated a more
pronounced amoeboid morphology and reduced ramification at advanced disease stages (Fig. 5d-e, 6a-d).
Overall, the statistical averaging of morphological parameters supports the classification of cells
based on the clinical status of the animals from which brain slices were derived. Myeloid cells with
similar mobility but different clinical backgrounds (e.g., normal vs. SSLOW low-mobility) exhibited
clear morphological distinctions, whereas cells with differing mobility within the same pathological
condition (SSLOW high- vs. low-mobility) were morphologically similar.

High-mobility myeloid cells exhibit sustained Ca²⁺ bursts

Calcium signaling plays a pivotal role in microglial activation and migration, particularly in
response to brain injury or neuronal damage [28-30]. Previous studies have demonstrated that activated
microglia exhibit sustained calcium bursts that correlate with their migratory behavior [31]. Using Calbryte-
590, we observed sustained calcium bursts in EGFP⁺ cells, especially in highly motile populations,
within brain slices from SSLOW-infected mice (Figure 7a).
Due to the time-lapse imaging interval of one frame every five minutes, we were unable to
precisely quantify the duration or frequency of individual calcium bursts. Nonetheless, high-mobility cells
consistently exhibited elevated calcium signals throughout the entire three-hour imaging period (Figure
7b,c; Videos S18–S20). These signals were primarily localized to the soma, although occasional bursts
were also observed within cellular processes (Videos S18–S20). EGFP⁺ cells engaged in neuronal
envelopment also showed calcium activity, albeit at lower intensities than highly motile cells (Figure 7b,c;
Videos S19, S21).

Inhibition of P2Y6 Receptor Reduces the Mobility of Reactive Myeloid Cells

Calcium transients in microglia are regulated by the P2Y6 receptor [30]. Activation of this receptor
by its endogenous ligand, UDP, has been implicated in multiple microglial functions, including migration,
phagocytosis of damaged neurons, and clearance of apoptotic debris and Aβ plaques [32,30,33-35]. To assess
the role of P2Y6 signaling in the dynamics of reactive myeloid cells, we acutely inhibited the receptor
using the selective antagonist MRS-2578. This was performed on brain slices obtained at the early clinical
disease stage, when the proportion of highly motile EGFP⁺ cells was maximal (Figure 3b).
MRS-2578 treatment reduced both mean cell speed and total distance traveled by EGFP⁺ cells in
SSLOW-infected slices compared to mock-treated controls (Figure 8a). However, mobility did not
decrease to the levels observed in uninfected slices, suggesting a partial rather than complete effect of
P2Y6 inhibition (Figure 8a). In MRS-2578-treated slices, the majority of reactive EGFP⁺ cells were
observed either enveloping neuronal somata or remaining closely associated with them (Videos S22, S23).
In some cases, individual EGFP⁺ cells were seen alternating between neighboring neuronal bodies,
sequentially enveloping each soma (Figure 8b; Video S23). These findings suggest that P2Y6 signaling
facilitates long-range migration of reactive myeloid cells. While its inhibition diminishes overall mobility,
it does not abolish localized movements or neuron-associated behaviors such as soma-to-soma
interactions.

Reactive myeloid cells retain elevated basal mobility *in vitro* in the absence of external stimuli

In homeostatic microglia, the extension of cellular processes and migration toward injury sites are driven by neuronal activity and extracellular cues such as ATP, ADP, and UDP. It is generally presumed that similar cues govern the dynamics of chronically reactive microglia; however, this remains unverified. To directly test whether environmental stimuli are required for the mobility of reactive myeloid cells, we examined their behavior *in vitro* under stimulus-free conditions.

Myeloid cells were acutely isolated from SSLOW-infected and uninfected Cx3cr1/EGFP mice using CD11b-coated magnetic beads. Basal motility was then assessed *in vitro* without any exogenous stimulation. Remarkably, cells from SSLOW-infected mice exhibited significantly higher basal mobility than those from control animals, as reflected by increased speed and displacement (Figure 9a, b, d).

Co-culture with N2a neuroblastoma cells did not alter the mobility of control myeloid cells. However, it modestly enhanced the mobility of reactive myeloid cells compared to monoculture conditions (Figure 9a, b, d). Notably, this increase was accompanied by a decrease in directional change rate, suggesting a shift toward more directed and less stochastic movement in the presence of N2a cells (Figure 9c). Together, these results indicate that reactive myeloid cells possess an intrinsically elevated basal mobility that does not require exogenous stimulation. Moreover, despite their heightened autonomous activity, these cells remain responsive to environmental cues that do not elicit similar behavioral changes in homeostatic microglia.

Discussion

In the healthy brain, microglia continuously monitor neuronal activity via highly dynamic processes that extend and retract to form purinergic junctions with neuronal somas [1-3,36]. These surveillance mechanisms in the homeostatic state are primarily tuned to detect acute injuries, as microglia respond to changes in neuronal activity and environmental signals such as ATP, ADP, and UDP released by damaged neurons [4,5,35,37-41]. In contrast to acute or focal injuries, neuronal dysfunction in chronic conditions evolves gradually and spreads across broader regions. It is commonly assumed that reactive microglia continue to rely on the same cues and surveillance strategies as in the homeostatic state. However, during the transition to a reactive phenotype in the context of chronic neurodegeneration, both the complexity and number of microglial processes involved in surveillance decrease, even as the demand for neuronal monitoring intensifies. Additionally, expression of P2Y12, a key receptor for sensing ATP and ADP at purinergic junctions, is significantly reduced [7,8,10-13]. This raises a critical question: how do reactive microglia adapt their surveillance strategies to meet the increasing demands of chronic neurodegeneration?

Our findings suggest that, in the reactive state, microglia adopt a distinct approach to neuronal monitoring. In the homeostatic brain, individual microglial cells occupy well-defined territories, extending their processes to simultaneously contact multiple neurons. By contrast, in the reactive state, myeloid cells lose this territorial organization, becoming highly mobile and migrating through the extracellular matrix. They pause intermittently to form direct body-to-body contacts, typically with one neuron at a time. This movement begins with process extension, followed by somal translocation along the path of the extended process.

On average, each myeloid cell surveyed one neuron per hour. However, the duration of contact varied considerably - from a few minutes to several hours - as did the extent of neuronal envelopment. Partial envelopment of neuronal somas was the most common form of interaction, though full envelopment was also observed. Notably, full envelopment was often reversible, transitioning back to partial envelopment or complete retraction. These ongoing morphological changes - including the appearance, disappearance, and reappearance of intercellular contacts - underscore the highly dynamic nature of communication between reactive myeloid cells and neuronal somas. Although individual contacts were fluid, they could be maintained for several hours. It is tempting to speculate that the duration and degree of envelopment may reflect both the extent of neuronal damage and the complexity of the underlying decision-making processes governing neuronal fate.

Reactive myeloid cells frequently interacted with two or three neuronal somas simultaneously,
exhibiting oscillatory or "wobbling" movements between them. This behavior suggests a capacity for
multitasking, enabling simultaneous surveillance of multiple neurons. Furthermore, within a three-hour
window, multiple myeloid cells were observed interacting with the same neuron, indicating a potential
collective or coordinated decision-making process regarding neuronal viability. Investigating the
molecular and cellular mechanisms underlying this decision-making process represents an important
direction for future research.
To navigate the extracellular matrix, an amoeboid morphology offers clear advantages over a
ramified one. In the reactive state, microglia upregulate matrix-degrading enzymes that facilitate
migration by clearing paths through the adhesive matrix [42]. The motility pattern of reactive myeloid cells
observed in our study resembles that of embryonic mouse microglia, which alternate between phases of
process extension and subsequent somal translocation along the axis of the extended processes [43]. This
mechanical similarity suggests that reactive microglia may engage migratory mechanisms akin to those
used during developmental stages to support their surveillance functions. However, unlike their
embryonic counterparts, reactive myeloid cells appear to actively scan their environment to select among
multiple potential neuronal targets, indicating a more decision-oriented migratory behavior.
In our study, the average speed of the highly mobile microglial population ranged from 13.2 μm/h
at the subclinical stage to 19.1 μm/h at the advanced disease stage. These velocities are lower than those
reported for hippocampal microglia at postnatal day 2 (~36 μm/h) and cortical microglia at embryonic day
17.5 (~25 μm/h) [44]. It is important to note that our measurements likely underestimate actual migration
speeds due to the low frequency of frame acquisition in time-lapse experiments and potential movements
occurring between frames that were not captured.
In this work, we classified cells into two categories - high and low mobility - based on their
behavior during the three-hour observation period. It remains unclear whether these categories represent
distinct phenotypic subpopulations. Given the limited time window, the observed mobility differences
may reflect transient stages within the dynamic surveying behavior, rather than stable phenotypic traits.
At the advanced disease stage, the proportion of highly mobile cells declined, potentially due to
prolonged interactions between myeloid cells and neurons. This observation raises compelling questions
for future investigation, including whether distinct phenotypes emerge among reactive myeloid cells as a
consequence of neuronal interactions, and what the functional implications of such differences might be.
The activation and migration of homeostatic microglia in response to acute neuronal injury are
known to be mediated by calcium signaling [28-31]. In the present study, sustained calcium bursts were
observed in association with the migratory activity of high-mobility myeloid cells, suggesting that
calcium signaling is crucial for migration in the reactive state too. However, unlike the brief calcium
transients typically observed in homeostatic microglia, which last up to a minute, reactive myeloid cells
exhibited elevated calcium levels for extended durations of at least three hours.
In microglia, calcium signaling is regulated in part by the metabotropic P2Y6 receptor [30].
Activation of P2Y6 by its endogenous ligand UDP has been implicated in various microglial functions,
including migration, phagocytosis of damaged neurons, and clearance of apoptotic bodies and Aβ plaques
[32,30,33-35]. In the present study, we found that the P2Y6 antagonist MRS-2578 partially suppressed the
migration of reactive myeloid cells but did not prevent their association with neurons. These findings
suggest that while P2Y6 signaling facilitates the motility of myeloid cells, it does not mediate their
physical interaction with neurons. This points to a specific role for P2Y6 in the surveillance behavior of
reactive myeloid cells under conditions of chronic neurodegeneration. Multiple purinergic receptors,
including P2Y1, P2Y2, P2Y4, P2Y6, P2Y12, P2X4, and P2X7, have been implicated in microglial
chemotaxis, phagocytosis, and directed movement in response to acute injury [25,45]. The observation that
MRS-2578 only partially inhibited cell movement supports the idea that multiple, potentially redundant
pathways govern microglial process motility and soma migration.
A particularly intriguing finding of this study is that reactive myeloid cells maintained elevated
mobility upon acute isolation *in vitro*, even in the absence of external stimuli or neuronal activity. This
indicates that heightened mobility is an intrinsic property of the reactive phenotype, in stark contrast to

homeostatic microglia, which typically require external cues to initiate soma migration. These results
imply that signaling mechanisms beyond purinergic pathways contribute to the enhanced mobility of
reactive myeloid cells. It is possible that this behavior is sustained through autocrine or paracrine
signaling involving proinflammatory factors secreted by reactive microglia themselves. Future studies
will be needed to determine whether such autonomous signaling mechanisms underlie the highly mobile
phenotype.
Quantitatively, the mean speed and total distance traveled by reactive myeloid cells *in vitro* were
two- to threefold greater than those observed in acute brain slices. This is not unexpected, given that in
situ, cells must navigate the extracellular matrix and often pause to establish direct contacts with neurons.
Notably, co-culturing reactive microglia with N2a neuronal cells further increased their mobility even
further. In the presence of N2a cells, reactive microglia exhibited more directed and less stochastic
movement patterns. These observations suggest that, in a reactive state, myeloid cells retain a heightened
responsiveness to environmental cues that do not elicit similar behavioral changes in homeostatic
microglia.
Several limitations of this study warrant discussion. In Cx3cr1/EGFP mice, all myeloid cells,
including monocytes and macrophages, express EGFP, raising questions about the specific identity of the
myeloid cells that establish extensive contacts with neurons. Previously, we demonstrated that myeloid
cells enveloping neurons are positive for P2Y12 [21], a marker generally associated with resident microglia.
Additionally, prior studies have shown that microglial expansion in prion diseases primarily results from
the proliferation of resident microglia, with minimal contribution from peripheral myeloid cell
recruitment [46,47]. Nevertheless, infiltration of peripheral myeloid cells and their potential role in neuronal
surveillance cannot be completely ruled out.
It is unclear whether reactive myeloid cells in other neurodegenerative diseases employ similar
strategies for neuronal surveillance. Furthermore, the molecular mechanisms and docking pathways
mediating the formation of close body-to-body contacts between reactive myeloid cells and neurons are
not yet understood. In earlier work, we found that knockout of CD11b, a component of complement
receptor 3 involved in the phagocytosis of newborn neurons during neurodevelopment, does not affect the
prevalence of neuronal envelopment or the progression of prion diseases [20]. Conversely, knockout of the
P2Y12 receptor, which mediates purinergic junctions between microglial processes and neuronal soma,
led to an increased prevalence of neuronal envelopment [21]. These findings suggest that P2Y12 is not
required for body-to-body contact formation; instead its absence facilitates neuronal envelopment by
microglia.
Cx3cr1 is a chemokine receptor expressed by microglia that binds its neuronal ligand Cx3cl1,
which exists in both membrane-bound and soluble forms [48]. The Cx3cr1-Cx3cl1 axis plays a role in
homeostatic functions such as immune surveillance, chemotaxis and phagocytosis [49]. In the present study,
extensive microglia–neuron body-to-body contacts were observed in Cx3cr1/EGFP mice lacking Cx3cr1,
indicating that this surveillance mechanism and neuronal docking do not depend on Cx3cr1-
Cx3cl1 signaling. While CX3CR1 is known to regulate microglial process dynamics and migration,
whether Cx3cr1 signaling reduce or accelerated microglial motility and mobility remains controversial
[50,51].
To summarize, the transformation of myeloid cells into a reactive phenotype involves a
fundamental reorganization of microglial surveillance strategies. In the homeostatic state, microglia
monitor a limited territory surrounding their location and require chemical cues to initiate migration
toward sites of injury. In contrast, reactive myeloid cells exhibit high intrinsic mobility and actively
search for neurons. These cells adopt a distinct surveillance pattern characterized by migration from one
neuron to another, pausing to form extensive, body-to-body contacts - typically with a single neuron at a
time. Prion-infected animals develop a lethal form of authentic prion disease. The observation of neuronal
envelopment by reactive myeloid cells in both individuals with sporadic CJD and prion-infected mice
underscores that this aspect of human disease pathology is faithfully recapitulated in prion-infected mice.
To our knowledge, this study represents the first report of microglial surveillance behavior in the context
of a bona fide neurodegenerative disease rather than a disease model.

Methods

Animals

B6.129P2(Cg)-*Cx3cr1*[tm1Litt]/J mice (Strain 005582, The Jackson Laboratory) were purchased from the Jackson laboratory and breed in house. B6.129P2(Cg)-*Cx3cr1*[tm1Litt]/J have an enhanced green fluorescent protein (EGFP) sequence replacing the first 390 bp of the coding exon of the chemokine (C-X3-C motif) receptor 1 (*Cx3cr1*) gene.

Brain-derived material for inoculations was prepared from terminally ill SSLOW- or 22L-infected wild type (WT, C57Bl/6J) mice as 10% (w/v) brain homogenate (BH) in PBS, pH 7.4, using glass/Teflon homogenizers attached to a cordless 12 V compact drill [52]. Immediately before inoculation, the inoculum was further dispersed by 30 sec indirect sonication at ~200 watts in a microplate horn of a sonicator (Qsonica, Newtown, CT) and diluted to 1% in PBS, pH 7.4. B6.129P2(Cg)-*Cx3cr1*[tm1Litt]/J and WT mice were inoculated with 200 μl 1% BH intraperitoneally under 3% isoflurane anesthesia. Alternatively, mice were inoculated with 20 μl 1 % BH intracranially (Figure S1a). Animals were scored weekly using four categories, each of which was graded using score '0 – 3', with '3' being the most severe impairment. The scoring categories were: clasping hind legs, posture (kyphosis, rigid tail, rearing difficulties), mobility (difficulties in ambulation and navigating the cage edge), and gate (keeping balance while walking, wobbly gate, disorientation, lethargy). The animals were deemed symptomatic when they displayed a consistent increase in the combined score starting from '4'. The mice were considered terminal when they were unable to rear and/or lost 20% of their weight. The following animal groups were used for main experiments: SSLOW-inoculated B6.129P2(Cg)-*Cx3cr1*[tm1Litt]/J mice at late subclinical (111-113 dpi, dpi), early clinical (125-128 dpi), and advanced stages(162-169 dpi), and non-infected 160-164 days-oldB6.129P2(Cg)-*Cx3cr1*[tm1Litt]/J mice.

Antibodies

Primary antibodies used for immunofluorescence, immunohistochemistry and immunoblotting were as follows: rabbit polyclonal anti-IBA1 (#013-27691, FUJIFILM Wako Chemicals USA; Richmond, VA); goat polyclonal anti-IBA1 (#NB100-1028, Novus, Centennial, CO); mouse monoclonal anti-NeuN, clone A60 (#MAB377, Millipore Sigma, Burlington, MA); mouse monoclonal anti-prion protein, clone SAF-84 (#189775, Cayman, Ann Arbor, MI); rabbit monoclonal anti-prion protein, clone 3D17 (#ZRB1268, Millipore Sigma). The secondary antibodies for immunofluorescence were Alexa Fluor 488-, 546-, and 647-labeled (ThermoFisher Scientific, Waltham, MA).

Acute brain slice preparation and *ex vivo* time-lapse imaging

Before euthanizing an animal, sterile PTFE hydrophilic membrane inserts (PICM0RG50, Sigma) of 0.4 μm pore size were kept in a 6-well plate supplemented with serum free culture media and incubated at 37 °C and 5% CO_2 for 1-2 hours in a standard cell culture incubator. For acute slice preparation, the whole mouse brain was removed from the skull and were immersed in ice cold oxygenated ACSF media (LRE-S-LSG10001, Ecocyte Bioscience LLC) saturated with 95% O_2 and 5% CO_2. The brain was then glued onto to the bottom of the tissue slicing chamber within the Vibratome (Leica VT1200) filled thereafter with oxygenated ACSF media, such that the ventral part of the brain is facing towards us and the dorsal region (surface of cortex) is facing the back of the vibratome. Subsequently, acute coronal cortical sections of 20 μm thickness were prepared using the vibratome at 4.5 mm amplitude, 84-86 Hz frequency and 2.0-2.5 mm/sec speed. Next, the membrane inserts were incubated with fresh complete growth medium containing 50% MEM (INV-42360032, Invitrogen), 25% BME (INV-21010046, Invitrogen), 5% heat inactivated horse serum (INV-26050070, Invitrogen), 10 ng/ml nerve growth factor (NGF) (INV-A42627, Invitrogen) and glial cell line-derived neurotrophic factor (GDNF) (INV-AF-450-44, Invitrogen). The cut sections were then transferred using a sterile pasteur pipette to the inserts. For nuclear staining, slices were stained with 0.5 mkM Hoechst 33342 (INV-H3570, Invitrogen) for 10 minutes followed by successive washing steps in ACSF media. The slices were then transferred to coverslip bottom petri dish (VWR-MSPP-P35G014C-CS, Mattek Corp MS) for their time-lapse imaging in Leica MICA fluorescent microscope (Leica Microsystems). In experiments on P2Y6 inhibition, acute brain slices were incubated for 30 minutes with 2 μM P2Y6 inhibitor MRS-2578 (SIG-M0319, SIGMA)

re-suspended in 0.1% DMSO with complete growth media, or mock alone (0.1 % DMSO). For Ca^{2+}
imaging, slices were incubated simultaneously with 0.5 mkM Hoechst and 0.5 mkM Calbryte 590 AM
(VWR-76484-390-EA, AAT Bioquest) for 45 minutes.
Time-lapse imaging
The time-lapse imaging experiments were performed using environmental climate setup in Leica MICA
for live cell imaging supplemented with 5% CO_2 and 37 °C to ensure slice viability. Image acquisition
parameters for all channels (EGFP, Hoechst and Calbryte 590AM) were kept at 100 ms exposure, with
the constant focusing on the green channel (EGFP) to avoid any focal drift. All the time-lapses were
subjected to a 3D z stacking of optical sections (1024*1024 pixels) with a total of 10 z-stacks at an
average step size of 2 μm from the upper layer of cortex until the bottom of the 20 μm thin cortical brain
slice. The entire images were captured by the 10X objective collecting the frames for every 5 mins time
interval for the complete 3 hour's duration. Background for the videos were then deduced from Leica
MICA using the thunder and lightning module. 3D representations were drawn by Leica MICA software
and image processing of the time lapse videos were further analyzed using ImageJ (FIJI).
Microglial cell tracking and analysis
Time-lapse videos generated by Leica MICA were loaded onto ImageJ (FIJI). Using Manual tracking
(Figure 3a) and MtrackJ plugin (Figure 2d), the individual microglial cells were tracked for 3 hours to
obtain the quantitative differential cell mobility parameters. Briefly, length was measured by the plugin as
the path-length between the first and last current point of the cell track. Accumulated distance was
calculated as the sum of the incremental distance measured from all the frames. Euclidean distance as the
straight-line distance between the frames. Displacement as the change in position from the first to the last
point of the cell track. Mean straight line speed as the total track displacement divided by the net tracking
time. Velocity measured by the ratio of accumulated distance to the total track migration time. Track
mean speed as the average of all the velocities linking the various tracks. Linearity of forward progression
by the ratio between the track mean straight line speed and track mean speed. Finally, the tracking
coordinates for each microglial cell for various groups were represented as ross plots by importing these
values into the Chemotaxis tool software (Ibidi GmbH, Germany).
Analysis of microglial cell mobility, morphology and calcium signal intensity using Trackmate
The datasets from time-lapse imaging in Figures S2, 2a, 2b, 2c, 3b, 3c, 3d, 3e, Figure 6, Figure 7a, 7b and
8a were analyzed using Trackmate 7.13.2 (FIJI). Microglial cell mobility, calcium signal intensity and
distance were detected by the thresholding detector, where auto generated threshold values were used for
each group. Spot filtering was carried out by the quality feature above 42.4. Each spot was linked by the
advanced Kalman tracker algorithm (max frame gap=1, alternative linking cost factor=1.05, Kalman
search radius = 20, linking max distance=5.0, gap closing max distance=15.0, merging max distance=15.0,
cutoff percentile=0.9). For differentiating high and low mobility microglia, track displacement cutoff of
15μm was set based on the average obtained from the mobility values of normal microglia (displacement
< 15μm; low and displacement ≥ 15μm; high). The mean directional change rate was calculated by
averaging the successive angle values for the entire frame. For each spot, the different morphological
parameters such as area, radius, perimeter and shape index were also generated. Track results were then
exported and further condensed for each cell by statistical analysis.
Confocal microscopy and 3D image reconstruction
Confocal images were acquired with Leica TCS SP8 microscope using laser lines 405, 488, 552, the
40×/1.30 oil immersion objective, the resolution of 1024×1024 pixels, and a scan speed of 400 Hz. For
3D reconstruction, the system-optimized number of steps was used. Images were processed using the
LAS X and ImageJ software.
Western blot
For Western blots, 10% (w/v) brain homogenates (BH) were prepared as previously described [52] using
RIPA Lysis Buffer (Millipore Sigma, St. Louis, MO). To analyze brain-derived PrP^Sc, BH aliquots were
diluted with RIPA buffer to achieve 5% BH final concentration and treated with 20 μg/ml proteinase K
(New England BioLabs) in the presence of 50 mM Tris, pH 7.5, and 2% Sarcosyl, for 30 min at 37°C. To
analyze other proteins, BH was diluted with RIPA buffer to 1% and proteinase digestion was omitted.

The resulting samples were supplemented with 4xSDS loading buffer and heated for 10 min in a boiling water bath before loading onto NuPAGE 12% Bis-Tris gels. Wet transfer onto PVDF membranes and probing of Western blots was done according to standard procedures. The signals were visualized by Immobilon Forte Western HRP Substrate (Millipore Sigma, Rockfield, MD) or SuperSignal West pico PLUS Chemiluminescent Substrate (Thermo Scientific, Rockford, IL) using Invitrogen iBright 1500 imager, and quantified with iBright Analysis software (Thermo Scientific, Rockford, IL). Intensity data were presented as normalized by actin, except for PrP^{Sc} Western blots treated with protease K.

Tracking and analysis of acutely isolated cells

Mouse microglial cells were isolated using Adult Brain Dissociation Kit with CD11b antibodies according to the manufacturer protocol (Miltenyi Biotec, #130-107-677). For non-infected B6.129P2(Cg)-*Cx3cr1^{tm1Litt}*/J mice, microglia were purified from pools of three cortices, which ensured enough cells. For animals infected with SSLOW (1% BH via i.p. route), microglia were purified from individual cortices of six mice at the clinical stage of the disease. Cells were counted and plated into wells of 24-well plates and left overnight to settle and recover from enzyme treatment. The next morning N2a cells were added to designated well at 1:1 ratio and live images were taken on green channel at 10x magnification for 6 hours with 5 min intervals. During the whole imaging session, cells were kept at 37 °C in an atmosphere of 5% CO_2. For microglia motility analysis, TrackMate plugin for Fiji ImageJ software was used. All individual tracks were manually validated for the accuracy of automated tracking and corrected if needed.

Study approval

The study was carried out in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. The animal protocol was approved by the Institutional Animal Care and Use Committee of the University of Maryland, Baltimore (Assurance Number: A32000-01; Permit Number: 0215002).

Statistical Analysis

All statistical analysis and graph plotting were performed using GraphPad Prism software, version 10.1.1 for Windows, as described in the figure legends.

Acknowledgements.
We thank Kara Molesworth for helping with animal procedures. Financial support for this study was provided by National Institute of Health Grants R01 NS045585 and R01 NS129502 to IVB.

References

Cserép, C. *et al.* Microglia monitor and protect neuronal function through specialized somatic purinergic junctions. *Science* **367**, 528-537, doi:10.1126/science.aax6752 (2020).
Cserép, C. *et al.* Microglial control of neuronal development via somatic purinergic junctions. *Cell Rep* **40**, 111369, doi:10.1016/j.celrep.2022.111369 (2022).
Nebeling, F. C. *et al.* Microglial motility is modulated by neuronal activity and correlates with dendritic spine plasticity in the hippocampus of awake mice. *eLife* **12**, e83176, doi:10.7554/eLife.83176 (2023).
Rotterman, T. M. & Alvarez, F. J. Microglia Dynamics and Interactions with Motoneurons Axotomized After Nerve Injuries Revealed By Two-Photon Imaging. *Sci Rep* **10**, 8648, doi:10.1038/s41598-020-65363-9 (2020).
Damani, M. R. *et al.* Age-related alterations in the dynamic behavior of microglia. *Aging Cell* **10**, 263-276, doi:10.1111/j.1474-9726.2010.00660.x (2011).
Makarava, N., Chang, J. C.-Y., Molesworth, K. & Baskakov, I. V. Region-specific glial homeostatic signature in prion diseases is replaced by a uniform neuroinflammation signature, common for brain regions and prion strains with different cell tropism. *Neurobiology of Disease* **137**, e104783, doi:https://doi.org/10.1016/j.nbd.2020.104783 (2020).

7   Butovsky, O. & Weiner, H. L. Microglial signatures and their role in health and disease. *Nature*
Reviews Neuroscience **19**, 622-635, doi:10.1038/s41583-018-0057-5 (2018).
8   Krasemann, S. *et al.* The TREM2-APOE Pathway Drives the Transcriptional Phenotype of
Dysfunctional Microglia in Neurodegenerative Diseases. *Immunity* **47**, 566-581.e569,
doi:https://doi.org/10.1016/j.immuni.2017.08.008 (2017).
9   Keren-Shaul, H. *et al.* A Unique Microglia Type Associated with Restricting Development of
Alzheimer's Disease. *Cell* **169**, 1276-1290.e1217, doi:https://doi.org/10.1016/j.cell.2017.05.018
(2017).
10   Zrzavy, T. *et al.* Loss of 'homeostatic' microglia and patterns of their activation in active multiple
sclerosis. *Brain* **140**, 1900-1913, doi:10.1093/brain/awx113 (2017).
11   Kenkhuis, B. *et al.* Co-expression patterns of microglia markers Iba1, TMEM119 and P2RY12 in
Alzheimer's disease. *Neurobiol Dis* **167**, 105684, doi:10.1016/j.nbd.2022.105684 (2022).
12   Holtman, I. R. *et al.* Induction of a common microglia gene expression signature by aging and
neurodegenerative conditions: a co-expression meta-analysis. *Acta Neuropathologica*
Communications **3**, 31, doi:10.1186/s40478-015-0203-5 (2015).
13   Maeda, J. *et al.* Distinct microglial response against Alzheimer's amyloid and tau pathologies
characterized by P2Y12 receptor. *Brain Communications* **3**, doi:10.1093/braincomms/fcab011
(2021).
14   Prusiner, S. B. Prions. *Proc Natl Acad Sci U S A* **95**, 13363-13383 (1998).
15   Legname, G. *et al.* Synthetic mammalian prions. *Science* **305**, 673-676 (2004).
16   Makarava, N., Savtchenko, R., Lasch, P., Beekes, M. & Baskakov, I. V. Preserving prion strain
identity upon replication of prions in vitro using recombinant prion protein. *Acta*
Neuropathologica Communications **6**, 92, doi:10.1186/s40478-018-0597-y (2018).
17   Makarava, N. *et al.* Region-Specific Homeostatic Identity of Astrocytes Is Essential for Defining
Their Response to Pathological Insults. *Cells* **12**, doi:10.3390/cells12172172 (2023).
18   Watts, J. C. & Prusiner, S. B. Mouse Models for Studying the Formation and Propagation of
Prions. *Journal of Biological Chemistry* **289**, 19841-19849, doi:10.1074/jbc.R114.550707 (2014).
19   Jeffrey, M. *et al.* Pathology of SSLOW, a transmissible and fatal synthetic prion protein disorder,
and comparison with naturally occurring classical transmissible spongoform encephalopathies.
Neuropath.Appl.Neurobiol. **40**, 296-310 (2014).
20   Makarava, N. *et al.* Reactive microglia partially envelop viable neurons in prion diseases. *J Clin*
Invest **134**, doi:10.1172/jci181169 (2024).
21   Makarava, N. *et al.* Knockout of P2Y12 receptor facilitates microglia-neuron body-to-body
interactions and accelerates prion disease. *bioRxiv*, 2025.2004.2007.647619,
doi:10.1101/2025.04.07.647619 (2025).
22   Striebel, J. F., Race, B., Carroll, J. A., Phillips, K. & Chesebro, B. Knockout of fractalkine
receptor Cx3cr1 does not alter disease or microglial activation in prion-infected mice. *J. Gen.*
Virol. **97**, 1481-1487 (2016).
23   Makarava, N., Chang, J. C.-Y., Molesworth, K. & Baskakov, I. V. Posttranslational modifications
define course of prion strain adaptation and disease phenotype. *The Journal of Clinical*
Investigation **130**, 4382-4395, doi:10.1172/JCI138677 (2020).
24   Makarava, N., Mychko, O., Chang, J. C.-Y., Molesworth, K. & Baskakov, I. V. The degree of
astrocyte activation is predictive of the incubation time to prion disease. *Acta Neuropathologica*
Communications **9**, 87, doi:10.1186/s40478-021-01192-9 (2021).
25   Smolders, S. M. *et al.* Microglia: Brain cells on the move. *Prog Neurobiol* **178**, 101612,
doi:10.1016/j.pneurobio.2019.04.001 (2019).
26   Delbridge, A. R. D. *et al.* Organotypic Brain Slice Culture Microglia Exhibit Molecular Similarity
to Acutely-Isolated Adult Microglia and Provide a Platform to Study Neuroinflammation. *Front*
Cell Neurosci **14**, 592005, doi:10.3389/fncel.2020.592005 (2020).

27 Ferrucci, L. *et al.* Time-dependent phenotypical changes of microglia drive alterations in
hippocampal synaptic transmission in acute slices. *Front Cell Neurosci* **18**, 1456974,
doi:10.3389/fncel.2024.1456974 (2024).
28 Eichhoff, G., Brawek, B. & Garaschuk, O. Microglial calcium signal acts as a rapid sensor of
single neuron damage in vivo. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*
1813, 1014-1024, doi:https://doi.org/10.1016/j.bbamcr.2010.10.018 (2011).
29 Umpierre, A. D. *et al.* Microglial calcium signaling is attuned to neuronal activity in awake mice.
eLife **9**, e56502, doi:10.7554/eLife.56502 (2020).
30 Umpierre, A. D. *et al.* Microglial P2Y(6) calcium signaling promotes phagocytosis and shapes
neuroimmune responses in epileptogenesis. *Neuron* **112**, 1959-1977.e1910,
doi:10.1016/j.neuron.2024.03.017 (2024).
31 Du, T., Zhou, X., Zhang, R. D. Y. & Du, X. F. Roles of Ca(2+) activity in injury-induced
migration of microglia in zebrafish in vivo. *Biochem Biophys Rep* **32**, 101340,
doi:10.1016/j.bbrep.2022.101340 (2022).
32 Neher, J. J., Neniskyte, U., Hornik, T. & Brown, G. C. Inhibition of UDP/P2Y6 purinergic
signaling prevents phagocytosis of viable neurons by activated microglia in vitro and in vivo.
Glia **62**, 1463-1475, doi:10.1002/glia.22693 (2014).
33 Puigdellívol, M. *et al.* The microglial P2Y(6) receptor mediates neuronal loss and memory
deficits in neurodegeneration. *Cell Rep* **37**, 110148, doi:10.1016/j.celrep.2021.110148 (2021).
34 Milde, S. *et al.* Inflammatory neuronal loss in the substantia nigra induced by systemic
lipopolysaccharide is prevented by knockout of the P2Y(6) receptor in mice. *J*
Neuroinflammation **18**, 225, doi:10.1186/s12974-021-02280-2 (2021).
35 Koizumi, S. *et al.* UDP acting at P2Y6 receptors is a mediator of microglial phagocytosis. *Nature*
446, 1091-1095, doi:10.1038/nature05704 (2007).
36 Berki, P. *et al.* Microglia contribute to neuronal synchrony despite endogenous ATP-related
phenotypic transformation in acute mouse brain slices. *Nat Commun* **15**, 5402,
doi:10.1038/s41467-024-49773-1 (2024).
37 Haynes, S. E. *et al.* The P2Y12 receptor regulates microglial activation by extracellular
nucleotides. *Nature Neuroscience* **9**, 1512-1519, doi:10.1038/nn1805 (2006).
38 Maeda, M., Tsuda, M., Tozaki-Saitoh, H., Inoue, K. & Kiyama, H. Nerve injury-activated
microglia engulf myelinated axons in a P2Y12 signaling-dependent manner in the dorsal horn.
Glia **58**, 1838-1846, doi:10.1002/glia.21053 (2010).
39 Fekete, R. *et al.* Microglia control the spread of neurotropic virus infection via P2Y12 signalling
and recruit monocytes through P2Y12-independent mechanisms. *Acta Neuropathol* **136**, 461-482,
doi:10.1007/s00401-018-1885-0 (2018).
40 Ohsawa, K. *et al.* Involvement of P2X4 and P2Y12 receptors in ATP-induced microglial
chemotaxis. *Glia* **55**, 604-616, doi:10.1002/glia.20489 (2007).
41 Eyo, U. B. *et al.* P2Y12R-Dependent Translocation Mechanisms Gate the Changing Microglial
Landscape. *Cell Rep* **23**, 959-966, doi:10.1016/j.celrep.2018.04.001 (2018).
42 Lively, S. & Schlichter, L. C. The microglial activation state regulates migration and roles of
matrix-dissolving enzymes for invasion. *J Neuroinflammation* **10**, 75, doi:10.1186/1742-2094-10-
75 (2013).
43 Smolders, S. M. *et al.* Age-specific function of α5β1 integrin in microglial migration during early
colonization of the developing mouse cortex. *Glia* **65**, 1072-1088, doi:10.1002/glia.23145 (2017).
44 Eyo, U. B., Miner, S. A., Weiner, J. A. & Dailey, M. E. Developmental changes in microglial
mobilization are independent of apoptosis in the neonatal mouse hippocampus. *Brain Behav*
Immun **55**, 49-59, doi:10.1016/j.bbi.2015.11.009 (2016).
45 Franco-Bocanegra, D. K., McAuley, C., Nicoll, J. A. R. & Boche, D. Molecular Mechanisms of
Microglial Motility: Changes in Ageing and Alzheimer's Disease. *Cells* **8**,
doi:10.3390/cells8060639 (2019).

46   Gomez-Nicola, D., Fransen, N. L., Suzzi, S. & Perry, V. H. Regulation of microglial proliferation
during chronic neurodegeneration. *J Neurosci* **33**, 2481-2493 (2013).
47   Gómez-Nicola, D., Schetters, S. T. & Perry, V. H. Differential role of CCR2 in the dynamics of
microglia and perivascular macrophages during prion disease. *Glia* **62**, 1041-1052,
doi:10.1002/glia.22660 (2014).
48   Bazan, J. F. *et al.* A new class of membrane-bound chemokine with a CX3C motif. *Nature* **385**,
640-644, doi:10.1038/385640a0 (1997).
49   Raoul, W. *et al.* Role of the chemokine receptor CX3CR1 in the mobilization of phagocytic
retinal microglial cells. *J Neuroimmunol* **198**, 56-61, doi:10.1016/j.jneuroim.2008.04.014 (2008).
50   Liang, K. J. *et al.* Regulation of dynamic behavior of retinal microglia by CX3CR1 signaling.
Invest Ophthalmol Vis Sci **50**, 4444-4451, doi:10.1167/iovs.08-3357 (2009).
51   Wagner, J. *et al.* CX3CR1 modulates migration of resident microglia towards brain injury.
bioRxiv, 2024.2009.2023.614458, doi:10.1101/2024.09.23.614458 (2024).
52   Makarava, N. *et al.* Stabilization of a prion strain of synthetic origin requires multiple serial
passages. *J Biol Chem* **287**, 30205-30214 (2012).
672
673

675
Figure 1. Iba1$^+$ cells envelop neurons in SSLOW-infected brains. a. Immunostaining for microglia
(IBA1, red) and neurons (NeuN, green) showing neuronal envelopment by myeloid cells in Cx3cr1/EGFP
mice infected by SSLOW via intraperitoneal route at the terminal stage of the disease. **b.** Immunostaining
for microglia (IBA1, red) and neurons (NeuN, green) of non-infected, age-matched Cx3cr1/EGFP mice.
Confocal microscopy imaging followed by 3D reconstruction was used for both **a** and **b**.
681

682
Figure 2. Reactive myeloid cells are highly mobile. Brain slices were prepared acutely using non-
infected Cx3cr1/EGFP (normal) mice, or Cx3cr1/EGFP mice infected with SSLOW via ip route and
examined at sub-clinical, early clinical or advanced stages of the disease. Analysis of mean speed (**a**) and
total distance (**b**) traveled by individual EGFP+ cells over 3-hour period. The midline of the box-and-
whisker plot denotes the median, the + represents the mean, and the ends of the box plot denote the 25th
and 75th percentiles. **c.** Examples of tracks recorded from EGFP+ cells. Colored lines represent tracks of
individual cells recorded within 3-hour period. **d.** Principal component analysis of mobility parameters.
n=70-90 cells per group, *p<0.05, **p<0.01, ****p<0.0001, ns - non-significant by non-parametric
Kruskal-Wallis test with Dunn's multiple comparison.
692

693
694
Figure 3. High- and low-mobility behavioral patterns of myeloid cells in prion-infected brains.
Acute brain slices were prepared using normal, non-infected Cx3cr1/EGFP mice, or SSLOW-infected
Cx3cr1/EGFP mice and examined at the sub-clinical, early clinical or advanced stages of the disease. **a**
Rose plot of individual cell trajectories. In brain slices from SSLOW-infected animals, EGFP⁺ cells
showed two behavioral patterns – with high and low mobility. **b** Change in the percentage of high-
mobility EGFP⁺ cells with the disease progression. N=3 animals per group. The data presented as Means
± SD, **p<0.01, ****p<0.0001, ns - non-significant by Tukeys' multiple comparisons test. **c, d, e**
Analysis of mean speed (**c**), total distance travelled over 3-hour period (**d**), and mean directional change
rate (**e**) for individual high- and low-mobility EGFP⁺ cells in slices from SSLOW-infected mice at three
disease stages, and normal mice. The midline of the box-and-whisker plot denotes the median, the +
represents the mean, and the ends of the box plot denote the 25th and 75th percentiles. n=25-30 cells per
group, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001, ns - non-significant by non-parametric Kruskal-
Wallis test with Dunn's multiple comparison.
708

709
710
Figure 4. Behavioral patterns of high-mobility reactive myeloid cells. Time-lapse imaging of acute
brain slices of SSLOW-infected Cx3cr1/EGFP mice captured at the early clinical (**a**), advanced (**b, d**),
and subclinical (**c**) stages of disease. **a** (Video S3): An EGFP[+] cell (indicated by an arrow) extends a
process toward neurons #1 and #2, migrates along this process, and subsequently envelops both neurons
while simultaneously extending processes toward neurons #3, #4, and #5. **b** (Video S10): EGFP[+] cell #1
migrates toward and surveys a neuron (circled), then departs. Subsequently, a second EGFP[+] cell (#2)
migrates to and interacts with the same neuron. White and blue curves trace the respective migration
paths of the two cells. **c** Upper panels: An EGFP[+] cell approaches, envelops, and then retracts from a
neuron. Middle panels: A migrating EGFP[+] cell (indicated by an arrow) surveys five distinct neurons
(circled), interacting simultaneously with neurons #1 and #2, followed by #3 and #4. Lower panels: An
EGFP[+] cell maintains prolonged contact with a neuron. **d** (Video S5): An EGFP[+] cell (arrow) migrates
across the field, initially extending processes (arrowhead) toward a neuron. The cell body then
translocates along these processes, briefly contacts one neuron, and continues movement toward another.

All videos were recorded over a 3-hour period with 5-minute intervals. Nuclei were visualized using
Hoechst staining. Scale bars = 20 μm.
726

727
728
Figure 5. Behavioral patterns of low-mobility reactive myeloid cells. Time-lapse imaging of acute
brain slices of SSLOW-infected Cx3cr1/EGFP mice, captured at early clinical (**a, b**) and advanced (**c**)
stages of disease. **a** (Video S8): An EGFP$^+$ cell exhibits prolonged interactions simultaneously with two
neurons. **b** (Video S3): An EGFP$^+$ cell shows sustained envelopment of neuron #1, then initiates
simultaneous envelopment of neuron #2 while maintaining contact with neuron #1 and possibly neuron
#3. **c** An EGFP$^+$ cell moves toward neurons #1 and #2, partially envelops neuron #1, and fully envelops
neuron #2. All videos were recorded over a 3-hour period with 5-minute intervals. Nuclei were visualized
using Hoechst staining. Scale bars = 20 μm. **d, e** 3D reconstructions from confocal microscopy images of
fixed brain slices immunostained for myeloid cells (IBA1, red) and neurons (NeuN, green) in WT mice

infected with SSLOW via the intraperitoneal route, analyzed at early clinical and advanced disease stages.
Scale bar = 25 μm.
740
741

742
743
Figure 6. Analysis of cell morphology. Acute brain slices were prepared from normal, non-infected
Cx3cr1/EGFP mice and from SSLOW-infected Cx3cr1/EGFP mice at the at sub-clinical, early clinical or
advanced stages of the disease. Analysis of cell radius (**a**), cell area (**b**), cell perimeter (**c**) and shape index
(**d**) of high- and low-mobility EGFP$^+$ cells in slices from SSLOW-infected mice at three disease stages,
and normal mice. The midline of the box-and-whisker plot denotes the median, the + represents the mean,
and the ends of the box plot denote the 25th and 75th percentiles. n=30-90 cells per group, *p<0.05,
**p<0.01, ***p<0.01, ****p<0.0001, ns - non-significant by non-parametric Kruskal-Wallis test with
Dunn's multiple comparison.

752
753
754
755

756
757
Figure 7. Sustained Ca²⁺ bursts in high-mobility myeloid cells. Acute brain slices were prepared from
non-infected Cx3cr1/EGFP mice and from SSLOW-infected Cx3cr1/EGFP mice at the advanced stage of
disease. **a** Quantification of signal intensity of Ca²⁺ puncta within high- or low-mobility EGFP⁺ cells.
Data from normal, non-infected mice are shown for reference. Sustained Ca²⁺ bursts were detected using
Calbryte-590 AM and averaged per cell over a 3-hour period. N = 24-28 cells per group. *p < 0.05, **p <
0.01, ns = not significant by non-parametric Kruskal-Wallis test with Dunn's multiple comparisons.
b Changes in Ca²⁺ signal intensity in individual EGFP⁺ cells over the 3-hour imaging session. Images
were acquired at 5-minute intervals. **c** Time-lapse imaging of acute brain slice from a SSLOW-infected
Cx3cr1/EGFP mouse recorded over 3 hours. Upper panels (Video S19): EGFP⁺ cell (#1) envelops a
neuronal soma and exhibits low Ca²⁺ activity. Lower panels: highly mobile EGFP⁺ cell (#2) displays
sustained somatic Ca²⁺ bursts. Scale bars = 50 μm.

770
771
772
Figure 8. Inhibition of the P2Y6 receptor reduces motility of reactive myeloid cells. Acute brain
slices were prepared from non-infected Cx3cr1/EGFP mice and from SSLOW-infected Cx3cr1/EGFP
mice at the early clinical stage of disease. **a, b** Quantification of mean speed (**a**) and total distance
traveled (**b**) by individual EGFP+ cells over a 3-hour period in brain slices from SSLOW-infected mice
treated with either MRS-2578 (2 μM) or vehicle control. Data from normal, non-infected mice are shown
for reference. n=100-250 cells per group. **p < 0.001, ***p < 0.0001 by Tukey's multiple comparisons
test. **c**, Time-lapse imaging of MRS-2578-treated acute brain slice from a SSLOW-infected Cx3cr1/EGFP
mouse recorded over 3 hours. The EGFP+ cell exhibits bidirectional movement between two neuronal
somas (labeled #1 and #2), sequentially enveloping each soma. Scale bars = 20 μm.
782

783
Figure 9. *In vitro* analysis of CD11b⁺ cell mobility. CD11b⁺ cells were acutely isolated
from Cx3cr1/EGFP mice at the clinical stage of SSLOW infection or from non-infected
Cx3cr1/EGFP mice. **a,b,c** Quantification of cell mobility parameters, including mean speed (**a**), total
track distance (**b**), and mean directional change rate (**c**) of CD11b⁺/EGFP⁺ cells over a 3-hour period, in
the presence or absence of N2a cells. In box-and-whisker plots, the midline indicates the median, the "x"
denotes the mean, and the box limits represent the 25^{th} and 75^{th} percentiles. *n* = 178-354 cells per group.
Statistical significance: p < 0.05; p< 0.01; *p < 0.001; **p < 0.0001; ns, not significant by non-parametric
Kruskal-Wallis test with Dunn's multiple comparison. (d) Representative cell tracks of CD11b⁺/EGFP⁺
cells over 3 hours. Colored lines indicate trajectories of individual cells.