Growth phase influences virulence in *Candidozyma auris* systemic infection models

Michael J. McFadden[1], Juliet A.E. Anku[1], Faith A. Davis[1], Catherine Luke[2], Andrea Obi[2], Teresa

R. O'Meara[1,†]

Affiliations:

1   Department of Microbiology and Immunology, University of Michigan, Ann Arbor, MI

48109, USA.

2   Section of Vascular Surgery, Department of Surgery, University of Michigan, Ann Arbor,

MI 48109, USA.

[†] Corresponding author: Correspondence to Teresa R. O'Meara (tromeara@umich.edu)

Abstract

Candidozyma auris is a growing public health concern, capable of causing long-term

contamination of healthcare settings, skin colonization, and life-threatening bloodstream

infections. However, *C. auris* pathogenesis is not well understood, which is exacerbated by

limitations and discrepancies in existing animal infection models. Further, the effects of *C. auris*

growth phase on virulence have not been examined, despite growth phase being linked to

virulence in many bacterial species. To address this question, and to develop an

immunocompetent murine model of infection, we directly compared log and stationary phase *C.*

auris systemic infection in immunocompetent C57BL/6J mice at high and low doses of infection.

Systemic infection with high dose log phase *C. auris* results in rapid mortality between 2 hours

and 1 day post infection, whereas stationary phase *C. auris* results in significantly extended

survival. However, at low doses of infection, there was no difference in mortality kinetics

between log and stationary phase cells. We observed that *C. auris* initially colonizes multiple

organs but is rapidly cleared from the lungs and spleen, while kidney fungal burdens remain

stable. Mice infected with high dose log phase *C. auris* had Fibrin-associated blood clotting in

multiple organs and decreased serum Fibrinogen levels, suggesting that coagulation may drive

rapid mortality. This was associated with increased β-glucan exposure and mannan abundance

in log phase *C. auris*. These results will inform the development of a more standardized animal

model of systemic *C. auris* infection, which can be used to reveal key aspects of *C. auris*

pathogenesis.

Importance

Despite its growing medical importance, there is limited understanding of *Candidozyma*

auris pathogenesis, due in part to limitations of existing laboratory models of infection. To

develop a more complete understanding of factors that contribute to *C. auris* pathogenesis, it

will be necessary to establish consistent parameters for animal models of infection. To address

this need, we directly compared log and stationary growth phases on *C. auris* pathogenesis in

immunocompetent C57BL/6J mice using a single virulent Clade I isolate. At a high dose of

infection, host survival was dramatically different between log or stationary phase *C. auris*,

suggesting that growth phase can affect *C. auris* pathogenesis. These differences correlated

with increased exposure of pathogen-associated molecular patterns in the *C. auris* cell wall in

log phase cells. These results will be instrumental in the future development of standardized

animal models to study *C. auris* pathogenesis.

Introduction

Candidozyma auris is an often multidrug resistant invasive species within healthcare

settings that also can cause systemic infection with mortality rates typically ranging from 30-

60% (1, 2). *C. auris* pathogenesis is not well understood due to its recent emergence and the

existence of comorbidities in many infected patients, as well as an incomplete understanding of

host-pathogen interactions and pathology within sites of *C. auris* infection (3–5). While animal

models have been of great utility in understanding the relative contribution of specific fungal

mutants to virulence (6–9), there is not currently a standard animal model for *C. auris*

disseminated disease. The immunocompromised model of *C. auris* was initially developed as a

physiologically relevant model of systemic infection, with a particular focus on its utility for

testing of antifungal compounds (10) or specific types of immunocompromise (11).

Immunocompetent models of *C. auris* infection have also been developed in ICR outbred mice

(12), BALB/c (13), and C57BL/6J (3, 4) backgrounds, with varying kinetics and rates of

mortality. Additionally, *C. auris*-specific factors, such as strain background are likely a major

source of variability, as evidenced by our recent work showing significant differences in

virulence between two closely related clade I isolates (6, 14).

Beyond strain differences, work in bacterial pathogenesis has demonstrated that growth

phase can also influence virulence of a single strain. However, the effects of growth phase are

species-specific and were determined empirically. For example, for *Legionella* and *Brucella*,

entry into stationary phase is associated with increases in virulence factor expression (15, 16),

but for *Salmonella* and *Streptococci*, it is the exponential growth phase that is associated with

virulence (17, 18). Therefore, we sought to determine the effects of *C. auris* growth phase and

dosage on host survival and pathology using a single strain of *C. auris* and a single genetically

tractable and immunocompetent murine model of systemic infection.

At a high dose of systemic *C. auris* infection, we observed that log/exponential phase

fungi cause rapid mortality compared to stationary phase fungi, which cause mortality over the

course of several days. These differences in mortality may stem from rapid extensive blood

clotting caused by log phase *C. auris*, as we observed decreased serum Fibrinogen levels and

increased presence of blood clotting and fibrin deposition in multiple organs. *In vitro*

exponentially growing *C. auris* cells had increased β-glucan exposure and mannan abundance,

potentially promoting detection by host cells and triggering blood clotting and rapid mortality.

However, differences in mortality observed at a high dose of infection were ablated at a low

dose of infection, in which mice survived over the course of multiple weeks. We recovered *C.*

auris from the lungs, spleen, and kidneys early after infection, but over time, only the kidneys

maintained a substantial fungal burden. Together, these results suggest that growth phase can

have dramatic effects on survival during systemic *C. auris* infection. Additionally, our work

provides new insight into *C. auris* disease progression and will be helpful in establishing more

standardized approaches to modeling *C. auris* infection in immunocompetent mouse models.

Results

Comparing the Effect of *Candidozyma auris* Growth Phase on Survival Kinetics and

Fungal Burden in Organs

To assess whether growth phase affects virulence during systemic *C. auris* infection*,* we

compared infection outcomes in an immunocompetent murine model using log or stationary

phase *C. auris* cultures at previously published high and low doses of infection (5×10^7 and

1×10^6, respectively) using the virulent AR0382 (B11109 clade 1) isolate (Fig. 1A). Following

intravenous infection, each group was monitored for several hours for the onset of disease

symptoms. A majority of the cohort infected with a high dose of log phase *C. auris* rapidly

declined in health, becoming moribund and showing labored breathing within 2 hours of

infection, resulting in humane sacrifice. None of the mice in this group survived beyond the first

day post-infection (Fig. 1B). In contrast, mice infected with the same high dose of stationary

phase *C. auris* survived significantly longer than their log phase-infected counterparts, with

onset of mortality starting at day 2 post-infection and full cohort mortality only occurring at day 6

post-infection (Fig. 1B). At the lower dose of infection, we did not observe any significance in

mortality between the log and stationary phase infection cohorts (Fig. 1C). These data suggest

that the growth phase of *C. auris* affects pathogenesis in murine infection models specifically at

a high dose of infection.

We measured the fungal burden in the lungs, spleen, and kidneys post-mortality in each

cohort. In the high dose log phase *C. auris*-infected cohort, which had very rapid mortality, we

observed a significantly higher fungal burden in the lungs than in the stationary phase cohort,

while there was no difference in burden in the kidneys or spleen (Fig 2A-C). However, we

hypothesized that the difference observed in lung fungal burden is likely a product of survival

kinetics, rather than colonization, as the fungal burden in the lungs appeared to sharply

decrease over time (Fig. 2D). To test this, we performed a matched comparison of *C. auris*

burden in the lungs at 2 hours post-high dose infection for both log and stationary phase and observed similar burden between both cohorts (Fig. 2G). Additionally, in low dose cohorts, which survived significantly longer than high dose cohorts, we recovered very few fungal colonies from the lungs from both cohorts, indicating that *C. auris* is effectively cleared from the lungs over time (Fig. 2H).

Similarly, fungal burden in the spleen showed a decreasing trend at day 7 in the high dose cohort (Fig. 2E) and was generally low at the time of mortality in the low dose cohorts (Fig. 2I), consistent with clearance from these organs over time. In contrast, fungal burden in the kidneys remained consistent over time in both the high dose (Fig. 2F) and low dose cohorts (Fig. 2J). Together, these data suggest that *C. auris* disseminates to multiple organs after systemic infection but is cleared from the lungs and spleen, while fungal burden remains steady over time in the kidneys.

High Dose Systemic Infection with Log Phase *C. auris* Causes Rapid Blood Clotting.

We next sought to further understand the differences in mortality observed between log and stationary phase *C. auris* during systemic infection at a high dose. We noted animals that rapidly succumbed within the log phase high dose cohort exhibited blood clotting while performing cardiac puncture. Coagulation is associated with septic shock, which has a high mortality rate in the absence of intensive care and is associated with increased mortality in candidemia patients (1, 19). ELISA data from the 2 hour post-infection mice showed decreased Fibrinogen levels in the plasma of animals infected with either high dose log or stationary phase *C. auris*, compared to control mice (Fig. 3A), consistent with Fibrinogen being converted to insoluble Fibrin to form clots (20). However, animals infected with stationary phase cells recovered from this initial response and showed higher Fibrinogen levels at the time of mortality (Fig. 3B). Additionally, using Martius Scarlet Blue (MSB) staining in which Fibrin stains red, erythrocytes stain yellow, and muscle and collagen stain blue, we observed increased Fibrin

deposition only in the log phase-infected animals, suggesting that the coagulation cascade

proceeded in this cohort to trigger systemic thrombosis (Fig. 3C-F; Fig. S1). Quantification of

Fibrin staining revealed that only log phase-infected kidneys showed significantly higher Fibrin

levels relative to erythrocyte abundance in MSB-stained kidney sections (Fig. 3G). The mock

and stationary-phase kidneys at both time points were not statistically different (Fig. 3G). Blood

clots could also be observed in MSB-stained lungs from the log phase infected cohort at the

time of mortality (Fig. S2), consistent with systemic thrombosis in log phase-infected animals,

rather than in a specific organ.

To determine whether a proinflammatory response drives mortality and coagulation in

the high dose log phase-infected cohort, we measured the levels of proinflammatory cytokines

in plasma samples from the high dose infection cohorts at a synchronized time of 2 hours post-

infection (hpi) and post-mortality. We did not observe consistent signs of systemic inflammation

in either log or stationary phase high dose infection cohorts at 2 hpi, as TNF levels were only

elevated in stationary phase infected mice (Fig. 4A), IL-6 levels were only elevated in log phase

infected mice (Fig. 4B), and IL-1β levels were unchanged compared to mock-infected control

mice (Fig. 4C). Proinflammatory cytokine levels were also similar at the time of mortality

between log and stationary phase high dose cohorts (Fig. 4D-F), again consistent with blood

clotting, rather than a cytokine storm, driving rapid mortality after systemic infection with log

phase *C. auris*. Hematoxylin and eosin (H&E)-stained lungs showed blood clots specifically in

log phase-infected lungs, but not dramatic immune cell recruitment (Fig. 4G), consistent with

these findings. Together, these data suggest that rapid and aberrant systemic coagulation

causes mortality after infection with high doses of log phase *C. auris*.

We then hypothesized that *C. auris* growth phase would affect the composition of the cell

wall, triggering differential recognition and leading to systemic coagulation. Therefore, we

measured the major cell wall components and fungal pathogen-associated molecular patterns

(PAMPs) chitin, mannan, and exposed β-glucan in log or stationary phase *C. auris*. Flow

cytometry analysis revealed that log phase *C. auris* showed higher β-glucan exposure than

stationary phase (Fig. 5A, D). Interestingly, there were two distinct populations of high and low

mannan cells in the log phase cells, compared to stationary phase cells, which showed only a

single peak with intermediate intensity (Fig. 5B, E). In contrast, chitin levels were similar

between log and stationary phase cells (Fig. 5C, F). Similar results were observed through

confocal microscopy, which revealed increased β-glucan exposure on log phase *C. auris* cells

and a subset of cells with increased mannan staining enriched on the cell periphery. Chitin

levels were heterogeneous between cells but overall levels did not differ between stationary and

log phase cells (Fig. 5G). These results reveal differences in the composition of the cell wall and

exposure of pathogen associated molecular patterns in log phase *C. auris* cells, which may

drive the immunopathological response and mortality observed during systemic infection at a

high dose.

Discussion

Here, we show that the growth phase of *C. auris* influences its virulence in a murine

model during bloodstream infection at a high dose of infection. Notably, the inoculum used in

our high dose cohort has been commonly used in previous reports for *in vivo* models of *C. auris*

pathogenesis and survival analyses (6, 8, 9). Therefore, growth phase will be an important

parameter to consider for future research on *C. auris* pathogenesis and a crucial consideration

in establishing animal models. Growth phase is often associated with virulence programs in

bacteria (15–18), and our results suggest that log phase growth in *C. auris* is associated with

increased exposure of mannan and β-glucan, which are major fungal PAMPs. These cell wall

changes correlate with initiation of blood clotting during bloodstream infection in a murine model

of infection, leading to rapid mortality, potentially caused by septic shock, systemic thrombosis,

and disseminated intravascular coagulation. Indeed, we found decreased Fibrinogen levels in

plasma samples from mice infected by log phase *C. auris* at a high dose, as well as increased

Fibrin deposition in the kidneys and lungs. These results suggest a systemic thrombosis

response specifically to log phase *C. auris* at a high dose, which we propose causes rapid

mortality.

Understanding the cause of the observed blood clotting response will be an important

future direction of this work. While we observed increased exposure of mannan and β-glucan,

additional work will be needed to establish these cell wall changes as the fungal determinant of

coagulation in murine models. Mannan binding lectin-associated serine proteases can trigger

blood clotting (21), revealing one possible mechanism by which the high mannan log phase *C.*

auris may trigger rapid mortality at a high dose. Whether these cell wall alterations are

conserved in other strains of *C. auris*, which can differ in cell wall composition and virulence

(22), and whether additional strains also cause thrombosis in log phase growth are also

significant outstanding questions. Additionally, elucidating the host cell populations responsible

for triggering coagulation will be of interest, as resident or circulating innate immune cells, such

as monocytes, macrophages, and neutrophils can detect fungal PAMPs and produce

coagulation factors (23–25). However, vascular endothelial cells have also been shown to

express PRRs that can recognize fungal cell wall components (26, 27), and thus may play an

important role in thrombosis during *C. auris* bloodstream infection.

By comparing fungal burden in the organs of mice post-mortem, we found that *C. auris*

burden is high in the lungs early after infection but is rapidly cleared, whereas *C. auris*

establishes a more stable burden in the kidneys. The kidneys are also considered the primary

target organ of *C. albicans* (28, 29), although it is not immediately clear why the kidneys are a

more permissive environment for *C. auris* colonization than the lungs. However, future work to

reveal how the lungs clear *C. auris* may lead to new understanding of effective organ-specific

immune responses against *C. auris* and perhaps fungal pathogens more broadly.

Full understanding of the clinical implications of this report is hindered by a lack of

knowledge of what growth state is common for *C. auris* when initiating bloodstream infections in

humans. However, septic shock may occur in more than 20% of patients with *C. auris*

candidemia and increases mortality, and thrombosis has also been reported during *C. auris*

candidemia (1, 30, 31). Therefore, understanding the molecular mechanisms by which *C. auris*

infection causes thrombosis during bloodstream infection may inform therapeutic strategies to

treat these high-mortality infections. Together this work establishes virulence differences

between stationary and log phase *C. auris* at high doses and indicates that the development of

a standardized murine infection system, including controlling for fungal growth phase, will be

important for future studies examining *C. auris* pathogenesis.

Limitations of the Study

This report has several limitations that will be important to address in future studies.

While this work focuses on virulence differences between log and stationary phase growth *C.*

auris cells during bloodstream infection at a high dose, these differences were not readily seen

at a lower dose of infection. Thus, it is unclear how these results translate to human infection,

although thrombosis has been reported in patients with *C. auris* candidemia. Additionally, this

study focuses primarily on a single strain of *C. auris* and the widely used C57BL6/J mouse

model, thus additional work is needed to determine whether the observed phenotypes are

generalizable across *C. auris* strains and clades and with different animal models. While we

report increased abundance of mannan and exposure of β-glucan in the *C. auris* cell wall during

log phase growth, which correlates with blood clotting observed during systemic infection, we

have not definitively shown whether these cell wall changes are responsible for triggering

coagulation. While our results support that systemic thrombosis is the likely cause of rapid

mortality in high dose log phase *C. auris*-infected mice, we have not extensively profiled tissue-

specific inflammation, thus we cannot definitively rule out a proinflammatory response within

specific organs. Finally, the host cells that initiate the coagulation response to log phase *C. auris*

infection are not identified in this work.

Methods

Candidozyma auris **growth conditions.** *C. auris* strain CDC-AR0382 (B11109) was grown in

YPD liquid media (1% yeast extract, 2% peptone, 2% dextrose) with constant agitation. After 16

hours, stationary phase cultures were sub-cultured by diluting to OD600 of 0.2 in fresh YPD and

grown at 30°C for 4 hours with constant agitation to establish log phase growth.

Survival analysis post-systemic infection. *C. auris* from log phase or stationary phase

cultures were pelleted by centrifugation (5000 rpm for 5min), washed once with sterile PBS, and

resuspended in sterile PBS to desired doses for infection. Immunocompetent 8-week old female

C57BL/6J mice were infected intravenously with *C. auris* from log phase or stationary phase

growth at high dose (5×10^7) or low dose (1×10^6) via retro-orbital injection in 100 µL volume.

Immediately following infection, mice were monitored for onset of disease symptoms for several

hours initially, then daily over the course of 21 days. Mice were sacrificed at a humane endpoint

defined as loss of 20 percent of initial bodyweight, or when severe disease symptoms were

observed, such as unresponsiveness and labored breathing, or severe neurological symptoms.

Analysis of fungal burden in organs post-mortality. After sacrifice, organs samples were

harvested to measure fungal burden. The right lung, right kidney, and spleen were harvested by

dissection and homogenized by bead beating with sterile ⅛ inch ball bearings (Grainger 4RJL3)

for 10 seconds. Serial dilutions were performed and plated on YPD agar with ampicillin (100

mg/mL) and gentamicin (50 mg/mL). Fungal colonies were grown for 2 days at 30°C and

counted, and fungal burdens per organ were calculated.

Histological analysis and Martius Scarlet Blue staining for Fibrin analysis and

quantification and in organs. After sacrifice, organs samples were harvested to perform

histological analysis. The left lung and left kidney were harvested by dissection and fixed in 10

percent formalin for 24 hours, then plunged in 70 percent ethanol prior to sectioning and

embedding on slides by the University of Michigan Orthopaedic Research Laboratories

Histology Core. For histological analysis, hematoxylin and eosin (H&E) staining was performed

by the University of Michigan Orthopaedic Research Laboratories Histology Core. Full organ

images were captured at 10x magnification with color brightfield imaging using a BioTek

Lionheart FX Automated Microscope. For Martius Scarlet Blue (MSB) staining, slides were

deparaffinized through xylene and gradient ethanol (EtOH), and rehydrated to distilled water

(diH2O). Postfix was performed in Bouin Fixative (Newcomer Supply, Inc., Middletown, WI) at

room temperature overnight (~16 hours), followed by a 15 minute wash in diH2O. MSB staining

was then performed, as previously described (32). Images of full organs were captured at 20x

magnification with color brightfield imaging using a BioTek Lionheart FX Automated Microscope.

For analysis of relative Fibrin levels, 6 representative erythrocyte-rich regions were blindly

chosen from each organ and a pixel classifier was trained in QuPath v0.6.0 (33), using the wand

tool to segment Fibrin-positive pixels, erythrocyte-positive pixels, and kidney tissue to ignore, in

order to quantify Fibrin levels relative to erythrocytes within each image. The pixel classifier was

then used for each image to quantify the Fibrin-positive pixel area, relative to erythrocyte-

positive pixel area.

Histological analysis in organs post-mortality. After sacrifice, organs samples were

harvested to perform histological analysis. The left lung and left kidney were harvested by

dissection and fixed in 10 percent formalin for 24 hours, then plunged in 70 percent ethanol prior

to sectioning and staining (H&E and PAS) by the University of Michigan Orthopaedic Research

Laboratories Histology Core. Slides were imaged using a BioTek Lionheart FX automated

microscope.

ELISA. After sacrifice, serum was collected by cardiac puncture, followed by isolation of serum

using centrifugation (8000g for 5 minutes) of lithium heparin serum collection tubes (Kent

Scientific KMIC-LIHEP). Serum samples were submitted to the University of Michigan Cancer

Center Immunology Core for quantification of Fibrinogen, TNF, IL-6, and IL-1β by ELISA.

Analysis of cell wall content. Log and stationary phase *C. auris* cells were pelleted by

centrifugation (3000 x g for 5 minutes), washed once in PBS, and fixed in 4% paraformaldehyde

for 15 minutes. Following fixation, *C. auris* cells were pelleted by centrifugation (3000 x g for 5

minutes), washed twice in PBS, and then stained for cell wall contents, followed by flow

cytometry analysis or confocal microscopy. To quantify mannan content, cells were stained with

5 µg/mL of FITC-Concanavalin A (MilliporeSigma, C7642) for 30 minutes. To quantify exposed

β-1,3-glucan, cells were blocked with 3% bovine serum albumin & 5% normal goat serum

(Invitrogen, 10000C) for 30 minutes. After blocking, cells were stained with 15 µg/mL of hDectin-

1a (InvivoGen, fc-hec1a-2) for 1 hour. Cells were washed twice with PBS before secondary

staining with 4 mg/mL of goat raised anti-human IgG antibody conjugated with Alexa Fluor 647

(Invitrogen A-21445) for 30 minutes. To quantify chitin content, cells were stained with 0.1 g/L of

Calcofluor White (MilliporeSigma, 18909-100ML-F) for 10 minutes. Following staining, cells

were washed with 500μL PBS three times and resuspended in 500μL PBS. Samples were

analyzed on a LSRFortessa Flow Cytometer (BD Bioscience, NJ, USA) using BD FACSDiva

Software. 10,000 events were recorded for each sample. FlowJo software was used to

determine mean fluorescence intensity. For confocal microscopy, images were captured on a

Yokogawa CellVoyager CQ1 automated confocal microscope at 40x magnification, and

maximum intensity projections were generated from confocal Z-stacks of 10 μm. Representative

images were processed using NIH Fiji/ImageJ (34).

Acknowledgements

We thank Joel Whitfield of the University of Michigan Cancer Center Immunology Core and

Emma Snyder-White of the University of Michigan Orthopaedic Research Laboratories

Histology Core (NIH P30 AR069620) for guidance and technical support for ELISA and

histological analysis, and members of the O'Meara lab for discussion. This work was supported

by National Institutes of Health grants U19AI181767 and the Burroughs Wellcome Fund

Investigators in Pathogenesis award 1173374 to TRO.

REFERENCES

1. Ortiz-Roa C, Valderrama-Rios MC, Sierra-Umaña SF, Rodríguez JY, Muñetón-López GA, Solórzano-Ramos CA, Escandón P, Alvarez-Moreno CA, Cortés JA. 2023. Mortality caused by Candida auris bloodstream infections in comparison with other Candida species, a multicentre retrospective cohort. J Fungi (Basel) 9.

2. Kim HY PhD, Nguyen TA MSc, Kidd S PhD, Chambers J MD, Alastruey-Izquierdo A PhD, Shin J-H MD, Dao A PhD, Forastiero A MD, Wahyuningsih R MD, Chakrabarti A MD, Beyer P, Gigante V PhD, Beardsley J PhD, Sati H PhD, Morrissey CO PhD, Alffenaar J-W PhD. 2024. Candida auris-a systematic review to inform the world health organization fungal priority pathogens list. Med Mycol 62.

3. Wang Y, Zou Y, Chen X, Li H, Yin Z, Zhang B, Xu Y, Zhang Y, Zhang R, Huang X, Yang W, Xu C, Jiang T, Tang Q, Zhou Z, Ji Y, Liu Y, Hu L, Zhou J, Zhou Y, Zhao J, Liu N, Huang G, Chang H, Fang W, Chen C, Zhou D. 2022. Innate immune responses against the fungal pathogen Candida auris. Nat Commun 13:3553.

4. Xin H, Mohiuddin F, Tran J, Adams A, Eberle K. 2019. Experimental mouse models of disseminated Candida auris infection. mSphere 4.

5. Eix EF, Nett JE. 2025. Candida auris: Epidemiology and antifungal strategy. Annu Rev Med 76:57–67.

6. Santana DJ, Anku JAE, Zhao G, Zarnowski R, Johnson CJ, Hautau H, Visser ND, Ibrahim AS, Andes D, Nett JE, Singh S, O'Meara TR. 2023. A Candida auris-specific adhesin, Scf1, governs surface association, colonization, and virulence. Science 381:1461–1467.

7. Iyer KR, Camara K, Daniel-Ivad M, Trilles R, Pimentel-Elardo SM, Fossen JL, Marchillo K, Liu Z, Singh S, Muñoz JF, Kim SH, Porco JA Jr, Cuomo CA, Williams NS, Ibrahim AS,

Edwards JE Jr, Andes DR, Nodwell JR, Brown LE, Whitesell L, Robbins N, Cowen LE.

2020. An oxindole efflux inhibitor potentiates azoles and impairs virulence in the fungal

pathogen Candida auris. Nat Commun 11:6429.

8. Singh S, Uppuluri P, Mamouei Z, Alqarihi A, Elhassan H, French S, Lockhart SR, Chiller T,

Edwards JE Jr, Ibrahim AS. 2019. The NDV-3A vaccine protects mice from multidrug

resistant Candida auris infection. PLoS Pathog 15:e1007460.

9. Zhao G, Lyu J, Veniaminova NA, Zarnowski R, Mattos E, Johnson CJ, Quintanilla D,

Hautau H, Hold LA, Xu B, Anku JAE, Dasgupta K, Hale JJ, Steltzer SS, Santana DJ,

Ibrahim AS, Snitkin ES, Andes D, Nett JE, Singh S, Abraham AC, Killian ML, Kahlenberg

JM, Wong SY, O'Meara TR. 2025. Adhesin Als4112 promotes Candida auris skin

colonization through interactions with keratinocytes and extracellular matrix proteins. Nat

Commun 16:5673.

10. Herrada J, Roberts K, Gamal A, Long L, Ghannoum MA. 2022. An immunocompromised

mouse model of Candida auris systemic infection. Methods Mol Biol 2517:317–328.

11. Torres SR, Pichowicz A, Torres-Velez F, Song R, Singh N, Lasek-Nesselquist E, De Jesus

M. 2020. Impact of Candida auris Infection in a Neutropenic Murine Model. Antimicrob

Agents Chemother 64.

12. Fakhim H, Vaezi A, Dannaoui E, Chowdhary A, Nasiry D, Faeli L, Meis JF, Badali H. 2018.

Comparative virulence of Candida auris with Candida haemulonii, Candida glabrata and

Candida albicans in a murine model. Mycoses 61:377–382.

13. Wang X, Bing J, Zheng Q, Zhang F, Liu J, Yue H, Tao L, Du H, Wang Y, Wang H, Huang G.

2018. The first isolate of Candida auris in China: clinical and biological aspects. Emerg

Microbes Infect 7:93.

14. Santana DJ, Zhao G, O'Meara TR. 2024. The many faces of Candida auris: Phenotypic and
strain variation in an emerging pathogen. PLoS Pathog 20:e1012011.

15. Hammer BK, Swanson MS. 1999. Co-ordination of legionella pneumophila virulence with
entry into stationary phase by ppGpp. Mol Microbiol 33:721–731.

16. Roop RM 2nd, Gee JM, Robertson GT, Richardson JM, Ng W-L, Winkler ME. 2003.
Brucella stationary-phase gene expression and virulence. Annu Rev Microbiol 57:57–76.

17. Wilson JA, Gulig PA. 1998. Regulation of the spvR gene of the Salmonella typhimurium
virulence plasmid during exponential-phase growth in intracellular salts medium and at
stationary phase in L broth. Microbiology 144 (Pt 7):1823–1833.

18. Ribardo DA, McIver KS. 2003. amrA encodes a putative membrane protein necessary for
maximal exponential phase expression of the Mga virulence regulon in Streptococcus
pyogenes: Exponential phase expression of Mga. Mol Microbiol 50:673–685.

19. Angus DC, van der Poll T. 2013. Severe sepsis and septic shock. N Engl J Med 369:840–
851.

20. Litvinov RI, Pieters M, de Lange-Loots Z, Weisel JW. 2021. Fibrinogen and fibrin. Subcell
Biochem 96:471–501.

21. Gulla KC, Gupta K, Krarup A, Gal P, Schwaeble WJ, Sim RB, O'Connor CD, Hajela K.
2010. Activation of mannan-binding lectin-associated serine proteases leads to generation
of a fibrin clot: MASPs generate a fibrin clot. Immunology 129:482–495.

22. Bruno M, Kersten S, Bain JM, Jaeger M, Rosati D, Kruppa MD, Lowman DW, Rice PJ,
Graves B, Ma Z, Jiao YN, Chowdhary A, Renieris G, van de Veerdonk FL, Kullberg B-J,
Giamarellos-Bourboulis EJ, Hoischen A, Gow NAR, Brown AJP, Meis JF, Williams DL,

Netea MG. 2020. Transcriptional and functional insights into the host immune response

against the emerging fungal pathogen Candida auris. Nat Microbiol 5:1516–1531.

23. Osterud B, Lindahl U, Seljelid R. 1980. Macrophages produce blood coagulation factors.

FEBS Lett 120:41–43.

24. Wilhelm G, Mertowska P, Mertowski S, Przysucha A, Strużyna J, Grywalska E, Torres K.

2023. The Crossroads of the coagulation system and the immune system: Interactions and

connections. Int J Mol Sci 24.

25. Netea MG, Joosten LAB, van der Meer JWM, Kullberg B-J, van de Veerdonk FL. 2015.

Immune defence against Candida fungal infections. Nat Rev Immunol 15:630–642.

26. Lowe EP, Wei D, Rice PJ, Li C, Kalbfleisch J, Browder IW, Williams DL. 2002. Human

vascular endothelial cells express pattern recognition receptors for fungal glucans which

stimulates nuclear factor kappaB activation and interleukin 8 production. Winner of the Best

Paper Award from the Gold Medal Forum. Am Surg 68:508–17; discussion 517-8.

27. Barker KS, Park H, Phan QT, Xu L, Homayouni R, Rogers PD, Filler SG. 2008.

Transcriptome profile of the vascular endothelial cell response to Candida albicans. J Infect

Dis 198:193–202.

28. Lehner T. 1964. Systemic candidiasis and renal involvement. Lancet 2:1414–1416.

29. Lionakis MS, Lim JK, Lee C-CR, Murphy PM. 2011. Organ-specific innate immune

responses in a mouse model of invasive candidiasis. J Innate Immun 3:180–199.

30. Sayeed MA, Farooqi J, Jabeen K, Awan S, Mahmood SF. 2019. Clinical spectrum and

factors impacting outcome of Candida auris: a single center study from Pakistan. BMC

Infect Dis 19:384.

31. Jacobs SE, Jacobs JL, Dennis EK, Taimur S, Rana M, Patel D, Gitman M, Patel G,

Schaefer S, Iyer K, Moon J, Adams V, Lerner P, Walsh TJ, Zhu Y, Anower MR, Vaidya MM,

Chaturvedi S, Chaturvedi V. 2022. Candida auris pan-drug-resistant to four classes of

antifungal agents. Antimicrob Agents Chemother 66:e0005322.

32. Obi AT, Sharma SB, Elfline MA, Luke CE, Dowling AR, Cai Q, Kimball AS, Hollinstat M,

Stanger L, Moore BB, Jaffer FA, Henke PK. 2023. Experimental venous thrombus

resolution is driven by IL-6 mediated monocyte actions. Sci Rep 13:3253.

33. Bankhead P, Loughrey MB, Fernández JA, Dombrowski Y, McArt DG, Dunne PD, McQuaid

S, Gray RT, Murray LJ, Coleman HG, James JA, Salto-Tellez M, Hamilton PW. 2017.

QuPath: Open source software for digital pathology image analysis. Sci Rep 7:16878.

34. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,

Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,

Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis.

Nat Methods 9:676–682.

Proinflammatory cytokine levels (2 hpi)

A TNF

B IL-6

C IL-1β

Proinflammatory cytokine levels (time of mortality)

D TNF

E IL-6

F IL-1β

G

Log

Stationary

Lung H&E Staining