Running Head

PAF and HFS

Title

Interindividual variations in peak alpha frequency do not predict the magnitude or extent of
secondary hyperalgesia induced by high-frequency stimulation

Authors

Louisien Lebrun[1], Gloria Ricci[2], Arthur Courtin[1,3], Emanuel N. van den Broeke[1,4], Cédric Lenoir[1], André
Mouraux[1]

Affiliations

[1] Institute of Neuroscience (IONS), UCLouvain, Brussels, Belgium

[2] Department of Neuroscience (DNS), University of Padova

[3] Centre for Functionally Integrative Neuroscience (CFIN), Aarhus University, Aarhus, Denmark

[4] Research Group Health Psychology, Faculty of Psychology and Educational Sciences, KU Leuven,
Belgium.

CORRESPONDING AUTHOR: louisien.lebrun@student.uclouvain.be

ADDITIONAL CO-CORESPONDING AUTHOR: andre.mouraux@uclouvain.be

Abstract (242 words)

Background

Previous studies have shown an association between interindividual variations in the frequency of

alpha-band EEG oscillations such as estimates of peak alpha frequency (PAF) and pain sensitivity.

Whether differences in PAF also influence the susceptibility to develop central sensitization (CS) is

unknown.

Objective

This study aimed to determine if the PAF of vision- and sensorimotor-related alpha-band activity is

associated with the magnitude and extent of secondary mechanical hyperalgesia induced by high-

frequency stimulation (HFS) of the skin, a surrogate marker of CS.

Methods

The EEG was recorded in 32 healthy participants at rest during eyes open and eyes closed conditions,

and during bilateral finger movements. Two methods were used to isolate vision- and sensorimotor-

related alpha-band activity based on sensitivity to eye closure and movement: one based on an

independent component analysis, the other on spectral subtraction. PAF was assessed using a

center-of-gravity approach or the Fitting Oscillations and One-Over-F (FOOOF) method, accounting

for the aperiodic EEG component. HFS was applied to the right forearm, and pinprick sensitivity was

assessed at both forearms, before and 40 minutes after HFS.

Results

Neither sensorimotor- nor vision-related PAF were significantly correlated with the magnitude or

extent of HFS-induced secondary hyperalgesia. Interestingly, at the non-sensitized forearm,

participants with a higher vision-related PAF exhibited greater habituation to pinprick stimuli,

suggesting that variations in PAF may relate to variations in sensory habituation.

Conclusion

Interindividual variations of PAF were not significantly associated with the susceptibility to develop

HFS-induced secondary hyperalgesia.

Keywords

High Frequency Stimulation (HFS), Peak Alpha Frequency, EEG Spectrum, alpha-band, Secondary

Hyperalgesia, Central Sensitization

New and Noteworthy (44 words):

Using several methods to estimate vision- and sensorimotor-related peak alpha frequency (PAF) in

the EEG frequency spectrum, we found no significant association between interindividual variations

in PAF and the susceptibility to develop secondary hyperalgesia following high-frequency stimulation

(HFS) of the skin in healthy participants.

INTRODUCTION

Alpha-band oscillatory activity (8-12 Hz) is the most prominent feature of the EEG recorded during rest in awake humans. The alpha rhythm was first described by Hans Berger, nearly a century ago (1). A distinctive feature of that rhythm is its predominance over posterior areas of the brain, and the fact that its magnitude markedly increases when participants close their eyes. The sensitivity to eye closure of posterior alpha oscillations has led most authors to consider that this oscillatory activity reflects a default state of brain areas "resting" or "idling" in the absence of input (2). However, alpha-band oscillations are not restricted to the visual cortex. Gastaut et al. (1952) described the Rolandic mu rhythm oscillating between 8-12 Hz over the scalp vertex and bilateral sensorimotor areas (3). Present at rest, the mu rhythm decreases during movement execution (3–5), most pronouncedly in the hemisphere contralateral to the executed movement (5). It is also suppressed during motor imagery (6,7), modulated by mental effort (8), attention and fatigue (9), and relatively unaffected by opening/closing of the eyes (10). Studies comparing the effects of upper and lower limb movements found that the movement-related suppression of the mu rhythm follows the mototopic and somatotopic organization of primary motor (M1) and somatosensory (S1) cortices (5).

Several studies have reported that the frequency at which the EEG spectrogram peaks within the alpha-band, referred to as the "peak alpha frequency" (PAF), tends to be slower in patients with chronic pain (11–16). Other studies have found that a slower PAF, recorded in healthy participants before they experienced experimental pain is associated with a greater sensitivity to phasic heat pain (17), tonic pain produced by topical capsaicin (18) and deep tissue pain produced by nerve growth factor injection (19,20), whereas an opposite trend has been observed in a previous study with tonic heat pain (17). The PAF has been shown to vary across individuals, but to remain relatively stable over time (17,20,22), suggesting that PAF may reflect an individual trait somehow related to pain sensitivity.

An important property of the nociceptive system is its propensity to sensitize when exposed to

repeated nociceptive input or following tissue injury (23–27). Sensitization of the nociceptive system

involves changes at the level of peripheral nociceptors leading to an increased sensitivity to noxious

stimuli (peripheral sensitization) (28) and changes at the level of the central nervous system leading

to an increased synaptic transmission of nociceptive input (central sensitization) described

extensively at the level of the dorsal horn in animal studies (29,30), but most likely also involving

higher-order synapses (31). A perceptual consequence of sensitization is hyperalgesia, i.e., an

increased sensitivity to pain (32). Following skin injury, hyperalgesia develops both within the injured

area (primary hyperalgesia) and in the surrounding uninjured skin (secondary hyperalgesia) (33,34).

While both peripheral and central sensitization may contribute to the development of primary

hyperalgesia, secondary hyperalgesia is thought to result exclusively from central sensitization

(34,35). A hallmark feature of secondary hyperalgesia is a prominent increase in sensitivity to

mechano-nociceptive input as compared to thermo-nociceptive input, enabling its characterization

by testing sensitivity to mechanical pinprick stimulation of the skin (36,37).

In a recent study conducted in patients scheduled for lateral thoracotomy, we found that

interindividual variations in the pre-operative susceptibility to develop secondary hyperalgesia

following high-frequency electrical stimulation of the skin (HFS) – a procedure used to induce central

sensitization experimentally – was associated with a greater likelihood post-thoracotomy pain at 2

months (38). The primary aim of the present study was to test whether interindividual variations in

PAF may be related to variations in this susceptibility to develop HFS-induced secondary

hyperalgesia.

Supporting this hypothesis, several studies have suggested that non-invasive brain stimulation of the

sensorimotor cortex can reduce pain sensitivity (39,40), and that this effect may be mediated

through a modulation of M1-connected brain structures involved in descending pain modulation

(41). Therefore, if variations of sensorimotor-related alpha band activities index variations in M1

function, interindividual differences in sensorimotor PAF could be associated with changes in the

state of descending pain modulation and the susceptibility to develop central sensitization. Finally,

since most previous reports of a relationship between PAF and sensitivity to pain have used models

of sustained pain which are known to induce some amount of sensitization, these observations might

be explained in part by a relationship between PAF and susceptibility to sensitize.

To test this hypothesis, the EEG was recorded in healthy participants with their eyes open at rest,

their eyes closed at rest, and with their eyes open while performing repetitive bilateral finger

movements. These three EEG recording conditions were used to functionally isolate, vision-related

alpha-band oscillations whose magnitude is known to increase following closure of the eyes and

sensorimotor-related alpha-band oscillations whose magnitude reduces during finger movements. As

in a multiverse analysis (42), several methods were used to (1) isolate vision- and sensorimotor-

related alpha-band activity and (2) estimate PAF. After the EEG recording, central sensitization was

induced in each participant using HFS delivered to the right volar forearm skin. The magnitude and

extent of HFS-induced secondary hyperalgesia was estimated by assessing the sensitivity to

mechanical pinprick stimuli delivered to the sensitized and non-sensitized forearms before and 40

minutes after HFS. The collected data allowed us to examine whether interindividual variations in

vision- and/or sensorimotor-related PAF predicts the susceptibility to develop HFS-induced

secondary mechanical hyperalgesia.

METHODS

Participants

Descriptive statistics are presented as mean ± standard deviation (SD), unless otherwise specified.

A total of 32 right-handed healthy volunteers (16 women and 16 men aged 25.3 ± 4.5 years [range:

19-38]) were included in this study between June and July 2020. Hand dominance was confirmed

using the Flinders Handedness Survey (43). The study was approved by the local ethics committee

(Comité d'Ethique Hospitalo-Facultaire des Cliniques universitaires Saint-Luc-UCLouvain;

B403201316436). All participants provided written informed consent and received financial

compensation. The experiment was conducted in agreement with the Declaration of Helsinki. The

decision to include 32 participants was guided by the correlation coefficients reported in previous

studies investigating the relationship between PAF and pain ratings in healthy volunteers (r = 0.74 in

(21) and r = 0.5 in (18). Based on these two studies, the estimated required sample sizes range from

12 to 38 participants to achieve a power of 0.90 with an alpha level of 0.05 (44).

Inclusion criteria were being healthy and aged 18-40 years. Exclusion criteria were (1) any history of

cardiac or neurological pathology, (2) regular practice of activities involving repeated stimulation of

the volar forearm skin (e.g., volleyball and handball players), (3) a dermatological condition involving

the forearm, (4) a history of traumatic injury of the upper limb and (5) regular tobacco consumption,

(6) having taken part in a previous experiment involving stimulation of the volar forearm(s).

Participants were asked to sleep a minimum of 8 hours the night before the experiment, to refrain

from using recreational drugs, psychotropics, analgesics, or other medications in the days leading up

to the study.

Experimental design

The time course of the experiment is summarized in Figure 1A. Participants were seated on a

comfortable chair with their arms resting on a desk, volar forearms facing upwards. Secondary

hyperalgesia was induced at the right volar forearm using HFS regardless of hand dominance. EEG

recordings were performed 25 minutes before HFS (T-25), 5 minutes before HFS (T-5), one minute

after HFS (T+1) and approximatively 35 minutes after HFS (T+35). Each EEG recording lasted 5

minutes and included three conditions: eyes open at rest, eyes closed at rest and eyes open while

performing repetitive bilateral flexion-extension movements with their index fingers. During the 30

minutes pause separating T+1 and T+35 EEG recordings, participants were instructed to read a book

or magazine of their choice. Sensitivity to mechanical pinprick stimuli was evaluated at both volar

forearms 20 minutes before (T-20) and approximatively 40 minutes after (T+40) HFS. The extent of

the HFS-induced area of secondary hyperalgesia was evaluated 40 minutes after HFS (T+40), at the

HFS-treated arm.

Figure 1. A. Timeline of the experiment. HFS was applied to the right volar forearm. EEG segments

lasting 5 minutes and including three conditions (eyes open at rest, eyes closed at rest, eyes open

while performing bilateral finger movements) were recorded 25 and 5 minutes before HFS (T-25 and

T-5), immediately after HFS, and approximately 35 minutes after HFS (T+35). Pinprick sensitivity of

the left and right volar forearms was assessed before HFS (T-20) and approximately 40 minutes after

HFS (T+40). The spatial extent of the HFS-induced area of secondary hyperalgesia was assessed

approximately 40 minutes after HFS (T+40). **B.** HFS was delivered to the forearm skin using a cathode

consisting of 10 blunt tungsten pins arranged on a circle with a diameter of 5 mm and a 24x20 mm

anode conductive gel pad placed next to the cathode.

Induction of secondary hyperalgesia using HFS

HFS was applied to the right volar forearm, 10 cm distal from the cubital fossa using an electrode

designed to preferentially activate cutaneous free nerve endings (EPS-P10, MRC Systems GmbH,

Heidelberg, Germany). The electrode consisted of a multi-pin cathode and a flat large-surface anode

(Figure 1B). The cathode consisted of 10 blunt tungsten pins arranged on a circle with a diameter of 5

mm. Each pin had a diameter of 0.25 mm and protruded by 0.65 mm over the base of the electrode.

The anode was a 24 x 20 mm conductive gel pad placed next to the cathode. The stimulation

consisted of 5 trains of 100 Hz charge-compensated biphasic electrical pulses (45) delivered using a

constant-current electrical stimulator (DS5; Digitimer, Welwyn Garden City, United Kingdom)

triggered by a National Instruments digital-analogue interface (NI6343, National Instruments, Austin,

Texas, USA). Each train lasted 1 second (100 pulses/train). The time interval between the onset of

each train was 10 s. The intensity of stimulation was set to 20 the detection threshold to a single

pulse, which was estimated in each participant at the beginning of the experiment using the method

of limits (0.126 ± 0.045 mA).

Assessment of mechanical pinprick sensitivity at the volar forearms

Before HFS (T-20) and after HFS (T+40), the sensitivity to mechanical pinprick stimuli was assessed at

the left and right volar forearms using a custom-made pinprick stimulator (0.25 mm probe diameter,

CATL, UCLouvain, Belgium) exerting a 128 mN force (26,27,45). At each time-point, three pinprick

stimuli were applied perpendicularly to the skin surrounding the area of HFS stimulation at the right

volar forearm and on the same location at the left volar forearm. After each stimulus, participants

were requested to report the intensity of the elicited sensation using a numerical rating scale ranging

from 0 (no perception) to 100 (maximal imaginable pain), with 50 marking the limit between non-

painful and painful sensations as used in several previous experiments (46–49). To avoid sensitization

of the skin by repeated mechanical pinprick stimulation, the probe was not applied twice on the

same location. The order of the testing (HFS-treated arm first or control arm tested first) was

counterbalanced across participants and remained constant across the two time points. The mean of

the three ratings obtained at each time point and arm was used as an index of pinprick .

Assessment of the area of HFS-induced secondary hyperalgesia

The same 128 mN pinprick probe was used to map, after HFS (T+40), the area of increased

mechanical pinprick sensitivity surrounding the site of HFS stimulation at the HFS-treated volar

forearm. The stimuli were applied along 4 axes and 8 directions separated by a 45-degrees angle and

each originating from skin location corresponding to the center of the HFS electrode and.

Participants were asked to close their eyes during the mapping procedure. Starting away from the

HFS-treated skin and moving towards the center, stimuli were applied every 5 mm. Participants were

asked to report when the sensation increased, and the corresponding stimulation location was

marked with a felt-tip pen (26,47,49–51). For each of the 8 directions, the distance between the

center of the HFS electrode location and the mark was measured, and the area of secondary

hyperalgesia was computed as the area of a piecewise polynomial fit of a 2D closed curve linking the

8 marked locations ('interpclosed' function; Santiago Benito 2023, Matlab Central File Exchange)

(26,27). Since this function cannot process null values, we assigned a value of 0.1 when no secondary

hyperalgesia area was present in this direction. The radius of the area of increased pinprick

sensitivity was then computed as the square root of the estimated area divided by π, as if the area

was in the shape of a circle. This transformation was performed prior to any analyses.

EEG recording

The EEG was recorded using 64 scalp electrodes with active shielding, placed according to the

International 10-20 system (ANT Neuro, Enschede, The Netherlands). The signals were digitized at

1,000 Hz using an average reference. All impedances were kept under 20 kΩ. At each time point (T-

25, T-5, T+1 and T+36), the recording consisted of 3 minutes at rest with eyes open (participants

were asked to relax and stare at a cross displayed 1 meter in front of them, at a 30° angle below eye

level), followed by 1 minute at rest with eyes closed, and 1 minute with eyes open while performing

repeated bilateral index finger movements at a rate of approximately 1 Hz while fixing their gaze on

the cross. These three conditions (EYES-OPEN-REST, EYES-CLOSED-REST, EYES-OPEN-MVT) were used

to differentially modulate vision-related alpha-band activity (which should be enhanced in the EYES-

CLOSED-REST condition as compared to the other conditions) and sensorimotor-related alpha-band

activity (which should be attenuated when performing finger movements in the EYES-OPEN-MVT

condition as compared to the other conditions).

EEG preprocessing

The EEG signals collected before HFS (T-25) were analyzed offline using the Letswave 6 toolbox

(http://letswave.org) running on MATLAB 2018a (Mathworks, USA). The EEG data collected at the

other time points was not analyzed in the present study.

First, the EEG recordings were cut in three segments: a 180-s EYES-OPEN-REST segment, a 60 s

segment EYES-CLOSED-REST and a 60 s segment EYES-OPEN-MVT (6). The three segments were then

filtered using a 0.5 - 47 Hz zero-phase Butterworth band-pass filter (order 4) and further segmented

into half-overlapping 5-s epochs. Each segment was demeaned, and a linear detrend was applied to

remove slow drifts of the signal. Noisy electrodes (1-2 electrodes in 10 out of 32 recordings) were

interpolated using the average of the three nearest electrodes. Epochs with clear artifacts (absolute

amplitude ≥200 µV) were rejected (2.4% of epochs from EYES-OPEN-REST condition, 2.8% of epochs

from EYES-CLOSED-REST condition and 2.8% of epochs from EYES-OPEN-MVT condition). Signals were

re-referenced to the average of all channels excluding the two mastoid electrodes (M1, M2).

Functional separation of vision- and sensorimotor-related alpha-band activity

In previous studies, PAF has been measured using various methods (17,18,21,52–54) and at different

scalp locations (17,55–59), with some studies interpreting it as a marker of sensorimotor activity

even when recorded over central or occipital regions or when clearly responding to closure of the

eyes (11,17–19). In this work, we aimed to explore multiple approaches to separate vision- and

sensorimotor-related alpha-band activity and estimate their PAF.

ICA-based method

In a first approach, an independent component analysis (ICA) was used to dissociate visual alpha-

band activity (VISUAL-ALPHA) and sensorimotor alpha-band activity (SM-ALPHA) based on their

sensitivity to closure of the eyes and movement of the hands.

For each participant, the 'runica' algorithm was used to separate the 62 EEG scalp signals into 61

independent components (ICs) having a fixed topographical distribution, and maximally independent

time courses. In the case channels were interpolated during preprocessing, the number of ICs was

lowered by the number of interpolated channels (60). Because the amplitude of VISUAL-ALPHA

should be selectively increased during the EYES-CLOSED-REST condition while SM-ALPHA should be

selectively decreased during bilateral moving of the fingers in the EYES-OPEN-MVT condition, ICA can

be expected to capture visual- and sensorimotor-related alpha-band activity into separate ICs.

For each participant and each IC time course, a Fast-Fourier Transform (FFT) was used to compute

the amplitude frequency spectra of all 5-s half-overlapping epochs, windowed using a Hanning

function. The spectra were then averaged per participant. The obtained frequency spectra were used

to identify ICs capturing alpha-band activity, and to categorize these ICs into the following three

categories: VISUAL-ALPHA (ICs capturing vision-related alpha-band activity that was increased in the

EYES-CLOSED-REST condition as compared to EYES-OPEN-REST condition and the EYES-OPEN-MVT

condition), SM-ALPHA (ICs capturing sensorimotor-related alpha-band activity that was decreased in

the EYES-OPEN-MVT condition as compared to the EYES-OPEN-REST condition and was not increased

in the EYES-CLOSED-REST condition as compared to the EYES-OPEN-REST condition) and MIXED-

ALPHA (ICs capturing alpha-band activity that was both sensitive to closure of the eyes – greater

activity in the EYES-CLOSED-REST condition compared to the EYES-OPEN-REST condition – and

sensitive to finger movements – reduced activity in the EYES-OPEN-MVT condition compared to the

EYES-OPEN-REST condition).

The classification was achieved for each IC of each participant as follows. First, using the average of

the frequency spectra obtained in all three conditions, the peak of potential alpha-band activity

captured by the IC was computed using the center of gravity (CoG) formula (19,61)

$$CoG = \frac{\sum_{i=1}^{n} f_i * a_i}{\sum_{i=1}^{n} a_i}$$

where f_i is the i^{th} frequency bin between 9 and 11 Hz, a_i is the spectral amplitude for f_i, and n is the

number of frequency bins between 9 and 11 Hz. This value was then used to define a 6 Hz frequency

window of interest centered on the estimated CoG. Second, the mean amplitude and standard error

of the mean (SEM: standard deviation of amplitudes divided by the square root of the number of

bins) were computed separately for each of the three EEG recording conditions using all amplitude

bins located within the selected window of interest. The mean and SEM were used to compute a

confidence interval corresponding to the mean ± 1.97*SEM, and the lower and upper bounds of

these confidence intervals (CI) were used to perform the categorization. ICs were classified as

capturing vision-related alpha-band activity (VISUAL-ALPHA) when the lower bound of the CI in the

EYES-CLOSED-REST condition exceeded the upper bound of the CI in the EYES-OPEN-REST condition

(indicating an increase of alpha during eye closure), whereas the CI in the EYES-OPEN-REST condition

overlapped the CI in the EYES-OPEN-MVT condition (indicating no strong effect of movement on

alpha-band amplitude). Similarly, ICs were classified as capturing movement-related alpha (SM-

ALPHA) when the lower bound of the CI in the EYES-OPEN-REST condition exceeded the upper bound

of the CI in the EYES-OPEN-MVT condition (indicating a movement-induced reduction of alpha),

whereas the CI in the EYES-OPEN-REST and EYES-CLOSED-REST conditions overlapped (indicating low

sensitivity to closure of the eyes). Finally, ICs were classified as MIXED-ALPHA when they were both

sensitive to closure of the eyes (lower bound of the CI in the EYES-CLOSED-REST condition > upper

bound of the CI in the EYES-OPEN-REST condition) and sensitive to the finger movements (lower

bound of the CI in the EYES-OPEN-REST condition > upper bound of the CI in the EYES-OPEN-MVT

condition). Examples of ICs capturing vision-related and sensorimotor-related alpha-band activity are

shown in Figure 2.

For each participant, the VISUAL-ALPHA, SM-ALPHA and MIXED-ALPHA ICs were then used to

reconstruct three separate EEG datasets maximally capturing vision-related, sensorimotor-related

and mixed alpha-band activity, respectively.

Figure 2. **A.** EEG frequency spectrum of a representative independent component (IC) categorized as

capturing vision-related alpha-band activity (VIS-ALPHA) that was increased during closure of the

eyes (EYES-CLOSED-REST condition) as compared to the eyes open conditions (EYES-OPEN-REST and

EYES-OPEN-MOVEMENT). **B.** EEG frequency spectrum of a representative IC categorized as capturing

sensorimotor-related alpha-band activity (SM-ALPHA) that was decreased during finger movements

(EYES-OPEN-MVT condition) as compared to rest (EYES-OPEN-REST and EYES-CLOSED-REST

conditions). Y-axis: arbitrary units. X-axis: frequency in Hz. The vertical grey lines represent the 6 Hz

frequency window used to assess the presence of alpha-band activity.

Subtraction-based method

Suppression of movement-related activity in the EYES-OPEN-MVT condition may be expected to

constitute the main difference between that condition and the EYES-OPEN-REST condition.

Therefore, computing the difference between the EYES-OPEN-REST and EYES-OPEN-MVT spectra

could constitute a simple alternative to efficiently isolate movement-related alpha-band activity.

Similarly, because closure of the eyes selectively enhances vision-related alpha oscillations,

subtraction of the EYES-OPEN-REST and EYES-CLOSED-REST spectra could constitute a means to

isolate vision-related alpha-band activity. As compared to the ICA-based approach, a possible

advantage of this alternative approach is that the subtraction can be expected to cancel out the

contribution of activity that is equally present in both of the subtracted spectra, including the 1/f

aperiodic component of the EEG power spectrum that can bias estimation of PAF (18).

Estimation of peak alpha frequency (PAF)

Vision-related alpha oscillations can be expected to predominate over occipital electrodes, whereas

sensorimotor-related alpha oscillations modulated by bilateral movements of the fingers can be

expected to spread over left and right central electrodes (5,10). Therefore, the PAF of vision-related

alpha-band activity was estimated in the average of the frequency spectra of occipital electrodes O1,

Oz and O2, using the eyes-closed derived VISUAL-ALPHA spectra of the ICA-based method and the

EYES-CLOSED-REST minus EYES-OPEN-REST spectra of the subtraction-based method. Similarly, the

PAF of movement-related alpha-band activity was estimated in the average of the frequency spectra

of central electrodes C3, Cz and C4, using the eyes open-derived SM-ALPHA spectra of the ICA-based

method and the EYES-OPEN-REST minus EYES-OPEN-MVT spectra of the subtraction-based method.

Finally, because the MIXED-ALPHA signals generated by the ICA-based method might have included

both vision- and sensorimotor-related alpha-band activity, its PAF was estimated both at occipital

and at central channels.

Two different methods were used to estimate the PAF: the center of gravity (CoG) method and the

Fitting Oscillations and One Over F method (FOOOF (62)).

For the CoG method, we used the formula described above and a 9-11 Hz interval as was used in

previous studies assessing the relationship between PAF and sensitivity to pain (18,19). As visual

inspection of the EEG frequency spectra indicated that the PAF of some individuals was located

outside the 9-11 Hz interval, the CoG was also estimated using a larger 7-13 Hz window. In the

subtracted spectra, some amplitude values within the 9-11 or 7-13 Hz windows were negative. Only

positive values were used for the CoG estimation.

For the FOOOF method, the spectrum was decomposed in its periodic and aperiodic components, as

described in Donoghue et al. (62). In this case, the alpha band was defined as the range from 7 to 14

Hz and default algorithm parameters were employed, in coherence with other studies that

performed similar analyses (63). When using this approach a larger frequency range is advised, as the

overlapping of the frequency borders with a spectral peak is associated with higher parameter

estimation errors (64). After the decomposition, only the periodic component was considered, as it is

expected to contain alpha-band oscillatory activities. A gaussian curve was used to fit the periodic

component of the spectrum, and its parameters were used to characterize the PAF (defined as the

center of the gaussian curve). As the FOOOF method is intended to suppress the 1/f aperiodic

component of the EEG, it was not applied to the subtracted spectra.

Statistical analysis

Statistical analyses were performed using JASP (JASP Version 0.14.1.0 for Windows). Statistical tests

were chosen depending on the distribution of the dependent variables, which was assessed by visual

examination of the histograms for skewness and kurtosis, and the Q-Q plots of the residuals.

HFS-induced change in pinprick sensitivity. Mechanical pinprick ratings followed a close-to-normal

distribution and we thus conducted a repeated-measures ANOVA with two within-subjects factors:

TIME (before vs. after HFS) and ARM (HFS-treated vs. control arm), and one between-subjects factor:

ORDER (whether the HFS-treated or control arm was tested first). Data were averaged per condition

for each participant prior to analysis.

Peak-alpha frequency. A correlation analysis was used to evaluate the relationship between

sensorimotor- and vision-related PAF and the changes in pinprick sensitivity after HFS (change in

pinprick sensitivity at the sensitized HFS-treated forearm: $\Delta NRS_{HFS}=NRS_{HFS}(T+40)$ - $NRS_{HFS}(T-20)$,

change in pinprick sensitivity at the non-sensitized forearm: $\Delta NRS_{NS}=NRS_{NS}(T+40)$ - $NRS_{NS}(T-20)$,

difference in the change in pinprick sensitivity at the two forearms: $\Delta\Delta NRS = \Delta NRS_{HFS}$ - ΔNRS_{NS}), and

between PAF and the radius of the area of secondary mechanical hyperalgesia (R). For each

estimated PAF, the analyses were performed using Spearman's correlations because normality was

not strictly respected for all variables. Yet, as it was respected for our primary aim (relationship

between sensorimotor PAF and secondary hyperalgesia), we displayed Pearson's correlation for

these analyses.

To examine a possible relationship between PAF and pain sensitivity, we also computed Spearman's

correlation between PAF and the detection threshold to single electrical stimuli, as well as the

average ratings of the intensity of the sensation elicited by the HFS trains. Finally, we also assessed

the correlation between inter-individual variations in vision- and sensorimotor-related PAF, and

between the estimates of PAF obtained using the different approaches (ICA vs subtraction-based

method to isolate vision- and sensorimotor-related alpha-band activity, CoG vs FOOOF method to

estimate PAF).

Because the primary aim of our study was to test a possible association between PAF of

sensorimotor alpha-band oscillations and the susceptibility to develop central sensitization, and

because previous studies showed that the extent of the area of secondary hyperalgesia may be a

more reliable readout as compared to changes in pinprick ratings (47), our primary outcome was the

correlation between the sensorimotor PAF and the radius of the HFS-induced area of increased

pinprick sensitivity. All other correlations were considered as exploratory, thus we did not apply

corrections for multiple comparisons.

RESULTS

Pinprick sensitivity before and after HFS

After HFS, all but one participant reported an increased sensitivity to pinprick stimuli at the sensitized

forearm as compared to the control arm (Figure 3A). The repeated-measures ANOVA showed a

significant TIME × ARM interaction (F(1, 30) = 48.414, p = <.001, η²ₚ = 0.617).

Post hoc comparisons confirmed a significant increase in pinprick ratings at the HFS-treated arm,

with a mean increase of 11.854 (t(31) = 5.424, p < 0.001, Cohen's d = 0.959 [0.533 to 1.374], 95% CI

for mean difference: 7.397 to 16.311). Conversely, there was a significant decrease in ratings at the

control arm, with a mean reduction of 3.312 (t(31) = −4.177, p < 0.001, Cohen's d = −0.738 [−1.126 to

−0.342], 95% CI for mean difference: −4.930 to −1.695).

Figure 3. **A**. Individual intensity ratings for pinprick stimuli delivered to the HFS-sensitized forearm

and the contralateral non-sensitized forearm, before and after HFS (T-20 and T+40). Note the clear

increase in pinprick ratings at the sensitized forearm due to HFS-induced sensitization, and the

decrease in pinprick ratings at the non-sensitized forearm possibly due to perceptual habituation. **B**.

Estimated area of HFS-induced secondary hyperalgesia (grey contours: individual participants; black

contour: group-level average).

HFS-induced area of secondary hyperalgesia

At the HFS-treated forearm, an area of increased pinprick sensitivity was found in all but one

participant (Figure 3B). Across participants showing an area of increased sensitivity, the extent was

101 ± 32 mm along the proximal-distal axis, 53 ± 20 mm along the lateral-medial axis, 77 ± 29 mm

and 66 ± 27 mm along the two diagonal axes. The grand average estimated area radius was 68 ± 22

mm.

Vision- and sensorimotor-related alpha-band activity isolated using the ICA-based method

After blind source separation using ICA, 19.4 ± 5.5 ICs capturing alpha-band activity were identified in

31 out of 32 participants, of which 11.9 ± 5.4 ICs were categorized as vision-related (VISUAL-ALPHA)

in 31 out of 32 participants, 5.5 ± 5 ICs were categorized as sensorimotor-related (SM-ALPHA) in 24

out of 32 participants and 4.5 ± 4.3 ICs were categorized as mixed (MIXED-ALPHA) in 22 out of 32

participants.

The reconstructed VISUAL-ALPHA signals clearly captured vision-related alpha-band activity that

increased during closure of the eyes, was only minimally affected by finger movements and, most

importantly, displayed an occipital scalp topography compatible with activity originating

predominantly from visual areas (Figure 4A). Similarly, the SM-ALPHA signals captured sensorimotor-

related alpha-band activity that decreased during movement of the fingers, was minimally influenced

by closure of the eyes, and exhibited a bilateral centroparietal scalp topography compatible with

activity originating from the representation of the left and right hands within the sensorimotor

cortex (Figure 4B). The reconstructed MIXED-ALPHA signal captured alpha-band activity that

predominated over posterior regions and tended to both increase during closure of the eyes and

decrease during finger movements (Figure S1).

Figure 4. **A.** Group-level average frequency spectra of the ICA-separated EEG signals capturing vision-

related alpha-band activity (VISUAL-ALPHA signals) at occipital electrodes in the eyes closed at rest

(blue), eyes open at rest (red) and eyes open during finger movements (green) conditions. Note the

clear increase in alpha-band amplitude during closure of the eyes. Also note the occipital topography

of the peak. **B.** Group-level average frequency spectra of the ICA-separated EEG signals capturing

sensorimotor-related alpha-band activity (SM-ALPHA signals) at central electrodes. Note the selective

decrease of amplitude during finger movements, and the bilateral central-parietal topography

compatible with activity originating from the left and right sensorimotor cortices.

Vision- and sensorimotor-related alpha-band activity isolated using the subtraction-based method

The group-level averages of the subtracted frequency spectra are shown in Figure 5. As expected, the

subtraction spectrum EYES-CLOSED-REST minus EYES-OPEN-REST exhibited a clear peak of alpha-

band activity that was maximal over occipital electrodes, compatible with alpha oscillations

originating predominantly from the visual cortex (Figure 5A).

The subtraction spectrum EYES-OPEN-REST minus EYES-OPEN-MVT also exhibited a peak of alpha-

band activity, with a topography maximal at central and bilateral parietal electrodes, compatible with

activity originating from the hand representation within the left and right sensorimotor cortices

(Figure 5B). As compared to the signals obtained using the ICA-based method, the contribution of 1/f

noise was minimal in these subtracted spectra.

Figure 5. **A.** Group-level average of the subtracted EEG frequency spectra EYES-CLOSED-REST minus

EYES-OPEN-REST conditions at occipital electrodes. Note the clear occipital topography of the peak

indicating that this approach effectively isolated vision-related alpha-band activity. **B.** Group-level

average of the subtracted frequency spectra EYES-OPEN-REST minus EYES-OPEN-MVT conditions at

central electrodes. Note the bilateral central topography of the peak indicating that this approach

succeeded in isolating sensorimotor-related alpha-band activity. Also note the lack of an 1/f decay

(i.e. the aperiodic component of the EEG).

 Baseline peak alpha frequency (PAF) estimated using the CoG method

Using the CoG approach and a 9-11 Hz window applied to the VISUAL-ALPHA, SM-ALPHA and MIXED-

ALPHA spectra obtained with the ICA-based method, the PAF of vision-related alpha-band activity

estimated at occipital electrodes in the VISUAL-ALPHA signals during closure of the eyes was 10.0 ±

0.2 Hz (visual PAF). The PAF of sensorimotor-related alpha-band activity estimated at central

electrodes in the eyes open condition at rest was 10.0 ± 0.1 Hz (sensorimotor PAF). The PAF

estimated in the MIXED-ALPHA spectra was 10.0 ± 0.1 Hz at occipital electrodes during closure of the

eyes, and 10.0 ± 0.1 Hz at central electrodes in the eyes open at rest condition. Using the same CoG

approach applied to the spectra obtained with the subtraction-based approach, data from one

participant was rejected for estimation of the visual PAF and data from three participants were

rejected for estimation of the sensorimotor PAF as their spectra did not exhibit a peak within the

frequency window. The visual PAF was 10.0 ± 0.3 Hz while the sensorimotor-related PAF was 9.9 ±

0.5 Hz. Average PAF estimates obtained using the larger 7-13 Hz frequency window are reported in

Table 1.

| | | | | Visual PAF | | | | | Sensorimotor PAF | | | | |
| | | | | ICA spectra | | | Subtracted spectra | | ICA spectra | | | Subtracted spectra | |
				CoG (9-11 Hz)	CoG (7-13 Hz)	FOOOF	CoG (9-11 Hz)	CoG (7-13 Hz)	CoG (9-11 Hz)	CoG (7-13 Hz)	FOOOF	CoG (9-11 Hz)	CoG (7-13 Hz)
Visual PAF	ICA spectra	CoG (9-11 Hz)	10.0 ±0.2 Hz		0,93	0,87	0,93	0,80	0,27	0,34	0,14	0,15	0,27
		CoG (7-13 Hz)	10.0±0.3 Hz			0,84	0,90	0,92	0,18	0,26	-0,02	0,07	0,16
		FOOOF	10.1 ±0.7 Hz				0,89	0,79	0,13	0,22	0,01	0,08	0,22
	Subtracted spectra	CoG (9-11 Hz)	10.0 ±0.3 Hz					0,89	0,16	0,29	0,25	0,10	0,38
		CoG (7-13 Hz)	10.1 ±0.7 Hz						0,08	0,18	0,15	0,08	0,32
Sensorimotor PAF	ICA spectra	CoG (9-11 Hz)	10.0 ±0.1 Hz							0,90	0,42	0,79	0,70
		CoG (7-13 Hz)	9.9 ±0.3 Hz								0,40	0,77	0,72
		FOOOF	9.7 ±1.4 Hz									0,56	0,73
	Subtracted spectra	CoG (9-11 Hz)	9.9 ±0.5 Hz										0,75
		CoG (7-13 Hz)	10.0 ±0.9 Hz										

Table 1. Spearman's correlation coefficients between visual and sensorimotor PAF estimates

obtained using the ICA-based or subtraction-based approaches to isolate visual- and sensorimotor-

related alpha-band activity, and the CoG (frequency windows: 9-11 Hz and 7-13 Hz) or FOOOF

methods to estimate peak frequency. Positive and negative correlations are coded in green and red,

respectively. Significant correlations (p<0.05) are highlighted with red borders

Estimation of Peak Alpha Frequency (PAF) using the FOOOF method

Using the FOOOF approach applied to the VISUAL-ALPHA, SM-ALPHA and MIXED-ALPHA spectra

obtained with the ICA-based method, the PAF of vision-related alpha-band activity was 10.1 ± 0.7 Hz,

while the PAF for sensorimotor-related alpha-band activity was 9.7 ± 1.4 Hz. The PAF estimated in

the MIXED-ALPHA spectra was 10.3 ± 1.0 Hz at occipital electrodes during closure of the eyes, and

10.4 ± 1.2 Hz at central electrodes in the eyes open at rest condition. Average PAF estimates

obtained using the larger 7-13 Hz frequency window are reported in Table 1.

Primary analysis: correlation between baseline sensorimotor PAF and the area radius of HFS-

induced secondary hyperalgesia

Using the data from the 24/32 participants in which the ICA-based approach succeeded in isolating

sensorimotor-related alpha-band activity, no significant correlation between the sensorimotor PAF

and the radius of the HFS-induced area of secondary hyperalgesia was found, neither when using the

CoG estimates of PAF (CoG window 9-11 Hz: Spearman's rho = 0.167, p = 0.434, Pearson's r = 0.267, p

= 0.207; CoG window 7 − 13 Hz: Spearman's rho = 0.207, p = 0.330, Pearson's r = 0.132, p = 0.539),

nor when using the FOOOF estimates of PAF (Spearman's rho = -0.203, p = 0.339, Pearson's r = -

0.252, p = 0.235) (Table 2 and Figure 6).

Similarly, when using the PAF estimates obtained from the subtraction-based spectra, there was no

significant correlation between the sensorimotor PAF and the radius of the HFS-induced area of

secondary hyperalgesia (CoG window 9-11 Hz: Spearman's rho=-0.068, p=0.729, Pearson's r = -0.092,

p = 0.641; CoG window 7-13 Hz: Spearman's rho=-0.095, p=0.624, Pearson's r = -0.107, p = 0.581)

(Table 2 and Figure 6).

	Visual PAF					Sensorimotor PAF				
	ICA spectra			Subtracted spectra		ICA spectra			Subtracted spectra	
	CoG (9-11 Hz)	CoG (7-13 Hz)	FOOOF	CoG (9-11 Hz)	CoG (7-13 Hz)	CoG (9-11 Hz)	CoG (7-13 Hz)	FOOOF	CoG (9-11 Hz)	CoG (7-13 Hz)
Secondary hyperalgesia radius	-0,23	-0,33	-0,23	-0,12	-0,28	0,17	0,21	0,20	-0,07	-0,10
Change in pinprick sensitivity at HFS forearm (ΔNRS_{HFS})	-0,07	-0,16	0,02	-0,25	-0,29	-0,03	-0,11	-0,02	0,00	-0,11
Change in pinprick sensitivity at non-sensitized forearm (ΔNRS_{NS})	-0,49	-0,42	-0,44	-0,48	-0,36	-0,17	-0,25	0,00	-0,01	-0,08
Change in pinprick sensitivity at HFS minus non-sensitized forearm ($\Delta\Delta NRS$)	0,09	-0,02	0,20	-0,08	-0,16	0,12	0,06	-0,02	0,07	-0,07
Baseline pinprick ratings	0,31	0,29	0,40	0,25	0,20	0,22	0,25	-0,34	-0,03	0,07
HFS pain ratings	0,20	0,09	0,31	0,15	0,01	0,36	0,39	-0,03	0,15	0,17
Electrical detection threshold	-0,29	-0,11	-0,17	-0,12	0,03	-0,13	-0,12	0,05	0,11	0,05

Table 2. Spearman's correlation coefficients between visual and sensorimotor PAF estimates and the following measures: the area radius of HFS-induced secondary hyperalgesia area, the changes in pinprick sensitivity at the HFS-sensitized and non-sensitized forearms, the detection threshold to a single electrical pulse averaged across the two forearms, the baseline pinprick ratings averaged across the two forearms, and the pain ratings reported during HFS. Positive and negative correlations are coded in green and red, respectively. Significant correlations (p<0.05) are highlighted with red borders.

Figure 6. Scatter plots showing the absence of a significant correlation between the radius of the HFS-induced area of secondary hyperalgesia and the sensorimotor peak alpha frequency (PAF), regardless of the method used to isolate sensorimotor-related activity (A: ICA-based approach; B: subtraction-based approach) or the method used to estimate the PAF (center of gravity using a narrow 9-11 Hz or wide 7-13 Hz frequency window, FOOOF algorithm).

Exploratory analyses

Correlation between baseline sensorimotor PAF and changes in pinprick sensitivity after HFS. Such as for the area of HFS-induced secondary hyperalgesia, there was no significant correlation between the sensorimotor PAF and (1) the magnitude of the HFS-induced increase in pinprick sensitivity at the sensitized forearm compared to the contralateral forearm ($\Delta\Delta NRS$), (2) the change in pinprick

sensitivity at the HFS-sensitized forearm (ΔNRS_{HFS}) and (3) the change in pinprick sensitivity at the

contralateral non-sensitized forearm (ΔNRS_{NS}) (Table 2).

Correlation between baseline visual PAF and the area of HFS-induced secondary hyperalgesia. Such as

for the sensorimotor PAF, there was no significant correlation between the visual PAF and the radius

of the area of HFS-induced secondary hyperalgesia (Table 2).

Correlation between baseline visual PAF and the changes in pinprick sensitivity after HFS. There was

also no significant correlation between the visual PAF and the increase in pinprick sensitivity at the

sensitized forearm compared to the contralateral forearm ($\Delta\Delta NRS$) or the change in pinprick

sensitivity at the HFS-sensitized forearm (ΔNRS_{HFS}). However, and regardless of the method used to

estimate the visual PAF, there was a significant negative correlation between the visual PAF and the

change in pinprick sensitivity at the non-sensitized forearm (ΔNRS_{NS}): participants with a higher visual

PAF at baseline exhibited a stronger perceptual habituation to pinprick stimulation at the non-

sensitized forearm (Table 2 and Figure 7).

Correlation between baseline PAF and baseline pain ratings. There was no significant correlation

between either the visual PAF nor the sensorimotor PAF and (1) the detection threshold to single

electrical pulses averaged across the two forearms, (2) the baseline pinprick ratings averaged across

the two forearms, or (3) the intensity of the pain during HFS (Table 2).

Correlation between vision- and sensorimotor-related PAF. Both visual and sensorimotor PAF

estimates derived using ICA-based and subtraction-based approaches to isolate visual- and

sensorimotor-related alpha-band activity and computed using the CoG and FOOOF methods were

highly correlated, indicating good consistency across the estimation techniques (Table 1). In contrast,

there was no significant correlation between the visual PAF and the sensorimotor PAF, indicating that

they likely reflect distinct oscillatory processes (Table 1). This highlights the importance of

considering these rhythms as functionally separate when assessing their relevance to perception.

Mixed alpha-band activity. Using the ICA-based approach, independent components (ICs) capturing

alpha-band activity that increased during closure of the eyes and decreased during finger movements

were identified in 22/32 participants. The PAF estimates of these MIXED-ALPHA signals were strongly

correlated with the visual PAF, but also showed significant correlation with the sensorimotor PAF

using the largest window, indicating that these signals may reflect the fact ICA did not fully dissociate

the two types of alpha-band activity (Table S1). There was no significant correlation between the PAF

of MIXED-ALPHA signals and the secondary hyperalgesia radius, the changes in pinprick sensitivity

after HFS, the electrical detection threshold and the HFS pain ratings. A weak correlation was

reported for the baseline pinprick ratings and mixed PAF at occipital electrodes (Table S2).

Figure 7. Scatter plots showing the correlation between the post-HFS decrease in pinprick sensitivity

at the non-sensitized forearm and the visual peak alpha frequency (PAF), present across methods

used to isolate visual-related activity (A: ICA-based approach; B: subtraction-based approach) and

methods to estimate the PAF (center of gravity using a narrow 9-11 Hz or wide 7-13 Hz frequency

window, FOOOF algorithm).

DISCUSSION

The results of the present study do not support our hypothesis that interindividual variations in peak-

alpha frequency are associated with the susceptibility to develop HFS-induced secondary

hyperalgesia. Indeed, in our sample of 32 healthy participants and using different approaches to

isolate sensorimotor- and vision-related alpha-band activity and estimate peak frequency in the

resting EEG, there was no significant correlation between sensorimotor or visual PAF estimates and

neither the spatial extent of the HFS-induced secondary hyperalgesia area, nor the magnitude of the

HFS-induced increase in pinprick sensitivity at the sensitized forearm.

Functional dissociation between vision- and sensorimotor-related alpha-band activity

562 Previous reports of a relationship between PAF and sensitivity to pain have often interpreted the

563 measured interindividual differences in PAF as reflecting differences in sensorimotor-related alpha-

564 band activity. However, the topographical distribution of the alpha-band in resting EEG predominates

565 over posterior areas, and its most prominent feature is its sensitivity to closure of the eyes. Hence,

566 measures of PAF in resting EEG may be expected to mainly reflect vision-related alpha-band

567 oscillations, even when the PAF is estimated at central electrodes overlying the sensorimotor cortex

568 (17,19,55). For this reason, several authors have attempted to isolate sensorimotor- from vision-

569 related alpha oscillations using an Independent Component Analysis and selecting components

570 displaying a central topography as compared to an occipital topography (18,52,54,65).

571 In the present study, we propose two functional approaches to isolate vision- and sensorimotor-

572 related alpha oscillations, based on the sensitivity of vision-related alpha oscillations to closure of the

573 eyes and the sensitivity of sensorimotor-related alpha oscillations to movement execution rather

574 than their topography. In the ICA-based approach, the functional properties of vision- and

575 sensorimotor-related alpha-band activity allowed separating alpha-band activity into visual-related

576 components that were sensitive to closure of the eyes and sensorimotor-related components that

577 were sensitive to the performance of bilateral finger movements. In the subtraction-based approach,

578 we subtracted the EEG spectra obtained during eyes open and eyes closed recordings at rest to

579 isolate the alpha-band increase occurring during eye closure, while we subtracted the EEG spectra

580 obtained during eyes open at rest and eyes open during bilateral finger movements to capture the

581 change in alpha-band activity occurring during movement. Both the ICA-based approach and the

582 subtraction-based approach showed that the alpha-band activity sensitive to closure of the eyes had

583 a clear occipital topography compatible with activity originating from visual areas, whereas the

584 alpha-band activity sensitive to moving of the fingers exhibited a bilateral central topography,

585 compatible with activity originating from left and right sensorimotor areas.

586 It should be noted that in the ICA-based approach, visual-related alpha-band activity was isolated in

587 31/32 participants, whereas sensorimotor-related alpha-band activity was isolated in 24/32

588 participants. This may be related to the fact that the sensorimotor mu rhythm is not always observed

589 (5).

Visual and sensorimotor PAF estimates

Previous studies using the center-of-gravity (CoG) method to estimate PAF have shown that the

choice of frequency window can markedly influence results (55,58,65). Some studies have employed

relatively narrow windows (9-11 Hz) to minimize the influence of the aperiodic 1/f component of the

EEG frequency spectrum (17–19,52), while other studies have used larger windows from 7 to 11 Hz

(57), 7.4 to 12 Hz (56) or 7.5 to 13 Hz (58). In the present study, the CoG was computed using both

narrow (9-11 Hz) and wide (7-13 Hz) windows to assess the robustness of the estimates. Additionally,

CoG estimates computed on the subtracted spectra are expected to be minimally influenced by the

1/f aperiodic component, as this component should be largely cancelled by the subtraction

procedure. Finally, we also used the FOOOF method to remove the aperiodic component and

estimate PAF using a Gaussian fit rather than computing the center of gravity (62).

Both for the visual PAF and the sensorimotor PAF, there was a strong correlation between the

different PAF estimates, indicating good consistency between estimation techniques.

Interestingly, and regardless of the method used to estimate PAF, the interindividual variations in

visual PAF were not significantly correlated with the interindividual variations in sensorimotor PAF.

This indicates that the two reflect at least partially distinct functional processes. Supporting this

distinction, previous studies have shown that the mu-rhythm and alpha rhythm can differ in

frequency and amplitude despite occupying the same frequency band (66). In light of these

observations, future research should investigate whether the previously reported association

between PAF and pain sensitivity (18,19,21,57,67) are predominantly driven by visual- versus

sensorimotor-related alpha-band activity.

Lack of association between PAF and the susceptibility to develop secondary hyperalgesia

Our study found no significant association between either visual or sensorimotor PAF and the

susceptibility to develop HFS-induced secondary hyperalgesia. This suggests that the frequency

characteristics of alpha-band activity do not play a modulatory role in the induction of central

sensitization. Notably, recent studies examining whether cognitive factors that are known to

modulate pain perception – such as attention and expectations – may also influence the

susceptibility to develop experimentally-induced central sensitization have reported mostly negative

results (68,69). Collectively, this suggests that the propensity to develop central sensitization may be

minimally influenced by cognitive factors and the functional state of the brain.

Whether this absence of association between PAF and secondary hyperalgesia is specific for HFS-

induced hyperalgesia in healthy volunteers, or whether it generalizes to other experimental methods

to induce hyperalgesia, as well as to hyperalgesia in patients remains an open question requiring

further studies.

Finally, it is important to acknowledge that the sample size in our study was based on prior work

reporting relatively large correlation coefficients. As such, our study was underpowered to detect

smaller effect sizes.

Association between visual PAF and habituation at the non-sensitized forearm

An incidental finding of the present study was a negative association between visual PAF and

perceptual habituation to pinprick stimulation. Specifically, participants with a higher visual PAF

showed a more pronounced decrease in pinprick ratings at the non-sensitized forearm. This raises an

intriguing possibility: could previous findings linking a lower PAF with greater pain sensitivity be

partially explained by this relationship between PAF and habituation? If individuals with a lower PAF

habituate less to repeated or sustained nociceptive stimuli as compared to individuals with a higher

PAF, they would also be expected to report higher average pain ratings when exposed to such

stimuli. Such an interpretation would also be compatible with the fact that the studies having shown

a relationship between PAF and sensitivity to pain have relied on methods inducing sustained pain

such as capsaicin-heat pain, NGF injection, and tonic heat pain (18–20,67), while studies employing

more phasic pain stimuli have reported no or only marginally-significant associations between PAF

and pain ratings (70).

CONCLUSION

In this study conducted in healthy volunteers, interindividual variations in visual or sensorimotor PAF

were not significantly associated with differences in the susceptibility to develop HFS-induced

secondary hyperalgesia. This suggests that PAF does not modulate the propensity to develop central

sensitization. However, exploratory analyses revealed that individuals with a higher visual PAF

tended to exhibit stronger perceptual habituation to pinprick stimulation. Further studies are needed

to confirm this incidental observation, and examine whether it might contribute to the previously

reported   association   between   PAF   and   pain   sensitivity.

DATA AVAILABILITY:

Source data are available at https://doi.org/10.6084/m9.figshare.29145725

SUPPLEMENTAL MATERIAL:

Supplemental Fig. S1 & Table S1-S2 DOI: https://doi.org/10.6084/m9.figshare.29145695

CURRENT ADDRESS:

Same as affiliations

GRANTS

This study was supported by a funding from the Innovative Medicines Initiative 2 Joint Undertaking under grant agreement No [777500]. This Joint Undertaking receives support from the European Union's Horizon 2020 research and innovation program and EFPIA. www.imi.europa.eu; www.imi-paincare.eu.

DISCLAIMERS

The statements and opinions presented here reflect the author's view and neither IMI nor the European Union, EFPIA, or any Associated Partners are responsible for any use that may be made of the information contained therein.

AUTHORS CONTRIBUTIONS

Louisien Lebrun[1]*, Gloria Ricci[2], Arthur Courtin[1,3], Emanuel N. van den Broeke[1,4], Cédric Lenoir[1], André Mouraux[1]

Louisien Lebrun: conceived and designed research, performed experiments, analyzed data, interpreted results of experiment, prepared figures, drafted manuscript, approved final version of manuscript

Gloria Ricci: conceived and designed research, analyzed data, edited and revised the manuscript, approved final version of manuscript

Arthur Courtin: conceived and designed research, edited and revised the manuscript, approved final version of manuscript

Emanuel N. van den Broeke: conceived and designed research, edited and revised the manuscript, approved final version of manuscript

Cédric Lenoir: conceived and designed research, edited and revised the manuscript, approved final

version of manuscript

André Mouraux: conceived and designed research, analyzed data, interpreted results of experiment,

prepared figures, edited and revised the manuscript, approved final version of manuscript

References

1.   Berger H. Ueber das Elektrenkephalogramm des Menschen. [Electrocephalography in man.]. J
Für Psychol Neurol. 1930;40:160–79.

2.   Omata K, Hanakawa T, Morimoto M, Honda M. Spontaneous Slow Fluctuation of EEG Alpha
Rhythm Reflects Activity in Deep-Brain Structures: A Simultaneous EEG-fMRI Study. PLoS ONE.
2013 Jun 18;8(6):e66869.

3.   Gastaut H. Etude electrocorticographique de la reactivite des rythmes rolandiques. Rev Neurol
Paris. 1952;87:176–82.

4.   Hughes SW, Crunelli V. Thalamic Mechanisms of EEG Alpha Rhythms and Their Pathological
Implications. The Neuroscientist. 2005 Aug 1;11(4):357–72.

5.   Pfurtscheller G, Neuper C, Andrew C, Edlinger G. Foot and hand area mu rhythms. Int J
Psychophysiol. 1997 Jun 1;26(1):121–35.

6.   Garakh Z, Novototsky-Vlasov V, Larionova E, Zaytseva Y. Mu rhythm separation from the mix
with alpha rhythm: Principal component analyses and factor topography. J Neurosci Methods.
2020 Dec 1;346:108892.

7.   Chatrian GE, Petersen MC, Lazarte JA. The blocking of the rolandic wicket rhythm and some
central changes related to movement. Electroencephalogr Clin Neurophysiol. 1959 Aug
1;11(3):497–510.

8.   Keil A, Mussweiler T, Epstude K. Alpha-band activity reflects reduction of mental effort in a
comparison task: A source space analysis. Brain Res. 2006 Nov 22;1121(1):117–27.

9.   Tran Y, Craig A, Craig R, Chai R, Nguyen H. The influence of mental fatigue on brain activity:
Evidence from a systematic review with meta-analyses. Psychophysiology. 2020;57(5):e13554.

10.   Kuhlman WN. Functional topography of the human mu rhythm. Electroencephalogr Clin
Neurophysiol. 1978 Jan 1;44(1):83–93.

11.   de Vries M, Wilder-Smith OH, Jongsma ML, van den Broeke EN, Arns M, van Goor H, et al.
Altered resting state EEG in chronic pancreatitis patients: toward a marker for chronic pain. J
Pain Res. 2013 Nov 25;6:815–24.

12.   Boord P, Siddall PJ, Tran Y, Herbert D, Middleton J, Craig A. Electroencephalographic slowing and
reduced reactivity in neuropathic pain following spinal cord injury. Spinal Cord. 2008
Feb;46(2):118–23.

13.   Sarnthein J, Stern J, Aufenberg C, Rousson V, Jeanmonod D. Increased EEG power and slowed
dominant frequency in patients with neurogenic pain. Brain. 2006 Jan;129(Pt 1):55–64.

14.   Walton KD, Dubois M, Llinas RR. Abnormal thalamocortical activity in patients with Complex
Regional Pain Syndrome (CRPS) type I. Pain. 2010 Jul;150(1):41–51.

15.   Lim M, Kim JS, Kim DJ, Chung CK. Increased Low- and High-Frequency Oscillatory Activity in the
Prefrontal Cortex of Fibromyalgia Patients. Front Hum Neurosci. 2016;10:111.

16. Kim JA, Bosma RL, Hemington KS, Rogachov A, Osborne NR, Cheng JC, et al. Neuropathic pain
and pain interference are linked to alpha-band slowing and reduced beta-band
magnetoencephalography activity within the dynamic pain connectome in patients with multiple
sclerosis. Pain. 2019 Jan;160(1):187–97.

17. Furman AJ, Prokhorenko M, Keaser ML, Zhang J, Chen S, Mazaheri A, et al. Sensorimotor Peak
Alpha Frequency Is a Reliable Biomarker of Prolonged Pain Sensitivity. Cereb Cortex. 2020 Jun
27;bhaa124.

18. Furman AJ, Meeker TJ, Rietschel JC, Yoo S, Muthulingam J, Prokhorenko M, et al. Cerebral peak
alpha frequency predicts individual differences in pain sensitivity. Neuroimage. 2018 Feb
15;167:203–10.

19. Furman AJ, Thapa T, Summers SJ, Cavaleri R, Fogarty JS, Steiner GZ, et al. Cerebral peak alpha
frequency reflects average pain severity in a human model of sustained, musculoskeletal pain. J
Neurophysiol. 2019 Oct 1;122(4):1784–93.

20. Chowdhury NS, Bi C, Furman AJ, Chiang AKI, Skippen P, Si E, et al. Predicting Individual Pain
Sensitivity Using a Novel Cortical Biomarker Signature. JAMA Neurol. 2025 Jan 27;82(3):237–46.

21. Nir RR, Sinai A, Raz E, Sprecher E, Yarnitsky D. Pain assessment by continuous EEG: association
between subjective perception of tonic pain and peak frequency of alpha oscillations during
stimulation and at rest. Brain Res. 2010 Jul 16;1344:77–86.

22. Grandy TH, Werkle-Bergner M, Chicherio C, Schmiedek F, Lövdén M, Lindenberger U. Peak
individual alpha frequency qualifies as a stable neurophysiological trait marker in healthy
younger and older adults. Psychophysiology. 2013;50(6):570–82.

23. Latremoliere A, Woolf CJ. Central sensitization: a generator of pain hypersensitivity by central
neural plasticity. J Pain. 2009 Sep;10(9):895–926.

24. Woolf CJ. Evidence for a central component of post-injury pain hypersensitivity. Nature. 1983
Dec;306(5944):686–8.

25. van den Broeke EN, Mouraux A. High-frequency electrical stimulation of the human skin induces
heterotopical mechanical hyperalgesia, heat hyperalgesia, and enhanced responses to
nonnociceptive vibrotactile input. J Neurophysiol. 2014 Apr;111(8):1564–73.

26. Lebrun L, Lenoir C, Leone C, Broeke EN van den, Caspani O, Schilder A, et al. Strength, extent and
duration of secondary hyperalgesia induced by high-frequency electrical stimulation of the foot
compared to the volar forearm of healthy human volunteers. PLOS ONE. 2025 Feb
7;20(2):e0318934.

27. Lebrun L, Stouffs A, Hermans E, van den Broeke EN, Leone C, Caspani O, et al. Repeated exposure
to cutaneous high-frequency electrical stimulation prolongs the time course of secondary
hyperalgesia. J Neurophysiol [Internet]. 2025 May 12 [cited 2025 May 15]; Available from:
https://journals.physiology.org/doi/abs/10.1152/jn.00130.2025

28. Basbaum AI, Bautista DM, Scherrer G, Julius D. Cellular and Molecular Mechanisms of Pain. Cell.
2009 Oct 16;139(2):267–84.

29. Ikeda H, Heinke B, Ruscheweyh R, Sandkühler J. Synaptic plasticity in spinal lamina I projection
neurons that mediate hyperalgesia. Science. 2003 Feb 21;299(5610):1237–40.

30. Kronschläger MT, Drdla-Schutting R, Gassner M, Honsek SD, Teuchmann HL, Sandkühler J.
Gliogenic LTP spreads widely in nociceptive pathways. Science. 2016 Dec 2;354(6316):1144–8.

31. Sandkühler J, Gruber-Schoffnegger D %J C opinion in pharmacology. Hyperalgesia by synaptic
long-term potentiation (LTP): an update. 2012;12(1):18–27.

32. Magerl W, Wilk SH, Treede RD. Secondary hyperalgesia and perceptual wind-up following
intradermal injection of capsaicin in humans. Pain. 1998 Jan;74(2):257–68.

33. Treede RD, Meyer RA, Raja SN, Campbell JN. Peripheral and central mechanisms of cutaneous
hyperalgesia. Prog Neurobiol. 1992 Apr;38(4):397–421.

34. Baumann TK, Simone DA, Shain CN, LaMotte RH. Neurogenic hyperalgesia: the search for the
primary cutaneous afferent fibers that contribute to capsaicin-induced pain and hyperalgesia. J
Neurophysiol. 1991 Jul;66(1):212–27.

35. Randic M, Jiang MC, Cerne R. Long-term potentiation and long-term depression of primary
afferent neurotransmission in the rat spinal cord. J Neurosci. 1993 Dec 1;13(12):5228–41.

36. Vollert J, Magerl W, Baron R, Binder A, Enax-Krumova EK, Geisslinger G, et al. Pathophysiological
mechanisms of neuropathic pain: comparison of sensory phenotypes in patients and human
surrogate pain models. PAIN. 2018 Jun;159(6):1090–102.

37. Leone CM, Lenoir C, van den Broeke EN. Assessing signs of central sensitization: A critical review
of physiological measures in experimentally induced secondary hyperalgesia. Eur J Pain. 2025
Mar;29(3):e4733.

38. Gousset S, Cappe M, Lenoir C, Steyaert A, Lavand'homme P, Mouraux A, et al. Preoperative
susceptibility to developing secondary hyperalgesia is associated with post-thoracotomy pain at
2 months. Eur J Pain Lond Engl. 2025 Jan;29(1):e4768.

39. Sacco P, Prior M, Poole H, Nurmikko T. Repetitive transcranial magnetic stimulation over primary
motor vs non-motor cortical targets; effects on experimental hyperalgesia in healthy subjects.
BMC Neurol. 2014 Sep 4;14(1):166.

40. Farina S, Valeriani M, Rosso T, Aglioti S, Tamburin S, Fiaschi A, et al. Transient inhibition of the
human motor cortex by capsaicin-induced pain. A study with transcranial magnetic stimulation.
Neurosci Lett. 2001 Nov 13;314(1–2):97–101.

41. Giannoni-Luza S, Pacheco-Barrios K, Cardenas-Rojas A, Mejia-Pando PF, Luna-Cuadros MA,
Barouh JL, et al. Noninvasive motor cortex stimulation effects on quantitative sensory testing in
healthy and chronic pain subjects: a systematic review and meta-analysis. PAIN. 2020
Sep;161(9):1955.

42. Steegen S, Tuerlinckx F, Gelman A, Vanpaemel W. Increasing Transparency Through a Multiverse
Analysis. Perspect Psychol Sci. 2016 Sep 1;11(5):702–12.

43. Nicholls ME, Thomas NA, Loetscher T, Grimshaw GM. The Flinders Handedness survey
(FLANDERS): a brief measure of skilled hand preference. Cortex. 2013 Dec;49(10):2914–26.

44. Kohn JS Michael. Correlation sample size | Sample Size Calculators [Internet]. [cited 2025 May 3].
Available from: https://sample-size.net/correlation-sample-size/

45. van den Broeke EN, Gousset S, Bouvy J, Stouffs A, Lebrun L, van Neerven SGA, et al.
Heterosynaptic facilitation of mechanical nociceptive input is dependent on the frequency of
conditioning stimulation. J Neurophysiol. 2019 Jul 10;

46. Cayrol T, Pitance L, Roussel N, Mouraux A, van den Broeke EN. No evidence of widespread
mechanical pressure hyperalgesia after experimentally induced central sensitization through skin
nociceptors. 2018;3(6):e691.

47. Cayrol T, Lebleu J, Mouraux A, Roussel N, Pitance L, Broeke EN. Within- and between-session
reliability of secondary hyperalgesia induced by electrical high-frequency stimulation. Eur J Pain.
2020 Jun 30;ejp.1613.

48. Lenoir C, Plaghki L, Mouraux A, van den Broeke EN. Quickly responding C-fibre nociceptors
contribute to heat hypersensitivity in the area of secondary hyperalgesia. 2018;596(18):4443–55.

49. van den Broeke EN, Lenoir C, Mouraux A. Secondary hyperalgesia is mediated by heat-insensitive
A-fibre nociceptors. 2016;594(22):6767–76.

50. Hansen MS, Wetterslev J, Pipper CB, Østervig R, Asghar MS, Dahl JB. The Area of Secondary
Hyperalgesia following Heat Stimulation in Healthy Male Volunteers: Inter- and Intra-Individual
Variance and Reproducibility. PLOS ONE. 2016 May 11;11(5):e0155284.

51. Werner MU, Petersen KL, Rowbotham MC, Dahl JB. Healthy Volunteers Can Be Phenotyped
Using Cutaneous Sensitization Pain Models. PLOS ONE. 2013 May 9;8(5):e62733.

52. De Martino E, Gregoret L, Zandalasini M, Graven-Nielsen T. Slowing in Peak-Alpha Frequency
Recorded After Experimentally-Induced Muscle Pain is not Significantly Different Between High
and Low Pain-Sensitive Subjects. J Pain. 2021 Dec 1;22(12):1722–32.

53. Valentini E, Halder S, McInnerney D, Cooke J, Gyimes IL, Romei V. Assessing the specificity of the
relationship between brain alpha oscillations and tonic pain. NeuroImage. 2022 Jul;255:119143.

54. Seminowicz DA, Thapa T, Furman AJ, Summers SJ, Cavaleri R, Fogarty JS, et al. Slow peak alpha
frequency and corticomotor depression linked to high pain susceptibility in transition to
sustained pain. 2018;

55. Furman AJ, Prokhorenko M, Keaser ML, Zhang J, Chen S, Mazaheri A, et al. Prolonged Pain
Reliably Slows Peak Alpha Frequency by Reducing Fast Alpha Power. bioRxiv. 2021;2021.07.
22.453260.

56. Raghuraman N, Wang Y, Schenk LA, Furman AJ, Tricou C, Seminowicz DA, et al. Neural and
behavioral changes driven by observationally-induced hypoalgesia. Sci Rep. 2019 Dec
24;9(1):19760.

57. Millard SK, Furman AJ, Kerr A, Seminowicz DA, Naidu B, Gao-Smith F, et al. Predicting Post-
operative Pain in Lung Cancer Patients using Pre-operative Peak Alpha Frequency. medRxiv.
2021;

58. McLain NJ, Yani MS, Kutch JJ. Analytic consistency and neural correlates of peak alpha frequency
in the study of pain. J Neurosci Methods. 2022 Feb 15;368:109460.

59. Cavaleri R, McLain NJ, Heindel M, Schrepf A, Rodriguez LV, Kutch JJ. Peak alpha frequency is
related to the degree of widespread pain, but not pain intensity or duration, among people with
urologic chronic pelvic pain syndrome. PAIN Rep. 2025 Apr;10(2):e1251.

60. Hu L, Zhang Z. EEG signal processing and feature extraction. Springer; 2019.

61. Klimesch W, Schimke H, Pfurtscheller G. Alpha frequency, cognitive load and memory
performance. Brain Topogr. 1993 Mar 1;5(3):241–51.

62. Donoghue T, Haller M, Peterson EJ, Varma P, Sebastian P, Gao R, et al. Parameterizing neural
power spectra into periodic and aperiodic components. Nat Neurosci. 2020 Dec;23(12):1655–65.

63. Trajkovic J, Sack AT, Romei V. EEG-based biomarkers predict individual differences in TMS-
induced entrainment of intrinsic brain rhythms. Brain Stimulat. 2024 Mar 1;17(2):224–32.

64. Gerster M, Waterstraat G, Litvak V, Lehnertz K, Schnitzler A, Florin E, et al. Separating Neural
Oscillations from Aperiodic 1/f Activity: Challenges and Recommendations. Neuroinformatics.
2022 Oct;20(4):991–1012.

65. Valentini E, Halder S, McInnersey D, Cooke J, Romei V. Assessing the specificity of the
relationship between brain alpha oscillations and tonic pain. bioRxiv. 2019;

66. Bender A, Voytek B, Schaworonkow N. Resting-state alpha and mu rhythms change shape across
development but lack diagnostic sensitivity for ADHD and autism [Internet]. bioRxiv; 2025 [cited
2025   May   9].   p.   2023.10.13.562301.   Available   from:
https://www.biorxiv.org/content/10.1101/2023.10.13.562301v3

67. Nir RR, Sinai A, Moont R, Harari E, Yarnitsky D. Tonic pain and continuous EEG: prediction of
subjective pain perception by alpha-1 power during stimulation and at rest. Clin Neurophysiol.
2012 Mar;123(3):605–12.

68. Gousset S, Cayrol T, Papleux M, Meulders A, Mouraux A, van den Broeke EN. Studying the Effect
of   Expectations   on   High-Frequency   Electrical   Stimulation-Induced   Pain   and   Pinprick
Hypersensitivity. J Pain. 2024 Dec 1;25(12):104682.

69. Della Porta D, Scheirman E, Legrain V. Top-down attention does not modulate mechanical
hypersensitivity consecutive to central sensitization: insights from an experimental analysis.
PAIN. 2024 Sep;165(9):2098.

70. May ES, Tiemann L, Gil Ávila C, Bott FS, Hohn VD, Gross J, et al. Assessing the predictive value of
peak alpha frequency for the sensitivity to pain. Pain. 2025 Mar 11;

Figure 1. A. Timeline of the experiment. HFS was applied to the right volar forearm. EEG segments lasting 5 minutes and including three conditions (eyes open at rest, eyes closed at rest, eyes open while performing bilateral finger movements) were recorded 25 and 5 minutes before HFS (T-25 and T-5), immediately after HFS, and approximately 35 minutes after HFS (T+35). Pinprick sensitivity of the left and right volar forearms was assessed before HFS (T-20) and approximately 40 minutes after HFS (T+40). The spatial extent of the HFS-induced area of secondary hyperalgesia was assessed approximately 40 minutes after HFS (T+40). **B.** HFS was delivered to the forearm skin using a cathode consisting of 10 blunt tungsten pins arranged on a circle with a diameter of 5 mm and a 24x20 mm anode conductive gel pad placed next to the cathode.

Figure 2. **A.** EEG frequency spectrum of a representative independent component (IC) categorized as capturing vision-related alpha-band activity (VIS-ALPHA) that was increased during closure of the eyes (EYES-CLOSED-REST condition) as compared to the eyes open conditions (EYES-OPEN-REST and EYES-OPEN-MOVEMENT). **B.** EEG frequency spectrum of a representative IC categorized as capturing sensorimotor-related alpha-band activity (SM-ALPHA) that was decreased during finger movements (EYES-OPEN-MVT condition) as compared to rest (EYES-OPEN-REST and EYES-CLOSED-REST conditions). Y-axis: arbitrary units. X-axis: frequency in Hz. The vertical grey lines represent the 6 Hz frequency window used to assess the alpha-band activity.

Figure 3. **A**. Individual intensity ratings for pinprick stimuli delivered to the HFS-sensitized forearm and the contralateral non-sensitized forearm, before and after HFS (T-20 and T+40). Note the clear increase in pinprick ratings at the sensitized forearm due to HFS-induced sensitization, and the decrease in pinprick ratings at the non-sensitized forearm possibly due to perceptual habituation. **B**. Estimated area of HFS-induced secondary hyperalgesia (grey contours: individual participants; black contour: group-level average). The origin (+) corresponds to the center of the cathode of the HFS electrode.

Figure 4. **A.** Group-level average frequency spectra of the ICA-separated EEG signals capturing vision-related alpha-band activity (VISUAL-ALPHA signals) at occipital electrodes in the EYES-CLOSED-REST (blue), EYES-OPEN-REST (red) and EYES-OPEN-MVT (green) conditions. Note the clear increase in alpha-band amplitude during closure of the eyes. Also note the occipital topography of the peak. **B.** Group-level average frequency spectra of the ICA-separated EEG signals capturing sensorimotor-related alpha-band activity (SM-ALPHA signals) at central electrodes. Note the selective decrease of amplitude during finger movements, and the bilateral central-parietal topography compatible with activity originating from the left and right sensorimotor cortices.

Figure 5. **A.** Group-level average of the subtracted EEG frequency spectra EYES-CLOSED-REST minus EYES-OPEN-REST conditions at occipital electrodes. Note the clear occipital topography of the peak indicating that this approach effectively isolated vision-related alpha-band activity. **B.** Group-level average of the subtracted frequency spectra EYES-OPEN-REST minus EYES-OPEN-MVT conditions at central electrodes. Note the bilateral central topography of the peak indicating that this approach succeeded in isolating sensorimotor-related alpha-band activity. Also note the lack of an 1/f decay (i.e. the aperiodic component of the EEG).

A. ICA-based approach: SM-ALPHA signals

B. Subtraction-based approach - EYES-OPEN-REST minus EYES-OPEN-MVT

Figure 6. Scatter plots showing the absence of a significant Spearman's correlation between the radius of the HFS-induced area of secondary hyperalgesia and the sensorimotor peak alpha frequency (PAF), regardless of the method used to isolate sensorimotor-related activity (A: ICA-based approach; B: subtraction-based approach) or the method used to estimate the PAF (center of gravity using a narrow 9-11 Hz or wide 7-13 Hz frequency window, FOOOF algorithm).

A. ICA-based approach: VISUAL-ALPHA signals

B. Subtraction-based approach - EYES-CLOSED-REST minus EYES-OPEN-REST

Figure 7. Scatter plots showing the Spearmans's correlation between the post-HFS decrease in pinprick sensitivity at the non-sensitized forearm and the visual peak alpha frequency (PAF), present across methods used to isolate visual-related activity (A: ICA-based approach; B: subtraction-based approach) and methods to estimate the PAF (center of gravity using a narrow 9-11 Hz or wide 7-13 Hz frequency window, FOOOF algorithm).