Developmental dynamic transcriptomics reveals multiple effectors and transcription factors critical for *Ditylenchus destructor* parasitism

Author: Yangyang Chen[1]†, Shurong Zhang[1]†, Yayi Zhou[1], Xueyu Wang[1], Dexin Bo[1], Yucheng Liao[1], Boyan Hu[1], Yali Zhang[1,2], Noha Mohamed Ashry[1,3], Donghai Peng[1], Ming Sun[1*], and Dadong Dai[1,2*]

Affiliations:
[1]State Key Laboratory of Agricultural Microbiology, Huazhong Agricultural University, Wuhan 430070, China; Hubei Hongshan Laboratory, Wuhan 430070, China.
[2]Department of Entomology and Nematology, University of California, Davis, California, USA.
[3]Agriculture Microbiology Department, Faculty of Agriculture, Benha University, Qalubia 13736, Egypt.

Corresponding author:

*Dadong Dai and Ming Sun.

Email: daidadong@mail.hzau.edu.cn; m98sun@mail.hzau.edu.cn

†These authors contributed equally to this work.

Abstract

Plant-parasitic nematodes (PPNs) pose a significant threat to global agricultural production, yet their fundamental biological research remains relatively limited. The origin and evolution of PPNs remain unresolved mysteries, largely due to the lack of chromosome-level genomic data. Among them, migratory PPNs are considered a key transitional stage from free-living to obligate parasitism, as they exhibit both plant parasitism and fungal feeding. In this study, we constructed the chromosome-level genome of the sweet potato rot nematode and confirmed its genome comprises four chromosomes through Hi-C and karyotype analysis. Comparative analysis with two migratory PPNs, *B. xylophilus* and *A. besseyi*, revealed that the Nigon elements of *B. xylophilus* are largely conserved with those of the model organism *C. elegans*, whereas *D. destructor* and *A. besseyi* underwent extensive Nigon element recombination. Notably, these recombination events were highly correlated with their colinearity of protein sequences. Furthermore, transcriptomic analysis across five developmental stages of *D. destructor* identified numerous stage-specific effectors and transcription factors. RNAi experiments demonstrated that these genes play crucial roles in embryonic development and parasitic capability. Overall, this study not only provides valuable genomic and transcriptomic resources for PPNs but also uncovers key aspects of their genome evolution and parasitic gene functions. By elucidating the genetic basis of their evolutionary transitions and host adaptation, our findings lay a crucial foundation for advancing the understanding of PPN biology and their impact on agricultural systems.

Introduction

Plant-parasitic nematodes pose a significant threat to global agriculture, causing substantial crop losses and economic burdens[1]. These different types of plant parasitic nematodes can damage the roots, stems, leaves, flowers and fruits of plants, resulting in reduced plant production, a decrease in product quality and yield, as well as co-infection with other pathogens and ultimately plant death[2]. Among them, *Ditylenchus destructor* is a migratory endoparasitic nematode that infects a wide range of host plants, including sweet potatoes and other economically important crops. Unlike sedentary nematodes, *D. destructor* does not induce and maintain permanent feeding sites and is infectious from larvae to adults[3]. Understanding the genomic and molecular basis of its parasitism is crucial for developing effective nematode management strategies.

The effector of a pathogen is an important weapon for it to infect its host[4]. Previous studies on *D. destructor* have found many effectors that are closely related to

parasitism. For example, RNAi of alpha-amylase can reduce the parasitic ability of nematodes by 76.6%[5]; The allergen-like proteins can promote the infection of *D. destructor* nematodes[6]; Dd-cpl-1 RNAi not only leads to a significant decrease in the nematode's ability to infect, but also causes delayed development of embryos and a decrease in the amount of eggs laid by female[7]. However, no studies have been reported on the function of *D. destructor* transcription factors to date. It is well known that transcription factors regulate the expression of numerous genes and serve as key cis-acting elements in biological processes. For instance, in fungi, the bZIP transcription factor MeaB plays a crucial regulatory role in multiple aspects of Aspergillus fumigatus, including virulence, nitrogen metabolism, biofilm formation, and cell wall integrity[8-10]. Therefore, through high quality genome annotation, exploring the role of these virulence genes and transcription factors in the process of nematode infection of the host will be a meaningful work for analyzing the mechanism of nematode parasitism of the host.

Chromosomal organization and transposable elements (TEs) play pivotal roles in shaping genome architecture, influencing gene expression, and driving genetic diversity[11]. However, the genomic evolution of migratory parasitic nematodes remains largely unexplored, necessitating a detailed investigation into their chromosomal dynamics and TE landscape. Despite the increasing availability of genomic data, the chromosomal evolution of plant-parasitic nematodes remains poorly understood. A key question in nematode genome evolution is how chromosomal rearrangements and transposable elements contribute to adaptation and parasitism. Previous studies have suggested that free-living nematodes, such as *Caenorhabditis elegans*, maintain relatively conserved chromosomal structures, whereas plant-parasitic nematodes exhibit extensive rearrangements[12]. However, the extent of these rearrangements, their impact on gene function, and their evolutionary drivers remain unclear. Furthermore, how these chromosomal changes correlate with parasitic adaptations in different nematode species requires further exploration. Complete genome assemblies are key to advancing studies of

chromosome evolution research.

In the early stage of our study, we assembled the genome of *D. destructor* through

second-generation sequencing[13], and on this basis, the related functional genes of *D.*

destructor were studied[5, 7]. At the same time, we established the transcriptome

landscape of 11 time points in *D. destructor* embryonic development[14].

Advancements in sequencing technologies have enabled the construction of

high-quality chromosome-level genomes, providing unprecedented insights into

nematode evolution and host adaptation. Among this, we utilized third-generation

sequencing and chromatin interaction sequencing (Hi-C) to assemble the unzipped

genome of the polyploid root-knot nematode, providing novel insights into nematode

genomics[12]. In this study, we present a chromosome-level genome assembly of *D.*

destructor and perform a comparative genomic analysis with other plant-parasitic

nematodes, including *B. xylophilus* and *A. besseyi*. Our long-term goal is to explore

fundamental questions regarding genome plasticity and adaptability in plant-parasitic

nematodes by investigating chromosome evolution, transposon activity, and the

functional roles of secreted proteins and transcription factors in nematode

development and parasitism. This research is a core component of all our programs.

Our study integrates transcriptomic and RNAi analyses and to provide novel insights

into the genetic mechanisms underlying nematode infection, highlighting potential

targets for nematode control strategies. This work contributes to a broader

understanding of the genomic and evolutionary basis of plant-parasitic nematode

adaptations.

Results

Chromosome-level genome assembly and annotation of *D. destructor*

The chromosomal evolution of plant-parasitic nematodes remains a relatively

understudied field, and the construction of chromosome-level genomes is crucial for

advancing research in this area. Firstly, we performed Illumina short-read sequencing

on the genomic DNA of *D. destructor* and assessed its ploidy using smudgeplot,

which indicated that *D. destructor* is diploid (Fig. S1A) with a haploid genome size of

109Mb (Fig. S1B). To further determine the chromosome number of *D. destructor*,

we conducted karyotype analysis, which revealed that *D. destructor* possesses 8

chromosomes (Fig. 1A), i.e., 2n = 8. We utilized 25 GB of PacBio Sequel I data to

assemble the genome of *D. destructor*, resulting in a genome size of 105 Mb, an N50

of 899 kb, and a total of 344 contigs. Following polishing with second-generation

sequencing data, we generated 40 GB of Hi-C sequencing data. Leveraging the Hi-C

data, we successfully anchored 99.92% of the contig sequence length onto four

chromosomes (Fig. 1B). Our assembled genome is approximately 30 Mb smaller than

the recently published version[15] but has similar BUSCO values (72.2% vs 73.8%).

Since both the larvae and adults of migratory parasitic nematodes have parasitic

abilities, current studies have used transcriptome data from mixed stage to annotate

their gene expression. In order to comprehensively study the changes in gene

expression of *D. destructor* during development, we used instar synchronization to

obtain 18 transcriptome data from six stages: egg, J2, J3J4, male, female, and mixed

stage. We identified a total of 20,494 protein coding genes, 563 tRNA, and 48 rRNA

in *D. destructor*'s genome. In addition, we annotated 93,855 transposon elements in

the genome, with a total length of 23.9 Mb, accounting for 15.9% of the genome size.

We then compared our assembled genome with the previously published one[15],

highlighting both similarities and differences. Our analysis revealed a high degree of

collinearity at the gene level between the two genomes (Fig. S2). Subsequently, we

compared the whole genome sequences of the two assemblies and confirmed their

high collinearity (Fig. 1C). Additionally, we found that the Dd-A1 genome was larger,

with each chromosome containing sequences absent from our assembly (Fig. 1C). In

addition, the comparison of the two genomes identified a total of 151,874 SNPs,

26,146 insertions, 23,368 deletions, 93 inversions, 60 translocations, and 42

copygains.

We analyzed various genomic features, including GC content, transposable element

(TE) abundance, gene density, and gene expression, and visualized them using

circular diagrams (Fig. 1D). To further investigate chromosome organization within

the nucleus, we utilized Hi-C data to reconstruct the *D. destructor* chromosome

structure and their spatial relationships (Fig. 1E). These findings may offer valuable

insights for future studies on gene expression differences and the interplay between

age transitions and chromatin dynamics.

Comparison of genomes among migratory parasitic nematodes

Comparison of chromosomes among plant-parasitic nematodes provides valuable

insights into their evolutionary trajectories. We conducted a comparative analysis of

chromosome sequences among *D. destructor*, *B. xylophilus* and *A. besseyi*, focusing

on their Nigon elements. To facilitate comparison with *Panagrolaimus*, a free-living

nematode more closely related to PPNs of the Tylenchomorpha clade[16], we

annotated the genome of *Panagrolaimus* sp. ES5. Nigon element analysis revealed

that its autosomes are primarily composed of single Nigon elements, whereas the X

chromosome represents a fusion of four major elements—NigonA, C, D, and X—

suggesting that it originated through fusion events involving two autosomes (Fig. 2A).

The Nigon originals of *D. destructor* and *A. besseyi* were fragmented (Fig. 2B, C),

suggesting that multiple chromosome divisions and fusions had occurred. To explore

the potential association between variations in Nigon elements and chromosomal

rearrangements, we performed a synteny analysis of protein sequences across the

chromosomes of these four nematode species. Syntenic blocks were identified only

when more than ten adjacent protein sequences exhibited similarity across species.

Our findings demonstrated a strong concordance between the syntenic blocks of

protein sequences and the relative positions of Nigon elements (Fig. 2D). For example,

the A Nigon element, which is primarily located on *B. xylophilus* chromosome 1[17],

is dispersed across four chromosomes in *D. destructor* but is concentrated on

chromosome 1 in *A. besseyi*. This pattern is consistent with the distribution of

syntenic blocks and their chromosomal locations. For each of the remaining

chromosomes, the positional relationships between Nigon elements and syntenic

blocks of protein sequences also exhibit a one-to-one correspondence (Fig. 2).

Synteny analysis between *Panagrolaimus* sp. ES5 and *D. destructor* also revealed a

high degree of concordance between Nigon element composition and patterns of

protein-coding gene collinearity (Fig. S3). Based on this evidence from both Nigon

and collinearity, we infer that chromosome 3 of *D. destructor* represents its X

chromosome. Our comparative analysis reveals that chromosomal rearrangements,

including fusions and fragmentations, have shaped the Nigon structures of PPNs, with

B. xylophilus retaining a more conserved architecture, while *D. destructor* and *A.*

besseyi exhibit extensive chromosomal reorganization.

Transposons elements (TEs) are important drivers of genome dynamics, promoting

genome rearrangement, chromosome structural changes, and the evolution of gene

regulatory networks[11]. To investigate the role of TEs in migratory plant-parasitic

nematodes genome evolution, we conducted a comparative analysis of TEs in these

three nematode species. First, we annotated the TE elements in their genomes,

revealing that, compared to *B. xylophilus*, *D. destructor* and *A. besseyi* possess a

greater diversity of TE families, particularly the LINE transposon family (Fig. S3A,

Table S1-3). By analyzing the age distribution of TEs using Kimura substitution rates,

we found that the peak age distribution of retrotransposons (LTR/LINE) in *D.*

destructor and *A. besseyi* is closer to 0 compared to *B. xylophilus* (Fig. S3B),

suggesting that these retrotransposons may have undergone recent active expansion.

Dynamic landscape of transcriptome at different stages of *D. destructor*

To study the differences in gene expression at different developmental stages of

migratory parasitic nematodes, we obtained transcriptome data of *D. destructor* at five

stages from egg to adult (Fig. 3A). We first performed differential gene expression

analysis between adjacent developmental stages and identified 8,702, 2,881, 4,527,

and 4,950 differentially expressed genes (DEGs) in egg_vs_J2, J2_vs_J3J4,

J3J4_vs_male, and J3J4_vs_female (Fig. S4), respectively. We conducted a functional

analysis of DEGs across developmental stages, providing a valuable resource for identifying key genes involved in plant-parasitic nematode development. The transition from the J3/J4 stage to adult is closely linked to nematode sex determination. Pfam enrichment analysis revealed that genes upregulated in J3/J4 compared to males were primarily associated with protein degradation, particularly ubiquitin ligases such as zf-RING_2, Skp1, Cullin_Nedd8, and CSN8_PSD8_EIF3K (Fig. S5A). Besides, many histone-related genes were enriched in this process, suggesting that sex determination of *D. destructor* may be associated with epigenetics. Moreover, these genes were significantly enriched in essential biological processes, including signal transduction, chromatin regulation, energy metabolism, transcription, DNA replication, and sperm motility (Fig. S5A). Interestingly, the DEGs downregulated in J3/J4 compared to males were primarily enriched in GPCRs, collagen, molting-related genes, and glycosyltransferase families (Fig. S5B). This may suggest that male nematodes enter a relatively stable mature stage, no longer undergoing intense developmental and metabolic remodeling. Instead, they focus more on reproduction while reducing investment in environmental signaling, molting, growth, and stress responses. Surprisingly, the DEGs upregulated or downregulated between J3/J4 and females showed similar functional enrichment patterns to the differential genes between males and females (Fig. S5 C, D). This suggests that J3/J4 larvae share similar genetic regulatory mechanisms during their development into both male and female nematodes. The J3/J4 larval stage appears to be a highly plastic stage, where developmental direction may be controlled by common regulatory networks, with sex determination being an additional fine-tuning of these mechanisms. Furthermore, the differentiation between males and females may rely on fewer key regulatory factors or be influenced by epigenetic regulation, as the DEGs in females did not show enrichment in histone-related genes.

Furthermore, we calculated the Transcripts Per Million (TPM) values of the transcriptomes at each period and performed cluster analysis on the expression levels at different periods. The results indicate that gene expression in *D. destructor*

undergoes highly dynamic changes during development (Fig. S6), reflecting finely

tuned regulatory mechanisms that adapt to the physiological demands of different

developmental stages. To gain deeper insights into gene regulation underlying growth,

development, and parasitic strategies, we performed a functional analysis of clustered

genes at each developmental stage. The results indicate that the functions of genes

enriched at each developmental stage largely correspond to stage-specific

characteristics (Fig. 3B). For example, genes highly expressed during the egg stage

are primarily enriched in embryonic development, tissue and organ development, and

neuronal differentiation. In contrast, genes highly expressed during the J2 to J3/J4

stages are enriched in muscle differentiation, feeding, and responses to environmental

stimuli. In the adult stage, enriched functions are related to reproduction, P granules,

and gametogenesis (Fig. 3B). These findings suggest that the gene expression patterns

of *D. destructor* at different developmental stages are closely linked to their specific

physiological functions, reflecting precise gene regulatory mechanisms during

development to accommodate the growth demands and survival strategies of each

stage.

Dynamic expression profile of effector, HGT and TF during development

We predicted genes containing secretory signal peptides and no transmembrane

domains as secreted proteins, including the nematode pathogenicity factor effector,

and annotated 1327 secreted proteins in total. On the other hand, we predicted 124

horizontally transferred genes (HGTs) in the *D. destructor* genome based on

heterologous feature analysis. We then analyzed the characteristics of secreted

proteins and HGTs in the *D. destructor* genome. In terms of chromosomal distribution,

secreted proteins are generally evenly distributed across the chromosomes, with a few

regions showing localized clustering (Fig. 4A). In contrast, HGTs exhibit distinct

hotspot distributions, particularly concentrated at the chromosomal termini (Fig. 4A).

The enrichment of HGTs at chromosome ends may be associated with the dynamic

rearrangement, high plasticity, and susceptibility of telomeric regions to foreign gene

integration. This distribution pattern may facilitate the retention and adaptive

evolution of HGT genes, making them more likely to be preserved by natural selection or further amplified. In addition, we found that 29 of the 124 HGT genes were secreted proteins. For example, all the pectate lyases of *D. destructor* were from HGT, and some of these HGT secreted proteins were concentrated on the chromosomes in hotspot (Fig. 4A).

Although *D. destructor* has the ability to infect at all stages except eggs, we found that most of its secretory proteins have the highest expression levels in the J2 larvae (Fig. 4B), suggesting that the nematode faces many challenges in the initial process of establishing infection. We further conducted functional enrichment analysis on these secretory proteins highly expressed in the J2 period and found that they were mainly enriched in functions such as sugar metabolism, protein degradation, lipid degradation, and immune response (Fig. 4C). These genes play a crucial role in the process of nematodes feeding on host nutrients and resisting immune defense responses, and are the key weapons for nematodes to first infect hosts and establish parasitism. The expression profile of HGT genes during nematode development closely resembles that of secretory proteins, with the majority exhibiting high expression in the J2 larval stage and relatively lower expression in the egg stage (Fig. S7A). In contrast, transcription factors (TFs) display the opposite pattern, with the highest expression observed in the egg stage, followed by the J2 stage (Fig. S7B). This suggests that HGT genes are more closely associated with nematode infection and parasitism, whereas TFs play a crucial regulatory role in embryonic development.

Secreted proteins and TFs play crucial roles in nematode parasitism
To determine whether these highly expressed secretory proteins and transcription factors (TFs) play a pivotal role in nematode development and host parasitism, we selected four secretory proteins and four TFs that are highly expressed during the J2 stage, along with one TFs that exhibit peak expression during the egg stage (Table S4). RNAi experiments were then conducted to investigate their biological functions (Fig. S8). We first conducted RNAi experiments on the transcription factor gene Dd_05507,

which contains a cold shock domain. The results showed that the hatching rate of eggs

treated with Dd_05507 dsRNA was significantly lower than that of the control group

(Fig. 5A). We then conducted RNAi experiments on secretory protein genes and

transcription factors that were highly expressed during the J2 stage. The results

revealed that, except for *Dd_09599* and *Dd_03680*, silencing these putative effector

genes significantly impaired the nematodes' ability to infect sweet potatoes (Fig. 5B,

C). The RNAi experiments on transcription factors further confirmed that these TFs,

belonging to the HMG_box, zf-C4, Homeodomain, and zf-LITAF-like families, are

highly expressed during the J2 stage and play a crucial role in facilitating J2

nematodes' successful colonization of sweet potato and establishment of parasitism.

(Fig. 5D, E). These findings highlight the essential role of secretory proteins and

transcription factors in nematode development and parasitism. The functional

validation of these genes provides key insights into the molecular mechanisms

underlying host infection, offering potential targets for disrupting nematode

parasitism and developing effective pest control strategies.

Disscusion

Recent studies on nematode evolution have shown that *B. xylophilus* is at the root of

the phylogenetic tree of plant-parasitic nematodes[16] and has the same chromosome

number as *C. elegans*, suggesting that *B. xylophilus* may not have undergone

chromosomal recombination, but rather has undergone processes such as inversion,

translocation, gene gain and loss, and mutation that enabled it to evolve the ability to

parasitize plants. The chromosome-level assembly of *D. destructor* provides a

valuable resource for studying the genome evolution of plant-parasitic nematodes.

Our comparative genomic analysis revealed significant chromosomal rearrangements

among *D. destructor*, *B. xylophilus*, and *A. besseyi*, with *B. xylophilus* maintaining a

more conserved Nigon structure similar to *C. elegans*, whereas *D. destructor* and *A.*

besseyi exhibit extensive chromosomal reorganization. These findings suggest that

structural variations, including fusions and fragmentations, have played a crucial role

in shaping the genome architecture of plant-parasitic nematodes. Transposons

elements play an extremely important role in the evolution of many organisms[18-21].

Moreover, TEs can not only regulate genome plasticity, but also act as mediators of

genetic variation in plant pathogens, affecting pathogenicity and adaptive

evolution[22]. Our transposon analysis indicated that *D. destructor* and *A. besseyi*

have undergone recent expansions of retrotransposons, particularly within the LINE

family, which may have contributed to genome plasticity and adaptive evolution.

Through years of experience in plant-parasitic nematodes, we have explored a method

to synchronize the stages of *D. destructor* and are able to collect nematodes of

different stages on a large scale. This work has laid the foundation for detailed studies

of the gene function and expression profile of *D. destructor*. The dynamic

transcriptomic landscape across developmental stages of *D. destructor* provided

insights into the regulatory mechanisms underlying nematode growth and parasitism.

Our differential expression analysis revealed stage-specific gene expression patterns,

with embryogenesis characterized by genes involved in tissue development and

neuronal differentiation, while J2 larvae exhibited upregulation of genes associated

with feeding, environmental responses, and host infection. The upregulation of

ubiquitin ligases and histone-related genes in J3/J4 suggests potential epigenetic

regulation of sex determination, highlighting a complex interplay between chromatin

modifications and nematode development.

We identified and characterized secreted proteins and horizontally transferred genes

(HGTs) in *D. destructor*, revealing their critical roles in nematode infection and

adaptation. Secreted proteins were found to be highly expressed in J2 larvae,

underscoring their importance in overcoming host defenses and establishing

parasitism. The chromosomal localization of HGTs at telomeric regions suggests their

potential involvement in genome plasticity and adaptive evolution[12]. Notably,

RNAi-mediated silencing of several highly expressed secreted proteins and

significantly impaired nematode infection, confirming their essential roles in

parasitism. Additionally, RNAi experiments on transcription factors demonstrated

their crucial involvement in J2 colonization and host establishment, providing evidence for the regulatory mechanisms governing nematode development and parasitism. Subsequent work can further explore the mechanism of action between these important effectors and plants and reveal their specific pathogenic processes. At the same time, our results show that TF play a more significant role in nematode colonization. Some transcription factors may regulate the expression of a class of effector proteins or participate in pathways that inhibit plant immunity. Studying the specific regulatory networks of these transcription factors will provide new research ideas for the interaction between PPNs and their hosts.

Our findings provide a comprehensive view of the genetic and regulatory factors that drive nematode parasitism, with implications for developing novel control strategies. The identification of key effectors and transcription factors offers potential targets for disrupting nematode-host interactions, paving the way for more effective pest management approaches.

Materials and methods

***D. destructor* nematode culture and collection**

Sterilize the surface of sweet potatoes using alcohol and UV in a clean bench for 15 minutes. Punch holes or cut triangular incisions in the sweet potatoes, with an average of two holes per potato, for nematode inoculation. Introduce the counted stem nematodes into the incisions or holes, inoculating 2,000 nematodes per hole. Each sweet potato receives two holes, totaling 4,000 nematodes per potato. Incubate at 25°C with 40%-60% humidity for 30-40 days. Nematodes can then be collected in large quantities using the shallow dish method.

Instar synchronization of *D. destructor* nematodes

1. **Sample Collection and Processing.** Sweet potatoes inoculated for 30–35 days are collected. The tubers are crushed using a homogenizer at 20% power, and the resulting mixture of nematodes and sweet potato slurry is transferred into a beaker.

2. **Filtration through Sieves.** The mixture is sequentially filtered through a series of standard sieves with mesh sizes of 10, 20, 40, 60, 100, and 200 (determined based on the size of nematodes and eggs), yielding a fine sweet potato particulate fraction mixed with nematodes and eggs.

3. **Density Gradient Centrifugation.** The obtained mixture is aliquoted and subjected to density gradient centrifugation using a 30% sucrose solution to rapidly separate nematodes, eggs, and other impurities. The supernatant, containing nematodes and eggs, is collected and further purified using a 3 µm membrane filter to remove bacteria and debris.

4. **Nematode Lysis and Egg Purification.** The nematode and egg suspension is treated with nematode lysis solution consisting of sodium hypochlorite (50%): sodium hydroxide: water at a ratio of 1:2:7 for 5 minutes, killing the nematodes while preserving viable eggs. Extending the lysis time to 45 minutes ensures complete degradation of nematode bodies, yielding purified, viable eggs.

5. **Egg Hatching and J2 Larval Collection.** The purified eggs are incubated in a 2000-mesh sieve device at 25°C for 72 hours, allowing synchronized hatching of second-stage juveniles (J2).

6. **Isolation of J3 and J4 Juveniles.** The hatched J2 nematodes are inoculated onto Botrytis cinerea plates and incubated at 25°C for 12 days. The nematodes are then collected, and adult nematodes (both male and female) are manually removed. The remaining nematodes consist of third-stage (J3) and fourth-stage (J4) juveniles.

7. **Isolation of male and female.** The female and male insects manually picked in step 6 were saved separately for transcriptome sequencing at the corresponding period.

DNA, RNA extraction and library construction

DNA samples used for PacBio and Illumina sequencing were obtained from egg stage of *D. destructor*. DNA extraction followed the CTAB method. First, the collected nematodes were centrifuged in a 35% sucrose solution to remove impurities, then suspended in 200 µL of lysis buffer. The nematodes, along with the buffer, were

transferred into a mortar containing liquid nitrogen. Once frozen, the nematodes were ground into a fine powder. The ground nematodes were transferred into a 1.5 mL centrifuge tube, followed by the addition of another 300 µL of lysis buffer and 500 µL of preheated (65°C) CTAB. The mixture was thoroughly mixed and incubated at 65°C for 30 minutes, with occasional inversion to ensure proper mixing. Proteins were removed using phenol, RNA was eliminated using RNase A, and DNA was precipitated with isopropanol. DNA intended for PacBio sequencing underwent an additional purification step using magnetic beads to further remove polysaccharide and protein contaminants. For RNA extraction, we first obtained nematodes at different developmental stages using the nematode synchronization method, collecting three biological replicates for each stage. RNA was then extracted using the TransZol Up Plus RNA Kit (TransGen Biotech, Beijinng, China), followed by library construction with the VAHTS Universal V10 RNA-seq Library Prep Kit for Illumina (NR606-01). Finally, RNA sequencing was performed on the Nova platform by NOVOGENE.

Hi-C library construction

The Hi-C library was constructed using *D. destructor* eggs. The construction method followed the previously established Hi-C library preparation protocol for root-knot nematodes[12]. Briefly, freshly collected eggs were crosslinked with 1.5% formaldehyde for 30 minutes, and the crosslinking was quenched with glycine. The eggs were then ground in a mortar to isolate nuclei. The chromatin was digested with the MboI restriction enzyme, followed by biotin labeling and ligation with T4 DNA ligase. The ligated DNA was then sheared by sonication, and biotin-labeled fragments were enriched using streptavidin magnetic beads. Library preparation was performed using the Vazyme ND606 kit, and the final Hi-C library was sequenced on the Illumina NovaSeq platform with 150 bp paired-end reads.

Chromosome-level genome assembly

For genome assembly, our workflow was as follows: First, we used Illumina

paired-end sequencing data to assess the ploidy of the Dd genome using

Smudgeplot[23]. After determining that the genome is diploid, we estimated the

genome size using GenomeScope2[23] with a k-mer size of 21, assuming a diploid

model. Following genome evaluation, we performed error correction on PacBio

Sequel I data using Canu$_{v1.9}$ [24], setting genome coverage at 40×. The corrected 40×

data was then assembled using Smartdenovo[25]. The resulting contigs were first

polished with gcpp (https://github.com/PacificBiosciences/gcpp) using PacBio data,

followed by multiple rounds of polishing with Pilon[26] using Illumina data. Next, we

used Juicer (v1.5.7) pipline[27] and 3D-DNA (v180419) pipline[28] to anchor the

polished genome to chromosomes based on Hi-C signals. Finally, the anchored

genome was manually refined in Juicebox (v1.13.01) [29], yielding the final

chromosome-level genome assembly. The three-dimensional structure of the genome

was reconstructed using miniMDS[30] software.

Genome annotation

We used transcriptome data from all developmental stages to assist in annotating the

genome of *Ditylenchus destructor*. First, we employed Trinity (v2.15.2)[31] to

perform de novo assembly of the transcriptome data, resulting in the "Trinity.fasta"

file. The assembled sequences were then used in PASA[32] for ORF prediction,

generating the "pasa_assemblies.gff3" file. Next, we masked the Dd genome using

RepeatMasker[33], followed by alignment of the transcriptome data to the masked

genome with HISAT2[34]. The resulting BAM files were used as input for

BRAKER3[35] to predict genes, generating the "braker.gff3" file. Additionally, the

BAM files from HISAT2 alignment were used for gene prediction with TransDecoder

and StringTie (v2.1.1)[36], yielding the "stringtie.gff3" file. Finally, we merged the

gene predictions from the three methods using EVidenceModeler, incorporating

weight-based merging to produce the final annotation file, "annotation.gff3". CDS

and protein sequences were extracted from this GFF3 file using GFFRead. The

genome annotation of *Panagrolaimus* sp. ES5 was performed by using the same

strategy, the genome and RNA-seq data were download from NCBI with accession
number GCA_964243005.1 and SRR5253562.

TE, HGT, TF, and secretory protein prediction

For TE prediction, we first used EDTA (v2.2.0) pipline[37] to predict the genome

with the parameters "--overwrite 1 --force 1 --sensitive 1 --anno 1 --evaluate 1" and

obtained "TEanno.gff3" and "TElib.fa". Then, using "TElib.fa" as a library file,

RepeatMasker[33] was used to re-predict TE in the genome, and

calcDivergenceFromAlign.pl was used to calculate Kimura substitution on the result

file "*.align". For HGT prediction, we use AvP pipline[38] with uniref90 database, the

candidate gene with HGT or HGT-NT lable was considered as HGT genes. For

transcription factors, we divided all protein sequences of *D. destructor* into two

groups of no more than 9000 genes and predict the TF by using AnimalTFDB v4.0

(https://guolab.wchscu.cn/AnimalTFDB4/#/TF_Predict). For secretory protein

prediction, we first used singalP(v5.0)[39] to predict genes with secretory signal

peptides, and then used tmhmm[40] to predict whether these genes had

transmembrane domains. After eliminating genes containing transmembrane domains,

they were potential secretory proteins.

Collinearity and genomic variation analysis

We used minimap2 with the parameters "-ax asm5 --eqx" to align the entire genome

between the *D. destructor* genome of this study and Dd-A1[15], and then used

Syri[41] to identify collinear regions, structural variations, insertion and deletion

mutations, and SNP information. The syntenic protein pairs between the *D. destructor*

genome of this study and Dd-A1 were identified by using JCVI pipeline[42].

RNA-seq analysis and expression profile determination

For the analysis of differentially expressed genes, we first aligned the transcriptome

data to the genome using HISAT2, followed by sorting with SAMtools[43] and

quantifying read counts with HTSeq[44]. Differential gene expression analysis was

then performed using DESeq2[45]. For TPM calculation, we first aligned the

transcriptome data to the genome using STAR[46] and then used RSEM[47] to calculate TPM values for each gene. Gene expression profile clustering across developmental stages was conducted using Mfuzz[48], while the expression pattern clustering of transcription factors, effectors, and other key genes at different stages was performed using pheatmap.

In vitro RNAi

Primers were designed for target genes, each incorporating a T7 promoter sequence. Then, the target genes were amplified from *D. destructor* cDNA. The PCR products were purified, ensuring DNA concentrations exceeded 100 ng/µl. Vazyme T7 RNAi Transcription Kit (TR102) was used for dsRNA synthesis. The final dsRNA concentration was adjusted to 1000 ng/µl. For the RNAi interference system preparation, the total reaction volume was 300 µl, supplemented with 15 µl of 1 mol/L octopamine and 0.5 µL of 1 mol/L spermidine. For the nematode and egg RNAi treatment, we use 6000 nematodes or 5000 eggs were for each gene-specific RNAi system. The interference reaction was conducted at 37°C for 48 hours. After RNAi treatment, 2000 nematodes were collected for RNA extraction and then used for qRT-PCR, the remaining nematodes were used for phenotypic analysis.

Assessment of nematode infectivity

Infection assay: Sweet potatoes of uniform diameter were punctured to a standardized depth (about 1cm). Each hole was inoculated with 500 nematodes, with 8 replicates per gene. After 30 days infection at 25°C of dark incubator, nematodes were collected and counted. The extent of infection was examined, and photographs were taken for documentation.

Egg hatching rate assay: 30–40 early-stage embryos were selected per treatment, with 5 replicates per gene. The eggs were incubated at 25°C for 3–7 days, hatching rates were recorded every 24 hours.

Nematode quantification: Harvesting nematodes from infected sweet potatoes, a 2 cm region surrounding the infection site was excised. The infected tissue was cut into 0.5

cm × 0.5 cm pieces and wrapped in a single layer of paper towel. The samples were submerged in water for 4 hours to allow nematodes to migrate into the liquid. Nematode suspensions were collected and counted at least three times to estimate the total population.

Negative Control Group: dsGFP synthesized using GFP gene as a template was used as a negative control for RNAi. The same infection ability and egg hatching rate assays were performed for comparison.

Data availability

All sequencing data have been deposited into the National Center for Biotechnology Information (NCBI) Sequence Read Archive. The different stages of RNA sequencing data of *D. destructor* are deposited in BioProject PRJNA1164248 (including 15 SRA); The PacBio Sequel I data, Hi-C data, and Illumina data of *D. destructor* are deposited in BioProject PRJNA1236705 (including 3 SRA). The genome of *B. xylophilus* and *A. besseyi* were download form wormbase parasite with accession number PRJEB40022 and PRJNA834627. The genome of *Panagrolaimus* sp. ES5 was download from GenBank with accession number GCA_964243005.1, and the RNA-seq data was download from SRA database with accession number SRR5253562. The assembled *D. destructor* genome and annotation files were uploaded to https://bmb.hzau.edu.cn/sjxz.htm.

Declaration of Competing Interest

The authors declare no conflicts of interest.

Acknowledgements

We thank Romnick Latina of University of California Davis for advice on nematodes evolution and the UC Davis farm cluster server for help. This research was supported by the National Natural Science Foundation of China (32271546, U20A2040 and 31970076) and Hubei Hongshan Laboratory (2022hszd012).

Author Contributions

D.D. and M.S. conceived and administration the project. M.S., acquisition the funding.

M.S. and D.P. manage this project. D.D. designed the experiments. Y.C. collected the

data and performed the RNAi experiment. S. Z. clone the gene. S. Z. and X.W.

constructed the RNA-seq libraries. D.D. and Y.Z. performed the data analysis. B.H.,

D.B., Y.L., N.A., and Y.L.Z. raised nematodes. D.D. wrote the manuscript. All authors

reviewed the manuscript.

References

1. Jones JT, Haegeman A, Danchin EG, Gaur HS, Helder J, Jones MG, Kikuchi T, Manzanilla-Lopez R, Palomares-Rius JE, Wesemael WM et al: **Top 10 plant-parasitic nematodes in molecular plant pathology**. *Mol Plant Pathol* 2013, **14**(9):946-961.

2. Parrado LM, Quintanilla M: **Plant-parasitic nematode disease complexes as overlooked challenges to crop production**. *Front Plant Sci* 2024, **15**:1439951.

3. Lu Y, Yang S, Chen W, Xie H, Xu C: **Advances in Migratory Plant Endoparasitic Nematode Effectors**. *Int J Mol Sci* 2024, **25**(12).

4. Lanver D, Tollot M, Schweizer G, Lo Presti L, Reissmann S, Ma LS, Schuster M, Tanaka S, Liang L, Ludwig N et al: **Ustilago maydis effectors and their impact on virulence**. *Nat Rev Microbiol* 2017, **15**(7):409-421.

5. Chen L, Xu M, Wang C, Zheng J, Huang G, Chen F, Peng D, Sun M: **Multi-copy alpha-amylase genes are crucial for Ditylenchus destructor to parasitize the plant host**. *PLoS One* 2020, **15**(10):e0240805.

6. Chang Q, Yang Y, Hong B, Zhao Y, Zhao M, Han S, Zhang F, Peng H, Peng D, Li Y: **A variant of the venom allergen-like protein, Dd VAP2, is required for the migratory endoparasitic plant nematode Ditylenchus destructor parasitism of plants**. *Frontiers in Plant Science* 2023, **14**:1322902.

7. Huang G, Cong Z, Liu Z, Chen F, Bravo A, Soberon M, Zheng J, Peng D, Sun M: **Silencing Ditylenchus destructor cathepsin L-like cysteine protease has negative pleiotropic effect on nematode ontogenesis**. *Sci Rep* 2024, **14**(1):10030.

8. Wagner D, Schmeinck A, Mos M, Morozov IY, Caddick MX, Tudzynski B: **The bZIP transcription factor MeaB mediates nitrogen metabolite repression at specific loci**. *Eukaryotic Cell* 2010, **9**(10):1588-1601.

9. Perez-Cuesta U, Guruceaga X, Cendon-Sanchez S, Pelegri-Martinez E, Hernando FL, Ramirez-Garcia A, Abad-Diaz-de-Cerio A, Rementeria A: **Nitrogen, iron, and zinc acquisition: key nutrients to Aspergillus fumigatus virulence**. *Journal of Fungi* 2021, **7**(7):518.

10. Chen Y, Gao F, Chen X, Tao S, Chen P, Lin W: **The basic leucine zipper transcription factor MeaB is critical for biofilm formation, cell wall integrity, and virulence in Aspergillus fumigatus**. *Msphere* 2024, **9**(2):e00619-00623.

11. Slotkin RK, Martienssen R: **Transposable elements and the epigenetic regulation of the genome**. *Nat Rev Genet* 2007, **8**(4):272-285.

12. Dai D, Xie C, Zhou Y, Bo D, Zhang S, Mao S, Liao Y, Cui S, Zhu Z, Wang X et al: **Unzipped chromosome-level genomes reveal allopolyploid nematode origin pattern as unreduced gamete hybridization**. *Nat Commun* 2023, **14**(1):7156.

13. Zheng J, Peng D, Chen L, Liu H, Chen F, Xu M, Ju S, Ruan L, Sun M: **The Ditylenchus destructor genome provides new insights into the evolution of plant parasitic nematodes**. *Proc Biol Sci* 2016, **283**(1835).

14. Wang X, Guo Z, Dai D, Xie C, Zhao Z, Zheng J, Sun M, Peng D: **High-resolution transcriptome datasets during embryogenesis of plant-parasitic nematodes**. *Sci Data* 2024, **11**(1):690.

15. Yang Y, Feng R, Hong B, Fang Y, Liu C, Wang K, Peng D, Li Y, Peng H, Chang Q: **Chromosome-level genome assembly of the sweet potato rot nematode Ditylenchus destructor**. *Sci Data* 2025, **12**(1):174.

16. Qing X, Zhang YM, Sun S, Ahmed M, Lo WS, Bert W, Holovachov O, Li H: **Phylogenomic**
Insights into the Evolution and Origin of Nematoda. *Syst Biol* 2024.
17. Gonzalez de la Rosa PM, Thomson M, Trivedi U, Tracey A, Tandonnet S, Blaxter M: **A**
telomere-to-telomere assembly of Oscheius tipulae and the evolution of rhabditid
nematode chromosomes. *G3 (Bethesda)* 2021, **11**(1).
18. Wicker T, Gundlach H, Spannagl M, Uauy C, Borrill P, Ramirez-Gonzalez RH, De Oliveira R,
International Wheat Genome Sequencing C, Mayer KFX, Paux E *et al*: **Impact of transposable**
elements on genome structure and evolution in bread wheat. *Genome Biol* 2018, **19**(1):103.
19. Kumar CS, Qureshi SF, Ali A, Satyanarayana ML, Rangaraju A, Venkateshwari A, Nallari P:
Hidden magicians of genome evolution. *Indian J Med Res* 2013, **137**(6):1052-1060.
20. Yu Z, Li J, Wang H, Ping B, Li X, Liu Z, Guo B, Yu Q, Zou Y, Sun Y *et al*: **Transposable elements in**
Rosaceae: insights into genome evolution, expression dynamics, and syntenic gene
regulation. *Hortic Res* 2024, **11**(6):uhae118.
21. Shao F, Han M, Peng Z: **Evolution and diversity of transposable elements in fish genomes**.
Sci Rep 2019, **9**(1):15399.
22. Mat Razali N, Cheah BH, Nadarajah K: **Transposable Elements Adaptive Role in Genome**
Plasticity, Pathogenicity and Evolution in Fungal Phytopathogens. *Int J Mol Sci* 2019, **20**(14).
23. Ranallo-Benavidez TR, Jaron KS, Schatz MC: **GenomeScope 2.0 and Smudgeplot for**
reference-free profiling of polyploid genomes. *Nat Commun* 2020, **11**(1):1432.
24. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM: **Canu: scalable and**
accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome*
Res 2017, **27**(5):722-736.
25. Liu H, Wu S, Li A, Ruan J: **SMARTdenovo: a de novo assembler using long noisy reads**.
GigaByte 2021, **2021**:gigabyte15.
26. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q,
Wortman J, Young SK *et al*: **Pilon: an integrated tool for comprehensive microbial variant**
detection and genome assembly improvement. *PLoS One* 2014, **9**(11):e112963.
27. Durand NC, Shamim MS, Machol I, Rao SS, Huntley MH, Lander ES, Aiden EL: **Juicer Provides**
a One-Click System for Analyzing Loop-Resolution Hi-C Experiments. *Cell Syst* 2016,
3(1):95-98.
28. Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, Shamim MS, Machol I,
Lander ES, Aiden AP *et al*: **De novo assembly of the Aedes aegypti genome using Hi-C yields**
chromosome-length scaffolds. *Science* 2017, **356**(6333):92-95.
29. Durand NC, Robinson JT, Shamim MS, Machol I, Mesirov JP, Lander ES, Aiden EL: **Juicebox**
Provides a Visualization System for Hi-C Contact Maps with Unlimited Zoom. *Cell Syst* 2016,
3(1):99-101.
30. Rieber L, Mahony S: **miniMDS: 3D structural inference from high-resolution Hi-C data**.
Bioinformatics 2017, **33**(14):i261-i266.
31. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,
Raychowdhury R, Zeng Q: **Trinity: reconstructing a full-length transcriptome without a**
genome from RNA-Seq data. *Nature biotechnology* 2011, **29**(7):644.
32. Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, White O, Buell CR, Wortman JR:
Automated eukaryotic gene structure annotation using EVidenceModeler and the Program
to Assemble Spliced Alignments. *Genome Biol* 2008, **9**(1):R7.

33.   Tarailo-Graovac M, Chen N: **Using RepeatMasker to identify repetitive elements in genomic**
sequences. *Curr Protoc Bioinformatics* 2009, **Chapter 4**:4 10 11-14 10 14.

34.   Kim D, Paggi JM, Park C, Bennett C, Salzberg SL: **Graph-based genome alignment and**
genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 2019, **37**(8):907-915.

35.   Gabriel L, Bruna T, Hoff KJ, Ebel M, Lomsadze A, Borodovsky M, Stanke M: **BRAKER3: Fully**
automated genome annotation using RNA-seq and protein evidence with GeneMark-ETP,
AUGUSTUS, and TSEBRA. *Genome Res* 2024, **34**(5):769-777.

36.   Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL: **StringTie enables**
improved reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol* 2015,
33(3):290-295.

37.   Ou S, Su W, Liao Y, Chougule K, Agda JRA, Hellinga AJ, Lugo CSB, Elliott TA, Ware D, Peterson T
et al: **Benchmarking transposable element annotation methods for creation of a**
streamlined, comprehensive pipeline. *Genome Biol* 2019, **20**(1):275.

38.   Koutsovoulos GD, Granjeon Noriot S, Bailly-Bechet M, Danchin EGJ, Rancurel C: **AvP: A**
software package for automatic phylogenetic detection of candidate horizontal gene
transfers. *PLoS Comput Biol* 2022, **18**(11):e1010686.

39.   Almagro Armenteros JJ, Tsirigos KD, Sonderby CK, Petersen TN, Winther O, Brunak S, von
Heijne G, Nielsen H: **SignalP 5.0 improves signal peptide predictions using deep neural**
networks. *Nat Biotechnol* 2019, **37**(4):420-423.

40.   Krogh A, Larsson B, von Heijne G, Sonnhammer EL: **Predicting transmembrane protein**
topology with a hidden Markov model: application to complete genomes. *J Mol Biol* 2001,
305(3):567-580.

41.   Goel M, Sun H, Jiao WB, Schneeberger K: **SyRI: finding genomic rearrangements and local**
sequence differences from whole-genome assemblies. *Genome Biol* 2019, **20**(1):277.

42.   Wang Y, Tang H, Debarry JD, Tan X, Li J, Wang X, Lee TH, Jin H, Marler B, Guo H *et al*: **MCScanX:**
a toolkit for detection and evolutionary analysis of gene synteny and collinearity. *Nucleic*
Acids Res 2012, **40**(7):e49.

43.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
Genome Project Data Processing S: **The Sequence Alignment/Map format and SAMtools**.
Bioinformatics 2009, **25**(16):2078-2079.

44.   Anders S, Pyl PT, Huber W: **HTSeq--a Python framework to work with high-throughput**
sequencing data. *Bioinformatics* 2015, **31**(2):166-169.

45.   Love MI, Huber W, Anders S: **Moderated estimation of fold change and dispersion for**
RNA-seq data with DESeq2. *Genome Biol* 2014, **15**(12):550.

46.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR:
STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 2013, **29**(1):15-21.

47.   Li B, Dewey CN: **RSEM: accurate transcript quantification from RNA-Seq data with or**
without a reference genome. *BMC Bioinformatics* 2011, **12**:323.

48.   Kumar L, M EF: **Mfuzz: a software package for soft clustering of microarray data**.
Bioinformation 2007, **2**(1):5-7.

Figures and Legends

Figure 1 The genome landscape of *D. destructor*. (A) The karyotype photo of *D.*
destructor shows that it has 4 pairs of chromosomes. (B) Chromatin interaction
signals mount the *D. destructor* genome onto four chromosomes (haploid). (C)
Syntenic analysis of *D. destructor* genome between this study and the published
Dd-A1 genome (ref14). (D) Circos plot of *D. destructor* haploid genomic features. (E)
Reconstructed 3D chromatin structure within the nucleus of *D. destructor* at the
mixed developmental stage based on Hi-C data.

Figure 2 Comparative genomic features of three migratory plant-parasitic nematodes. (A) The distribution characteristics of Nigon elements on the Panagrolaimus sp. ES5 genome shows that each element is generally concentrated on one autosome, while Nigon A, C, D, and X fused the X chromosome. (B) The distribution characteristics of Nigon elements on the *D. destructors* genome. (C) The distribution characteristics of Nigon elements on the *A. besseyi* genome. (D) Synteny of protein sequences among three migratory plant-parasitic nematodes. The results showed that the collinearity of protein sequences was highly consistent with the distribution characteristics of Nigon elements.

727
Figure 3 Gene expression profile characteristics and functional analysis of _D._
destructor nematodes at different stages. The heat map on the left shows the
clustering of highly expressed genes in each stage after soft clustering of
transcriptome data of _D. destructor_ at five different developmental stages. The right
shows the GO functional enrichment results of genes highly expressed at each
developmental stages. The length of the bar plot indicates the number of genes, and
the color indicates the p-value of the enrichment degree.

Figure 4 Distribution characteristics of secretory proteins and HGT on the *D. destructor* genome and their expression characteristics among different developmental stages. (A) The distribution characteristics of secretory proteins and HGT on the *D. destructor* genome showed that HGT was distributed in a hotspot pattern, especially concentrated at the chromosome ends. (B) The expression of putative effector (secretory proteins) genes at different developmental ages was clustered, and most of the effectors were highly expressed in the J2 stage. (C) The functional enrichment of putative effectors that were highly expressed in the J2 period was performed through the pfam database. The results showed that a variety of functional genes highly related to nematode feeding, such as those involved in sugar metabolism, protein degradation, and lipid degradation, were enriched.

750
Figure 5 The influence of effector and TF genes in *D. destructor* on its development and infection ability. (A) The hatching rate of eggs was significantly reduced after transcription factor Dd_05507 RNAi. (B) Phenotypes of sweet potato infected with *D. destructor* after the putative effector gene RNAi. (C) Statistics of *D. destructor* reproductive ability after RNAi of putative effector genes. (D) Phenotypes of sweet potato infected with *D. destructor* after the putative TF genes RNAi. (E)

757   Statistics of *D. destructor* reproductive ability after RNAi of putative TF genes. The
758   number of replicates for each treatment in Figures C and E was 4-7, and the
759   significance was analyzed using t-test, ns means no significance, ** means p < 0.01,
760   *** means p < 0.001.
761
762

Figures and Legends

Figure 1 The genome landscape of *D. destructor*. (A) The karyotype photo of *D. destructor* shows that it has 4 pairs of chromosomes. (B) Chromatin interaction signals mount the *D. destructor* genome onto four chromosomes (haploid). (C) Syntenic analysis of *D. destructor* genome between this study and the published Dd-A1 genome (ref14). (D) Circos plot of *D. destructor* haploid genomic features. (E) Reconstructed 3D chromatin structure within the nucleus of *D. destructor* at the mixed developmental stage based on Hi-C data.

Figure 2 Comparative genomic features of three migratory plant-parasitic nematodes. (A) The distribution characteristics of Nigon elements on the *Panagrolaimus* sp. ES5 genome show that each element is generally concentrated on one autosome. (B) The distribution characteristics of Nigon elements on the *D. destructors* genome. (C) The distribution characteristics of Nigon elements on the *A. besseyi* genome. (D) Synteny of protein sequences among three migratory plant-parasitic nematodes. The results showed that the collinearity of protein sequences was highly consistent with the distribution characteristics of Nigon elements.

Figure 3 Gene expression profile characteristics and functional analysis of *D. destructor* nematodes at different stages. The heat map on the left shows the clustering of highly expressed genes in each stage after soft clustering of transcriptome data of *D. destructor* at five different developmental stages. The right shows the GO functional enrichment results of genes highly expressed at each developmental stages. The length of the bar plot indicates the number of genes, and the color indicates the p-value of the enrichment degree.

Figure 4 Distribution characteristics of secretory proteins and HGT on the *D. destructor* genome and their expression characteristics among different developmental stages. (A) The distribution characteristics of secretory proteins and HGT on the *D. destructor* genome showed that HGT was distributed in a hotspot pattern, especially concentrated at the chromosome ends. (B) The expression of putative effector (secretory proteins) genes at different developmental ages was clustered, and most of the effectors were highly expressed in the J2 stage. (C) The functional enrichment of putative effectors that were highly expressed in the J2 period was performed through the pfam database. The results showed that a variety of functional genes highly related to nematode feeding, such as those involved in sugar metabolism, protein degradation, and lipid degradation, were enriched.

Figure 5 The influence of effector and TF genes in *D. destructor* on its development and infection ability. (A) The hatching rate of eggs was significantly reduced after transcription factor Dd_05507 RNAi. (B) Phenotypes of sweet potato infected with *D. destructor* after the putative effector gene RNAi. (C) Statistics of *D. destructor* reproductive ability after RNAi of putative effector genes. (D) Phenotypes of sweet potato infected with *D. destructor* after the putative TF genes RNAi. (E)

Statistics of *D. destructor* reproductive ability after RNAi of putative TF genes. The number of replicates for each treatment in Figures C and E was 4-7, and the significance was analyzed using t-test, ns means no significance, ** means $p < 0.01$, *** means $p < 0.001$.