3 An optimized rabies vaccine vehicle for

4 orotopical administration to wild vampire bats

Authors

Cole Knuese[1], Elsa M. Cárdenas-Canales[1], Travis McDevitt-Galles[2], María Magdalena

Ramírez Martínez[3], Daniel Limonta[1], Lisa E. Powers[2], Daniel P. Walsh[4], Daniel G.

Streicker[5,6], Jorge E. Osorio[1,7], Mostafa Zamanian[1], Tonie E. Rocke[2*]

Affiliations

[1] School of Veterinary Medicine, University of Wisconsin, Madison, Wisconsin, USA.

[2] US Geological Survey National Wildlife Health Center, Madison, Wisconsin, USA.

[3] Departamento de Ciencias de la Salud y Ecología Humana, Centro Universitario de la

Costa Sur, Universidad de Guadalajara, Autlán, Jalisco, México.

[4] U.S. Geological Survey, Montana Cooperative Wildlife Research Unit, Wildlife Biology

Program, University of Montana, Missoula, Montana, 59812, USA.

[5] School of Biodiversity, One Health and Veterinary Medicine, University of Glasgow,

Glasgow, UK.

[6] MRC-University of Glasgow Centre for Virus Research, Glasgow, UK.

[7] Global Health Institute, University of Wisconsin, Madison, Wisconsin, USA.

* trocke@usgs.gov

Abstract

Rabies vaccination of vampire bats (*Desmodus rotundus*) has been proposed as a superior control method to culling but has yet to be implemented. Success of rabies vaccination depends on a topical vehicle that spreads through a bat colony via allogrooming while additionally preserving vaccine immunogenicity. This work describes the *in vitro* and *in vivo* optimization of a new orotopical gel for rabies vaccine delivery to vampire bats. Autonomous transferability of our carboxymethyl cellulose (CMC) gel vaccine delivery formulation was tested in a microchipped vampire bat colony in rural Jalisco, Mexico. Intra-colony gel uptake was traced using the fluorescent biomarker rhodamine B. Importantly, application of topical treatment of ~20% of the bat colony resulted in estimated uptake by over 85% of the colony. The *in vitro* stability of our raccoon poxviral-vectored rabies vaccine candidate within CMC was measured at time points up to 3 months at 40 °C, 23 °C, and 4 °C. Extended storage at 4 °C and short exposure at higher temperatures of potential vampire bat environments preserved vaccine titers within CMC. Furthermore, physical properties of our CMC formulation were compared to the previously used glycerin jelly at 40 °C, 20 °C, and 0 °C using rheological tests. These tests indicate that CMC exhibits optimal properties for topical application to bats even at extreme temperatures possible during field vaccination. This study advances our rabies vaccination strategy for vampire bats by providing a topical vehicle suitable for field application that may additionally be employed for other significant bat diseases.

Key words: rabies, vaccine, vampire bat, topical, carboxymethyl cellulose, biomarker, rhodamine B

1. Introduction

In Latin America, the vampire bat (*Desmodus rotundus*) is the primary reservoir of the deadly rabies virus (Rhabdoviridae, Lyssavirus)[1]. First reported in the early 1900's, vampire bat-transmitted rabies (VBR) poses significant health risk to humans[2,3] and, especially, to livestock[4]. Each year, VBR causes an estimated economic burden of $50 million USD to the livestock industry[5]. Additionally, the suitable habitat range for vampire bats is expected to spread due to changing climates[6,7], and annual reports of VBR are trending upwards in several countries[1,8]. Recent models predict future expansion of the bats into the southern United States[7] posing risk of extensive damages in this region with high populations of livestock[9]. Across the US, an estimated 40,000 individuals annually receive rabies prophylaxis post-exposure to natively found bat species[10], and 90% of fatal rabies cases acquired from 2000-2022 were of bat origin[11].

Attempts to mitigate VBR rely on vaccination of livestock and proactive or reactive culling of bat colonies via application of "vampiricide" – a topical paste containing poisons that spread by the allogrooming behavior of the bats[12]. However, since their inception in the 1970's, these methods have been insufficient in preventing the expansion of VBR. Reactive application of vampiricide has been shown to exacerbate rabies virus transmission by provoking dispersal of infected bats to new roosts[13,14].

Furthermore, the poison may transfer to multiple non-target species in the wild, causing

unintended mortality[1,12].

The ingestion of topically applied (orotopical) vaccines has been previously proposed as

a superior method for bolstering current control strategies[15,16]. Mathematical simulations

predict that transferrable vaccines that similarly leverage bat allogrooming behavior

outperform culling practices in reducing the probability, size, and duration of rabies

outbreaks[17]. In earlier work, members of our group developed and tested a recombinant

raccoon poxvirus (Poxviridae: Orthopoxvirus) expressing a mosaic lyssavirus

glycoprotein (RCN-MoG)[18,19]. This vaccine was shown to produce protective immunity

intradermally in A/J mice (*Mus musculus*)[18], and orotopically in big brown bats

(*Eptesicus fuscus*)[18]. Notably, orotopical vaccination blocked shedding of rabies virus in

vampire bats[19]. Our team investigated glycerin jelly (GJ) as a topical vehicle to deliver

the RCN-MoG vaccine in these captive bat colonies[18,19] and to assess topical vehicle

uptake in the wild[17]. However, the highly temperature dependent physical properties of

GJ preclude its practicality for use in the field throughout Latin America, where tropical

and subtropical climates commonly exceed its melting point[20,21]. Additionally,

crystallization of GJ was noted in a separate field trial in Jalisco, Mexico[22].

In this work, we describe the optimization of a novel carboxymethyl cellulose (CMC) gel

vehicle for orotopical administration of the RCN-MoG vaccine to vampire bats. We

compared physical properties of CMC gel and GJ with rheological tests to emulate

application at environmentally relevant temperatures. Vaccine preservation was

assessed by tracking viral titers over three-months across a range of temperatures.

Lastly, we applied CMC gel mixed with the fluorescent biomarker rhodamine B (RB) to a

wild colony of microchipped vampire bats in rural Jalisco, Mexico to investigate gel

transfer and uptake of CMC in a field setting.

2. Methods

Characterization of Physical Properties

Gel Preparation

Precise weights of CMC powder (Sigma-Aldrich, cat. no. 419281; St. Louis, Missouri,

USA) were added to ultrapure water (Modulab® High Flow, Evoqua, Pittsburgh,

Pennsylvania, USA) to prepare bulk solutions of 10%, 12.5%, and 15% (w/v). CMC gels

were dissolved at room temperature overnight.

GJ (Carolina Biological Supply Co, cat. no. 865495; Burlington, North Carolina, USA)

was melted at 37 °C and combined with ultrapure water at a ratio of 55.5:44.5 as was

previously described[17]. GJ was solidified and kept at room temperature overnight.

Rheological Measurements

For characterization of rheological properties, an ARES LS2 Advanced research grade

rheometer (TA Instruments, New Castle, Delaware, USA) was fitted with 50 mm cone

and plate geometry. An FS18-HD Refrigerated/Heating Circulator (Julabo, Allentown,

Pennsylvania, USA) and Peltier Plate were used for temperature control. Two different

rheological tests were performed at 0 °C, 20 °C, and 40 °C with at least 5 min allowed

for temperature equilibration. Testing was performed the day after gel preparation.

The first test, a steady rate sweep test, was used to record viscosity over a shear ramp

from 0.05 s^{-1} to 200 s^{-1}. Low shear rates characterize the near-zero shear rate and

mimic behavior at rest. The range of 20 s^{-1} to 200 s^{-1} has previously been described as

the shear rate to investigate spreading properties[23,24].

Secondly, a dynamic strain sweep test characterized the viscoelastic properties of CMC

and GJ**.** The elastic (storage) modulus G' and viscous (loss) modulus G" were

measured at a frequency of 1 hz with an initial strain of 0.015% up to 1000%.

RCN-MoG Vaccine Stability Assay

Cells and Viruses

Vero cells (ATCC CCL-81) were cultured in Dulbecco's Modified Eagle Media (DMEM;

Sigma-Aldrich, St. Louis, Missouri, USA) supplemented with 5% fetal bovine serum and

1x antibiotics-antimycotics and incubated at 37°C with 5% CO_2. The previously

described RCN-MoG vaccine[18] was used in the viral stability assay.

Vaccine Gel Preparation

To prepare RCN-MoG vaccine in CMC gel, RCN-MoG virus was homogenized in sterile

10mM Tris hydrochloride (Tris-HCl) buffer (pH 9.0) at a final concentration of 10^8

PFU/mL in 4 mL total volume. CMC powder was then added at a final concentration of

12.5% (0.5 g) and stirred with a sterile microspatula.

For GJ preparation, RCN-MoG virus was added to sterile 10mM Tris-HCl buffer (pH

9.0). Then melted glycerin jelly was added such that the ratio of Tris buffer to glycerin

jelly was 55.5:44.5 and the final concentration of RCN-MoG was 10^8 plaque-forming

units (PFU)/mL in 4 mL total volume. After homogenizing, 100 µL aliquots of the GJ

preparation were transferred with a needle-free syringe into sterile 1.5 mL Eppendorf

tubes.

Both the GJ aliquots and CMC gel were stored at 4°C overnight to allow the CMC

powder to dissolve completely. Next, the CMC gel was briefly mixed and likewise

aliquoted (100 µL) into sterile 1.5 mL Eppendorf tubes with a needle-free syringe.

Aliquots were then placed at their respective storage temperatures (4 °C, 23 °C, or 40

°C). At each timepoint (24 hr, 48 hr, 96 hr, 1 wk, 2 wk, 1 mo, 3 mo), two aliquots were

moved from each temperature and stored at -80 °C until titration by viral plaque assay.

Titration of RCN-MoG Vaccine by Viral Plaque Assay

Viral plaque assays were performed with Vero cells in 24-well plates (TPP, Trasadingen,

Switzerland). Gel samples were thawed at 4 °C and dissolved by adding 900 µL of 1x

DMEM to each tube with 1 hr incubation at 37 °C. Samples were mixed by pipetting

every 15 min to facilitate dissolution. Cells were inoculated with 100 µL of ten-fold

dilutions of the dissolved gels. After 1 hr of infection at 37 °C with 5% CO_2, the inoculum

was removed and 1 mL of plaque assay media (1x DMEM, 1.5% CMC, 2% FBS, 1x

antibiotics-antimycotics) was added. Plates were fixed with 10% formaldehyde after 3

days of infection and stained with 1% crystal violet and 20% ethanol. Log fold change in

viral titer was standardized to the initial time point (T=0).

Western Blot

Six well plates of Vero cells were infected with 100 µL of dissolved CMC gel. After 18

hours, when >80% CPE was present, the supernatant and cells were harvested and

centrifuged at 3,000 g for 10 minutes. After decanting, the cell pellet was resuspended

in 100 µL of 4X Laemmli Sample Buffer (Bio-Rad, Richmond, California, USA)

containing 10% 2-mercaptoethanol (Sigma Aldrich, Germany). Samples were denatured

at 95°C for 5 min. SDS-PAGE resolved samples were then transferred to a

nitrocellulose membrane (Bio-Rad, Richmond, California, USA). The membrane blocked

was at 23 °C in 5% milk in PBS-T (blocking buffer) for 2 hours. Rabies virus glycoprotein

polyclonal antibody (Invitrogen, Waltham, Massachusetts, USA) was used at 1:1000 in

blocking buffer with 5% rabbit serum for 90 minutes at 23 °C. After three washes with

PBS-T, Goat anti-Rabbit IgG (H+L) Secondary Antibody, HRP (Invitrogen, Waltham,

Massachusetts, USA) was used at 1:4000 in blocking buffer with 5% goat serum for 90

minutes at 23 °C. After a final three washes, the membrane was developed with Pierce

1-step Ultra TMB Blotting solution (Thermo Fisher Scientific, Waltham, Massachusetts,

USA). Membranes were scanned with an Epson Scanner (EPSON Perfection 4490

Photo) using the Epson Scan Utility v3.24 software.

Bat Field Study

Ethics Statement

The Institutional Animal Care and Use Committee (IACUC) at the University of

Wisconsin – Madison, WI, USA approved all animal work (Protocol No. V006701-A02).

Permit number SPARN/DGVS/05928/24 was obtained from Secretaria de Medio

Ambiente y Recursos Naturales (SEMARNAT, Mexico City, Mexico) for capturing and

collecting hair samples from vampire bats.

Study Site

The field study was performed at a vampire bat colony roosting in a long-abandoned

ranch-house ~10 km outside the Casimiro Castillo township, Jalisco, Mexico

(19.580572, -104.534957). This colony had been monitored since October 2023 with

follow-up captures every six months. During these previous captures, subcutaneous

radio frequency identification (RFID) microchips were implanted in all captured bats to

track bat movement at the roost through detections by a solar powered electromagnetic

antenna and RFID detection system (Biomark, Boise, ID).

On October 31 and November 3 and 7, 2024, we performed diurnal captures. Two

personnel entering the roost captured bats using butterfly nets, and a mist net was

installed at the lone exit of the roost to ensnare bats attempting to fly out of the roost

during capture efforts. Captured bats were immediately removed from the nets and kept

in cages until processing. On each day of capturing, three capture sessions of 10 min

spaced 1.5 hr apart were performed. Each captured bat was sexed and aged, with age

determined by assessing fusion of the phalangeal epiphyses[25]. For analysis, subadults

were grouped as juveniles.

Twenty of the bats were euthanized during captures 2 and 3 by local authorities as part

of the activities of the Mexican rabies campaign for rabies prevention in livestock

species due to equine rabies cases reported nearby.

Gel Application and Sample Collection

The day prior to the first bat capture, 0.225 g (final concentration 0.3% w/v) RB was

dissolved in 75 mL water. Then, 9.375 g CMC was added and manually stirred for a final

concentration of 12.5%. The gel dissolved overnight.

During bat capture 1 (October 31, 2024), 1 mL of the gel was topically applied to the

dorsal fur of bats with a needle free syringe. These bats were also given 100 μL orally to

act as gel consumption controls for hair sampling. This dose of RB was previously

determined to clearly mark vampire bat hair samples. The bats were then released back

to the colony. On captures 2 and 3 (November 3 and 7, 2024), hair samples were

plucked with tweezers from captured bats and stored in envelopes protected from light

for later visualization with a fluorescence microscope.

RB Visualization in Bat Hair

Uptake of RB within the hair samples was qualitatively assessed with a Zeiss Imager A2

microscope (Oberkochen, Baden-Württemberg, Germany) with a Zeiss 424931

Microscope Fluorescence Filter Cube (excitation 546 nm / emission 590 nm).

Observations were performed and assessed separately by two individuals. Samples

were only recorded as positive if fluorescence was noted by both observers.

Bat Colony Size Estimation

The colony size was estimated during capture 1 by using the Lincoln-Petersen method[26]

with the microchipped bats detected in the colony the night prior as the "marked"

population. Each captured bat was scanned with an RFID scanner (AVID, Norco,

California, USA) for microchip presence. This estimation method was used to determine

the number of bats to topically treat during capture 1.

$$Colony\ size = \frac{(\#\ of\ captured\ bats) * (\#\ of\ RFIDs\ detected\ previous\ night)}{(\#\ of\ captured\ bats\ with\ RFID)}$$

Statistical Analysis

To assess variation in biomarker coverage across sex and age cohorts in our field trail

we used a generalized linear model with a binomial distribution and a logit link function.

For our response variable we used the presence of the biomarker (0/1) in the hair

samples collected in unique bats from capture sessions 2 and 3. As our predictor

variables, we included a grouping variable for both sex and age resulting in four distinct

bat cohorts: adult female, adult male, juvenile female and juvenile male. The model was

fit using the BRMS package[22] in the R statistical language (R Core Team 2024). R

version 4.4.2 was used to generate estimated biomarker coverage within the colony

based on the positivity of collected hair samples. A hypothesis test built into the BRMS

package was used to determine significant differences in the rates of coverage within

the cohorts of non-treated bats.

For our covariate estimates, we used weakly informative priors with a normal

distribution with a mean of 0 and standard deviation of 2. The model was run for 8,000

iterations with 2,000 iterations for warmup. To assess model convergence, we used the

Gelman-Rubin test statistic, with a value <1.2 indicating convergence[27] and visually

inspected the trace plots for signs of divergent transitions.

Additionally, viral titer decay rates were modeled by linear regression and statistical

comparisons were made using emtrends function of the emmeans R package[28]. A Chi-

squared test was used to analyze the age and sex distribution of bats across the three

days of captures. Fisher's exact test was used to analyze differences in RB positivity of

collected hair samples of bat cohorts stratified by the combination of age and sex.

3. Results

CMC-based vaccine formulation exhibits optimal properties for topical application

To compare the consistency of CMC and GJ across a range of temperatures,

rheological testing was performed. The temperatures tested (0 °C – 40 °C) represent

extreme conditions under which a vaccine gel might be applied to vampire bats for

rabies vaccination or to other bat species in temperate climates.

A steady rate sweep test was performed to measure viscosity across a range of shear

rates, or rates of deformation. As the viscosity of CMC solutions is modulated by

concentration (Figure 1A; Figure S1), a concentration of 12.5% was considered to have

the optimal consistency. The apparent zero-shear viscosity of 12.5% CMC, estimated by

the Newtonian plateau and an approximation of gel behavior at rest, exhibited much

variation with changes in temperature compared to GJ (Figure 1B). Zero-shear viscosity

for 12.5% CMC ranged from approximately 110-400 Pa*s from 0 to 40 °C compared to

0.05 – 1,070 Pa*s for GJ. At all temperatures, the viscosity of CMC and GJ decreased

as shear rate increased, indicating shear-thinning behavior where viscosity decreases

as deformation rates increase.

At shear rates within the range of 20-200 s^{-1} (Figure 1C), shear rates associated with

spreading[23,24], CMC was more viscous than GJ and varied minimally with changes in

temperature. In the dynamic strain sweep test, CMC gel exhibited viscoelastic

properties and acted as a viscoelastic liquid even at 40 °C (Figure 1D). GJ which

displayed properties of an elastic solid at 0 °C and 20 °C (Figure S2) but became

liquified at 40 °C (Figure 1D).

[Figure 1]

CMC gel preserves viral vaccine titers

We measured viral titers at time intervals up to three months to assess preservation of

our RCN-MoG vaccine within CMC and GJ under different temperature conditions.

Stability of the RCN-MoG vaccine titers was highly temperature dependent within both

CMC and GJ formulations (Figure 2A-C). High titers were preserved up to 3 months at 4

°C (Figure 2B) with minimal decay of RCN-MoG in both formulations (CMC: -0.0385 $\pm$

.0221 PFU/mL per week; GJ: -0.00311 $\pm$ 0.0191 PFU/mL per week; Figure 2C). At 23

°C, viral titer decreased in CMC at a moderate rate (-0.463 $\pm$ 0.062 Log10 PFU/mL per

week), while in GJ decay was slower (-0.157 $\pm$ 0.041 Log10 PFU/mL per week). RCN-

MoG viral titer rapidly decreased at 40 °C in both CMC (-3.50 $\pm$ 1.03 Log10 PFU/mL per

week) and GJ (-1.50 $\pm$ 0.28 Log10 PFU/mL per week). RCN-MoG was no longer

detectable at 40 °C after 30 days in CMC and after 90 days in GJ. Of note, slightly more

virus was recovered from dissolved GJ (Log10 8.02 +/- 0.123 PFU/mL) than CMC

(Log10 7.64 +/- 0.136 PFU/mL) at the initial time point (Figure S3).

Cells infected with RCN-MoG stored in CMC at 4 °C for three months still produced

antigen detected by western blot (Figure 2D). While replication levels of samples stored

at 23 °C reduced drastically over time, vaccine poxviral DNA was preserved throughout

the study (Table S1-S2; Figure S4).

[Figure 2]

High uptake rate of CMC-based delivery formulation within a wild vampire bat

colony

CMC gel uptake by wild vampire bats was measured using the fluorescent biomarker

rhodamine B (RB) within a microchipped colony in Jalisco, Mexico (Figure 3A). The bats

inhabited an abandoned ranch-house (Figure 3B) in which we previously installed an

RFID system for microchip detection (Figure 3C). Over three capture sessions (Figure

3D), 156 total *Desmodus rotundus* bats were captured (Figure 3E), representing 95

unique bats. No bats of other species were captured.

During capture 1, 47/89 (52.8%) bats had RFID microchips from previous trips to the

site (Figure 3E), while 62 microchipped bats were detected the night before capturing

(Figure 3F). This results in an estimated colony size of 117.4 ($\pm$ 5.625 SE) by the

Lincoln-Petersen method. One milliliter of 12.5% CMC with 0.3% rhodamine B (RB) was

topically applied to the dorsal fur of 24 bats (13 females and 11 males), representing an

estimated 20.5% of the colony. Forty-two new microchips were implanted during capture

1, resulting in the observed increase in detected RFIDs (Figure 3F).

Age and sex demographics within the entire colony were determined from the

proportions of unique bats captured belonging to each subpopulation across the three

captures. Of the bats in the colony, 57/95 were female (60.00%), and 66/95 were adults

(69.47%) (Table 1). RFID data shows a decline in the number of microchipped bats

detected each night within the colony beginning after capture 2 (Figure 3F). The age

and sex distribution of bats captured during captures 2 and 3 were not statistically

different from capture 1 (Chi-squared test, p = 0.165).

[Table 1]

[Figure 3]

During bat capture sessions 2 and 3, 34 and 27 hair samples were collected,

respectively. Accounting for recaptures, hair samples were obtained from 48 unique

bats, including 13 that were initially topically treated with CMC gel (Figure 4A-D). Hair

samples were visualized with a fluorescent microscope (Figure 4E-F), and all 13 treated

bats that were recaptured were RB-positive. Of the nontreated bats, 29/35 (82.86%)

were RB positive. Notably, the only negative hair samples belonged to adult male bats

resulting in a statistically different proportion of positive samples compared to the other

cohorts (Fisher's exact test, p = 0.002). Three adult male bats that were captured during

both captures for hair sample collection had positive hair samples on capture 2 but

negative samples on capture 3 (Figure 4A). These individuals were still considered RB

positive for analysis.

Based on the proportion of positive and negative hair samples and colony demographic

estimates, total gel uptake by the colony was modeled using Bayesian regression

modelling with the BRMS package[29]. Gel uptake within non-treated and non-captured

bats was predicted using the brm model (Figure 4G). Near 100% rates of uptake were

predicted in non-treated adult female (97.4%; confidence interval (CI): 85.7% - 100%),

juvenile female (94.5%; CI: 73.3% - 100%) and juvenile male (98.8%; CI: 92.3% -

100%) cohorts. Notably, an estimated uptake of 48.4% (CI: 30.4%-69.6%) was found in

non-treated adult males. Across the entire colony, uptake was estimated at 88.0%

(103/117; 90 - 110) (CI: 76.9% - 94.0%). Based on this projection, 103 bats consumed

the gel, with each bat consuming on average ~200 µL of the total 24 mL of topically

applied gel (1 mL applied to 24 bats), assuming all the gel was consumed.

[Figure 4]

4. Discussion

While rabies vaccination of vampire bats in captivity has been shown to reduce rabies

virus shedding[19] the delivery of the vaccine to wild vampire bats presents a major

barrier to the implementation of this strategy as a rabies control measure. Although

effective in humans and domesticated animals, parenteral vaccination is impractical for

bats. Likewise, oral baits successfully used for other wildlife species[30] are not suitable

for bats due to differing feeding behavior. Because topically administered poisons

demonstrate high rates of oral uptake within bat colonies[31,32], orotopical vaccination of

vampire bats has been proposed but lacks an optimal delivery vehicle. In this work, we

describe the development of a topical vehicle that maintains optimal physical properties

even at extreme temperatures, preserves our viral vector vaccine candidate, and

exhibits high rates of transfer and oral uptake between treated and untreated bats in a

wild bat colony.

Previous bat studies tested petroleum jelly[33] – a paste commonly used for anticoagulant

poison delivery – or glycerin jelly (GJ)[17–19] as the topical vaccine delivery vehicles.

Petroleum jelly proved difficult to work with resulting in dilution of the vaccine and

substantial loss of vaccine paste[33]. Meanwhile, the consistency of GJ is highly

temperature dependent as demonstrated by our study. We found the mixture of GJ in

water previously used[17] has a melting temperature around 30 °C. While vampire bats

typically roost where temperatures remain below 37 °C[34] and do not well tolerate

temperatures above 33 °C[35], environmental temperatures may exceed 40 °C during

heat waves in some regions[20].

Characterization of physical properties of a topical vehicle under environmental

temperatures is necessary for proper application as revealed by the shortcomings of

GJ. Rheology, measuring how a material sample responds to deformation, presents a

useful method for measuring physical properties of gel-like materials[36]. The rheological

properties of CMC solutions are well studied and vary with the molecular weight and

degree of substitution of the CMC powder used[37–39]. The CMC gel preparation used in

our study demonstrates shear thinning behavior and viscoelastic properties

corresponding with these previous reports. In a steady rate sweep test, CMC

demonstrates less variation than GJ in zero-shear viscosity over the studied range of

temperature. At 40 °C, the near-zero shear rate of CMC was approximately 110 Pa*s,

demonstrating high viscosity at high temperatures. This improves on the GJ formulation

which has a near-zero shear rate approaching 0 Pa*s at 30 °C when it becomes

completely liquified. In the characterization of viscoelastic materials, the storage

modulus G' characterizes the elastic component (energy stored during deformation) and

the loss modulus G" characterizes the viscous component (energy dissipated during

deformation). In CMC, G" is greater than G' in the linear viscoelastic region which

suggests CMC acts as a viscoelastic liquid that flows slowly over time. This may

contribute to improved surface wettability and to the enhanced stickiness of CMC that

we observed compared to GJ. In GJ, G' > G", indicating primarily elastic solid-like

behavior. Notably at 0 °C, G' becomes greater than G" in CMC indicating trends toward

solidification at temperatures near freezing, while still exhibiting viscoelastic properties.

Additional described properties of CMC support its use as a topical vaccine vehicle. The

muco-adhesion of CMC[40] may enhance viral entry into mucosal tissue. Mucoadhesive

cellulose-based films have been previously used to facilitate entry of Modified Vaccinia

Ankara – a commonly used poxviral vaccine vector - into the buccal mucosa of mice[41].

CMC is also commonly used in drug delivery[42], is non-toxic at high quantities[43], and was

found to mix easily with buffer containing RB.

To facilitate vaccination strategies, the vaccine must be readily shipped and stored

without dependence on cold-chains in regions where reliability of infrastructure may be

unpredictable[44]. Our assessment of viral stability in CMC under different temperature

conditions provides insight into vaccine preservation under storage conditions and

persistence in field-relevant conditions. Minimal decay of viral titer within CMC was

evident at refrigeration temperatures (4 °C) with only a 0.63 log fold decrease in viral

titer after 3 months of storage. This will enable shipment and storage of the vaccine

when challenges in infrastructure prevent maintaining temperatures below freezing. If

refrigeration is lost, vaccine titer may also be preserved up to 1 week at 23 °C with only

a 0.46 log fold decrease in titer and 24 hrs at 40 °C with 0.54 log fold decrease.

Although virus stability was generally better preserved in GJ;, the viral titer initially

decreased more rapidly in GJ than in CMC at 40 °C with a log fold decrease of 0.82

after 24 hours. Other additives could be considered to improve stability within the CMC

gel formulation such as sucrose, which has been shown to enhance poxviral vaccine

vector stability[45]. The implications of these vaccine preservation data depend on the

determination of the minimum effective dose of RCN-MoG in vampire bats; however,

during field application, we expect gel to be consumed within hours after application.

Like applications of vampiricide, orotopical vaccination utilizes the natural propensity of

vampire bats to allogroom to facilitate the transfer of vaccine between treated and

untreated bats. Ideally, only a small portion of a colony would need to be treated to

effectively manage rabies due to difficulties in locating and treating entire roosts.

Vampire bat captures are commonly performed at ranches where bats are known to be

feeding to apply vampiricide, rather than at the roost istelf[31]. This highlights the

importance of understanding the rates of vaccine transfer and uptake within colonies

and between cohorts at low rates of application. The RB biomarker used in this study is

a common proxy for measuring uptake of treatment in wildlife species given its

properties to stably mark the hair follicle and shaft with a fluorescent signal[17,46,47]. Based

on assumptions about population size and demographic distribution of bats in this trial,

an estimated gel uptake by 88.0% (CI: 76.9% - 94.0%) of the colony was measured

after initial treatment of 20.5% of the colony. This corresponds to gel uptake in 3.29 bats

(CI: 2.75 – 3.58) per initial bat treated which improves upon the transfer reported in

Bakker et al. (2019) of 1.8-2.1[17]. This difference is likely related to the increase in

volume of gel applied per bat from 0.45 mL to 1 mL and the higher proportion of

remaining non-treated bats available for gel uptake. Given the near 100% uptake

observed in adult females and juveniles in our study, gel transfer rates may be even

higher given the finite number of untreated bats within the colony.

Importantly, adult male bats are disproportionately implicated in the dispersal of VBR[48].

While our field trial suggests that adult male bats may be less likely to consume

orotopically applied vaccines vehicles, we still demonstrate predicted uptake by 48.4%

(CI: 30.4%-69.6%) of non-treated adult male bats. It has also previously been

documented that males may be less effective than females at spreading vampiricide

within a colony[32]. However, this result differs from Bakker et al. (2019) who found a

nonsignificant difference in the proportion of males that showed uptake of RB compared

to females[17]. This previous finding could be associated with the high proportion of

treatment (up to 50%) in the colonies reported, variation in the social dynamics within

the colonies between the studies, or potential regional or seasonal influences. Vampire

bat colonies are differentiated as either maternity colonies with resident and non-

resident males or primarily-male bachelor roosts[49,50]. Social grooming interactions and

thus topical vehicle uptake may vary between different sites with different structures and

sex and age ratios[51]. Notably, 2.3-9 times the proportion of juveniles (31.5%) were

captured in the roost in the present study compared to the three roosts studied in

Bakker et al. (2019) (3.5%, 8.7% and 13.6%) during the first diurnal capture[17].

While RFID detection remained stable between captures 1 and 2 in this study, the

number detected declined by around 50% a week later. In addition to the multiple

captures at this site, disturbance was caused by the euthanasia of 20 of the bats from

our colony carried out by authorities of the Mexican Rabies Campaign as equine rabies

cases were reported near the roost site. As previously reported[13,17], vampire bats are

prone to roost abandonment when disturbed, relocating to nearby manmade and natural

roosts. For example, five months before the initiation of this biomarker study, we visited

a large colony of over 100 bats within an abandoned facility 10 km away as a potential

study site. However, before captures could be initiated, the colony relocated with only a

few bats remaining at the site.

Epidemiological models indicate that vaccination significantly reduces duration and

frequency of rabies outbreaks compared to culling even at low rates of vaccination[17].

The estimated gel consumption rates by nearly 100% of adult females, and male and

female juveniles found in this study will likely aid in preventing rabies virus introduction

to naive colonies. As a lower proportion of adult males consumed the gel, targeting

males as juveniles may result in higher rates of vaccination; however, this will require

investigation of the development and duration of the immune response in vaccinated

juvenile bats. Further field trials are needed to understand gel uptake dynamics within

multiple bat colonies to maximize vaccine uptake by adult males. Additional models may

also investigate the effect of differing rates of vaccination uptake between males and

females. Given that vampire bat rabies has been known to spread among colonies in a

slow wavelike fashion[1,52,53], strategies of ring vaccination around areas of predicted

spread may be effective at containing expansion of VBR[17].

CMC gel may also work well as a topical vehicle for vaccine delivery to other bat

species. In the United States, an estimated 40,000 individuals receive rabies

prophylaxis each year after exposure to various bat species[10]. Furthermore, 36 of 40

(90%) fatal rabies cases acquired in the United States between 2000 and 2022 were of

bat origin[11]. CMC could additionally be used to deliver a white nose syndrome (WNS)

vaccine[54] or to scale up deployment of a herpesvirus-vectored rabies vaccine[55]. As

WNS-affected bats often inhabit colder regions in the United States and Canada, it is

notable that CMC additionally maintains its physical properties at colder temperatures.

However, vampire bats participate in allogrooming at higher rates than other bat

species[56], and field studies would be needed to determine gel transfer within colonies of

other targeted bat species.

The identification of vampire bats within 50 km of the southern border of the United

States[57] and the expectation that their geographic range will continue to expand

emphasizes the need for superior control methods[7]. While topical application of

vaccines presents an appealing alternative to culling practices that may exacerbate

rabies transmission, further studies are needed to understand the dimorphism of uptake

rates in adult male and female bats observed in this study. Likewise, additional studies

are needed to determine the impact of vaccination on vampire bat colony sizes,

although it is proposed to include contraceptives as a method for population control

while minimizing unintended ecological impacts of anticoagulant poisons[58].

Conclusion

This study details the characterization of a carboxymethyl cellulose (CMC) gel vehicle

for orotopical rabies vaccine delivery to vampire bats. The new CMC formulation

maintains optimal physical properties across a broad range of temperatures improving

on a past topical vehicle. CMC gel is also able to sufficiently preserve our recombinant

viral-vector rabies vaccine candidate. A field study in Jalisco, Mexico demonstrates the

capacity for high rates of uptake of the CMC-based delivery formulation within a wild

vampire bat colony at a low level of application. In summary, CMC gel provides an

orotopical vehicle for vaccine delivery to bat species that socially groom and may be

tested to treat other diseases and species beyond rabies in vampire bats.

Data Availability

The data and code that support the findings of this study are openly available in github

at https://github.com/zamanianlab/Bat_vaccine_gel-ms.

Acknowledgements

We are very grateful for the support of team members from the Universidad de

Guadalajara - Centro Universitario de la Costa Sur (CUCSUR) and Comité Estatal para

el Fomento y Protección Pecuaria Jalisco (CEFPPJ) who aided in the field work. We

would also like to thank the Centro Nacional de Servicios de Diagnóstico en Salud

Animal (CENASA) in Tecamac, Mexico inviting us to work with their fluorescence

microscope and assisting with visualizing the hair samples.

The authors gratefully acknowledge the use of facilities and instrumentation supported

by NSF through the University of Wisconsin Materials Research Science and

Engineering Center (DMR-1121288, 0079983, 0520057, 0832760, and 0425880) with

special thanks to staff of the Soft Materials Characterization Lab for providing training

on rheometer use and data analysis.

The project was funded by the NSF/BBSRC Ecology and Evolution of Infectious

Diseases Program (DEB 2011069, BB/V003798/1) and the University of Wisconsin

Madison Comparative Biomedical Sciences Student Travel Award (C.K.). D.G.S. was

funded by a Wellcome Trust Senior Research Fellowship (217221/Z/19/Z). This

research received support from the UK Medical Research Council through core funding

of the MRC-University of Glasgow Centre for Virus Research (Virus Cross Species

Transmission Programme: MC_UU_00034/3).

Any use of trade, firm, or product names is for descriptive purposes only and does not

imply endorsement by the U.S. Government.

Author Contributions

Conceptualization, C.K., E.M.C.C., D.P.W., D.G.S., J.E.O., M.Z., T.E.O.; data curation,

C.K., T.M.G., M.Z.; formal analysis, C.K., T.M.G., funding acquisition, C.K., D.P.W.,

D.G.S., J.E.O., M.Z., T.E.O.; investigation C.K., E.M.C.C., T.M.G., M.M.R.M., D.L., L.P.,

D.P.W., D.G.S, T.E.O.; methodology, C.K., E.M.C.C., D.L., M.Z., T.E.O.; project

administration, C.K., E.M.C.C., M.M.R.M., D.P.W., D.G.S., M.Z., T.E.O.; resources,

M.M.R.M, J.E.O., M.Z., T.E.O.; software, C.K., T.M.G.; supervision, E.M.C.C., M.M.R.M.,

M.Z., T.E.O.; validation, C.K., E.M.C.C., M.Z., T.E.O.; visualization, C.K., T.M.G., D.L.;

writing – original draft preparation, C.K.; writing – review & editing, C.K., E.M.C.C.,

T.M.G., M.M.R.M., D.L., L.P., D.P.W., D.G.S., J.E.O., M.Z., T.E.O.

References

1. Johnson, N., Aréchiga-Ceballos, N. & Aguilar-Setien, A. Vampire bat rabies: ecology,

epidemiology and control. *Viruses* **6**, 1911–1928 (2014).

2. Meske, M. *et al.* Evolution of Rabies in South America and Inter-Species Dynamics

(2009-2018). *Trop Med Infect Dis* **6**, 98 (2021).

3. Schneider, M. C. *et al.* Rabies transmitted by vampire bats to humans: an emerging

zoonotic disease in Latin America? *Rev Panam Salud Publica* **25**, (2009).

4. Soler-Tovar, D. & Escobar, L. E. Rabies transmitted from vampires to cattle: An

overview. *PLoS One* **20**, e0317214 (2025).

5. Belotto, A., Leanes, L. F., Schneider, M. C., Tamayo, H. & Correa, E. Overview of

rabies in the Americas. *Virus Research* **111**, 5–12 (2005).

6. Lee, D. N., Papeş, M. & Van den Bussche, R. A. Present and potential future

distribution of common vampire bats in the Americas and the associated risk to cattle.

PLoS One **7**, e42466 (2012).

7. Van de Vuurst, P., Gohlke, J. M. & Escobar, L. E. Future climate change and the

distributional shift of the common vampire bat, Desmodus rotundus. *Sci Rep* **15**, 5989

(2025).

8. Benavides, J. A. *et al.* Defining New Pathways to Manage the Ongoing Emergence of

Bat Rabies in Latin America. *Viruses* **12**, 1002 (2020).

9. Anderson, A., Shwiff, S., S. & Shwiff, S., A. Economic Impact of the Potential Spread

of Vampire Bats into South Texas. *vertebrate_pest_conference* **26**, (2014).

10.   Kunkel, A. *et al. Notes from the Field:* Three Human Rabies Deaths Attributed to

Bat Exposures — United States, August 2021. *MMWR Morb. Mortal. Wkly. Rep.* **71**,

31–32 (2022).

11.   Ma, X. *et al.* Rabies surveillance in the United States during 2022. *javma* **262**,

1518–1525 (2024).

12.   Ávila-Vargas, L., Soler-Tovar, D., Dong, Q. & Escobar, L. E. Anticoagulants for

the Control of the Common Vampire Bat (Desmodus rotundus). *Zoonoses Public*

Health **72**, 101–116 (2025).

13.   Viana, M. *et al.* Effects of culling vampire bats on the spatial spread and spillover

of rabies virus. *Sci Adv* **9**, eadd7437 (2023).

14.   Streicker, D. G. *et al.* Ecological and anthropogenic drivers of rabies exposure in

vampire bats: implications for transmission and control. *Proc Biol Sci* **279**, 3384–3392

(2012).

15.   Almeida, M. F., Martorelli, L. F. A., Aires, C. C., Barros, R. F. & Massad, E.

Vaccinating the vampire bat Desmodus rotundus against rabies. *Virus Research* **137**,

275–277 (2008).

16.   Sétien, A. A. *et al.* Experimental rabies infection and oral vaccination in vampire

bats (Desmodus rotundus). *Vaccine* **16**, 1122–1126 (1998).

17.   Bakker, K. M. *et al.* Fluorescent biomarkers demonstrate prospects for

spreadable vaccines to control disease transmission in wild bats. *Nat Ecol Evol* **3**,

1697–1704 (2019).

18.   Stading, B. *et al.* Protection of bats (Eptesicus fuscus) against rabies following

topical or oronasal exposure to a recombinant raccoon poxvirus vaccine. *PLoS Negl*

Trop Dis **11**, e0005958 (2017).

19.   Cárdenas-Canales, E. M. *et al.* A recombinant rabies vaccine that prevents viral

shedding in rabid common vampire bats (Desmodus rotundus). *PLoS Negl Trop Dis*

16, e0010699 (2022).

20.   El Niño and climate change impacts slam Latin America and Caribbean in 2023.

World Meteorological Organization https://wmo.int/news/media-centre/el-nino-and-

climate-change-impacts-slam-latin-america-and-caribbean-2023 (2024).

21.   South America - Climate, Geography, Altitude | Britannica.

https://www.britannica.com/place/South-America/Factors-influencing-climate (2025).

22. Cárdenas-Canales, E. M. Vaccinating vampire bats against rabies□: studies of

vaccine efficacy, field application, and social perceptions in México. (University of

Wisconsin - Madison, 2021).

23. Boylan, J. C. Rheological Study of Selected Pharmaceutical Semisolids. *Journal*

of Pharmaceutical Sciences **55**, 710–715 (1966).

24. Mahdi, M. H., Conway, B. R., Mills, T. & Smith, A. M. Gellan gum fluid gels for

topical administration of diclofenac. *International Journal of Pharmaceutics* **515**, 535–

542 (2016).

25. *Ecological and Behavioral Methods for the Study of Bats.* (Johns Hopkins

University Press, Baltimore, 2009).

26. Otis, D. L., Burnham, K. P., White, G. C. & Anderson, D. R. *Statistical Inference*

from Capture Data on Closed Animal Populations. (Wiley, 1978).

27. Gelman, A. & Rubin, D. B. Inference from Iterative Simulation Using Multiple

Sequences. *Statist. Sci.* **7**, (1992).

28. Lenth, R. Estimated marginal means, aka least-squares means. R package

version 1.7.2.

29. Bürkner, P.-C. **brms**□: An *R* Package for Bayesian Multilevel Models Using *Stan.*

J. Stat. Soft. **80**, (2017).

30. Maki, J. *et al.* Oral vaccination of wildlife using a vaccinia-rabies-glycoprotein

recombinant virus vaccine (RABORAL V-RG®): a global review. *Vet Res* **48**, 57

(2017).

31. Linhart, S. B., Flores Crespo, R. & Mitchell, G. C. [Control of vampire bats by

means of an anticoagulant]. *Bol Oficina Sanit Panam* **73**, 100–109 (1972).

32.   Gomes, M. N., Uieda, W. & Latorre, M. D. R. D. D. O. Influência do sexo de

indivíduos da mesma colônia no controle químico das populações do morcego

hematófago Desmodus rotundus (Phyllostomidae) no estado de São Paulo. *Pesq.*

Vet. Bras. **26**, 38–43 (2006).

33.   Almeida, M. F., Martorelli, L. F. A., Aires, C. C., Sallum, P. C. & Massad, E.

Indirect oral immunization of captive vampires, Desmodus rotundus. *Virus Research*

111, 77–82 (2005).

34.   Ribeiro, R. *et al.* Incorporating environmental heterogeneity and observation

effort to predict host distribution and viral spillover from a bat reservoir. *Proc. R. Soc.*

B. **290**, 20231739 (2023).

35.   Lyman, C. P. & Wimsatt, W. A. Temperature Regulation in the Vampire Bat,

Desmodus rotundus. *Physiological Zoology* **39**, 101–109 (1966).

36.   Rueda, M. M. *et al.* Rheology and applications of highly filled polymers: A review

of current understanding. *Progress in Polymer Science* **66**, 22–53 (2017).

37.   Dong, S. *et al.* Factors influencing the adhesive behavior of carboxymethyl

cellulose-based hydrogel for food applications. *International Journal of Biological*

Macromolecules **179**, 398–406 (2021).

38.   Benchabane, A. & Bekkour, K. Rheological properties of carboxymethyl cellulose

(CMC) solutions. *Colloid Polym Sci* **286**, 1173–1180 (2008).

39.   Edali, M., Esmail, M. N. & Vatistas, G. H. Rheological properties of high

concentrations of carboxymethyl cellulose solutions. *J. Appl. Polym. Sci.* **79**, 1787–

1801 (2001).

40. Cook, S. L., Woods, S., Methven, L., Parker, J. K. & Khutoryanskiy, V. V.

Mucoadhesive polysaccharides modulate sodium retention, release and taste

perception. *Food Chem* **240**, 482–489 (2018).

41. Esih, H. *et al.* Mucoadhesive film for oral delivery of vaccines for protection of the

respiratory tract. *Journal of Controlled Release* **371**, 179–192 (2024).

42. Javanbakht, S. & Shaabani, A. Carboxymethyl cellulose-based oral delivery

systems. *Int J Biol Macromol* **133**, 21–29 (2019).

43. Mondal, Md. I. H. & Yeasmin, Mst. S. Toxicity study of food-grade carboxymethyl

cellulose synthesized from maize husk in Swiss albino mice. *International Journal of*

Biological Macromolecules **92**, 965–971 (2016).

44. Seaman, C. P. *et al.* Controlled temperature chain for vaccination in low- and

middle-income countries: a realist evidence synthesis. *Bull World Health Organ* **100**,

491–502 (2022).

45. Eilts, F. *et al.* An investigation of excipients for a stable Orf viral vector

formulation. *Virus Research* **336**, 199213 (2023).

46. Fernandez, J. R.-R. & Rocke, T. E. Use of Rhodamine B as a Biomarker for Oral

Plague Vaccination of Prairie Dogs. *Journal of Wildlife Diseases* **47**, 765–768 (2011).

47. Cagnacci, F., Massei, G., Coats, J., De Leeuw, A. & Cowan, D. P. Long-lasting

Systemic Bait Markers for Eurasian Badgers. *Journal of Wildlife Diseases* **42**, 892–

896 (2006).

48. Streicker, D. G. *et al.* Host–pathogen evolutionary signatures reveal dynamics

and future invasions of vampire bat rabies. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 10926–

10931 (2016).

49.   Delpietro, H. A., Russo, R. G., Carter, G. G., Lord, R. D. & Delpietro, G. L.

Reproductive seasonality, sex ratio and philopatry in Argentina's common vampire

bats. *R. Soc. open sci.* **4**, 160959 (2017).

50.   Delpietro, H. A. & Russo, R. G. Observations of the common vampire bat

(Desmodus rotundus) and the hairy-legged vampire bat (Diphylla ecaudata) in

captivity. *Mammalian Biology* **67**, 65–78 (2002).

51.   Ávila-Vargas, L., Soler-Tovar, D., Dong, Q. & Escobar, L. E. Anticoagulants for

the Control of the Common Vampire Bat (Desmodus rotundus). *Zoonoses Public*

Health **72**, 101–116 (2025).

52.   Delpietro, H., De Díaz, A. M., Fuenzalida, E. & Bell, J. F. [Determination of the

rate of rabies attack in bats]. *Bol Oficina Sanit Panam* **73**, 222–230 (1972).

53.   Benavides, J. A., Valderrama, W. & Streicker, D. G. Spatial expansions and

travelling waves of rabies in vampire bats. *Proc. R. Soc. B.* **283**, 20160328 (2016).

54.   Rocke, T. E. *et al.* Virally-vectored vaccine candidates against white-nose

syndrome induce anti-fungal immune response in little brown bats (Myotis lucifugus).

Sci Rep **9**, 6788 (2019).

55.   Griffiths, M. E., Meza, D. K., Haydon, D. T. & Streicker, D. G. Inferring the

disruption of rabies circulation in vampire bat populations using a betaherpesvirus-

vectored transmissible vaccine. *Proc. Natl. Acad. Sci. U.S.A.* **120**, e2216667120

(2023).

56.   Carter, G. & Leffer, L. Social Grooming in Bats: Are Vampire Bats Exceptional?

PLoS One **10**, e0138430 (2015).

57.   Hayes, M. A. & Piaggio, A. J. Assessing the potential impacts of a changing

climate on the distribution of a rabies virus vector. *PLoS ONE* **13**, e0192887 (2018).

58.   Serrano, H., Pérez-Rivero, J.-J., Aguilar-Setién, A., de-Paz, O. & Villa-Godoy, A.

Vampire bat reproductive control by a naturally occurring phytooestrogen. *Reprod.*

Fertil. Dev. **19**, 470 (2007).

Figure and Table Legends

Figure 1. Rheological properties of carboxymethyl cellulose (CMC) gel and

glycerin jelly (GJ). A) The viscosity of 10%, 12.5% and 15% solutions of

carboxymethyl cellulose (CMC) at 40 °C and **B)** the viscosity of 12.5 % CMC and GJ at

0 °C, 20 °C, and 40 °C were measured by a steady rate sweep test. Panel **C)** displays

the viscosity of CMC and GJ depicted over shear rates associated with spreading (20 –

200 s^{-1}). **D)** Viscoelastic properties of CMC and GJ at 40 °C were measured by a

dynamic strain sweep test. G' represents the elastic modulus – energy stored during

deformation, while G" represents the loss modulus – energy dissipated during

deformation. Error bars represent standard error of three experimental replicates.

Figure 2. Viral stability of a recombinant raccoonpox vaccine expressing a mosaic

lyssavirus glycoprotein (RCN-MoG) stored within carboxymethyl cellulose (CMC)

and glycerin jelly (GJ) formulations at 4 °C, 23 °C, and 40 °C. Log fold change in

vaccine titer measured over **A)** seven days and **B)** three months of storage by plaque

assay in Vero CCL-81 cells at each time point after three days of infection. Error bars

depict the standard error of three experimental replicates. **C)** Weekly decay rates of

RCN-MoG titer within CMC and GJ at each storage condition were determined by a linear regression model. Statistical comparisons were made using emtrends (ns = no significance; * = $p < 0.05$; *** = $p < 0.001$). **D)** MoG antigen expression of cell lysates infected with RCN-MoG after storage in CMC at 4 °C for one and three months was detected by western blot. Cells were infected with 100 µL of dissolved gel and protein was collected 18 hours post-infection.

Figure 3. Study design and demographics of captured vampire bats (*Desmodus rotundus*) during three capture sessions in Jalisco, Mexico. A) In Mexico, the field study location is marked with the yellow star, while regions with the northernmost sightings of vampire bats are labeled with red circles (map retrieved from Google Earth v 10.80.0.1 on May 15, 2025). **B)** The vampire bat colony inhabited the photographed abandoned ranch-house. Photograph by C.K. on May 28, 2024. **C)** In the ranch-house, a radio frequency identification (RFID) system was installed (overhead schematic) to detect movement of microchipped bats. **D)** Three captures were performed for gel application with rhodamine B biomarker during capture 1 and hair collection during captures 2 and 3. **E)** During each capture, bats were sexed and aged. Lines represent the number of recaptures from the previous capture session, or in the case of capture 1, bats with RFID microchips from previous captures at this site. **F)** Each night, the RFID system detected the movement of microchipped bats into and out of the roost. The y-axis represents the number of unique RFIDs detected each night (x-axis).

Figure 4. Estimated gel uptake within a vampire bat (*Desmodus rotundus*) colony

in Jalisco, Mexico, after approximately 20.5% of the colony was treated topically

with 1 mL of carboxymethyl cellulose (CMC) gel containing 0.3% rhodamine B

(RB). Panels **A** (adult males), **B** (adult females), **C** (juvenile males), and **D** (juvenile

females) depict capture and treatment status during capture 1 and collection and

positivity of hair samples of bats captured during captures 2 and 3. During capture 1,

orange squares represent topically treated bats, while open circles were captured, but

not treated. The presence of orange fluorescence was used to distinguish between **E)**

positive and **F)** negative hair samples. Hair samples were visualized with a Zeiss

Imager A2 microscope (Oberkochen, Baden-Württemberg, Germany) and a Zeiss

424931 Microscope Fluorescence Filter Cube (excitation 546 nm / emission 590 nm).

G) A Bayesian model was used to calculate estimates and confidence intervals of

estimated coverage within each subpopulation of bats. Statistical significance indicates

difference in coverage of non-treated bats and was determined by brms hypothesis

testing.

Table 1. Age and sex distribution of unique captured vampire bats (*Desmodus*

***rotundus*) during three capture sessions in Jalisco, Mexico.** Age was determined

by fusion of the phalangeal epipyhses. Subadults were grouped with juveniles.

Cohort	Unique Bats Captured	Percentage of Captures
Adult Female	40	42.11%
Adult Male	26	27.37%
Juvenile Female	17	17.89%
Juvenile Male	12	12.63%
Total	95	100%