Examining the Role of Extrachromosomal DNA in 1,216 Lung Cancers

Azhar Khandekar[1,2,3,4], Phuc H. Hoang[1], Jens Luebeck[5], Marcos Díaz-Gay[2,3,4,6], Wei Zhao[1], John
P. McElderry[1], Caleb Hartman[1], Mona Miraftab[1], Olivia W. Lee[1], Kara M. Barnao[1], Erik N.
Bergstrom[2,3,4], Yang Yang[7], Martin A. Nowak[8,9], Nathaniel Rothman[1], Robert Homer[10], Soo-
Ryum Yang[11], Qing Lan[1], David C. Wedge[12,13], Lixing Yang[7,14,15], Stephen J. Chanock[1],
Tongwu Zhang[1*], Ludmil B. Alexandrov[2,3,4,16*], Maria Teresa Landi[1*]

[1]Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD, USA,
[2]Department of Cellular and Molecular Medicine, University of California San Diego, La Jolla,
CA, USA, [3]Department of Bioengineering, University of California San Diego, La Jolla, CA,
USA, [4]Moores Cancer Center, University of California San Diego, La Jolla, CA, USA,
[5]Department of Computer Science, University of California San Diego, La Jolla, CA, USA,
[6]Digital Genomics Group, Structural Biology Program, Spanish National Cancer Research
Center (CNIO), Madrid, Spain, [7]Ben May Department for Cancer Research, The University of
Chicago, Chicago, IL, USA, [8]Department of Mathematics, Harvard University, Cambridge, MA,
USA, [9]Department of Organismic and Evolutionary Biology, Harvard University, Cambridge,
MA, USA, [10]Department of Pathology, Yale School of Medicine, New Haven, CT, USA,
[11]Department of Pathology and Laboratory Medicine, Memorial Sloan Kettering Cancer Center,
New York, NY, USA, [12]Manchester Cancer Research Centre, The University of Manchester,
Manchester, UK, [13]Manchester NIHR Biomedical Research Centre, Manchester, UK,
[14]Department of Human Genetics, The University of Chicago, Chicago, IL, USA, [15]The
University of Chicago Medicine Comprehensive Cancer Center, The University of Chicago,
Chicago, IL, USA, [16]Sanford Stem Cell Institute, University of California San Diego, La Jolla,
CA, USA

26
27   [*]Correspondence should be addressed to tongwu.zhang@nih.gov,
28   L2alexandrov@health.ucsd.edu and landim@nih.gov

29
30

ABSTRACT

The role of extrachromosomal DNA (ecDNA) in lung cancer, particularly in subjects who never

smoked (LCINS), remains unclear. Examination of 1,216 whole-genome-sequenced lung cancers

identified ecDNA in 18.9% of patients. Enrichment of *MDM2* and other oncogenes'

amplification *via* ecDNA possibly drives a LCINS subset. Tumors harboring ecDNA or other

focal amplifications showed similarly poor survival. A strong association with whole-genome

doubling suggests most ecDNA reflects genomic instability in treatment-naïve lung cancer.

MAIN

Extrachromosomal DNA (ecDNA) is a potent mechanism for oncogene amplification in human

cancer[1], exhibiting unique characteristics such as a circular structure[2], non-Mendelian

inheritance[3,4], and an altered epigenetic and transcriptional landscape[5]. Prior studies have linked

ecDNA to worse overall survival in pan-cancer analyses, highlighting its clinical significance[6-8]

and potential as a therapeutic target[9]. Recent advances in computational techniques[10] have

enabled large-scale detection of ecDNA from whole-genome sequencing (WGS) data. However,

only three pan-cancer studies have examined ecDNA in primary lung cancer cohorts with over

100 samples, focusing predominantly on patients of European descent[6-8]. Two studies[6,7] analyzed

lung cancer samples from the United States, reporting ecDNA prevalence rates of about 21% in

36 lung adenocarcinomas (LUAD) and 27% in 47 lung squamous cell carcinomas (LUSC).

Additionally, a UK lung cancer study[8] used a similar methodology to analyze 14,778 patients

across 39 tumor types, including 718 LUAD tumors and 378 LUSC tumors, detecting ecDNA in

8.2% of LUAD and 22.4% of LUSC cases. All three studies[6-8] reported lung cancer findings as

part of broader pan-cancer analyses and did not distinguish between lung cancers in subjects who

have never-smoked (LCINS) and subjects who have smoked (LCSS). Notably, based on the high

prevalence of the tobacco-associated[11] mutational signature SBS4 in these studies, more than

84% of the analyzed lung cancer cases were from individuals with a history of smoking.

Despite these prior studies, the clinical relevance of ecDNA in lung cancer remains unclear,

including its prevalence across histologies, geographical areas, genetic ancestries, and biological

sexes. Its role in LCINS, LCSS, and subjects who were exposed to passive smoking also requires

further exploration. To address these questions, here, we analyzed WGS data from 1,216

treatment-naïve lung cancers from diverse patient populations (**Fig. 1a**; **Supplementary Fig. 1**).

This large international cohort enabled a comprehensive evaluation of ecDNA's association with

genomic alterations, as well as its prevalence across smoking status, biological sexes, ancestries,

and 18 areas defined primarily by country of residence. The dataset includes 871 LCINS,

sourced from the Sherlock-Lung[12] (n=817), the Environment And Genetics in Lung Cancer

Etiology (EAGLE[13], n=25), the Cancer Genome Atlas (TCGA[14],n=15), and other publicly

available cohorts (n=14)[15-19], representing patients from 17 geographical areas (**Supplementary**

Fig. 2a). Additionally, it comprises 345 LCSS, drawn from EAGLE[13] (n=221) and TCGA[14]

(n=83) and other publicly available cohorts (n=41)[15-19], spanning five geographical areas

(**Supplementary Fig. 2b**). The majority of tumors were LUAD (n=1,023; 84.1%) or LUSC

(n=67; 5.5%), with carcinoids (n=61; 5%, exclusively in LCINS) and other rarer histological

lung cancer subtypes (n=65; 5.3%) also represented. Amongst Sherlock-*Lung* LCINS, 250 were

exposed to secondhand tobacco smoke, while 208 had documented non-exposure (**Fig. 1a**).

To detect and reconstruct ecDNA across the cohort, a computational pipeline that accounts for

both tumor purity and ploidy was developed (**Methods**). A total of 231 samples harbored at least

one ecDNA (18.9%), with similar prevalence between LCINS (17%) and LCSS (23%; q-value:

0.21; **Fig. 1b**). No statistically significant differences were observed in the prevalence of ecDNA

across various histological subtypes, biological sexes, ancestries or age at diagnosis (**Fig. 1c, 1e**),

except for carcinoids, which showed a lower prevalence compared to LUAD in the overall

analysis (OR=0.07; q=0.04). ecDNA was present at similar rates across all geographic locations

(**Fig. 1d**; **Supplementary Fig. 2c-d**), as indicated by a one-versus-all multivariate logistic

regression for each country. Although there was no difference in ecDNA prevalence between

LCSS and LCINS (**Fig.1c**), in LUAD cases from the United States, ecDNA was more prevalent

in LCSS compared to LCINS (OR=2.26; p=0.03; **Supplementary Fig. 2e**). ecDNA was not

associated with tumor stage (**Supplementary Fig. 3a&b**), subsequent development of metastases

(**Supplementary Fig. 3c&d**) or passive smoking status (**Supplementary Fig. 3e**). These

findings remained consistent when analyses were restricted to only LUAD cases in either LCSS

or LCINS (**Supplementary Fig. 4**).

To examine the association between ecDNA and genomic features previously linked to genome

instability[20-23], we evaluated the relationship of ecDNA with whole-genome doubling (WGD),

chromothripsis, tumor mutational burden, mitochondrial copy number, overall genome instability

index (wGII), and telomere length tumor/normal ratio . Amongst these, only WGD showed a

significant association with ecDNA in both LCSS and LCINS (**Fig. 2a**), as determined by

logistic regression adjusted for age, sex, ancestry, histology, and tumor purity. Specifically,

ecDNA was present in 7.1% of LCINS and 8.9% of LCSS without WGD, but its prevalence

increased markedly in WGD-positive tumors from LCINS (24.9%, OR=4.0; q=1.24 x 10^{-8}) and

LCSS (30.3%, OR=5.85; q=8.06 x 10^{-4}). The same association was observed when only LUAD

was examined (**Supplementary Fig. 5a**). Moreover, this association was further validated in a

pan-cancer dataset of 974 samples with high-confidence WGD annotation[14], using logistic

regression adjusted for age, purity, sex, and tissue type (OR=2.81; q=1.63 x 10^{-6};

Supplementary Fig. 5b).

In addition to WGD, chromothripsis was the only other feature associated with ecDNA, but this

was exclusively observed in LCINS (OR=1.76; q=0.048; **Fig. 2a**). Prior studies have indicated

that chromothripsis may contribute to the formation of certain ecDNA[23]. In this study, 115 out of

151 LCINS with ecDNA (76.1%) and 60 out of 80 LCSS with ecDNA (75%) exhibited

chromothripsis (**Supplementary Tables 1-2**). Nevertheless, among samples where both ecDNA

and chromothripsis were detected, the overlap between ecDNA segments and chromothripsis

regions was observed in only 31 out of 115 LCINS and 4 out of 60 LCSS (OR=5.01, p=0.02

comparing LCINS *vs.* LCSS, **Supplementary Fig. 5*c***). These findings suggest that while the

majority of ecDNA emerges in genomically unstable lung cancers that also exhibit

chromothripsis, the actual genomic overlap between ecDNA and chromothripsis regions is

limited.

To evaluate the relationship between ecDNA presence and lung cancer driver genes, we analyzed

driver genes altered in at least 10 samples or with hotspot mutations occurring at the same

genomic position in at least three samples. Using multivariate logistic regression, adjusted for

age, sex, tumor purity, ancestry, and histology, no significant associations were found between

ecDNA and individual driver gene alterations (**Fig. 2*b***). However, hotspot analysis showed an

enrichment of *EGFR* L858R mutations (OR=1.78; q=0.05) and a depletion of *KRAS* G12V

mutations both in LCSS with ecDNA (OR=0.18; q=0.03; **Fig. 2*c***; **Supplementary Fig. 5*d***).

Mutational signature analysis provides insights into both the endogenous and exogenous

mutational processes that have been active throughout the lineage of a cancer genome[11]. We

investigated the association between ecDNA presence and mutational signatures. As expected[24],

multiple copy-number and structural variant signatures were associated with ecDNA presence

similarly in LCSS and LCINS (**Supplementary Fig. 6a-b**). Additionally, in LCINS, ecDNA-

positive samples were enriched for clock-like signatures (SBS1, SBS5, ID1, ID2) and, as

previously reported[19,25], APOBEC-associated signatures (SBS2, SBS13; **Supplementary Fig.**

6a). In LCSS, enrichment was observed only for SBS3 (**Supplementary Fig. 6b**), a signature

previously associated with homologous recombination deficiency[26].

Previous studies have shown that ecDNA can harbor oncogenes, regulatory elements, and

immunomodulatory genes in varying configurations[3,8,27,28]. In our cohort, 151 (17%) LCINS and

80 (23%) LCSS harbored ecDNA, with a total of 236 ecDNAs in LCINS and 123 in LCSS,

reflecting multiple ecDNAs in some tumors. Among these, ecDNA in LCINS were enriched with

oncogenes (p<0.001) and immunomodulatory genes (p=0.008) compared to LCSS (**Fig. 2d,**

Supplementary Fig. 7a-b). Further, ecDNA containing oncogenes exhibited elevated copy

numbers (CN) compared to other ecDNA without oncogenes in LCINS (p=1.5 x 10^{-3}) and LCSS

(5.8 x 10^{-3}), suggesting greater selective pressure for these ecDNAs (**Supplementary Fig. 7c**), as

previously reported in pan-cancer[8]. *MDM2* was the most frequently amplified oncogene on

ecDNA in both LCINS (*n*=36) and LCSS (*n*=5). Given its prominence, we examined the

relationship between ecDNA-driven *MDM2* amplification and *TP53* alterations, as MDM2

negatively regulates TP53 through ubiquitination[29]. We found that *MDM2* amplification on

ecDNA was significantly mutually exclusive with *TP53* alterations (OR = 7.51, p = 0.01). In

LCINS, the next four most prevalent oncogenes that did not co-occur with *MDM2* were *TERT*

(n=10), *CCND1* (*n*=8), *MYC* (*n*=8), and *EGFR* (*n*=7; **Supplementary Table 3**). In LCSS, no

oncogene besides *MDM2* was amplified on ecDNA in more than two samples (**Supplementary**

Table 4).

Prior pan-cancer studies have shown that cancers harboring ecDNA amplifications exhibit worse

overall survival[6-8]. To evaluate this association in the lung cancer cohort, survival analysis was

conducted on 693 LCINS and 333 LCSS with available stage and covariate data. Lung cancers

were categorized as harboring ecDNA, other focal amplifications, or no focal amplifications.

Cox multivariate analysis revealed that ecDNA-positive lung cancers had worse overall survival

than those without any focal amplifications in LCINS (HR=1.95; p=0.004; **Fig. 2*e***). However,

ecDNA did not confer a greater risk than other focal amplifications in either LCINS or LCSS

(**Fig. 2*e***).

In summary, we examined the full spectrum of ecDNA in the largest dataset of lung cancers to

date, from many geographical regions and ancestry, and with unique information on tobacco

smoking status. ecDNA was present in 17% of LCINS and 23% of LCSS. ecDNA in LCINS

were enriched with oncogenes and immunomodulatory genes in comparison to LCSS. In LCINS,

the *MDM2* locus was often amplified through ecDNA, while the *EGFR* L858R mutation showed

a weak association with ecDNA presence. No associations were identified between presence of

ecDNA and histological subtypes, biological sexes, smoking status, or ancestries. Nevertheless, a

strong association was observed between ecDNA and WGD, and this observation was validated

in an independent pan-cancer cohort. These results suggest that while ecDNA carrying

oncogenes could contribute to driving LCINS, most ecDNA are likely to be a byproduct of

genomic instability in treatment-naïve lung cancer.

METHODS

Criteria for sample inclusion

A total of 1217 samples were available for analysis, but one sample with unknown smoking

status was excluded, resulting in a total of 1216 samples across the cohort.

Determination of sample purity and ploidy

The Battenberg algorithm (v.2.2.9)[30] was used to determine the sample purity and ploidy. Any

somatic copy number variations (CNV) profile determined to have low-quality after manual

inspection underwent a refitting process. This process required new tumor purity and ploidy

inputs, either estimated by Ccube (v.1.0)[31] or recalculated from local copy number status. The

Battenberg refitting procedures were iteratively executed until the final CNV profile was

established and met the criteria of manual validation check.

ecDNA detection and characterization

CNVKit v.0.9.6[32] was run in tumor-normal mode to call somatic CNVs against the matched

normal whole-genome sequenced samples for each patient, using the log ratio of tumor to normal

read depths (logR) and the circular binary segmentation (CBS) method. Note that this method, by

default, assumes a sample purity of 1 and a sample ploidy of 2 as it does not correct for purity

and ploidy. This limitation can lead to high copy number segments being missed due to low

purity[33]. Due to the wide range of tumor purities in our dataset, consistent with prior reports in

201 lung cancer[33], we rescaled the total copy number estimate using the purity and ploidy estimated

202 from Battenberg according to the following equation:

$$Equation\ 1.1.\ TCN\ =\ \frac{(\psi\ *\ (2^{logR}))\ -\ 2\ *\ (1\ -\ \rho)}{\rho}$$

where ψ is the tumor ploidy, ρ is the tumor purity, and logR represents the log ratio of the tumor

read depth to the normal read depth (**Supplementary Fig. 8a**). These copy number calls were

used to identify putative 'seed' regions. We then used AmpliconArchitect v.1.3[10] to reconstruct

the architecture of the amplicons and AmpliconClassifier (v.1.3.1) to classify amplicons

according to the most probable mechanism of formation (ecDNA, BFB, complex non-cyclic, or

linear). Visual inspection and verification were performed for all ecDNA amplicon regions. This

pipeline allowed for samples with low purities to be utilized as it completely removed the effect

of tumor purity in ecDNA detection (**Supplementary Fig. 8b**). As a result, more ecDNA[+]

samples were detected (**Supplementary Fig. 8b**) and all 1216 samples could be used for ecDNA

analysis (**Supplementary Fig. 8c**). Of the additional ecDNA detected, eight of these were

EGFR-ecDNA (seven in LCINS and one in LCSS) and each of these ecDNA except for one also

harbored a driver mutation (**Supplementary Table 5**). Among the eight EGFR-ecDNA+

samples, five had RNA-seq data, which showed very high EGFR expression (**Supplementary**

Fig. 8d).

Determination of whole genome doubling status

As previously done[34], samples were considered to have a whole genome doubling event if the

major copy number was ≥3 for more than 50% of copy number segments.

Chromothripsis detection and characterization

To detect chromothripsis regions in genomes, we applied the computational algorithm ShatterSeek[35], which employs a set of statistical criteria to identify chromothripsis given an input of copy number and structural variants. We used the most stringent criteria corresponding to high confidence chromothripsis calls. Specifically, these criteria were: *(i)* a cluster of structural variants (>6 DUP/DEL/h2hINV/t2tINV), *(ii)* oscillating CNV between two states (>7 CNV events), *(iii)* chromosomal enrichment and distribution of DNA breakpoints (p-value<0.05), *(iv)* randomness of fragment joins (p-value>0.05) and/or ≥4 inter-chromosomal rearrangements between multiple chromosomes. The input CNV calls were generated using CNVkit in tumor-normal mode with default parameters, and the input structural variants calls were generated using a union of Meerkat (v.0.189) and Manta (v.1.6.0) calls with the recommended filtering.

Association of ecDNA with demographic and clinical features

Demographic and clinical features where information was available for all samples (age, sex, ancestry, and tumor histologyand purity) served as input variables in a multivariate logistic regression model with ecDNA presence or absence as the dependent outcome variable, and these models were constructed separately for LCINS and LCSS. In addition, separate models for passive smoking, tumor stage, and presence of metastases were run along with the rest of the variables in both LCINS and LCSS.

Determination of telomere length tumor/normal lung tissue ratio

We estimated telomere length (TL) in kilobases using TelSeq (v.0.0.2)[36]. As was previously

done[37], we used seven as the threshold for the number of TTAGGG/CCCTAA repeats in a read

for the read to be considered telomeric. The TelSeq calculation was done individually for each

read group within a sample, and the total number of reads in each read group was used as weight

to calculate the average TL for each sample. The telomere length ratio ($\log_2$ scale) between

tumor and normal tissue was then calculated for each patient and served as the independent

variable in multivariable logistic regressions.

Association of mutational signatures and ecDNA status

Mutational signature attributions were determined as was previously done, based on the

methodology in Díaz-Gay et. al[37]. SigProfilerMatrixGenerator[38] was used to create the

mutational matrix for all types of somatic mutations. The deconvolution of mutational signatures

was performed by SigProfilerExtractor (v.1.1.3)[39] using the whole-genome sequencing setting

and the COSMIC mutational signatures (v3.2) as reference. Signature extractions were

performed separately for LCINS and LCSS. Signature attributions were input into a multivariate

logistic regression model as either present or absent if the signature was present in less than 50%

of the samples. If the signature was present in ≥50% samples (*e.g.*, SBS1 and SBS5), it was input

as either above median or below median. For SBS1 and SBS5, the signature attributions were

also input as continuous numeric variables to confirm the significant associations found with

inputting them as above median or below median.

Association of driver genes and ecDNA status

The IntOGen pipeline (v.2020.02.0123)[40], which combines seven state-of-the-art computational

methods, was employed with default parameters to detect signals of positive selection in the

mutational patterns of driver genes across the cohort. The status of each driver gene (altered or

unaltered) was then input into a logistic regression model with covariates age, sex, and tumor

purity, and ecDNA status was the output. These sets of genes were considered altered if there

was a focal deletion overlapping the gene, an inactivating (stop or missense) mutation, or an

activating gain of function mutation.

Association of driver mutations and ecDNA status

To identify driver mutations within the set of identified driver genes, we implemented a rigorous

and multifaceted strategy, considering multiple criteria: *(i)* the presence of truncating mutations

specifically in genes annotated as tumor suppressors, *(ii)* the recurrence of missense mutations in

a minimum of 3 samples, *(iii)* mutations designated as "Likely drivers" with a boostDM score[41]

exceeding 0.5, *(iv)* mutations categorized as either "Oncogenic" or "Likely Oncogenic" based on

the criteria established by OncoKB[42], an expert-guided precision oncology knowledge base, *(v)*

mutations previously recognized as drivers in the TCGA MC3 drivers paper[43], and *(vi)* missense

mutations characterized as "likely pathogenic" in genes that are annotated as tumor suppressors,

as described in Cheng et al.[44] Any mutation meeting one or more of these criteria was recognized

as a potential driver mutation.

Calculation of genome instability index score

To calculate the genome instability index score (wGII), the method devised in Bailey et al.[8] was

used. The overall ploidy was calculated as the length-weighted average of all copy number

segments. The proportion of segments that were aberrant (where total copy number was greater

than the ploidy or less than the ploidy) was computed for each chromosome, these proportions

were summed, and the total was divided by 22.

Identifying regulatory regions on ecDNA

A set of lung enhancer regions were downloaded from EnhancerAtlas[45], and each ecDNA region

was overlapped with these enhancer regions with the requirement that the entire enhancer had to

be fully contained (100% overlap) on the ecDNA. A set of lung cancer promoter regions was

extracted using a R script which downloaded regions 1000 base pairs upstream of transcription

start sites for all genes.

Defining a set of immunomodulatory genes

ecDNAs were considered immunomodulatory if a gene from the significant gene sets mapped to

an ecDNA that did not contain an oncogene, and that significant gene set had an

immunomodulatory function (GO terms: 0006968, 0002228, 0042267, 0001906, 0001909,

0002698, 0001910, 0031341, 0002367, 0002695, 0050866, 0051250, 0050777), following the

procedure in Bailey et. al[8]. This resulted in a set of 376 genes.

Mitochondrial copy number estimation

Mitochondrial copy number was estimated per sample using the ratio of mitochondrial read

depth to nuclear genome depth, adjusted for tumor ploidy and purity, following the procedure in

Reznik *et al.*[46]

Statistical analysis

All statistical analyses and graphic displays were performed using the R software v4.2.3 1011

(https://www.r-project.org/). Logistic regression with adjustment for covariates was used for the

enrichment analyses of categorical variables. For the comparison of numerical variables across

groups, we used non-parametric Mann-Whitney (Wilcoxon rank sum) tests. P-values <0.05 were

considered statistically significant. If multiple hypothesis testing was required, we used a false-

discovery rate correction based on the Benjamini-Hochberg method and reported FDR. FDR

<0.05 were considered statistically significant.

Figure 1: Landscape of extrachromosomal DNA in lung cancer across histologies, sex, ancestry, countries, and smoking status. *a)* Epidemiological features of the LCINS (top) and LCSS (bottom) cohorts. Total number of samples in each category is indicated. *b)* The number and proportion of tumors harboring ecDNA among LCINS and LCSS, stratified by histology. Samples with 1 or more ecDNA were classified as ecDNA$^+$, and samples with no ecDNA were classified as ecDNA$^-$. *c)* Forest plot of the full cohort (LCINS and LCSS) from a multivariate logistic regression analysis of key epidemiological and clinical features and the presence of ecDNA in the tumors. *d)* Volcano plot showing $\log_2$ odds ratio (x-axis) and $\log_{10}$ FDR corrected q-value (y-axis) for a one-vs-all statistical comparison of ecDNA prevalence by region. *e)* Forest plots for multivariate logistic regression models testing key epidemiological and clinical features with the presence of ecDNA in the tumors, stratified by smoking status. NA/EU: North America or Europe region, AS: Asia region EAS: East Asian genetic ancestry super-sample, EUR: European genetic ancestry super-sample, LUAD: lung adenocarcinoma, LUSC: lung squamous cell carcinoma.

Figure 2. Genomic and clinical characteristics associated with ecDNA. *a)* Volcano plot of logistic regression models of genomic features in association with ecDNA status for LCINS (left) and LCSS (right). *b)* Volcano plot of a logistic regression model of driver gene alterations in a sample in association with ecDNA status for LCINS (left) and LCSS (right). *c)* Volcano plot of a logistic regression model of hotspot driver mutations in a sample in association with ecDNA status for LCINS (left) and LCSS (right). In all volcano plots, the x-axes reflect the $\log_2$ odds ratio, and the y-axes correspond to the $\log_{10}$ FDR q-value. An FDR q-value threshold of 0.05 is

indicated with the dashed red line, an FDR q-value threshold of 0.01 is indicated with the dashed

blue line. *d)* Bar plot indicating the proportion of ecDNA containing a particular genomic

element in LCINS vs LCSS. *e)* Kaplan–Meier survival curves for 5-year overall survival

stratified by the mode of amplification status for lung cancers in LCINS (left) and LCSS (right).

P-values for significance and HRs of the difference were calculated using two-sided Cox

proportional-hazards regression with adjustment for age, sex, tumor stage, ancestry and histology

and are indicated within each plot.

SUPPLEMENTARY FIGURE LEGENDS

Supplementary Fig 1. Additional cohort information. Sankey diagrams showing the numbers

of samples with clinical variables (stage, metastasis) as well as survival data available for *a)*

LCINS and *b)* LCSS

Supplementary Fig 2. Distribution of ecDNA harboring lung cancers in LCINS and LCSS

by country. Maps showing the worldwide prevalence of lung cancers harboring ecDNA

(ecDNA$^+$) by country for the *a)* LCINS cohort and *b)* LCSS cohort. Volcano plot of a

multivariate one-vs-all logistic regression model for each country, with ecDNA status as the

outcome, for *c)* LCINS and *d)* LCSS. *e)* Bar plot depicting the prevalence of ecDNA in LCINS

versus LCSS from the United States. Statistical results from a logistic regression model are

shown above.

Supplementary Fig 3. Analysis of the association between stage, metastasis, and passive

smoking with ecDNA status in all LCINS and LCSS. Forest plots of logistic regression model

with ecDNA status as the outcome and the variables for stage in **a)** LCINS and **b)** LCSS;

metastasis in **c)** LCINS and **d)** LCSS; and passive smoking in **e)** LCINS. The total number of

samples with information for all variables is indicated above each plot.

Supplementary Fig 4. Analysis of the association between stage, metastasis, and passive

smoking with ecDNA status in lung adenocarcinomas. Forest plots of logistic regression

model with ecDNA status as the outcome and the variables for stage in **a)** LCINS and **b)** LCSS;

metastasis in **c)** LCINS and **d)** LCSS; and passive smoking in **e)** LCINS. The total number of

samples with information for all variables is indicated above each plot.

Supplementary Fig 5. Genomic features associated with the presence of ecDNA. *a)* Volcano

plot of logistic regression models of genomic features in association with ecDNA status for lung

adenocarcinoma in LCINS (left) and lung adenocarcinoma in LCSS (right). In all volcano plots,

the x-axes reflect the $\log_2$ odds ratio, and the y-axes correspond to the $\log_{10}$ FDR q-value. An

FDR q-value threshold of 0.05 is indicated with the dashed red line, an FDR q-value threshold of

0.01 is indicated with the dashed blue line. *b)* Forest plot of multivariate logistic regression

modeling the presence or absence of ecDNA in relation to whole-genome doubling in a pan-

cancer dataset comprising four different tumor types from the PCAWG project, adjusted for the

covariates of age, purity, sex, and tissue type. All significant associations are indicated in red. *c)*

Bar plot indicating the proportion of samples that contained overlapping chromothripsis and

ecDNA regions (red) and the proportion of all samples that did not contain any overlap (gray) for

all samples that had both an ecDNA and chromothripsis event detected. *d)* Bar plot showing the

proportion of samples with an *EGFR* L858R mutation, stratified by ecDNA status.

Supplementary Fig 6. Association of mutational signatures with ecDNA status. Volcano plots of logistic regression models of the activities of Single base substitution (SBS), Indel (ID), copy number (CN) and structural variant (SV) signatures in association with ecDNA in lung cancers of *a)* LCINS and *b)* LCSS. The logistic regression models were adjusted for age, sex, ancestry, histology, and purity. In all volcano plots, the x-axes reflect the $\log_2$ odds ratio, and the y-axes correspond to the $\log_{10}$ FDR q-value. An FDR q-value threshold of 0.05 is indicated with the dashed red line, an FDR q-value threshold of 0.01 is indicated with the dashed blue line.

Supplementary Fig 7. Genomic content of ecDNA

a) Upset plot indicating the number of ecDNA with oncogenes, immunomodulatory genes, promoter or enhancer regions, other genes that are not oncogenes, or unidentified genomic elements (Unknown) or any combination thereof for LCINS and LCSS. **b)** Bar plot indicating the proportion of ecDNA containing a particular genomic element in LCINS vs LCSS. **c)** $\log_2$ median copy number (y-axis) for each class of ecDNA (x-axis) in LCINS and LCSS.

Supplementary Fig 8. Highly sensitive pipeline for ecDNA detection that accounts for tumor purity and ploidy.

a) Schematic depicting the methodology for the rescaled pipeline that considers tumor purity and ploidy. The logR track for the region containing the *EGFR* gene on chr7 and the purity and ploidy computed from Battenberg is indicated in one tumor as example (left). The exact location of the *EGFR* gene is shown by the dashed red lines. The equation that simultaneously accounts

for sample purity and ploidy when computing the total copy number is shown, as well as the total

copy number estimate when assuming a sample purity of 1.0 and ploidy of 2.0 (Uncorrected CN)

vs. considering the purity and ploidy (Corrected CN). The Sashimi plot indicates the structural

variants (colored arcs) and smoothed coverage around the *EGFR* region, which was annotated as

ecDNA by AmpliconClassifier. **b)** Distribution of tumor purities for samples with an ecDNA

annotation for both the default (left) and rescaled (right) pipelines. P-value for a Mann-Whitney

test between ecDNA$^+$ and ecDNA$^-$ samples is indicated. The total number of samples in each

group is also indicated. **c)** The ecDNA prevalence (y-axis), which is the proportion of ecDNA$^+$

samples over the total number of samples, for all possible tumor purity cutoffs applied to

samples (x-axis). The total number of samples available for analysis after each tumor purity

cutoff is applied is also indicated. **d)** Gene expression (log$_2$ CPM) of the EGFR gene for all

samples with RNA-seq data available. LCINS samples with EGFR-ecDNA are indicated in red.

SUPPLEMENTARY MATERIAL LEGENDS

Supplementary Table 1. Co-occurrence of ecDNA genetic cargo in LCINS

Supplementary Table 2. Co-occurrence of ecDNA genetic cargo in LCSS

Supplementary Table 3. Recurrently amplified oncogenes on ecDNA in LCINS across

unique genomic loci

Supplementary Table 4. Recurrently amplified oncogenes on ecDNA in LCSS across

unique genomic loci

Supplementary Table 5. Additional *EGFR*-ecDNA detected by modified pipeline

Ethics declarations

Since the National Cancer Institute only received de-identified samples and data from collaborating centers, had no direct contact or interaction with the study participants, and did not use or generate identifiable private information, Sherlock-*Lung* has been determined to constitute "Not Human Subject Research (NHSR)" based on the federal Common Rule (45 CFR 46; https://www.ecfr.gov/cgi-bin/ECFR?page=browse).

DATA AVAILABILITY

Normal and tumor-paired CRAM files for the WGS subjects of the Sherlock-Lung study and the EAGLE study have been deposited in dbGaP under the accession numbers phs001697.v2.p1 and phs002992.v1.p1, respectively.

Author Contributions

Conceptualization, MTL, TZ, LBA; Methodology, AK, TZ; Formal Analysis, AK, TZ, MDG, JL, WZ; Data Curation, TZ, PH, JM, CH, MM, OWL, MAN, NR, RH, SY, QL; Writing – Original Draft, AK; Writing – Review & Editing, LBA, TZ, PHH, MTL; Visualization: AK; Supervision, LBA, TZ, MTL.

ACKNOWLEDGEMENTS

This work was supported by the Intramural Research Program of the National Cancer Institute, US National Institute of Health (NIH) (project ZIACP101231 to MTL), and by the Anne Wojcicki Foundation (Grant Number: LC009). Part of this work was supported by the US National Institute of Health grants R01ES032547, R01CA269919, P01CA281819, and U01CA290479 to LBA as well as by LBA's Packard Fellowship for Science and Engineering. MD-G was awarded with a fellowship within the "Generación D" initiative, Red.es, Ministerio para la Transformación Digital y de la Función Pública, for talent attraction (C005/24-ED CV1), funded by the European Union NextGenerationEU funds, through PRTR. The research in this study was also supported by UC San Diego Sanford Stem Cell Institute. This work utilized the computational resources of the NIH HPC Biowulf cluster (https://hpc.nih.gov). The funders had no roles in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

COMPETING INTERESTS

LBA is a co-founder, CSO, scientific advisory member, and consultant for io9, has equity and receives income. The terms of this arrangement have been reviewed and approved by the University of California, San Diego in accordance with its conflict of interest policies. LBA is a compensated member of the scientific advisory board of Inocras. LBA's spouse is an employee of Hologic, Inc. LBA declares U.S. provisional applications with serial numbers: 63/289,601; 63/269,033; 63/366,392; 63/412,835 as well as international patent application PCT/US2023/010679. LBA is also an inventor of a US Patent 10,776,718 for source identification by non-negative matrix factorization. LBA and MD-G further declare a European patent application with application number EP25305077.7. Soo-Ryum Yang has received

REFERENCES

Benner, S. E., Wahl, G. M. & Von Hoff, D. D. Double minute chromosomes and homogeneously staining regions in tumors taken directly from patients versus in human tumor cell lines. *Anticancer Drugs* **2**, 11-25 (1991). https://doi.org/10.1097/00001813-199102000-00002

Nielsen, J., Walsh, J. T., Degen, D. R., Drabek, S. M. & McGill, J. R. Evidence of gene amplification in the form of double minute chromosomes is frequently observed in lung cancer. *Cancer Genetics* **65**, 120-124 (1993). https://doi.org/https://doi.org/10.1016/0165-4608(93)90219-C

Hung, K. L. *et al.* Coordinated inheritance of extrachromosomal DNAs in cancer cells. *Nature* **635**, 201-209 (2024). https://doi.org/10.1038/s41586-024-07861-8

Lange, J. T. *et al.* The evolutionary dynamics of extrachromosomal DNA in human cancers. *Nat Genet* **54**, 1527-1533 (2022). https://doi.org/10.1038/s41588-022-01177-x

Wu, S. *et al.* Circular ecDNA promotes accessible chromatin and high oncogene expression. *Nature* **575**, 699-703 (2019). https://doi.org/10.1038/s41586-019-1763-5

Kim, H. *et al.* Mapping extrachromosomal DNA amplifications during cancer progression. *Nat Genet* **56**, 2447-2454 (2024). https://doi.org/10.1038/s41588-024-01949-7

Kim, H. *et al.* Extrachromosomal DNA is associated with oncogene amplification and poor outcome across multiple cancers. *Nat Genet* **52**, 891-897 (2020). https://doi.org/10.1038/s41588-020-0678-2

Bailey, C. *et al.* Origins and impact of extrachromosomal DNA. *Nature* **635**, 193-200 (2024). https://doi.org/10.1038/s41586-024-08107-3

Tang, J. *et al.* Enhancing transcription-replication conflict targets ecDNA-positive cancers. *Nature* **635**, 210-218 (2024). https://doi.org/10.1038/s41586-024-07802-5

Deshpande, V. *et al.* Exploring the landscape of focal amplifications in cancer using AmpliconArchitect. *Nat Commun* **10**, 392 (2019). https://doi.org/10.1038/s41467-018-08200-y

Alexandrov, L. B. *et al.* The repertoire of mutational signatures in human cancer. *Nature* **578**, 94-101 (2020). https://doi.org/10.1038/s41586-020-1943-3

Landi, M. T. *et al.* Tracing Lung Cancer Risk Factors Through Mutational Signatures in Never-Smokers. *Am J Epidemiol* **190**, 962-976 (2021). https://doi.org/10.1093/aje/kwaa234

Consonni, D. *et al.* Outdoor particulate matter (PM10) exposure and lung cancer risk in the EAGLE study. *PLoS One* **13**, e0203539 (2018). https://doi.org/10.1371/journal.pone.0203539

Consortium, I. T. P.-C. A. o. W. G. Pan-cancer analysis of whole genomes. *Nature* **578**, 82-93 (2020). https://doi.org/10.1038/s41586-020-1969-6

Lee, J. J. *et al.* Tracing Oncogene Rearrangements in the Mutational History of Lung Adenocarcinoma. *Cell* **177**, 1842-1857 e1821 (2019). https://doi.org/10.1016/j.cell.2019.05.013

Imielinski, M. *et al.* Mapping the hallmarks of lung adenocarcinoma with massively parallel sequencing. *Cell* **150**, 1107-1120 (2012). https://doi.org/10.1016/j.cell.2012.08.029

17 Lee, J. K. *et al.* Clonal History and Genetic Predictors of Transformation Into Small-Cell
Carcinomas From Lung Adenocarcinomas. *J Clin Oncol* **35**, 3065-3074 (2017).
https://doi.org/10.1200/JCO.2016.71.9096
18 Leong, T. L. *et al.* Deep multi-region whole-genome sequencing reveals heterogeneity
and gene-by-environment interactions in treatment-naive, metastatic lung cancer.
Oncogene **38**, 1661-1675 (2019). https://doi.org/10.1038/s41388-018-0536-1
19 Bergstrom, E. N. *et al.* Mapping clustered mutations in cancer reveals APOBEC3
mutagenesis of ecDNA. *Nature* **602**, 510-517 (2022). https://doi.org/10.1038/s41586-
022-04398-6
20 Chen, J. *et al.* MYC-driven increases in mitochondrial DNA copy number occur early
and persist throughout prostatic cancer progression. *JCI Insight* **8** (2023).
https://doi.org/10.1172/jci.insight.169868
21 Bielski, C. M. *et al.* Genome doubling shapes the evolution and prognosis of advanced
cancers. *Nat Genet* **50**, 1189-1195 (2018). https://doi.org/10.1038/s41588-018-0165-1
22 O'Sullivan, R. J. & Karlseder, J. Telomeres: protecting chromosomes against genome
instability. *Nat Rev Mol Cell Biol* **11**, 171-181 (2010). https://doi.org/10.1038/nrm2848
23 Shoshani, O. *et al.* Chromothripsis drives the evolution of gene amplification in cancer.
Nature **591**, 137-141 (2021). https://doi.org/10.1038/s41586-020-03064-z
24 Steele, C. D. *et al.* Signatures of copy number alterations in human cancer. *Nature* **606**,
984-991 (2022). https://doi.org/10.1038/s41586-022-04738-6
25 Hadi, K. *et al.* Distinct Classes of Complex Structural Variation Uncovered across
Thousands of Cancer Genome Graphs. *Cell* **183**, 197-210 e132 (2020).
https://doi.org/10.1016/j.cell.2020.08.006
26 Davies, H. *et al.* HRDetect is a predictor of BRCA1 and BRCA2 deficiency based on
mutational signatures. *Nat Med* **23**, 517-525 (2017). https://doi.org/10.1038/nm.4292
27 Hung, K. L. *et al.* ecDNA hubs drive cooperative intermolecular oncogene expression.
Nature **600**, 731-736 (2021). https://doi.org/10.1038/s41586-021-04116-8
28 Zhu, Y. *et al.* Oncogenic extrachromosomal DNA functions as mobile enhancers to
globally amplify chromosomal transcription. *Cancer Cell* **39**, 694-707.e697 (2021).
https://doi.org/10.1016/j.ccell.2021.03.006
29 Chene, P. Inhibiting the p53-MDM2 interaction: an important target for cancer therapy.
Nat Rev Cancer **3**, 102-109 (2003). https://doi.org/10.1038/nrc991
30 Nik-Zainal, S. *et al.* The life history of 21 breast cancers. *Cell* **149**, 994-1007 (2012).
https://doi.org/10.1016/j.cell.2012.04.023
31 Dharanipragada, P. *et al.* Blocking Genomic Instability Prevents Acquired Resistance to
MAPK Inhibitor Therapy in Melanoma. *Cancer Discov* **13**, 880-909 (2023).
https://doi.org/10.1158/2159-8290.CD-22-0787
32 Talevich, E., Shain, A. H., Botton, T. & Bastian, B. C. CNVkit: Genome-Wide Copy
Number Detection and Visualization from Targeted DNA Sequencing. *PLoS Comput*
Biol **12**, e1004873 (2016). https://doi.org/10.1371/journal.pcbi.1004873
33 Aran, D., Sirota, M. & Butte, A. J. Systematic pan-cancer analysis of tumour purity. *Nat*
Commun **6**, 8971 (2015). https://doi.org/10.1038/ncomms9971
34 Zhang, T. *et al.* Deciphering lung adenocarcinoma evolution and the role of LINE-1
retrotransposition. *bioRxiv* (2025). https://doi.org/10.1101/2025.03.14.643063

35   Cortes-Ciriano, I. *et al.* Comprehensive analysis of chromothripsis in 2,658 human
582   cancers using whole-genome sequencing. *Nat Genet* **52**, 331-341 (2020).
583   https://doi.org/10.1038/s41588-019-0576-7
584   36   Ding, Z. *et al.* Estimating telomere length from whole genome sequence data. *Nucleic*
585   *Acids Res* **42**, e75 (2014). https://doi.org/10.1093/nar/gku181
586   37   Diaz-Gay, M. *et al.* The mutagenic forces shaping the genomic landscape of lung cancer
587   in never smokers. *medRxiv* (2024). https://doi.org/10.1101/2024.05.15.24307318
588   38   Khandekar, A. *et al.* Visualizing and exploring patterns of large mutational events with
589   SigProfilerMatrixGenerator. *BMC Genomics* **24**, 469 (2023).
590   https://doi.org/10.1186/s12864-023-09584-y
591   39   Islam, S. M. A. *et al.* Uncovering novel mutational signatures by de novo extraction with
592   SigProfilerExtractor. *Cell Genom* **2**, 100179 (2022).
593   https://doi.org/10.1016/j.xgen.2022.100179
594   40   Gonzalez-Perez, A. *et al.* IntOGen-mutations identifies cancer drivers across tumor types.
595   *Nat Methods* **10**, 1081-1082 (2013). https://doi.org/10.1038/nmeth.2642
596   41   Muinos, F., Martinez-Jimenez, F., Pich, O., Gonzalez-Perez, A. & Lopez-Bigas, N. In
597   silico saturation mutagenesis of cancer genes. *Nature* **596**, 428-432 (2021).
598   https://doi.org/10.1038/s41586-021-03771-1
599   42   Chakravarty, D. *et al.* OncoKB: A Precision Oncology Knowledge Base. *JCO Precis*
600   *Oncol* **2017** (2017). https://doi.org/10.1200/PO.17.00011
601   43   Bailey, M. H. *et al.* Comprehensive Characterization of Cancer Driver Genes and
602   Mutations. *Cell* **174**, 1034-1035 (2018). https://doi.org/10.1016/j.cell.2018.07.034
603   44   Cheng, F. *et al.* Comprehensive characterization of protein-protein interactions perturbed
604   by disease mutations. *Nat Genet* **53**, 342-353 (2021). https://doi.org/10.1038/s41588-020-
605   00774-y
606   45   Gao, T. & Qian, J. EnhancerAtlas 2.0: an updated resource with enhancer annotation in
607   586 tissue/cell types across nine species. *Nucleic Acids Res* **48**, D58-D64 (2020).
608   https://doi.org/10.1093/nar/gkz980
609   46   Reznik, E. *et al.* Mitochondrial DNA copy number variation across human cancers. *Elife*
610   **5** (2016). https://doi.org/10.7554/eLife.10769
611

Fig 1.

a

Never-smokers (n=871)

| Histology | Ancestry | Region | Sex | Passive smoking |

LUSC n = 31 · LUAD n = 737 · Carcinoid n = 61 · Other n = 42
EUR n = 488 · EAS n = 349 · Other n = 34
NA/EU n = 541 · AS n = 309 · Other n = 21
Male n = 183 · Female n = 688
YES n = 250 · NO n = 208 · Unknown n = 413

Smokers (n=345)

| Histology | Ancestry | Region | Sex |

LUSC n = 36 · LUAD n = 286 · Other n = 23
EUR n = 329 · EAS n = 11 · Other n = 5
NA/EU n = 308 · Unknown. n = 37
Male n = 259 · Female n = 86

b

Never-smokers / Smokers

Proportion of tumors

ecDNA count
≥1 (ecDNA+)
0 (ecDNA−)

Never-smokers: Total (n=871) 17%/83%, LUAD (n=737) 18%/82%, LUSC (n=31) 23%/77%, Carcinoid (n=61) 2%/98%, Other (n=42) 21%/79%

Smokers: Total (n=345) 23%/77%, LUAD (n=286) 21%/79%, LUSC (n=36) 36%/64%, Other (n=23) 30%/70%

c

All 1216 samples

Smoker (Ref=Never−Smoker) OR = 1.43, q= 0.21
Sex:Female (Ref=Male) OR = 0.75, q= 0.24
Histology:Other (Ref=LUAD) OR = 1.25, q= 0.57
Histology:LUSC (Ref=LUAD) OR = 1.21, q= 0.57
Histology:Carcinoid (Ref=LUAD) OR = 0.07, q= 0.04
Ancestry:Other (Ref=EUR) OR = 1.22, q= 0.62
Ancestry:EAS (Ref=EUR) OR = 1.26, q= 0.33
Age OR = 1.01, q= 0.57

log2(Odds Ratio)

d

ecDNA status by region (All 1216 samples)
Adjusted by age, sex, ancestry, histology, and smoking

−log10(FDR)

Germany · Hong Kong S.A.R. · United States of America · South Korea · Poland · France · USA · Canada · Czech Republic · Italy · Taiwan · Russia · Romania · Republic of Serbia

log2(Odds Ratio)

e

Never-smokers (n=871)

Sex:Female (Ref=Male) OR = 0.6, q= 0.06
Histology:Other (Ref=LUAD) OR = 1.4, q= 0.58
Histology:LUSC (Ref=LUAD) OR = 0.89, q= 0.81
Histology:Carcinoid (Ref=LUAD) OR = 0.09, q= 0.06
Ancestry:Other (Ref=EUR) OR = 1.2, q= 0.77
Ancestry:EAS (Ref=EUR) OR = 1.36, q= 0.24
Age OR = 1.01, q= 0.5

log2(Odds Ratio)

Smokers (n=345)

Sex:Female (Ref=Male) OR = 1.02, q= 0.99
Histology:Other (Ref=LUAD) OR = 0.92, q= 0.99
Histology:LUSC (Ref=LUAD) OR = 1.3, q= 0.99
Ancestry:Other (Ref=EUR) OR = 1.46, q= 0.99
Ancestry:EAS (Ref=EUR) OR = 0.63, q= 0.99
Age OR = 1, q= 0.99

log2(Odds Ratio)

Fig 2.

Supplementary Data Fig. 1

a

Never-smokers (n=871) – Clinical Data

Stage   Metastasis   Overall Survival

b

Smokers (n=345) – Clinical Data

Stage   Metastasis   Overall Survival

Supplementary Data Fig. 2

Supplementary Data Fig. 3

Supplementary Data Fig. 4

Supplementary Data Fig. 5

a
Genomic features associated with ecDNA

Never-smoker LUAD (n=737)
Adjusted by age, sex, ancestry, histology & purity

- WGD
- Chromothripsis
- TMB
- MT Copy Number
- Telomere T/N Ratio

Y-axis: −log10(FDR)
X-axis: log2(Odds Ratio)

Depleted in ecDNA+ ← → Enriched in ecDNA+

Smoker LUAD (n=286)
Adjusted by age, sex, ancestry, histology & purity

- WGD
- Telomere T/N Ratio
- TMB
- MT Copy Number
- Chromothripsis

Y-axis: −log10(FDR)
X-axis: log2(Odds Ratio)

Depleted in ecDNA+ ← → Enriched in ecDNA+

b
Pan-cancer analysis of ecDNA status
n=974

Purity	OR = 28.97, q= 3.77e−04
WGD (Ref= No WGD)	OR = 2.81, q= 1.63e−06
Sex:Female (Ref=Male)	OR = 1.22, q= 6.46e−01
Ancestry:EUR (Ref=Other)	OR = 1.2, q= 6.53e−01
Tissue:Breast (Ref=Bladder)	OR = 1.11, q= 7.69e−01
Tissue:Head/Neck (Ref=Bladder)	OR = 1.09, q= 7.69e−01
Age	OR = 1.01, q= 1.55e−01
Ancestry:ASN (Ref=Other)	OR = 0.7, q= 7.26e−01
Tissue:Stomach (Ref=Bladder)	OR = 0.69, q= 6.46e−01
Tissue:Uterine (Ref=Bladder)	OR = 0.58, q= 2.15e−01
Tissue:Skin (Ref=Bladder)	OR = 0.57, q= 1.91e−01

X-axis: log2(Odds Ratio)

c
Chromothripsis and ecDNA co-occurrence
OR=5.01, p=0.02, Adjusted by age, sex, ancestry, histology, purity

Never-Smoker (n=115): ecDNA & chromothripsis overlapping event 27%, distinct events 73%
Smoker (n=60): ecDNA & chromothripsis overlapping event 6.7%, distinct events 93.3%

Y-axis: Percentage

Annotation
- ecDNA & chromothripsis, overlapping event (red)
- ecDNA & chromothripsis, distinct events (grey)

d
EGFR L858R mutation in never−smokers
OR=1.78, q=0.05, Adjusted by age, sex, ancestry, histology, purity

Non-Mutated (n=678): ecDNA+ 15.5%, ecDNA− 84.5%
Mutated (n=192): ecDNA+ 24%, ecDNA− 76%

Y-axis: Percentage

ecDNA.
- ecDNA+ (red)
- ecDNA− (grey)

Mutational signatures associated with ecDNA

a

Supplementary Data Fig. 7

Supplementary Data Fig. 8

a

NSLC-0841-T01

logR = 0.95

Purity: 0.39
Ploidy: 3.86

$$TCN = \frac{(\psi * (2^{logR})) - 2 * (1 - \rho)}{\rho}$$

Uncorrected CN: ~3.88
Corrected CN: ~15.6

NSLC-0841-T01

Forward–Forward (FF) — Inversion
Reverse–Reverse (RR) — Inversion

EGFR
chr7

b

Default Pipeline

P = 2.5e−07

ecDNA−
n=1004

ecDNA+
n=213

ecDNA Status

Rescaled Pipeline

P = 0.45

ecDNA−
n=963

ecDNA+
n=254

ecDNA Status

c

ecDNA Prevalence vs Tumor Purity Cutoff (n=1216)

n= 1216   n= 1216   n= 965   n= 743   n= 548   n= 361   n= 233   n= 131   n= 68   n= 20

No cutoff   >= 0.1   >= 0.2   >= 0.3   >= 0.4   >= 0.5   >= 0.6   >= 0.7   >= 0.8   >= 0.9

Tumor Purity Cutoff

d

NSLC−0841−T01
NSLC−0366−T01
NSLC−0752−T01
NSLC−0607−T01
NSLC−0859−T01

EGFR Classification

EGFR−ecDNA
EGFR−non−ecDNA

EGFR gene