Examining the Role of Extrachromosomal DNA in 1,216 Lung Cancers

Azhar Khandekar[1,2,3,4], Phuc H. Hoang[1], Jens Luebeck[5], Marcos Díaz-Gay[2,3,4,6], Wei Zhao[1], John P. McElderry[1], Caleb Hartman[1], Mona Miraftab[1], Olivia W. Lee[1], Kara M. Barnao[1], Erik N. Bergstrom[2,3,4], Yang Yang[7], Martin A. Nowak[8,9], Nathaniel Rothman[1], Robert Homer[10], Soo-Ryum Yang[11], Qing Lan[1], David C. Wedge[12,13], Lixing Yang[7,14,15], Stephen J. Chanock[1], Tongwu Zhang[1*], Ludmil B. Alexandrov[2,3,4,16*], Maria Teresa Landi[1*]

[1]Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD, USA, [2]Department of Cellular and Molecular Medicine, University of California San Diego, La Jolla, CA, USA, [3]Department of Bioengineering, University of California San Diego, La Jolla, CA, USA, [4]Moores Cancer Center, University of California San Diego, La Jolla, CA, USA, [5]Department of Computer Science, University of California San Diego, La Jolla, CA, USA, [6]Digital Genomics Group, Structural Biology Program, Spanish National Cancer Research Center (CNIO), Madrid, Spain, [7]Ben May Department for Cancer Research, The University of Chicago, Chicago, IL, USA, [8]Department of Mathematics, Harvard University, Cambridge, MA, USA, [9]Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA, [10]Department of Pathology, Yale School of Medicine, New Haven, CT, USA, [11]Department of Pathology and Laboratory Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA, [12]Manchester Cancer Research Centre, The University of Manchester, Manchester, UK, [13]Manchester NIHR Biomedical Research Centre, Manchester, UK, [14]Department of Human Genetics, The University of Chicago, Chicago, IL, USA, [15]The University of Chicago Medicine Comprehensive Cancer Center, The University of Chicago, Chicago, IL, USA, [16]Sanford Stem Cell Institute, University of California San Diego, La Jolla, CA, USA

[*]Correspondence should be addressed to tongwu.zhang@nih.gov, L2alexandrov@health.ucsd.edu and landim@nih.gov

ABSTRACT

The role of extrachromosomal DNA (ecDNA) in lung cancer, particularly in subjects who never

smoked (LCINS), remains unclear. Examination of 1,216 whole-genome-sequenced lung cancers

identified ecDNA in 18.9% of patients. Recurrent amplification of *MDM2* and other oncogenes

via ecDNA possibly drives a LCINS subset. Tumors harboring ecDNA showed worse overall

survival than tumors harboring other focal amplifications. A strong association with whole-

genome doubling suggests most ecDNA reflects genomic instability in treatment-naïve lung

cancer.

MAIN

Extrachromosomal DNA (ecDNA) is a potent mechanism for oncogene amplification in human

cancer[1], exhibiting unique characteristics such as a circular structure[2], non-Mendelian

inheritance[3,4], and an altered epigenetic and transcriptional landscape[5]. Prior studies have linked

ecDNA to worse overall survival in pan-cancer analyses, highlighting its clinical significance[6-8]

and potential as a therapeutic target[9]. Recent advances in computational techniques[10] have

enabled large-scale detection of ecDNA from whole-genome sequencing (WGS) data. However,

only three pan-cancer studies have examined ecDNA in primary lung cancer cohorts with over

100 samples, focusing predominantly on patients of European descent[6-8]. Two studies[6,7] analyzed

lung cancer samples from the United States, reporting ecDNA prevalence rates of about 21% in

36 lung adenocarcinomas (LUAD) and 27% in 47 lung squamous cell carcinomas (LUSC).

Additionally, a UK lung cancer study[8] analyzed 14,778 patients across 39 tumor types, including

718 LUAD tumors and 378 LUSC tumors, detecting ecDNA in 8.2% of LUAD and 22.4% of

LUSC cases. All three studies[6-8] reported lung cancer findings as part of broader pan-cancer

analyses and did not distinguish between lung cancers in subjects who have never-smoked

(LCINS) and subjects who have smoked (LCSS). Notably, based on the high prevalence of the

tobacco-associated[11] mutational signature SBS4 in these studies, more than 84% of the analyzed

lung cancer cases were from individuals with a history of smoking.

Despite these prior studies, the clinical relevance of ecDNA in lung cancer remains unclear,

including its prevalence across histologies, geographical areas, genetic ancestries, and biological

sexes. Its role in LCINS, LCSS, and subjects who were exposed to passive smoking also requires

further exploration. To address these questions, here, we analyzed WGS data from 1,216

treatment-naïve lung cancers from diverse patient populations (**Fig. 1a**; **Supplementary Fig. 1**).

This large international cohort enabled a comprehensive evaluation of ecDNA's association with

genomic alterations, as well as its prevalence across smoking status, biological sexes, ancestries,

and 18 areas defined primarily by country of residence. The dataset includes 871 LCINS,

sourced from the Sherlock-Lung[12] (*n*=817), the Environment And Genetics in Lung Cancer

Etiology (EAGLE[13], *n*=25), the Cancer Genome Atlas (TCGA[14], *n*=15), and other publicly

available cohorts (*n*=14)[15-19], representing patients from 17 geographical areas (**Supplementary**

Fig. 2a). Additionally, it comprises 345 LCSS, drawn from EAGLE[13] (*n*=221) and TCGA[14]

(*n*=83) and other publicly available cohorts (*n*=41)[15-19], spanning five geographical areas

(**Supplementary Fig. 2b**). The majority of tumors were LUAD (*n*=1,023; 84.1%) or LUSC

(*n*=67; 5.5%), with carcinoids (*n*=61; 5%, exclusively in LCINS) and other rarer histological

lung cancer subtypes (*n*=65; 5.3%) also represented. Amongst Sherlock-*Lung* LCINS, 250 were

exposed to secondhand tobacco smoke, while 208 had documented non-exposure (**Fig. 1a**).

To detect and reconstruct ecDNA across the cohort, a computational pipeline that accounts for

both tumor purity and ploidy was developed (**Methods**). A total of 231 samples harbored at least

one ecDNA (18.9%), with similar prevalence between LCINS (17%) and LCSS (23%; q-value:

0.21; **Fig. 1b**). No statistically significant differences were observed in the prevalence of ecDNA

across various histological subtypes, biological sexes, ancestries or age at diagnosis (**Fig. 1c&e**),

except for carcinoids, which showed a lower prevalence compared to LUAD in the overall

analysis (OR=0.07; q=0.04). ecDNA was present at similar rates across all geographic locations

(**Fig. 1d**; **Supplementary Fig. 2c-d**), as indicated by a one-versus-all multivariate logistic

regression for each country. There was also no difference in ecDNA prevalence between LCSS

and LCINS (**Fig.1c**). ecDNA was not associated with tumor stage (**Supplementary Fig. 3a&b**),

subsequent development of metastases (**Supplementary Fig. 3c&d**) or passive smoking status

(**Supplementary Fig. 3e**). These findings remained consistent when analyses were restricted to

only LUAD cases in either LCSS or LCINS (**Supplementary Fig. 4**).

To examine the association between ecDNA and genomic features previously linked to genome

instability[20-23], we evaluated the relationship of ecDNA with whole-genome doubling (WGD),

chromothripsis, tumor mutational burden, mitochondrial copy number, overall genome instability

index (wGII), and telomere length tumor/normal ratio. Amongst these, only WGD showed a

significant association with ecDNA in both LCSS and LCINS (**Fig. 2a**), as determined by

logistic regression adjusted for age, sex, ancestry, histology, and tumor purity. Specifically,

ecDNA was present in 7.1% of LCINS and 8.9% of LCSS without WGD, but its prevalence

increased markedly in WGD-positive tumors from LCINS (24.9%, OR=4.0; q=1.24 x 10^{-8}) and

LCSS (30.3%, OR=5.85; q=8.06 x 10^{-4}). The same association was observed when only LUAD

was examined (**Supplementary Fig. 5a**). Moreover, this association was further validated in a

pan-cancer dataset of 974 samples with high-confidence WGD annotation[14], using logistic

regression adjusted for age, purity, sex, and cancer type (OR=2.81; q=1.63 x 10^{-6};

Supplementary Fig. 5b).

In addition to WGD, chromothripsis was the only other feature associated with ecDNA, but this

was exclusively observed in LCINS (OR=1.76; q=0.048; **Fig. 2a**). Prior studies have indicated

that chromothripsis may contribute to the formation of certain ecDNA[23]. In this study, 115 out of

151 LCINS with ecDNA (76.2%) and 60 out of 80 LCSS with ecDNA (75%) exhibited

chromothripsis (**Supplementary Tables 1-2**). Nevertheless, in samples where both ecDNA and

chromothripsis were detected, overlap between ecDNA segments and chromothripsis regions

was rare, occurring more frequently in LCINS—observed in 31 of 115 LCINS cases compared to

4 of 60 LCSS (OR=5.01, p=0.02 comparing LCINS *vs.* LCSS; **Supplementary Fig. 5c**). This

suggests that there could be potentially different mechanisms of ecDNA formation between

LCINS and LCSS. However, while the majority of ecDNA emerges in genomically unstable

lung cancers that also exhibit chromothripsis, the actual genomic overlap between ecDNA and

chromothripsis regions is limited in both lung cancer types.

To evaluate the relationship between ecDNA presence and lung cancer driver genes, we analyzed

driver genes altered in at least 10 samples as well as driver hotspot mutations occurring at the

same genomic position in at least three samples. Using multivariate logistic regression, adjusted

for age, sex, tumor purity, ancestry, and histology, no significant associations were found

between ecDNA and individual driver gene alterations (**Fig. 2b**). However, hotspot analysis

showed an enrichment of *EGFR* L858R mutations (OR=1.78; q=0.05; **Fig. 2c**; **Supplementary**

Fig. 5d) in LCINS with ecDNA and a depletion of *KRAS* G12V in LCSS with ecDNA

(OR=0.18; q=0.03; **Fig. 2c**).

Mutational signature analysis provides insights into both the endogenous and exogenous

mutational processes that have been active throughout the lineage of a cancer genome[11]. We

investigated the association between ecDNA presence and mutational signatures. As expected[24],

multiple copy-number and structural variant signatures were associated with ecDNA presence

similarly in LCSS and LCINS (**Supplementary Fig. 6a-b**). Additionally, in LCINS, ecDNA-

positive samples were enriched for clock-like signatures (SBS1, SBS5, ID1, ID2) and, as

previously reported[19,25], APOBEC-associated signatures (SBS2, SBS13; **Supplementary Fig.**

6a). In LCSS, enrichment was observed only for SBS3 (**Supplementary Fig. 6b**), a signature

previously associated with homologous recombination deficiency[26].

Previous studies have shown that ecDNA can harbor oncogenes, regulatory elements, and

immunomodulatory genes in varying configurations[3,8,27,28]. In our cohort, 151 (17%) LCINS and

80 (23%) LCSS harbored ecDNA, with a total of 236 ecDNAs in LCINS and 123 in LCSS,

reflecting multiple ecDNAs in some tumors. Among these, ecDNA in LCINS were enriched with

oncogenes (p<0.001) and immunomodulatory genes (p=0.008) compared to LCSS (**Fig. 2d**;

Supplementary Fig. 7a-b). Further, ecDNA containing oncogenes exhibited elevated copy

numbers (CN) compared to other ecDNA without oncogenes in LCINS (p=1.5 x 10^{-3}) and LCSS

(5.8 x 10^{-3}), suggesting greater selective pressure for these ecDNAs (**Supplementary Fig. 7c**), as

previously reported in pan-cancer[8]. *MDM2* was the most frequently amplified oncogene on

ecDNA in both LCINS (*n*=36) and LCSS (*n*=5). Given its prominence, we examined the

relationship between ecDNA-driven *MDM2* amplification and *TP53* alterations, as MDM2

negatively regulates TP53 through ubiquitination[29]. We found that *MDM2* amplification on

ecDNA was mutually exclusive with *TP53* alterations (OR=7.51, p=0.01). In LCINS, the next

four most prevalent oncogenes that did not co-occur with *MDM2* were *TERT* (n=10), *CCND1*

(*n*=8), *MYC* (*n*=8), and *EGFR* (*n*=7; **Supplementary Table 3**). In LCSS, no oncogene besides

MDM2 was amplified on ecDNA in more than two samples (**Supplementary Table 4).**

Prior pan-cancer studies have shown that cancers harboring ecDNA amplifications exhibit worse

overall survival[6-8]. To evaluate this association in the lung cancer cohort, survival analysis was

conducted on 693 LCINS and 333 LCSS with available stage and covariate data. Lung cancers

were categorized as harboring ecDNA, other focal amplifications, or no focal amplifications.

Multivariate Cox proportional-hazards regression analysis revealed that ecDNA-positive lung

cancers had worse overall survival than those with only chromosomal amplifications; this

association was statistically significant in LCINS (hazard ratio=2.17; p=0.025; **Fig. 2***e*).

In summary, we examined the full spectrum of ecDNA in the largest dataset of whole-genome

sequenced lung cancers to date, from many geographical regions and ancestry, and with unique

information on tobacco smoking status. ecDNA was present in 17% of LCINS and 23% of

LCSS. ecDNA in LCINS were enriched with oncogenes and immunomodulatory genes in

comparison to LCSS. In LCINS, the *MDM2* locus was often amplified through ecDNA, while

the *EGFR* L858R mutation showed a weak association with ecDNA presence. No associations

were identified between presence of ecDNA and histological subtypes, biological sexes, smoking

status, or ancestries. However, tumors carrying ecDNA showed worse overall survival in

comparison to those with focal chromosomal amplification, particularly in LCINS. A strong

association was observed between ecDNA and WGD, and this observation was validated in an

independent pan-cancer cohort, suggesting that most ecDNA in treatment-naïve lung cancer are

likely to be a byproduct of genomic instability. Notably, ecDNA carrying oncogenes could

contribute to driving LCINS and affecting prognosis.

METHODS

Criteria for sample inclusion

A total of 1217 samples were available for analysis, but one sample with unknown smoking

status was excluded, resulting in a total of 1216 samples across the cohort.

Determination of sample purity and ploidy

The Battenberg algorithm (v.2.2.9)[30] was used to determine the sample purity and ploidy. Any

somatic copy number variations (CNV) profile determined to have low-quality after manual

inspection underwent a refitting process. This process required new tumor purity and ploidy

inputs, either estimated by Ccube (v.1.0)[31] or recalculated from local copy number status. The

Battenberg refitting procedures were iteratively executed until the final CNV profile was

established and met the criteria of manual validation check.

ecDNA detection and characterization

CNVKit v.0.9.6[32] was run in tumor-normal mode to call somatic CNVs against the matched

normal whole-genome sequenced samples for each patient, using the log ratio of tumor to normal

read depths (logR) and the circular binary segmentation (CBS) method. Note that this method, by

default, assumes a sample purity of 1 and a sample ploidy of 2 as it does not correct for purity

and ploidy. This limitation can lead to high copy number segments being missed due to low

purity[33]. Due to the wide range of tumor purities in our dataset, consistent with prior reports in

lung cancer[33], we rescaled the total copy number estimate using the purity and ploidy estimated

from Battenberg according to the following equation:

$$Equation\ 1.1.\ TCN = \frac{(\psi * (2^{logR})) - 2 * (1 - \rho)}{\rho}$$

where ψ is the tumor ploidy, ρ is the tumor purity, and logR represents the log ratio of the tumor

read depth to the normal read depth (**Supplementary Fig. 8a**). These copy number calls were

used to identify putative 'seed' regions. We then used AmpliconArchitect (v.1.3)[10] to reconstruct

the architecture of the amplicons and AmpliconClassifier (v.1.3.1) to classify amplicons

according to the most probable mechanism of formation (ecDNA, BFB, complex non-cyclic, or

linear). Visual inspection and verification were performed for all ecDNA amplicon regions. This

pipeline allowed for samples with low purities to be utilized as it completely removed the effect

of tumor purity in ecDNA detection (**Supplementary Fig. 8b**). As a result, more ecDNA$^+$

samples were detected (**Supplementary Fig. 8b**) and all 1216 samples could be used for ecDNA

analysis (**Supplementary Fig. 8c**). Of the additional ecDNA detected, eight of these were

EGFR-ecDNA (seven in LCINS and one in LCSS) and each of these ecDNA except for one also

harbored a driver mutation (**Supplementary Table 5**). Among the eight EGFR-ecDNA+

samples, five had RNA-seq data, which showed very high EGFR expression (**Supplementary**

Fig. 8d).

Determination of whole genome doubling status

As previously done[34], samples were considered to have a whole genome doubling event if the

major copy number was ≥3 for more than 50% of copy number segments.

Chromothripsis detection and characterization

To detect chromothripsis regions in genomes, we applied the computational algorithm

ShatterSeek[35], which employs a set of statistical criteria to identify chromothripsis given an input

of copy number and structural variants. We used the most stringent criteria corresponding to high

confidence chromothripsis calls. Specifically, these criteria were: *(i)* a cluster of structural

variants (>6 DUP/DEL/h2hINV/t2tINV), *(ii)* oscillating CNV between two states (>7 CNV

events), *(iii)* chromosomal enrichment and distribution of DNA breakpoints (p-value<0.05), *(iv)*

randomness of fragment joins (p-value>0.05) and/or ≥4 inter-chromosomal rearrangements

between multiple chromosomes. The input CNV calls were generated using CNVkit in tumor-

normal mode with default parameters, and the input structural variants calls were generated using

a union of Meerkat (v.0.189) and Manta (v.1.6.0) calls with the recommended filtering.

Association of ecDNA with demographic and clinical features

Demographic and clinical features where information was available for all samples (age, sex,

ancestry, and tumor histology and purity) served as input variables in a multivariate logistic

regression model with ecDNA presence or absence as the dependent outcome variable, and these

models were constructed separately for LCINS and LCSS. In addition, separate models for

passive smoking, tumor stage, and presence of metastases were run along with the rest of the

variables in both LCINS and LCSS.

Determination of telomere length tumor/normal lung tissue ratio

We estimated telomere length (TL) in kilobases using TelSeq (v.0.0.2)[36]. As was previously

done[37], we used seven as the threshold for the number of TTAGGG/CCCTAA repeats in a read

for the read to be considered telomeric. The TelSeq calculation was done individually for each

read group within a sample, and the total number of reads in each read group was used as weight

to calculate the average TL for each sample. The telomere length ratio ($\log_2$ scale) between

tumor and normal tissue was then calculated for each patient and served as the independent

variable in multivariable logistic regressions.

Association of mutational signatures and ecDNA status

Mutational signature attributions were determined as was previously done, based on the

methodology in Díaz-Gay et. al[37]. SigProfilerMatrixGenerator[38,39] was used to create the

mutational matrix for all types of somatic mutations. The deconvolution of mutational signatures

was performed by SigProfilerExtractor (v.1.1.3)[40] using the whole-genome sequencing setting

and the COSMIC mutational signatures (v3.2) as reference. Signature extractions were

performed separately for LCINS and LCSS. Signature attributions were input into a multivariate

logistic regression model as either present or absent if the signature was present in less than 50%

of the samples. If the signature was present in ≥50% samples (*e.g.*, SBS1 and SBS5), it was input

as either above median or below median. For SBS1 and SBS5, the signature attributions were

also input as continuous numeric variables to confirm the significant associations found with

inputting them as above median or below median.

Association of driver genes and ecDNA status

The IntOGen pipeline (v.2020.02.0123)[41], which combines seven state-of-the-art computational

methods, was employed with default parameters to detect signals of positive selection in the

mutational patterns of driver genes across the cohort. The status of each driver gene (altered or

unaltered) was then input into a logistic regression model with covariates age, sex, and tumor

purity, and ecDNA status was the output. These sets of genes were considered altered if there

was a focal deletion overlapping the gene, an inactivating (stop or missense) mutation, or an

activating gain of function mutation.

Association of driver mutations and ecDNA status

To identify driver mutations within the set of identified driver genes, we implemented a rigorous

and multifaceted strategy, considering multiple criteria: *(i)* the presence of truncating mutations

specifically in genes annotated as tumor suppressors, *(ii)* the recurrence of missense mutations in

a minimum of 3 samples, *(iii)* mutations designated as "Likely drivers" with a boostDM score[42]

exceeding 0.5, *(iv)* mutations categorized as either "Oncogenic" or "Likely Oncogenic" based on

the criteria established by OncoKB[43], an expert-guided precision oncology knowledge base, *(v)*

mutations previously recognized as drivers in the TCGA MC3 drivers paper[44], and *(vi)* missense

mutations characterized as "likely pathogenic" in genes that are annotated as tumor suppressors,

as described in Cheng et al.[45] Any mutation meeting one or more of these criteria was recognized

as a potential driver mutation.

Calculation of genome instability index score

To calculate the genome instability index score (wGII), the method devised in Bailey et al.[8] was

used. The overall ploidy was calculated as the length-weighted average of all copy number

segments. The proportion of segments that were aberrant (where total copy number was greater

than the ploidy or less than the ploidy) was computed for each chromosome, these proportions

were summed, and the total was divided by 22.

Identifying regulatory regions on ecDNA

A set of lung enhancer regions were downloaded from EnhancerAtlas[46], and each ecDNA region

was overlapped with these enhancer regions with the requirement that the entire enhancer had to

be fully contained (100% overlap) on the ecDNA. A set of lung cancer promoter regions was

extracted using a R script which downloaded regions 1000 base pairs upstream of transcription

start sites for all genes.

Defining a set of immunomodulatory genes

ecDNAs were considered immunomodulatory if a gene from the significant gene sets mapped to

an ecDNA that did not contain an oncogene, and that significant gene set had an

immunomodulatory function (GO terms: 0006968, 0002228, 0042267, 0001906, 0001909,

0002698, 0001910, 0031341, 0002367, 0002695, 0050866, 0051250, 0050777), following the

procedure in Bailey et. al[8]**.** This resulted in a set of 376 genes.

Mitochondrial copy number estimation

Mitochondrial copy number was estimated per sample using the ratio of mitochondrial read

depth to nuclear genome depth, adjusted for tumor ploidy and purity, following the procedure in

Reznik *et al.*[47]

Statistical analysis

All statistical analyses and graphic displays were performed using the R software v4.2.3 1011

(https://www.r-project.org/). Standard logistic regression with adjustment for covariates was

used for the enrichment analyses of categorical variables. We applied Firth's penalized logistic

regression using the logistf package in R to model the association between ecDNA status and

geographic regions, in order to reduce small-sample bias and address potential issues of data

separation. For the comparison of numerical variables across groups, we used non-parametric

Mann-Whitney (Wilcoxon rank sum) tests. P-values <0.05 were considered statistically

significant. Survival analyses were performed using Cox proportional-hazards regression

adjusted for age at diagnosis, sex, tumor stage, tumor histology, ancestry, and wGII. If multiple

hypothesis testing was required, we used a false-discovery rate correction based on the

Benjamini-Hochberg method and reported FDR. FDR <0.05 were considered statistically

significant.

 FIGURE LEGENDS

Figure 1: Landscape of extrachromosomal DNA in lung cancer across histologies, sex,

ancestry, countries, and smoking status. *a)* Epidemiological features of the LCINS (top) and

LCSS (bottom) cohorts. Total number of samples in each category is indicated. *b)* The number

and proportion of tumors harboring ecDNA among LCINS and LCSS, stratified by histology.

Samples with 1 or more ecDNA were classified as ecDNA^{+}, and samples with no ecDNA were

classified as ecDNA^{-}. *c)* Forest plot of the full cohort (LCINS and LCSS) from a multivariate

logistic regression analysis of key epidemiological and clinical features and the presence of

ecDNA in the tumors. *d)* Volcano plot showing $\log_2$ odds ratio (x-axis) and $\log_{10}$ FDR corrected

q-value (y-axis) for a one-vs-all statistical comparison of ecDNA prevalence by region. *e)* Forest

plots for multivariate logistic regression models testing key epidemiological and clinical features

with the presence of ecDNA in the tumors, stratified by smoking status. NA/EU: North America

or Europe region, AS: Asia region EAS: East Asian genetic ancestry super-sample, EUR:

European genetic ancestry super-sample, LUAD: lung adenocarcinoma, LUSC: lung squamous

cell carcinoma.

Figure 2. Genomic and clinical characteristics associated with ecDNA. *a)* Volcano plot of

logistic regression models of genomic features in association with ecDNA status for LCINS

(left) and LCSS (right). *b)* Volcano plot of a logistic regression model of driver gene alterations

in a sample in association with ecDNA status for LCINS (left) and LCSS (right). *c)* Volcano plot

of a logistic regression model of hotspot driver mutations in a sample in association with ecDNA

status for LCINS (left) and LCSS (right). In all volcano plots, the x-axes reflect the $\log_2$ odds

ratio, and the y-axes correspond to the $\log_{10}$ FDR q-value. An FDR q-value threshold of 0.05 is

indicated with the dashed red line, an FDR q-value threshold of 0.01 is indicated with the dashed

blue line. *d)* Bar plot indicating the proportion of ecDNA containing a particular genomic

element in LCINS vs LCSS. *e)* Kaplan–Meier survival curves for 5-year overall survival

stratified by the mode of amplification status for lung cancers in LCINS (left) and LCSS (right).

P-values for significance and HRs of the difference were calculated using two-sided Cox

proportional-hazards regression with adjustment for age, sex, tumor stage, ancestry and histology

and are indicated within each plot.

SUPPLEMENTARY FIGURE LEGENDS

Supplementary Fig 1. Additional cohort information. Sankey diagrams showing the numbers

of samples with clinical variables (stage, metastasis) as well as survival data available for *a)*

LCINS and *b)* LCSS

Supplementary Fig 2. Distribution of ecDNA harboring lung cancers in LCINS and LCSS

by country. Maps showing the worldwide prevalence of lung cancers harboring ecDNA

(ecDNA$^+$) by country for the *a)* LCINS cohort and *b)* LCSS cohort. Volcano plot of a

multivariate one-vs-all logistic regression model for each country, with ecDNA status as the

outcome, for *c)* LCINS and *d)* LCSS.

Supplementary Fig 3. Analysis of the association between stage, metastasis, and passive

smoking with ecDNA status in all LCINS and LCSS. Forest plots of logistic regression model

with ecDNA status as the outcome and the variables for stage in **a)** LCINS and **b)** LCSS;

metastasis in **c)** LCINS and **d)** LCSS; and passive smoking in **e)** LCINS. The total number of

samples with information for all variables is indicated above each plot.

Supplementary Fig 4. Analysis of the association between stage, metastasis, and passive

smoking with ecDNA status in lung adenocarcinomas. Forest plots of logistic regression

model with ecDNA status as the outcome and the variables for stage in **a)** LCINS and **b)** LCSS;

metastasis in **c)** LCINS and **d)** LCSS; and passive smoking in **e)** LCINS. The total number of

samples with information for all variables is indicated above each plot.

Supplementary Fig 5. Genomic features associated with the presence of ecDNA. *a)* Volcano

plot of logistic regression models of genomic features in association with ecDNA status for lung

adenocarcinoma in LCINS (left) and lung adenocarcinoma in LCSS (right). In all volcano plots,

the x-axes reflect the $\log_2$ odds ratio, and the y-axes correspond to the $\log_{10}$ FDR q-value. An

FDR q-value threshold of 0.05 is indicated with the dashed red line, an FDR q-value threshold of

0.01 is indicated with the dashed blue line. *b)* Forest plot of multivariate logistic regression

modeling the presence or absence of ecDNA in relation to whole-genome doubling in a pan-

cancer dataset comprising four different tumor types from the PCAWG project, adjusted for the

covariates of age, purity, sex, and tissue type. All significant associations are indicated in red. *c)*

Bar plot indicating the proportion of samples that contained overlapping chromothripsis and

ecDNA regions (red) and the proportion of all samples that did not contain any overlap (gray) for

all samples that had both an ecDNA and chromothripsis event detected. *d)* Bar plot showing the

proportion of samples with an *EGFR* L858R mutation, stratified by ecDNA status.

Supplementary Fig 6. Association of mutational signatures with ecDNA status. Volcano

plots of logistic regression models of the activities of Single base substitution (SBS), Indel (ID),

copy number (CN) and structural variant (SV) signatures in association with ecDNA in lung

cancers of *a)* LCINS and *b)* LCSS. The logistic regression models were adjusted for age, sex,

ancestry, histology, and purity. In all volcano plots, the x-axes reflect the $\log_2$ odds ratio, and the

y-axes correspond to the $\log_{10}$ FDR q-value. An FDR q-value threshold of 0.05 is indicated with

the dashed red line, an FDR q-value threshold of 0.01 is indicated with the dashed blue line.

Supplementary Fig 7. Genomic content of ecDNA

a) Upset plot indicating the number of ecDNA with oncogenes, immunomodulatory genes,

promoter or enhancer regions, other genes that are not oncogenes, or unidentified genomic

elements (Unknown) or any combination thereof for LCINS and LCSS. **b)** Bar plot indicating

the proportion of ecDNA containing a particular genomic element in LCINS vs LCSS. **c)** $\log_2$

median copy number (y-axis) for each class of ecDNA (x-axis) in LCINS and LCSS.

Supplementary Fig 8. Highly sensitive pipeline for ecDNA detection that accounts for

tumor purity and ploidy.

a) Schematic depicting the methodology for the rescaled pipeline that considers tumor purity and

ploidy. The logR track for the region containing the *EGFR* gene on chr7 and the purity and

ploidy computed from Battenberg is indicated in one tumor as example (left). The exact location

of the *EGFR* gene is shown by the dashed red lines. The equation that simultaneously accounts

for sample purity and ploidy when computing the total copy number is shown, as well as the total

copy number estimate when assuming a sample purity of 1.0 and ploidy of 2.0 (Uncorrected CN)

vs. considering the purity and ploidy (Corrected CN). The Sashimi plot indicates the structural

variants (colored arcs) and smoothed coverage around the *EGFR* region, which was annotated as

ecDNA by AmpliconClassifier. **b)** Distribution of tumor purities for samples with an ecDNA

annotation for both the default (left) and rescaled (right) pipelines. P-value for a Mann-Whitney

test between ecDNA$^+$ and ecDNA$^-$ samples is indicated. The total number of samples in each

group is also indicated. **c)** The ecDNA prevalence (y-axis), which is the proportion of ecDNA$^+$

samples over the total number of samples, for all possible tumor purity cutoffs applied to

samples (x-axis). The total number of samples available for analysis after each tumor purity

cutoff is applied is also indicated. **d)** Gene expression ($\log_2$ CPM) of the EGFR gene for all

samples with RNA-seq data available. LCINS samples with EGFR-ecDNA are indicated in red.

SUPPLEMENTARY TABLE LEGENDS

Supplementary Table 1. Co-occurrence of ecDNA genetic cargo in LCINS

Supplementary Table 2. Co-occurrence of ecDNA genetic cargo in LCSS

Supplementary Table 3. Recurrently amplified oncogenes on ecDNA in LCINS across

unique genomic loci

Supplementary Table 4. Recurrently amplified oncogenes on ecDNA in LCSS across

unique genomic loci

Supplementary Table 5. Additional *EGFR*-ecDNA detected by modified pipeline

Ethics declarations

Since the National Cancer Institute only received de-identified samples and data from collaborating centers, had no direct contact or interaction with the study participants, and did not use or generate identifiable private information, Sherlock-*Lung* has been determined to constitute "Not Human Subject Research (NHSR)" based on the federal Common Rule (45 CFR 46; https://www.ecfr.gov/cgi-bin/ECFR?page=browse).

DATA AVAILABILITY

Normal and tumor-paired CRAM files for the WGS subjects of the Sherlock-Lung study and the EAGLE study have been deposited in dbGaP under the accession numbers phs001697.v2.p1 and phs002992.v1.p1, respectively.

Author Contributions

Conceptualization, MTL, TZ, LBA; Methodology, AK, TZ; Formal Analysis, AK, TZ, MDG, JL, WZ; Data Curation, TZ, PH, JM, CH, MM, OWL, MAN, NR, RH, SY, QL; Writing – Original Draft, AK; Writing – Review & Editing, LBA, TZ, PHH, MTL; Visualization: AK; Supervision, LBA, TZ, MTL.

ACKNOWLEDGEMENTS

This work was supported by the Intramural Research Program of the National Cancer Institute, US National Institute of Health (NIH) (project ZIACP101231 to MTL), and by the Anne Wojcicki Foundation (Grant Number: LC009). Part of this work was supported by the US National Institute of Health grants R01ES032547, R01CA269919, P01CA281819, and

U01CA290479 to LBA as well as by LBA's Packard Fellowship for Science and Engineering. MD-G was awarded with a fellowship within the "Generación D" initiative, Red.es, Ministerio para la Transformación Digital y de la Función Pública, for talent attraction (C005/24-ED CV1), funded by the European Union NextGenerationEU funds, through PRTR. The research in this study was also supported by UC San Diego Sanford Stem Cell Institute. This work utilized the computational resources of the NIH HPC Biowulf cluster (https://hpc.nih.gov). The funders had no roles in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

COMPETING INTERESTS

LBA is a co-founder, CSO, scientific advisory member, and consultant for io9, has equity and receives income. The terms of this arrangement have been reviewed and approved by the University of California, San Diego in accordance with its conflict of interest policies. LBA is a compensated member of the scientific advisory board of Inocras. LBA's spouse is an employee of Hologic, Inc. LBA declares U.S. provisional applications with serial numbers: 63/289,601; 63/269,033; 63/366,392; 63/412,835 as well as international patent application PCT/US2023/010679. LBA is also an inventor of a US Patent 10,776,718 for source identification by non-negative matrix factorization. LBA and MD-G further declare a European patent application with application number EP25305077.7. Soo-Ryum Yang has received speaking fees from Medscape, OncLive, Cure Today, Medical Learning Institute, PRIME Education, AstraZeneca, Roche; and consulting fees from AstraZeneca, AbbVie, Merus, Eli Lilly, Boehringer Ingelheim, Roche, Amgen, Sanofi. All other authors declare no conflicts of interest.

REFERENCES

Benner, S. E., Wahl, G. M. & Von Hoff, D. D. Double minute chromosomes and homogeneously staining regions in tumors taken directly from patients versus in human tumor cell lines. *Anticancer Drugs* **2**, 11-25 (1991). https://doi.org/10.1097/00001813-199102000-00002

Nielsen, J., Walsh, J. T., Degen, D. R., Drabek, S. M. & McGill, J. R. Evidence of gene amplification in the form of double minute chromosomes is frequently observed in lung cancer. *Cancer Genetics* **65**, 120-124 (1993). https://doi.org/https://doi.org/10.1016/0165-4608(93)90219-C

Hung, K. L. *et al.* Coordinated inheritance of extrachromosomal DNAs in cancer cells. *Nature* **635**, 201-209 (2024). https://doi.org/10.1038/s41586-024-07861-8

Lange, J. T. *et al.* The evolutionary dynamics of extrachromosomal DNA in human cancers. *Nat Genet* **54**, 1527-1533 (2022). https://doi.org/10.1038/s41588-022-01177-x

Wu, S. *et al.* Circular ecDNA promotes accessible chromatin and high oncogene expression. *Nature* **575**, 699-703 (2019). https://doi.org/10.1038/s41586-019-1763-5

Kim, H. *et al.* Mapping extrachromosomal DNA amplifications during cancer progression. *Nat Genet* **56**, 2447-2454 (2024). https://doi.org/10.1038/s41588-024-01949-7

Kim, H. *et al.* Extrachromosomal DNA is associated with oncogene amplification and poor outcome across multiple cancers. *Nat Genet* **52**, 891-897 (2020). https://doi.org/10.1038/s41588-020-0678-2

Bailey, C. *et al.* Origins and impact of extrachromosomal DNA. *Nature* **635**, 193-200 (2024). https://doi.org/10.1038/s41586-024-08107-3

Tang, J. *et al.* Enhancing transcription-replication conflict targets ecDNA-positive cancers. *Nature* **635**, 210-218 (2024). https://doi.org/10.1038/s41586-024-07802-5

Deshpande, V. *et al.* Exploring the landscape of focal amplifications in cancer using AmpliconArchitect. *Nat Commun* **10**, 392 (2019). https://doi.org/10.1038/s41467-018-08200-y

Alexandrov, L. B. *et al.* The repertoire of mutational signatures in human cancer. *Nature* **578**, 94-101 (2020). https://doi.org/10.1038/s41586-020-1943-3

Landi, M. T. *et al.* Tracing Lung Cancer Risk Factors Through Mutational Signatures in Never-Smokers. *Am J Epidemiol* **190**, 962-976 (2021). https://doi.org/10.1093/aje/kwaa234

Consonni, D. *et al.* Outdoor particulate matter (PM10) exposure and lung cancer risk in the EAGLE study. *PLoS One* **13**, e0203539 (2018). https://doi.org/10.1371/journal.pone.0203539

Consortium, I. T. P.-C. A. o. W. G. Pan-cancer analysis of whole genomes. *Nature* **578**, 82-93 (2020). https://doi.org/10.1038/s41586-020-1969-6

Lee, J. J. *et al.* Tracing Oncogene Rearrangements in the Mutational History of Lung Adenocarcinoma. *Cell* **177**, 1842-1857 e1821 (2019). https://doi.org/10.1016/j.cell.2019.05.013

Imielinski, M. *et al.* Mapping the hallmarks of lung adenocarcinoma with massively parallel sequencing. *Cell* **150**, 1107-1120 (2012). https://doi.org/10.1016/j.cell.2012.08.029

17 Lee, J. K. *et al.* Clonal History and Genetic Predictors of Transformation Into Small-Cell
Carcinomas From Lung Adenocarcinomas. *J Clin Oncol* **35**, 3065-3074 (2017).
https://doi.org/10.1200/JCO.2016.71.9096
18 Leong, T. L. *et al.* Deep multi-region whole-genome sequencing reveals heterogeneity
and gene-by-environment interactions in treatment-naive, metastatic lung cancer.
Oncogene **38**, 1661-1675 (2019). https://doi.org/10.1038/s41388-018-0536-1
19 Bergstrom, E. N. *et al.* Mapping clustered mutations in cancer reveals APOBEC3
mutagenesis of ecDNA. *Nature* **602**, 510-517 (2022). https://doi.org/10.1038/s41586-
022-04398-6
20 Chen, J. *et al.* MYC-driven increases in mitochondrial DNA copy number occur early
and persist throughout prostatic cancer progression. *JCI Insight* **8** (2023).
https://doi.org/10.1172/jci.insight.169868
21 Bielski, C. M. *et al.* Genome doubling shapes the evolution and prognosis of advanced
cancers. *Nat Genet* **50**, 1189-1195 (2018). https://doi.org/10.1038/s41588-018-0165-1
22 O'Sullivan, R. J. & Karlseder, J. Telomeres: protecting chromosomes against genome
instability. *Nat Rev Mol Cell Biol* **11**, 171-181 (2010). https://doi.org/10.1038/nrm2848
23 Shoshani, O. *et al.* Chromothripsis drives the evolution of gene amplification in cancer.
Nature **591**, 137-141 (2021). https://doi.org/10.1038/s41586-020-03064-z
24 Steele, C. D. *et al.* Signatures of copy number alterations in human cancer. *Nature* **606**,
984-991 (2022). https://doi.org/10.1038/s41586-022-04738-6
25 Hadi, K. *et al.* Distinct Classes of Complex Structural Variation Uncovered across
Thousands of Cancer Genome Graphs. *Cell* **183**, 197-210 e132 (2020).
https://doi.org/10.1016/j.cell.2020.08.006
26 Davies, H. *et al.* HRDetect is a predictor of BRCA1 and BRCA2 deficiency based on
mutational signatures. *Nat Med* **23**, 517-525 (2017). https://doi.org/10.1038/nm.4292
27 Hung, K. L. *et al.* ecDNA hubs drive cooperative intermolecular oncogene expression.
Nature **600**, 731-736 (2021). https://doi.org/10.1038/s41586-021-04116-8
28 Zhu, Y. *et al.* Oncogenic extrachromosomal DNA functions as mobile enhancers to
globally amplify chromosomal transcription. *Cancer Cell* **39**, 694-707.e697 (2021).
https://doi.org/10.1016/j.ccell.2021.03.006
29 Chene, P. Inhibiting the p53-MDM2 interaction: an important target for cancer therapy.
Nat Rev Cancer **3**, 102-109 (2003). https://doi.org/10.1038/nrc991
30 Nik-Zainal, S. *et al.* The life history of 21 breast cancers. *Cell* **149**, 994-1007 (2012).
https://doi.org/10.1016/j.cell.2012.04.023
31 Dharanipragada, P. *et al.* Blocking Genomic Instability Prevents Acquired Resistance to
MAPK Inhibitor Therapy in Melanoma. *Cancer Discov* **13**, 880-909 (2023).
https://doi.org/10.1158/2159-8290.CD-22-0787
32 Talevich, E., Shain, A. H., Botton, T. & Bastian, B. C. CNVkit: Genome-Wide Copy
Number Detection and Visualization from Targeted DNA Sequencing. *PLoS Comput*
Biol **12**, e1004873 (2016). https://doi.org/10.1371/journal.pcbi.1004873
33 Aran, D., Sirota, M. & Butte, A. J. Systematic pan-cancer analysis of tumour purity. *Nat*
Commun **6**, 8971 (2015). https://doi.org/10.1038/ncomms9971
34 Zhang, T. *et al.* Deciphering lung adenocarcinoma evolution and the role of LINE-1
retrotransposition. *bioRxiv* (2025). https://doi.org/10.1101/2025.03.14.643063

35   Cortes-Ciriano, I. *et al.* Comprehensive analysis of chromothripsis in 2,658 human
cancers using whole-genome sequencing. *Nat Genet* **52**, 331-341 (2020).
https://doi.org/10.1038/s41588-019-0576-7
36   Ding, Z. *et al.* Estimating telomere length from whole genome sequence data. *Nucleic
Acids Res* **42**, e75 (2014). https://doi.org/10.1093/nar/gku181
37   Diaz-Gay, M. *et al.* The mutagenic forces shaping the genomic landscape of lung cancer
in never smokers. *medRxiv* (2024). https://doi.org/10.1101/2024.05.15.24307318
38   Khandekar, A. *et al.* Visualizing and exploring patterns of large mutational events with
SigProfilerMatrixGenerator. *BMC Genomics* **24**, 469 (2023).
https://doi.org/10.1186/s12864-023-09584-y
39   Bergstrom, E. N. *et al.* SigProfilerMatrixGenerator: a tool for visualizing and exploring
patterns of small mutational events. *BMC Genomics* **20**, 685 (2019).
https://doi.org/10.1186/s12864-019-6041-2
40   Islam, S. M. A. *et al.* Uncovering novel mutational signatures by de novo extraction with
SigProfilerExtractor. *Cell Genom* **2**, 100179 (2022).
https://doi.org/10.1016/j.xgen.2022.100179
41   Gonzalez-Perez, A. *et al.* IntOGen-mutations identifies cancer drivers across tumor types.
Nat Methods **10**, 1081-1082 (2013). https://doi.org/10.1038/nmeth.2642
42   Muinos, F., Martinez-Jimenez, F., Pich, O., Gonzalez-Perez, A. & Lopez-Bigas, N. In
silico saturation mutagenesis of cancer genes. *Nature* **596**, 428-432 (2021).
https://doi.org/10.1038/s41586-021-03771-1
43   Chakravarty, D. *et al.* OncoKB: A Precision Oncology Knowledge Base. *JCO Precis
Oncol* **2017** (2017). https://doi.org/10.1200/PO.17.00011
44   Bailey, M. H. *et al.* Comprehensive Characterization of Cancer Driver Genes and
Mutations. *Cell* **174**, 1034-1035 (2018). https://doi.org/10.1016/j.cell.2018.07.034
45   Cheng, F. *et al.* Comprehensive characterization of protein-protein interactions perturbed
by disease mutations. *Nat Genet* **53**, 342-353 (2021). https://doi.org/10.1038/s41588-020-
00774-y
46   Gao, T. & Qian, J. EnhancerAtlas 2.0: an updated resource with enhancer annotation in
586 tissue/cell types across nine species. *Nucleic Acids Res* **48**, D58-D64 (2020).
https://doi.org/10.1093/nar/gkz980
47   Reznik, E. *et al.* Mitochondrial DNA copy number variation across human cancers. *Elife*
5 (2016). https://doi.org/10.7554/eLife.10769
610

Fig 1.

Fig 2.