Pancreatic cancer cachexia is mediated by PTHrP-driven disruption of adipose *de novo* lipogenesis

Nikita Bhalerao[1,2], Yamini Ogoti[1,2], Jessica Peura[1,2], Calvin Johnson[1,2], Qingbo Chen[3], Ekaterina D. Korobkina[3], Faith N. Keller[4], Maximilian Wengyn[5], Robert J. Norgard[5], Claire Shamber[1,2], Kelsey Klute[6], Dominick DiMaio[7], Karine Sellin[8], Paul M. Grandgenett[9], Rui Li[10], Michael A. Hollingsworth[9], Adilson Guilherme[11], Michael P. Czech[11], Richard Kremer[8], Lihua Julie Zhu[2,3,10], Emma V. Watson[4], Marcus Ruscetti[2], David A. Guertin[3], Jason R. Pitarresi[1,2]

[1] Division of Hematology-Oncology, Department of Medicine, University of Massachusetts Chan Medical School, Worcester, MA, USA

[2] Department of Molecular, Cell and Cancer Biology, University of Massachusetts Chan Medical School, Worcester, MA, USA

[3] Program in Molecular Medicine, University of Massachusetts Chan Medical School, Worcester, MA, USA

[4] Department of Systems Biology, University of Massachusetts Chan Medical School, Worcester, MA, USA

[5] Abramson Family Cancer Research Institute and Department of Medicine, University of Pennsylvania, Philadelphia, PA, USA

[6] Department of Internal Medicine, University of Nebraska Medical Center, Omaha, NE, USA

[7] Department of Pathology and Microbiology, College of Medicine, University of Nebraska Medical Center, Omaha, NE, USA

8 Division of Endocrinology and Metabolism, Department of Medicine, McGill University and McGill University Health Centre, Montréal, Quebec, Canada.

9 Eppley Institute for Research in Cancer and Allied Diseases, Fred and Pamela Bufferr Cancer Center, University of Nebraska Medical Center, Omaha, NE, USA

10 Department of Genomics and Computational Biology, University of Massachusetts Chan Medical School, Worcester, MA, USA

11 Program in Molecular Medicine, Diabetes Center of Excellence, University of Massachusetts Chan Medical School, Worcester, MA, USA

Corresponding author: J.R. Pitarresi; Jason.Pitarresi@umassmed.edu

Conflict of interest statement: The authors declare no potential conflicts of interest. RK is the inventor of the therapeutic monoclonal anti-PTHrP antibodies used in this study and authorized for use by Miramab Bioscience. RK holds an equity stake in Miramab Bioscience.

Running Title: PTHrP drives pancreatic cancer cachexia

Keywords: Pancreatic cancer, PTHrP, PTHLH, cancer cachexia, adipose wasting, *de novo* lipogenesis, DNL

ABSTRACT

Pancreatic cancer patients have the highest rates and most severe forms of cancer cachexia, yet cachexia etiologies remain largely elusive, leading to a lack of effective intervening therapies. PTHrP has been clinically implicated as a putative regulator of cachexia, with serum PTHrP levels correlating with increased weight loss in PDAC patients. Here we show that cachectic PDAC patients have high expression of tumor PTHrP and use a genetically engineered mouse model to functionally demonstrate that deletion of *Pthlh* (encoding the PTHrP protein) blocks cachectic wasting, dramatically extending overall survival. The re-expression of PTHrP in lowly cachectic models is sufficient to induce wasting and reduce survival in mice, which is reversed by the conditional deletion of the PTHrP receptor, *Pth1r*, in adipocytes. Mechanistically, tumor-derived PTHrP suppresses *de novo* lipogenesis in adipocytes, leading to a molecular rewiring of adipose depots to promote wasting in the cachectic state. Finally, the pharmacological disruption of the PTHrP-PTH1R signaling axis abrogates wasting, highlighting that a targeted disruption of tumor-adipose crosstalk is an effective means to limit cachexia.

STATEMENT OF SIGNIFICANCE

Pancreatic ductal adenocarcinoma (PDAC) is the prototypical cancer type associated with cancer cachexia, a debilitating wasting syndrome marked by adipose tissue loss and muscle atrophy. Herein, we establish that PTHrP is a tumor-derived factor that facilitates cachexia by downregulating *de novo* lipogenesis in adipocytes and that blocking PTHrP is an effective means to limit wasting in preclinical mouse models.

INTRODUCTION

Pancreatic cancer ranks as the third most fatal cancer in the United States, surpassed only by lung cancer and colorectal cancer [1]. The lack of robust treatment options and early detection opportunities for Pancreatic Ductal Adenocarcinoma (PDAC) patients contribute to a dismal 5-year survival rate of ~13% [2]. In addition, PDAC patients often present with cancer-associated cachexia, a systemic metabolic syndrome marked by profound adipose tissue and muscle wasting, which impacts body condition and tolerance to available treatment options, thus contributing to reduced survival rates. A striking >75% of PDAC patients suffer from cachexia [3], with nearly one-third of PDAC-associated deaths being attributable to cachexia rather than tumor burden [4]. The perturbed metabolic state in cachectic patients across cancer types has been proposed to be mediated by various tumor-derived factors, such as IL-6 [5], Parathyroid Hormone-Related Protein (PTHrP) [6], TNFα [7], and GDF15 [8, 9], that likely act in concert to alter tumor-host interactions and facilitate wasting.

Clinically, cachexia presents as a multifactorial metabolic syndrome characterized by loss of body weight due to wasting of adipose tissue and muscle mass [10]. Despite patients often presenting with anorexia and anemia, nutritional supplementation does little to curtail weight loss and extend survival [11]. Thus, it is clear from clinical evidence and preclinical models that cachexia is a comorbidity in cancer and that a deeper understanding of the molecular basis of cancer-associated cachexia is required to find new ways to treat cachectic patients and, ultimately, extend survival. Additionally, how tumors communicate with peripheral tissues such as adipose and muscle is largely

unexplored and represents an untapped area to apply targeted therapy to block cachectic

48   wasting.

50   Adipose tissue loss has been shown to precede muscle loss in cachectic PDAC patients

[12], nominating tumor-adipose crosstalk as a potential point of therapeutic intervention.

52   Under normal physiological conditions, white adipocytes have unilocular lipid droplets and

are involved in energy storage and mobilization. During cachexia, some white adipocytes

54   shift their metabolism towards an energy-expending state that resembles thermogenesis

[12]. This shift from a primarily energy-storing organ to an energy-consuming organ is a

56   hallmark of adipose tissue wasting in cancer cachexia, yet few tumor-derived factors have

been nominated as putative regulators of this crosstalk. Canonically, this metabolic shift

58   has been attributed to an upregulation of adipocyte lipolysis, or the breaking down of fatty

acids [13-15]. However, emerging work suggests that decreased *de novo* lipogenesis

60   (DNL) may also contribute to adipose tissue loss in this context [16].

62   To better understand the molecular etiology of cancer cachexia and associated adipose

tissue wasting, we employ a murine model of pancreatic cancer that is accompanied by

64   severe cachexia. We identify PTHrP as a PDAC tumor-derived factor that promotes

cancer-associated adipose wasting marked by profound downregulation of *de novo*

66   lipogenesis, and provide pre-clinical evidence to support anti-PTHrP therapy as a

cachexia blocking strategy.

RESULTS

Tumor cell-specific Pthlh deletion extends overall survival and reduces PDAC-associated cachexia

The conditional expression of $Kras^{LSL\text{-}G12D}$ and $Trp53^{LSL\text{-}R172H}$ were targeted to the murine pancreas using the *Pdx1-Cre* allele, as previously described [17]. The addition of a

Rosa26$^{LSL\text{-}YFP}$ lineage label allows for the tracking of tumor cells derived from the Pdx1 lineage [18]. The resulting **K**$ras^{LSL\text{-}G12D}$; *Tr**p**53*$^{LSL\text{-}R172H}$; *Pdx1-**C**re*; Rosa26$^{LSL\text{-}\underline{Y}FP}$ (KPCY)

mice were combined with the *Pthlh*^{LoxP} allele (*Pthlh* gene encodes the PTHrP protein) to delete *Pthlh* specifically in tumor cells (Supplemental Figure 1A). Our prior work revealed

a genetic linkage between *Kras* and *Pthlh* (these genes are directly adjacent in the mouse and human genome), which precludes the inheritance of homozygous *Pthlh*$^{LoxP/LoxP}$

alleles in combination with $Kras^{LSL\text{-}G12D}$ [19]. Nonetheless, the heterozygous deletion of *Pthlh* in KPCY mice (KPCY-Pthlh^{HET}) dramatically extended survival compared to KPCY

controls, with KPCY-Pthlh^{HET} mice living ~2.5 months longer (Figure 1A). KPCY-Pthlh^{HET} mice survived a median of 187 days, a ~68% extension of survival relative to the KPCY

median survival of 111 days, which we had previously attributed to reduced metastatic ability [19]. Upon closer examination of the overall body condition of KPCY-Pthlh^{HET}

animals, we also observed that they were less cachectic than their KPCY counterparts. To quantify cachexia, body weights were tracked in an age-matched cohort of KPCY and

KPCY-Pthlh^{HET} mice in the 15 days prior to sacrifice. KPCY mice underwent significant cachectic wasting over time, while KPCY-Pthlh^{HET} mice, on average, continued to gain or

maintain their body mass (Figure 1B). At endpoint, the mouse carcass weights were increased in KPCY-Pthlh^{HET} mice relative to KPCY controls (Figure 1C).

To profile more deeply the degree of cachectic wasting in these animals, inguinal white adipose tissue (iWAT), epididymal white adipose tissue (eWAT), and quadricep (quad)

muscle were collected across two cohorts; the first set of tissue was collected when mice reached endpoint (Figure 1D-F), and the second cohort at a set endpoint of 5 months

(Figure 1G-I). At endpoint, KPCY-Pthlh[HET] mice showed reduced iWAT, eWAT, and quad wasting (Figure 1D-F). Similarly, in the age-matched 5-month-old cohort, KPCY-Pthlh[HET]

mice showed reduced iWAT, eWAT, and quad wasting (Figure 1G-I). Strikingly, KPCY-Pthlh[HET] adipose and muscle tissue masses were more similar to those of non-tumor

bearing (NTB) animals, suggesting that the loss of tumor cell-derived PTHrP blocked cachectic wasting. Histopathologically, the adipose tissue of KPCY mice was highly

cachectic, with marked remodeling of the adipose architecture and a reduction in adipocyte lipid droplet size (Figure 1D-E and 1G-H). In contrast, iWAT and eWAT from

KPCY-Pthlh[HET] mice more closely resembled that of NTB hosts and were primarily composed of large unilocular adipocytes (Figure 1D-E and 1G-H). Quad muscle

morphology was also perturbed in KPCY mice, while muscle from KPCY-Pthlh[HET] mice was more similar to NTB (Figure 1F and 1I). Cachectic muscle from KPCY mice had

increased expression of the E3 ubiquitin ligases *Trim63* (MURF1) and *Fbxo32* (ATROGIN1) that mediate muscle wasting [20], which were lost in KPCY-Pthlh[HET] mice

(Supplemental Figure 1B). To precisely quantify the cachectic burden in adipose tissue, we measured adipocyte size, a proxy for the lipid storing capacity of adipocytes.

Adipocyte size was reduced in the iWAT from *KPCY* mice (indicative of adipose wasting), while KPCY-Pthlh[HET] adipocytes were larger and more similar in size to NTB animals

(Figure 1J; Supplemental Figure 1C). Fluorescent staining of iWAT with the adipocyte lipid

116   droplet marker PLIN1 indicated that overall lipid size was reduced in cachectic KPCY

mice, which returned to normal in non-cachectic KPCY-Pthlh[HET] mice (Figure 1K).

Importantly, overall tumor burden did not correlate with the degree of adipose

120   (Supplemental Figure 1D; Pearson R^2=0.002) or muscle wasting (Supplemental Figure

1E; Pearson R^2=0.015) across our cohort of KPCY orthotopic and GEMM animals,

122   consistent with prior reports demonstrating that tumor burden does not dictate the degree

of cancer-associated cachexia in other models [21, 22]. In fact, we often observed

124   relatively small tumors (~500-750mg) that were highly cachectic and much larger tumors

(>2 grams) that showed no evidence of cachectic wasting (Supplemental Figure 1D-E).

126   Taken together, the varying degree of PDAC-associated cachexia in genetically identical

KPCY tumor bearing animals indicates that the ability to induce cachexia is a tumor cell-

128   intrinsic trait, and we have identified highly cachectic and lowly cachectic KPCY tumor

cell clones to test putative cachexia-inducing tumor-derived factors such as PTHrP.

Prior work has established that cachectic PDAC patients have higher serum PTHrP

132   relative to non-cachectic patients [23]. We hypothesized, based on our prior work, that

PTHrP is coming from PDAC tumor cells [19], and set out to determine if high PTHrP

134   tumor expression corresponds with cachectic burden in a cohort of cachectic and non-

cachectic PDAC patients. Indeed, levels of PTHrP in primary tumors from cachectic

136   PDAC patients were increased relative to non-cachectic PDAC patients (Figure 1L), and

we have previously demonstrated that increased PTHrP staining in primary tumors

138   correlates with decreased survival of PDAC patients [19]. Collectively, these data show

that PTHrP mediates adipose and muscle wasting cachectic phenotypes in genetically

140   engineered mouse models of pancreatic cancer and that PTHrP is increased in cachectic

PDAC patients, setting the stage for mechanistic studies to determine the role of tumor-

142   derived PTHrP in the induction of pancreatic cancer cachexia.

Pthlh overexpression is sufficient to induce cachexia

146   The genetic deletion of PTHrP reduced cachexia phenotypes, demonstrating that it is

necessary for PDAC-associated cachectic wasting. To determine if PTHrP is sufficient to

148   induce wasting, we took a lowly cachectic KPCY cell line with low basal expression of

Pthlh (Supplemental Figure 2A) and stably re-expressed *Pthlh* (PTHrP^{OE}) via lentiviral-

150   mediated transduction (Supplemental Figure 2B). The resulting KPCY-PTHrP^{OE} tumor

cells had a modest, but statistically significant increase in tumor cell proliferation *in vitro*

152   relative to empty vector expressing cells (KPCY-EV; Supplemental Figure 2C), consistent

with our prior reports [19]. Upon orthotopic implantation, the re-expression of *Pthlh*

154   significantly reduced overall survival (Figure 2A) and was sufficient to induce a dramatic

cachectic phenotype, effectively converting a non-cachectic PDAC cell line into a

156   cachectic one (Figure 2B). Consistent with the induction of cachexia, iWAT architecture

was altered upon *Pthlh* overexpression (Figure 2C), and iWAT mass was greatly reduced

158   in KPCY-PTHrP^{OE} tumor bearing animals relative to isogenic KPCY-EV controls (Figure

2D). Adipocyte size was also reduced (Figure 2E; Supplemental Figure 2D) with a

160   concomitant reduction in PLIN1-positive lipid droplet size in iWAT from KPCY-PTHrP^{OE}

mice (Figure 2F). Similarly, muscle architecture was perturbed, and muscle mass was

162 reduced in KPCY-PTHrP^{OE} tumor bearing mice (Figure 2G-H). Murf1 (*Trim63*) and

Atrogin1 (*Fbxo32*) expression were increased in cachectic muscle from KPCY-PTHrP^{OE}

164 tumor bearing hosts relative to those with KPCY-EV tumors (Supplemental Figure 2E).

166 KPCY-PTHrP^{OE} tumors were larger than KPCY-EV tumors (Supplemental Figure 2F),

potentially complicating the interpretation that PTHrP is driving wasting phenotypes. To

168 control for this, we repeated the experiment in an independent cohort where primary

tumors were size matched (Supplemental Figure 2G), by harvesting mice at day 18 for

170 KPCY-EV tumors and day 10 for KPCY-PTHrP^{OE} tumors. Despite having less time to

induce cachexia, KPCY-PTHrP^{OE} tumor bearing animals showed increased adipose

172 wasting phenotypes in iWAT (Supplemental Figure 2H). At this early time point, quad

weights were not significantly different (Supplemental Figure 2I), highlighting that adipose

174 wasting precedes muscle wasting in this model and that PTHrP directly mediates adipose

wasting. Finally, to demonstrate that PTHrP acts directly on adipose tissue, we injected

176 NTB C57BL/6 animals with recombinant PTHrP (rPTHrP), which was sufficient to

decrease iWAT mass after just 5 days of treatment (Figure 2I-J). These results indicate

178 that tumor cell-derived PTHrP is sufficient to induce PDAC-associated cachexia, primarily

by facilitating adipose wasting.

182 ***Adipocyte-specific deletion of Pth1r attenuates PDAC-associated cachexia***

To decouple tumor growth and energy consumption cachectic phenotypes from direct

184   tumor-adipocyte crosstalk mediated wasting, we set out to disrupt PTHrP signaling by

conditionally deleting its cognate receptor, Parathyroid Hormone 1 Receptor (*Pth1r*), in

186   cachectic tissue. Analysis of the Genotype-Tissue Expression (GTEX) database indicated

that amongst tissues that undergo wasting in cachexia (adipose, skeletal muscle, heart,

188   etc.), adipose tissue has the highest levels of *PTH1R* expression, with very little to no

expression in skeletal muscle (Figure 3A; all tissues shown in Supplemental Figure 3A).

190   To confirm this phenomenon in mice, we isolated RNA from multiple organs of non-tumor

bearing C57BL/6 mice and performed qPCR for *Pth1r*, which indicated that adipose tissue

192   depots were the most highly expressing organ that undergoes cachectic wasting (Figure

3B). Of note, *Pth1r* levels are highest in the kidney, consistent with previous studies [24]

194   and likely unrelated to cachectic wasting. Additionally, analysis of a single-cell atlas of

human adipose tissue [25] indicated that *PTH1R* is most highly expressed in adipocytes

196   and pre-adipocytes (adipose stem and progenitor cells or ASPCs; Figure 3C), suggesting

that adipocyte lineages are the most likely cell in adipose tissue to respond to tumor-

198   derived PTHrP. Similar results were seen in the murine adipose tissue single-cell atlas

[25], with *Pth1r* being most highly expressed in adipocytes and pre-adipocytes

200   (Supplemental Figure 3B). Immunofluorescent staining of adipose tissue indicated that

PTH1R is expressed in adipocytes and is increased in cachectic animals (Figure 3D).

202   Collectively, this indicated that PTH1R is highly expressed in adipocytes and is likely to

mediate PTHrP-PTH1R tumor to adipose crosstalk.

To functionally interrogate the role of the PTHrP-PTH1R signaling axis in cachectic

206   wasting, we deleted *Pth1r* specifically in adipocytes (i.e. *Adiponectin-Cre; Pth1r^LoxP/LoxP^*

mice, herein *Pth1r^Adipo-cKO^*). When implanted with KPCY-PTHrP^OE^ tumors, adipose tissue

208   wasting was reduced in *Pth1r^Adipo-cKO^* mice relative to WT hosts, with the loss of *Pth1r* in

adipocytes blocking tumor-induced morphological changes to the adipose tissue (Figure

210   3E-F). *Pth1r^Adipo-cKO^* tumor bearing hosts exhibited increased lipid droplet size (Figure 3G-

H; Supplemental Figure 3C), indicating that cachectic adipose tissue wasting was

212   reduced when PTHrP-PTH1R signaling in adipocytes is disrupted. Remarkably, blocking

PTHrP-PTH1R tumor to adipocyte signaling preserved muscle architecture and reduced

214   wasting (Figure 3I-J). In KPCY-PTHrP^OE^ tumor bearing mice, Murf1 (*Trim63*) and Atrogin1

(*Fbxo32*) expression were increased in cachectic muscle from WT hosts, which was

216   reduced in *Pth1r^Adipo-cKO^* hosts (Supplemental Figure 3D). Thus, tumor cell-secreted

PTHrP initiates PTH1R-mediated adipose and muscle wasting.

PTHrP-mediated adipose tissue wasting downregulates de novo lipogenesis

220   To determine the downstream mechanism by which PTHrP-PTH1R signaling axis

mediates cachectic adipose wasting, we performed bulk RNA-sequencing (RNA-seq)

222   analysis on iWAT from cachectic (KPCY-PTHrP^OE^) and non-cachectic (KPCY-EV) tumor

bearing mice (Figure 4A). Cachectic adipose tissue from KPCY-PTHrP^OE^ tumor bearing

224   animals and non-cachectic adipose tissue from KPCY-EV clustered separately by

differential gene expression analysis (Supplemental Figure 4A) and Principal Component

226   Analysis (PCA; Supplemental Figure 4B), indicating broad transcriptomic changes

between the groups, consistent with the striking morphological changes we observed in

228   the cachectic tissues. The top 50 differentially expressed downregulated genes in KPCY-

PTHrP^OE tumor bearing hosts relative to KPCY-EV tumor bearing hosts were comprised

230   of known regulators of adipocyte cell biology, including *Fasn*, *Pnpla3*, *Nnat*, and *Orm1*

(Supplemental Figure 4C; Supplemental Table 1). The top upregulated genes consisted

232   of immune cell markers such as *Marco*, *Ccr6*, and *Cxcr5* (Supplemental Figure 4C;

Supplemental Table 1).

Gene set enrichment analysis (GSEA) indicated that adipose from highly cachectic KPCY-

236   PTHrP^OE tumor bearing animals upregulated hallmark mediators of cancer cachexia,

such as the TNF-α and IL-6 pathways (Figure 4B-C) [10], consistent with the cachexia-

238   inducing ability of PTHrP [26] . Surprisingly, in addition to these known pro-cachectic

pathways, the adipose from KPCY-PTHrP^OE tumor bearing animals dramatically

240   downregulated adipogenesis and fatty acid metabolism pathways (Figure 4B-C),

suggesting a perturbed regulation of lipid homeostasis in adipocytes during cachexia.

242   Consistent with this pathway analysis, key fatty acid metabolism genes were markedly

downregulated in the adipose of highly cachectic animals, including a downregulation of

244   *de novo* lipogenesis (DNL) enzymes (*Acss2*, *Acly*, *Acaca*, and *Fasn*), lipases (*Lep*, *Lipe*,

Pbpla2, *Plin1/2*), and fatty acid metabolism genes (*FABPs*, *Prdm16*, *Ppargc1a*, *Elovl6*)

246   (Figure 4D). The vast majority of DNL pathway components were downregulated in the

adipose from cachectic KPCY-PTHrP^OE mice relative to non-cachectic KPCY-EV control

248   mice (Figure 4E; red arrows indicate downregulated genes). Notably, *Fasn* was the most

significantly downregulated DNL gene (Figure 4A) and one of the top 15 overall

250   downregulated genes (Supplemental Figure 4C). Thus, tumor-derived PTHrP significantly

downregulates adipose DNL to putatively facilitate cachectic wasting.

Adipocyte specific disruption of de novo lipogenesis sensitizes adipose tissue to
254   ***wasting***

To confirm that DNL pathway components were downregulated in response to PTHrP, we

256   performed qPCR on adipose tissue from PTHrP-overexpressing tumor bearing animals,

which showed lower *Acss2*, *Acly*, *Acaca*, and *Fasn* expression (Figure 5A). Western

258   blotting analysis further confirmed the downregulation of DNL enzymes in cachectic iWAT

(Figure 5B). Finally, qPCR demonstrated that master transcriptional regulators of DNL,

260   *Mlxipl* (ChREBPα and CHREBPβ isoforms) and *Srebf1c* (SREBP), were downregulated

in the adipose of PTHrP-overexpressing tumor bearing mice (Figure 5C).

To determine the functional role of DNL in PDAC-associated adipose tissue wasting, we

264   implanted the non-cachectic KPCY PDAC line in *Adiponectin-Cre; Fasn^{LoxP/LoxP}* mice

(herein *Fasn^{Adipo-cKO}*; knockout confirmed in iWAT in Supplemental Figure 5A). In this

266   context, the loss of adipocyte *Fasn* resulted in perturbed tissue histology and significantly

reduced adipose tissue mass (Figure 5D-E), demonstrating that the disruption of DNL in

268   adipose tissue is sufficient to prime the cachectic state. In addition to worsened adipose

tissue wasting, *Fasn^{Adipo-cKO}* tumor bearing animals had reduced adipocyte size (Figure

270   5F; Supplemental Figure 5B) and smaller PLIN1-positive lipid droplets (Figure 5G). Loss

of adipose *Fasn* did not alter primary tumor growth (Supplemental Figure 5C) or quad

272   wasting or pathology (Supplemental Figure 5D-E). Collectively, this data shows a

previously unappreciated role for DNL in regulating PDAC-associated adipose tissue

274   wasting.

276   ***Anti-PTHrP therapy blocks PDAC-associated cachexia***

No effective therapies currently exist to treat PDAC-associated cachexia. Given our

278   genetic data indicating that the deletion of tumor cell *Pthlh* or adipocyte *Pth1r* reduces

cachexia, we set out to determine if anti-PTHrP therapy is an effective means to disrupt

280   PTHrP-PTH1R tumor-adipose crosstalk and block cachectic wasting in PDAC. We have

previously demonstrated that monoclonal neutralizing antibodies directed against PTHrP

282   (anti-PTHrP) effectively block PTHrP signaling in PDAC tumor cells, reducing primary

tumor burden and metastatic ability [19]. To test the efficacy of anti-PTHrP as a cachexia-

284   blocking therapy in autochthonous KPCY tumors, we enrolled mice when their tumors

reached 75 mm^3, at which point we began anti-PTHrP or anti-IgG antibody treatment.

286   Anti-PTHrP therapy normalized adipose tissue architecture (Figure 6A), reduced adipose

wasting (Figure 6B), and increased adipocyte lipid size (Figure 6C; Supplemental Figure

288   6A) in KPCY mice. Anti-PTHrP treatment of KPCY mice trended towards increased quad

mass and normalized morphology (Figure 6D-E). Murf1 (*Trim63*) and Atrogin1 (*Fbxo32*)

290   were mildly induced in the muscles of tumor-bearing animals at this stage, with anti-

PTHrP antibody reducing their expression (Supplemental Figure 6B). Similar to the

292   GEMMs, the treatment of orthotopic KPCY tumors (from a cachectic clonal KPCY cell

line) with anti-PTHrP therapy showed a similar reduction in cachexia phenotypes, with

294   normalization of the adipose tissue architecture (Figure 6F), reduced adipose tissue

wasting (Figure 6G), and increased lipid droplet size (Figure 6H; Supplemental Figure

6C). Additionally, anti-PTHrP treated orthotopic tumor-bearing animals had reduced muscle wasting phenotypes (Figure 6I-J). The muscles of anti-IgG-treated orthotopic mice

showed evidence of modest Murf1 (*Trim63*) and Atrogin1 (*Fbxo32*) induction, which was reduced in anti-PTHrP-treated mice (Supplemental Figure 6D). Both KPCY GEMMs and

orthotopic tumor bearing animals treated with anti-PTHrP had increased lipid droplet size as visualized by PLIN1 staining of iWAT (Figure 6K-L). Finally, RNA-seq was performed

on iWAT from anti-PTHrP (or anti-IgG control) treated orthotopic KPCY tumor bearing animals. Similar to the genetic disruption of PTHrP-PTH1R signaling, anti-PTHrP

treatment reduced pro-inflammatory TNF-α signaling and enhanced adipogenesis and fatty acid metabolism signaling pathways in adipose tissue (Figure 6M). Similarly, iWAT

from anti-PTHrP treated animals upregulated *Fasn* (Figure 6N) and other DNL enzymes (Supplemental Figure 6E) relative to anti-IgG. Thus, anti-PTHrP monoclonal antibody

therapy is an effective means to reduce cachexia in multiple PDAC mouse models, primarily through a reduction in adipose tissue wasting phenotypes by restoring DNL

pathway expression.

DISCUSSION

Pancreatic cancer remains the lowest 5-year survival rate amongst the common cancers, in large part due to the highest prevalence and severity of cancer cachexia [3]. Cachectic wasting of both the adipose and muscle reduces PDAC patients' quality of life and overall survival [4, 27], and there are very few treatment options for the clinical management of cancer cachexia. To effectively manage cachexia in the clinic, new therapeutic targets must be identified that directly mediate wasting phenotypes. In our prior work, we identified the oncogenic and pro-metastatic molecule PTHrP as a tumor-derived secreted factor that correlates with decreased survival in human PDAC patients and whose genetic deletion or pharmacological inhibition in PDAC mouse models greatly enhances survival [19]. Herein, we expand upon these initial findings to show that PTHrP is responsible for PDAC-associated adipose tissue wasting in cachexia (Figure 7). Thus, targeting PTHrP therapeutically may be a multi-pronged attack to simultaneously block cancer cachexia, tumor growth, and metastasis. We show that cachectic patients have higher tumor PTHrP expression relative to non-cachectic patients, and that the genetic deletion or pharmacological inhibition of PTHrP reduces cachectic wasting and enhances overall survival in mouse models of PDAC-associated cancer cachexia. Induced expression of *Pthlh* in a lowly cachectic PDAC tumor cell line was sufficient to induce adipose and muscle wasting, ultimately leading to reduced overall survival in mice. Disruption of PTHrP-PTH1R tumor to adipose crosstalk via adipocyte-specific conditional deletion of *Pth1r* reduced adipose and muscle wasting, demonstrating that direct tumor cell to adipocyte signaling networks mediate adipose wasting that ultimately fuels muscle wasting. Mechanistically, we found that cachectic adipose tissues lose core DNL signaling

components while concomitantly upregulating pro-cachectic factors such as TNF-α and

336   IL-6. Finally, the adipose-specific deletion of *Fasn* sensitized animals towards adipose

wasting, demonstrating that loss of DNL signaling primes the cachectic state. As such,

338   our results reveal a previously unappreciated PTHrP-PTH1R-DNL signaling axis that

drives cancer-associated cachexia and provides a promising target to ameliorate

340   cachectic wasting in PDAC patients.

342   Cancer cachexia involves complex multi-organ metabolic crosstalk [10], with tumor cells

sending and receiving multiple inputs from systemic paracrine/endocrine signaling

344   networks. Several mechanisms have been identified to date to indicate how tumor cells

communicate with the muscle to facilitate muscle wasting and catabolism, but few studies

346   have identified mediators of direct tumor to adipocyte crosstalk as drivers of adipose

tissue wasting and subsequent muscle wasting [28]. This is important clinically, as

348   adipose tissue wasting precedes muscle wasting in PDAC patients and can occur up to

1.5 years prior to diagnosis [29]. Similar phenomena have been observed in other cancers

350   as well [14, 30, 31]. Our adipocyte-specific deletion of *Pth1r* interestingly halted both

adipose wasting and muscle wasting, suggesting that adipose wasting in this model may

352   fuel muscle wasting. Future studies will seek to identify and functionally characterize how

these adipose tissue-secreted signaling molecules, called "adipokines", mediate muscle

354   wasting, either through direct catabolism or by altering the metabolic status of the host.

356   Adipose tissue wasting during cancer cachexia has primarily been attributed to the

induction of a UCP1-driven thermogenic program that promotes adipocyte browning [32].

358   Indeed, PTHrP is known to trigger adipose tissue browning in the Lewis Lung Carcinoma

        (LLC) model of cancer cachexia [6], and we find a modest induction of adipose UCP1 in

360   our PDAC-associated cachexia models (Figure 4D). However, we found a more profound

        downregulation of *de novo* lipogenesis in our system that suppresses lipid production in

362   adipocytes and primes them for cachectic wasting. Thus, harnessing new therapies to

        block or reverse the loss of DNL hold promise as a means to reverse cachectic wasting

364   of adipose tissue depots. One such approach that shows promise in preclinical models is

        the Peroxisome proliferator-activated receptor alpha (PPARα) agonist fenofibrate, which

366   has been shown to rescue reduced *Acly*, *Fasn*, *Acsl1*, and *Acsl5* in hepatic *de novo*

        lipogenesis, thus restoring lipid homeostasis in the liver and improving overall survival

368   [16]. Importantly, our data indicate that PPARα (encoded by *Ppara*) is still expressed,

        albeit at slightly lower levels, in the adipose of PTHrP tumor bearing animals, and thus its

370   reactivation is an attractive therapeutic target to pursue in future studies.

372   Prior correlative studies in a small cohort consisting of cachectic (weight losing) and non-

        cachectic (weight stable) patients suggested that enhanced adipocyte lipolysis, but *not*

374   reduced lipogenesis was involved in adipose tissue loss in cancer cachexia [33].

        Provocatively, other work has demonstrated that whole body lipolytic rate in cancer

376   patients is not different from healthy controls, suggesting that the loss of body fat in cancer

        cachexia is due to reduced lipogenesis rather than augmented lipolysis [34]. The latter is

378   supported by clinical data to show that lipogenic enzymes such as FASN are reduced in

        adipose from cachectic patients [35, 36], consistent with our experiments showing that

380   PTHrP mediates loss of DNL in cachectic adipose tissue. Nonetheless, subsequent

functional studies in the field have primarily focused on the role of enhanced lipolysis, and

382   not reduced DNL, in mediating adipose tissue loss. This work has confirmed that lipases

are enriched in the cachectic state, which can be blocked through the inhibition of key

384   lipases such as adipose triglyceride lipase (ATGL) or hormone-sensitive lipase (HSL) to

limit adipose wasting [14]. Surprisingly, in PTHrP-driven adipose wasting, the expression

386   of most cachexia-associated lipases, such as ATGL (*Pnpla2*), HSL (*Lipe*), and

Monoglyceride Lipase (*Mgll*), were unchanged or even reduced in the cachectic state,

388   suggesting that they are not driving the adipose tissue wasting phenotype in this model

or that their regulation occurs post-transcriptionally. Rather, our work demonstrates that

390   the loss of *de novo* lipogenesis, as opposed to the activation of lipases, in adipocytes

primes the host for cachectic adipose tissue wasting. Consistent with this working model,

392   prior work has demonstrated that the decrease in adipose tissue mass is due to reduced

adipocyte size and not from a reduction in the total number of adipocytes (i.e. adipocyte

394   death) [33, 37]. It is likely that the balance between adipose tissue lipolysis and

lipogenesis is altered during progressive cachexia-associated adipose wasting, and thus

396   different models may find varying functional roles for these enzymes during different

stages of cachexia progression. Nonetheless, drugs targeting lipases and DNL

398   components will need to achieve a balance to restore lipid homeostasis and block adipose

wasting.

PDAC patients with higher serum levels of PTHrP lose significantly more weight, have

402   reduced handgrip strength, and score lower on the Karnofsky Performance Status (KPS)

scale relative to those with lower serum PTHrP, indicative of enhanced cachectic burden

[23]. Additionally, whole body fat oxidation is increased in PDAC patients with high PTHrP, suggesting a perturbed homeostasis between lipolysis and lipogenesis [23], in agreement

with our preclinical experiments in mice. Critically, in our models, disrupting direct tumor-to-adipocyte crosstalk by deleting *Pth1r* on adipocytes was sufficient to block cachectic

wasting, thereby setting the stage for PTHrP- or PTH1R-targeted therapeutic interventions such as the anti-PTHrP monoclonal antibody therapy that we used. This

result, combined with our prior work showing that PTHrP facilitates tumor growth and metastasis, demonstrates that blocking PTHrP-PTH1R signaling axis is a potent means

to block many of the confounding factors in PDAC that lead to its dismal overall survival rate. Interestingly, PTHrP/*PTHLH* is a squamous identity gene and is highly expressed in

many squamous carcinomas, including the squamous/basal-like subtype of PDAC [19], lung squamous cell carcinoma [38], head and neck squamous cell carcinoma [39], oral

squamous cell carcinoma [40], and cervical squamous cell carcinoma [41]. Many of these squamous cancers also have high levels of cancer-associated cachexia, suggesting that

targeting PTHrP in other cancers to block cachectic wasting warrants further preclinical testing. In a recent pan-cancer analysis, patients with high serum PTHrP levels lost ~7%

of their body weight independent of tumor burden, while those without PTHrP in their serum lost only ~1% [42], highlighting that PTHrP may also drive wasting in other cancers.

Notably, the Lewis lung cancer model of cancer cachexia was similarly responsive to anti-PTHrP therapy [6], perhaps pointing towards a more broadly applicable mechanism of

PTHrP-induced cachectic wasting across cancer types.

METHODS

426 **_Mice_**

$Kras^{LSL-G12D}$; $Trp53^{LSL-R172H}$; $Pdx1$-Cre; $Rosa26^{LSL-YFP}$ (KPCY) and $Kras^{LSL-G12D}$; $Trp53^{LSL-}$

428 ^{R172H}; $Pdx1$-Cre; $Rosa26^{LSL-YFP}$; $Pthlh^{Loxp/wt}$ (KPCY-Pthlh^{Het}) mice have been described

previously [18, 19]. $Pthlh^{LoxP}$ mice were a generous gift from Dr. Andrew Karaplis (McGill

430 University) [43]. $Pth1r^{LoxP}$ mice were a generous gift from Dr. Henry Kronenberg (Harvard

University) [44]. Fasn^{LoxP} mice were generously provided by Dr. Clay Semenkovich

432 (Washington University at St. Louis) [45]. Adiponectin-Cre and wild-type C57BL/6J mice

were purchased from the Jackson Laboratory (Stock # 028020 and #000664,

434 respectively). $Fasn^{LoxP}$ or $Pthlh^{LoxP}$ mice were crossed with Adiponectin-Cre mice to

generate $Adiponectin$-Cre; $Fasn^{LoxP/LoxP}$ (Fasn$^{Adipo-cKO}$) and $Adiponectin$-Cre; $Pth1r^{LoxP/LoxP}$

436 (Pth1r$^{Adipo-cKO}$) lines, respectively. For survival analysis, mice were palpated for tumors

and examined for evidence of morbidity twice weekly and sacrificed when moribund. Mice

438 treated with monoclonal neutralizing antibody against PTHrP (anti-PTHrP; generously

provided by Dr. Richard Kremer, McGill University; 200µg/injection) or anti-IgG control

440 were injected three times per week every 48 hours for the indicated time. Mice were

treated with rPTHrP (1mg/kg; Bachem T-4512) for 5 consecutive days, and tissues were

442 harvested within 1 hour of the final dosing. All vertebrate animals were maintained, and

experiments were conducted in compliance with the NIH guidelines for animal research

444 and approved by the University of Massachusetts Chan Medical School Institutional

Animal Care and Use Committee.

Genotyping

448   Mice were genotyped by automated genotyping services via TransnetYX for the following

alleles: *Kras*^{LSL-G12D}, *Trp53*^{LSL-R172H}, *Pdx1-Cre*, *Rosa26*^{LSL-YFP}, *Pth1r*, *Adiponectin-Cre*,

450   and *Pthlh*^{Loxp}. For Fasn^{Adipo-cKO} mice, conventional PCR genotyping reactions were

performed. Briefly, tails were cut and immersed in lysis buffer (40% NaOH + 0.5M EDTA

452   pH 8) for one hour and incubated at 95°C, followed by addition of neutralization buffer

(40mM Tris-HCl). PCR reactions were carried out with the following primers:

454   *Fasn forward: 5' - GGA TAG CTG TGT AGT GTA ACC AT -3'*

Fasn reverse: 5'- GGT CAT CGT GAT AAC CAC ACA T -3'

456   *Adipo-Cre transgene forward: 5'- ACG GAC AGA AGC ATT TTC CA -3'*

Adipo-Cre transgene reverse: 5'- GGA TGT GCC ATG TGA GTC TG - 3'

458   *Adipo-Cre internal control forward: 5'- CTA GGC CAC AGA ATT GAA AGA TCT -3'*

Adipo-Cre internal control reverse: 5'- GTA GGT GGA AAT TCT AGC ATC ATC C -3'

Orthotopic injection

462   Orthotopic injection was performed as previously described [46]. Briefly, C57BL/6 mice

were anesthetized using isoflurane. A small incision was made on the upper left quadrant

464   of the mouse, and the pancreas was exteriorized on a sterile field. 1.0×10^5 cells were

injected into the pancreas in 100ul sterile PBS via an insulin syringe. A cotton swab was

466   held over the site of injection. After injection, the pancreas was placed back into the

peritoneal cavity and the incision was sutured with 4-0 Vicryl sutures. The mice received

468   meloxicam post-surgery and were monitored daily following surgery to assess post-

surgery recovery. For each experiment, the sex of mice was matched to the sex of the

470   KPCY cell line.

Immunohistochemistry and immunofluorescence staining

For staining of mouse tissues, zinc formalin-fixed paraffin embedded (FFPE) tissues were

474 dewaxed by baking in the oven at 60°C for an hour. Following dewaxing, tissues were

rehydrated by serial immersions in xylene-substituent histoclear and ethanol gradients.

476 Antigen retrieval was performed using citric acid-based antigen retrieval buffer (Vector

laboratories H-3300). Plin1 (Cell signaling technologies 9349), PTHrP (Sigma AV33885),

478 and PTH1R (Abcam ab189924) primary antibodies were used at 1:1000 dilution overnight

at 4°C. Tissues were incubated with either HRP-tagged secondary antibody (Vector

480 laboratories MP-7401) or donkey anti-rabbit Alexa fluor 594 (Fisher Scientific Cat

#A21207) at room temperature for 1 hour and followed with hematoxylin counterstain or

482 mounted with DAPI mounting media. Images were acquired on a Leica Thunder Imager

or EVOS M5000.

Human specimens

486 Primary PDAC patient tumor samples and associated omental adipose tissues were

obtained from decedents who had previously been diagnosed with pancreatic ductal

488 adenocarcinoma from the University of Nebraska Medical Center's Tissue Bank through

the Rapid Autopsy Program (RAP) in compliance with IRB 091-01. Non-cancer tissues

490 are collected in a manner similar to RAP specimens through the UNMC Normal Organ

Recovery (NORs) Program in association with LiveOn Nebraska. To ensure specimen

492 quality, organs were harvested within three hours post-mortem and the specimens flash

frozen in liquid nitrogen or placed in formalin for immediate fixation. Sections were cut

494   from paraffin blocks of formalin fixed tissue into 4-micron thick sections and mounted on

charged slides. Pancreatic cancer patients with a known history of body weight loss, as

496   confirmed by the pathologist, were binned into cachectic and non-cachectic cohorts. The

primary tumors were processed and stained for PTHrP immunohistochemistry as

498   described above. The slides were scanned by the UMass Chan SCOPE core facility using

a SL Slide Scanner, followed by PTHrP H Score quantification using QuPath software.

Cell lines and culture conditions

502   Murine PDAC cell lines were derived from KPCY mice that were backcrossed onto the

C57BL/6J strain [47]. KPCY-EV and KPCY-PTHrP^{OE} cells were described previously [19].

504   All cell lines were maintained in Dulbecco's modified eagle medium (DMEM) with 10%

fetal bovine serum (FBS) and 1% penicillin/streptomycin (P/S) at 5% CO_2 at 37°C. Cell

506   lines were tested regularly for *Mycoplasma*.

Cell proliferation assay

508   The WST-1 cell proliferation assay (ab65475, Abcam) was performed per the

510   manufacturer's instructions. Briefly, 2.0×10^3 cells were plated on 96-well plates and

grown in DMEM supplemented with 10% FBS and 1% P/S. Fresh media containing 10%

512   WST solution was incubated on the cells for 2 hours at 37°C. After shaking the plate, the

absorbance was read at 420 to 480 nm with a reference wavelength of 650 nm. Final

514   absorbance was obtained by subtracting the 650-nm reference wavelength from the 420-

to 480-nm read.

Western blot analysis

Inguinal white adipose tissues were snap-frozen in liquid nitrogen immediately after retrieving from orthotopic and GEM mice. The iWAT's were transferred to 2ml Eppendorf

tubes with pre-cooled metal bead and homogenized using Cryomill (Retsch) for 3 minutes to get a fine powder. The powderized iWAT's were then lysed in RIPA buffer containing

protease and phosphatase inhibitors (Sigma 04693124001 and Roche 04906837001). The samples were centrifuged at 15,000 g for 15 min at 4°C, and the supernatant was

collected for western blotting. Lysates were run on NuPAGE gels (Thermo Fisher NP0336BOX), transferred to PVDF membranes, and stained with primary antibodies:

Fasn (Cell signaling technologies 3180), Acly (Proteintech 15421-1-1 AP), Acaca (Cell signaling technologies 3662) and beta-actin (Sigma A5316). Membranes were washed

and stained with secondary antibodies: donkey anti-mouse IgG IRDye 800 (LICORbio 926-32212) or donkey anti-rabbit IgG IRDye 800 (LICORbio 926-32213) or donkey anti-

rabbit IgG IRDye 680 (LICORbio 926-68073) (1:15,000), washed, and imaged on a LI-COR Odyssey imaging system.

RNA extraction and qPCR

534   RNA was extracted from cryomilled adipose tissues using QIAzol/chloroform and Qiagen RNeasy kit (Cat #74106) following the manufacturer's protocol. For extracting RNA from

536   the iWAT, powderized tissues were resuspended in 1ml QIAzol per sample and lysed using TissueLyser II (Qiagen Cat#) and pre-cooled metal beads. The lysed sample was

538   mixed with chloroform and mixed vigorously, followed by centrifugation at 4°C for 10 minutes at 10,000g. The upper aqueous phase was mixed with 70% ethanol in 1:1 ratio

540 and transferred to the RNA spin columns from the Qiagen RNeasy kit and spun for 20

seconds at 9000g. All the remaining steps were followed as per manufacturer's

542 guidelines. Extracted RNA was quantified using Nanodrop and cDNA was prepared using

100ng-1000ng RNA using the New England Biolabs LunaScript RT supermix (Cat#

544 M3010L). Quantitative PCR was setup using Luna Universal qPCR Master Mix (Cat

#M3003E). The following primers were used:

546 *Pth1r forward: 5'- TCT GCA ATG GTG AGG TGC AG - 3'*

Pth1r reverse: 5'- GCT ACT CCC ACT TCG TGC TT - 3'

548 *Fasn forward: 5'- GGA GGT GGT GAT AGC CGG TAT - 3'*

Fasn reverse: 5'- TGG GTA ATC CAT AGA GCC CAG - 3'

550 *Acly forward: 5'- ACC CTT TCA CTG GGG ATC ACA -3'*

Acly reverse: 5'- GAC AGG GAT CAG GTA TTC CTT G - 3'

552 *Acaca forward: 5'- ATT GTG GCT CAA ACT GCA GGT - 3'*

Acaca reverse: 5'- GCC AAT CCA CTC GAA GAC CA -3'

554 *Murf1 forward: 5'- GGA CGG AAA TGC TAT GGA GAA CC CC-3'*

Murf1 reverse: 5'- GAT GGC TGT TTC CAC AAG CTT GG - 3'

556 *Atrogin1 forward: 5'- CCT CAG CAG TTA CTG CAA CAA GGA -3'*

Atrogin 1 reverse: 5'- CAT CTT CTT CCA ATC CAG CTG C - 3'

558 *Chrebp-a forward: 5'- CGA CAC TCA CCC ACC TCT TC -3'*

Chrebp-a reverse: 5' - TTG TTC AGC CGG ATC TTG TC - 3'

560 *Chrebp-b forward: 5' - TCT GCA GAT CGC GTG GAG -3'*

Chrebp-b reverse: 5'- CTT GTC CCG GCA TAG CAA C -3'

562 *Srebf1c forward: 5'- AAG CAA ATC ACT GAA GGA CCT GG - 3'*

Srebf1c reverse: 5'- AAA GAC AAG CTA CTC TGG GAG -3'

564   *18S RNA forward: 5'- CGA ACG TCT GCC CTA TCA ACT T - 3'*

        18S RNA reverse: 5'- CCG GAA TCG AAC CCT GAT T - 3'

RNA-seq library prep and GSEA

568   Frozen iWAT tissues were cryomilled and RNA was isolated using Qiazol and Qiagen

        RNAeasy mini kits. A quantity of 500ng total RNA was used for poly(A) mRNA purification

570   using the NEBNext poly(A) mRNA magnetic isolation module (E7490L). Libraries were

        prepared using the NEBNext Ultra II Directional Library Prep Kit for Illumina (E7760L) as

572   per the manufacturer's recommendations and were barcoded using NEBNext Multiplex

        Oligos for Illumina (Dual Index Primers set 2 E7780S) during PCR amplification. Final

574   libraries were quantified using the NEBNext Library Quant Kit for Illumina (E7630L).

        Sequencing was performed using an Illumina NextSeq2000 with 115 bp single-end reads,

576   plus 8 cycles for barcodes. Fastq files were generated with Illumina DRAGEN BCL

        Convert v3.8.4. Heatmaps were generated by Morpheus (Broad Institute).

        The RNA-seq analysis was performed using OneStopRNAseq [48]. Briefly, FastQC and

580   MultiQC were used for raw reads quality control and QoRTs for post-alignment quality

        control [49, 50]. Reads were aligned to the reference genome assembly mm10 with

582   star_2.7.5a and annotated with gencode.vM25.primary_assembly [51, 52]. Aligned exon

        reads were counted toward gene expression with featureCounts_2.0.0 with default

584   settings [53]. Differential expression (DE) analysis was performed with DESeq2_1.28.1

        [54]. Significantly differentially expressed genes (DEGs) were filtered with the criteria FDR

586   < 0.05 (|LFC|) > 0.585. Gene set enrichment analysis were performed with GSEA [55].

Heatmaps of TPM values were generated by Morpheus (Broad Institute).

Adipocyte size quantification

590   Inguinal white adipose tissues were harvested, sectioned, and stained with hematoxylin

and eosin. H&E stained tissues were imaged at 20X magnification and adipocyte size

592   (area in μm^2) was quantified using Adiposoft plugin in ImageJ software [56]. All of the

adipocyte areas counted by Adiposoft were included in the analysis and are represented

594   as relative frequencies.

596   ***Lentiviral transduction***

KPCY-PTHrP^{OE} cells were previously generated by lentiviral-mediated overexpression of

598   *Pthlh* [19]. Briefly, KPCY cells were transduced with pLenti-C-Myc-DDK-P2A-Puro

Lentiviral Particles with EV (Origene PS100092) or Pthlh ORF (Origene MR201519L3)

600   and stably selected with puromycin (8 µg/ml). Stable KPCY-PTHrP^{OE} cells were

maintained with puromycin after selection.

Statistical analysis

604   All data points are biological replicates and represent individual mice, tissues, or cell

samples. The data was first tested for normal distribution, followed by the appropriate

606   statistical test as indicated in the figure legends. Data are represented as mean ±

standard deviation; **P<0.05*; non-significant *P* values are indicated in the figure text or

608   legend. All statistical analysis was performed using GraphPad Prism.

610 ***Data and code availability***

All sequencing data have been deposited in the Gene Expression Omnibus under the

612 series GSE298308.

ACKNOWLEDGEMENTS

616   This work was supported by an American Gastroenterology Association Bern Schwartz

Research Scholar Award in Pancreatic Cancer (JRP); National Institute of Health (NIH)

618   K99-R00 CA252153 (JRP); US Department of Defense (DOD) Pancreatic Cancer

Research Program (PCARP) Idea Development Award PA230161P1 (JRP and DG);

620   Hopper-Belmont Inspiration Award (NB); Pancreatic Cancer Alliance (PCA) Postdoctoral

Fellowship (NB); NIH Innate Immunity Training Program T32 AI095213 (JP); NIH Institute

622   for Maximizing Student Development T32 GM135751 (CJ); Canadian Institutes of Health

Research (CIHR) MOP142287 (RK); Pancreatic Cancer Detection Consortium,

624   U01CA210240; NCI Cancer Center Support Grant, P30CA36727; NCI Research

Specialist, R50CA211462; NIH DK130852 (MPC); Breast Cancer Alliance Young

626   Investigator Grant (EVW); Massachusetts Life Sciences Center Bits to Bytes grant (to

Drs. Dori Schafter and Christina Baer); and Massachusetts Life Sciences Center

628   Research Infrastructure grant (to Drs. Kate Fitzgerald and Christina Baer). We

acknowledge and thank the Sanderson Center for Optical Experimentation (SCOPE; Dr.

630   Christina Baer and Jill McConnell) and Morphology core facilities at the University of

Massachusetts Chan Medical School; and Pancreatic Cancer Mouse Hospital and the

632   Molecular Pathology and Imaging Core at the University of Pennsylvania.

634   RK is the inventor of the therapeutic monoclonal anti-PTHrP antibodies used in this

study and authorized for use by Miramab Bioscience. RK holds an equity stake in

636   Miramab Bioscience.

REFERENCES

1.   Siegel, R.L., A.N. Giaquinto, and A. Jemal, *Cancer statistics, 2024.* CA Cancer J Clin, 2024. **74**(1): p. 12-49.

2.   *American Cancer Society: Cancer Facts & Figures 2025.* Atlanta: American Cancer Society, 2025

3.   Baracos, V.E., L. Martin, M. Korc, D.C. Guttridge, and K.C.H. Fearon, *Cancer-associated cachexia.* Nat Rev Dis Primers, 2018. **4**: p. 17105.

4.   Henderson, S.E., N. Makhijani, and T.A. Mace, *Pancreatic Cancer-Induced Cachexia and Relevant Mouse Models.* Pancreas, 2018. **47**(8): p. 937-945.

5.   Narsale, A.A. and J.A. Carson, *Role of interleukin-6 in cachexia: therapeutic implications.* Curr Opin Support Palliat Care, 2014. **8**(4): p. 321-7.

6.   Kir, S., J.P. White, S. Kleiner, L. Kazak, P. Cohen, et al., *Tumour-derived PTH-related protein triggers adipose tissue browning and cancer cachexia.* Nature, 2014. **513**(7516): p. 100-4.

7.   Patel, H.J. and B.M. Patel, *TNF-alpha and cancer cachexia: Molecular insights and clinical implications.* Life Sci, 2017. **170**: p. 56-63.

8.   Ling, T., J. Zhang, F. Ding, and L. Ma, *Role of growth differentiation factor 15 in cancer cachexia (Review).* Oncol Lett, 2023. **26**(5): p. 462.

9.   Groarke, J.D., J. Crawford, S.M. Collins, S. Lubaczewski, E.J. Roeland, et al., *Ponsegromab for the Treatment of Cancer Cachexia.* N Engl J Med, 2024. **391**(24): p. 2291-2303.

10.   Ferrer, M., T.G. Anthony, J.S. Ayres, G. Biffi, J.C. Brown, et al., *Cachexia: A systemic consequence of progressive, unresolved disease.* Cell, 2023. **186**(9): p. 1824-1845.

11.   Fearon, K., F. Strasser, S.D. Anker, I. Bosaeus, E. Bruera, et al., *Definition and classification of cancer cachexia: an international consensus.* Lancet Oncol, 2011. **12**(5): p. 489-95.

12.   Weber, B.Z.C., D.H. Arabaci, and S. Kir, *Metabolic Reprogramming in Adipose Tissue During Cancer Cachexia.* Front Oncol, 2022. **12**: p. 848394.

13.   Petruzzelli, M., M. Schweiger, R. Schreiber, R. Campos-Olivas, M. Tsoli, et al., *A switch from white to brown fat increases energy expenditure in cancer-associated cachexia.* Cell
Metab, 2014. **20**(3): p. 433-47.

    14.   Das, S.K., S. Eder, S. Schauer, C. Diwoky, H. Temmel, et al., *Adipose triglyceride lipase
666   contributes to cancer-associated cachexia.* Science, 2011. **333**(6039): p. 233-8.

    15.   Fearon, K.C., D.J. Glass, and D.C. Guttridge, *Cancer cachexia: mediators, signaling, and
668   metabolic pathways.* Cell Metab, 2012. **16**(2): p. 153-66.

    16.   Yang, X., J. Wang, C.Y. Chang, F. Zhou, J. Liu, et al., *Leukemia inhibitory factor suppresses
670   hepatic de novo lipogenesis and induces cachexia in mice.* Nat Commun, 2024. **15**(1): p. 627.

672   17.   Hingorani, S.R., L. Wang, A.S. Multani, C. Combs, T.B. Deramaudt, et al., *Trp53R172H and KrasG12D cooperate to promote chromosomal instability and widely metastatic
674   pancreatic ductal adenocarcinoma in mice.* Cancer Cell, 2005. **7**(5): p. 469-83.

    18.   Rhim, A.D., E.T. Mirek, N.M. Aiello, A. Maitra, J.M. Bailey, et al., *EMT and dissemination
676   precede pancreatic tumor formation.* Cell, 2012. **148**(1-2): p. 349-61.

    19.   Pitarresi, J.R., R.J. Norgard, A.M. Chiarella, K. Suzuki, B. Bakir, et al., *PTHrP Drives
678   Pancreatic Cancer Growth and Metastasis and Reveals a New Therapeutic Vulnerability.* Cancer Discov, 2021. **11**(7): p. 1774-1791.

680   20.   Gumucio, J.P. and C.L. Mendias, *Atrogin-1, MuRF-1, and sarcopenia.* Endocrine, 2013. **43**(1): p. 12-21.

682   21.   Rupert, J.E., A. Narasimhan, D.H.A. Jengelley, Y. Jiang, J. Liu, et al., *Tumor-derived IL-6 and trans-signaling among tumor, fat, and muscle mediate pancreatic cancer cachexia.* J Exp
684   Med, 2021. **218**(6).

22.   Goncalves, M.D., S.K. Hwang, C. Pauli, C.J. Murphy, Z. Cheng, et al., *Fenofibrate prevents*
686   *skeletal muscle loss in mice with lung cancer.* Proc Natl Acad Sci U S A, 2018. **115**(4): p. E743-E752.

688   23.   Iresjo, B.M., S. Kir, and K. Lundholm, *Parathyroid hormone related protein (PTHrP) in patients with pancreatic carcinoma and overt signs of disease progression and host tissue*
690   *wasting.* Transl Oncol, 2023. **36**: p. 101752.

24.   Urena, P., X.F. Kong, A.B. Abou-Samra, H. Juppner, H.M. Kronenberg, et al., *Parathyroid*
692   *hormone (PTH)/PTH-related peptide receptor messenger ribonucleic acids are widely distributed in rat tissues.* Endocrinology, 1993. **133**(2): p. 617-23.

694   25.   Emont, M.P., C. Jacobs, A.L. Essene, D. Pant, D. Tenen, et al., *A single-cell atlas of human and mouse white adipose tissue.* Nature, 2022. **603**(7903): p. 926-933.

696   26.   De Miguel, F., P. Martinez-Fernandez, C. Guillen, A. Valin, A. Rodrigo, et al., *Parathyroid hormone-related protein (107-139) stimulates interleukin-6 expression in human*
698   *osteoblastic cells.* J Am Soc Nephrol, 1999. **10**(4): p. 796-803.

27.   Tuca, A., P. Jimenez-Fonseca, and P. Gascon, *Clinical evaluation and optimal management*
700   *of cancer cachexia.* Crit Rev Oncol Hematol, 2013. **88**(3): p. 625-36.

28.   Schmidt, S.F., M. Rohm, S. Herzig, and M. Berriel Diaz, *Cancer Cachexia: More Than*
702   *Skeletal Muscle Wasting.* Trends Cancer, 2018. **4**(12): p. 849-860.

29.   Sah, R.P., A. Sharma, S. Nagpal, S.H. Patlolla, A. Sharma, et al., *Phases of Metabolic and*
704   *Soft Tissue Changes in Months Preceding a Diagnosis of Pancreatic Ductal Adenocarcinoma.* Gastroenterology, 2019. **156**(6): p. 1742-1752.

706   30.   Rohm, M., M. Schafer, V. Laurent, B.E. Ustunel, K. Niopek, et al., *An AMP-activated protein kinase-stabilizing peptide ameliorates adipose tissue wasting in cancer cachexia in mice.*
708   Nat Med, 2016. **22**(10): p. 1120-1130.

31.   Agustsson, T., M. Ryden, J. Hoffstedt, V. van Harmelen, A. Dicker, et al., *Mechanism of*
710   *increased lipolysis in cancer cachexia.* Cancer Res, 2007. **67**(11): p. 5531-7.

32.   Kir, S. and B.M. Spiegelman, *Cachexia & Brown Fat: A Burning Issue in Cancer.* Trends
712   Cancer, 2016. **2**(9): p. 461-463.

33.   Ryden, M., T. Agustsson, J. Laurencikiene, T. Britton, E. Sjolin, et al., *Lipolysis--not*
714   *inflammation, cell death, or lipogenesis--is involved in adipose tissue loss in cancer cachexia.* Cancer, 2008. **113**(7): p. 1695-704.

716   34.   Jeevanandam, M., G.D. Horowitz, S.F. Lowry, and M.F. Brennan, *Cancer cachexia and the rate of whole body lipolysis in man.* Metabolism, 1986. **35**(4): p. 304-10.

718   35.   Notarnicola, M., A. Miccolis, V. Tutino, D. Lorusso, and M.G. Caruso, *Low levels of lipogenic enzymes in peritumoral adipose tissue of colorectal cancer patients.* Lipids, 2012. **47**(1): p.
720   59-63.

36.   Fang, R., L. Yan, and Z. Liao, *Abnormal lipid metabolism in cancer-associated cachexia and*
722   *potential therapy strategy.* Front Oncol, 2023. **13**: p. 1123567.

724 37. Bing, C., S. Russell, E. Becket, M. Pope, M.J. Tisdale, et al., *Adipose atrophy in cancer cachexia: morphologic and molecular analysis of adipose tissue in tumour-bearing mice.* Br J Cancer, 2006. **95**(8): p. 1028-37.

726 38. Hastings, R.H., F. Araiza, D.W. Burton, and L.J. Deftos, *Role of parathyroid hormone-related protein in lung cancer cell survival.* Chest, 2004. **125**(5 Suppl): p. 150S.

728 39. Chang, W.M., Y.F. Lin, C.Y. Su, H.Y. Peng, Y.C. Chang, et al., *Parathyroid Hormone-Like Hormone is a Poor Prognosis Marker of Head and Neck Cancer and Promotes Cell Growth*
730  *via RUNX2 Regulation.* Sci Rep, 2017. **7**: p. 41131.

 40. Lv, Z., X. Wu, W. Cao, Z. Shen, L. Wang, et al., *Parathyroid hormone-related protein serves*
732  *as a prognostic indicator in oral squamous cell carcinoma.* J Exp Clin Cancer Res, 2014. **33**(1): p. 100.

734 41. Kitazawa, S., R. Kitazawa, M. Fukase, T. Fujimori, and S. Maeda, *Immunohistochemical evaluation of parathyroid hormone-related protein (PTHrP) in the uterine cervix.* Int J
736  Cancer, 1992. **50**(5): p. 731-5.

 42. Hong, N., H.J. Yoon, Y.H. Lee, H.R. Kim, B.W. Lee, et al., *Serum PTHrP Predicts Weight Loss*
738  *in Cancer Patients Independent of Hypercalcemia, Inflammation, and Tumor Burden.* J Clin Endocrinol Metab, 2016. **101**(3): p. 1207-14.

740 43. He, B., R.A. Deckelbaum, D. Miao, M.L. Lipman, M. Pollak, et al., *Tissue-specific targeting of the pthrp gene: the generation of mice with floxed alleles.* Endocrinology, 2001. **142**(5):
742  p. 2070-7.

 44. Kobayashi, T., U.I. Chung, E. Schipani, M. Starbuck, G. Karsenty, et al., *PTHrP and Indian*
744  *hedgehog control differentiation of growth plate chondrocytes at multiple steps.* Development, 2002. **129**(12): p. 2977-86.

746 45. Chakravarthy, M.V., Z. Pan, Y. Zhu, K. Tordjman, J.G. Schneider, et al., *"New" hepatic fat activates PPARalpha to maintain glucose, lipid, and cholesterol homeostasis.* Cell Metab,
748  2005. **1**(5): p. 309-22.

 46. Aiello, N.M., A.D. Rhim, and B.Z. Stanger, *Orthotopic Injection of Pancreatic Cancer Cells.*
750  Cold Spring Harb Protoc, 2016. **2016**(1): p. pdb prot078360.

 47. Li, J., K.T. Byrne, F. Yan, T. Yamazoe, Z. Chen, et al., *Tumor Cell-Intrinsic Factors Underlie*
752  *Heterogeneity of Immune Cell Infiltration and Response to Immunotherapy.* Immunity, 2018. **49**(1): p. 178-193 e7.

754 48. Li, R., K. Hu, H. Liu, M.R. Green, and L.J. Zhu, *OneStopRNAseq: A Web Application for Comprehensive and Efficient Analyses of RNA-Seq Data.* Genes (Basel), 2020. **11**(10).

756 49. Ewels, P., M. Magnusson, S. Lundin, and M. Kaller, *MultiQC: summarize analysis results for multiple tools and samples in a single report.* Bioinformatics, 2016. **32**(19): p. 3047-8.

758 50. Hartley, S.W. and J.C. Mullikin, *QoRTs: a comprehensive toolset for quality control and data processing of RNA-Seq experiments.* BMC Bioinformatics, 2015. **16**(1): p. 224.

760 51. Dobin, A., C.A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1): p. 15-21.

762 52. Harrow, J., A. Frankish, J.M. Gonzalez, E. Tapanari, M. Diekhans, et al., *GENCODE: the reference human genome annotation for The ENCODE Project.* Genome Res, 2012. **22**(9):
764  p. 1760-74.

 53. Liao, Y., G.K. Smyth, and W. Shi, *featureCounts: an efficient general purpose program for*
766  *assigning sequence reads to genomic features.* Bioinformatics, 2014. **30**(7): p. 923-30.

54.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.* Genome Biol, 2014. **15**(12): p. 550.

55.   Subramanian, A., P. Tamayo, V.K. Mootha, S. Mukherjee, B.L. Ebert, et al., *Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles.* Proc Natl Acad Sci U S A, 2005. **102**(43): p. 15545-50.

56.   Galarraga, M., J. Campion, A. Munoz-Barrutia, N. Boque, H. Moreno, et al., *Adiposoft: automated software for the analysis of white adipose tissue cellularity in histological sections.* J Lipid Res, 2012. **53**(12): p. 2791-6.

FIGURE LEGENDS

Figure 1.

(A) Kaplan–Meier overall survival plots of KPCY and KPCY-Pthlh[HET] mice. Log-rank $P <$ 0.0001. **(B)** Percent change from baseline bodyweight in KPCY and KPCY-Pthlh[HET] mice.

(C) Carcass weights at endpoint from **(B)**. **(D-F)** Representative H&E images and quantification of normalized tissue masses for iWAT **(D)**, eWAT **(E)**, and quads **(F)** from

non-tumor bearing (NTB), KPCY, and KPCY-Pthlh[HET] mice at endpoint. Scale bars, 100 µm. **(G-I)** Representative H&E images and quantification of normalized tissue masses for

iWAT **(G)**, eWAT **(H)**, and quads **(I)** from NTB, KPCY, and KPCY-Pthlh[HET] mice at a set endpoint of 5 months of age. Scale bars, 100 µm. **(J)** Histograms of adipocyte size in

iWAT from NTB, KPCY, and KPCY-Pthlh[HET] mice. **(K)** Immunofluorescence staining of PLIN1 in iWAT from NTB, KPCY, and KPCY-Pthlh[HET] mice. Scale bars, 50 µm. **(L)**

Representative images and H Score quantification of PTHrP IHC in primary tumors from cachectic (N=21) and non-cachectic (N=22) PDAC patients. Scale bars, 100 µm.

Statistical analysis by Mann–Whitney U test (B, C, and D) or Student unpaired t test (D, E, F, G, H, I and L), with significance indicated (*, $P < 0.05$). Error bars indicate standard

deviation (SD).

Figure 2.

(A) Kaplan–Meier overall survival plots of C57BL/6 mice orthotopically implanted with KPCY-EV (N=9) or KPCY-PTHrP[OE] (N=8) cells. Log-rank $P < 0.0001$. **(B)** Representative

dissection images of iWAT from KPCY-EV or KPCY-PTHrP[OE] orthotopic tumor-bearing mice. Scale bars, 1 cm. **(C-D)** Representative H&E images and quantification of

normalized tissue masses for iWAT from KPCY-EV or KPCY-PTHrP[OE] orthotopic tumor-bearing mice. Scale bars, 100 µm. **(E)** Histograms of adipocyte size in iWAT from KPCY-

EV or KPCY-PTHrP[OE] orthotopic tumor-bearing mice. **(F)** Immunofluorescence staining of PLIN1 in iWAT from KPCY-EV or KPCY-PTHrP[OE] orthotopic tumor-bearing mice. Scale

bars, 50 µm. **(G-H)** Representative H&E images and quantification of normalized tissue masses for quads from KPCY-EV or KPCY-PTHrP[OE] orthotopic tumor-bearing mice.

Scale bars, 100 µm. **(I-J)** Schematic depicting rPTHrP treatment once per day over five days and quantification of normalized tissue masses of iWAT from rPTHrP- or sham-

injected C57BL/6 mice. Representative Statistical analysis by Student unpaired t test (D and H) with significance indicated (*, $P < 0.05$). Error bars indicate standard deviation

(SD).

Figure 3.

(A) Human *PTH1R* expression from the GTEX database in the indicated tissues. Adipose

816 and muscle tissues are highlighted. **(B)** qPCR for murine *Pth1r* expression in the indicated tissues; inguinal Brown Adipose Tissue (iBAT), subcutaneous Brown Adipose Tissue

818 (sBAT). Adipose and muscle tissues are highlighted. **(C)** Single-cell RNA-seq clustering plots (left) and *PTH1R* expression (right) from the Emont *et al.*, human adipocyte single

820 cell atlas [25]. **(D)** Immunofluorescence staining of PTH1R in iWAT from non-tumor bearing or KPCY mice. Scale bars, 100 µm. **(E-F)** H&E images and quantification of

822 normalized tissue masses for iWAT from KPCY-EV or KPCY-PTHrP[OE] orthotopic tumor-

bearing mice with the indicated host genotypes. Scale bars, 100 µm. **(G)** Histograms of adipocyte size in iWAT from KPCY-PTHrP^{OE} orthotopic tumor-bearing WT or *Pth1r$^{Adipo-cKO}$* host mice. **(H)** Immunofluorescence staining of PLIN1 in iWAT from KPCY-PTHrP^{OE} orthotopic tumor-bearing WT or *Pth1r$^{Adipo-cKO}$* host mice. Scale bars, 50 µm. **(I-J)** H&E images and quantification of normalized tissue masses for quads from KPCY-EV or KPCY-PTHrP^{OE} orthotopic tumor-bearing mice with the indicated host genotypes. Scale bars, 100 µm. Statistical analysis by Student unpaired *t* test (F and J), with significance indicated (*, *P* < 0.05). Error bars indicate standard deviation (SD).

Figure 4.

(A) Volcano plot of differentially expressed genes from RNA-seq of iWAT from KPCY-EV or KPCY-PTHrP^{OE} orthotopic tumor-bearing mice in C57BL/6 hosts; DNL genes highlighted in blue. **(B)** Significantly altered gene set enrichment pathways enriched in iWAT from KPCY-EV (blue) or KPCY-PTHrP^{OE} (red) tumor-bearing animals. **(C)** GSEA plots of iWAT from KPCY-EV (blue) or KPCY-PTHrP^{OE} (red) tumor-bearing animals. **(D)** Heatmap of altered genes in the DNL, Lipase, Fatty Acid (FA) metabolism, TNF, and IL-6 pathways. **(E)** Schematic representation of the DNL pathway with red arrows indicating all downregulated genes in adipose from KPCY-PTHrP^{OE} tumor-bearing animals relative to KPCY-EV tumor-bearing animals. Created in BioRender. Pitarresi, J. (2025) https://BioRender.com/vapcw88

Figure 5.

(A) qPCR of *Ascc2*, *Acly*, *Acaca*, and *Fasn* in iWAT from KPCY-EV or KPCY-PTHrP^{OE} orthotopic tumor-bearing hosts. **(B)** Western blot analysis of FASN, ACACA, ACLY, and ACTIN in iWAT from KPCY-EV or KPCY-PTHrP^{OE} orthotopic tumor-bearing hosts. **(C)** qPCR of *Chrebp-a*, *Chrebp-b*, and *Srebf1c* in iWAT from KPCY-EV or KPCY-PTHrP^{OE} orthotopic tumor-bearing hosts. **(D-E)** Representative H&E images and quantification of normalized tissue masses for iWAT from KPCY-EV orthotopic tumor-bearing C57BL/6 or *Fasn$^{Adipo-cKO}$* hosts. Scale bars, 100 µm. **(F)** Histograms of adipocyte size in iWAT from KPCY-EV orthotopic tumor-bearing C57BL/6 or *Fasn$^{Adipo-cKO}$* hosts. **(G)** Immunofluorescence staining of PLIN1 in iWAT from KPCY-EV orthotopic tumor-bearing C57BL/6 or *Fasn$^{Adipo-cKO}$* hosts. Scale bars, 50 µm. Statistical analysis by Student unpaired *t* test (A, C, and E), with significance indicated (*, *P* < 0.05). Error bars indicate standard deviation (SD).

Figure 6.

(A-B) Representative H&E images and quantification of normalized tissue masses for iWAT from anti-IgG or anti-PTHrP treated KPCY mice. Scale bars, 100 µm. **(C)** Histograms of adipocyte size in iWAT from anti-IgG or anti-PTHrP treated KPCY mice. **(D-E)** Representative H&E images and quantification of normalized tissue masses for quads from anti-IgG or anti-PTHrP treated KPCY mice. Scale bars, 100 µm. **(F-G)** Representative H&E images and quantification of normalized tissue masses for iWAT from anti-IgG or anti-PTHrP treated C57BL/6 mice orthotopically implanted with KPCY cells. Scale bars, 100 µm. **(H)** Histograms of adipocyte size in iWAT from anti-IgG or anti-PTHrP treated C57BL/6 mice orthotopically implanted with KPCY cells. **(I-J)** Representative H&E images and quantification of normalized tissue masses for quads

from anti-IgG or anti-PTHrP treated C57BL/6 mice orthotopically implanted with KPCY cells. Scale bars, 100 μm. **(K-L)** Immunofluorescence staining of PLIN1 in iWAT from KPCY mice treated with anti-IgG or anti-PthrP (K) or C57BL/6 mice orthotopically implanted with KPCY cells and treated with anti-IgG or anti-PTHrP (L). Scale bars, 50 μm. **(M)** GSEA plots of iWAT from animals orthotopically implanted with KPCY cells and treated with anti-PTHrP (red) or anti-IgG (blue) tumor-bearing animals. **(N)** qPCR of *Fasn* in iWAT from KPCY orthotopic tumor-bearing hosts treated with anti-IgG or anti-PTHrP therapy. Scale bars, 50 μm. Statistical analysis by Student unpaired *t* test (B, E, G, J, and N), with significance indicated (*, *P* < 0.05). Error bars indicate standard deviation (SD).

Figure 7.

Schematic of PTHrP-induced adipose tissue wasting. Tumor-derived PTHrP binds to PTH1R on adipocytes, which downregulates *de novo* lipogenesis in adipose tissue to facilitate wasting. Created in BioRender. Pitarresi, J. (2025) https://BioRender.com/wbw5raq

FIGURE 1

FIGURE 4

FIGURE 6