1
VEGFA critically controls neurovascular invasion and chronic low back pain during
intervertebral disc degeneration

Authors: *Ryan S. Potter[1,2], *Hong J. Moon[2], *Sade W. Clayton[2], Remy E. Walk[3], Addison L.
Liefer[3], Liufang Jing[3], Evan G. Buettmann[2,3], Jennifer A. McKenzie[2], Alec T. Beeve[3], Erica L.
Scheller[3,4,5], Matthew J. Silva[1,2,3], Amber N. Stratman[4], Lori A. Setton[1,2,3], Munish C. Gupta[2], Simon
Y. Tang[1,2,3]
***These authors led the study equally.**

Affiliations:
[1] Department of Mechanical Engineering and Materials Science, Washington University in St. Louis
[2] Department of Orthopaedic Surgery, Washington University School of Medicine
[3] Department of Biomedical Engineering, Washington University in St. Louis
[4] Department of Cell Biology and Physiology, Washington University School of Medicine
35
36

Corresponding author: Simon Tang (simon.tang@wustl.edu)

One Sentence Summary:
VEGFA is a key mediator of neurovascular infiltration in the degenerating intervertebral disc and an
essential driver of chronic low back pain, whose ablation prevents pain-related behaviors.

Keywords:

VEGFA, intervertebral disc injury, low back pain, neurovascular features, preclinical model

Abstract

Despite its enormous burden on patients and society, chronic low back pain (LBP) has no effective therapeutic options. Innervation of the degenerating intervertebral disc (IVD) is suspected to cause discogenic LBP, but the mechanisms that orchestrate the IVD's neo-innervation and subsequent symptoms of LBP remain unknown. We hypothesize that Vascular Endothelial Growth Factor-A (VEGFA) critically mediates the neurite invasion in the IVD and contributes to chronic LBP. Initiating IVD degeneration through a mechanical injury, we evaluated the progression of neurovascular features into the IVD, as well as ensuring LBP symptoms and locomotive impairments at acute (3-weeks) and chronic (12-weeks) timepoints following the IVD injury. To determine the role of VEGFA, we utilized a mouse model with ubiquitously inducible recombination of the floxed *Vegfa* allele (UBC-Cre^{ERT2}; *Vegfa*$^{fl/fl}$). The ablation of VEGFA after an IVD injury attenuated *de novo* neurite and vessel infiltration and impeded the expression of TRPA1, a nociceptive ion channel, in the dorsal root ganglion. The VEGFA-null animals, despite IVD degeneration, exhibited alleviated mechanical allodynia and improved locomotive performance. To determine the effects of IVD-derived VEGFA on endothelial cells and neurons, we co-cultured HMEC-1 endothelial cells and SH-SY5Y neurons with *VEGFA*-silenced human primary IVD cells. The endothelial cells co-cultured with *VEGFA*-silenced IVD cells exhibited reduced vessel growth and shifted their transcriptome and secretome from angiogenic to lymphangiogenic. The neurons co-cultured *VEGFA*-silenced IVD cells showed slowed growth and attenuated transcriptional programs for growth and elongation. These results show that VEGFA directs the growth of intradiscal vessels and neurites that cause low back pain and impaired function, and the inhibition of IVD-derived VEGFA during degeneration may be sufficient to prevent chronic pain behavior and motor impairment associated with discogenic low back pain.

INTRODUCTION

Low back pain (LBP) is a prevalent, debilitating, costly, and multifactorial condition worldwide, imposing substantial global and economic burdens[1-3]. LBP affects as much as 80% of the population, and the elderly experience greater rates of LBP that exacerbate their functional decline, frailty, and loss of independence[4]. LBP is the leading cause of years lived with disability and contributes an estimated $50 to $100 billion USD annually in healthcare costs[4-6]. Chronic LBP is commonly associated with intervertebral disc (IVD) degeneration, which occurs with aging, genetic predisposition, and injury[7-10]. Degeneration leads to structural and biochemical changes in the IVD, including the loss of hydration, inflammation, and matrix breakdown[11-14]. Despite our understanding of the features of IVD degeneration, the specific degenerative features that are mechanistically responsible for the eventual low back pain remain unknown[15]. While the healthy IVD is relatively aneural, the painful human intervertebral discs are innervated, suggesting a role for aberrant nerve growth in discogenic pain[16]. VEGFA (Vascular Endothelial Growth Factor A) is a potent signaling protein that promotes the proliferation, migration, and survival of endothelial and neural cells[17-19]. While VEGFA has known roles in both vascular and neural support, its function following an IVD injury, and how it initiates or sustains discogenic pain remains unresolved. We hypothesized that VEGFA is crucial for the neurite growth and vascularization in the injured IVD to mediate acute and chronic discogenic pain.

To investigate the role of VEGFA in IVD degeneration and low back pain, we utilized a mouse model with an inducible global ablation of VEGFA. To model discogenic low back pain, we created an injury in the lumbar IVD to initiate degeneration[20-22]. Following the IVD injury, we administered tamoxifen to induce the ablation of the *Vegfa* gene. We subsequently measured pain-related symptoms such as

mechanical and thermal sensitivity as well as locomotor function[20]. Because VEGFA is also a critical

regulator of vascular growth, we assessed neo-vascularization into the injured IVD along with neural

infiltration to better understand the relationship between vascular and neural response in disc

degeneration. To evaluate whether there was peripheral nervous system adaptation, we evaluated pain-

related ion channels expression in the lumbar dorsal root ganglia (DRGs). We performed co-cultures

of human cells to determine the role of IVD-produced VEGFA on neuron and endothelial cell growth.

Our findings show that VEGFA plays a central role in promoting neurovascular ingrowth into the

degenerating intervertebral disc. The ablation of VEGFA reduced neurite and blood vessel infiltration

into the degenerated IVD and alleviated the mechanical allodynia and locomotor deficits in mice.

VEGFA ablation also attenuated pain-related ion-channel expression in the DRGs of the injured

animals. Diminished endothelial and neuronal growth *in vitro* with VEGFA-silenced IVD cells

highlights the key role of IVD cell derived VEGFA in promoting vascular and neurite infiltration.

These results demonstrate that targeted inhibition of VEGFA can suppress intradiscal vessels and

neurites that cause peripheral system sensitization, low back pain symptoms, and impaired function.

RESULTS

Removal of VEGFA Blunts Intradiscal Invasion of Neurovascular Structures

In WT (Cre−;*Vegfa*^{fl/fl} with tamoxifene administration) animals, the injured IVD was infiltrated by

PGP9.5+ neurons and by Endomucin+ vessels. VEGFA ablation significantly blunted these neuronal

and vascular features. Immunofluorescence images from 50µm thick sections (Figure 1A-B) showed

fewer neural and vascular structures in VEGFA-null mice compared to WT particularly in the region

around the outer annulus fibrosus (AF) near the injury site. In a validation study, at least 94% of the *de*

novo PGP9.5 positive structures within the IVD were positive for $Na_v1.8$ (Figure S3), a marker for

sensory neurons[60]. Quantitative measurements revealed that VEGFA removal diminished neurite

length (p=0.01; Mean difference = -57.34, 95%CI:-103−-12.1[µm]), vessel length (p=0.004; Mean

difference = -177, 95%CI:-295−-59.4[µm]), neurite penetration depth (p=0.002; Mean difference = -

15.25, 95%CI: -24.9−-5.72[µm]), and vessel penetration depth (p=0.047; Mean difference = -18.80,

95%CI:-37.3−-0.284[µm]) (Figure 1C-F). Post hoc analysis reveals differences between the WT and

VEGFA-null IVDs in neurite depth at 3 weeks (p=0.001) and in vessel length at 12 weeks (p=0.01).

These findings show that VEGFA is crucial for instigating the neurite growth and sustaining vessel

propagation in IVDs following injury. No sex-differences were observed (Table S1).

Fig. 1: Removal of VEGFA Blunts the Invasion of Neurovascular Structures on the

Intervertebral disc dorsal surface Following lumbar IVD Injury. A-B) IHC with the nuclei

(DAPI) in blue, neuronal features (PGP9.5) in green and vascular features (Endomucin) in red. C-F)

VEGFA-null animals had diminished neurite lengths (p=0.01; Mean difference = -57.34, 95%CI:-

103−-12.1[μm]) and vessel lengths (p=0.004; Mean difference = -177, 95%CI:-295−-59.4[μm]).

Likewise, the neurite penetration depth (p=0.002; Mean difference = -15.25, 95%CI: -24.9 − -

5.72[μm]) and vessel penetration depth depth (p=0.047; Mean difference = -18.80, 95%CI:-37.3−-

0.284[μm]) are blunted by VEGFA removal. Post hoc analysis reveals differences between the WT

and VEGFA-null IVDs in neurite depth at 3 weeks (p=0.001) and in vessel length at 12 weeks

(p=0.01). Mean ± standard deviation are shown in the graphs.

VEGFA Is Essential for the Spatial Colocalization of Intradiscal Nerves and Vessels

The spatial proximity of intradiscal nerves and vessels are disrupted by VEGFA ablation.

Immunofluorescence images reveal the distribution of neural (PGP9.5, green) and vascular

(Endomucin, red) structures highlighting the regions of interest around the outer annulus fibrosus (AF)

and overlapping lengths relative to neurites (Figure 2A-B). The colocalization analysis shows that the WT injured IVDs have neurites with a large amount of vessels in their proximity. The removal of VEGFA in the IVDs 3 weeks following injury protected more IVDs from neurites penetration (Figure 2C;p =0.02; OR=4.275, 95%CI:1.44−12.6). The loss of VEGFA also protected more IVDs from the penetration of endomucin+ vessels (Figure 2D; p=0.03; OR=10.67, 95%CI:1.69−122). The spatial overlap analysis between neurites and vessels shows that VEGFA ablation dramatically reduces the amount of neurites near vessels (Figure 2E; p=0.007; Mean difference = -48.8, 95%CI: -84.0—-13.69[μm]). After adjusting for amount of neurites in each animal, this observation remains robustly independent of the basal level of neurites (Figure 2F p=0.008; Mean difference=-36.1, 95%CI: -0.943—-71.2[μm]). The lack of difference over time in the depth and the length ratios between the neurites and vessels indicates that once initiated, VEGFA does not appear to affect the propagation of these features (Figure 2G-H). These findings highlight the critical role of VEGFA in the *de novo* initiation of neural and vascular features in the IVD.

Fig. 2: VEGFA is Essential for the Colocalization of Intradiscal Nerves and Vessels and for *de novo* neurovascular invasion. A) An injured WT IVD and the ROI for the analysis neurovascular

features. B) Graphical representation of how spatial colocalization analyses quantified the number of neurites that are with a 30-µm proximity of vessels. C-D) Neurite (p =0.02; OR=4.275, 95%CI:1.44−12.6) and vessel (p =0.03; OR=10.67, 95%CI:1.69−122) infiltration proportions were both significantly higher in WT animals than in the VEGFA-null groups. E) VEGFA ablation suppressed vessel-adjacent neurites (p=0.007; Mean difference = -48.8, 95%CI: -84.0—-13.69[µm]). F) The loss of VEGFA prevented neurite-vessel spatial coupling even when accounting for varying intradiscal neurite lengths (p=0.008; Mean difference=-36.1,95%CI: -0.943—-71.2[µm]), confirming that this phenomenon is conserved even when accounting for heterogenous vessel sprouting across animals. G-H) The lack of an effect of time in the depth of penetration and length elongation ratios suggest that once the neurites and vessels have initiated, the loss of VEGFA does not appear to affect their propagation. Statistical model C-D: Fisher's exact test. Statistical model E-H: 2 factor ANOVA with Fisher LSD post-hoc comparisons. E-H Data are presented as mean ± standard deviation.

Depletion of VEGFA Suppresses the Expression of Nociceptive Ion Channels in the DRG

Quantitative measurements of corrected total cell fluorescence (CTCF), averaged from the left and the right sides, of the L2-L3 lumbar dorsal root ganglia (DRGs; Figure 3B-D) revealed a significant reduction in transient receptor potential ankyrin 1 (TRPA1) fluorescence (Figure 3D; p=0.037;Mean difference=-16040, 95%CI: -30800—-1280) due to VEGFA removal following injury. No significant changes were measured in tyrosine kinase receptor A (TrKA) or transient receptor potential vanilloid 1 (TRPV1) CTCF levels due to the loss of VEGFA ablation.

174
Fig. 3: VEGFA Inhibition Reduces the Expression of Pain-related Ion Channels in the DRG. A)

TrKA (top row), TRPV1 (middle row), and TRPA1 (bottom row) in green and DAPI in blue were

quantified. B-C) No effects due to VEGFA were observed in TrKA or TRVP1 expression measured by

Corrected total cell fluorescence (CTCF). D) VEGFA ablation suppressed TRPA1 expression

(p=0.037;Mean difference=-16040,95%CI:-30800 — -1280). TRPA1 expression is canonically

associated with nociceptive sensitization.

Ablation of VEGFA Alleviates Mechanical Sensitivity and Locomotive Function

Loss of VEGFA Attenuates Mechanical, but Not Hot or Cold, Sensitivity

WT animals exhibited mechanical sensitivity measured by the Von Frey filament test at both 3 and 12

weeks after the IVD injury compared to the pre-surgery baseline. Ablation of VEGFA prevented this

impairment (Figure 4A; p=0.01, Mean difference=0.490, 95%CI: 0.110—0.860). The loss of VEGFA

reversed the mechanical allodynia at 12 weeks. The electronic Von Frey filament test also revealed

that there is a greater mechanical sensitivity on the side the surgical approach (e.g. left). VEGFA

removal specifically alleviated the mechanical sensitivity of the injured side, without modifying the

contralateral side (Figure S1G). No notable changes were observed in heat or cold sensitivity at any

post-operative time point (Figure 4B-C; Table S1).

VEGFA Ablation Improves Rotarod Performance, but Does Not Affect Passive Function

We longitudinally evaluated locomotive performance through an active challenge of physical function,

measured through the rotarod assay, every three weeks up to 12 weeks after injury. The longitudinal

rotarod test reveals that VEGFA-null mice exhibited longer latencies to fall compared to WT (Figure

4D; p=0.004, Mean difference=15.13, 95%CI:5.35—24.9), in contrast to the progressive decline of

motor coordination over time in the WT animals (p=0.003). Remarkably, at 9- and 12- weeks, the

initial decline of rotarod performance in VEGFA null animals was recovered. In the Open Field test,

total distance traveled (Figure 4E) and rearing counts (Figure 4F) were not different between

genotypes, sex, tamoxifen, or due to surgery (Table S1), suggesting comparable levels of exploratory

behavior and general activity between all groups.

Fig. 4: VEGFA Ablation Improves Mechanical Sensitivity and Performance A) The loss of
VEGFA mitigated the mechanical sensitivity of the animals with injured IVDs (p=0.01;Mean
difference=0.490, 95%CI: 0.110—0.860). In particular, the VEGFA-deficient animals recovered from

the initial mechanical sensitivity at 9- and 12- weeks while the WT animals do not recover (p<0.01).

B-C) No differences in hot or cold sensitivity were observed. D) There were significant differences in

longitudinal measures of performance that included rotarod (genotype, p=0.004; time, p=0.003),

indicative of a time-dependent VEGFA effect on the progression of performance. E-F) There was no

differences in any measures of open field with genotype or time point. G-I) There was no differences

in the inverted screen endurance or hind paw grip strength with genotype and at any time point.

Statistical model A-F, I: Linear mixed effects with genotype and time as factors with post-hoc LSM

with Sidak correction. G-H Mantel-Cox test. A-F, I Data are presented as mean ± standard deviation.

VEGFA Removal Does Not Affect Measures of Strength

To determine the impacts on musculoskeletal function following VEGFA ablation, strength

assessments were conducted, including inverted screen endurance and hind paw grip strength. The

inverted screen endurance test, presented using Kaplan-Meier curves, showed that VEGFA-null mice

have similar probabilities of survival (e.g. hanging on the wired mesh) compared to WT at both 3

weeks (Figure 4G) and 12 weeks (Figure 4H). Measures of hind paw grip strength (Figure 4I) were

also found to be similar across genotypes, indicating no compromise in muscle function from VEGFA

ablation.

Overall, the behavioral and locomotive function analyses show that VEGFA-null mice have

improved recovery of mechanical sensitivity. Further, the VEGFA-null animals averted the transition

to chronic sensitivity, while also avoiding the performance deficits observed in WT at the chronic time

points.

Ablation of VEGFA does not affect structure, function, or degeneration of the IVD

Loss of VEGFA Does Not Affect the Histopathologic Degeneration of the IVD

CEμCT analysis of the IVDs 3 weeks after injury reveals significant deterioration of the IVD structure. By using a contrast enhancing agent, Ioversol, the injury was observed on the left ventral side of the outer AF as progressively higher attenuating regions that mark the fibrous scar tissues (Figure 5A). The injured IVDs also exhibit a loss of NP attenuation, measured by NI/DI, demonstrating the loss of the water retaining ability of the NP (Figure 5B; $p=0.03$, Mean difference$=-0.0102$, 95%CI:-0.0190—-0.00138). The ablation of VEGFA does not affect injury-mediated degeneration. A narrowing of the intradiscal space measured by disc height ratio (DHR), a hallmark of IVD degeneration, is observed in all of the injured discs (Figure 5C; $p=0.01$, Mean difference$=-0.0246$, 95%CI:-0.0431—-0.00605) with no differences among genotypes or with post-operative time. There was not a change in NP volume either due to injury or VEGFA ablation (Figure 5D). Injury increases the degenerative features including decreased NP hydration and a narrowing DHR; VEGFA ablation did not appear to alleviate these tissue-level changes in the IVD.

Injury to the IVD Does Not Impair Its Mechanical Function

Dynamic compression of the IVDs 3 weeks post injury shows no distinct mechanical responses to injury or impact from VEGFA ablation. The stiffness measurements show no significant differences between WT and VEGFA-null groups, both in sham and injured conditions (Figure 5E). Likewise, the dissipation factor indicates no appreciable changes from injury or VEGFA loss (Figure 5F). These findings show that the degeneration that occurs 3 weeks after injury in this model does not yet impair the IVD's mechanical function.

VEGFA Loss Does Not Rescue the Degenerative Changes Caused by Injury

Histopathological scoring (Figure 5G-H) revealed the damaging effects of the needle stab injury on the compartments in the disc, including decreased NP cell density and changes in cellular morphology, disruption of the AF concentric lamellar structures with increased cellularity and presence of clefts, along with thinning and disruptions in the endplate cartilage, and structural disorganization in the boundary regions. Scores for injured groups averaged near 26 out of a possible 36 for maximal degeneration, while sham controls averaged 11 (Figure 5G,H; p<0.0001; Mean difference=15.2, 95%CI:13.5—16.5). Effects of VEGFA removal were not detected. IVD injury dramatically changed all compartments of the disc with the largest effects on the AF followed by NP while sham scores indicate mild degeneration from the surgical exposure of the disc and likely disruption of surrounding soft tissues.

Fig. 5: Loss of VEGFA Does Not Alter Structural, Mechanical, or Histopathologic Degeneration of the Injured IVD. A) IVD structure analyses via CEμCT show that B) injury decreased NP hydration (p=0.03, Mean difference=-0.0102, 95%CI:-0.0190—-0.00138), indicated by the decreased attenuation of the NP compared to sham. C) Injury decreased the IVD's disc height ratio, DHR, (p=0.01, Mean difference=-0.0246, 95%CI:-0.0431—-0.00605) compared to sham. D) No effect of injury was observed on NP volume. Changes to mechanical behavior were not observed in E) stiffness and F) dissipation factor, indicating that tissue level mechanics were not changed by injury or VEGFA removal at 3-weeks post injury. G) Histological images stained with Safranin-O and Fast Green show IVD degeneration. H) Histopathologic scoring shows a strong effect of injury (p<0.0001;Mean difference=15.2, 95%CI:13.5—16.5) with no statistically distinguishable differences due to VEGFA loss in nucleus pulposus, annulus fibrosus, end plate, or the interface/boundary demonstrating the pathophysiological response from needle puncture. Data are presented as mean ± standard deviation.

Silencing *VEGFA* RNA in Human IVD Cells Attenuates the Growth of Co-Cultured Neuronal Cells (NC) and Endothelial Cells (ECs)

Transfecting human IVD cells with VEGFA-specific small interfering RNA (siRNA) suppressed their IL-1β-stimulated VEGFA production by 57.4% (Figure S6). After removing the IVD cells from the IL-1β laden media, co-culturing these VEGFA-siRNA transfected IVD cells with SH-SY5Y neuroblastoma-derived neuronal cells (aka NCs) maintained the NCs' morphology but blunted the axonal growth (Figure 6A). The total neurite length reveals declined neurite growth in NCs co-cultured VEGFA-siRNA transfected IVD cells (Figure 6B-D). RNA sequencing revealed that NCs co-cultured with VEGFA-knockdown IVD cells differentially downregulated 39 genes and upregulated 107 genes (aka DEGs) compared to those co-cultured with VEGFA-intact IVD cells (Figure 6E). Gene

ontology analysis determined the most enriched biological process from the DEGs were involved

GPCR signaling pathways, known for neurovascular development (*EDNRA, CHGA, RAMP3*),

regulation of growth (*DCC, GREM1, STC2, MYOZ1, MT1F, BST2, IGFBP3, CXCR4, MT1X, IGFBP7,*

BCL2), and the negative regulation of cell population proliferation (*GREM1, PTPN14, DHRS2, SSTR2,*

SCG2, SERPINF1, IGFBP3, OVOL1, ADM, IGFBP7, BCL2, S100A11). Notably, several DEGs were

specific to cell growth inhibition (*SSTR2, S100A11),* immune cells and inflammation (*CXCR4, FGFB2,*

IL32), and neuronal cell functions (*DCC, SDK2*, *ASTN2*).

We also co-cultured VEGFA-siRNA transfected IVD cells, stimulated a priori with IL-1β, with the

HMEC-1 endothelial cells (aka ECs). The ECs co-cultured with VEGFA-silenced human IVD cells

formed fewer closed tubes, a measure of vessel formation, after 6 hours of culturing (Figure 7A-B;

p=0.04). RNAseq uncovered 175 downregulated and 0 upregulated DEGs when ECs are co-cultured

with IVD cells transfected with VEGFA siRNA compared with scram (Figure 7C). Gene ontology

analysis determined the most enriched biological processes were involved in vessel and vasculature

related processes. These included lymphatic vessel development (*NR2F2, TBX1, SOX18, EFNB2,*

HEG1), blood vessel morphogenesis (*TNFAIP2, EFNB2, HEG1, PRKACA, NRARP, TBX1, SOX4,*

NR2F2, AKT1, QKI, SMAD7, SOX18, VEGFB), and regulation of neuron projection development

(*EFNB2, CDK5R1, RYK, RAPGEF1, CARM1, AKT1, PLZNA1, NCS1, DVL1, SLK, STK24, BRSK2,*

ADNP). Interestingly, the canonical WNT signaling pathway was also an enriched biological pathway.

WNT signaling plays an important role in endothelial cell proliferation, survival, differentiation and

response to mechanical forces due to shear stress from blood flow.[23,24]

Fig. 6: Co-culturing SH-SY5Y neuronal cells with VEGFA-silenced human primary IVD cells suppressed *in vitro* neurite growth and transcriptional processes related to growth and elongation. A-D) The neurite extension assay after co-coculturing neuronal cells with IL-1β stimulated VEGFA siRNA transfected human IVD cells shows reduced total neurite lengths. E) Volcano plot of RNA-seq analysis showing 146 differentially expressed genes and F) key biological processes regulated by VEGFA siRNA. NCs co-cultured with IL1β naïve disc cells showed undifferentiated cells possessing few short projections which cluster together, while NCs co-cultured with IL1β+SCRAM siRNA and NCs co-cultured with IL1β+VEGFA siRNA showed differentiated cells having many extensive projections.

314
Fig. 7: Co-culturing with VEGFA-silenced human primary IVD cells suppressed *in vitro*

endothelial cell growth and shifts the endothelial cell transcriptional program from angiogenic

to lymphangiogenic. A) Matrigel tube forming assay images of endothelial cells (ECs) cocultured

with human IVD cells treated with siRNA showing initiation of vessel formation after 6 hours. B) The

silencing of VEGFA siRNA in IVD cells reduced tube formation in EC co-cultures, compared to IVD

cells with Scrambled siRNA (p=0.04). C) The ECs co-cultured with VEGFA-silenced IVD cells

downregulated 176 genes. Key angiogenic genes were suppressed including *HEG1*, *NARAP*, and

NCS1. There is also a suppression of endothelial cell-derived neurogenic genes including *BRSK2* and

CRLF1. D) The RNAseq analyses of the HMEC-1 also show an ontological shift to

lymphangiogenesis, matrix remodeling, and suppression of neurovascular regulating pathways

following co-culture with VEGFA-silenced IVD cells.

The endothelial cells (ECs) co-cultured with the VEGFA-silenced IL-1β-stimulated IVD cells

produce less angiogenic and inflammatory factors and more non-VEGFA proangiogenic

compensatory factors

Following co-culture with VEGFA-silenced, IL-1β-stimulated IVD cells, the ECs exhibited a distinct

cytokine profile that indicate a shift in cellular function. Notably, there was a reduction in secretion of

VEGFA (Figure 8A) and CCL3 (Figure 8K) protein in the ECs co-cultured with VEGFA-silenced

IVD cells, indicating decreased angiogenic and inflammatory signaling. The ECs exposed to VEGFA-

silenced IVD cells also produced more VEGFC protein, a canonical driver of lymphangiogenesis and

lymphatic endothelial cell proliferation[25,26]. Simultaneously, ECs increased protein production of

PDGF-AA (Figure 8J), a mitogenic factor that can compensate for VEGFA loss by promoting vessel

stabilization and pericyte recruitment[27]. Increases in MMP1, MMP3, MMP10, and MMP13 protein

levels (Figure 8C-F) reflect heightened extracellular matrix remodeling, a process essential for vessel

pruning and structural reorganization typically associated with either vessel regression or tissue

remodeling rather than active angiogenesis[28,29]. The elevation of granulocyte colony-stimulating factor

(G-CSF) and granulocyte-macrophage colony-stimulating factor (GM-CSF) supports a role for

immune cell recruitment and activation, especially of myeloid lineages that modulate inflammation

and tissue repair (Figure 8G-H)[30,31]. Similarly, upregulation of CXCL1 and CCL5 (RANTES)

reinforces the potential for leukocyte chemotaxis and immune surveillance (Figure 8I,L)[32,33].

Fig. 8: The secreted factors from HMEC-1 (aka ECs) co-cultured with IVD cells showed

compensatory factors for angiogenic homeostasis and shifts to lymphanogenesis. Despite having a

functional *VEGFA* gene, the ECs produced less (A) VEGFA and (K) CCL3 protein when co-cultured

with *VEGFA*-silenced primary human IVD cells. The ECs also produced more (B) VEGFC and (J)

PDGF-AA that are both positive regulators of lymphangiogenesis; MMPs (C-F; MMP1, MMP3,

MMP10, MMP13) involved in matrix remodeling; chemokines including (G) G-CSF, (H) GM-CSF,

(I) CXCL1, and (L) CCL5; and (J) PDGF-AA, a compensatory angiogenic growth factor. Units are

pg/ml.

Taken together, this cross talk supports a mechanism in which endothelial cells, in the presence of IL-

1β-stimulated secretome of VEGFA-silenced IVD cells, pivot toward a reparative state characterized

by elevated lymphangiogenesis, increased immune cell engagement, and enhanced matrix

remodeling—rather than the robust angiogenic response seen with IL-1β-stimulated VEGFA-intact

IVD cells. These findings align with the vessel-pruning and tissue remodeling hypotheses seen in

regenerative contexts and suggest that VEGFA inhibition not only reduces angiogenesis but promotes

a more immunomodulatory and matrix-adaptive endothelial phenotype. In addition, several mediators

known to support vessel homeostasis (e.g., Angiopoietin-2, Endoglin, Endothelin-1, PlGF) remained

unchanged in co-culture with the VEGFA-silenced IVD cells (Table S2) and likely continue to support

the homeostatic functions of existing vessels.

DISCUSSION

This is the first study to demonstrate that intradiscal innervation contributes to low back pain behavior and locomotive performance in a model of intervertebral disc degeneration. While studies in the past have shown progressive associations between neurovascular structures with degeneration and pain[34-38], none have been able to block or prevent these structures to prevent chronic pain and maintain locomotive performance. From this work, we also demonstrate that VEGFA may be a therapeutic target for a disease modifying therapy of low back pain. VEGFA ablation had a profound effect on neural and vascular infiltration and the subsequent mechanical sensitivity and locomotive performance following degeneration, independently of the structural and mechanical properties of the IVDs.

VEGFA plays a crucial role in limiting nerve and vessel infiltration post-injury. IVD degeneration in VEGFA-null mice exhibited reduced neurite and vessel lengths and the depths of their penetration, particularly in the outer annulus fibrosus (AF). VEGFA ablation inhibited the penetration and elongation of neural and vascular structures in the normally avascular IVD, preventing nociceptive sensory neurons from entering injured areas. The synergistic relationship between nerve and vessel infiltration highlights VEGFA's importance in neurovascular coupling within the IVD, with its absence leading to diminished neurite and vessel growth, which may limit pathological nociceptive signals that propagate into chronic pain.

The loss of VEGFA also affected neuropeptide-related expression in the Dorsal Root Ganglion (DRG). The increase expression of TrKA, TRPV1, and TRPA1 in the DRG have been observed in chronic low back pain[39,40], as well as in DRG that are sensitive to capsaicin stimulation[41]. VEGFA removal also decreased TRPA1 expression, whose expression is associated with nociception, in the DRGs. In contrast, TRPV1 expression, commonly associated with heat sensitivity[42], did not change between VEGFA-null mice and wild-type mice. Indeed, the VEGFA-null mice and wild-type mice did

not differ in heat sensitivity to the hot plate. VEGFA-null mice recovered from the early mechanical

sensitivity at three-weeks and avoided the chronic sensitivity observed in WT mice. VEGFA-null mice

did not develop the motor performance impairments exhibited by WT mice at later points (9-12

weeks), indicating a protective effect of VEGFA ablation against chronic pain and impaired

performance. The asymmetric, injury-induced hypersensitivity was avoided by VEGFA inhibition,

indicating that VEGFA loss mitigated the effects of IVD injury without altering overall pain

sensitivity. Together, these findings suggest that VEGFA modulation is disease modifying rather than

producing analgesic effects.

Neurovascular Elongation vs Sprouting

Injuries to the intervertebral disc primarily trigger nerve and blood vessel sprouting as an adaptive

response to inflammation and hypoxia[43,44]. This sprouting—stimulated by factors like NGF and

inflammatory cytokines—leads to increased neurovascular infiltration into normally aneural and

avascular disc regions, contributing to chronic pain and ongoing inflammation[45-47]. Consistent with

this, we observed neurites and vessels infiltrating into the degenerating IVD at 3-weeks following

injury. However, once sprouted, the neurites did not appear to elongate further, as evidenced by the

relatively similar lengths and depths between 3- and 12- weeks. In this animal model, VEGFA is

ubiquitously ablated from the whole animal, and it is not possible to determine the source of the

VEGFA. However, our *in vitro* experiments support that blocking IVD-produced VEGFA is sufficient

to prevent the growth of neurites and the formation of blood vessels. RNAseq analysis suggested that

the reduced neural outgrowth is driven by the suppression of cell proliferation. We observed a

phenotypic shift in the endothelial cells co-cultured with VEGFA-silenced IL-1β−stimulated IVD cells

that secreted higher levels of VEGFC and PDGF-AA, positive regulators of lymphangiogenesis,

alongside increased MMPs, and immune cell recruiting chemokines. Such changes suggest that, in the

absence of VEGFA-driven angiogenesis, endothelial cells promote a microenvironment conducive to

tissue remodeling and immune-mediated repair. These immune cell recruiting chemokines are likely to

be important for the intrinsic healing of the IVD[48]. RNAseq analyses of the HMEC-1 cells confirms

the ontological shift to lymphangiogenesis and matrix remodeling with a suppression of key

angiogenic genes including *Heg1, Narap*, and *Ncs1*. There is also a suppression of endothelial cell-

derived neurogenic factors including *Brsk2* and *Crlf1*.

Needle stab injury models in animals are frequently utilized to study the pathophysiology of

intervertebral disc (IVD) degeneration and to develop treatments for chronic lower back pain (CLBP).

These models effectively replicate critical aspects of human disc degeneration, such as mechanical

damage, inflammatory responses, and biochemical changes. For example, they simulate the

mechanical damage caused by traumatic events and the inflammatory cascade observed in human

discs, which are crucial for understanding the progression of disc degeneration[49]. However, these

models have limitations, especially in replicating the chronic, multifactorial nature of human CLBP

and the complexity of human pain perception[50,51]. Human IVD degeneration occurs over years, with

the herniation of the IVD being one of the most apparent instigating factors. Despite the varied

etiologies of discogenic pain, the chronic symptoms and eventual degenerative presentation, such as

the deterioration of IVD structure, are quite similar. Therefore, the needle injury model in murine IVD

is employed to understand the mechanisms following IVD injury that eventually lead to low back pain

symptoms. The needle injury in a mouse induces progressive, but relatively rapid degeneration,

occurring over weeks and months, compared to human degeneration, which spans years and decades.

Despite being traumatic relative to the human scale, this model is a necessary caveat for developing

feasible and mechanistic experiments[51]. A single level lumbar injury utilized here may also

underestimate the magnitude of our findings, since three-level injuries appear to show more

pronounced effects[52]. Even with a single-level injury, however, the shift to painful behavior were

obvious with longitudinal measurements. Notably, neurovascular ingrowth was more pronounced on

the dorsal side opposite the ventral injury location,[34], likely due to its proximity to the spinal cord and

increased availability of neural and vascular sources. These findings have translational promise since it

is possible to intervene early upon an acute IVD injury (e.g. a herniation) in the sequalae of chronic

low back pain development. Since patients typically seek treatment only after an injury or an acute

episode of pain, the timely post-injury inhibition of VEGFA may be able to prevent the transition to

chronic low back pain.

Overall, the loss of VEGFA protected against mechanical sensitivity and locomotive

performance and decreased intradiscal innervation. These findings show a causative relationship

between neurovascular pathoanatomy and pain, while demonstrating VEGFA as a critical mediator of

chronic pain development through the modulation of neurovascular features.

MATERIALS AND METHODS

Experimental Design

The Postnatal Inducible Ablation of Vegfa in Adult Mice

Young adult, 4–6-month-old, ubiquitin (UBC)-Cre^{ERT2} ; *Vegfa*$^{fl/fl}$; Ai9-tdTomato[53] mice on a C57BL/6J background[54-56] allowed for a temporally controllable ablation of *Vegfa* in all tissues upon tamoxifen administration, along with an Ai9 fluorescence reporter to confirm recombination. The *Cre* element is activated when a mutant estrogen receptor (ERT2) renders *Cre* active upon tamoxifen binding and promotes Cre entering the nucleus to exise the *Vegf*a-encoding regions flanked by the *LoxP* sites. This approach avoids VEGFA-deficiency-mediated developmental defects and embryonic lethality, especially in pathways that are essential to organogenesis. All procedures (breeding, drug administration, procedures, and tissue extraction) were performed with WUSM IACUC (Institutional Animal Care and Use Committee) approval.

The Lumbar Puncture Intervertebral Disc Injury Model

Male and female mice of 4-6 months of age were randomly allocated into respective experimental groups. The animals undergoing surgery were anesthetized, shaven, and sterilized prior to surgery. Under microscopy guidance, the lateral lumbar spine was surgically exposed by mobilizing the peritoneum and retracting the psoas anteriorly. Using the superior pelvic margin to identify the L6 vertebrae, and then the L5/6 and L6/S1 IVDs are surgically exposed. The L5/6 IVD was injured using a 30G beveled hypodermic needle with three unilateral annular injuries. The sham animals received the same surgical exposure with no injury[20,22,34].

On Days 3–4 following the surgery, Cre+;*Vegfa*$^{fl/fl}$ and Cre−;*Vegfa*$^{fl/fl}$ littermates received tamoxifen by gavage to induce *Vegfa* ablation in the Cre+ animals. (aka 'VEGFA-null') with

Cre−;Vegfa[fl/fl] littermates as WT animals. Data were collected at baseline, within 7 days, and at 3, 6, 9,

and 12 weeks (behavioral assays); body weight at baseline/3/12 weeks; survival tracked for Kaplan–

Meier analysis. Acute 3-week post-op and chronic 12-week post-op time points were chosen with the

acute injury response allowing for the initial surgical wound healing recovery, while chronic LBP is

defined as pain lasting 12 weeks or longer[57,58]. Additional experimental details are provided in the

Supplemental Materials.

Histological analyses

Sample Preparation

Spines from the euthanized animals were fixed in 4% paraformaldehyde, demineralized in 14%

EDTA, and serially infiltrated with sucrose solutions prior to optimal cutting temperature (OCT)

infiltration and freezing for cryosectioning.

480
Histopathological Evaluation of Degeneration

Histopathological scoring for degeneration in four compartments of the IVD including the NP, AF,

end plate, and interface/boundary regions on 10µm-thick sections[59]. Mean degeneration scores from

three independent investigators (n=3/sex/genotype/surgery, N=24 total) were used for analysis.

Structural Quantification of Nerve and Vascular Features by Immunohistochemistry

Two to three 50µm-thick serial sections were used to evaluate the neuronal and vascular morphology

in 3-dimensions. Protein Gene Product 9.5 (PGP9.5), Endomucin, and DAPI counter stain[60] were co-

stained with respective antibodies to evaluate the nerve and vascular structures. DAPI, PGP9.5, and

Endomucin helped visualize, respectively, cell nuclei, broad neural populations, and endothelial cells

of vasculature via a confocal microscope at 2.5 µm focal depth increments with 10x magnification.

Excitation and emission wavelengths were set as follows: DAPI excitation 405nm and emission 414-

450 nm, PGP9.5 excitation 488nm and emission 498-530 nm, and Endomucin excitation 635nm and

emission 650-725 nm. Neurite and vessel quantification of the outer AF were performed on max

projections of a Z-stack with Fiji (ImageJ), with lengths on or within the AF and the surrounding

fibrous tissue calculated from the Neuroanatomy SNT plugin and with depths calculated normal to the

peripheral edge of the AF. PGP9.5 staining showed some non-specific regions so was only not

included if neuronal morphology was apparent. To quantify spatial coupling of neurites and vessels, an

automated MATLAB script was developed to measure lengths and depths and calculate colocalization

lengths and a colocalization ratio based on distance from the rare neurite features, over a 30 µm

distance. We also confirmed that 94% of *de novo* PGP9.5+ neurites that infiltrate the IVD are positive

for $Na_v1.8$, a sensory neuron marker (Figure S3).

Neuropeptide and Ion Channel Immunostaining of the Dorsal Root Ganglia

Frozen sections of isolated Doral Roo Ganglia (DRG) from L1-L3 levels were processed for

immunohistochemistry for anti-TrkA, anti-TRPA1, and anti-TRPV1. Nuclear counterstaining was

performed using DAPI, and slides were mounted prior to imaging. DRG immunohistochemical images

were quantified using the custom Python software, DRGVisual. For each cell, the corrected total cell

fluorescence (CTCF) was calculated by subtracting the background intensity from each pixel's

intensity within the cell region and summing the total fluorescence across the cells for each ion

channel.

3D Structural Evaluation of Intervertebral Disc Structure Using Contrast-enhanced µCT

At 3 weeks post lumbar intervertebral disc injury, *in vivo* µCT was used to evaluate structural changes

of the intradiscal space using disc height[20]. The structural changes at this 3 week post injury time point

were further evaluated using the e*x vivo* contrast-enhanced micro-computed tomography (CEµCT).

Ioversol was used as a contrast agent to highlight the water-rich nucleus pulposus[61]. Total disc ROI

and nucleus pulposus ROI were contoured using the MATLAB graphical user interface from

Washington University Musculoskeletal Image Analyses (github.com). NP hydration, a ratio of

nucleus attenuation to total disc attenuation (NI/DI), was calculated along with volumes and disc

height ratio, DHR, as an average of 5 height measures in a midsagittal slice over the disc width[62].

Biomechanical Assessment of the Intervertebral Disc

Mechanical behavior of the IVDs using dynamic compression testing under displacement control[63].

FSUs were affixed to aluminum platens and immersed in a phosphate-buffered saline bath at room

temperature and preloaded to 0.25N. A sinusoidal compressive waveform at 10% strain and 1 Hz

frequency was applied for 20 cycles. The average stiffness was calculated from the second through the

final loading cycles and dissipation factor (aka the loss tangent), was calculated from the phase shift

between load and displacement data.

Behavioral Assessments

Habituation measures were taken to reduce animal stress during behavioral assessments[64]. All tests

were performed by the same laboratory personnel who was blinded to genotype and surgical procedure.

Baseline data was collected one week prior to surgery and post-operative data collection included a

cross-sectional group at baseline, 3, and 12 weeks and a longitudinal group at 3, 6, 9, and 12 weeks.

Mechanical Sensitivity Measured Using the Von Frey and E-Von Frey

Mechanical sensitivity was measured using the up-down method of the Von Frey test using Semmes

Weinstein filaments and an electronic Von Frey (EVF; BIOSEB, France)[65,66]. Prior to baseline

behavioral testing, mice were habituated to the elevated mesh grid chamber for 1 hr. Calibrated

filaments were applied to the plantar surface of the hind paw to elicit paw withdrawal or filament

buckling[67]. Filament Von Frey was done on the left hind paw to focus on the injury response. EVF

consisted of maximum force values at withdrawal from the median of the result of 3 trials which were

conducted bilaterally, with at least 5 min between testing opposite paws, and at least 10 min between

testing the same paw. Post-op measures were normalized to the pre-surgery baseline.

Hot and Cold Nociceptive Sensitivity

Mice were subjected to thermal stimulus to measure the temperature sensitivity response[68,69]. A

hot/cold plate (Bioseb, France) was set to 55±0.5°C or 5±0.5°C and mice were placed individually

onto the plate with a transparent enclosure to prevent escape and allow for visibility. The latency to the

first sign of nociceptive behavior, either lick of the hind paws or jumping for hot plate; and lifting,

licking, or shaking of the hind paws for cold plate was recorded with a cutoff time of 20 seconds for

hot or 30 seconds for cold to prevent tissue damage. Three repeated measures were collected with >15

minutes between tests to allow for home cage recovery.

Functional Assessments of Performance

<u>Locomotor Function</u>

Rotarod

Assessing motor coordination and balance, the Rotarod performance test challenged mice to a rotating

rod apparatus (Bioseb, France)[66,70,71]. Prior to testing, mice were trained on the Rotarod at a slow 4

rpm for 2 minutes to ensure baseline capabilities. On a subsequent day, mice were tested on an

accelerating bar from 4 to 40 rpm over a 2-minute period. Latency to fall was recorded for each mouse.

Each mouse underwent 5 trials with a >5-minute rest period between trials to reduce fatigue. The best

performance of the 5 trials was used for data analysis and normalized to baseline.

Open Field Testing

In the open field test, designed to evaluate exploratory passive behavior, mice were individually

introduced into the center of an open field arena (40 cm x 40 cm, with 30 cm high walls) equipped

with an automated tracking system (Omnitech Electronics, Columbus, OH)[66,70,71]. Total distance

traveled and instances of vertical rearing were recorded for 60 minutes.

Strength

Inverted Wire Hang Endurance

In the wire hang behavioral test, mice were evaluated for strength and endurance[72]. The apparatus

consisted of a wire mesh (2 mm diameter) suspended above a soft bedding layer to ensure a safe

landing for the mice. Each mouse was gently held by the base of the tail and allowed to grasp the wire

at a 60° incline before the mesh was inverted to 180°. The duration the mouse remained suspended

inverted was recorded up to a maximum of 2 minutes. A minimum 5-minute rest was given between

trials. The maximum time of three trials was recorded.

Grip Strength

Grip strength was measured using a uniaxial grip force tester (Bioseb, France)[73]. Hind paw grip

strength assessed mechanical strength as the mouse was gently restrained and allowed to freely grab

and hold a bar with both hind paws. Maximum force was recorded as the tail was pulled in a steady

manner to avoid rate-dependent effects. A brief rest period, >15 sec, was given between each of the 5

repeated trials. Average strength was reported normalized to body weight at the time of testing and

normalized to pre-surgery baseline.

Cell Culture

Human Primary IVD Cell Culture

Primary human intervertebral disc cells were isolated from the disc tissue of patients undergoing elective discectomy surgeries (Table S3). Human biospecimens were obtained under IRB Exemption Category 4, in accordance with institutional and federal guidelines for research using de-identified, previously collected samples[74,75]. Isolated cells were centrifuged, resuspended, plated, and incubated at 37°C, 5% CO_2, 20% O_2 to grow to 90% confluence.

Vascular Endothelial Growth Factor (VEGF)-siRNA Transfection

Isolated human IVD cells were seeded in 6-well plates at a density of $3x10^5$ cells per well 24 hours prior to transfection. VEGFA-specific siRNA (sc-29620; Santa Cruz Biotechnology, Dallas, Tx) and scrambled control siRNA- FITC (sc-36869; Santa Cruz Biotechnology) were transfected into IVD cells using Lipofectamine™ RNAiMAX Transfection Reagent (Catalog #: 13778100, Thermo Fisher) following the manufacturer's protocol. The final concentration of siRNA in the culture medium was 50 pM. The effectiveness of transfection was evaluated by checking the amount of secreted VEGFA in media using ELISA (DY293; R&D system, Minneapolis, MN) and confirming transfection of control siRNA by FITC fluorescence with a Nikon Eclipse Ti2 inverted scope at 10x.

Co-culturing VEGFA-silenced IVD cells with Immortalized Human Microvascular Endothelial Cell (HMEC-1) and Neuroblastoma Cell (SH-SY5Y) cultures

IVD cells were plated in a 6 well plate and transfected with either control, scrambled siRNA or with VEGFA siRNA. Medium with 1% FBS and recombinant human interleukin-1β (IL-1β; R&D Systems)

at 1 ng/mL was added to stimulate Scram siRNA (IL-1β+Scram siRNA) and VEGFA siRNA (IL-

1β+VEGFA siRNA) transfected IVD cells for 24 hours. After stimulation, HMEC-1 cells "ECs" that

were plated for 24 hours on 1 µm pore co-culture inserts (Thincert cell culture insert; Greiner, Monroe,

NC) at a density of $3x10^5$ cells were inserted to the wells containing Scram siRNA or VEGFA siRNA

transfected IVD cells with MCDB media plus 1% FBS.

HMEC-1 (aka Endothelial cells - ECs) and SH-SY5Y (aka Neuronal cells - NCs) cells were obtained

from the suppliers, cultured and passaged 5-10 times, and then grown to 90% confluence. ECs co-

cultured with stimulated Scram siRNA treated IVD cells are denoted as $EC_{IL-1β+Scram\ siRNA}$ and ECs co-

cultured with stimulated VEGFA siRNA treated IVD cells are denoted as $EC_{IL-1β+VEGF\ siRNA}$. Naïve

ECs that did not undergo co-culturing were used as a control group. Following coculture, the ECs

underwent a Matrigel tube formation assay[76], RNA-sequencing, and a multiplex ELISA of the EC-

derived conditioned media (Figure S4). NCs were plated at a density of 1.5×10^5 cells per well in a 6

well plate and cultured for 48 hours prior to the co-culture experiment. For co-culturing, non-

stimulated IVD cells "naive", IL-1β+Scram siRNA, and IL-1β+VEGFA siRNA transfected IVD cells

were plated on the co-culture insert at a density of 3.0×10^5 cells and co-cultured for 48 hours with

NCs.   Following co-culture, the NCs underwent a neurite outgrowth assay and RNA-sequencing

(Figure S5).

RNA Sequencing and Multiplex protein ELISA

Post co-culture experiments, ECs and NCs were lysed, the total RNA was extracted and sequenced on

an Illumina NovaSeq X Plus. RNA-seq data were processed using standard pipelines for base calling,

alignment, quantification, and normalization. Quality control ensured accurate read alignment,

expression quantification, and removal of low-quality or low-expression genes. Normalized expression

values were then used for downstream differential analysis. The raw data files are available on the

Gene Expression Omnibus database: GSE294577. The conditioned media from naive ECs and those

co-cultured with IL-1β+Scram siRNA and IL-1β+VEGFA siRNA transfected IVD cells was collected

and analyzed by multiplex protein assay.

Statistical and Bioinformatics Analyses

Statistical comparisons on repeated measures, eg the longitudinal behavioral assays, were done using a

linear mixed effect model to assess surgery (Sham/Injured), genotype (WT/VEGFA-null), 3- and 12-

weeks after surgery (3/12), sex (M/F), and surgery (Naïve/Sham). Linear mixed effect models were

initiated with all factors and reduced in complexity as factors were found to be not significant.

Additionally, focused questions were modeled to assess injured groups with genotype and time. A 2 or

3 factor ANOVA was used to assess effects of injury, genotype, and sex on CEμCT, biomechanics,

histopathologic and immunohistochemical outcomes. ANOVA pairwise comparisons were evaluated

with the Fisher's Least Significant Difference method. Survival curves implemented Mantel-Cox test

while binary penetration results used a Fisher's exact test. Simple pairwise comparisons were done

using a Welch's t-test when appropriate. Appropriate adjustments for unequal variances and non-

normal distributions were made where appropriate. The ROUT outlier test with Q=0.1% removed

definitive outliers within groups[77]. R and GraphPad Prism were used to model the results. For gene

ontology analyses, biological pathways enriched by the upregulated and downregulated DEGs were

identified by using the Panther Classification System with version Panther 19.0 by using the statistical

overrepresentation test. For ECs, an adjusted p value of $p < 0.05$ and logFC of 0.41 was used to

identify statistically significant DEGs with at least a 50% increase in fold change of the treatment

group when compared to the control. For neuronal cells, an adjusted p value of $p < 0.05$ and logFC of

0.41 was used to identify statistically significant DEGs with at least a 50% increase in fold change of

the treatment group when compared to the controls. The full list of all DEGs and GO terms for the

endothelial cell's comparisons are listed in Tables S8A and S8B and for the neuronal cell comparisons

are listed in Tables S9A and S9B. Additional details on experimental methods can be found in the

Supplemental Materials.

SUPPLEMENTARY MATERIALS

Supplementary Materials.docx
DEGs Endothelial Cells.xlsx
DEGs Neuronal Cells.xlsx
Gene Ontology Neuronal Cells.xlsx
Gene Ontology Endothelial Cells.xlsx

REFERENCES

1.   Airaksinen O, Brox JI, Cedraschi C, et al. Chapter 4. European guidelines for the management of chronic nonspecific low back pain. *Eur Spine J*. Mar 2006;15 Suppl 2(Suppl 2):S192-300. doi:10.1007/s00586-006-1072-1

2.   Hartvigsen J, Hancock MJ, Kongsted A, et al. What low back pain is and why we need to pay attention. *Lancet*. Jun 9 2018;391(10137):2356-2367. doi:10.1016/S0140-6736(18)30480-X

3.   Yang H, Haldeman S. Behavior-Related Factors Associated With Low Back Pain in the US Adult Population. *Spine (Phila Pa 1976)*. Jan 1 2018;43(1):28-34. doi:10.1097/BRS.0000000000001665

4.   Hoy D, March L, Brooks P, et al. The global burden of low back pain: estimates from the Global Burden of Disease 2010 study. *Ann Rheum Dis*. Jun 2014;73(6):968-74. doi:10.1136/annrheumdis-2013-204428

5.   Shokri P, Zahmatyar M, Falah Tafti M, et al. Non-spinal low back pain: Global epidemiology, trends, and risk factors. *Health Sci Rep*. Sep 2023;6(9):e1533. doi:10.1002/hsr2.1533

6.   Collaborators GBDLBP. Global, regional, and national burden of low back pain, 1990-2020, its attributable risk factors, and projections to 2050: a systematic analysis of the Global Burden of Disease Study 2021. *Lancet Rheumatol*. Jun 2023;5(6):e316-e329. doi:10.1016/S2665-9913(23)00098-X

7.   Rannou F, Lee TS, Zhou RH, et al. Intervertebral disc degeneration: the role of the mitochondrial pathway in annulus fibrosus cell apoptosis induced by overload. *Am J Pathol*. Mar 2004;164(3):915-24. doi:10.1016/S0002-9440(10)63179-3

8.   Feng Y, Egan B, Wang J. Genetic Factors in Intervertebral Disc Degeneration. *Genes Dis*. Sep 2016;3(3):178-185. doi:10.1016/j.gendis.2016.04.005

9.   Wang F, Cai F, Shi R, Wang XH, Wu XT. Aging and age related stresses: a senescence mechanism of intervertebral disc degeneration. *Osteoarthritis Cartilage*. Mar 2016;24(3):398-408. doi:10.1016/j.joca.2015.09.019

10.   Luoma K, Riihimäki H, Luukkonen R, Raininko R, Viikari-Juntura E, Lamminen A. Low Back Pain in Relation to Lumbar Disc Degeneration:. *Spine*. 02/2000 2000;25(4):487-492. doi:10.1097/00007632-200002150-00016

11.   Urban JP. Swelling pressure of the lumbar intervertebral discs: influence of age, spinal level, composition, and degeneration. *Spine*. Feb 01 1988;13(2):179-187.

12.   Buckwalter JA. Aging and degeneration of the human intervertebral disc. *Spine (Phila Pa 1976)*. Jun 1 1995;20(11):1307-14. doi:10.1097/00007632-199506000-00022

13.   Adams MA, Freeman BJ, Morrison HP, Nelson IW, Dolan P. Mechanical initiation of intervertebral disc degeneration. *Spine (Phila Pa 1976)*. Jul 1 2000;25(13):1625-36. doi:10.1097/00007632-200007010-00005

14.   Setton LA, Chen J. Mechanobiology of the intervertebral disc and relevance to disc degeneration. *The Journal of Bone and Joint Surgery American Volume*. 2006-04 2006;88 Suppl 2:52-57. doi:10.2106/JBJS.F.00001

15.   Binch ALA, Fitzgerald JC, Growney EA, Barry F. Cell-based strategies for IVD repair: clinical progress and translational obstacles. *Nat Rev Rheumatol*. Mar 2021;17(3):158-175. doi:10.1038/s41584-020-00568-w

16.   Binch ALA, Cole AA, Breakwell LM, et al. Nerves are more abundant than blood vessels in the degenerate human intervertebral disc. *Arthritis Research & Therapy*. 12/2015 2015;17(1)doi:10.1186/s13075-015-0889-6

17.   Sato J, Inage K, Miyagi M, et al. Inhibiting Vascular Endothelial Growth Factor in Injured
Intervertebral Discs Attenuates Pain-Related Neuropeptide Expression in Dorsal Root Ganglia in Rats.
Asian spine journal. Aug 2017;11(4):556-561. doi:10.4184/asj.2017.11.4.556
18.   Carmeliet P, Ruiz de Almodovar C. VEGF ligands and receptors: implications in
neurodevelopment and neurodegeneration. *Cell Mol Life Sci*. May 2013;70(10):1763-78.
doi:10.1007/s00018-013-1283-7
19.   Falk T, Gonzalez RT, Sherman SJ. The yin and yang of VEGF and PEDF: multifaceted
neurotrophic factors and their potential in the treatment of Parkinson's Disease. *Int J Mol Sci*. Aug 5
2010;11(8):2875-900. doi:10.3390/ijms11082875
20.   Walk RE, Moon HJ, Tang SY, Gupta MC. Contrast-enhanced microCT evaluation of
degeneration following partial and full width injuries to the mouse lumbar intervertebral disc. *Sci Rep*.
Sep 16 2022;12(1):15555. doi:10.1038/s41598-022-19487-9
21.   Millecamps M, Stone LS. Delayed onset of persistent discogenic axial and radiating pain after
a single-level lumbar intervertebral disc injury in mice. *PAIN®*. Sep 01 2018;159(9):1843.
doi:10.1097/j.pain.0000000000001284
22.   Leimer EM, Gayoso MG, Jing L, Tang SY, Gupta MC, Setton LA. Behavioral Compensations
and Neuronal Remodeling in a Rodent Model of Chronic Intervertebral Disc Degeneration. *Scientific
Reports*. 2019-03-06 2019;9(1):3759. doi:10.1038/s41598-019-39657-6
23.   Rickman M, Ghim M, Pang K, et al. Disturbed flow increases endothelial inflammation and
permeability via a Frizzled-4-beta-catenin-dependent pathway. *J Cell Sci*. Mar 15
2023;136(6)doi:10.1242/jcs.260449
24.   Dejana E. The role of wnt signaling in physiological and pathological angiogenesis. *Circ Res*.
Oct 15 2010;107(8):943-52. doi:10.1161/CIRCRESAHA.110.223750
25.   Lohela M, Saaristo A, Veikkola T, Alitalo K. Lymphangiogenic growth factors, receptors and
therapies. *Thromb Haemost*. Aug 2003;90(2):167-84. doi:10.1160/TH03-04-0200
26.   Tammela T, Alitalo K. Lymphangiogenesis: Molecular mechanisms and future promise. *Cell*.
Feb 19 2010;140(4):460-76. doi:10.1016/j.cell.2010.01.045
27.   Cao R, Brakenhielm E, Li X, et al. Angiogenesis stimulated by PDGF-CC, a novel member in
the PDGF family, involves activation of PDGFR-alphaalpha and -alphabeta receptors. *FASEB J*. Oct
2002;16(12):1575-83. doi:10.1096/fj.02-0319com
28.   Page-McCaw A, Ewald AJ, Werb Z. Matrix metalloproteinases and the regulation of tissue
remodelling. *Nat Rev Mol Cell Biol*. Mar 2007;8(3):221-33. doi:10.1038/nrm2125
29.   Kessenbrock K, Plaks V, Werb Z. Matrix metalloproteinases: regulators of the tumor
microenvironment. *Cell*. Apr 2 2010;141(1):52-67. doi:10.1016/j.cell.2010.03.015
30.   Panopoulos AD, Watowich SS. Granulocyte colony-stimulating factor: molecular mechanisms
of action during steady state and 'emergency' hematopoiesis. *Cytokine*. Jun 2008;42(3):277-88.
doi:10.1016/j.cyto.2008.03.002
31.   Hamilton JA. Colony-stimulating factors in inflammation and autoimmunity. *Nat Rev
Immunol*. Jul 2008;8(7):533-44. doi:10.1038/nri2356
32.   Zlotnik A, Yoshie O. The Chemokine Superfamily Revisited. *Immunity*. 05/2012
2012;36(5):705-716. doi:10.1016/j.immuni.2012.05.008
33.   Appay V, Rowland-Jones SL. RANTES: a versatile and controversial chemokine. *Trends
Immunol*. Feb 2001;22(2):83-7. doi:10.1016/s1471-4906(00)01812-3
34.   Lee S, Millecamps M, Foster DZ, Stone LS. Long-term histological analysis of innervation and
macrophage infiltration in a mouse model of intervertebral disc injury-induced low back pain. *J
Orthop Res*. Jun 2020;38(6):1238-1247. doi:10.1002/jor.24560

35.	Freemont AJ, Peacock TE, Goupille P, Hoyland JA, O apos Brien J, Jayson MI. Nerve
ingrowth into diseased intervertebral disc in chronic back pain. *Lancet*. Jul 19 1997;350(9072):178-
181. doi:10.1016/s0140-6736(97)02135-1
36.	Brisby H. Pathology and possible mechanisms of nervous system response to disc
degeneration. *J Bone Joint Surg Am*. Apr 2006;88 Suppl 2:68-71. doi:10.2106/JBJS.E.01282
37.	Lama P, Le Maitre CL, Harding IJ, Dolan P, Adams MA. Nerves and blood vessels in
degenerated intervertebral discs are confined to physically disrupted tissue. *J Anat*. Jul
2018;233(1):86-97. doi:10.1111/joa.12817
38.	Krock E, Rosenzweig DH, Chabot-Dore AJ, et al. Painful, degenerating intervertebral discs up-
regulate neurite sprouting and CGRP through nociceptive factors. *J Cell Mol Med*. Jun
2014;18(6):1213-25. doi:10.1111/jcmm.12268
39.	Chen X, Chen Z, Ma G, et al. Reactive Oxygen Species Induced Upregulation of TRPV1 in
Dorsal Root Ganglia Results in Low Back Pain in Rats. *J Inflamm Res*. 2024;17:2245-2256.
doi:10.2147/JIR.S446841
40.	Kameda T, Zvick J, Vuk M, et al. Expression and Activity of TRPA1 and TRPV1 in the
Intervertebral Disc: Association with Inflammation and Matrix Remodeling. *Int J Mol Sci*. Apr 10
2019;20(7)doi:10.3390/ijms20071767
41.	Shuba YM. Beyond Neuronal Heat Sensing: Diversity of TRPV1 Heat-Capsaicin Receptor-
Channel Functions. *Front Cell Neurosci*. 2020;14:612480. doi:10.3389/fncel.2020.612480
42.	Rosenbaum T, Simon SA. TRPV1 Receptors and Signal Transduction. In: Liedtke WB, Heller
S, eds. *TRP Ion Channel Function in Sensory Transduction and Cellular Signaling Cascades*. 2007.
Frontiers in Neuroscience.
43.	Freemont AJ, Peacock TE, Goupille P, Hoyland JA, O'Brien J, Jayson MI. Nerve ingrowth into
diseased intervertebral disc in chronic back pain. *Lancet*. Jul 19 1997;350(9072):178-81.
doi:10.1016/s0140-6736(97)02135-1
44.	Risbud MV, Shapiro IM. Role of cytokines in intervertebral disc degeneration: pain and disc
content. Review. *Nature reviews Rheumatology*. Jan 2014;10(1):44-56. doi:10.1038/nrrheum.2013.160
45.	Burke JG, Watson RWG, McCormack D, Dowling FE, Walsh MG, Fitzpatrick JM.
Intervertebral discs which cause low back pain secrete high levels of proinflammatory mediators. *The
Journal of bone and joint surgery British volume*. 2002;84(2):196-201.
46.	Freemont AJ. The cellular pathobiology of the degenerate intervertebral disc and discogenic
back pain. *Rheumatology (Oxford)*. Jan 2009;48(1):5-10. doi:10.1093/rheumatology/ken396
47.	Grunhagen T, Wilde G, Soukane DM, Shirazi-Adl SA, Urban JPG. Nutrient Supply and
Intervertebral Disc Metabolism. *Bone and Joint Surgery*. 2006 2006;88-A:30-35.
doi:10.2106/JBJS.E.01290
48.	Clayton SW, Walk RE, Mpofu L, Easson GWD, Tang SY. Sex-specific divergences in the
types and timing of infiltrating immune cells during the intervertebral disc acute injury response and
their associations with degeneration. *Osteoarthritis Cartilage*. Feb 2025;33(2):247-260.
doi:10.1016/j.joca.2024.10.002
49.	Johnson ZI, Schoepflin ZR, Choi H, Shapiro IM, Risbud MV. Disc in flames: Roles of TNF-
alpha and IL-1beta in intervertebral disc degeneration. *Eur Cell Mater*. Sep 21 2015;30:104-16;
discussion 116-7. doi:10.22203/ecm.v030a08
50.	Millecamps M, Tajerian M, Naso L, Sage EH, Stone LS. Lumbar intervertebral disc
degeneration associated with axial and radiating low back pain in ageing SPARC-null mice. *PAIN®*.
Jun 2012;153(6):1167-1179. doi:10.1016/j.pain.2012.01.027

51.   Oehme D, Goldschlager T, Ghosh P, Rosenfeld JV, Jenkin G. Cell-Based Therapies Used to
Treat Lumbar Degenerative Disc Disease: A Systematic Review of Animal Studies and Human
Clinical Trials. *Stem Cells Int*. 2015;2015:946031. doi:10.1155/2015/946031
52.   Barbe MF, Chen FL, Loomis RH, et al. Characterization of pain-related behaviors in a rat
model of acute-to-chronic low back pain: single vs. multi-level disc injury. *Front Pain Res
(Lausanne)*. 2024;5:1394017. doi:10.3389/fpain.2024.1394017
53.   Buettmann EG, McKenzie JA, Migotsky N, et al. VEGFA from Early Osteoblast Lineage Cells
(Osterix+) is Required in Mice for Fracture Healing. *Journal of bone and mineral research : the
official journal of the American Society for Bone and Mineral Research*. May 13 2019:jbmr.3755.
doi:10.1002/jbmr.3755
54.   Ruzankina Y, Pinzon-Guzman C, Asare A, et al. Deletion of the developmentally essential
gene ATR in adult mice leads to age-related phenotypes and stem cell loss. *Cell Stem Cell*. Jun 7
2007;1(1):113-126. doi:10.1016/j.stem.2007.03.002
55.   Gerber HP, Hillan KJ, Ryan AM, et al. VEGF is required for growth and survival in neonatal
mice. *Development*. Mar 1999;126(6):1149-59. doi:10.1242/dev.126.6.1149
56.   Madisen L, Zwingman TA, Sunkin SM, et al. A robust and high-throughput Cre reporting and
characterization system for the whole mouse brain. *Nat Neurosci*. Jan 2010;13(1):133-40.
doi:10.1038/nn.2467
57.   Andersson GB. Epidemiological features of chronic low-back pain. *Lancet*. Aug 14
1999;354(9178):581-5. doi:10.1016/S0140-6736(99)01312-4
58.   Atlas SJ, Deyo RA. Evaluating and managing acute low back pain in the primary care setting. *J
Gen Intern Med*. Feb 2001;16(2):120-31. doi:10.1111/j.1525-1497.2001.91141.x
59.   Melgoza IP, Chenna SS, Tessier S, et al. Development of a standardized histopathology
scoring system using machine learning algorithms for intervertebral disc degeneration in the mouse
model-An ORS spine section initiative. *JOR Spine*. Jun 2021;4(2):e1164. doi:10.1002/jsp2.1164
60.   Scheller E, Brazill JM, Lorenz M. Spatial mapping and contextualization of axon subtypes
innervating the long bones of C3H and B6 mice. protocolsio2021.
61.   Walk RE, Tang SY. In vivo contrast☐enhanced microCT for the monitoring of mouse thoracic,
lumbar, and coccygeal intervertebral discs. *JOR Spine*. Jun 01 2019;2(2)doi:10.1002/jsp2.1058
62.   Lin KH, Wu Q, Leib DJ, Tang SY. A novel technique for the contrast-enhanced microCT
imaging of murine intervertebral discs. *Journal of the Mechanical Behavior of Biomedical Materials*.
10/2016 2016;63:66-74. doi:10.1016/j.jmbbm.2016.06.003
63.   Easson GWD, Savadipour A, Anandarajah A, et al. Modulation of TRPV4 protects against
degeneration induced by sustained loading and promotes matrix synthesis in the intervertebral disc.
FASEB J. Feb 2023;37(2):e22714. doi:10.1096/fj.202201388R
64.   Goh J, Ladiges W. Voluntary Wheel Running in Mice. *Curr Protoc Mouse Biol*. Dec 2
2015;5(4):283-290. doi:10.1002/9780470942390.mo140295
65.   Chaplan SR, Bach FW, Pogrel JW, Chung JM, Yaksh TL. Quantitative assessment of tactile
allodynia in the rat paw. *J Neurosci Methods*. Jul 1994;53(1):55-63. doi:10.1016/0165-
0270(94)90144-9
66.   Sheahan TD, Siuda ER, Bruchas MR, et al. Inflammation and nerve injury minimally affect
mouse voluntary behaviors proposed as indicators of pain. *Neurobiol Pain*. Aug-Dec 2017;2:1-12.
doi:10.1016/j.ynpai.2017.09.001
67.   Duraku LS, Hossaini M, Hoendervangers S, et al. Spatiotemporal dynamics of re-innervation
and hyperinnervation patterns by uninjured CGRP fibers in the rat foot sole epidermis after nerve
injury. *Mol Pain*. Aug 30 2012;8:61. doi:10.1186/1744-8069-8-61

68. Allen KD, Griffin TM, Rodriguiz RM, et al. Decreased physical function and increased pain
sensitivity in mice deficient for type IX collagen. *Arthritis Rheum*. Sep 2009;60(9):2684-93.
doi:10.1002/art.24783
69. Kamisli S, Karakurt A, Uyumlu AB, et al. Evaluation of the effects of novel nafimidone
derivatives on thermal hypoalgesia in mice with diabetic neuropathy. *Balkan Med J*. Mar
2013;30(1):94-8. doi:10.5152/balkanmedj.2012.067
70. Samineni VK, Yoon J, Crawford KE, et al. Fully implantable, battery-free wireless
optoelectronic devices for spinal optogenetics. *Pain*. Nov 2017;158(11):2108-2116.
doi:10.1097/j.pain.0000000000000968
71. Jain S, Golden JP, Wozniak D, Pehek E, Johnson EM, Jr., Milbrandt J. RET is dispensable for
maintenance of midbrain dopaminergic neurons in adult mice. *J Neurosci*. Oct 25 2006;26(43):11230-
8. doi:10.1523/JNEUROSCI.1876-06.2006
72. Deacon RM. Measuring the strength of mice. *J Vis Exp*. Jun 2 2013;(76)doi:10.3791/2610
73. Meyer OA, Tilson HA, Byrd WC, Riley MT. A method for the routine assessment of fore- and
hindlimb grip strength of rats and mice. *Neurobehav Toxicol*. Fall 1979;1(3):233-6.
74. Barcellona MN, Speer JE, Fearing BV, et al. Control of adhesive ligand density for modulation
of nucleus pulposus cell phenotype. *Biomaterials*. Aug 2020;250:120057.
doi:10.1016/j.biomaterials.2020.120057
75. Bridgen DT, Fearing BV, Jing L, et al. Regulation of human nucleus pulposus cells by peptide-
coupled substrates. *Acta Biomater*. Jun 2017;55:100-108. doi:10.1016/j.actbio.2017.04.019
76. Moon HJ, Yurube T, Lozito TP, et al. Effects of secreted factors in culture medium of annulus
fibrosus cells on microvascular endothelial cells: elucidating the possible pathomechanisms of matrix
degradation and nerve in-growth in disc degeneration. *Osteoarthritis Cartilage*. Feb 2014;22(2):344-
54. doi:10.1016/j.joca.2013.12.008
77. RE MHaB. Detecting outliers when fitting data with nonlinear regression – a new method
based on robust nonlinear regression and the false discovery rate. *BMC Bioinformatics*. 2006;7(123)

List of Abbreviations
AF – Annulus Fibrosus
Ai9 – Molecular reporter expressing tdTomato fluorescent protein (Rosa-CAG-LSL-tdTomato)
ANOVA – Analysis of Variance
CEμCT – Contrast-Enhanced Micro-Computed Tomography
cLBP – Chronic Low Back Pain
CM – Conditioned Media
CTCF – Corrected Total Cell Fluorescence
C57BL/6J – Common inbred mouse strain
DAPI – 4′,6-Diamidino-2-Phenylindole (DNA stain)
DEG – Differentially Expressed Gene
DHR – Disc Height Ratio
DRG – Dorsal Root Ganglion
EC – Endothelial Cell (HMEC-1)
EDTA – Ethylenediaminetetraacetic Acid (chelating agent)
ELISA – Enzyme-Linked Immunosorbent Assay
EMCN – Endomucin (endothelial cell marker)
EMEM, ATCC – Eagle's Minimum Essential Medium, from American Type Culture Collection
ERT2 – Modified Estrogen Receptor Ligand Binding Domain
EVF – Electronic Von Frey
FBS – Fetal Bovine Serum
FITC – Fluorescein Isothiocyanate dye
FSU – Functional Spinal Unit
HBSS – Hanks' Balanced Salt Solution
HMEC-1 – Human Microvascular Endothelial Cell Line
IL1β – Interleukin 1 Beta
IVD – Intervertebral Disc
LBP – Low Back Pain
LoxP – Locus of X-over P1 (recombination sites for Cre-Lox systems)
LSD – Least Significant Difference (statistical post hoc test)
MCDB – Molecular, Cell, and Developmental Biology
Nav1.8 – Voltage-Gated Sodium Channel (gene SCN10A) expressed in nociceptive neurons
NC – Neuron cells (SH-SY5Y)
NFkB – Nuclear Factor Kappa B
NGF – Nerve Growth Factor
NI/DI – Nucleus Attenuation Intensity/Disc Attenuation Intensity
NP – Nucleus Pulposus
OCT – Optimal Cutting Temperature Compound
PAP – Hydrophobic barrier pen
PBS – Phosphate-Buffered Saline
PGP9.5 – Protein Gene Product 9.5 (neuronal marker)
P/S – Penicillin/Streptomycin
ROI – Region of Interest
SCRAM – Scrambled siRNA (negative control)
SH-SY5Y – Human Neuroblastoma Cell Line
siRNA – Small Interfering RNA

SNT – Simple Neurite Tracer (ImageJ plugin)
TNFα – Tumor Necrosis Factor Alpha
TNT – Tris-NaCl-Tween buffer
TrKA – Tropomyosin Receptor Kinase A (NGF receptor)
TRPA1 – Transient Receptor Potential Ankyrin 1
TRPV1 – Transient Receptor Potential Vanilloid 1
UBC – Ubiquitin
VEGFA – Vascular Endothelial Growth Factor Type A
WT – Wild Type
925

Acknowledgments:
We acknowledge the support of the Washington University Musculoskeletal Research Center
Structure and Function Core (Michael Brodt); Histology Core (Crystal Idleburg and Samantha
Coleman); and Animal Behavior and Function Core. We also acknowledge Kelsey Collins and
Washington University Pain Center Judith Golden for their training in behavioral assessments of pain.
We thank the Washington University Genomics Technology Access Core (GTAC), and the Eve-tech
for the multiplex ELISA analysis.
933

Funding:
This work was conducted with funding support from National Institutes of Health. The content is
solely the responsibility of the authors and does not necessarily represent the official views of the
National Institutes of Health.
National Institutes of Health grant R21AR081517 (ST/MG)
National Institutes of Health grant R01AR074441 (ST)
National Institutes of Health grant R01AR077678 (LS/ST)
National Institutes of Health grant P30AR074992 (MS)
National Institutes of Health grant S10OD028573 (MS)

Author contributions:
Conceptualization: RSP, MJS, LAS, MCG, SYT
Methodology: RSP, HJM, SWC, REW, ALL, ANS, LJ, EGB, JAM, ATB, LAS, MCG, SYT
Data Interpretation: Methodology: RSP, REW, HJM, ALL, ANS, SWC, LJ, LAS, MCG, SYT
Investigation: RSP, HJM, SWC, ALL, LJ, LAS, MCG, SYT
Visualization: RSP, HJM, SWC, ALL, SWC
Funding acquisition: LAS, MCG, SYT
Project administration: RSP, SYT
Supervision: SYT
Writing – original draft: RSP, HJM, SWC, ALL
Writing – review & editing: RSP, SWC, SYT
954

955 **Competing interests:**
956 Provisional Patent: 020859/US
957 Title: VEGFA INHIBITORS AS A TREATMENT FOR CHRONIC MECHANICAL ALLODYNIA
958 Co-inventors: RSP, HJM, MCG, SYT
959

Data and materials availability:

All data, code, and materials used will be made available to other researchers upon reasonable request
for the proposes of reproducing or extending the analysis. Vegfa$^{fl/fl}$ mice were obtained via a materials
transfer agreements (MTAs) with Genentech Inc and provided by MJS. The raw data files of the
endothelial cell and neuron gene expression analyses are available on the Gene Expression Omnibus
database under accession number GSE294577. Supplemental materials contains additional methods
and data in support of this study.