1

1    Dual inhibition of EGFR and PI3Kinase signaling in EGFR amplified triple negative breast

2                    cancer cells induces apoptosis and reduces tumor growth

David J. Wisniewski[1], Donna Voeller[1], Yonit A. Addissie[1], Sachin Kumar Deshmukh[2], Sharon

Wu[2], Maryam B. Lustberg[3], Darawalee Wangsa[4], Danny Wangsa[4], Kerstin Heselmeyer-

Haddad[4], George W. Sledge Jr.[2] and *Stanley Lipkowitz[1]

[1]Women's Malignancies Branch, Center for Cancer Research, National Cancer Institute,

Bethesda MD

[2]Caris Life Sciences, Phoenix AZ

[3]Yale Cancer Center, New Haven CT

[4]Genetics Branch, Center for Cancer Research, National Cancer Institute, Bethesda MD

*Corresponding Author

**Keywords: epidermal growth factor receptor amplification, PI3Kinase pathway mutations,**

**triple negative breast cancer**

**Abstract**

**Background**: Epidermal Growth Factor Receptor (EGFR) family signaling is commonly dysregulated in cancer by amplification or activating mutations. We investigated the incidence of EGFR amplification with or without PI3Kinase pathway mutations in breast cancer and whether patients with genetic abnormalities in both pathways can be targeted by dual EGFR/PI3K inhibition. **Methods:** EGFR amplification and PI3K pathway mutations were studied through data sets from cBioPortal and Caris. We used the triple negative breast cancer (TNBC) cell lines BT20 (amplified EGFR, PIK3CA activating mutation), MDA-MB-468 (amplified EGFR, PTEN deletion), and MDA-MB-231 (no amplification of EGFR or PI3K pathway mutations). EGFR amplification was determined by immunoblot and fluorescent in situ hybridization, and PI3K mutations by sequencing. Signaling was determined by immunoblot, drug synergy by cell viability, cell death by propidium iodide staining, cell cycle analysis by flow cytometry, and animal studies through xenografts. **Results:** EGFR amplification is found in approximately 1-2.5% of breast cancer patients, more frequently in TNBC (2.45-6.7%) and ER-/HER2+ (1.3-6.5%) breast cancers, and in the molecular basal (2.33-8.1%) and HER2 enriched (1.87-5.4%) subtypes. Overall survival is shorter among patients compared to those with unamplified EGFR. Up to 71% of EGFR amplified tumors have activating mutations in the PI3K pathway. Combination of EGFR and PI3K inhibitors more dramatically reduced mTOR and AKT signaling in the BT20 and MDA-MB-468 cells, whereas the inhibition of downstream signaling was less significant in MDA-MB-231 cells. Combination of EGFR and PI3K inhibitors reduced cell viability in these three cell lines, but inhibition was greater, statistically significant, and synergistic in MDA-MB-468 and BT20 compared to MDA-MB-231. Only MDA-MB-468 and BT20 cells had an increased fraction of apoptotic cells. EGFR or PI3K inhibition alone in a

45 BT20 xenograft model reduced tumor volume, however the combination was the only

46 statistically significant reduction in tumor volume when compared to vehicle control.

47 **Conclusions:** EGFR/PI3K inhibitor combination causes apoptosis and reduction in tumor growth

48 in cells with EGFR amplification and PI3K alteration. Dual inhibition of EGFR/PI3K presents as

49 a potential targeted therapy in patients with EGFR amplification and aberrant PI3K signaling.

**Background**

The ErbB family is comprised of four receptor tyrosine kinases, Epidermal Growth

Factor Receptor (EGFR), HER2 (ErbB2), HER3 (ErbB3) and HER4 (ErbB4).  Typically, upon

ligand binding, the receptors either hetero or homodimerize which allows them to activate

downstream signaling.  Of note, HER2 does not bind any known ligands but instead is activated

by forming hetero or homodimers with other ligand bound EGFR family members (1). EGFR is

typically activated upon binding of a ligand such as epidermal growth factor (EGF), resulting in

EGFR homodimerization (or heterodimerization with other EGFR receptor tyrosine kinases), and

trans-phosphorylation of multiple tyrosine residues on the receptors.  Upon activation, EGFR

signals through various downstream pathways, including phosphoinositide 3-kinases (PI3Ks) or

mitogen-activated protein kinases (MAPKs), resulting in increased proliferation, inhibition of

apoptosis, migration, invasion, and other pro-tumor signaling (2-5).

Mutation or overexpression of these family members are common drivers of various

cancer types (6).  For example, HER2 amplification is a driver in approximately 12-20% of

breast cancers, correlates with poor survival, and has been effectively targeted to improve

outcomes in patients with HER2 amplified breast cancer (7-10).  HER2 activating mutations in

the tyrosine kinase domain are found in 2-3% of breast cancers and are likely to behave similarly

to HER2 amplification, and HER2 mutants are known to dimerize with EGFR, and induce

phosphorylation of EGFR (11).

EGFR can be aberrantly activated through mutations or amplification/overexpression,

which drives pro-tumor effects in cancers such as non-small cell lung cancer (NSCLC),

pancreatic cancer and head and neck cancer (2, 3, 12-15).  Due to the driving effect of EGFR on

cancer progression, various therapies have been approved for use in patients with EGFR

abnormalities, such as the tyrosine kinase inhibitors (TKI's) gefitinib, erlotinib, afatinib,

dacomitinib, osimertinib, and vandetanib, and the monoclonal antibodies cetuximab,

panitumumab and necitumumab (2). EGFR has been shown to be expressed in basal triple

negative breast cancer (TNBC), however there are various reports which define the degree of

EGFR overexpression differently with varying methods (16). In general, EGFR has some degree

of elevated expression by immunohistochemistry in breast cancer samples (17-23). Further,

EGFR expression was associated with tumor size, poor differentiation, worse relapse-free and

overall survival, and shorter metastasis free survival (24, 25). Depending on the study, EGFR

gene amplification has been observed in 0.8-14% of breast cancer patients and is associated with

high grade but has not always been associated with survival (20, 26-28). Lastly, mutation of

EGFR in breast cancer can range from approximately 2-11% (22). Relatively small studies

looking specifically at metaplastic breast carcinomas have observed amplification of EGFR in

33-37% of patients (29, 30). Despite these data, clinical trials investigating EGFR inhibitors

have not shown great promise in breast cancer, with studies finding partial response rates of 0-

6% (18, 20, 22, 31, 32). It is important to note that these studies were typically in heavily

pretreated patients, and patients were not stratified by EGFR amplification status. Therefore,

investigation of combination therapy targeting EGFR and other pathways may elucidate the

functional benefits of targeting a subset of breast cancer patients with EGFR amplification and

provide insight into optimal therapy.

In this study we evaluated the frequency and subtype distribution of EGFR amplification

and found that EGFR is amplified in approximately 1-2.5% of breast cancer cases, with the most

frequent amplification in TNBC, ER-, and ER-/HER2+ tumors, HER2 enriched, basal, and

normal subtypes, depending on the dataset studied. Further, we found that up to 71% of EGFR

amplified breast cancers had an activating alteration of the PI3K pathway. Using TNBC models

of EGFR amplification and PI3K alteration, we demonstrate that dual inhibition of EGFR and

PI3K significantly reduced P70S6K, AKT and S6 signaling and cell viability.  Further, dual

inhibition caused G1 accumulation, and reduction in S phase and G2/M phase, with a caspase-

dependent increase in cancer cell death.  Lastly, dual inhibition statistically significantly reduced

tumor volume *in vivo* as compared to control.

## Methods

### Materials

Z-VAD-FMK (S7023), Erlotinib (S7786), BKM120 (S2247), Afatinib (S1011), Alpelisib

(S2814) and sodium carboxymethyl cellulose (S6703) were purchased from Selleck Chemicals.

Fetal bovine serum (FBS) (631106) was obtained from Takara Bio. 4-20% precast

polyacrylamide gels (catalog #5671094) were purchased from Bio-Rad. Immobilon PVDF

transfer membrane (IPVH00010), mini protease inhibitor cocktail (11836153001) and propidium

iodide (P4864) was purchased from Millipore-Sigma. PBS (21-031-CV), EGF (354042) and

Phenol Red-Free LDEV-Free Matrigel (356237) was purchased from Corning. Sodium

orthovanadate (P0758S) was from Fisher Chemicals. NP40 Cell Lysis Buffer (FNN0021),

Penicillin-Streptomycin (15140-122), tissue extraction reagent (FNN0071) and RPMI 1640

medium (11875-093) was obtained from ThermoFisher Scientific. were purchased from Selleck

Chemicals (Houston, TX).

### Antibodies

Anti-phospho-HER3 Y1289 (4791S), anti-HER3 (12708S), anti-EGFR (2232), anti-

PTEN (9559S), anti-phospho-EGFR Y1068 (2234), anti-phospho-AKT S473 (9271S), anti-AKT

(9272S), anti-phospho-MAPK T202/Y204 (9101), anti-phospho-p70S6K T389 (9234S), anti-

p70S6K (9202S), anti-phospho-S6 S235/236 (4856S), anti-S6 (2217S) and anti-PI3K (4249S)

were purchased from Cell Signaling Technology. anti-HSC70 (sc-7298) and anti-ERK2 (D-2,

sc-1647) antibodies were purchased from Santa Cruz Biotechnology. Goat Anti-Mouse IgG

HRP conjugate (172-1011) and Goat Anti-Rabbit IgG HRP conjugate (1721019) were purchased

from BioRad.

**Cell Lines**

MDA-MB-468, BT20, MDA-MB-231 were obtained from ATCC, were maintained in RPMI

1640 medium with 10% FBS and 1% penicillin/streptomycin, and grown at $37^{\circ}C$, 5% $CO_2$.

**Caris dataset analysis**

*Next-generation sequencing methods (WES, Hybrid)*

Because this study utilized samples spanning several years, the methods (including materials and

software) used for DNA and RNA profiling evolved over time. Initially, DNA was profiled using

whole exome sequencing (WES) with targeted enrichment of 720 clinically relevant genes in

2020; whole transcriptome sequencing (WTS) was performed separately. In 2023, this approach

was replaced by MI Tumor Seek Hybrid, which analyzes RNA and DNA from the same total

nucleic acid extraction. Nucleic acid was extracted using appropriate FFPE kits for RNA, DNA,

or total nucleic acid. DNA sequencing was performed using by whole exome sequencing with

enrichment of 720 clinically relevant genes on the NovaSeq 6000 platform (RRID:SCR_016387)

(Illumina, Inc., San Diego, CA). WTS was performed using the Illumina Novaseq 6000 platform

to an average of 60M reads. Raw WTS data was demultiplexed by Illumina Dragen BioIT

accelerator, trimmed, counted, PCR-duplicates removed, and aligned to human reference genome

(hg19/hg38) by STAR aligner (RRID:SCR_004463) (33). For transcription counting, transcripts

per million (TPM) molecules were generated using the Salmon expression pipeline

(RRID:SCR_017036) (34). For MI Tumor Seek Hybrid, RNA was labeled during first strand

cDNA synthesis by adapter sequences on the 5' end of the cDNA primers and next-generation

sequencing was performed (720 clinically relevant genes were sequenced at high-depth).

Sequencing data was extracted into split FASTQ files (RNA and DNA) for processing. DNA

variants detected were mapped to reference genome hg38, and well-established bioinformatics

tools such as Burrows-Wheel Aligner (BWA 0.7.17) (RRID:SCR_010910), SamTools

(RRID:SCR_002105), Pindel (RRID:SCR_000560), and snpFF (RRID:SCR_005191) were

incorporated to perform variant calling functions; germline variants were filtered with various

germline databases, including dbSNP (RRID:SCR_002338). Genetic variants identified were

interpreted by board-certified molecular geneticists and categorized as 'pathogenic,' 'likely

pathogenic,' 'variant of unknown significance,' 'likely benign,' or 'benign,' according to the

American College of Medical Genetics and Genomics (ACMG) standards. Pathogenic and

likely-pathogenic variants are counted as "reportable". A copy number (CN) cutoff of $CN \geq 6$

was used to define gene amplification. The CN cutoff of 6 for amplification was determined

internally at Caris as a standard (based on MYC/ERBB2 and validated with IHC).

*Real-world overall survival analysis*

Real-world overall survival (OS) information was obtained from insurance claims data. OS was

calculated from time of tissue collection as a surrogate for diagnosis until last contact. Patients

without contact/claims data for a period of at least 100 days were presumed deceased.

Conversely, patients with a documented clinical activity within 100 days prior to the latest data

update were censored in the analysis. Kaplan-Meier estimates were calculated for molecularly

defined patient cohorts. Hazard ratios (HR) were determined by Cox Proportional Hazards model

and *p*-values by log-rank test. Significance was determined as $p < 0.05$.

**cBioPortal dataset analysis**

Datasets from TCGA (Cell 2015), Metabric (Nature 2012 and Nature Communications 2016)

and MSKCC (Cell 2018) were searched for EGFR amplification, as well as mutations in

PIK3CA, PIK3R1, PIK3R2, PIK3R3, PTEN, AKT1, AKT2, AKT3, RPS6KB1, RPS6KB2 and

RPS6 in cBioPortal (cbioportal.org) (35-37). Amplification was defined by GISTIC2.0 score = 2

which corresponds to high levels of amplification (copy number > 5) (38).

**Western Blot**

Cell lysates were collected, prepared and immunoblotted as previously described (39). Every

immunoblot was repeated at least 3 times, with densitometric analysis of band intensities being

calculated using ImageJ.  Data were presented as an average ± SEM.

**Fluorescence *in situ* Hybridization (FISH)**

To prepare metaphase chromosomes, cells were treated with 0.02 mg/ml Colcemid

(Invitrogen, Grand Island, NY) for one hour, suspended in a hypotonic solution for 20 minutes,

and then fixed using a methanol/acetic acid mixture (1:3 ratio). The cell suspension was dropped

onto slides in a cytogenetic drying system (Thermotron, Holland, MI).

Dual color FISH was conducted using a contig of three overlapping bacterial artificial

chromosome (BAC) clones containing the EGFR gene on chromosome band 7p11.2, labeled in

red, and a centromere probe for chromosome 7 (CEP7) labeled in aqua. The EGFR and CEP7

probes were custom manufactured to our specifications by Cytotest (Rockville, MD), with dyes

sourced from Dyomics (Jena, Germany).

In brief, the FISH hybridization protocol included incubating the slides in 1× PBS for 10

minutes, followed by treatment in a pre-warmed pepsin solution (final concentration: 0.02 µg/µl

pepsin and 0.01N HCl in dH$_2$O) at 37°C for 2 to 5 minutes. The slides were then washed in 1x

PBS, dehydrated in an ethanol series, and left to air-dry. Co-denaturation was performed at 72°C

for 2 minutes, followed by overnight hybridization at 37°C on a ThermoBrite (Abbott Molecular,

Des Plaines, IL). The detection protocol involved washing the slides in 2× SSC/0.3% Nonidet P-

40 for 2 minutes at 48°C, followed by a wash in 2× SSC/0.1% Nonidet P-40 for 1 minute at

room temperature. Finally, an antifade solution (Vector Laboratories, Burlingame, CA)

containing the nuclear counterstain 4,6-diamidino-2-phenylindole was applied, and coverslips

were added. Images were captured using a Leica DM-RXA fluorescence microscope (Leica,

Wetzlar, Germany) equipped with a 40 X objective and custom optical filters (Chroma, Bellows

Falls, VT).

**Sanger Sequencing**

Sanger DNA sequencing of genomic DNA from the MDA-MB-231, MDA-MB-468 and BT20

cell lines for PIK3CA mutations was performed using the Big Dye Terminator v.1.1 Cycle

Sequencing Kit (Applied Biosystems, CA) according to the manufacturer's specifications and

run on an ABI 3130xl or 3730 Genetic Analyzer (Applied Biosystems, CA). Sequencing was

conducted at the CCR Genomics Core at the National Cancer Institute, NIH, Bethesda, MD.

Primers were synthesized by (Eurofins Genomics) as follows:

exon 9 sequencing primer (AAGATTTGCTGAACCCTATTG);

exon 20 sequencing primer (TCGTCACAATAGTAACATCATGG).

**Cell Death**

CytoTox-Glo (G9291) was performed as directed by Promega.  Briefly, cells were pretreated

with 20 µM ZVAD for 1 hour, followed by co-treatment with DMSO, 10 µM erlotinib, 1 µM

BKM120 or the combination for 48 hours.  Dead cell percentage was then calculated according

to the manufacturer instructions.

**Cell Viability**

Viability was determined by CellTiter-Glo 2.0 (Catalog #G9242) as directed by Promega.

Briefly, 10,000 cells were plated in a 96-well white walled plate (Corning #3610) and treated the

following day with experimental conditions for 48 hours in 1% FBS RPMI. Reagent was added

to each well, the plate was shaken for two minutes and then incubated at room temperature for 10

minutes. Luminescence was then recorded using a plate reader.

**Synergy Scores**

Calculated using SynergyFinder.org, using a matrix format with data calculated as percent

viability. Data was interpreted as previously described (40).

**Cell Death**

Cell death was determined by using propidium iodide (PI) staining, imaging and quantification

using the BioTek Cytation 1 Imaging Reader from Agilent (Santa Clara, CA). Briefly, 5,000

cells were plated in a 96-well plate, and treated the next day with experimental conditions in 1%

FBS RPMI, along with PI (1:3000 dilution). Time 0 images were acquired, and then cells were

incubated at normal cell culture conditions for 72 hours, followed by a second image acquisition.

Dead cell percentage was calculated by dividing dead cell number (PI stained) by brightfield cell

count number and multiplying by 100. Dead cell percentage was normalized relative to the Day

0 dead cell percentage to account for variation among wells. PI staining positive and brightfield

cell quantification was optimized and performed in a cell-line specific manner, using Gen5

Image Prime 3.14 software (Agilent). In experiments using ZVAD-FMK, cells were pretreated

+/- 20 µM ZVAD-FMK for 1 hour, followed by co-treatment with experimental conditions.

**Cell cycle**

Cells were plated in 6-well plates ($3\times10^5$ cells/well) and treated with EGFR and/or PI3K

inhibitors. After 48 hours, cells were harvested, washed with PBS and fixed in 70% ethanol for

at least 1 hour at 4°C. Cells were then washed twice with cold PBS after which 50 ul of 100

ug/ml RNAse A and 200 ul of 50 µg/ml propidium iodide were added. After incubation for 1

hour at room temperature, cell cycle data was acquired using a BD LSRFortessa™ (BD

Biosciences) flow cytometer and analyzed using FlowJo® software (FlowJo LLC).

**Animal Studies**

Sixty female athymic nude mice between 5-6 weeks in age were obtained (Charles River

Laboratories) and were housed and observed according to approved NCI-ACUC guidelines. 5

million BT20 cells suspended in matrigel (1:1 with PBS) were injected subcutaneously into the

mammary fat pad with a 25G needle (Terumo Medical Care Solutions; SS-01T2516). Once the

tumors grew to 100 mm$^3$ the mice were randomized into four groups of 15. 0.5% sodium

carboxymethyl cellulose (CMC) in water (w/v) was used as control, and afatinib or alpelisib

were suspended in 0.5% CMC. Control or 20 mg/kg alpelisib or afatinib, or the combination of

both was given by oral gavage (1 inch, 22G needle) once a day for five days each week for the

duration of the study. After the first four days of treatment, 5 mice per group were sacrificed,

tumors were excised and proteins were harvested in Tissue Extraction Reagent I (Invitrogen-

FNN0071) according to the manufacturers protocol. Tumor caliper measurements and body

weight were measured twice a week for the duration of the study, as required by humane

endpoints for all mice. Tumor volumes were calculated according to the formula $V = \frac{1}{2}$

(length $\times$ width$^2$). All aspects of this study were approved by the NCI Animal Care and Use

Committee (IACUC number WMB-004) and performed in accordance with the approved

protocol.  Maximum tumor size for each animal was 20 mm in any direction, and this maximum

tumor size was not exceeded.  As soon as tumors reached maximum size, the mice were

humanely euthanized.

**Statistical Analysis**

Student's *t*-test with 2 tailed comparisons assuming equal variance, one-way or two-way

ANOVA were performed as indicated.  P-values of $\leq 0.05$ were considered significant.

**Results**

**Incidence of EGFR amplification in breast cancer**

The frequency of EGFR amplification in breast cancer was evaluated in the TCGA,

METABRIC and MSKCC databases in cBioPortal (35-37) . EGFR amplification was observed

in 1.2-2.4% of cases, depending on the dataset (Figure 1A-C). The TCGA and METABRIC

databases contain primarily early-stage primary breast cancers while the MSKCC dataset is

approximately 50% metastatic/recurrent breast cancers. We analyzed EGFR amplification by

subtype and found that amplification is more common in invasive ductal carcinomas (IDC) (1.4-

2.9%) compared to invasive lobular carcinomas (ILC) (0-0.6%). Further, the highest rate of

amplification was in TNBC (3.2-6.7%), ER- (2.8-5.4%) and ER-/HER2+ tumors (1.3-6.5%),

whereas EGFR amplification was relatively less common in ER+ breast cancers (0.8-1.4%).

Using PAM50 classification, EGFR amplification is the highest in basal (4.3-8.1%) and HER2

enriched (3.9-5.4%) molecular subtypes, compared to luminal A or luminal B subtypes (0.7-1%

and 1.5-1.6%, respectively) (41). To determine the prognosis of patients with EGFR

amplification, we combined the TCGA, METABRIC and MSKCC datasets due the small

number of cases in each and evaluated overall survival (OS) (Figure 1D). There is a statistically

significant lower OS in patients with amplified EGFR, with a difference of 53.5 months (111.4

vs 164.6 months) (Figure 1D).

To confirm these data in a larger dataset with clinical annotation, we evaluated the

incidence of EGFR amplification in patients with breast cancer profiled by Caris Life Sciences

(hereafter referred to as Caris), where the majority of patients profiled had metastatic breast

cancer (Figure 1E). EGFR amplification was found in 1.03% of the breast cancers profiled. The

lower frequency of EGFR amplification seen in the Caris data set may be due to higher cutoff

used for calling amplification, since cBioportal uses a GISTIC2.0 score of 2 to indicate

amplification which corresponds to CN ≥4-5 while Caris used a CN ≥ 6 as the cutoff for

amplification.  As in the other datasets, EGFR amplification was more prevalent in IDC (1.29%)

compared to ILC (0.12%). EGFR amplification was more prevalent in TNBC (2.45%), and in the

molecular HER2 enriched (1.87%), basal (2.33%) and normal (1.43%) subtypes.  In the entire

cohort of patients, OS was statistically significantly shorter in patients with EGFR amplification

compared to those with wild type EGFR (21.6 vs 33 months; Figure 1F).  Overall, this data

shows that there is a subset of breast cancer patients with EGFR amplification that is most

frequent in TNBC and ER-/HER2 amplified tumors and is associated with a poor prognosis.

Since EGFR amplification is enriched in patients with TNBC or Basal cancers, we

investigated the differences in overall survival in patients with EGFR amplification depending on

subtype using the Caris dataset (Additional File 1).  There was no OS difference in TNBC or

Basal cancers with EGFR amplification (Additional File 1 A and B).  HER2 enriched patients

with EGFR amplification had a trend for a slightly worse overall survival (difference of 6.7

months), however this was not statistically significant (p=0.17) (Additional File 1 C).  In

contrast, patients with ER+/HER2- tumors had a statistically significant worse OS in EGFR

amplified patients (difference 22.9 months; p<0.00001) (Additional File 1 D).

**_In vitro_ modeling of EGFR amplification and PI3K alterations**

To evaluate the role of EGFR amplification in cell models, we used two cell lines with

amplified EGFR (MDA-MB-468 and BT20) and one cell line without amplified EGFR (MDA-

MB-231).  As evidenced in Figure 2A, MDA-MB-468 and BT20 both have elevated EGFR

protein levels as compared to MDA-MB-231.  Fluorescence _in situ_ hybridization (FISH) analysis

confirmed the amplification of EGFR in both MDA-MB-468 and BT20, and the lack of EGFR

amplification in MDA-MB-231 (Figure 2B). MDA-MB-468 has loss of PTEN protein, which

supports previously published studies showing loss of PTEN expression in these cells (Figure

2A) (42, 43). As previously reported, the BT20 TNBC cell line has two activating mutations in

PIKCA (P539R and H1047R), which previously have been shown to be in cis (43-45) (Figure

2C). MDA-MB-468 and MDA-MB-231 have wildtype PIK3CA. Therefore, MDA-MB-468 and

BT20 were the two cell lines with EGFR amplification and PI3Kinase pathway activating

mutations, and MDA-MB-231 was the control cell line without amplified EGFR or PI3Kinase

pathway mutations.

**PI3K pathway mutations in EGFR amplified breast cancer**

Considering the co-incidence of mutations or alterations in PI3K signaling in these EGFR

amplified cell lines, we investigated how often this occurs in patients. Once again using

cBioPortal, in a total of 75 patients with EGFR amplification from the TCGA, METABRIC and

MSKCC cohorts, we discovered that approximately 53 patients (71%) with EGFR amplified

tumors have at least one genetic alteration which can potentially activate PI3K signaling (Figure

3A and Additional File 2). 36% have an activating mutation in PIK3CA, 5.3% have a loss of

PTEN, 12% have mutations in PIK3R1, R2 or R3, and 29.3% have amplification of AKT1, 2, or

3. Also, there are 16% of patients who have amplifications downstream of mTOR (either in

ribosomal protein S6 kinase B1 or B2, or ribosomal protein S6). Of note, multiple tumors had

two or more alterations in this pathway, which can be observed in the oncoplot of the EGFR

amplified tumors (Additional File 2).

Similarly in the Caris dataset, 54.3% of EGFR amplified tumors have mutation or

amplification in the PI3K pathway (including PIK3CA, pTEN, PIK3R1, PIK3R2, or AKT 1, 2 or

3) (Figure 3B). The largest percentage of alterations are in PIK3CA (39%). Of note, the Caris

dataset does not include alterations in the genes downstream of mTOR, which contributes to the

difference comparing the sums of Caris and the cBioPortal data.  This data confirms that the

majority of tumors with EGFR amplification are likely to have co-incident aberrant PI3K

signaling.

**EGFR and PI3K dual inhibition significantly reduce downstream signaling in EGFR**

**amplified and PI3K altered breast cancer**

To elucidate the effect of EGFR signaling in these two EGFR amplified cell lines, we

treated the cells with epidermal growth factor (EGF) and the tyrosine kinase inhibitor (TKI)

afatinib and observed that EGFR phosphorylation was decreased upon afatinib treatment in both

MDA-MB-468 and BT20 cell lines, indicating that the drug inhibited EGFR signaling upon EGF

stimulation (Figure 4A).  We confirmed this observation by using another TKI, erlotinib

(Additional File 3 A).  To determine the effects of PI3K and EGFR dual inhibition on basal

signaling mechanisms, we looked at pathways downstream of EGFR signaling, specifically

focusing on MAPK, AKT, p70S6K and S6K signaling.  In the EGFR amplified cell lines MDA-

MB-468 and BT20, afatinib significantly inhibited basal MAPK signaling, but had no significant

effect on MAPK signaling in MDA-MB-231 cells (Figure 4B and C). Therefore, EGFR

amplified cells have basally activated MAPK signaling downstream of EGFR. Afatanib alone

also down regulated AKT signaling in the MDA-MB-468 cells but not in the BT20 or the MDA-

MB-231 cell lines (Figure 4B and D). The PIK3CA inhibitor alpelisib significantly down

regulated AKT activation in all three cell lines but did not affect pMAPK (Figure 4B and D,

Additional File 3 B). The combination of afatanib and alpelisib led to a greater reduction in in

AKT phosphorylation in the EGFR amplified cell lines (MDA-MB-468 and BT20) than either

drug alone (Figure 4B and D).  MDA-MB-231 cells, which do not have EGFR amplification or

PI3K alterations, did not have a significant reduction in AKT signaling when comparing the drug

combination to either drug alone, which can be seen with a darker exposure western blot (Figure

4D and Additional File 3 B).  All three cell lines had measurable p-p70S6K and an pS6, however

the drug combination only elicited a significant difference compared to DMSO control in BT20

and MDA-MB-468 (Figure 4B, E and F).  It is noteworthy that afatinib or alpelisib alone in

BT20 cells reduced pS6 in a statistically significant manner, however the combination of the two

drugs reduced pS6 to near undetectable levels, providing evidence that dual inhibition is the most

effective treatment at reducing S6 signaling (Figure 4B and 4F).  No significant effects on p-

p70S6K or pS6 were seen in MDA MB-231 treated with either drug alone or the combination.

To confirm that this downregulation of MAPK, AKT, p70S6K and S6K signaling was

due to a class effect and not these specific inhibitors, we also used EGFR TKI erlotinib and PI3K

inhibitor BKM120 and observed similar results (Additional File 3 C-F).  Overall, this shows that

the inhibition of pMAPK and a synergistic downregulation of pAKT, p-p70S6K and pS6K

signaling occurs in response to dual EGFR and PI3K inhibition in cells with EGFR amplification

and dysregulated PI3K in response to the dual inhibition of EGFR and PI3K.  The greatest effect

was observed in BT20 cells which have EGFR amplification and PIK3CA activating mutations.

It has been previously described that EGFR inhibition by cetuximab for 24 hours can

increase expression of HER3, leading to heterodimerization of EGFR and HER3, which in turn

overrides the EGFR inhibition by cetuximab (46).  In our work we did not see changes in HER3

or phospho-HER3 levels after 24 hours of exposure to the TKI erlotinib.  Further, the

phosphorylation of HER3 was inhibited by erlotinib in both of the EGFR amplified cell lines

(Additional File 4).  HER3 levels were very low, and phosphorylation was not detected in the

MDA-MB-231 cells.  It is likely that increased expression of HER3 cannot overcome inhibition

by the TKI as it does for cetuximab since the kinase activity of EGFR is necessary for signaling

of the EGFR/HER3 heterodimer (47).

**Dual EGFR and PI3K inhibition synergistically reduce cell viability in breast cancer cells**

**with EGFR amplification and PI3K alteration**

Next, we tested the effects of afatinib and alpelisib in combination on cell viability, to

determine whether this disruption in signaling had a functional effect.  In MDA-MB-468 cells,

which have EGFR amplification and PTEN deletion, each drug alone significantly inhibited

viability but the combination showed statistical significance when comparing the combination of

drugs to either drug alone, and exhibited synergy using all four calculations as determined by

SynergyFinder, with values over ten (Figure 5A) (40).  In BT20 cells, which have EGFR

amplification and PI3K mutations, the individual drugs again significantly inhibited viability and

the combination demonstrated the most significant effect of synergy with the drug combinations

completely inhibiting cell viability, as exemplified by the highest synergy score values (Figure

5B).  In MDA-MB-231 cells, which do not have EGFR amplification or PI3K alterations, there

was not significant viability inhibition by alpelisib alone, and there was only an inhibition of

viability to a high dose of afatinib (Figure 5C).  The combination of the two drugs did not

significantly affect viability, and synergy scores calculated resulted in values indicating an

additive effect of the two drugs together (score between -10 and 10) (Figure 5C).  Overall, EGFR

and PI3K inhibitors in combination exhibit a synergistic effect on cell viability in cell lines with

EGFR amplification and PI3K alterations, and only an additive effect in a cell line without

EGFR amplification or PI3K alterations.

We confirmed these effects to be a class effect and not a drug specific effect by testing

the combinations of erlotinib and BKM-120 (Additional File 5 A-C).  Erlotinib reduced the

viability of all three cell lines, however there is a much more significant decrease in cell viability

in the two EGFR amplified cells. When combined with BKM120, there is a further reduction of

cell viability in the EGFR amplified cells. Specifically, in BT20 cells, a combination of erlotinib

and BKM120 resulted in 90% reduction in cell viability. Overall, this data confirms that EGFR

amplified and aberrant PI3K cell lines MDA-MB-468 and BT20 have an increased reliance on

EGFR and PI3K signaling for cell viability regardless of EGFR TKI or PI3K inhibitor.

**Dual EGFR and PI3K inhibition induce cell cycle arrest and apoptosis in EGFR amplified**

**and PI3K altered breast cancer cells**

To evaluate the reason for the reduction in cell viability, we performed cell cycle analysis

and observed that there is an increase in cells in the G1 phase of the cell cycle when EGFR and

PI3K are inhibited by the combination of alpelisib and afatinib in both MB468 and BT20 cells

(Figure 6A and B). In addition to the G1 accumulation, the drug combination also exhibited

statistically significant reductions in both the S phase and G2/M cell populations. Further, cell

cycle was not significantly altered in MB231 cells which do not have EGFR amplification or

PI3K alteration (Figure 6C). EGFR and PI3K inhibition with erlotinib and BKM120

combination had similar effects on cell cycle, however there was also a statistically significant

increase in sub-G1 population with the drug combination in both MDA-MB-468 and BT20,

however there was no effect in control MDA-MB-231 (Additional File 5 D-F).

To determine whether EGFR and PI3K inhibitor combinations induced cell death and

apoptosis, we quantified dead cell percentages using propidium iodide (PI) staining of cells

treated with afatanib, alpelisib, and the combination in the presence or absence of Z-VAD-FMK

(ZVAD), which is a pan-caspase inhibitor. In control MDA-MB-231 cells, drugs alone or in

combination did not show an increase in dead cell percentage, consistent with our viability data

showing a lack of synergy for inducing cell death with these drugs in cells without EGFR

amplification and PI3K mutation (Figure 6D).  In contrast, in the BT20 cells with EGFR

amplification and PI3K mutation, the combination of afatanib and alpelisib resulted in a

dramatic, statistically significant increase in dead cells while each drug alone did not induce cell

death (Figure 6E).  Pre-treatment of the cells with ZVAD blocked this death, showing that the

cell death observed is caspase dependent.  We similarly confirmed that induction of cell death

was caspase-dependent in BT20 and MDA-MB-468 with erlotinib and BKM-120 being rescued

by ZVAD treatment (Additional File 6). It should be noted that erlotinib and BKM-120 alone did

increase caspase-dependent cell death, however the greatest and most significant increases in cell

death were when the drugs were combined. The MDA-MB-231 cells only showed a slight

increase in cell death with each erlotinib alone and no further increase in the effect of the

combination when compared to erlotinib alone (Additional File 6).

**Dual EGFR and PI3K inhibition is most effective at reducing tumor volume *in vivo***

In a mammary fat pad xenograft model of BT20 cells in nude mice, mice were

randomized 56 days post-implantation (indicated by R) when tumors reached 100 mm$^3$, and

treatment (indicated by Tx) was started 63 days post-implantation (Figure 7A).   Mice were

treated by oral gavage with control (0.5% carboxymethylcellulose), 20 mg/kg afatinib, 20 mg/kg

alpelisib, or a combination of afatinib and alpelisib for 5 days a week (Monday through Friday).

After the first four days of treatment, four mice from each group were sacrificed, and the tumors

were harvested for protein analysis.  Afatinib induced a statistically significant reduction in

pEGFR when compared to controls and alpelisib induced a statistically significant reduction in

pS6 compared to controls (Figure 7B).  Afatanib induced a non-significant reduction in pS6 and

alpelisib induced a non-significant reduction in pEGFR.  Of note, the combination of afatinib and

alpelisib did not further reduce downstream signaling as measured by pS6 compared to alpelisib

alone.

Single agents alone reduced tumor volume after 36 days of treatment, however neither

were statistically significant when compared to control due the slow and large variability in

growth of the BT20 cells *in vivo* (Figure 7A and C).  However, the reduction in tumor size by the

combination of both afatinib and alpelisib were statistically significant when compared to the

control.  Drug treatment was terminated after 36 days of treatment due to the requirement of

sacrificing control mice which had developed tumors larger than 20mm in any direction.

**Discussion**

The goal of this study was to determine the incidence of EGFR amplification in breast cancer

patients and to examine the benefit of molecularly targeted agents in models of EGFR

amplification.  Two cell lines with EGFR amplification and PI3K pathway mutations were

compared to a control cell line with wild-type EGFR and PI3K signaling.  EGFR and PI3K

inhibitor combinations were tested on EGFR/PI3K signaling, cancer cell viability, cancer cell

death, cell cycle, and mouse xenograft tumor growth.

We found EGFR amplification in approximately 1-2.5% of breast cancer patients with worse

outcomes compared to patients without EGFR amplification (Figure 1).  This EGFR

amplification was enriched in TNBC, HER2 amplified, and ER negative patients.  Importantly,

TNBC patients are difficult to treat due to lack of defined molecular targets. Determination of

targeted therapies aimed at a subset of breast cancer patients may provide therapeutic benefit in

those who previously may have had poor prognosis and could help shift the paradigm from

treatment of large populations with general chemotherapy, to treating smaller populations with a

focused and specific molecular inhibition strategy to provide clinical benefit.  This study

provides evidence that dual EGFR and PI3K inhibition in TNBC with EGFR amplification and

PI3K aberrations act synergistically in reducing cell viability, as well as inducing apoptosis *in*

*vitro*, and reducing tumor growth *in vivo*.  Further, we provide evidence of a patient population

which can be determined by molecular profiling and would benefit from treatment with this

therapy. For example, of the 272 EGFR amplified patients in the Caris dataset, only three of

them received the EGFR inhibitor Neratinib.  This study presents evidence for potential efficacy

of EGFR and PI3K inhibitors in these patients, and patients are currently not receiving this type

of therapy in the clinic.  This warrants clinical trial investigation into whether these patients

would benefit from this therapy.

It is interesting that a higher percentage of patients with EGFR amplification in TNBC

(2.45%) do not have an altered survival (Figure 1E and Additional File 1 A).  This likely reflects

that TNBC or Basal tumors generally have a poor prognosis.  None-the-less, our data using cell

models of EGFR amplification and PI3K pathway mutations suggest that these patients may

benefit from targeted EGFR and PI3K pathway inhibition.  Further, this study characterized a

model looking at TNBC, however considering HER2 enriched had a trend of worse OS, and

ER+HER2- patients had a statistically significantly worse OS, there are likely other breast cancer

patient populations with EGFR amplification and PI3K alteration who can benefit from this dual

therapy (Additional File 1).

The two cell lines with EGFR amplification and PI3K alterations (BT20 and MDA-MB-468)

exhibited significant inhibition of downstream signaling with the combination of EGFR

inhibitors and PI3K inhibitors, however the MDA-MB-231 cells without the alterations were less

affected by the drug combination (Figure 4, Additional File 3).  Further, the combination of

drugs had a synergistic effect on cell viability and cell cycle arrest of the cell lines with EGFR

amplification and PI3K alterations (Figure 5 and 6, Additional File 5).  In cells with EGFR

amplification and PI3K mutation, the combination of EGFR and PI3K inhibitors induced

apoptosis (Figure 6D and E, Additional File 6).  Lastly, in a xenograft mouse model, the

combination of EGFR and PI3K inhibitors reduced downstream signaling in the EGFR and PI3K

pathways and inhibited the tumor growth in a statistically significant manner when compared to

vehicle control.  Either drug alone did not reduce the tumor volume in a statistically significant

manner, indicating that the most effective treatment for these tumors with EGFR amplification

and PI3K mutation is the combination of both an EGFR inhibitor and PI3K inhibitor (Figure 7).

Between 20-50% of breast cancers exhibit PIK3CA mutations, with the most common

being hormone receptor positive or HER2 amplified breast cancer (48-53). It is noteworthy that

PIK3CA alterations found in this study (36-39%) in EGFR amplified patients is similar to the

percentage observed in HER2 amplified patients (30%) (51-53).  Combining HER2 targeted

therapy with PI3K inhibitors has been previously studied, and our study presents evidence that

there is a similar population of EGFR amplified patients who could benefit from combination

therapy (54, 55). Depending on the dataset, this study identified PI3K pathway alterations in 54-

71% of patients with EGFR amplification (Figure 3).

Alpelisib, which targets PIK3CA, has been FDA approved for treatment in hormone

receptor positive breast cancer with PIK3CA mutations in combination with endocrine therapy,

as it significantly improved progression-free survival (56-58).  In TNBC, alpelisib monotherapy

showed no overall response rate or clinical benefit rate in patients with PI3K alterations, and the

trial recruitment was ended due to lack of efficacy (53).  Therefore, alpelisib may benefit from

combination with other targeted therapies in breast cancer patients, especially in TNBC.

Capivasertib, which targets AKT, has been approved for treatment of metastatic breast cancers in

combination with ER targeted therapies, and improves patient survival (59, 60),

EGFR and PI3K inhibitor combination has been studied in NSLC,  head and neck

squamous cell carcinoma (HNSCC), glioblastoma, ovarian cancer and breast cancer, however

these studies did not monitor EGFR amplification or PI3K pathway mutation, and did not use

EGFR amplified and PI3K mutated cell lines as models (61-69). Further, a recent study looking

at PDX models with PI3K mutations found through a high throughput synergy model that

afatinib and alpelisib were synergistic, which confirms our data, however they also did not

monitor EGFR status (70).    Our study indicates that it is important to determine the subset of

patients who would benefit from this combination, by stratifying patients with EGFR

amplification and PI3K pathway alterations. We also do not rule out the possibility of

combining these drugs with HER2 targeted therapies, as we found that up to 5.8% of EGFR

amplification is found in breast cancer patients with HER2 amplification or HER2 enrichment

(Figure 1). In fact, there is a study showing synergy with BKM120 and the HER2 inhibitor

Trastuzumab, which further provides evidence that combining PI3K inhibitors with EGFR

family inhibitors may provide clinical benefit in patients with HER2 and EGFR amplification

and PI3K mutation (71).

We studied multiple EGFR (erlotinib or afatinib) and PI3K (BKM120 or alpelisib) inhibitors

(Figures 4, 5, 6, 7 and Additional Files 3, 5 and 6). It is important to note that afatinib inhibits

wild-type (WT) EGFR at the lowest IC50, and then at increasing IC50s it inhibits L858R EGFR,

HER2 and HER4 (72, 73), and BKM120 also inhibits mTOR, VPS34 and DNAPK (74).

Alpelisib is a more specific PI3K inhibitor, without observed off target effects. Since we tested

different EGFR and PI3K inhibitors and observed similar reductions in downstream signaling,

cell viability, cell cycle, and apoptosis when the drugs were combined, it is likely that the

observed effects were due to a class effect of the drugs. Taken together with the *in vivo* data in

this study, drugs from these classes warrant investigation in combination in breast cancer patients

who exhibit EGFR amplification and PI3K alterations.

Dual EGFR and PI3K inhibition *in vitro* exhibited significant synergy in this study,

however the *in vivo* data was more modest. The modest effects are likely due to the short half-

life of alpelisib (approximately 1.5 hours in mice), whereas afatinib has a longer half-life

(approximately 21 hours in mice) (75, 76).  Compared to the *in vitro* work where cells were

treated continuously for 48-72 hours with the drugs, this could account for the difference in

observed results when comparing *in vitro* and *in vivo* results in this study.  The half-life of

afatinib in human patients is approximately 30-40 hours and alpelisib is approximately 7.5 hours,

suggesting with the daily dosing used, there would be better steady state levels than in the mouse

(77, 78).

**Conclusions**

Approximately 1-2.5% of breast cancer patients have EGFR amplification, with a worse

prognosis.  Up to 71% of the EGFR amplified population exhibits co-incidence of PI3K pathway

activating alterations.  Dual targeting of these pathways in cells which express EGFR

amplification and PI3K alteration with chemical inhibitors demonstrated synergy *in vitro*, and

their combination reduced tumor growth in a xenograft model more than either drug alone when

compared to control.  Although rare, our data suggests that breast cancers that exhibit EGFR

amplification and PI3K pathway alterations are a poor prognosis and should be approached with

targeted therapy similar to those with HER2 amplification. Taken together, this study

demonstrated that the combination of EGFR and PI3K pathway inhibitors should be tested in

breast cancer patients with EGFR amplification and PI3K pathway alteration.  This will shift the

paradigm for the treatment regimen in this patient population, especially triple negative breast

cancer patients with fewer targeted therapy options.

**Figures**

**Figure 1. Incidence of EGFR amplification in breast cancer.** A) Breast cancer subtype

analysis of patients in TCGA dataset with EGFR amplification. B) Similar analysis of

METABRIC dataset, and C) MSKCC dataset. D) Combination of A-C datasets to analyze

overall survival (OS) of patients comparing EGFR amplification (red line) to patients without

EGFR amplification (blue line). E) Breast cancer subtype analysis of patients in Caris dataset

with EGFR amplification. F) OS of patients with EGFR amplification (red line) compared to

patients without EGFR amplification (blue line).

**Figure 2. EGFR protein expression and gene amplification, and PI3K mutations in MDA-**

**MB-231, MDA-MB-468 and BT20 cells.** A) Western blot showing protein expression of

EGFR, PIK3CA, pTEN and HSC70 (loading control) in MDA-MB-231, MDA-MB-468 and

BT20 cell lines. B) FISH analysis showing EGFR (red) and CEP7 (aqua, control) in MDA-MB-

231, MDA-MB-468 and BT20 cell lines. C) DNA sequencing for PIK3CA mutation status in

BT20, MDA-MB-468 and MDA-MB-231 cell lines.

**Figure 3. PI3K pathway mutations in EGFR amplified breast cancer.** A) Datasets from

TCGA, METABRIC and MSKCC showing alterations in PI3K pathway genes in 75 patients

with EGFR amplification. B) Dataset from Caris showing alterations in PI3K pathway genes in

272 breast cancers with EGFR amplification.

**Figure 4. EGFR/PI3K dual inhibition significantly reduces downstream signaling in**

**EGFR amplified and PI3K altered breast cancer.** A) BT-20 and MDA-MB-468 cells were

pretreated for 24 hours with 0, 10 or 100 nM afatinib, and then starved in serum-free RPM1

media for 3 hours prior to stimulation with or without 25 ng/mL EGF for fifteen minutes. Cells

were lysed and immunoblotted for pEGFR (Y1068), EGFR, pMAPK (T202/Y204), ERK2 and

HSC70 (loading control). B) MDA-MB-231, MDA-MB-468 and BT20 cells were treated with

100 nM alpelisib (Alp), 1 µM afatinib (Afat) or a combination of both for 24 hours, and were

then lysed and immunoblotted for pAKT (S473), AKT, pMAPK (T202/Y204), ERK2, p-p70S6K

(T389), p70S6K, pS6 (S235/236), S6 and HSC70 (loading control).  In at least three independent

experiments, the band density of the phosphorylated protein relative to total protein was

averaged ±SEM for C) pMAPK/ERK2, D) pAKT/AKT, E) p-p70S6K/p70S6K and F) pS6/S6.

Two-way ANOVA was performed, where ns indicates not statistically significant, * indicates

$p < 0.05$, ** indicates $p < 0.01$, *** indicates $p < 0.001$ and **** indicates $p < 0.0001$.

**Figure 5. Dual EGFR/PI3K inhibition synergistically reduce cell viability in TNBC with**

**EGFR amplification and PI3K alteration.**  A) MDA-MB-468, B) BT20 and C) MDA-MB-231

cells were treated with the indicated doses for 48 hours, and then viability was determined by

Promega CellTiter-Glo 2.0.  Data represents the average ±SEM of at least three independent

experiments, and statistical analysis was performed as Two-Way ANOVA, and the p value

indicates interaction, where ns indicates not statistically significant, ** indicates $p < 0.01$, ***

indicates $p < 0.001$ and **** indicates $p < 0.0001$.  The averages were also subjected to four

different synergy score analyses through synergyfinder.org, where a score of $> +10$ indicates

synergy, -10 to +10 indicates additive effects, and $< -10$ indicates antagonism.

**Figure 6. EGFR/PI3K inhibition induce cell cycle arrest and apoptosis in EGFR amplified**

**and PI3K altered TNBC.**  A) MDA-MB-468 were treated with DMSO (control), 1 µM afatinib,

10 µM alpelisib, or the combination of both B) BT20 were treated with DMSO (control), 100

nM afatinib, 1 µM alpelisib, or the combination of both and C) MDA-MB-231 were treated with

DMSO (control), 1 µM afatinib, 10 µM alpelisib, or the combination of both for 48 hours, and

then were harvested, washed, fixed and stained with propidium iodide before being analyzed by

flow cytometry and FlowJo for cell cycle changes. Data represents the average ±SEM of at least

three independent experiments, and Two-Way ANOVA statistical analysis was performed where

* indicates $p<0.05$, ** indicates $p<0.01$, *** indicates $p<0.001$ and **** indicates $p<0.0001$. D)

MDA-MB-231 and E) BT20 cells were pretreated with 20 µM Z-VAD-FMK (ZVAD) for one

hour, and then were treated with doses DMSO, 1 µM alpelisib, 1 µM afatinib, or the combination

of alpelisib and afatinib with continued ZVAD treatment in 1% FBS RPM1 media containing

propidium iodide for 72 hours. Imaging of the cells was performed by BioTek Cytation 1, and

dead cell percentage was calculated by dividing PI positive cells by the total brightfield cell

count and multiplying by 100. Data represents the average ±SEM of at least three independent

experiments. Statistical analysis was performed by Student's t-test, with *** indicating $p<0.001$.

**Figure 7. Dual EGFR/PI3K inhibition is most effective at reducing tumor volume *in vivo*.**

A) BT20 cells were injected subcutaneously into the mammary fat pad of nude mice. 56 days

post implantation, when the average size was 100 mm$^3$, mice were randomized (R) to 15 mice

into four groups and treated starting on day 63 post implantation (Tx) with control (0.5%

carboxymethylcellulose), 20 mg/kg afatinib, 20 mg/kg alpelisib, or the combination of both

afatinib and alpelisib Monday through Friday for 36 days. Statistical analysis was performed by

Student's t-test, and * indicates $p<0.05$ when comparing afatinib and alpelisib to control on day

94 and day 98. B) After four days of treatment, four mice from each group were euthanized and

the protein from the tumors was collected and immunoblotted for pEGFR (Y1068), EGFR, pS6

(S235/236), S6, and HSC70 (loading control). Data represents the average of four mice from

each group ±SEM and statistical analysis was performed by Student's t-test, where ns indicates

not statistically significant, * indicates $p<0.05$ and ** indicates $p<0.01$. C) Bar graph of data

from A on day 98 (36 days of drug treatment).  Statistical analysis was performed by Student's t-

test, where * indicates p<0.05 when compared to control.  No other comparisons were

statistically significant.

**List of Abbreviations**

EGFR- Epidermal Growth Factor Receptor

PI3K- Phosphoinositide 3-kinase

TNBC- Triple Negative Breast Cancer

FISH- fluorescent in situ hybridization

PI- propidium iodide

ER- Estrogen Receptor

HER2- Human epidermal growth factor receptor 2

mTOR- mechanistic target of rapamycin

AKT- Protein Kinase B

EGF- epidermal growth factor

MAPK- mitogen activated protein kinases

TKI- tyrosine kinase inhibitor

P70S6K- ribosomal protein S6 kinase beta 1

SEM- standard error of the mean

TCGA- The Cancer Genome Atlas Program

MSKCC- Memorial Sloan Kettering Cancer Center

IDC- invasive ductal carcinoma

ILC- Invasive lobular carcinoma

OS- overall survival

PTEN- phosphatase and tensin homolog

DMSO- dimethyl sulfoxide

NSCLC- non-small cell lung cancer

HNSCC- head and neck squamous cell carcinoma

PDX- patient derived xenograft

WT- wild-type

DNAPK- DNA-dependent protein kinase

**Declarations**

**Ethics approval and consent to participate:** All animal aspects of this study were approved by

the NCI Animal Care and Use Committee (IACUC number WMB-004).  All studies in the Caris

dataset were conducted in accordance with guidelines of the Declaration of Helsinki, Belmont

report, and U.S. Common rule, and in keeping with 45 CFR 46.101(b)(4), was performed

utilizing retrospective, deidentified clinical data. Per WCG IRB, it was considered institutional

review board (IRB) exempt with waiver of patient consent. 15,247 breast cancer samples

underwent molecular testing at Caris Life Sciences (Phoenix, AZ, USA), a CLIA/CAP-certified

laboratory. No specific inclusion criteria or exclusion criteria were applied aside from the

requirement for successful sequencing results.

**Consent for publication:** not applicable

**Availability of data and materials:** The MSKCC, TCGA and Metabric datasets analyzed in this

study are available on cBioPortal.org (35-37).  The Caris deidentified sequencing data are owned

by Caris Life Sciences and cannot be publicly shared due to the data usage agreement in place.

These data will be made available to researchers for replication and verification purposes

through our letter of intent process, which are generally fulfilled within 6 months. For more

information on how to access this data, please contact Joanne Xiu at jxiu@carisls.com.  All other

data generated in this study are included in this published article and its supplementary

information.

**Competing interests:** The authors declare that they have no competing interests.

**Funding:** This research was supported by the Intramural Research Program of the National

Cancer Institute, National Institutes of Health (ZIA BC 010977)

**Authors' contributions:** DJW designed and performed *in vitro* and *in vivo* laboratory

experiments, helped with project conceptualization, data analysis and interpretation, contributed

to the bioinformatic analysis of the patient data from cBioPortal and Caris Life Sciences and

wrote the manuscript. DV contributed to experimental design and performed *in vitro* laboratory

experiments and contributed to data analysis and interpretation. YA contributed to experimental

design, *in vitro* laboratory experiments, and data analysis. SD, SW, ML and GS performed the

bioinformatic analysis and interpretation of the CARIS Life Sciences datasets. DW, DW and

KH performed and analyzed the FISH experiments. SL contributed to the project

conceptualization, bioinformatic analysis of the cBioPortal dataset and interpretation of Caris

Life Sciences datasets, data analysis and interpretation, and manuscript editing. All authors read

and edited the manuscript and approved the final manuscript.

**Acknowledgements:** We thank all members of the Lipkowitz lab in the Women's Malignancies

Branch, Center for Cancer Research, National Cancer Institute for their support and review of

this manuscript. We would like to thank Dr. Thomas Ried for his assistance with the planning of

the FISH experiment. We would like to acknowledge the CCR Genomics Core for their

assistance with sequencing, and the veterinarians and animal care staff at NIH for their assistance

with animal care, drug delivery and tumor measurement. The content is solely the responsibility

of the authors and does not necessarily represent the official views of the National Institutes of

Health.

**References**

1.      Normanno N, De Luca A, Bianco C, Strizzi L, Mancino M, Maiello MR, et al. Epidermal growth factor receptor (EGFR) signaling in cancer. Gene. 2006;366(1):2-16.

2.      Uribe ML, Marrocco I, Yarden Y. EGFR in Cancer: Signaling Mechanisms, Drugs, and Acquired Resistance. Cancers (Basel). 2021;13(11).

3.      Voldborg BR, Damstrup L, Spang-Thomsen M, Poulsen HS. Epidermal growth factor receptor (EGFR) and EGFR mutations, function and possible role in clinical trials. Ann Oncol. 1997;8(12):1197-206.

4.      Zhang X, Gureasko J, Shen K, Cole PA, Kuriyan J. An allosteric mechanism for activation of the kinase domain of epidermal growth factor receptor. Cell. 2006;125(6):1137-49.

5.      Levantini E, Maroni G, Del Re M, Tenen DG. EGFR signaling pathway as therapeutic target in human cancers. Seminars in Cancer Biology. 2022;85:253-75.

6.      Tebbutt N, Pedersen MW, Johns TG. Targeting the ERBB family in cancer: couples therapy. Nat Rev Cancer. 2013;13(9):663-73.

7.      Zeillinger R, Kury F, Czerwenka K, Kubista E, Sliutz G, Knogler W, et al. HER-2 amplification, steroid receptors and epidermal growth factor receptor in primary breast cancer. Oncogene. 1989;4(1):109-14.

8.      Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A, McGuire WL. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. Science. 1987;235(4785):177-82.

9.      Sjögren S, Inganäs M, Lindgren A, Holmberg L, Bergh J. Prognostic and predictive value of c-erbB-2 overexpression in primary breast cancer, alone and in combination with other prognostic markers. J Clin Oncol. 1998;16(2):462-9.

10.     Swain SM, Shastry M, Hamilton E. Targeting HER2-positive breast cancer: advances and future directions. Nat Rev Drug Discov. 2023;22(2):101-26.

11.     Bose R, Kavuri SM, Searleman AC, Shen W, Shen D, Koboldt DC, et al. Activating HER2 Mutations in HER2 Gene Amplification Negative Breast Cancer. Cancer Discovery. 2013;3(2):224-37.

12.     Mitsudomi T, Yatabe Y. Epidermal growth factor receptor in relation to tumor development: EGFR gene and cancer. Febs j. 2010;277(2):301-8.

13.     da Cunha Santos G, Shepherd FA, Tsao MS. EGFR mutations and lung cancer. Annu Rev Pathol. 2011;6:49-69.

14.     Nedaeinia R, Avan A, Manian M, Salehi R, Ghayour-Mobarhan M. EGFR as a potential target for the treatment of pancreatic cancer: dilemma and controversies. Curr Drug Targets. 2014;15(14):1293-301.

15.     Tathineni P, Joshi N, Jelinek MJ. Current State and Future Directions of EGFR-Directed Therapy in Head and Neck Cancer. Curr Treat Options Oncol. 2023;24(6):680-92.

16.     Rakha EA, Elsheikh SE, Aleskandarany MA, Habashi HO, Green AR, Powe DG, et al. Triple-negative breast cancer: distinguishing between basal and nonbasal subtypes. Clin Cancer Res. 2009;15(7):2302-10.

17.     Klijn JGM, Berns PMJJ, Schmitz PIM, Foekens JA. The Clinical Significance of Epidermal Growth Factor Receptor (EGF-R) in Human Breast Cancer: A Review on 5232 Patients*. Endocrine Reviews. 1992;13(1):3-17.

18.     Masuda H, Zhang D, Bartholomeusz C, Doihara H, Hortobagyi GN, Ueno NT. Role of
epidermal growth factor receptor in breast cancer. Breast Cancer Research and Treatment.
2012;136(2):331-45.
19.     Zhao Y, Ma J, Fan Y, Wang Z, Tian R, Ji W, et al. TGF-β transactivates EGFR and
facilitates breast cancer migration and invasion through canonical Smad3 and ERK/Sp1
signaling pathways. Mol Oncol. 2018;12(3):305-21.
20.     Burness ML, Grushko TA, Olopade OI. Epidermal Growth Factor Receptor in Triple-
Negative and Basal-Like Breast Cancer: Promising Clinical Target or Only a Marker? The
Cancer Journal. 2010;16(1):23-32.
21.     Rakha EA, El-Sayed ME, Green AR, Lee AH, Robertson JF, Ellis IO. Prognostic
markers in triple-negative breast cancer. Cancer. 2007;109(1):25-32.
22.     Li X, Zhao L, Chen C, Nie J, Jiao B. Can EGFR be a therapeutic target in breast cancer?
Biochimica et Biophysica Acta (BBA) - Reviews on Cancer. 2022;1877(5):188789.
23.     DiGiovanna MP, Stern DF, Edgerton SM, Whalen SG, Moore D, 2nd, Thor AD.
Relationship of epidermal growth factor receptor expression to ErbB-2 signaling activity and
prognosis in breast cancer patients. J Clin Oncol. 2005;23(6):1152-60.
24.     Sainsbury JR, Farndon JR, Needham GK, Malcolm AJ, Harris AL. Epidermal-growth-
factor receptor status as predictor of early recurrence of and death from breast cancer. Lancet.
1987;1(8547):1398-402.
25.     Spyratos F, Delarue JC, Andrieu C, Lidereau R, Champème MH, Hacène K, Brunet M.
Epidermal growth factor receptors and prognosis in primary breast cancer. Breast Cancer Res
Treat. 1990;17(2):83-9.
26.     Al-Kuraya K, Schraml P, Torhorst J, Tapia C, Zaharieva B, Novotny H, et al. Prognostic
relevance of gene amplifications and coamplifications in breast cancer. Cancer Res.
2004;64(23):8534-40.
27.     Bhargava R, Gerald WL, Li AR, Pan Q, Lal P, Ladanyi M, Chen B. EGFR gene
amplification in breast cancer: correlation with epidermal growth factor receptor mRNA and
protein expression and HER-2 status and absence of EGFR-activating mutations. Mod Pathol.
2005;18(8):1027-33.
28.     Ro J, North SM, Gallick GE, Hortobagyi GN, Gutterman JU, Blick M. Amplified and
overexpressed epidermal growth factor receptor gene in uncultured primary human breast
carcinoma. Cancer Res. 1988;48(1):161-4.
29.     Reis-Filho JS, Milanezi F, Carvalho S, Simpson PT, Steele D, Savage K, et al.
Metaplastic breast carcinomas exhibit EGFR, but not HER2, gene amplification and
overexpression: immunohistochemical and chromogenic in situ hybridization analysis. Breast
Cancer Res. 2005;7(6):R1028-35.
30.     Reis-Filho JS, Pinheiro C, Lambros MB, Milanezi F, Carvalho S, Savage K, et al. EGFR
amplification and lack of activating mutations in metaplastic breast carcinomas. J Pathol.
2006;209(4):445-53.
31.     O'DONOVAN N, CROWN J. EGFR and HER-2 Antagonists in Breast Cancer.
Anticancer Research. 2007;27(3A):1285-94.
32.     Tan AR, Yang X, Hewitt SM, Berman A, Lepper ER, Sparreboom A, et al. Evaluation of
biologic end points and pharmacokinetics in patients with metastatic breast cancer after treatment
with erlotinib, an epidermal growth factor receptor tyrosine kinase inhibitor. J Clin Oncol.
2004;22(15):3080-90.

33.    Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.
34.    Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-
aware quantification of transcript expression. Nat Methods. 2017;14(4):417-9.
35.    Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer
genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer
Discov. 2012;2(5):401-4.
36.    Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, et al. Integrative
analysis of complex cancer genomics and clinical profiles using the cBioPortal. Sci Signal.
2013;6(269):pl1.
37.    de Bruijn I, Kundra R, Mastrogiacomo B, Tran TN, Sikina L, Mazor T, et al. Analysis
and Visualization of Longitudinal Genomic and Clinical Data from the AACR Project GENIE
Biopharma Collaborative in cBioPortal. Cancer Res. 2023;83(23):3861-7.
38.    Mermel CH, Schumacher SE, Hill B, Meyerson ML, Beroukhim R, Getz G. GISTIC2.0
facilitates sensitive and confident localization of the targets of focal somatic copy-number
alteration in human cancers. Genome Biol. 2011;12(4):R41.
39.    Liyasova MS, Ma K, Voeller D, Ryan PE, Chen J, Klevit RE, Lipkowitz S. Cbl interacts
with multiple E2s in vitro and in cells. PLoS One. 2019;14(5):e0216967.
40.    Zheng S, Wang W, Aldahdooh J, Malyutina A, Shadbahr T, Tanoli Z, et al.
SynergyFinder Plus: Toward Better Interpretation and Annotation of Drug Combination
Screening Datasets. Genomics, Proteomics & Bioinformatics. 2022;20(3):587-96.
41.    Prat A, Parker JS, Fan C, Perou CM. PAM50 assay and the three-gene model for
identifying the major and clinically relevant molecular subtypes of breast cancer. Breast Cancer
Res Treat. 2012;135(1):301-6.
42.    Zhang J, Gao X, Schmit F, Adelmant G, Eck MJ, Marto JA, et al. CRKL Mediates
p110β-Dependent PI3K Signaling in PTEN-Deficient Cancer Cells. Cell Reports.
2017;20(3):549-57.
43.    Kao J, Salari K, Bocanegra M, Choi YL, Girard L, Gandhi J, et al. Molecular profiling of
breast cancer cell lines defines relevant tumor models and provides a resource for cancer gene
discovery. PloS one. 2009;4(7):e6146.
44.    Stemke-Hale K, Gonzalez-Angulo AM, Lluch A, Neve RM, Kuo WL, Davies M, et al.
An integrative genomic and proteomic analysis of PIK3CA, PTEN, and AKT mutations in breast
cancer. Cancer Res. 2008;68(15):6084-91.
45.    Vasan N, Razavi P, Johnson JL, Shao H, Shah H, Antoine A, et al. Double PIK3CA
mutations in cis increase oncogenicity and sensitivity to PI3Kα inhibitors. Science.
2019;366(6466):714-23.
46.    Tao JJ, Castel P, Radosevic-Robin N, Elkabets M, Auricchio N, Aceto N, et al.
Antagonism of EGFR and HER3 enhances the response to inhibitors of the PI3K-Akt pathway in
triple-negative breast cancer. Sci Signal. 2014;7(318):ra29.
47.    Alroy I, Yarden Y. The ErbB signaling network in embryogenesis and oncogenesis:
signal diversification through combinatorial ligand-receptor interactions. FEBS Lett.
1997;410(1):83-6.
48.    Armaghani AJ, Han HS. Alpelisib in the Treatment of Breast Cancer: A Short Review on
the Emerging Clinical Data. Breast Cancer (Dove Med Press). 2020;12:251-8.

49. Bailey MH, Tokheim C, Porta-Pardo E, Sengupta S, Bertrand D, Weerasinghe A, et al.
Comprehensive Characterization of Cancer Driver Genes and Mutations. Cell. 2018;173(2):371-
85.e18.
50. Pereira B, Chin SF, Rueda OM, Vollan HK, Provenzano E, Bardwell HA, et al. The
somatic mutation profiles of 2,433 breast cancers refines their genomic and transcriptomic
landscapes. Nat Commun. 2016;7:11479.
51. Razavi P, Chang MT, Xu G, Bandlamudi C, Ross DS, Vasan N, et al. The Genomic
Landscape of Endocrine-Resistant Advanced Breast Cancers. Cancer Cell. 2018;34(3):427-
38.e6.
52. Comprehensive molecular portraits of human breast tumours. Nature.
2012;490(7418):61-70.
53. Savas P, Lo LL, Luen SJ, Blackley EF, Callahan J, Moodie K, et al. Alpelisib
Monotherapy for PI3K-Altered, Pretreated Advanced Breast Cancer: A Phase II Study. Cancer
Discov. 2022;12(9):2058-73.
54. Rodon J, Pérez-Fidalgo A, Krop IE, Burris H, Guerrero-Zotano A, Britten CD, et al.
Phase 1/1b dose escalation and expansion study of BEZ235, a dual PI3K/mTOR inhibitor, in
patients with advanced solid tumors including patients with advanced breast cancer. Cancer
Chemother Pharmacol. 2018;82(2):285-98.
55. Loibl S, de la Pena L, Nekljudova V, Zardavas D, Michiels S, Denkert C, et al.
Neoadjuvant buparlisib plus trastuzumab and paclitaxel for women with HER2+ primary breast
cancer: A randomised, double-blind, placebo-controlled phase II trial (NeoPHOEBE). Eur J
Cancer. 2017;85:133-45.
56. André F, Ciruelos E, Rubovszky G, Campone M, Loibl S, Rugo HS, et al. Alpelisib for
PIK3CA-Mutated, Hormone Receptor-Positive Advanced Breast Cancer. N Engl J Med.
2019;380(20):1929-40.
57. André F, Ciruelos EM, Juric D, Loibl S, Campone M, Mayer IA, et al. Alpelisib plus
fulvestrant for PIK3CA-mutated, hormone receptor-positive, human epidermal growth factor
receptor-2-negative advanced breast cancer: final overall survival results from SOLAR-1. Ann
Oncol. 2021;32(2):208-17.
58. Copur MS. Alpelisib to treat breast cancer. Drugs Today (Barc). 2020;56(6):357-63.
59. Howell SJ, Casbard A, Carucci M, Ingarfield K, Butler R, Morgan S, et al. Fulvestrant
plus capivasertib versus placebo after relapse or progression on an aromatase inhibitor in
metastatic, oestrogen receptor-positive, HER2-negative breast cancer (FAKTION): overall
survival, updated progression-free survival, and expanded biomarker analysis from a
randomised, phase 2 trial. Lancet Oncol. 2022;23(7):851-64.
60. Turner NC, Oliveira M, Howell SJ, Dalenc F, Cortes J, Gomez Moreno HL, et al.
Capivasertib in Hormone Receptor-Positive Advanced Breast Cancer. N Engl J Med.
2023;388(22):2058-70.
61. Yang Z, Liao J, Schumaker L, Carter-Cooper B, Lapidus RG, Fan X, et al.
Simultaneously targeting ErbB family kinases and PI3K in HPV-positive head and neck
squamous cell carcinoma. Oral Oncol. 2022;131:105939.
62. Guo T, Wu C, Zhang J, Yu J, Li G, Jiang H, et al. Dual blockade of EGFR and PI3K
signaling pathways offers a therapeutic strategy for glioblastoma. Cell Commun Signal.
2023;21(1):363.

63. Sidorov M, Dighe P, Woo RWL, Rodriguez-Brotons A, Chen M, Ice RJ, et al. Dual Targeting of EGFR and MTOR Pathways Inhibits Glioblastoma Growth by Modulating the Tumor Microenvironment. Cells. 2023;12(4).

64. Glaysher S, Bolton LM, Johnson P, Atkey N, Dyson M, Torrance C, Cree IA. Targeting EGFR and PI3K pathways in ovarian cancer. Br J Cancer. 2013;109(7):1786-94.

65. Li P, Zhang Q, Torossian A, Li ZB, Xu WC, Lu B, Fu S. Simultaneous inhibition of EGFR and PI3K enhances radiosensitivity in human breast cancer. Int J Radiat Oncol Biol Phys. 2012;83(3):e391-7.

66. El Guerrab A, Bamdad M, Bignon YJ, Penault-Llorca F, Aubel C. Co-targeting EGFR and mTOR with gefitinib and everolimus in triple-negative breast cancer cells. Scientific reports. 2020;10(1):6367.

67. Glaysher S, Bolton LM, Johnson P, Torrance C, Cree IA. Activity of EGFR, mTOR and PI3K inhibitors in an isogenic breast cell line model. BMC Res Notes. 2014;7:397.

68. Deng M, Wang J, Chen Y, Zhang L, Liu D. Combination of SF1126 and gefitinib induces apoptosis of triple-negative breast cancer cells through the PI3K/AKT-mTOR pathway. Anticancer Drugs. 2015;26(4):422-7.

69. She QB, Gruvberger-Saal SK, Maurer M, Chen Y, Jumppanen M, Su T, et al. Integrated molecular pathway analysis informs a synergistic combination therapy targeting PTEN/PI3K and EGFR pathways for basal-like breast cancer. BMC Cancer. 2016;16:587.

70. Boyd DC, Zboril EK, Olex AL, Leftwich TJ, Hairr NS, Byers HA, et al. Discovering Synergistic Compounds with BYL-719 in PI3K Overactivated Basal-like PDXs. Cancers. 2023;15(5):1582.

71. Yu F, Zhao J, Hu Y, Zhou Y, Guo R, Bai J, et al. The combination of NVP-BKM120 with trastuzumab or RAD001 synergistically inhibits the growth of breast cancer stem cells in vivo. Oncol Rep. 2016;36(1):356-64.

72. Li D, Ambrogio L, Shimamura T, Kubo S, Takahashi M, Chirieac LR, et al. BIBW2992, an irreversible EGFR/HER2 inhibitor highly effective in preclinical lung cancer models. Oncogene. 2008;27(34):4702-11.

73. Ferrarotto R, Gold KA. Afatinib in the treatment of head and neck squamous cell carcinoma. Expert Opin Investig Drugs. 2014;23(1):135-43.

74. Burger MT, Pecchi S, Wagman A, Ni Z-J, Knapp M, Hendrickson T, et al. Identification of NVP-BKM120 as a Potent, Selective, Orally Bioavailable Class I PI3 Kinase Inhibitor for Treating Cancer. ACS Medicinal Chemistry Letters. 2011;2(10):774-9.

75. Hedges CP, Boix J, Jaiswal JK, Shetty B, Shepherd PR, Merry TL. Efficacy of Providing the PI3K p110α Inhibitor BYL719 (Alpelisib) to Middle-Aged Mice in Their Diet. Biomolecules. 2021;11(2).

76. Wang J, Su G, Yin X, Luo J, Gu R, Wang S, et al. Non-small cell lung cancer-targeted, redox-sensitive lipid-polymer hybrid nanoparticles for the delivery of a second-generation irreversible epidermal growth factor inhibitor-Afatinib: In vitro and in vivo evaluation. Biomed Pharmacother. 2019;120:109493.

77. Murakami H, Tamura T, Takahashi T, Nokihara H, Naito T, Nakamura Y, et al. Phase I study of continuous afatinib (BIBW 2992) in patients with advanced non-small cell lung cancer after prior chemotherapy/erlotinib/gefitinib (LUX-Lung 4). Cancer Chemother Pharmacol. 2012;69(4):891-9.

78. Juric D, Rodon J, Tabernero J, Janku F, Burris HA, Schellens JHM, et al. Phosphatidylinositol 3-Kinase α-Selective Inhibition With Alpelisib (BYL719) in PIK3CA-

Altered Solid Tumors: Results From the First-in-Human Study. J Clin Oncol. 2018;36(13):1291-
9.

**Additional Files**

**Additional File 1**

File Format: .tiff

Title of Data: Overall survival of patients with EGFR amplification by subtype.

Description of Data: Using the Caris dataset, overall survival of breast cancer patients with

EGFR amplification (red) compared to no amplification (blue) in A) Triple negative breast

cancer, B) Basal, C) HER2 enriched, or D) ER+ HER2-.

**Additional File 2**

File Format: .tiff

Title of Data:  Oncoplot for PI3K pathway alterations in EGFR amplified breast cancer.

Description of Data: Patient data from cbioportal.org plotting varying mutations or

amplifications by gene in the PI3K pathway.

**Additional File 3**

File Format: .tiff

Title of Data:  EGFR/PI3K dual inhibition significantly reduce downstream signaling in EGFR

amplified and PI3K altered TNBC.

Description of Data: A) MDA-MB-468 and BT20 cells were starved in serum-free RPM1 media

for 1 hour with 0, 1 or 10 µM erlotinib, and then stimulated with or without 25 ng/mL EGF for

10 minutes.  Cells were lysed and immunoblotted for pEGFR (Y1068) and EGFR (loading

control). B) Darker exposure for MDA-MB-231 panel from Figure 4B which was used for

quantification.  C) MDA-MB-231, MDA-MB-468 and BT20 cells were treated with 10 µM

erlotinib, 1 µM BKM120 or a combination of both for 24 hours, and were then lysed and

immunoblotted for pAKT (S473), AKT, pMAPK (T202/Y204), ERK2, p-p70S6K (T389),

p70S6K and HSC70 (loading control).  In at least three independent experiments, the band

density of the phosphorylated protein relative to total protein was averaged ±SEM for D)

pMAPK/ERK2, E) pAKT/AKT and F) p-p70S6K/p70S6K.  Student's t-test was performed,

where ns indicates not statistically significant, * indicates $p<0.05$, ** indicates $p<0.01$, ***

indicates $p<0.001$ and **** indicates $p<0.0001$.

**Additional File 4**

File Format: .tiff

Title of Data: Erlotinib treatment does not activate pHER3.

Description of Data: MDA-MB-231, MDA-MB-468 and BT20 cells were treated with 10 µM

erlotinib, 1 µM BKM120 or the combination of both for 24 hours before the cells were lysed and

immunoblotted for pHER3 (Y1289) and HER3 (loading control).

**Additional File 5**

File Format: .tiff

Title of Data: EGFR/PI3K inhibition reduces viability, induces cell cycle arrest in TNBC with

EGFR amplification, PI3K alteration.

Description of Data: A) MDA-MB-231, B) MDA-MB-468 and C) BT20 cells were treated with

the indicated doses for 48 hours, and then viability was determined by Promega CellTiter-Glo

2.0.  Data represents the average ±SEM of at least three independent experiments, and statistical

analysis was performed as Two-Way ANOVA, and the p value indicates interaction, where ns

indicates not statistically significant, * indicates $p<0.05$, *** indicates $p<0.001$ and ****

indicates $p<0.0001$. D) MDA-MB-231, E) MDA-MB-468 and F) BT20 cells were treated with

DMSO control, 1 µM erlotinib or 10 µM BKM120 or the combination of erlotinib and BKM120

for 48 hours, and then were harvested, washed, fixed and stained with propidium iodide before

being analyzed by flow cytometry and FlowJo for cell cycle changes.  Data represents the

average ±SEM  of at least three independent experiments, and Two-Way ANOVA statistical

analysis was performed where * indicates $p<0.05$, ** indicates $p<0.01$ and *** indicates

$p<0.001$.

**Additional File 6**

File Format: .tiff

Title of Data:  Dual EGFR/PI3K inhibition induce apoptosis in EGFR amplified and PI3K

altered TNBC.

Description of Data:  A) MDA-MB-231, B) MDA-MB-468 and C) BT20 cells were pretreated

with 20 µM Z-VAD-FMK (ZVAD) for one hour, and then were treated with DMSO, 10 µM

erlotinib, 1 µM BKM120 or the combination of both, with continued ZVAD treatment for 48

hours.  Dead cell percentage was calculated by using CytoTox-Glo by Promega.  Data represents

the average ±SEM of at least three independent experiments.  Statistical analysis was performed

by One-way ANOVA, with ns indicating not statistically significant, * indicating $p<0.05$, **

indicating $p<0.01$ and *** indicating $p<0.001$.

1011

1012

**A**

| Breast Cancer Subtype | Samples # | EGFR Amplification (%) |
|---|---|---|
| All Patients | 816 | 14 (1.7%) |
|  |  |  |
| **Histology** |  |  |
| IDC | 489 | 10 (2.0%) |
| Mixed IDC/ILC | 88 | 0 (0%) |
| Lobular | 127 | 0 (0%) |
| Other Histologies | 112 | 4 (3.6%) |
|  |  |  |
| **ER and HER2 Status** |  |  |
| ER+ | 600 | 5 (0.0%) |
| ER+/HER2- | 326 | 3 (0.9%) |
| ER Negative | 175 | 8 (4.6%) |
| HER2 Positive | 121 | 3 (2.5%) |
| ER+/HER2+ | 89 | 1 (1.1%) |
| ER-/HER2+ | 31 | 2 (6.5%) |
| TNBC | 90 | 6 (6.7%) |
|  |  |  |
| **Molecular Subtype** |  |  |
| Total Profiled | 587 | 10 (1.7%) |
| Luminal A (IDC) | 201 | 2 (1.0%) |
| Luminal A (ILC) | 106 | 0 (0%) |
| Luminal B | 122 | 2 (1.6%) |
| HER2 Enriched | 51 | 2 (3.9%) |
| Basal | 107 | 4 (4.3%) |

Table title: TCGA (Cell 2015)

**B**

| Breast Cancer Subtype | Samples # | EGFR Amplification (%) |
|---|---|---|
| All Patients | 2173 | 52 (2.4%) |
|  |  |  |
| **Histology** |  |  |
| IDC | 1660 | 48 (2.9%) |
| Mixed IDC/ILC | 218 | 2 (0.9%) |
| Lobular | 172 | 1 (0.6%) |
| Other Histologies | 123 | 1 (0.8%) |
|  |  |  |
| **ER and HER2 Status** |  |  |
| ER+ | 1617 | 23 (1.4%) |
| ER+/HER2- | 1398 | 19 (1.4%) |
| ER Negative | 523 | 28 (5.4%) |
| HER2 Positive | 247 | 9 (3.6%) |
| ER+/HER2+ | 108 | 1 (0.9%) |
| ER-/HER2+ | 139 | 8 (5.8%) |
| TNBC | 335 | 19 (5.7%) |
|  |  |  |
| **Molecular Subtype** |  |  |
| Total Profiled | 1980 | 47 (2.4%) |
| Luminal A- | 700 | 5 (0.7%) |
| Luminal B | 475 | 7 (1.5%) |
| HER2 Enriched | 224 | 12 (5.4%) |
| Basal | 209 | 17 (8.1) |
| Claudin Low | 218 | 3 (1.4%) |
| Normal | 148 | 3 (2.0%) |

Table title: Metabric (Nature 2012 and Nat Comm 2016)

**C**

| Breast Cancer Subtype | Samples # | EGFR Amplification (%) |
|---|---|---|
| All Patients | 1918 | 23 (1.2%) |
|  |  |  |
| **Histology** |  |  |
| IDC | 1473 | 21 (1.4%) |
| Mixed IDC/ILC | 89 | 0 (0%) |
| Lobular | 299 | 0 (0%) |
| Other Histologies | 57 | 2 (3.5%) |
|  |  |  |
| **ER and HER2 Status** |  |  |
| ER+ | 1512 | 13 (0.9%) |
| ER+/HER2- | 1357 | 11 (0.8%) |
| ER Negative | 327 | 9 (2.8%) |
| HER2 Positive | 240 | 3 (1.3%) |
| ER+/HER2+ | 76 | 1 (1.3%) |
| ER-/HER2+ | 79 | 1 (1.3%) |
| TNBC | 251 | 8 (3.2%) |

Table title: MSKCC Profile (Cell 2018)

**D**

EGFR WT
EGFR Amplification
p=0.0269

| Number at risk (n) | | | | | | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| EGFR Amplified | 87 | 71 | 50 | 46 | 36 | 31 | 24 | 19 | 14 | 12 | 10 | 9 | 8 | 7 | 5 | 3 | 1 | 9 | 5 | 2 | 1 |
| EGFR Wild Type | 4461 | 3455 | 2787 | 2301 | 1924 | 1575 | 1279 | 1079 | 832 | 666 | 482 | 357 | 251 | 152 | 66 | 31 | 17 | 9 | 5 | 2 | 1 |

EGFR WT median survival = 164.60 months
EGFR amplification survival = 111.10 months

**E**

| Breast Cancer Subtype | Samples # | EGFR Amplification (%) |
|---|---|---|
| All patients | 26360 | 272 (1.03%) |
|  |  |  |
| **Histology** |  |  |
| IDC | 9789 | 126 (1.29%) |
| ILC | 1677 | 2 (0.12%) |
| Other/unclear | 14894 | 144 (0.97%) |
|  |  |  |
| **ER and HER2 Status** |  |  |
| ER+ | 15320 | 61 (0.40%) |
| ER+/HER2- | 11892 | 49 (0.41%) |
| ER Negative | 7818 | 181 (2.32%) |
| HER2 Positive | 1531 | 11 (0.72%) |
| ER+/HER2+ | 830 | 2 (0.24%) |
| ER-/HER2+ | 684 | 9 (1.32%) |
| TNBC | 6125 | 150 (2.45%) |
|  |  |  |
| **Molecular Subtypes** |  |  |
| Luminal A | 2204 | 3 (0.14%) |
| Luminal B | 6430 | 9 (0.14%) |
| HER2 Enriched | 2672 | 50 (1.87%) |
| Basal | 3869 | 90 (2.33%) |
| Normal | 140 | 2 (1.43%) |

**F**

EGFR WT
EGFR Amplification
p<0.0001

| | | | | | | |
|---|---|---|---|---|---|---|
| EGFR Wild Type | 21983 | 1271 | 148 | 18 | 4 | 1 |
| EGFR Amplified | 267 | 8 | 1 | 0 | 0 | 0 |

EGFR Wild Type median survival = 32.999 months
EGFR amplification median survival = 21.648 months

**A**

| Gene | Potential PI3K Pathway Activating Alterations | Potential PI3KPathway Activating Alterations (%) |
|---|---|---|
| PIK3CA | 27 | 36 |
| PIK3R1 | 3 | 4 |
| PIK3R2 | 3 | 4 |
| PIK3R3 | 3 | 4 |
| PTEN | 4 | 5.3 |
| AKT1 | 4 | 5.3 |
| AKT2 | 5 | 6.7 |
| AKT3 | 13 | 17.3 |
| RPS6KB1 | 6 | 8 |
| RPS6KB2 | 3 | 4 |
| RPS6 | 3 | 4 |
| **Patients with at least one PI3K activation** | **53** | **70.7** |

**B**

| Gene | Potential PI3KPathway Activating Alterations (%) |
|---|---|
| PIK3CA | 38.97 |
| PIK3R1 | 6.04 |
| PIK3R2 | 2.21 |
| PIK3R3 | Not Available |
| PTEN | 3.79 |
| AKT1 | 0.37 |
| AKT2 | 2.93 |
| AKT3 | 0 |
| RPS6KB1 | Not Available |
| RPS6KB2 | Not Available |
| RPS6 | Not Available |
| **Any PI3K Activation** | **54.31** |