1

1   EGFR amplification and PI3K pathway mutations identify a subset of breast cancers that

2   synergistically respond to EGFR and PI3K inhibition

David J. Wisniewski[1], Donna Voeller[1], Yonit A. Addissie[1], Sachin Kumar Deshmukh[2], Sharon

Wu[2], Maryam B. Lustberg[3], Darawalee Wangsa[4], Danny Wangsa[4], Kerstin Heselmeyer-

Haddad[4], Yoshimi Endo Greer[1], George W. Sledge Jr.[2] and *Stanley Lipkowitz[1]

[1]Women's Malignancies Branch, Center for Cancer Research, National Cancer Institute,

Bethesda MD

[2]Caris Life Sciences, Phoenix AZ

[3]Yale Cancer Center, New Haven CT

[4]Genetics Branch, Center for Cancer Research, National Cancer Institute, Bethesda MD

*Corresponding Author (lipkowis@navmed.nci.nih.gov)

Competing Interests Statement: The authors declare that they have no competing interests.

Keywords: epidermal growth factor receptor amplification, PI3Kinase pathway mutations,

triple negative breast cancer

Abstract

EGFR family receptor tyrosine kinase signaling is commonly dysregulated in cancer by

amplification or activating mutations. Although studies have investigated dual EGFR/PI3K

inhibition in breast cancer, they have not determined biomarkers which predict success. We

present evidence of a patient subset with EGFR amplification and PI3Kinase pathway mutations

in breast cancer which can be synergistically targeted by dual EGFR/PI3K inhibition. This study

identified that EGFR amplification occurs in approximately 1-5% of breast cancer patients with

shorter overall survival compared to unamplified patients. Up to 71% of EGFR amplified tumors

have activating mutations in the PI3K pathway. Dual EGFR/PI3K inhibition more dramatically

reduced mTOR and AKT signaling in BT20 and MDA-MB-468 cells which both have EGFR

amplification and PI3K pathway activating mutations, compared to control cells. Dual inhibition

synergistically reduced cell viability and increased apoptosis in MDA-MB-468 and BT20

compared to control. Single agent therapy in a BT20 xenograft model reduced tumor volume,

however only the combination statistically significantly reduced tumor volume compared to

control. We conclude that EGFR amplification with co-incident PI3K pathway mutations are

driver mutations in a subset of breast cancers and present a subgroup of breast cancers that are

more likely to respond to dual targeted therapy.

Introduction

The ErbB family is comprised of four receptor tyrosine kinases, Epidermal Growth Factor Receptor (EGFR), HER2 (ErbB2), HER3 (ErbB3) and HER4 (ErbB4). Upon ligand binding, the receptors hetero or homodimerize allowing them to activate downstream signaling, including phosphoinositide 3-kinases (PI3Ks) or mitogen-activated protein kinases (MAPKs), resulting in increased proliferation, inhibition of apoptosis, migration, invasion, and other pro-tumor signaling (1-4). Mutation or overexpression of these family members are common drivers of various cancer types (5). For example, HER2 amplification is a driver in 12-20% of breast cancers, correlates with poor survival, and has been effectively targeted to improve outcomes in patients with HER2 amplified breast cancer (6-9).

EGFR can be aberrantly activated through mutations or amplification/overexpression, which drives pro-tumor effects in cancers such as non-small cell lung cancer, pancreatic cancer and head and neck cancer, leading to approval of EGFR targeted therapies including tyrosine kinase inhibitors (TKI's) gefitinib, erlotinib, afatinib, dacomitinib, osimertinib, and vandetanib, and the monoclonal antibodies cetuximab, panitumumab and necitumumab (1, 2, 10-13). EGFR is often expressed in basal triple negative breast cancer (TNBC), and is associated with tumor size, poor differentiation, worse relapse-free and overall survival, and shorter metastasis free survival (14, 15). Depending on the study, EGFR gene amplification has been observed in 0.8-14% of breast cancer patients (16-19). Mutation of EGFR in breast cancer can range from approximately 2-11% (20). Relatively small studies looking specifically at metaplastic breast carcinomas have observed amplification of EGFR in 33-37% of patients (21, 22). Despite these data, clinical trials investigating EGFR inhibitors have not shown great promise in breast cancer,

with studies finding partial response rates of 0-6% (16, 20, 23-25). These studies were typically

in heavily pretreated patients, and patients were not stratified by EGFR amplification status.

Although previous studies in various *in vitro* cancer models have evaluated EGFR and

PI3K inhibitor combination, these studies did not investigate the role of EGFR amplification in

addition to PI3K mutation status as biomarkers for drug synergy (26-37). Further, they did not

characterize a potential patient population which would most benefit from this treatment

regimen. Here, we evaluated the frequency and subtype distribution of EGFR amplification, the

clinical outcomes of patients with EGFR amplified tumors, the frequency of coincident

activating mutations in the PI3K pathway and studied the effects of dual inhibition of EGFR and

the PI3K pathways in TNBC cell line models in vitro and in vivo. Our data provides evidence

that breast cancers with EGFR amplification and PI3K pathway activating mutations may be

more likely to respond to therapies targeting these pathways.

Methods

Materials

Z-VAD-FMK (S7023), Erlotinib (S7786), BKM120 (S2247), Afatinib (S1011), Alpelisib

(S2814) and sodium carboxymethyl cellulose (S6703) were purchased from Selleck Chemicals.

Fetal bovine serum (FBS) (631106) was obtained from Takara Bio. 4-20% precast

polyacrylamide gels (5671094) were purchased from Bio-Rad. Immobilon PVDF transfer

membrane (IPVH00010), mini protease inhibitor cocktail (11836153001) and propidium iodide

(P4864) were purchased from Millipore-Sigma. PBS (21-031-CV), EGF (354042) and Phenol

Red-Free LDEV-Free Matrigel (356237) was purchased from Corning. Sodium orthovanadate

(P0758S) was from Fisher Chemicals. NP40 Cell Lysis Buffer (FNN0021), Penicillin-

Streptomycin (15140-122), tissue extraction reagent (FNN0071) and RPMI 1640 medium

(11875-093) was obtained from ThermoFisher Scientific.

Antibodies

Anti-phospho-HER3 Y1289 (4791S), anti-HER3 (12708S), anti-EGFR (2232), anti-PTEN

(9559S), anti-phospho-EGFR Y1068 (2234), anti-phospho-AKT S473 (9271S), anti-AKT

(9272S), anti-phospho-MAPK T202/Y204 (9101), anti-phospho-P70S6K T389 (9234S), anti-

P70S6K (9202S), anti-phospho-S6 S235/236 (4856S), anti-S6 (2217S) and anti-PI3K (4249S)

were purchased from Cell Signaling Technology. anti-HSC70 (sc-7298) and anti-ERK2 (D-2,

sc-1647) antibodies were purchased from Santa Cruz Biotechnology. Goat Anti-Mouse IgG

HRP conjugate (172-1011) and Goat Anti-Rabbit IgG HRP conjugate (1721019) were purchased

from BioRad.

Cell Lines

MDA-MB-468, BT20, MDA-MB-231 were obtained from ATCC, maintained in RPMI 1640

medium with 10% FBS and 1% penicillin/streptomycin, and grown at 37^{o}C, 5% CO_2. These cell

lines were recently authenticated and tested for mycoplasma contamination.

Caris data

Next-generation sequencing (NGS)

NGS was performed on genomic DNA using the NextSeq (592-whole gene targets; Agilent

Technologies, Santa Clara, CA, USA) or NovaSeq 6000 (>700 genes at high coverage/read-

depth and >20 000 genes at lower depth; Agilent Technologies, Santa Clara, CA, USA)

platforms (Illumina, Inc., San Diego, CA, USA) as described previously (38). The copy number

alteration (CNA) of each exon is determined by normalizing the sequencing depth of each exon

divided by the average sequencing depth of the sample and comparing it to the pre-calibrated

mean of normalized values in the training data (The mean values are re-calibrated every 60 days

with up to 10,000 samples). A gene was defined as amplified by having segment level of copies

$\geq$4.5.

Survival analysis

Caris CodeAI™ clinico-genomic database containing insurance claims data were used to

calculate real-world overall survival (OS) from tissue collection to last contact. Kaplan-Meier

curves were generated to calculate molecularly defined patient cohorts as previously described

(39, 40).

cBioPortal data

Datasets from TCGA (Cell 2015), METABRIC (Nature 2012 and Nature Communications 2016)

and MSKCC (Cell 2018) were searched for EGFR amplification, as well as mutations in

PIK3CA, PIK3R1, PIK3R2, PIK3R3, PTEN, AKT1, AKT2, AKT3, RPS6KB1, RPS6KB2 and

RPS6 in cBioPortal (cbioportal.org) (41-43). Amplification was defined by GISTIC2.0 score = 2

(CN $\geq$4-5) (44).

Western Blot

Cell lysates were collected, prepared and immunoblotted as previously described, and repeated at

least 3 times (45). Densitometric analysis of band intensities was calculated using ImageJ and

data are presented as average $\pm$ SEM.

Fluorescence *in situ* Hybridization (FISH)

To prepare metaphase chromosomes, cells were treated with 0.02 mg/ml Colcemid (Invitrogen,

Grand Island, NY) for one hour, suspended in a hypotonic solution for 20 minutes, fixed using a

methanol/acetic acid mixture (1:3 ratio) and dropped onto slides in a cytogenetic drying system

(Thermotron, Holland, MI). Dual color FISH was performed with custom probes for EGFR and

centromere 7 from Cytotest (Rockville, MD), labeled with Dyomics dyes (Jena, Germany).

Slides were pepsin treated, washed in 1x PBS, dehydrated in an ethanol series, and left to air-dry.

Co-denaturation was performed at 72°C for 2 minutes, followed by overnight hybridization at

37°C. The slides were detected in 2× SSC/0.3% Nonidet P-40 for 2 minutes at 48°C, and 2×

SSC/0.1% Nonidet P-40 for 1 minute at room temperature and were mounted with DAPI

antifade solution (Vector Laboratories, Burlingame, CA). Images were captured using a Leica

DM-RXA fluorescence microscope (Leica, Wetzlar, Germany) equipped with a 40 X objective

and custom optical filters (Chroma, Bellows Falls, VT).

Sanger Sequencing

148 Sanger DNA sequencing of genomic DNA from the MDA-MB-231, MDA-MB-468 and BT20

149 cell lines for PIK3CA mutations was performed using the Big Dye Terminator v.1.1 Cycle

150 Sequencing Kit (Applied Biosystems, CA) according to the manufacturer's specifications and

151 run on an ABI 3130xl or 3730 Genetic Analyzer (Applied Biosystems, CA) at the CCR

152 Genomics Core at the National Cancer Institute.

Cell Death

Cell death was determined by CytoTox-Glo (G9291) as directed by Promega or by propidium

iodide uptake. Propidium iodide (PI) staining, imaging and quantification was performed using

the BioTek Cytation 1 Imaging Reader from Agilent (Santa Clara, CA) and Gen5 Image Prime

3.14 software (Agilent). For CytoTox Glo, cells were pretreated with 20 µM ZVAD for 1 hour,

followed by co-treatment with DMSO, 10 µM erlotinib, 1 µM BKM120 or the combination for

48 hours. Dead cell percentage was then calculated according to the manufacturer instructions.

For PI, 5000 cells were plated in a 96-well plate and treated the next day with experimental

conditions in 1% FBS RPMI, along with PI (1:3000 dilution). Dead cell percentage was

calculated by dividing dead cell number (PI stained) by brightfield cell count number and

multiplying by 100 and normalized relative to the Day 0 dead cell percentage. In experiments

using ZVAD-FMK, cells were pretreated +/- 20 µM ZVAD-FMK for 1 hour, followed by co-

treatment with experimental conditions.

Cell Viability

170   10 000 cells were plated in a 96-well white walled plate (3610, Corning) and treated the

171   following day with experimental conditions for 48 hours in 1% FBS RPMI. Viability was

172   determined by CellTiter-Glo 2.0 (Catalog #G9242) as directed by Promega.

Cell cycle

174   Cells were plated in 6-well plates ($3x10^5$ cells/well) and treated with EGFR and/or PI3K

175   inhibitors for 48 hours. DNA content was measured by flow cytometry as previously described,

176   using a BD LSRFortessa™ (BD Biosciences) flow cytometer and analyzed using FlowJo®

177   software (FlowJo LLC) (46, 47).

Animal Studies

Sixty female athymic nude mice between 5-6 weeks in age obtained from Charles River

Laboratories were housed and treated according to approved NCI-ACUC guidelines and in

accordance with an approved NCI Animal Care and Use Committee (IACUC number WMB-

004). 5 million BT20 cells suspended in matrigel (1:1 with PBS) were injected subcutaneously

into the mammary fat pad with a 25G needle (Terumo Medical Care Solutions; SS-01T2516).

When tumors were 100 mm^3 the mice were randomized into four groups of 15. The mice were

randomized so each group included similar average tumor size, with similar standard deviation.

0.5% sodium carboxymethyl cellulose (CMC) in water (w/v) was used as control, and 20 mg/kg

afatinib or 20 mg/kg alpelisib were suspended in 0.5% CMC. Control or 20 mg/kg alpelisib, 20

mg/kg afatinib, or the combination of both was given by oral gavage (1 inch, 22G needle) once a

day for five days each week for the duration of the study. Animals were not included in

treatment or analysis if the tumors did not reach 100 mm^3 at the time of randomization.   After

the first four days of treatment, four mice per group were sacrificed, tumors were excised and

proteins were harvested in Tissue Extraction Reagent I (Invitrogen-FNN0071) according to the

manufacturers protocol. Tumor caliper measurements and body weight were measured twice a

week for the duration of the study, as required by humane endpoints for all mice. Investigators

were blinded to the tumor measurements. Tumor volumes were calculated according to the

formula V= ½ (length×width^2). When maximum tumor size for each animal was reached (20

mm in any direction) the mice were humanely euthanized.

Statistical Analysis

Student's *t*-test with 2 tailed comparisons assuming equal variance, and one-way or two-way

ANOVA were performed as indicated. P-values of ≤0.05 were considered significant. Synergy

scores were calculated using SynergyFinder.org (48). *In vitro* experiments consisted of

triplicates per treatment group, with at least three experimental repeats performed (with specific

repeat numbers indicated in figure legends). *In vivo* experiments contained n=15 per treatment

group.

Results

Incidence of EGFR amplification in breast cancer

EGFR amplification was observed in 1.2-2.4% of cases in the TCGA, METABRIC and

MSKCC databases in cBioPortal (41-43), depending on the dataset (Table 1). The TCGA and

METABRIC databases contain mostly early-stage primary breast cancers while the MSKCC

dataset is approximately 50% metastatic/recurrent breast cancers. EGFR amplification is more

common in invasive ductal carcinomas (IDC) (1.4-2.9%) compared to invasive lobular

carcinomas (ILC) (0-0.6%). Further, the highest rate of amplification was in TNBC (3.2-6.7%),

ER- (2.8-5.4%) and ER-/HER2+ tumors (1.3-6.5%), whereas EGFR amplification was relatively

less common in ER+ breast cancers (0.8-1.4%). Using PAM50 classification, EGFR

amplification is the highest in basal (3.7-8.1%) and HER2 enriched (3.9-5.4%) molecular

subtypes, compared to luminal A or luminal B subtypes (0.7% and 1.5-1.6%, respectively) (49).

We combined the TCGA, METABRIC and MSKCC datasets due the small number of EGFR

amplified cases in each and found that there is a statistically significant lower OS in patients with

amplified EGFR, with a difference of 53.5 months (111.4 vs 164.6 months) (Fig. 1A).

To confirm these data in a larger dataset with clinical annotation, we evaluated the

incidence of EGFR amplification ($\geq$4.5 CN) in patients with breast cancer profiled by Caris Life

Sciences (Caris), where the majority of patients profiled had metastatic breast cancer (Table 1).

EGFR amplification was found in 5.1% of the breast cancers profiled. As in the other datasets,

EGFR amplification was more prevalent in IDC (5.2%) compared to ILC (2.1%). EGFR

amplification was more prevalent in TNBC (8.4%), and in the molecular HER2 enriched (7.7 %),

basal (6.8%) and normal (3.8%) subtypes. The EGFR amplification seen in TNBC in this dataset

(8.4%) is similar to previously published data, which found 8.9% EGFR amplification in TNBC

(50). OS was statistically significantly shorter by 9.1 months in patients with EGFR

amplification compared to those with unamplified EGFR (23.9 vs 33 months; p<0.00001: Fig.

1B). The shorter survival in the Caris dataset is likely due to the patients being metastatic, while

the majority of the patients in the dataset for Figure 1A were early-stage patients. This data

shows that there is a subset of breast cancer patients with EGFR amplification that is most

frequent in TNBC and ER-/HER2 amplified tumors and is associated with a poor prognosis.

Since EGFR amplification is enriched in patients with TNBC or Basal cancers, we

investigated the differences in overall survival in patients with EGFR amplification depending on

subtype using the Caris dataset (Fig. S1). There was a borderline worse overall survival in

patients with EGFR amplification (difference of 1.7 months; p=0.09) in TNBC patients (Fig.

S1A). Patients with molecularly defined Basal tumors with EGFR amplification also exhibited a

borderline worse overall survival (difference of 1.3 months; p=0.06) (Fig. S1B). There was no

significant difference in the survival of HER2 enriched patients with EGFR amplification (Fig.

S1C). In contrast, overall survival was statistically significantly worse in patients with

ER+/HER2- tumors that had EGFR amplification compared to those with WT EGFR (difference

9.6 months; p<0.001) (Fig. S1D).

***In vitro* modeling of EGFR amplification and PI3K alterations**

To evaluate the role of EGFR amplification in cell models, we used two cell lines with

amplified EGFR (MDA-MB-468 and BT20) and one cell line without amplified EGFR (MDA-

MB-231). MDA-MB-468 and BT20 have elevated EGFR protein levels as compared to MDA-

MB-231 and FISH confirmed the amplification of EGFR in both MDA-MB-468 and BT20, and

the lack of EGFR amplification in MDA-MB-231 (Fig. 2A and B). MDA-MB-468 has loss of

PTEN protein, which supports previously published studies showing loss of PTEN expression in

these cells (Fig. 2A) (51, 52). As previously reported, the BT20 TNBC cell line has two

activating mutations in PIKCA (P539R and H1047R), which previously have been shown to be

in cis (Fig. 2C) (52-54). MDA-MB-468 and MDA-MB-231 have wildtype PIK3CA. Therefore,

MDA-MB-468 and BT20 were the two cell lines with EGFR amplification and PI3Kinase

pathway activating mutations, and MDA-MB-231 was the control cell line without amplified

EGFR or PI3Kinase pathway mutations.

PI3K pathway mutations in EGFR amplified breast cancer

Considering the co-incidence of mutations in PI3K signaling in these EGFR amplified

cell lines, using cBioPortal we investigated the incidence in patients. In 75 patients with EGFR

amplification from the TCGA, METABRIC and MSKCC cohorts, 53 patients (approximately

71%) with EGFR amplified tumors have at least one genetic alteration which can potentially

activate PI3K signaling (Table 2 and Fig S2). 36% have an activating mutation in PIK3CA,

5.3% have a loss of PTEN, 12% have mutations in PIK3R1, R2 or R3, and 29.3% have

amplification of AKT1, 2, or 3. There are 16% of patients who have amplifications downstream

of mTOR (either in ribosomal protein S6 kinase B1 or B2, or ribosomal protein S6). Of note,

multiple tumors had two or more alterations in this pathway (Fig. S2).

In the Caris dataset, 50.8% of EGFR amplified tumors have mutation or amplification in

the PI3K pathway (including PIK3CA, pTEN, PIK3R1, PIK3R2, or AKT 1, 2 or 3) (Table 2).

The largest percentage of alterations are in PIK3CA (36.5%). The Caris dataset does not include

alterations in the genes downstream of mTOR, contributing to the difference comparing the sums

of Caris and the cBioPortal data. This data confirms that many tumors with EGFR amplification

are likely to have co-incident aberrant PI3K signaling.

274 **EGFR and PI3K dual inhibition significantly reduce downstream signaling in EGFR**

275 **amplified and PI3K altered breast cancer**

276   Epidermal growth factor (EGF) and tyrosine kinase inhibitor (TKI) treatment showed that

277 EGFR phosphorylation was decreased upon afatinib or erlotinib treatment in both MDA-MB-468

278 and BT20 cell lines, indicating that the TKI inhibited EGFR signaling upon EGF stimulation

279 (Fig. 3A, Fig. S3A). To determine the effects of EGFR and PIK3CA dual inhibition on steady

280 state signaling, we looked at MAPK, AKT, P70S6K and S6K activation. In the EGFR amplified

281 cell lines MDA-MB-468 and BT20, afatinib significantly inhibited basal MAPK activation, but

282 had no significant effect on MAPK signaling in MDA-MB-231 cells (Fig. 3B and C). Therefore,

283 EGFR amplified cells have activated MAPK signaling downstream of EGFR. Afatanib alone

284 also down regulated AKT signaling in the MDA-MB-468 cells but not in the BT20 or the MDA-

285 MB-231 cell lines (Fig. 3B and D, Fig. S3B). The PIK3CA inhibitor alpelisib significantly down

286 regulated AKT activation in all three cell lines but did not affect pMAPK (Fig. 3B and D, Fig.

287 S3B). Combination of afatanib and alpelisib led to a greater reduction in in AKT

288 phosphorylation in the EGFR amplified cell lines (MDA-MB-468 and BT20) than either drug

289 alone (Fig. 3B and D). MDA-MB-231 cells did not have a significant reduction in AKT

290 signaling when comparing the drug combination to either drug alone (Fig. 3D and Fig. S3B). It

291 should be noted that phosphorylated AKT levels were undetectable in MB231 at the exposure

292 shown in Figure 3B, and the quantification of phosphorylated AKT levels in MB231 used in 3D

293 were from immunoblots with longer exposures, such as the representative blot seen in

294 Supplemental Figure S3B. All three cell lines had measurable p-P70S6K and pS6, however the

295 drug combination only elicited a significant decrease in BT20 and MDA-MB-468 (Fig. 3B, E

296 and F). In BT20 cells, afatinib or alpelisib alone reduced pS6 in a statistically significant

manner, however the combination of the two drugs reduced pS6 to near undetectable levels (Fig.

3B and F). No significant effects on p-P70S6K or pS6 were seen in MDA MB-231. Similar

results were observed using the EGFR TKI erlotinib and PI3K inhibitor BKM120 (Fig. S3B-F).

Overall, this shows that the inhibition of pMAPK and a synergistic downregulation of pAKT, p-

P70S6K and pS6K signaling occurs in response to dual EGFR and PI3K inhibition in cells with

EGFR amplification and dysregulated PI3K. The greatest effect was observed in BT20 cells

which have EGFR amplification and PIK3CA activating mutations.

It has been previously described that EGFR inhibition by cetuximab can increase

expression of HER3, leading to heterodimerization of EGFR and HER3, in turn overriding the

EGFR inhibition by cetuximab (55). We did not observe changes in HER3 or pHER3 levels

after exposure to erlotinib. Further, the phosphorylation of HER3 was inhibited by erlotinib in

the EGFR amplified cell lines (Fig. S4). HER3 levels were very low, and pHER3

phosphorylation was not detected in the MDA-MB-231 cells. It is likely that increased

expression of HER3 cannot overcome inhibition by the TKI as it does for cetuximab since the

kinase activity of EGFR is necessary for signaling of the EGFR/HER3 heterodimer (56).

Dual EGFR and PI3K inhibition synergistically reduce cell viability in breast cancer cells

with EGFR amplification and PI3K alteration

We next tested the effects of afatinib and alpelisib alone and in combination on cell

viability. In MDA-MB-231 there was not significant viability inhibition by alpelisib alone, there

was only an inhibition of viability at a high dose of afatinib, the combination did not

significantly affect viability and synergy scores indicated an additive effect (Fig. 4A). In

contrast, in BT20 and MDA-MB-468 cells, each drug alone significantly inhibited viability, but

the combination showed statistical significance when comparing the combination of drugs to

either drug alone, and exhibited synergy using all four calculations as determined by

SynergyFinder (Fig. 4B and C) (48). Similar results were seen with erlotinib and BKM-120 (Fig.

S5A-C). Overall, EGFR and PI3K inhibitors in combination exhibit a synergistic effect on cell

viability in cell lines with EGFR amplification and PI3K alterations, and only an additive effect

in a cell line without EGFR amplification or PI3K alterations.

Dual EGFR and PI3K inhibition induce cell cycle arrest and apoptosis in EGFR amplified

and PI3K altered breast cancer cells

Cell cycle was not significantly altered in MB231 cells (Fig. 5A). In both MB468 and

BT20, cell cycle analysis showed an increase in cells in the G1 phase of the cell cycle and

statistically significant reductions in both the S phase and G2/M cell populations when EGFR

and PI3K are inhibited by the combination of alpelisib and afatinib (Fig. 5B and C). Similar

effects on the cell cycle were seen with erlotinib and BKM120, however there was also a

statistically significant increase in sub-G1 population with the drug combination in both MDA-

MB-468 and BT20 but not in MDA-MB-231 (Fig. S5 D-F).

We quantified dead cell percentages using propidium iodide (PI) staining of cells treated

with afatanib, alpelisib, and the combination in the presence or absence of the pan-caspase

inhibitor Z-VAD-FMK (ZVAD). The combination of afatanib and alpelisib in both MB468 and

BT20 cells resulted in a dramatic, statistically significant increase in dead cells which was

rescued by ZVAD treatment, while the drugs alone or in combination did not increase the

percentage of dead cells in MDA-MB-231 (Fig. 5D-F). Similarly, dual inhibition with erlotinib

and BKM-120 increased the dead cells in BT20 and MDA-MB-468, which was rescued by

ZVAD treatment (Fig. S6). Erlotinib and BKM-120 alone did increase caspase-dependent cell

death in MDA-MB-468 and BT20 cells, but to a lesser extent (Fig. S6). MDA-MB-231 cells only

showed a slight increase in cell death with erlotinib alone and no further increase in the

percentage dead cells by the combination (Fig. S6A).

Dual EGFR and PI3K inhibition reduce tumor volume *in vivo*

346   We conducted an orthotopic mammary fat pad xenograft model of BT20 cells in nude

mice treated with control, afatanib, alpelisib or the combination (Fig 6A). Afatinib induced a

statistically significant reduction in pEGFR when compared to controls and alpelisib induced a

statistically significant reduction in pS6 compared to controls (Fig. 6B). Afatanib non-

significantly reduced pS6 and alpelisib non-significantly reduced pEGFR. The combination of

afatinib and alpelisib did not further reduce downstream signaling as measured by pS6 compared

to alpelisib alone.

353   Single agents alone reduced tumor volume after 36 days of treatment, however neither

were statistically significant when compared to control due the slow and large variability in

growth in vivo (Fig. 6A and C). The reduction in tumor size by the combination of both afatinib

and alpelisib was statistically significant when compared to the control.

Discussion

This studies goal was to determine the incidence of EGFR amplification in breast cancer

patients and to examine the benefit of molecularly targeted agents in models of EGFR

amplification. EGFR amplification occurs in approximately 1-5% of breast cancer patients with

worse outcomes compared to patients without EGFR amplification (Table 1 and Fig. 1). This

EGFR amplification was enriched in TNBC, HER2 amplified, and ER negative patients.

Importantly, TNBC patients are difficult to treat due to lack of defined molecular targets.

Determination of targeted therapies aimed at a subset of breast cancer patients may provide

therapeutic benefit in those who previously may have had poor prognosis and could help shift the

paradigm to treating identifiable populations with a focused and specific molecular inhibition

strategy.

Dual EGFR and PI3K inhibition in TNBC with EGFR amplification and PI3K aberrations act

synergistically in reducing cell viability, as well as inducing apoptosis in vitro, and reducing

tumor growth in vivo. Our data identifies a patient population which can be determined by

molecular profiling and would benefit from treatment with this therapy. This study presents

evidence for potential efficacy of EGFR and PI3K inhibitors in these patients and warrants

clinical trial investigation into whether these patients would benefit from this therapy.

TNBC patients with EGFR amplification have a borderline worse survival compared to

TNBC without EGFR amplification (Table 1 and Fig. S1A). Our data suggest that these patients

may benefit from targeted EGFR and PI3K pathway inhibition. Further, this study characterized

a model looking at TNBC, however ER+HER2- patients had a statistically significantly worse

OS, therefore there are likely other breast cancer patient populations with EGFR amplification

and PI3K alteration who might benefit from this dual therapy (Fig. S1).

The two cell lines with EGFR amplification and PI3K alterations (BT20 and MDA-MB-468)

exhibited significant signaling pathway and biological effects of EGFR inhibitors and PI3K

inhibitors while the MDA-MB-231 cells without the alterations were less affected by the drug

combination (Fig. 3 and 4, Fig. S3). In a xenograft mouse model, the combination of EGFR and

PI3K inhibitors reduced downstream signaling in the EGFR and PI3K pathways and inhibited

the tumor growth in a statistically significant manner when compared to vehicle control (Fig. 6).

Dual EGFR and PI3K inhibition in vitro exhibited significant synergy in this study, however

the in vivo data was more modest. The modest effects are likely due to the short half-life of

alpelisib (approximately 1.5 hours in mice), whereas afatinib has a longer half-life

(approximately 21 hours in mice) (57, 58). In the in vitro work cells were treated continuously

for 48-72 hours with the drugs which could account for the difference in observed results when

comparing in vitro and in vivo results in this study. The half-life of afatinib in human patients is

approximately 30-40 hours and alpelisib is approximately 7.5 hours, suggesting with the daily

dosing used there would be better steady state levels than in the mouse (59, 60).

Between 20-50% of breast cancers exhibit PIK3CA mutations, with the most common being

hormone receptor positive or HER2 amplified breast cancer (61-66). It is noteworthy that

PIK3CA alterations found in this study (~36%) in EGFR amplified patients is similar to the

percentage observed in HER2 amplified patients (30%) (64-66). Combining HER2 targeted

therapy with PI3K inhibitors has been previously studied, and our study presents evidence that

there is a similar population of EGFR amplified patients who could benefit from combination

targeted therapy (67, 68).

EGFR and PI3K inhibitor combination has been studied in head and neck squamous cell

carcinoma, glioblastoma, ovarian cancer and breast cancer preclinical cell line models, however

these studies did not compare the synergistic effects of EGFR amplified and PI3K mutant cells to

non-amplified and non-mutant cells (26-34, 36, 37). Further, a recent high throughput screen

using breast cancer PDX models with PI3K mutations found that afatinib and alpelisib were

synergistic, which confirms our data, however they did not compare the synergy of models with

both EGFR amplification and PI3K mutations to those without (35). Our study indicates that it

is important to determine the subset of patients who would benefit from this combination, by

stratifying patients with EGFR amplification and PI3K pathway alterations. We also do not rule

out the possibility of combining these drugs with HER2 targeted therapies, as we found that up

to 7.7% of HER2 amplified/HER2 enriched tumors have amplified EGFR (Table 1). A study

showing synergy with BKM120 and the HER2 inhibitor Trastuzumab in patients provides a

proof of principle for combining PI3K inhibitors with EGFR family inhibitors may provide

clinical benefit in patients with EGFR family amplification and PI3K mutation (69).

We studied multiple EGFR (erlotinib or afatinib) and PI3K (BKM120 or alpelisib) inhibitors

(Fig. 3-6; Fig. S3, S5 and S6). Afatinib inhibits wild-type (WT) EGFR with the lowest IC50,

and then at increasing IC50s it inhibits L858R EGFR, HER2 and HER4 (70, 71). In addition to

PIK3CA, BKM120 also inhibits mTOR, VPS34 and DNAPK (72). Alpelisib is a more specific

PIK3CA inhibitor, without observed off target effects. Since we tested different EGFR and PI3K

inhibitors and observed similar reductions in downstream signaling, cell viability, cell cycle, and

apoptosis when the drugs were combined, it is likely that the observed effects were due to a class

effect of the drugs targeting EGFR and PIK3CA.

This study identifies that there is a subset of breast cancer patients with EGFR

amplification with co-incident PI3K pathway mutations who are more likely to respond to this

combination therapy. While this population represents a rare subset of metastatic breast cancer

patients, they are identified by genomic analyses that are routinely performed on breast cancer

patients as part of their clinical assessment. These analyses highlight approved drugs and

potential clinical trials available to these patients but currently there are no clinical trial data or

ongoing trials to support using EGFR and PI3K inhibitors in these patients and they are not

treated with these agents. Therefore, the combination of EGFR and PI3K pathway inhibitors

should be tested in breast cancer patients with EGFR amplification and PI3K pathway alteration.

In order to test if there is activity, a phase I/II trial of 30-50 patients would likely be needed to

see the response rate as this was sufficient to see activity of lapatinib in patients with metastatic

HER2-amplified breast cancers (73). Currently, there are ~150,000 patients living with

metastatic breast cancer in the US (74). Based on our analysis shown in Table 1 and 2, with

~1.2-5.1% of all breast cancers having an EGFR amplification, there would be ~ 1800-7650

patients whose tumor has EGFR amplification, with approximately half having a mutation in the

PI3K pathway. Gating only on TNBC, the prevalence of patients with metastatic TNBC is

approximately 15% of the total, or about 22,500 cases (75). Based on our data showing the

incidence of EGFR amplification between 3.2-8.4% in TNBC this would mean that that there

would be approximately 720-1890 cases with EGFR amplification, again with about half having

a PI3K pathway mutation. Thus, such a trial should be feasible. This ultimately could shift the

paradigm for the treatment regimen in this patient population with EGFR amplification,

especially triple negative breast cancer patients who have fewer targeted therapy options.

List of Abbreviations

EGFR- Epidermal Growth Factor Receptor

PI3K- Phosphoinositide 3-kinase

TNBC- Triple Negative Breast Cancer

FISH- fluorescent in situ hybridization

PI- propidium iodide

ER- Estrogen Receptor

HER2- Human epidermal growth factor receptor 2

mTOR- mechanistic target of rapamycin

AKT- Protein Kinase B

EGF- epidermal growth factor

MAPK- mitogen activated protein kinases

TKI- tyrosine kinase inhibitor

P70S6K- ribosomal protein S6 kinase beta 1

SEM- standard error of the mean

TCGA- The Cancer Genome Atlas Program

MSKCC- Memorial Sloan Kettering Cancer Center

IDC- invasive ductal carcinoma

ILC- Invasive lobular carcinoma

OS- overall survival

PTEN- phosphatase and tensin homolog

DMSO- dimethyl sulfoxide

PDX- patient derived xenograft

WT- wild-type

DNAPK- DNA-dependent protein kinase

Acknowledgements: We thank all members of the Lipkowitz lab in the Women's Malignancies

Branch, Center for Cancer Research, National Cancer Institute for their support and review of

this manuscript. We would like to thank Dr. Thomas Ried for his assistance with the planning of

the FISH experiment. We would like to acknowledge the CCR Genomics Core for their

assistance with sequencing, and the veterinarians and animal care staff at NIH for their assistance

with animal care, drug delivery and tumor measurement.

Authors' contributions: DJW performed immunoblotting, cell viability assays, cytotoxicity

assays, xenograft assays, helped with project conceptualization, experimental design, data

analysis and interpretation, and wrote the manuscript. DV performed immunoblotting, cell

viability assays, cytotoxicity assays, prepared samples for FISH and sequencing, performed flow

cytometry, helped with project conceptualization and experimental design. YA performed

immunoblotting and flow cytometry, helped with experimental design and data analysis. SD

analyzed Caris datasets and helped with data analysis and interpretation. SW ML and GS

assisted with the Caris dataset. Darawalee Wangsa, Danny Wangsa and KH performed FISH

analysis. YEG helped with experimental design, data analysis and interpretation, and animal

studies. SL analyzed cBioPortal datasets, helped with project conceptualization, data analysis

and interpretation, and manuscript editing. All authors read and approved the final manuscript.

Ethics approval and consent to participate: All animal aspects of this study were approved by

the NCI Animal Care and Use Committee (IACUC number WMB-004). All studies in the Caris

dataset were conducted in accordance with guidelines of the Declaration of Helsinki, Belmont

report, and U.S. Common rule, and in keeping with 45 CFR 46.101(b)(4), was performed

utilizing retrospective, deidentified clinical data. Per WCG IRB, it was considered institutional

review board (IRB) exempt with waiver of patient consent. 15,247 breast cancer samples

underwent molecular testing at Caris Life Sciences (Phoenix, AZ, USA), a CLIA/CAP-certified

laboratory. No specific inclusion criteria or exclusion criteria were applied aside from the

requirement for successful sequencing results.

Data Availability Statement: The MSKCC, TCGA and METABRIC datasets analyzed in this

study are available on cBioPortal.org (41-43). The Caris deidentified sequencing data are owned

by Caris Life Sciences and cannot be publicly shared due to the data usage agreement in place.

These data will be made available to researchers for replication and verification purposes

through the Caris letter of intent process, which are generally fulfilled within 6 months. For more

information on how to access this data, please contact Joanne Xiu at jxiu@carisls.com. All other

data generated in this study are included in this published article and its supplementary

information.

Competing interests: The authors declare that they have no competing interests.

Funding: This research was supported by the Intramural Research Program of the National

Institutes of Health (NIH) (ZIA BC 010977). The contributions of the NIH authors are

considered Works of the United States Government. The findings and conclusions presented in

this paper are those of the authors and do not necessarily reflect the views of the NIH or the U.S.

Department of Health and Human Services.

References

1.	Uribe ML, Marrocco I, Yarden Y. EGFR in Cancer: Signaling Mechanisms, Drugs, and Acquired Resistance. Cancers (Basel). 2021;13(11).

2.	Voldborg BR, Damstrup L, Spang-Thomsen M, Poulsen HS. Epidermal growth factor receptor (EGFR) and EGFR mutations, function and possible role in clinical trials. Ann Oncol. 1997;8(12):1197-206.

3.	Zhang X, Gureasko J, Shen K, Cole PA, Kuriyan J. An allosteric mechanism for activation of the kinase domain of epidermal growth factor receptor. Cell. 2006;125(6):1137-49.

4.	Levantini E, Maroni G, Del Re M, Tenen DG. EGFR signaling pathway as therapeutic target in human cancers. Seminars in Cancer Biology. 2022;85:253-75.

5.	Tebbutt N, Pedersen MW, Johns TG. Targeting the ERBB family in cancer: couples therapy. Nat Rev Cancer. 2013;13(9):663-73.

6.	Zeillinger R, Kury F, Czerwenka K, Kubista E, Sliutz G, Knogler W, et al. HER-2 amplification, steroid receptors and epidermal growth factor receptor in primary breast cancer. Oncogene. 1989;4(1):109-14.

7.	Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A, McGuire WL. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. Science. 1987;235(4785):177-82.

8.	Sjögren S, Inganäs M, Lindgren A, Holmberg L, Bergh J. Prognostic and predictive value of c-erbB-2 overexpression in primary breast cancer, alone and in combination with other prognostic markers. J Clin Oncol. 1998;16(2):462-9.

9.	Swain SM, Shastry M, Hamilton E. Targeting HER2-positive breast cancer: advances and future directions. Nat Rev Drug Discov. 2023;22(2):101-26.

10.	Mitsudomi T, Yatabe Y. Epidermal growth factor receptor in relation to tumor development: EGFR gene and cancer. Febs j. 2010;277(2):301-8.

11.	da Cunha Santos G, Shepherd FA, Tsao MS. EGFR mutations and lung cancer. Annu Rev Pathol. 2011;6:49-69.

12.	Nedaeinia R, Avan A, Manian M, Salehi R, Ghayour-Mobarhan M. EGFR as a potential target for the treatment of pancreatic cancer: dilemma and controversies. Curr Drug Targets. 2014;15(14):1293-301.

13.	Tathineni P, Joshi N, Jelinek MJ. Current State and Future Directions of EGFR-Directed Therapy in Head and Neck Cancer. Curr Treat Options Oncol. 2023;24(6):680-92.

14.	Sainsbury JR, Farndon JR, Needham GK, Malcolm AJ, Harris AL. Epidermal-growth-factor receptor status as predictor of early recurrence of and death from breast cancer. Lancet. 1987;1(8547):1398-402.

15.	Spyratos F, Delarue JC, Andrieu C, Lidereau R, Champème MH, Hacène K, Brunet M. Epidermal growth factor receptors and prognosis in primary breast cancer. Breast Cancer Res Treat. 1990;17(2):83-9.

16.	Burness ML, Grushko TA, Olopade OI. Epidermal Growth Factor Receptor in Triple-Negative and Basal-Like Breast Cancer: Promising Clinical Target or Only a Marker? The Cancer Journal. 2010;16(1):23-32.

17.	Al-Kuraya K, Schraml P, Torhorst J, Tapia C, Zaharieva B, Novotny H, et al. Prognostic relevance of gene amplifications and coamplifications in breast cancer. Cancer Res. 2004;64(23):8534-40.

18.	Bhargava R, Gerald WL, Li AR, Pan Q, Lal P, Ladanyi M, Chen B. EGFR gene amplification in breast cancer: correlation with epidermal growth factor receptor mRNA and protein expression and HER-2 status and absence of EGFR-activating mutations. Mod Pathol. 2005;18(8):1027-33.

19.   Ro J, North SM, Gallick GE, Hortobagyi GN, Gutterman JU, Blick M. Amplified and overexpressed
epidermal growth factor receptor gene in uncultured primary human breast carcinoma. Cancer Res.
1988;48(1):161-4.
20.   Li X, Zhao L, Chen C, Nie J, Jiao B. Can EGFR be a therapeutic target in breast cancer? Biochimica
et Biophysica Acta (BBA) - Reviews on Cancer. 2022;1877(5):188789.
21.   Reis-Filho JS, Milanezi F, Carvalho S, Simpson PT, Steele D, Savage K, et al. Metaplastic breast
carcinomas exhibit EGFR, but not HER2, gene amplification and overexpression: immunohistochemical
and chromogenic in situ hybridization analysis. Breast Cancer Res. 2005;7(6):R1028-35.
22.   Reis-Filho JS, Pinheiro C, Lambros MB, Milanezi F, Carvalho S, Savage K, et al. EGFR amplification
and lack of activating mutations in metaplastic breast carcinomas. J Pathol. 2006;209(4):445-53.
23.   Masuda H, Zhang D, Bartholomeusz C, Doihara H, Hortobagyi GN, Ueno NT. Role of epidermal
growth factor receptor in breast cancer. Breast Cancer Research and Treatment. 2012;136(2):331-45.
24.   O'DONOVAN N, CROWN J. EGFR and HER-2 Antagonists in Breast Cancer. Anticancer Research.
2007;27(3A):1285-94.
25.   Tan AR, Yang X, Hewitt SM, Berman A, Lepper ER, Sparreboom A, et al. Evaluation of biologic
end points and pharmacokinetics in patients with metastatic breast cancer after treatment with
erlotinib, an epidermal growth factor receptor tyrosine kinase inhibitor. J Clin Oncol. 2004;22(15):3080-
90.
26.   Yang Z, Liao J, Schumaker L, Carter-Cooper B, Lapidus RG, Fan X, et al. Simultaneously targeting
ErbB family kinases and PI3K in HPV-positive head and neck squamous cell carcinoma. Oral Oncol.
2022;131:105939.
27.   Guo T, Wu C, Zhang J, Yu J, Li G, Jiang H, et al. Dual blockade of EGFR and PI3K signaling
pathways offers a therapeutic strategy for glioblastoma. Cell Commun Signal. 2023;21(1):363.
28.   Sidorov M, Dighe P, Woo RWL, Rodriguez-Brotons A, Chen M, Ice RJ, et al. Dual Targeting of
EGFR and MTOR Pathways Inhibits Glioblastoma Growth by Modulating the Tumor Microenvironment.
Cells. 2023;12(4).
29.   Glaysher S, Bolton LM, Johnson P, Atkey N, Dyson M, Torrance C, Cree IA. Targeting EGFR and
PI3K pathways in ovarian cancer. Br J Cancer. 2013;109(7):1786-94.
30.   Li P, Zhang Q, Torossian A, Li ZB, Xu WC, Lu B, Fu S. Simultaneous inhibition of EGFR and PI3K
enhances radiosensitivity in human breast cancer. Int J Radiat Oncol Biol Phys. 2012;83(3):e391-7.
31.   El Guerrab A, Bamdad M, Bignon YJ, Penault-Llorca F, Aubel C. Co-targeting EGFR and mTOR
with gefitinib and everolimus in triple-negative breast cancer cells. Scientific reports. 2020;10(1):6367.
32.   Glaysher S, Bolton LM, Johnson P, Torrance C, Cree IA. Activity of EGFR, mTOR and PI3K
inhibitors in an isogenic breast cell line model. BMC Res Notes. 2014;7:397.
33.   Deng M, Wang J, Chen Y, Zhang L, Liu D. Combination of SF1126 and gefitinib induces apoptosis
of triple-negative breast cancer cells through the PI3K/AKT-mTOR pathway. Anticancer Drugs.
2015;26(4):422-7.
34.   She QB, Gruvberger-Saal SK, Maurer M, Chen Y, Jumppanen M, Su T, et al. Integrated molecular
pathway analysis informs a synergistic combination therapy targeting PTEN/PI3K and EGFR pathways for
basal-like breast cancer. BMC Cancer. 2016;16:587.
35.   Boyd DC, Zboril EK, Olex AL, Leftwich TJ, Hairr NS, Byers HA, et al. Discovering Synergistic
Compounds with BYL-719 in PI3K Overactivated Basal-like PDXs. Cancers. 2023;15(5):1582.
36.   Yi YW, Hong W, Kang HJ, Kim HJ, Zhao W, Wang A, et al. Inhibition of the PI3K/AKT pathway
potentiates cytotoxicity of EGFR kinase inhibitors in triple-negative breast cancer cells. Journal of
Cellular and Molecular Medicine. 2013;17(5):648-56.
37.   Garcia AR, Mendes A, Custódia C, Faria CC, Barata JT, Malhó R, et al. Abrogating Metastatic
Properties of Triple-Negative Breast Cancer Cells by EGFR and PI3K Dual Inhibitors. Cancers.
2023;15(15):3973.

38.	Hadadi A, Krause HB, Elliott A, Brown JT, Nazha B, Harik LR, et al. The Genomic Landscape of Urothelial Carcinoma with High and Low ERBB2 Expression. Cancers (Basel). 2023;15(24).

39.	Toboni MD, Wu S, Farrell A, Xiu J, Ribeiro JR, Oberley MJ, et al. Differential outcomes and immune checkpoint inhibitor response among endometrial cancer patients with MLH1 hypermethylation versus MLH1 "Lynch-like" mismatch repair gene mutation. Gynecol Oncol. 2023;177:132-41.

40.	Darabi S, Xiu J, Samec T, Kesari S, Carrillo J, Aulakh S, et al. Capicua (CIC) mutations in gliomas in association with MAPK activation for exposing a potential therapeutic target. Med Oncol. 2023;40(7):197.

41.	Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer Discov. 2012;2(5):401-4.

42.	Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, et al. Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. Sci Signal. 2013;6(269):pl1.

43.	de Bruijn I, Kundra R, Mastrogiacomo B, Tran TN, Sikina L, Mazor T, et al. Analysis and Visualization of Longitudinal Genomic and Clinical Data from the AACR Project GENIE Biopharma Collaborative in cBioPortal. Cancer Res. 2023;83(23):3861-7.

44.	Mermel CH, Schumacher SE, Hill B, Meyerson ML, Beroukhim R, Getz G. GISTIC2.0 facilitates sensitive and confident localization of the targets of focal somatic copy-number alteration in human cancers. Genome Biol. 2011;12(4):R41.

45.	Liyasova MS, Ma K, Voeller D, Ryan PE, Chen J, Klevit RE, Lipkowitz S. Cbl interacts with multiple E2s in vitro and in cells. PLoS One. 2019;14(5):e0216967.

46.	Darzynkiewicz Z, Juan G. DNA Content Measurement for DNA Ploidy and Cell Cycle Analysis. Current Protocols in Cytometry. 1997;00(1):7.5.1-7.5.24.

47.	Nair JR, Huang T-T, Sunkara A, Pruitt MR, Ibanez KR, Chiang C-Y, et al. Distinct effects of sacituzumab govitecan and berzosertib on DNA damage response in ovarian cancer. iScience. 2024;27(12).

48.	Zheng S, Wang W, Aldahdooh J, Malyutina A, Shadbahr T, Tanoli Z, et al. SynergyFinder Plus: Toward Better Interpretation and Annotation of Drug Combination Screening Datasets. Genomics, Proteomics & Bioinformatics. 2022;20(3):587-96.

49.	Prat A, Parker JS, Fan C, Perou CM. PAM50 assay and the three-gene model for identifying the major and clinically relevant molecular subtypes of breast cancer. Breast Cancer Res Treat. 2012;135(1):301-6.

50.	Jacot W, Mollevi C, Fina F, Lopez-Crapez E, Martin PM, Colombo PE, et al. High EGFR protein expression and exon 9 PIK3CA mutations are independent prognostic factors in triple negative breast cancers. BMC Cancer. 2015;15:986.

51.	Zhang J, Gao X, Schmit F, Adelmant G, Eck MJ, Marto JA, et al. CRKL Mediates p110β-Dependent PI3K Signaling in PTEN-Deficient Cancer Cells. Cell Reports. 2017;20(3):549-57.

52.	Kao J, Salari K, Bocanegra M, Choi YL, Girard L, Gandhi J, et al. Molecular profiling of breast cancer cell lines defines relevant tumor models and provides a resource for cancer gene discovery. PloS one. 2009;4(7):e6146.

53.	Stemke-Hale K, Gonzalez-Angulo AM, Lluch A, Neve RM, Kuo WL, Davies M, et al. An integrative genomic and proteomic analysis of PIK3CA, PTEN, and AKT mutations in breast cancer. Cancer Res. 2008;68(15):6084-91.

54.	Vasan N, Razavi P, Johnson JL, Shao H, Shah H, Antoine A, et al. Double PIK3CA mutations in cis increase oncogenicity and sensitivity to PI3Kα inhibitors. Science. 2019;366(6466):714-23.

55.   Tao JJ, Castel P, Radosevic-Robin N, Elkabets M, Auricchio N, Aceto N, et al. Antagonism of EGFR
and HER3 enhances the response to inhibitors of the PI3K-Akt pathway in triple-negative breast cancer.
Sci Signal. 2014;7(318):ra29.
56.   Alroy I, Yarden Y. The ErbB signaling network in embryogenesis and oncogenesis: signal
diversification through combinatorial ligand-receptor interactions. FEBS Lett. 1997;410(1):83-6.
57.   Hedges CP, Boix J, Jaiswal JK, Shetty B, Shepherd PR, Merry TL. Efficacy of Providing the PI3K
p110α Inhibitor BYL719 (Alpelisib) to Middle-Aged Mice in Their Diet. Biomolecules. 2021;11(2).
58.   Wang J, Su G, Yin X, Luo J, Gu R, Wang S, et al. Non-small cell lung cancer-targeted, redox-
sensitive lipid-polymer hybrid nanoparticles for the delivery of a second-generation irreversible
epidermal growth factor inhibitor-Afatinib: In vitro and in vivo evaluation. Biomed Pharmacother.
2019;120:109493.
59.   Murakami H, Tamura T, Takahashi T, Nokihara H, Naito T, Nakamura Y, et al. Phase I study of
continuous afatinib (BIBW 2992) in patients with advanced non-small cell lung cancer after prior
chemotherapy/erlotinib/gefitinib (LUX-Lung 4). Cancer Chemother Pharmacol. 2012;69(4):891-9.
60.   Juric D, Rodon J, Tabernero J, Janku F, Burris HA, Schellens JHM, et al. Phosphatidylinositol 3-
Kinase α-Selective Inhibition With Alpelisib (BYL719) in PIK3CA-Altered Solid Tumors: Results From the
First-in-Human Study. J Clin Oncol. 2018;36(13):1291-9.
61.   Armaghani AJ, Han HS. Alpelisib in the Treatment of Breast Cancer: A Short Review on the
Emerging Clinical Data. Breast Cancer (Dove Med Press). 2020;12:251-8.
62.   Bailey MH, Tokheim C, Porta-Pardo E, Sengupta S, Bertrand D, Weerasinghe A, et al.
Comprehensive Characterization of Cancer Driver Genes and Mutations. Cell. 2018;173(2):371-85.e18.
63.   Pereira B, Chin SF, Rueda OM, Vollan HK, Provenzano E, Bardwell HA, et al. The somatic
mutation profiles of 2,433 breast cancers refines their genomic and transcriptomic landscapes. Nat
Commun. 2016;7:11479.
64.   Razavi P, Chang MT, Xu G, Bandlamudi C, Ross DS, Vasan N, et al. The Genomic Landscape of
Endocrine-Resistant Advanced Breast Cancers. Cancer Cell. 2018;34(3):427-38.e6.
65.   Comprehensive molecular portraits of human breast tumours. Nature. 2012;490(7418):61-70.
66.   Savas P, Lo LL, Luen SJ, Blackley EF, Callahan J, Moodie K, et al. Alpelisib Monotherapy for PI3K-
Altered, Pretreated Advanced Breast Cancer: A Phase II Study. Cancer Discov. 2022;12(9):2058-73.
67.   Rodon J, Pérez-Fidalgo A, Krop IE, Burris H, Guerrero-Zotano A, Britten CD, et al. Phase 1/1b
dose escalation and expansion study of BEZ235, a dual PI3K/mTOR inhibitor, in patients with advanced
solid tumors including patients with advanced breast cancer. Cancer Chemother Pharmacol.
2018;82(2):285-98.
68.   Loibl S, de la Pena L, Nekljudova V, Zardavas D, Michiels S, Denkert C, et al. Neoadjuvant
buparlisib plus trastuzumab and paclitaxel for women with HER2+ primary breast cancer: A randomised,
double-blind, placebo-controlled phase II trial (NeoPHOEBE). Eur J Cancer. 2017;85:133-45.
69.   Yu F, Zhao J, Hu Y, Zhou Y, Guo R, Bai J, et al. The combination of NVP-BKM120 with trastuzumab
or RAD001 synergistically inhibits the growth of breast cancer stem cells in vivo. Oncol Rep.
2016;36(1):356-64.
70.   Li D, Ambrogio L, Shimamura T, Kubo S, Takahashi M, Chirieac LR, et al. BIBW2992, an
irreversible EGFR/HER2 inhibitor highly effective in preclinical lung cancer models. Oncogene.
2008;27(34):4702-11.
71.   Ferrarotto R, Gold KA. Afatinib in the treatment of head and neck squamous cell carcinoma.
Expert Opin Investig Drugs. 2014;23(1):135-43.
72.   Burger MT, Pecchi S, Wagman A, Ni Z-J, Knapp M, Hendrickson T, et al. Identification of NVP-
BKM120 as a Potent, Selective, Orally Bioavailable Class I PI3 Kinase Inhibitor for Treating Cancer. ACS
Medicinal Chemistry Letters. 2011;2(10):774-9.

73.   Johnston S, Trudeau M, Kaufman B, Boussen H, Blackwell K, LoRusso P, et al. Phase II study of
predictive biomarker profiles for response targeting human epidermal growth factor receptor 2 (HER-2)
in advanced inflammatory breast cancer with lapatinib monotherapy. J Clin Oncol. 2008;26(7):1066-72.
74.   Gallicchio L, Devasia TP, Tonorezos E, Mollica MA, Mariotto A. Estimation of the Number of
Individuals Living With Metastatic Cancer in the United States. J Natl Cancer Inst. 2022;114(11):1476-83.
75.   Waks AG, Winer EP. Breast Cancer Treatment: A Review. Jama. 2019;321(3):288-300.

Table Legends

Table 1. Incidence of EGFR amplification in breast cancer. Breast cancer subtype analysis of

patients in TCGA, METABRIC, MSKCC and Caris dataset with EGFR amplification. IDC,

invasive ductal carcinoma; ILC, invasive lobular carcinoma; nd, not done.

Table 2. PI3K pathway mutations in EGFR amplified breast cancer. Datasets from TCGA,

METABRIC and MSKCC combined showing alterations in PI3K pathway genes in 75 patients

with EGFR amplification. Dataset from Caris showing alterations in PI3K pathway genes in

1,177 breast cancers with EGFR amplification.

Figure Legends

Figure 1. Survival of patients with EGFR amplification in breast cancer. A) Combination of

TCGA, METABRIC, MSKCC datasets to analyze overall survival (OS) of patients comparing

EGFR amplification (red line) to patients without EGFR amplification (blue line). B) OS of

patients from Caris dataset with EGFR amplification (red line) compared to patients without

EGFR amplification (blue line).

Figure 2. EGFR protein expression and gene amplification, and PI3K mutations in MDA-

MB-231, MDA-MB-468 and BT20 cells. A) Western blot showing protein expression of

EGFR, PIK3CA, pTEN and HSC70 (loading control) in MDA-MB-231, MDA-MB-468 and

BT20 cells. B) FISH analysis showing EGFR (red) and CEP7 (aqua, control) in MDA-MB-231,

MDA-MB-468 and BT20 cell lines. Average EGFR signal per nuclei was quantified using

ImageJ, and statistical analysis was performed by Student's t-test, with **** indicating p<0.0001

and ** indicating p<0.01 compared to MB231. C) DNA sequencing for PIK3CA mutation status

in BT20, MDA-MB-468 and MDA-MB-231 cell lines.

Figure 3.   EGFR/PI3K dual inhibition significantly reduces downstream signaling in

EGFR amplified and PI3K altered breast cancer. A) BT-20 and MDA-MB-468 cells were

pretreated for 24 hours with 0, 10 or 100 nM afatinib, and then starved in serum-free RPM1

media for 3 hours prior to stimulation with or without 25 ng/mL EGF for fifteen minutes. Cells

were lysed and immunoblotted for pEGFR (Y1068), EGFR, pMAPK (T202/Y204), ERK2 and

HSC70 (loading control). B) MDA-MB-231, MDA-MB-468 and BT20 cells were treated with

100 nM alpelisib (Alp), 1 µM afatinib (Afat) or a combination of both for 24 hours, and were

then lysed and immunoblotted for pAKT (S473), AKT, pMAPK (T202/Y204), ERK2, p-P70S6K

(T389), P70S6K, pS6 (S235/236), S6 and HSC70 (loading control). In three independent

experiments, the band density of the phosphorylated protein relative to total protein was

averaged ±SEM for C) pMAPK/ERK2, D) pAKT/AKT, E) p-P70S6K/P70S6K and F) pS6/S6.

Two-way ANOVA was performed, where ns indicates not statistically significant, * indicates

p<0.05, ** indicates p<0.01, *** indicates p<0.001 and **** indicates p<0.0001.

Figure 4. Dual EGFR/PI3K inhibition synergistically reduce cell viability in TNBC with

EGFR amplification and PI3K alteration. A) MDA-MB-231 in three independent

experiments, B) MDA-MB-468 in four independent experiments and C) BT20 cells in three

independent experiments were treated with the indicated doses for 48 hours, and then viability

was determined by Promega CellTiter-Glo 2.0. Data represents the average ±SEM of the

independent experiments, and statistical analysis was performed as Two-Way ANOVA, and the

p value indicates interaction, where ns indicates not statistically significant, ** indicates p<0.01,

*** indicates p<0.001 and **** indicates p<0.0001. The averages were also subjected to four

different synergy score analyses through synergyfinder.org, where a score of >+10 indicates

synergy, -10 to +10 indicates additive effects, and <-10 indicates antagonism.

Figure 5. EGFR/PI3K inhibition induce cell cycle arrest and apoptosis in EGFR amplified

and PI3K altered TNBC. A) MDA-MB-231 were treated with DMSO (control), 1 µM afatinib,

10 µM alpelisib, or the combination of both B) MDA-MB-468 were treated with DMSO

(control), 1 µM afatinib, 10 µM alpelisib, or the combination of both C) BT20 were treated with

DMSO (control), 100 nM afatinib, 1 µM alpelisib, or the combination of both and for 48 hours,

and then were harvested, washed, fixed and stained with propidium iodide before being analyzed

by flow cytometry and FlowJo for cell cycle changes. Data represents the average ±SEM of at

least three independent experiments, and Two-Way ANOVA statistical analysis was performed

where * indicates p<0.05, ** indicates p<0.01, *** indicates p<0.001 and **** indicates

p<0.0001. D) MDA-MB-231 (1 µM alpelisib, 1 µM afatinib), E) MDA-MB-468 (3 µM alpelisib,

10 µM afatinib) or F) BT20 cells (1 µM alpelisib, 1 µM afatinib) were pretreated with 20 µM Z-

VAD-FMK (ZVAD) for one hour, and then were treated with indicated drugs with continued

ZVAD treatment in 1% FBS RPM1 media containing propidium iodide for 72 hours. Imaging of

the cells was performed by BioTek Cytation 1, and dead cell percentage was calculated by

dividing PI positive cells by the total brightfield cell count and multiplying by 100. Data

represents the average ±SEM of three independent experiments. Statistical analysis was

performed by Student's t-test, with *** indicating p<0.001.

Figure 6. Dual EGFR and PI3K inhibition is most effective at reducing tumor volume *in*

vivo. A) BT20 cells were injected subcutaneously into the mammary fat pad of nude mice. 56

days post implantation, when the average size was 100 mm^3 , mice were randomized (R) to 15

mice into four groups and treated starting on day 63 post implantation (Tx) with control (0.5%

carboxymethylcellulose) (black), 20 mg/kg afatinib (red), 20 mg/kg alpelisib (green), or the

combination of both afatinib and alpelisib (blue) Monday through Friday for 36 days. Zoomed

inset shows day 63 to 100. Statistical analysis was performed by Student's t-test, and *

indicates p<0.05 when comparing afatinib and alpelisib to control on day 94 and day 98. B)

After four days of treatment, four mice from each group were euthanized and the protein from

the tumors was collected and immunoblotted for pEGFR (Y1068), EGFR, pS6 (S235/236), S6,

and HSC70 (loading control). Data represents the average of four mice from each group ±SEM

and statistical analysis was performed by Student's t-test, where ns indicates not statistically

significant, * indicates p<0.05 and ** indicates p<0.01. C) Bar graph of data from A on day 98

(36 days of drug treatment). Statistical analysis was performed by Student's t-test, where *

indicates p<0.05 when compared to control. No other comparisons were statistically significant.

Supplemental Figures

Figure S1

File Format: .tiff

Title of Data: Overall survival of patients with EGFR amplification by subtype.

Figure S1 Legend: Using the Caris dataset, overall survival of breast cancer patients with EGFR

amplification (red) vs WT (blue) in A) Triple negative breast cancer, B) Basal, C) HER2

enriched, or D) ER+ HER2-.

Figure S2

File Format: .tiff

Title of Data: Oncoplot for PI3K pathway alterations in EGFR amplified breast cancer.

Figure S2 Legend: Patient data from cbioportal.org plotting varying mutations or amplifications

by gene in the PI3K pathway.

Figure S3

File Format: .tiff

Title of Data: EGFR/PI3K dual inhibition significantly reduce downstream signaling in EGFR

amplified and PI3K altered TNBC.

Figure S3 Legend: A) MDA-MB-468 and BT20 cells were starved in serum-free RPM1 media

for 1 hour with 0, 1 or 10 µM erlotinib, and then stimulated with or without 25 ng/mL EGF for

10 minutes. Cells were lysed and immunoblotted for pEGFR (Y1068) and EGFR (loading

control). B) Darker exposure for MDA-MB-231 panel from Figure 4B which was used for

quantification. C) MDA-MB-231, MDA-MB-468 and BT20 cells were treated with 10 µM

erlotinib, 1 µM BKM120 or a combination of both for 24 hours, and were then lysed and

immunoblotted for pAKT (S473), AKT, pMAPK (T202/Y204), ERK2, p-P70S6K (T389),

P70S6K and HSC70 (loading control). In three independent experiments, the band density of the

phosphorylated protein relative to total protein was averaged ±SEM for D) pMAPK/ERK2, E)

pAKT/AKT and F) p-P70S6K/P70S6K. Student's t-test was performed, where ns indicates not

statistically significant, * indicates $p<0.05$, ** indicates $p<0.01$, *** indicates $p<0.001$ and ****

indicates $p<0.0001$.

Figure S4

File Format: .tiff

Title of Data: Erlotinib treatment does not activate pHER3.

Figure S4 Legend: MDA-MB-231, MDA-MB-468 and BT20 cells were treated with 10 µM

erlotinib, 1 µM BKM120 or the combination of both for 24 hours before the cells were lysed and

immunoblotted for pHER3 (Y1289) and HER3 (loading control).

Figure S5

File Format: .tiff

Title of Data: EGFR/PI3K inhibition reduces viability, induces cell cycle arrest in TNBC with

EGFR amplification, PI3K alteration.

Figure S5 Legend: A) MDA-MB-231, B) MDA-MB-468 and C) BT20 cells were treated with

the indicated doses for 48 hours, and then viability was determined by Promega CellTiter-Glo

2.0. Data represents the average ±SEM of three independent experiments, and statistical analysis

was performed as Two-Way ANOVA, and the p value indicates interaction, where ns indicates

not statistically significant, * indicates $p < 0.05$, *** indicates $p < 0.001$ and **** indicates

$p < 0.0001$. D) MDA-MB-231, E) MDA-MB-468 and F) BT20 cells were treated with DMSO

control, 1 µM erlotinib or 10 µM BKM120 or the combination of erlotinib and BKM120 for 48

hours, and then were harvested, washed, fixed and stained with propidium iodide before being

analyzed by flow cytometry and FlowJo for cell cycle changes. Data represents the average

±SEM of at least three independent experiments, and Two-Way ANOVA statistical analysis was

performed where * indicates $p < 0.05$, ** indicates $p < 0.01$ and *** indicates $p < 0.001$.

Figure S6

File Format: .tiff

Title of Data: Dual EGFR/PI3K inhibition induce apoptosis in EGFR amplified and PI3K

altered TNBC.

Figure S6 Legend: A) MDA-MB-231, B) MDA-MB-468 and C) BT20 cells were pretreated

with 20 µM Z-VAD-FMK (ZVAD) for one hour, and then were treated with DMSO, 10 µM

erlotinib, 1 µM BKM120 or the combination of both, with continued ZVAD treatment for 48

hours. Dead cell percentage was calculated by using CytoTox-Glo by Promega. Data represents

the average ±SEM of three independent experiments. Statistical analysis was performed by One-

way ANOVA, with ns indicating not statistically significant, * indicating $p < 0.05$, ** indicating

$p < 0.01$ and *** indicating $p < 0.001$.

A MB231

B MB468

C BT20

MB231			
ZIP	Loewe	HSA	Bliss
4.93	8.04	9.94	6.08

Synergy Scores

MB468			
ZIP	Loewe	HSA	Bliss
10.67	16.13	18.62	10.53

Synergy Scores

BT20			
ZIP	Loewe	HSA	Bliss
24.49	24.6	30.42	24.31

Breast Cancer Subtype	TCGA		METABRIC		MSKCC		Caris	
	Samples #	EGFR Amplification (%)	Samples #	EGFR Amplification (%)	Samples #	EGFR Amplification (%)	Samples #	EGFR Amplification (%)
All Patients	816	14 (1.7%)	2173	52 (2.4%)	1918	23 (1.2%)	23160	1177 (5.1%)
Histology								
IDC	489	10 (2.0%)	1660	48 (2.9%)	1473	21 (1.4%)	8411	434 (5.2%)
Mixed IDC/ILC	88	0 (0%)	218	2 (0.9%)	89	0 (0%)	nd	nd
ILC	127	0 (0%)	172	1 (0.6%)	299	0 (0%)	1311	27 (2.1%)
Other Histologies	112	4 (3.6%)	123	1 (0.8%)	57	2 (3.5%)	13438	716 (5.3%)
ER and HER2 Status								
ER+	600	5 (0.8%)	1617	23 (1.4%)	1512	13 (0.9%)	10802	363 (3.4%)
ER+/HER2-	326	3 (0.9%)	1398	19 (1.4%)	1357	11 (0.8%)	9887	342 (3.5%)
ER Negative	175	8 (4.6%)	523	28 (5.4%)	327	9 (2.8%)	6312	512 (8.1%)
HER2 Positive	121	3 (2.5%)	247	9 (3.6%)	240	3 (1.3%)	1625	73 (4.5%)
ER+/HER2+	89	1 (1.1%)	108	1 (0.9%)	76	1 (1.3%)	916	21 (2.3%)
ER-/HER2+	31	2 (6.5%)	139	8 (5.8%)	79	1 (1.3%)	709	52 (7.3%)
TNBC	90	6 (6.7%)	335	19 (5.7%)	251	8 (3.2%)	5335	446 (8.4%)
Molecular Subtype								
Total Profiled	587	10 (1.7%)	1980	47 (2.4%)	nd	nd	12872	611 (4.6%)
Luminal A	307	2 (0.7%)	700	5 (0.7%)	nd	nd	1764	51 (2.9%)
Luminal B	122	2 (1.6%)	475	7 (1.5%)	nd	nd	5402	153 (2.8%)
HER2 Enriched	51	2 (3.9%)	224	12 (5.4%)	nd	nd	2233	173 (7.7%)
Basal	107	4 (3.7%)	209	17 (8.1)	nd	nd	3367	230 (6.8%)
Claudin Low	nd	nd	218	3 (1.4%)	nd	nd	nd	nd
Normal	nd	nd	148	3 (2.0%)	nd	nd	106	4 (3.8%)

Gene	TCGA, METABRIC, MSKCC combined Potential PI3KPathway Alterations (%)	Caris Potential PI3KPathway Alterations (%)
PIK3CA	36	36.5
PIK3R1	4	4.32
PIK3R2	4	0
PIK3R3	4	Not Available
PTEN	5.3	10.5
AKT1	5.3	5.62
AKT2	6.7	0
AKT3	17.3	0
RPS6KB1	8	Not Available
RPS6KB2	4	Not Available
RPS6	4	Not Available
Patients with at least one PI3K activation	70.7	50.8