Drivers of genetic diversity across the marine tree of life

Rachel H. Toczydlowski,[1,2*] Reid S. Brennan,[3,4] Eric D. Crandall,[5]
Joanna L. Kelley,[6] James M. Pringle,[7] Cynthia Riginos,[8,9] John P. Wares,[10]
Gideon S. Bradburd[11*]

[1] Department of Integrative Biology, Michigan State University, East Lansing, MI, 48824, U.S.
[2] Northern Research Station, USDA Forest Service, Rhinelander, WI, 54501, U.S.
[3] Marine Evolutionary Ecology, GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, 24105, Germany
[4] Marine Mammal and Turtle Division, Southeast Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Miami, FL, U.S.
[5] Department of Biology, The Pennsylvania State University, State College, PA, 16802, U.S.
[6] Ecology and Evolutionary Biology, University of California Santa Cruz, Santa Cruz, CA, 95064, U.S.
[7] Ocean Process Analysis Laboratory, University of New Hampshire, Durham, NH, 03824, U.S.
[8] School of the Environment, The University of Queensland, Brisbane, QLD, 4072, Australia
[9] Australian Institute of Marine Science, Cape Cleveland, QLD, 4810, Australia
[10] Odum School of Ecology, Department of Genetics, University of Georgia, Athens, GA, 30602, U.S.
[11] Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI, 48109, U.S.

*To whom correspondence may be addressed. E-mail: bradburd@umich.edu or rhtoczydlowski@gmail.com.

Author Contributions: All authors contributed to conceptualization and writing. R.H.T. and R.S.B. designed the bioinformatics workflow and processed the genomic datasets. E.D.C., C.R., and J.P.W. amassed and curated biotic trait data. J.M.P. and R.H.T. calculated abiotic predictors. G.S.B. conceived of population genetic analyses. R.H.T and G.S.B provided supervision and leadership. We hold the time we had to work with and learn from E. D. Crandall fondly and with gratitude; more about his life and career can be read at https://rcn-ecs.github.io/.

Competing Interests Statement: The authors have no competing interests to declare.

Classification: Biological Sciences\Evolution

Keywords: conservation, comparative phylogeography, genetic diversity, life history, macro-genetics, marine biodiversity, spatial population genomics

This PDF file includes:

Main Text
Figures 1 to 3
Supporting Information

Abstract

Why do some species have more genetic diversity than others? This question is one of the greatest remaining mysteries in evolutionary biology, yet we still know little about what factors predict genetic diversity among species. Efforts to quantify diversity or understand its distribution have been hampered by a lack of genomic data analyzed in a consistent way with statistical approaches that can move beyond the population level to generate species-wide estimates of genetic diversity. We address this critical gap by generating standardized estimates of genomic-level genetic diversity for 93 species sampled over 9,000 localities. Because marine species have evolved a strikingly diverse array of biological traits that have long been hypothesized to affect species-level genetic diversity, we focus our study on marine taxa. For each species, we aggregated biotic traits related to life history and abiotic features of the species' range. We show that genetic diversity increases with species range extent and planktonic dispersal. We hypothesize that these traits increase a species' ability to avoid or recover from bottlenecks, thereby maintaining genetic diversity. Our findings provide insights into how biotic and abiotic factors interact to shape genomic variation in the ocean, and offer a predictive framework for understanding marine biodiversity in the face of global change.

Significance statement

Species exhibit vast trait diversity that allows populations to persist. However, diversity in these traits — e.g., how organisms reproduce, disperse, and feed — is not easily linked to what is known of genomic diversity, even in well-studied organisms. We investigate a diverse assemblage of marine species to begin to build a predictive framework for how species' traits drive genomic diversity across the world's oceans. We find that species with larger geographic ranges have higher genomic diversity. More dispersive species also harbor higher diversity,

suggesting that greater dispersal facilitates resilience to environmental change. As available datasets increase, our ability to understand fundamental links between trait and genetic diversity and how they shape species' responses to environmental change will continue to improve.

Main Text

Introduction

Why does genetic diversity vary among species? This is a central question in population genetics (*1–4*), and may inform a species' adaptive potential, extinction risk, or conservation priority (*5–9*). Little is known about the factors that predict the observed variation in diversity across species (*10*). Indeed, for most species, we have no estimate of genetic diversity at all.

Species-level genetic diversity, which we define as the total level of genetic polymorphism within a species, is ultimately the outcome of the relative rates of alleles entering and leaving the species gene pool, and is thus controlled by population genetic and ecological processes that affect those rates (*10*), such as selection, the mutation rate, and population demography. Although selection, including linked and background selection, can affect patterns of genetic diversity across the genome (*11, 12*), evidence from both theoretical and empirical studies suggests that selection is unlikely to be the major determinant of variation in genetic diversity among species (*4, 13*). And, while data on mutation rates are rare for most non-model organisms (*14*), previous work suggests that variation in mutation rate alone is also unlikely to explain patterns of genetic diversity across species (*3*).

Population genetic theory offers several hypotheses about the mechanisms by which different factors should increase or decrease genetic diversity within a species. First, the genetic diversity of a population is expected to increase with the census number of reproductive individuals (*15*). Second, for a continuously distributed species, the total amount of genetic diversity is

expected to decrease with a species' dispersal ability and increase with its range extent (*16, 17*). Dispersal homogenizes genetic variation that has arisen locally due to drift or selection (*18*), and therefore, can reduce the total diversity found across all demes of a structured metapopulation (*17, 19*). Third, bottlenecks, whether local (i.e., confined to a local deme) or range-wide, are expected to decrease a species' genetic diversity (*20–22*). Therefore, traits that impact a species' population size, structure, and demography, have the potential to drive patterns of genetic diversity across species.

Specific biological traits that have been hypothesized to predict genetic diversity include dispersal traits, morphological characteristics, and life history traits (*23–25*), as well as abiotic factors such as habitat stability and range location (*26–29*). In addition, factors that emerge as a function of the interaction between biotic and abiotic traits, like range size and geographic complexity (*30*), can also influence both recent and historical population demography, and thereby affect genetic diversity. Understanding why levels of genetic diversity vary within and across species has received considerable attention (*31, 32*) since the first molecular measurements of genetic variation were made (*33, 34*), but despite these clear theoretical predictions and hypothesized relationships, empirically testing these ideas has proven challenging.

Several factors have hindered empirical progress in understanding drivers of genetic diversity across species. First, quantifying genetic diversity remains difficult, particularly in species for which field sampling is laborious or costly. Estimates made from limited genetic information, such as mitochondrial DNA or microsatellites, might give noisy (*35*) or biased (*36, 37*) estimates of diversity, but generating genomic datasets remains expensive. Second, even with genomic sequence data in hand, there are myriad bioinformatic pipelines and choices used to transform raw sequence data into assembled loci and called genotypes. Each bioinformatic decision point can have large impacts on downstream analyses (*38*). To make meaningful comparisons of genetic diversity, it is therefore vital that the methods used for analyzing sequencing

data and subsequently estimating genetic diversity are standardized across species, a bioinformatics challenge that has rarely been achieved at scale for genomic data. Finally, in comparative population genetic studies, estimates of genetic diversity are implicitly treated as species-level traits, which requires correcting for phylogenetic non-independence (*39*). Implementing such a phylogenetic correction necessitates an accurate, time-calibrated phylogeny that includes all species under consideration. Our understanding of phylogenetic relationships across the tree of life continues to grow, but many species still lack a branch tip, constraining the scope of comparative population genetic analyses that can be conducted.

Heterogeneity in sampling effort across species, and specifically in the spatial pattern and extent of sampling, poses another hurdle to a comparative analysis of species-level genetic diversity. Because many (if not most) species exhibit a pattern of isolation by distance (*40, 41*) – as well as possible hierarchical or cryptic spatial structure – the geographic extent of sampling may often be a strong predictor of the estimated genetic diversity in a species. Even if sampling effort were standardized, variation in density, dispersal, and recent demography across species could result in biased estimates of a species' genetic diversity (*42*). These factors impede our ability to accurately quantify genetic diversity within species and test hypotheses about its predictors across species.

In the context of global environmental change and the sixth mass extinction event currently occurring, as well as from a basic science perspective, the ability to determine the effect of different factors on genetic diversity is critical (*43, 44*). Here, we begin to address this by building a predictive framework using a novel, unified, spatially and phylogenetically explicit approach to test hypotheses about drivers of genetic diversity. We focus on marine biodiversity because oceans contain a taxonomically and functionally diverse biotic assemblage (*45*). How this marine biodiversity is generated and maintained in a dynamic, fluid environment, without obvious barriers to gene flow, continues to be a fundamental problem in evolutionary biology

(*46–48*). Moreover, marine biota exhibit tremendous variation in range size and life history traits that govern connectivity across those ranges (*24*). These biological traits interact with oceanic processes, both physical and chemical, to generate spatial patterns of genomic diversity, which in turn feedback to affect speciation and biogeography (*49–51*). The marine environment can impact all levels of biological processes, including huge variation in population size (*4, 52*), the importance of passive gamete or larval dispersal (*53, 54*), and dispersal affected by ocean current movements (*55, 56*). This diversity of traits makes marine organisms well-suited for testing long-standing hypotheses about how biological traits and abiotic environments shape species-wide genetic diversity. There is also a critical need to understand and generalize such diversity patterns because oceanic biodiversity is being rapidly affected by climate change (*57–61*) and other anthropogenic threats, including overfishing (*62*), pollution (*63*), disease (*64*), and habitat change. Our approach addresses these gaps and gives insight into the drivers of genetic diversity across marine species while also establishing a template for future studies to test hypotheses about why genetic diversity varies across diverse taxa.

Results

We identified and downloaded all georeferenced, publicly available reduced-representation and whole-genome datasets of marine species present in the International Nucleotide Sequence Database Collaboration as of October 2020. We included as broad of a phylogenetic diversity of species as we could in order to maximize the generality of our findings, but we excluded datasets containing laboratory-reared, freshwater, or captive populations, as well as those with individuals sampled outside their native range. The final metadataset of 93 sampled species covered a significant proportion of the phylogenetic diversity of the eukaryotic tree of life (Fig 1), including representatives from vascular plants, invertebrate animals — including cnidarians, crustaceans, and molluscs — and vertebrate animals — including cartilaginous fishes, mam-

mals, birds, reptiles, and ray-finned fishes. These datasets also met our constraint that they contain a minimum of 10 individuals (mean: 101, range: $8-366$) sampled from at least 2 distinct geographic locations (mean: 18, range: $2-121$). Starting from raw sequence reads, we filtered, cleaned, and assembled each of these 93 genomic datasets using multiple parameter combinations, then picked the best set using an objective optimality criterion (see Methods). This resulted in a final genomic metadataset of 93 species (Figure 1a); each species was represented by a mean of 101 individuals (range: $8-366$), 3.6 million raw sequence reads per individual (range: $0.4-30.1$ million), 3.7 million aligned nucleotide base pairs (range: $5,800-22.3$ million), and 37,463 loci (range: $86-187,685$).

We then employed the novel parametric modeling approach introduced by Hancock *et. al* (*65*) to generate estimates of species-level genetic diversity that are robust to heterogeneity in sampling area and intensity. Briefly, we fit an asymptotic curve to the observed Isolation by Distance (IBD) pattern of pairwise homozygosity (*40, 66*) in each species. We calculated geographic distance here as the distance traveling via the sea between the locations of sampled individuals; pairwise homozygosity is the probability that a pair of alleles sampled at random from two individuals are the same. The asymptote of this IBD curve — a parameter estimated as part of the model-fitting procedure — captures the extrapolated genetic diversity contained within the entire species range, which we refer to as π. By including multiple individuals per species and estimating diversity in an explicitly geographic statistical framework, this approach represents an advance on previous estimates made from a small number of "representative individuals" that are treated as experimental replicates. Our estimates of π in the world's oceans varied greatly across species, spanning two orders of magnitude from 3.369×10^{-4} to 3.001×10^{-2}, with a mean of 5.224×10^{-3} (phylogenetically corrected mean: 5.335×10^{-3}) (Figure 1c and S3).

To test hypotheses about which biotic and abiotic traits predict genetic diversity across ma-

rine species, we assembled a time-calibrated phylogeny and aggregated biotic trait data as well as abiotic factors for all 93 species in our dataset (Figure 1b). Genetic diversity had significant phylogenetic structure (Pagel's λ=0.697, $p = 0.039$; Blomberg's K=0.098, $p = 0.016$) [67–69]; the correlation between phylogenetic relatedness and similarity in genetic diversity was significant over the last 490My, and the strength of the correlation increased monotonically as it approached the present (Figure S6). Genetic diversity is also affected by species-level characteristics and mechanisms that emerge as the result of the interaction of biotic and abiotic processes, such as effective dispersal ability, reproductive census population size, and the frequency, magnitude, duration, and geography of bottlenecks. These characteristics and mechanisms are difficult to directly observe, so instead we collated biotic and abiotic traits that are hypothesized to affect them. The suite of biotic traits is comprised of body size, egg size, spawning, pelagic larval duration, philopatry, iteroparity, planktotrophy (larval feeding behavior), benthicity (degree of association with seafloor), and planktonicity (qualitative association with drifting development as a component of dispersal). Abiotic predictors consisted of mean species' range latitude, the number of ecoregions within the species' range (ecological complexity and its potential to act as a barrier to gene flow), and overall range extent. All traits are further described in the Supplemental Methods and Appendices.

Range extent, number of ecoregions, and planktonicity had significant positive effects on species-level genetic diversity (Figures 2 and 3); species with planktonic life stages, as well as species with larger range extents or ranges that included more ecological regions, all had higher expected π (Table S1). Range extent, defined as the greatest overwater distance between any pair of individuals of that species in the Global Biodiversity Information Facility (GBIF) – with filtering to ensure incidental range-edge observations did not inflate this value, had a mean effect of 0.53 (95% CI: 0.06–0.95). Number of ecoregions, defined as the number of ecoregion polygons from Spalding et al. (2007) [70] that contained filtered GBIF observation

points for that species, had a mean effect of 0.007 (95% CI: -0.0002 – 0.011). Note that we use ecoregions to quantify the number of discontinuities in biodiversity and abiotic features within a species' range that may limit gene flow (and thereby increase species-level genetic diversity). Planktonicity, defined as whether the species was planktonic at any point in their life cycle, had a mean effect of 0.560 (95% CI: -0.055–1.17). Planktotrophy and free-spawning had a strongly positive estimated effect on genetic diversity, and the effects of body size, iteroparity, and philopatry were strongly negative. However, these effects' credible intervals overlapped zero and were not inferred to be significant (see Methods), so those results should be interpreted with caution. Range extent and number of ecoregions were highly correlated (Kendall's τ = 0.78; Figure S2), and subsequent analyses (see Methods) suggest that the inferred effect of ecoregions on genetic diversity is at least partially driven by the range extent. Planktonicity and range extent were very weakly correlated (Kendall's τ = -0.05; Figure S2), indicating that the significance levels of their effects are independent. The Bayesian R^2 values (*71*) for the models that included range extent and planktonicity as biological predictors (Figure 3) were 0.55 (95%CI: 0.29-0.76) and 0.56 (95%CI: 0.34-0.75) respectively.

Discussion

Our analyses reveal enormous variation in the amount of genetic diversity harbored within marine species across the tree of life. Estimates of species-level genetic diversity varied over almost two orders of magnitude. Such variation offers immense opportunity to test hypotheses about what factors predict genetic diversity across species. We found strong phylogenetic signal in genetic diversity, suggesting that diversity may be reasonably predicted in species of known phylogenetic position but lacking population genomic sampling resources. We also found strong evidence that range extent and planktonicity were positively correlated with genetic diversity (Figure 3), in line with previous findings about the roles of range size and dispersal (*3*). Unex-

pectedly, we found that all dispersal-related traits yielded positive effects on genetic diversity, contrary to what is expected from theory; we propose that these relationships reflect the effects of dispersal-related traits on species' capacity to avoid or recover from bottlenecks. Below, we relate the biotic and abiotic traits we collected and their estimated effects on genetic diversity back to the evolutionary mechanisms shaping variation in genetic diversity among species.

Based on population genetic theory, the genetic diversity of a species is expected to increase with its census size of reproductive individuals (*15*). In lieu of estimates of census size, which would be unreliable for the majority of taxa included in our study, we included two factors likely to be correlated with census population size: body size and range extent. Body size is expected to be negatively correlated with census size (*72*) and we find that body size has a negative effect on genetic diversity. Conversely, range extent is expected to be positively correlated with census population size (*73, 74*), and, consistent with predictions from theory, we find a positive effect of range extent on diversity (Figure 2). Although neither body size nor range extent is a perfect predictor of population census size, their mean estimated effect sizes are nonetheless consistent with the predicted effects of census size on diversity.

Range extent is expected to be correlated with census size, and thereby with species-wide genetic diversity, but it can also influence a species' genetic diversity via its impact on spatial population genetic processes. Range extent is expected to be positively correlated with genetic diversity (*16, 17*) because lineages separated by a larger distance must wait longer before coalescing, and the geographic distance between a pair of lineages can be larger in a larger area. Thus, larger ranges increase the expected time to most recent common ancestor of a sample and thereby also increase the amount of standing genetic diversity. Ecological complexity within a range is also expected to increase the overall amount of genetic diversity, as ecological heterogeneity and discontinuities in abiotic features and biotic communities may act as a direct barriers to dispersal or decrease effective migration due to selection against locally maladaptive

alleles and linked variation (*75, 76*). However, because the total amount of ecological complexity contained within a range is inextricably linked to the range size (e.g., Figure S2), it is difficult to tease apart the relative contributions of range extent and ecological complexity to total genetic diversity.

Intriguingly, all four dispersal-related traits were estimated to have the opposite of their predicted effects on species-level genetic diversity. Dispersal homogenizes local genetic variation that has arisen due to drift, selection, and mutation; therefore, dispersal is expected to reduce the total amount of genetic differentiation found across all demes in a spatially structured metapopulation (*19*). In lieu of direct estimates of the effective parent-offspring dispersal distance (*77*), we included biological traits related to effective dispersal: planktonicity, planktotrophy, spawning (internal vs. nesting vs. free-spawning), and philopatry. We note that planktonicity and planktotrophy are not perfect predictors of dispersal ability (e.g., even direct developers can disperse long distances (*78*)). However, in general, planktonic dispersal and longer planktonic durations lead to longer dispersal distances (*79, 80*) and there is abundant evidence that the mean potential for dispersal is substantively greater for planktotrophs than non-planktotrophs (*49, 81, 82*).

Planktonicity, planktotrophy, and more dispersive spawning modes were positively correlated with diversity, while philopatry was negatively correlated. Planktonicity and planktotrophy were strongly correlated with each other (Kendall's τ = 0.76), and both were negatively correlated with philopatry (τ = -0.646 and -0.431 respectively; Figure S2), so it is possible that some of the signal in the estimated effects of these predictors are due to those correlations. Nonetheless, we see no evidence for the prediction, supported by theory, that an increase in dispersal leads to lower levels of genetic diversity within a species.

The associations of dispersal-related traits with a species' ability to recover quickly after a local bottleneck might explain why their estimated effects were the opposite of what is predicted

via dispersal's direct effect on diversity. Population bottlenecks will reduce the genetic diversity of a species, to an extent that is determined by their longevity and severity (*20*), so species traits that reduce the prevalence or severity of bottlenecks, or that increase the rate of recovery after a bottleneck, are predicted to increase species-level genetic diversity. Predictors associated with higher dispersal (e.g., planktonicity, planktotrophy, dispersive spawning modes) might lead to faster recovery from local bottlenecks, as extirpated demes may be recolonized more quickly if there is greater connectivity among demes (*82, 83*). And, conversely, philopatry, which decreases the connectivity between demes, might lead to slower recovery from local bottlenecks and decreased genetic diversity (*84, 85*).

Other life history traits are also predicted to affect the capacity of a species to recover from a bottleneck, and thereby genetic diversity. In addition to being highly dispersive, species with higher fecundity are also expected to recover more quickly from bottlenecks and thus to harbor more genetic diversity. In their comparative population genomic analysis, Romiguier *et al.* (*3*) found strong support for this hypothesis, with more fecund species harboring more genetic diversity, as well as those species with smaller propagules. Likewise, the multiple reproductive events of iteroparous species may speed their recovery to local bottlenecks, especially bottlenecks resulting from a generation of failed reproduction. In line with these predictions, we found egg size (which is often inversely correlated with fecundity (*86, 87*)) to have a negative inferred effect on diversity and iteroparity a strong positive one.

Lastly, while dispersal and fecundity are expected to affect species' responses to bottlenecks, geographic position may influence how frequently species experience bottlenecks. Populations closer to the equator are hypothesized to experience greater abiotic stability (*88, 89*) and may therefore be subjected to fewer extreme climatic events and associated bottlenecks. In addition, due to their geological history, lower latitude regions have had more similar abiotic environments for longer (*90*). Both patterns have been proposed as explanations for the latitudinal

gradient in species diversity (*91*) (but see (*92*)). These hypotheses also suggest predictions for *intra*-specific diversity: namely that genetic diversity should be higher in lower latitudes. Consistent with this prediction, we find that species with lower mean range latitudes have higher genetic diversity.

There are a number of important caveats to our results. First, the lack of information on variation in mutation rate across taxa precluded employing mutation rate as a predictor in our models. However, assuming mutation rate is phylogenetically correlated (*93, 94*), variation in genetic diversity across species due to variation in mutation rate will have been captured by the phylogenetic covariance structure of our regression analyses. Second, we focused on deep evolutionary time, yet the lability of many life history traits can be quite high even within families or genera (*95*). Therefore, we hope to see complementary comparative population genomic analyses that focus on more recent phylogenetic scales. Only a minute subset of marine diversity has been studied intensely enough to support such analyses (*96*), but this will change as large-scale datasets become increasingly available. Finally, though we have controlled for phylogenetic dependence on the associations studied, still greater insights could be gained from removing as much of the phylogenetic effect as possible (*97*).

Conclusions

No single factor explained the majority of variation in genetic diversity across taxa (as in, e.g., (*3*)). Instead, we found two factors that were significantly predictive of genetic diversity in the world's oceans; species with larger range extents had higher genetic diversity, as did species with at least one planktonic life-stage. The effects of other predictors, though not significant, suggest that species traits that facilitate greater resilience to, or quicker recovery from, bottlenecks have higher diversity. These results should be digested with the caveat that we cannot rule out the possibility that some portion of the estimated effect sizes for these pre-

dictors may be driven by their correlations with other predictors in our dataset or with other traits not included in our study. Taken together, our work sheds light on the mystery of heterogeneity in genetic variation across species and suggests exciting future avenues of comparative genomic research. Predicting spatial patterns of diversity based on life history traits and spatial distribution represents a significant step forward in both fundamental research and applied conservation.

Materials and methods

Locating genomic datasets

We located and downloaded all georeferenced, publicly available, reduced-representation and whole-genome sequence datasets of marine species present in the International Nucleotide Sequence Database Collaboration (INSDC) as of October 2020. We included an additional 48 datasets for which we retrieved sampling coordinates external to INSDC (*44*). We retained datasets that had $\geq$10 georeferenced individuals and $\geq$2 sampled locations (Figure S1). We excluded samples from non-native ranges, freshwater/inland locales, captive or lab populations, quantitative genetic studies, ancient time, pooled genetic sequences, known species hybrids, and salmonids (due to philopatry to freshwater habitats).

Bioinformatic processing of genomic datasets

Starting from the raw sequence reads, we filtered, cleaned, and assembled each genomic dataset *de novo* using a standardized, custom pipeline and Stacks2 (*98*). Post-Stacks, we dropped low coverage individuals (mean number of genotyped and/or co-genotyped base pairs <30% of median for dataset) and low coverage loci (scored in <50% of remaining individuals). We then used the r80 method to pick the optimal *de novo* assembly for each dataset of nine different standardized Stacks2 parameter combinations tested (*99*); we used the optimal assembly for

each dataset (unique species) to estimate species-level genetic diversity.

Estimating species-level genetic diversity

We estimated species-level genetic diversity for each of the 93 species in this study using a Bayesian parameterization of the Wright-Malécot isolation by distance model (IBD) (*65*). This approach allowed us to generate comparable estimates of genetic diversity across datasets that differed in sampling extent (number of individuals, absolute geographic area, and proportion of species' range). We used pairwise homozygosity between sequenced individuals and pairwise sea-wise distances (the distance in kilometers between sampling locations traveling via water and avoiding land) as model inputs. We ran five independent chains of the Wright-Malécot IBD model per dataset and reported results from the chain with the highest mean log posterior probability (5,000 iterations/chain, 2500 iteration burn-in, chains thinned to every 500 iterations). In all downstream analyses, we used an estimate of π (post burn-in mean of the marginal distribution from the best chain) as our species-level estimate of genetic diversity. This parameter is best understood as long-term diversity; it describes the expectation of the maximum genetic divergence between any pair of samples in a species, regardless of where they are sampled in the species range. This quantity is related to the long-term effective population size, and is determined by the species mutation rate, dispersal rate, and the geometry of its range (*17*).

Collecting biotic, abiotic, and phylogenetic predictors

We collected a suite of biotic traits, abiotic traits, and phylogenetic divergence times for all species in the study to test the extent to which they predict species-level genetic diversity. We selected biotic traits hypothesized to influence levels of genetic diversity via their associations with population size and/or dispersal. These include: iteroparity (whether individuals can mate multiple times before death, or once); philopatry (habitual return to birthplace to mate); spawn-

ing mode (i.e., internal vs. nesting vs. free-spawning, which relate to the degree of parent-progeny dispersal); benthicity (degree to which the adult species is associated with the ocean floor); planktonicity (whether the species has a planktonic stage); planktotrophy (larval feeding behavior, which is related to early-life dispersal and r- vs K-selection); pelagic larval duration (PLD - time spent drifting in the plankton); egg size; and body size. See Appendix S4 for further descriptions of how these biotic traits were defined.

We tested abiotic predictors related to the size and ecological complexity of each species' range, which may influence species-level genetic diversity via associations with population size and levels of genetic divergence. Specifically, we quantified: range extent (maximum linear extent of range); mean range latitude (absolute value of mean latitude of range); and the number of unique ecoregions present within each species' range (Marine Ecoregions of the World (*100*)). We used species occurrence records from the Global Biodiversity Information Facility (GBIF) (*101*) as input for estimating these three abiotic predictors. Finally, we used the January 2022 beta release of TimeTree 5 (*102*) to obtain estimates of the pairwise divergence times between all 93 species in the study. For the 26 species that were not present in Time-Tree, we assigned them randomly to a tip within the clade representing the lowest/shallowest taxonomic level possible or formed a polytomy.

Modeling effects of predictors on genetic diversity

We tested for an effect of phylogenetic relatedness on similarity in π using Blomberg's K (*68*) and Pagel's λ (*67*). We tested for a correlation across evolutionary time between phylogenetic relatedness and similarity in π using a phylocorrelogram (*103*), using a permutation approach to assess significance. We tested hypotheses about the effects of biotic and abiotic predictors on π, using a custom Bayesian phylogenetic multiple regression model with beta-distributed errors (*104*) implemented in Rstan (*105*). To reduce model complexity, avoid issues with mul-

ticollinearity, and increase the interpretability of estimated coefficients, we estimated the effect of each biotic and abiotic factor on π one at a time, in each case incorporating a phylogenetic variance-covariance structure (*106*) as well as four technical "nuisance" predictors: (1) number of individuals in the species' dataset; (2) the mean raw read count; (3) the mean read length; and (4) the mean locus depth. By including these nuisance predictors in each analysis and jointly estimating their effects with those of the biotic and abiotic factors in which we are interested, we ensured that technical artifacts are not having a significant impact on the results of this study. We calculated the Bayesian R^2 (*71*) of each model (including the four nuisance parameters and each of the biological parameters in turn).

Acknowledgments

This research is the product of a working group formed at the National Science Foundation Research Coordination Network for Evolution in Changing Seas Synthesis Workshop (NSF-OCE 1764316, awarded to Katie E. Lotterhos). We thank the authors of Toczydlowski et al. 2021 (*43*) and Crandall et al. 2023 (*44*) for locating geospatial coordinates that were absent in INSDC for 48 of the genomic datasets included in this study. We are grateful to Bruce Martin for writing code to assist with phylogentic analyses and the Spilhaus map projection. Ricardo Lemos also provided code for implementing the Spilhaus projection. Pat Bills, Yaniv Brand-vain, Doc Edge, Steve Goldstein, Morgan Kelly, Katie Lotterhos, Bruce Martin, Misha Matz, Marjorie Weber, Will Wetzel, Ben Winger, Max Witynski, and members of the Bradburd lab all provided helpful feedback and scientific discussion. Computational work was performed using computational resources and services provided by the Institute for Cyber-Enabled Research at Michigan State University. **Funding**: R. H. T. was supported in part by NSF-OCE 1764316 and the U.S. Department of Agriculture, Forest Service. This research was also supported by the National Institute of General Medical Sciences of the National Institutes of Health

References

1. Lewontin RC (1974) *The genetic basis of evolutionary change*. (Columbia University Press New York), pp. xiii, 346 p.

2. Leffler EM, et al. (2012) Revisiting an old riddle: What determines genetic diversity levels within species? *PLOS Biology* 10(9):1–9.

3. Romiguier J, et al. (2014) Comparative population genomics in animals uncovers the determinants of genetic diversity. *Nature* 515(7526):261–263.

4. Buffalo V (2021) Quantifying the relationship between genetic diversity and population size suggests natural selection cannot explain lewontin's paradox. *eLife* 10.

5. Author FA (1992) Conservation genetics and conservation biology: A troubled marriage in *Conservation of Biodiversity for Sustainable Development*, eds. Sandlund OT, Hindar K, Brown AHD. (Oxford University Press), pp. 55–69.

6. Kardos M, et al. (2021) The crucial role of genome-wide genetic variation in conservation. *Proc Natl Acad Sci USA* 118:e2104642118.

7. Webster MS, et al. (2017) Who should pick the winners of climate change? *Trends Ecol Evol* 32:167–173.

8. Stange M, Barrett RDH, Hendry AP (2021) The importance of genomic variation for biodiversity, ecosystems, and people. *Nat Rev Genet* 22:89–105.

9. Mathur S, Mason AJ, Bradburd GS, Gibbs HL (2023) Functional genomic diversity is correlated with neutral genomic diversity in populations of an endangered rattlesnake. *Proceedings of the National Academy of Sciences* 120(43).

10. Ellegren H, Galtier N (2016) Determinants of genetic diversity. *Nature Reviews Genetics* 17(7):422–433.

11. Begun DJ, Aquadro CF (1992) Levels of naturally occurring DNA polymorphism correlate with recombination rates in d. melanogaster. *Nature* 356(6369):519–520.

12. Elyashiv E, et al. (2016) A genomic map of the effects of linked selection in Drosophila. *PLOS Genetics* 12(8):e1006130.

13. Coop G (2016) Does linked selection explain the narrow range of genetic diversity across species? *bioRxiv*.

14. Popovic I, et al. (2024) High germline mutation rates, but not extreme population outbreaks, influence genetic diversity in a keystone coral predator. *PLOS Genetics* 20(2):e1011129.

15. Kimura M, Crow JF (1963) The measurement of effective population number. *Evolution* 17(3):279.

16. Maruyama T (1972) Rate of decrease of genetic variability in a two-dimensional continuous population of finite size. *Genetics* 70(4):639–651.

17. Wilkins JF (2004) A separation-of-timescales approach to the coalescent in a continuous population. *Genetics* 168(4):2227–2244.

18. Slatkin M (1985) Gene flow in natural populations. *Annual Review of Ecology and Systematics* 16:393–430.

19. Kimura M, Weiss GH (1964) The stepping stone model of population structure and the decrease of genetic correlation with distance. *Genetics* 49(4):561–576.

20. Nei M, Maruyama T, Chakraborty R (1975) The bottleneck effect and genetic variability in populations. *Evolution* 29(1):1–10.

21. Pannell JR, Charlesworth B (1999) Neutral genetic diversity in a metapopulation with recurrent local extinction and recolonization. *Evolution* 53(3):664–676.

22. Alcala N, Vuilleumier S (2014) Turnover and accumulation of genetic diversity across large time-scale cycles of isolation and connection of populations. *Proceedings of the Royal Society B: Biological Sciences* 281(1794):20141369.

23. Hines A (1986) Larval problems and perspectives in life histories of marine invertebrates. *Bulletin of Marine Science* 39:506–525.

24. Grosberg R, Cunningham C (2001) Genetic structure in the sea: From populations to communities in *Marine community ecology*, eds. Bertness MD, Gaines SD, Hay ME. (Sinauer Associates Sunderland), pp. 61–84.

25. Marshall D, Morgan S (2011) Ecological and evolutionary consequences of linked life-history stages in the sea. *Current Biology* 21:R718–R725.

26. Figuerola-Ferrando L, et al. (2023) Global patterns and drivers of genetic diversity among marine habitat-forming species. *Global Ecology and Biogeography* 32(7):1218–1229.

27. Pope LC, Riginos C, Ovenden J, Keyse J, Blomberg SP (2015) Population genetic diversity in the australian 'seascape': A bioregion approach. *PLOS ONE* 10(9):e0136275.

28. Wares JP, Strand AE, Sotka EE (2021) Diversity, divergence and density: How habitat and hybrid zone dynamics maintain a genomic cline in an intertidal barnacle. *Journal of Biogeography* 48(9):2174–2185.

29. Pegan TM, et al. (2025) Long-distance seasonal migration to the tropics promotes genetic diversity but not gene flow in boreal birds. *Nature Ecology &; Evolution*.

30. Jablonski D (2008) Species selection: Theory and data. *Annual Reviews in Ecology Evolution and Systematics* 39:501–524.

31. De Kort H, et al. (2021) Life history, climate and biogeography interactively affect worldwide genetic diversity of plant and animal populations. *Nature Communications* 12(1).

32. Clark RD, Pinsky ML (2024) Global patterns of nuclear and mitochondrial genetic diversity in marine fishes. *Ecology and Evolution* 14(5).

33. Hubby JL, Lewontin RC (1966) A molecular approach to the study of genic heterozygosity in natural populations. i. the number of alleles at different loci in drosophila pseudoobscura. *Genetics* 54(2):577–594.

34. Lewontin RC, Hubby JL (1966) A molecular approach to the study of genic heterozygosity in natural populations. ii. amount of variation and degree of heterozygosity in natural populations of drosophila pseudoobscura. *Genetics* 54(2):595–609.

35. Patterson N, Price AL, Reich D (2006) Population structure and eigenanalysis. *PLoS Genetics* 2(12):e190.

36. Birky CW, Maruyama T, Fuerst P (1983) An approach to population and evolutionary genetic theory for genes in mitochondria and chloroplasts, and some results. *Genetics* 103(3):513–527.

37. Rand DM (2001) The units of selection on mitochondrial dna. *Annual Review of Ecology and Systematics* 32(1):415–448.

38. Hemstrom W, Grummer JA, Luikart G, Christie MR (2024) Next-generation data filtering in the genomics era. *Nature Reviews Genetics* 25(11):750–767.

39. Felsenstein J (1985) Phylogenies and the comparative method. *The American Naturalist* 125(1):1–15.

40. Wright S (1943) Isolation by distance. *Genetics* 28(2):114–138.

41. Meirmans PG (2012) The trouble with isolation by distance. *Molecular Ecology* 21(12):2839–2846.

42. Exposito-Alonso M, et al. (2022) Genetic diversity loss in the anthropocene. *Science* 377(6613):1431–1435.

43. Toczydlowski RH, et al. (2021) Poor data stewardship will hinder global genetic diversity surveillance. *Proceedings of the National Academy of Sciences* 118(34).

44. Crandall ED, et al. (2023) Importance of timely metadata curation to the global surveillance of genetic diversity. *Conservation Biology* 37(4):e14061.

45. Chaudhary C, Costello M (2023) Marine species turnover but not richness peaks at the equator. *Progress in Oceanography* 210:102941.

46. Hutchinson G (1961) The paradox of the plankton. *The American Naturalist* 95:137–145.

47. Palumbi S (1994) Genetic divergence, reproductive isolation, and marine speciation. *Annual Reviews in Ecology and Systematics* 25:547–572.

48. Knope ML, Bush AM, Frishkoff LO, Heim NA, Payne JL (2020) Ecologically diverse clades dominate the oceans via extinction resistance. *Science* 367(6481):1035–1038.

49. Wares JP, Gaines S, Cunningham CW (2001) A comparative study of asymmetric migration events across a marine biogeographic boundary. *Evolution* 55:295–306.

50. Ewers-Saucedo C, et al. (2016) The oceanic concordance of phylogeography and biogeography: a case study in *Notochthamalus*. *Ecology and Evolution* 6(13):4403–4420.

51. Álvarez-Noriega M, et al. (2020) Global biogeography of marine dispersal potential. *Nature Ecology &; Evolution* 4(9):1196–1203.

52. McCauley DJ, et al. (2015) Marine defaunation: Animal loss in the global ocean. *Science* 347(6219).

53. SOTKA EE, WARES JP, BARTH JA, GROSBERG RK, PALUMBI SR (2004) Strong genetic clines and geographical variation in gene flow in the rocky intertidal barnaclebalanus glandula. *Molecular Ecology* 13(8):2143–2156.

54. Pringle J, Wares J (2007) Going against the flow: maintenance of alongshore variation in allele frequency in a coastal ocean. *Marine Ecology Progress Series* 335:69–84.

55. Becker BJ, Levin LA, Fodrie FJ, McMillan PA (2007) Complex larval connectivity patterns among marine invertebrate populations. *Proceedings of the National Academy of Sciences* 104(9):3267–3272.

56. Riginos C, Liggins L (2013) Seascape genetics: Populations, individuals, and genes marooned and adrift. *Geography Compass* 7(3):197–216.

57. Sunday JM, Bates AE, Dulvy NK (2012) Thermal tolerance and the global redistribution of animals. *Nature Climate Change* 2(9):686–690.

58. Poloczanska ES, et al. (2013) Global imprint of climate change on marine life. *Nature Climate Change* 3(10):919–925.

59. Poloczanska ES, et al. (2016) Responses of marine organisms to climate change across oceans. *Frontiers in Marine Science* 3.

60. Pinsky ML, Eikeset AM, McCauley DJ, Payne JL, Sunday JM (2019) Greater vulnerability to warming of marine versus terrestrial ectotherms. *Nature* 569(7754):108–111.

61. Boyce DG, et al. (2022) A climate risk index for marine life. *Nature Climate Change* 12(9):854–862.

62. Pinsky M, Palumbi S (2014) Meta-analysis reveals lower genetic diversity in overfished populations. *Molecular Ecology* 23:29–39.

63. Rochman CM, et al. (2015) Anthropogenic debris in seafood: Plastic debris and fibers from textiles in fish and bivalves sold for human consumption. *Scientific Reports* 5(1).

64. Harvell CD, Lamb JB (2020) 141Disease outbreaks can threaten marine biodiversity in *Marine Disease Ecology*, ed. Behringer DC. (Oxford University Press).

65. Hancock ZB, Toczydlowski RH, Bradbur GS (2024) A spatial approach to jointly estimate Wright's neighborhood size and long-term effective population size. *Genetics* p. iyae094.

66. Malécot G (1969) *The Mathematics of Heredity*. (Freeman). Translated from the French edition, 1948.

67. Pagel M (1999) Inferring the historical patterns of biological evolution. *Nature* 401(6756):877–884.

68. Blomberg SP, Garland T, Ives AR (2003) Testing for phylogenetic signal in comparative data: Behavioral traits are more labile. *Evolution* 57(4):717–745.

69. Revell LJ (2024) phytools 2.0: an updated R ecosystem for phylogenetic comparative methods (and other things). *PeerJ* 12:e16505.

70. Spalding MD, et al. (2007) Marine Ecoregions of the World: A Bioregionalization of Coastal and Shelf Areas. *BioScience* 57(7):573–583.

71. Gelman A, Goodrich B, Gabry J, Vehtari A (2019) R-squared for bayesian regression models. *The American Statistician* 73(3):307–309.

72. Brown J (1995) *Macroecology*. (University of Chicago Press).

73. James J, Eyre-Walker A (2020) Mitochondrial DNA Sequence Diversity in Mammals: A Correlation between the Effective and Census Population Sizes. *Genome Biology and Evolution* 12(12):2441–2449.

74. Rosenfield J (2002) Pattern and process in the geographical ranges of freshwater fishes. *Global Ecology & Biogeography* 11:323–332.

75. Wang IJ, Bradburd GS (2014) Isolation by environment. *Molecular Ecology* 23(23):5649–5662.

76. Sobel JM, Chen GF, Watt LR, Schemske DW (2010) The biology of speciation. *Evolution* 64(2):295–315.

77. Bradburd GS, Ralph PL (2019) Spatial population genetics: It's about time. *Annual Review of Ecology, Evolution, and Systematics* 50(1):427–449.

78. Spencer HG, Fraser CI, Poulin E, González-Wevar CA (2025) Towards a more nuanced understanding of long-distance rafting: Case studies from the southern ocean. *Global Ecology and Biogeography* 34(2).

79. Shanks AL (2009) Pelagic larval duration and dispersal distance revisited. *The Biological Bulletin* 216(3):373–385.

80. Pringle JM (2023) Are coastal marine larvae dispersed less than would be expected by ocean currents? *The Biological Bulletin* 245(3):129–138.

81. Chust G, et al. (2016) Dispersal similarly shapes both population genetics and community patterns in the marine realm. *Scientific Reports* 6(1).

82. Schiebelhut LM, Gaylord B, Grosberg RK, Jurgens LJ, Dawson MN (2022) Species' attributes predict the relative magnitude of ecological and genetic recovery following mass mortality. *Molecular Ecology* 31(22):5714–5728.

83. Schiebelhut LM, et al. (2022) Minor genetic consequences of a major mass mortality: short-term effects in *Pisaster ochraceus*. *The Biological Bulletin* 243(3):328–338.

84. Kawecki TJ, Holt RD (2002) Evolutionary consequences of asymmetric dispersal rates. *The American Naturalist* 160:333–347.

85. Waples RS (2025) The idiot's guide to effective population size. *Molecular Ecology*.

86. Marshall DJ, Keough MJ (2007) *The Evolutionary Ecology of Offspring Size in Marine Invertebrates*. (Elsevier), p. 1–60.

87. Kasimatis K, Riginos C (2016) A phylogenetic analysis of egg size, clutch size, spawning mode, adult body size, and latitude in reef fishes. *Coral Reefs* 35:387–397.

88. Willig MR, Kaufman DM, Stevens RD (2003) Latitudinal gradients of biodiversity: pattern, process, scale, and synthesis. *Annual review of ecology, evolution, and systematics* 34(1):273–309.

89. Rohde K (1992) Latitudinal gradients in species diversity: the search for the primary cause. *Oikos* pp. 514–527.

90. Brodie JF, Mannion PD (2023) The hierarchy of factors predicting the latitudinal diversity gradient. *Trends in Ecology &; Evolution* 38(1):15–23.

91. Fine P (2015) Ecological and evolutionary drivers of geographic variation in species diversity. *Annu.Rev.Ecol.Evol.Syst* 46:369–392.

92. Chaudhary C, Saeedi H, Costello MJ (2016) Bimodality of latitudinal gradients in marine species richness. *Trends in Ecology & Evolution* 31(9):670–676.

93. Drummond AJ, Suchard MA, Xie D, Rambaut A (2012) Bayesian phylogenetics with beauti and the beast 1.7. *Molecular Biology and Evolution* 29(8):1969–1973.

94. Ramos-Almodovar F, Gao Z, Voight BF, Mathieson I (2025) Methylation-associated mutagenesis underlies variation in the mutation spectrum across eukaryotes. *bioRxiv*.

95. Hart MW, Byrne M, Smith MJ (1997) Molecular phylogenetic analysis of life-history evolution in asterinid starfish. *Evolution* 51(6):1848–1861.

96. Dietrich MR, Ankeny RA, Crowe N, Green S, Leonelli S (2020) How to choose your research organism. *Studies in History and Philosophy of Science Part C: Studies in History and Philosophy of Biological and Biomedical Sciences* 80:101227.

97. Zakas C, Deutscher JM, Kay AD, Rockman MV (2018) Decoupled maternal and zygotic genetic effects shape the evolution of development. *eLife* 7:e37143.

98. Rochette NC, Rivera-Colón AG, Catchen JM (2019) Stacks 2: Analytical methods for paired-end sequencing improve radseq-based population genomics. *Molecular ecology* 28(21):4737–4754.

99. Paris JR, Stevens JR, Catchen JM (2017) Lost in parameter space: a road map for stacks. *Methods in Ecology and Evolution* 8(10):1360–1373.

100. Spalding MD, et al. (2007) Marine Ecoregions of the World: A Bioregionalization of Coastal and Shelf Areas. *BioScience* 57(7):573–583.

101. GBIF: The Global Biodiversity Information Facility (2020) What is GBIF? Available from https://www.gbif.org/what-is-gbif.

102. Kumar S, et al. (2022) Timetree 5: an expanded resource for species divergence times. *Molecular biology and evolution* 39(8):msac174.

103. Keck F, Rimet F, Bouchez A, Franc A (2016) phylosignal: an r package to measure, test, and explore the phylogenetic signal. *Ecology and Evolution* 6(9):2774–2780.

104. Ferrari S, Cribari-Neto F (2004) Beta regression for modelling rates and proportions. *Journal of Applied Statistics* 31(7):799–815.

105. Stan Development Team (2024) RStan: the R interface to Stan. R package version 2.32.5.

106. Garland, Jr T, Ives AR (2000) Using the past to predict the present: confidence intervals for regression equations in phylogenetic comparative methods. *The American Naturalist* 155(3):346–364.

107. R Core Team (2023) *R: A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, Vienna, Austria).

108. Winter DJ (2017) rentrez: An R package for the NCBI eUtils API. *The R Journal* 9(2):520–526.

109. WoRMS 2021 Editorial Board (2021) World Register of Marine Species (WoRMS). Available from https://www.marinespecies.org.

110. Holstein J (2018) worms: Retriving aphia information from world register of marine species. *R package version 0.2.2*.

111. Korunes KL, Samuk K (2021) pixy: Unbiased estimation of nucleotide diversity and divergence in the presence of missing data. *Molecular ecology resources* 21(4):1359–1368.

112. Kodama Y, Shumway M, Leinonen R (2012) The sequence read archive: explosive growth of sequencing data. *Nucleic acids research* 40(D1):D54–D56.

113. Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet. journal* 17(1):10–12.

114. Battey CJ, Ralph PL, Kern AD (2020) Space is the Place: Effects of Continuous Spatial Structure on Analysis of Population Genetic Data. *Genetics* 215(1):193–214.

115. Bradburd GS, Coop GM, Ralph PL (2018) Inferring Continuous and Discrete Population Genetic Structure Across Space. *Genetics* 210(1):33–52.

116. Exposito-Alonso M, et al. (2022) Genetic diversity loss in the anthropocene. *Science* 377(6613):1431–1435.

117. Amante C, Eakins BW (2009) *ETOPO1 arc-minute global relief model: procedures, data sources and analysis*. (National Oceanic and Atmospheric Administration).

118. Pante E, Simon-Bouhet B (2013) marmap: a package for importing, plotting and analyzing
bathymetric and topographic data in r. *PLoS one* 8(9):e73051.

119. Strathmann MF (1987) *Reproduction and Development of Marine Invertebrates of the*
Northern Pacific Coast: Data and Methods for the Study of Eggs, Embryos, and Larvae.
(University of Washington Press).

120. R. Froese, D. Pauly, Editors (2023) FishBase. World Wide Web electronic publication.
Available from https://www.fishbase.org, Accessed July 2021–December 2023.

121. M. L. D. Palomares, D. Pauly. Editors (2023) SeaLifeBase. World Wide Web electronic
publication. Available from https://www.sealifebase.org, Accessed July 2021–December
2023.

122. Cowen RK, Sponaugle S (1997) Relationships between early life history traits and recruit-
ment among coral reef fishes in *Early Life History and Recruitment in Fish Populations*,
eds. Chambers RC, Trippel EA. (Springer Netherlands, Dordrecht), pp. 423–449.

123. Chamberlain S, Ram K, Barve V, Mcglinn D, Chamberlain MS (2017) rgbif: Interface to
the global biodiversity information facility api. *R package version 3.5.0.*

124. Costello MJ, et al. (2017) Marine biogeographic realms and species endemicity. *Nature*
communications 8(1):1057.

125. Paradis E, Schliep K (2019) ape 5.0: an environment for modern phylogenetics and evo-
lutionary analyses in R. *Bioinformatics* 35:526–528.

126. Gelman A, Meng XL, Stern H (1996) Posterior predictive assessment of model fitness via
realized discrepancies. *Statistica sinica* pp. 733–760.

 # Figures

Figure 1. Species-level genetic diversity varies by orders of magnitude in the world's oceans. (A) Time-calibrated phylogeny of all species (*N* = 93) and map of all samples (*N* = 9,430) included in this study. The map uses the Spilhaus projection which, by emphasizing the oceans and their connectivity, rather than land masses, represents the world as a marine organism might see it. (B) Set of discrete (gray-scale) and continuous (blue-scale) biotic and abiotic predictors collated for each species. Colors represent the trait state or value for each species. (C) Species-level genetic diversity estimated using a spatially explicit model and genomic sequence data.

Figure 2. The estimated effects of biotic and abiotic predictors on species' genetic diversity vary in their magnitude and direction. Estimated effect sizes (presented via their marginal posterior distributions) are standardized by the standard deviation of their predictor and are ordered from top to bottom by mean effect on species-level genetic diversity. Significant predictors – those for which the 95% equal-tailed credible interval of the marginal distribution of effect sizes did not overlap zero (gray, vertical dashed line) - are denoted with an asterisk.

Figure 3. Species-level genetic diversity is most significantly predicted by species range extent and the presence of a planktonic life stage. (A) Species with larger range extents have higher genetic diversity. Shaded polygons represent 50%, 80%, and 95% model prediction credible intervals. Mean Bayesian R^2 = 0.55 (95%CI: 0.29-0.76). (B) Species that have at least one planktonic life stage have higher genetic diversity than species that are never planktonic. Mean Bayesian R^2=0.56 (95%CI: 0.34-0.75). Each point represents a unique species (N = 93) and is colored by taxonomic clade (as in Figure 1).

Supporting Information for

Drivers of genetic diversity across the marine tree of life

Rachel H. Toczydlowski, Reid S. Brennan, Eric D. Crandall, Joanna L. Kelley, James M. Pringle, Cynthia Riginos, John P. Wares, and Gideon S. Bradburd

To whom correspondence may be addressed: bradburd@umich.edu or rhtoczydlowski@gmail.com.

This PDF file includes:

Supplementary Methods

Supplementary Appendices S1 to S6

Figs. S1 to S5

Supplementary Methods

Locating genomic datasets

We searched the International Nucleotide Sequence Database Collaboration (INSDC) to locate publicly available genomic-level sequence data for marine organisms. We accessed the INSDC from the United States National Center for Biotechnology Information (NCBI) data center using R (*107*) and the *rentrez* R package (v1.2.3, (*108*)). We used the entrez_search function to locate entries in INSDC and entrez_summary and/or entrez_fetch to download the associated metadata. We first identified all non-human, non-viral, non-metagenomic, and non-bacterial datasets (Bio-Projects) that were focused on DNA sequencing and had publicly available sequence data. Next, we downloaded all metadata for the individuals in each of these datasets from the Sequence Read Archive (SRA) and BioSample databases. We retained sequences with library strategies: WGS, WCS, WGA, RAD-Seq, or OTHER. We then queried the remaining organism names (assigned in the BioSample database) in the World Register of Marine Species (*109*) to identify marine organisms. We used the *worms* R package (v0.2.2, (*110*)) with no fuzzy name matching to execute this query.

We then filtered the remaining potentially marine genomic datasets to those that included at least 10 unique individuals (BioSamples) and 2 unique collection locations (based on the geospatial coordinates provided in the BioSamples database). Finally, we manually dropped samples (or datasets) from non-native ranges, captive or lab populations, quantitative genetic studies, ancient time, and pooled genetic sequences. We also dropped known species hybrids, salmonids (due to philopatry to freshwater habitats), and freshwater/inland samples. In cases where there were multiple genomic datasets for a species, we retained the dataset with the broadest geographic sampling extent (as our goal was to estimate species-level genetic diversity). We manually recovered latitude/longitude from associated scientific publications for 48 datasets that met all of our filtering criteria but lacked spatial coordinates in INSDC (*44*); we dropped those where latitude/longitude were unrecoverable. We performed this search strategy on October 28, 2019 and again on October 19, 2020. See Appendix S1 for the exact search strategy and Appendix S2 for the citations of all 93 genomic datasets used in this study.

Bioinformatic processing of genomic datasets

We started with the raw reads published in the SRA for each relevant dataset, then cleaned, filtered, and assembled them into loci using a standardized bioinformatic pipeline. Applying the same bioinformatic steps to all datasets allowed us to make meaningful comparisons across datasets. Furthermore, many studies associated with the publication of these datasets only published diversity metrics (and/or derived files) for polymorphic loci. As a result, we could not calculate absolute measures of diversity without re-assembling the reads to obtain the number of both polymorphic and nonpolymorphic base pairs sequenced (*111*). The following steps were all performed on the Michigan State Institute for Cyber-Enabled Research High Performance Computing Cluster (https://icer.msu.edu/).

We downloaded raw reads from the SRA using the fasterq-dump function in the SRA Toolkit v2.10.7 (*112*). We treated all paired-end datasets as single-end datasets, retaining all forward reads (regardless of whether they had a reverse pair) and discarding all reverse reads (SRA Tools fasterq-

dump –split-files option). Next, we used the Stacks2 process_radtags module v2.54 (*98*) to remove reads with uncalled bases and/or where the mean Phred score in a sliding window 15% of the read length dropped below 15. After removing low quality reads, we used cutadapt v2.1 (*113*) to search for and remove adapter sequences (e.g. the Illumina common adapter) from the 3' end of reads. See Appendix S3 for the complete list of next-generation sequence adapters that we searched for. Datasets with adapters were easy to identify by the presence of adapter hits in >25% of the reads (compared to ~5% of reads in clean datasets). About a fifth of the datasets (20 of 93) contained adapters. We confirmed that no adapter sequences remained in cleaned, trimmed reads by running a final adapter search. Lastly, we trimmed all reads to a standardized length within each dataset; *de novo* loci assembly is more robust when all raw reads are a uniform length within each dataset (*98*). For datasets with variable length reads, we selected a uniform read length for each dataset by visually assessing the distribution of read lengths for the dataset and choosing a length that retained long reads in a majority of samples.

We assembled each dataset *de novo* using Stacks2 v2.54 (*98*). For each dataset, we identified a set of optimal assembly parameters following guidelines from (*99*) and (*98*). We used a custom parallel workflow to assemble each dataset with 9 different parameter combinations of *M* (ustacks) and *n* (cstacks): $\{M = 3, n = 2\}$, $\{M = 3, n = 3\}$, $\{M = 3, n = 4\}$, $\{M = 6, n = 5\}$, $\{M = 6, n = 6\}$, $\{M = 6, n = 7\}$, $\{M = 9, n = 8\}$, $\{M = 9, n = 9\}$, and $\{M = 9, n = 10\}$. These are the parameters that most strongly influence assembly behavior (*98*). The *M* parameter controls the number of mismatches allowed between reads within individuals and *n* controls the number of mismatches allowed between loci among individuals. We used default values for all other parameters.

For each assembled dataset and parameter combination, we dropped individuals with a mean number of genotyped and/or co-genotyped base pairs <30% of the median for the dataset (including nonpolymorphic and polymorphic sites). These individuals with few final assembled loci also commonly had few raw reads (suggesting they were poorly sequenced, rather than poorly assembled). We then dropped any loci that were scored in less than 50% of the remaining individuals. We performed this filtering after the populations module using the populations.samples.fa and populations.snps.vcf Stacks output files and custom R scripts. Finally, we selected the assembly parameter combination (of the nine standardized combinations tested) that yielded the highest number of polymorphic loci scored in at least 80% of individuals for each dataset (r80 method, (*99*)). The r80 method has been demonstrated to identify the set of assembly parameters that yield the maximum number of conservatively plausible polymorphic loci for a genomic dataset.

Estimating species-level genetic diversity

Estimates of species-level genetic diversity can be heavily influenced by the number of individuals sampled, where in the species' range those samples are located, and what proportion of the species' range is sampled. Such biases result from spatial genetic structure (i.e., correlations between genetic ancestry and geographic location) within a species' range (*114–116*). To generate unbiased estimates of diversity from datasets with heterogeneous sampling, we used the Bayesian parameterization of the Wright-Malécot model of isolation by distance (IBD) (*40, 66*) implemented in (*65*) for each of the 93 species in our study. Briefly, this method implements the theoretical model of isolation by distance introduced by Wright (*40*) and Malécot (*66*) to describe the decay of genetic relatedness between individuals as an asymptotic function of the geographic

distance separating them. The asymptote – a parameter estimated by the model-fitting procedure – captures the mean maximum genetic divergence between individuals of a species, regardless of their geographic separation (*65*). In Hancock *et al.*, the estimate of the value of this asymptote is designated as collecting phase π, or π_c, in reference to the collecting phase of the spatial coalescent (*17*); for simplicity, we refer to it as π throughout this manuscript. Our estimate of species-level genetic diversity is informative about long-term inbreeding N_e and relatively insensitive to recent demography (*65*). Because this approach explicitly models the rate of isolation-by-distance, our estimates of species-level diversity are robust to both the number of samples and their spatial distribution (and comparable across studies with heterogeneous sampling schemes).

To calculate species-level genetic diversity, the Hancock et al. (*65*) model implementation requires as inputs – a matrix of pairwise homozygosity (identity-by-state) and a corresponding matrix of pairwise geographic distances between genotyped individuals. We calculated pairwise homozygosity as 1-D_{XY} using custom R scripts following (*65*); these scripts accounted for missing genotypes by defining the denominator for D_{XY} as the total number of base pairs co-genotyped for each pair of individuals (within a dataset). We calculated pairwise geographic distance as the distance between each pair of individuals traveling via the sea (avoiding crossing land). We used the National Oceanic and Atmospheric Administration ETOPO1 global bathymetric data with a resolution of 4 arc-minutes (*117*) and the *trans.mat* and *lc.dist* marmap functions of the *marmap* R package v1.0.6 (*118*) to calculate these sea-wise distances. For individuals that fell on land (e.g., near coastlines), we used the *dist2isobath* marmap function with isobath = 0 to relocate each "land point" to the closest point in water. We calculated each sea-wise distance using two resistance matrices (one with longitudes from -180 to 180, the other with longitudes from 0 to 360) and retained the shorter distance.

For each species dataset, we then ran five independent chains of the model in (*65*) for 5,000 iterations each. We discarded the first 2,500 chains as burn-in and sampled the remaining chain ever 500th iteration to reduce autocorrelation. We discarded chains that mixed poorly and identified the single best chain for each species as that with the highest mean log posterior probability. We used the mean of the marginal distribution of π from the best chain for each dataset as our estimate of species-level genetic diversity in downstream analyses.

To quantify variation in diversity across populations within species, we also calculated π for all individuals sampled at each unique geographic location included in the sampling dataset. We report the standard deviation of those location-specific estimates of π in Figure S4, and we show the relationship between that metric variance in diversity and our global estimates of diversity for each species in Figure S5. The comparison between variation in location-specific estimates of diversity and the species global estimate shows a positive correlation, which is an artifact of the mean-variance relationship.

Collecting biotic, abiotic, and phylogenetic predictors

Biotic predictors   We focused on nine biotic traits related to reproduction and dispersal: philopatry, iteroparity, spawning, egg size, benthicity, planktonicity, planktotrophy, pelagic larval duration [PLD], and body size. We selected these traits because they have demonstrated associations with intraspecific spatial genetic diversity and divergence or are expected to be correlated with genetic diversity via their associations with population size and/or dispersal. We defined each discrete trait (i.e., all but PLD, egg size, and body size, which are continuous) as "present", "intermediate", or

"absent" for each species (Figure 1).

Specifically, we defined iteroparity as "present" for species that can breed multiple times per lifetime, and "absent" for species that can only breed once. We defined philopatry as "present" for species where individuals return to their natal breeding area to reproduce and "absent" for species that do not perform such migrations. We defined three categories for spawning to capture variation in how far embryos disperse from parents. Embryos can be internally fertilized and remain attached to the parent ("absent"), laid in nests or other benthically attached structures that are often (but not always) dissociated from the parents ("intermediate"), or free-floating, typically the product of broadcast spawning ("present"). Note this is a distinct life history facet from the presence or absence of planktonic larvae (see below). We defined the trait benthicity to characterize the degree to which species have a relationship with the ocean floor (or other fixed location). Benthicity was "present" for species that maintain a strong connection to the bottom throughout their lifetime (e.g., hydrozoans and anemonefishes that lack planktonic larvae), "absent" for species that are never associated with the bottom (e.g., dolphins that give birth at sea to swimming young), and "intermediate" for species that have both benthic and pelagic life stages (e.g., fish with planktonic larvae and benthic adults). We used planktonicity to distinguish species that have at least one planktonic – floating or drifting – life stage ("present") from those that are never planktonic ("absent").

Planktotrophy is a predictor of the amount of time larvae spend drifting in the water column feeding, which is related to the level of provisioning provided by the egg as well as the capacity for external feeding. We coded planktotrophy as "absent" for species that lack a (planktonic) larval stage and instead develop directly from embryo into young that resemble miniature adults. We coded planktotrophy as "intermediate" for species that produce planktonic larvae that feed on yolk from well-provisioned eggs to facilitate metamorphosis into juveniles (i.e., lecithotrophic larvae); as these larvae are planktonic, they drift in the water column while developing, but develop relatively quickly given the well-provisioned eggs. Lastly, we coded planktotrophy as "present" for species that produce planktonic larvae that must feed on particulates in the water column to fuel metamorphosis into juveniles given they derive from minimally-provisioned eggs (i.e., planktotrophic species); development usually takes longer than planktonic larvae derived from well-provisioned eggs (*119*). We defined pelagic larval duration (PLD) as the mean (or midpoint) number of days that marine larvae spend in the plankton. Finally, we defined egg size as the largest dimension of the egg at hatching (or progeny at birth) and body size as the largest dimension of an adult individual. Extended trait definitions are available in Appendix S4.

When possible, we used trait values reported in FishLifeBase (*120*), SeaLifeBase (*121*), and large compilations (e.g., (*122*) for reef fishes). For most values, however, we performed Boolean searches in Web of Science or Google Scholar (e.g., <species name> AND egg size). We were able to obtain trait values for $> 90\%$ of the species in our study for all traits except egg size (72% coverage across species; Figure S8). All species had values for $> 77\%$ of the nine biotic traits (Figure S9). Full citations for all biotic trait values are listed in Appendix S5.

Abiotic predictors We used three abiotic predictors to characterize the geographic extent and amount of ecological complexity encompassed by each species' range – range extent, mean range latitude, and the number of ecoregions encompassed within the species' range. We estimated these three quantities for each species using species occurrence records from the Global Biodiver-

sity Information Facility (GBIF) (*101*). We first downloaded all GBIF presence records for each species that were not fossils, not managed, and did not have identified geospatial issues using the occ_download() function of the *rgbif* R package (v3.5.0, (*123*)) on September 8, 2021 (see Appendix S6 for exact search parameters and full citations for all downloaded GBIF data). We then visually examined maps of the occurrence records returned for each species and manually removed spatial outliers (that were unlikely to be correctly identified records) and observations within non-native ranges. We used a combination of literature, expert opinion, and published range maps to decide which additional GBIF occurrence records, if any, to manually drop for each species.

We defined the range extent as the longest axis of the species' range "as the fish swims", that is, traveling via the sea and avoiding crossing land. For each species, we calculated all pairwise "sea-wise distances" between the final, cleaned GBIF occurrence records using the same approach that we employed for the genotyped individuals (see *Estimating species-level genetic diversity*). We then used the value at the 97.5^{th} percentile of the resulting pairwise "sea-wise distance" distance matrix (between species occurrence records) as the final range extent; using the 97.5^{th} percentile minimized the influence of any one individual occurrence record, especially those near the range edges. For computational efficiency, we thinned occurrence datasets that had more than 15,000 points using a raster proportional to the spatial extent of the dataset before generating the pairwise "sea-wise distance" distance matrix.

We defined mean range latitude as the absolute value of the mean latitude of all the final, cleaned GBIF occurrence records for each species. We used original locations in this calculation, (rather than relocating "land points" to the nearest ocean), but mean latitudes were highly correlated between the two approaches (Pearson's $R = 0.98$). We did not include the genetic sample locations in the range extent or distance from equator calculations because they were nested within the spatial extent of, and often duplicated in, the GBIF occurrence records.

Finally, we characterized the amount of ecological complexity contained within each species' range by calculating the total number of unique Marine Ecoregions of the World (MEOW) ecoregions that at least one GBIF occurrence record or genetic sample fell within (*70, 100*). We refer to this predictor as "number of ecoregions". MEOW ecoregions are biogeographic units, developed in large part to facilitate conservation priority setting and planning, whose boundaries reflect synethetic discontinuities in taxonomic assemblages, dominant habitats, ocean currents and temperature, and geomorphological features (*70*). MEOW ecoregions classify coastal and shelf systems, where the majority of (known) marine species diversity resides (*70*). Note, the number of ecoregions was highly correlated with the spatially coarser MEOW realms and provinces as well as the Costello et al. (*124*) (Pearson's R all > 0.87). Transitions between ecoregions may act as barriers to gene flow within a taxon, so even though a given species may not be found in every ecoregion contained within their range, the number of ecoregions may nonetheless be predictive of total population genetic diversity of the species.

Phylogenetic relationships   We derived estimates of the pairwise divergence times between all species in the study using the January 2022 beta release of TimeTree 5 (*102*). We first identified species that were present in TimeTree (67/93 species in study). When a focal species was not in TimeTree but other closely related species were (more closely related than any other species in our study, based on published taxonomy), we assigned the focal species randomly to a tip within the clade representing the lowest/shallowest taxonomic level possible (15 species assigned to a

representative tip in their genus, 1 species assigned at level of family). This approach is sensible because any tip within the focal clade will give the same divergence time to species outside of the clade, however, it assumes that the oldest species defining the focal clade is present in TimeTree. Divergence times for focal species in clades where this assumption is not met will be upwardly biased. Ten focal species were missing from TimeTree and shared a clade with other species in this study (meaning we could not assign them to a random tip within the clade). For 3 of these 10 species, we were able to identify the most closely related species present in TimeTree (using published literature) and use this tip. For the remaining 7 species absent from TimeTree, we added a tip to the tree that formed a polytomy with the lowest taxonomic level of classification present in TimeTree.

Phylogenetic Comparative Analyses

We evaluated the strength and significance of the phylogeny in predicting patterns of similarity in diversity using three complementary approaches. We first estimated the overall significance of phylogenetic structure in genetic diversity using Blomberg's K (68) and Pagel's λ (67), both implemented with the function *phylosig* in the R package *phytools* 2.0 (69). We tested for a correlation between phylogenetic relatedness and similarity in π using the R package *phylosignal* 1.3.1 (103). We determined significance of correlation through time using a permutation approach: we first randomly permuted tip diversity values and, for each permutation, we estimated the correlogram using the *phylosignal* package. We repeated this permutation approach 500 times and used the equal-tailed 95% confidence interval of these "null" simulations to determine the significance of the observed correlogram for diversity.

Modeling effects of predictors on genetic diversity

To test hypotheses about the effects of biotic and abiotic predictors on π, we used a custom Bayesian phylogenetic multiple regression model with Beta-distributed errors (104) implemented in Rstan v2.32.3 (105). Specifically, we used the mean/precision parameterization of the Beta distribution shown below:

$$\pi \sim \text{Beta}(a, b) \tag{1}$$

$$a = \mu \times \phi \tag{2}$$

$$b = (1 - \mu) \times \phi \tag{3}$$

$$\mu_i = \frac{1}{1 + \exp(-\theta_i)} \tag{4}$$

$$\vec{\theta} \sim \mathcal{N}\left(\gamma + \sum_M \beta_m \vec{X}_m \ , \ \alpha \Psi\right) \tag{5}$$

where a and b are the shape parameters of the Beta distribution, and μ and ϕ function as the mean and precision. The mean of the Beta distribution, μ, is given by the inverse-logit transformed parameter θ, which is modeled as a draw from a multivariate normal distribution. The mean of that multivariate normal distribution is given by the expression $\gamma + \sum_M \beta_m \vec{X}_m$, where γ is a

global mean of θ, M is the number of predictor variables, $\vec{X}_m$ is the vector of predictor variable values for the mth predictor, and β_m is the effect size of the mth predictor. The covariance of the multivariate normal distribution on θ is given by $\alpha\Psi$, where Ψ is a phylogenetic correlation matrix (i.e. a phylogenetic variance-covariance matrix scaled so that all values fall between 0 and 1), and α is the estimated Brownian rate parameter (i.e. the macroevolutionary heritability) (*106*). We generated the phylogenetic variance-covariance matrix from our ultrametric phylogeny using the R package *ape* v5.0 (*125*).

The priors on our parameters were: $\alpha \sim \mathcal{N}(0, 10)$, $\phi \sim \text{Exp}(\mathbb{V}[\pi], 10)$, $\gamma \sim \mathcal{N}(\log(\mathbb{E}[\pi]/(1 - \mathbb{E}[\pi])), 10)$, and $\beta \sim \mathcal{N}(0, 10)$, where $\mathbb{V}[\pi]$ and $\mathbb{E}[\pi]$ denote the empirical variance and mean, respectively, of observed π across species. Each model contained one focal predictor of interest, the phylogenetic variance-covariance matrix, and four nuisance predictors (number of individuals in the species dataset; the mean raw read count; the mean read length; and the mean locus depth).

For each model, we included all species that were scored for the focal trait of interest (excluding species with missing data or for which the trait was not applicable). We ran each MCMC chain for 20,000 iterations, discarding the first half of the iterations as burn-in and thinning the remaining samples by retaining every 40th iteration. We assessed model convergence by examining parameter trace plots, marginal distributions, and joint marginal distributions. For each predictor discussed in the text, we report the 95% equal-tailed credible interval from the thinned, burned-in marginal distribution.

We assessed the significance of each parameter by determining whether the 95% equal-tailed credible interval contained the value 0; if it did, the parameter was deemed not significant. To compare effect sizes between predictors, we standardized them by multiplying the estimated effect size by the standard deviation of the predictor. In order to evaluate model adequacy (i.e., the ability of our model to describe the data), we used posterior predictive sampling (*126*). For each model run, we randomly sampled 500 MCMC iterations, with replacement, from the burned-in, thinned, posterior distribution. For each sampled iteration, we recorded the estimated parameter values for the shape parameters of the Beta distribution used to model our response variable a and b. We then simulated draws from a Beta distribution parameterized by a_i and b_i for sampled iteration i. Those posterior predicted values of π were compared to the observed value of π for each species to determine whether the model as a whole was effectively capturing patterns in the data, as well as to determine whether diversity for specific species was well or poorly predicted by our model (Figure S7).

Because range extent and number of ecoregions were both significant predictors of genetic diversity (Figure 2), but were also highly correlated (Fig S2), we ran an additional analysis to try to get closer to ascertaining causality. Number of ecoregions is necessarily correlated with range extent, so we were concerned that the significance of ecoregions was being driven by its correlation with range extant. To test whether this suspicion was correct, we analyzed diversity as a function of number of ecoregions *per* unit range extent. The mean estimated effect size of this hybrid predictor was -0.003, with a 95% equal-tailed posterior credible interval of -0.0072 – 0.0003 which was not significantly different from zero. As a result, we cannot conclusively rule out the possibility that the effect of ecoregions on diversity is entirely being driven by the correlation of ecoregions with range extent, and we thus drop ecoregions from further investigation.

To assess the explanatory power of each model, we calculate a Bayesian R^2 following the recommendations of Gelman *et al.* (2017) (*71*). Specifically, we work in the 'link' space of the model and, for each sampled iteration in the MCMC, we calculate the ratio of the variance in the

fitted means to the sum of the variance of fitted means and residual means. For a given iteration s, this is as follows:

$$\mathbb{E}[\theta_s] = \gamma_s + \sum_M \beta_{sm}\vec{X}_m \qquad (6)$$

$$\mathbb{V}_{\text{fit}} = \text{Var}\left(\mathbb{E}[\theta_s]\right) \qquad (7)$$

$$\mathbb{V}_{\text{resid}} = \text{Var}\left(\theta_s - \mathbb{E}[\theta_s]\right) \qquad (8)$$

$$R_s^2 = \frac{\mathbb{V}_{\text{fit}}}{\mathbb{V}_{\text{fit}} + \mathbb{V}_{\text{resid}}} \qquad (9)$$

Table S1. Biotic and abiotic predictors of species-level genetic diversity included in analysis. "Trait values" describes the range of possible values species could have for each predictor. "Significant" describes whether the effect size of the predictor was inferred to be significantly different from zero. The final column gives the mean (unstandardized) effect size of the predictor, as well as the lower and upper bounds of the 95% equal-tailed credible interval of the marginal distribution of the effect size.

Predictor	Trait values	Significant	Mean effect (95% CI)
Benthicity	absent/intermediate/present	no	-4.401e-02 (-3.683e-01 – 2.881e-01)
Body size	continuous	no	-5.366e-02 (-1.588e-01 – 4.708e-02)
Ecoregions	integer	yes	7.627e-03 (-1.954e-04 – 1.154e-02)
Egg size	continuous	no	4.396e-04 (-1.228e-02 – 9.218e-03)
Iteroparity	absent/present	no	-3.795e-01 (-9.386e-01 – 2.721e-01)
Latitude	continuous	no	-1.066e-03 (-1.302e-02 – 6.846e-03)
Pelagic larval duration	continuous	no	1.314e-03 (-4.466e-03 – 3.453e-03)
Philopatry	absent/present	no	-4.665e-01 (-9.773e-01 – 4.253e-02)
Planktonicity	absent/present	yes	5.625e-01 (-5.532e-02 – 1.169e+00)
Planktotrophy	absent/intermediate/present	no	1.650e-01 (-9.903e-02 – 4.216e-01)
Spawning	absent/intermediate/present	no	1.071e-01 (-7.655e-02 – 2.765e-01)
Range extent	continuous	yes	5.291e-01 (5.943e-02 – 9.515e-01)

Figure S1. Spatial distribution and sampling intensity of sequenced individual marine organisms included in this study (N = 9,430 total). Colors represent the number of sequenced individuals within each grid square. Map is in the Spilhaus projection.

Figure S2. Kendall rank correlations (using pairwise complete observations) between all biotic and abiotic predictors analyzed. Blue and red colors indicate positive and negative correlations, respectively, and color intensity is proportional to the absolute value of the correlation.

Figure S3. Species-level genetic diversity estimated for each species included in analysis. Species diversity estimates are colored by broad taxonomic group and sorted by value.

Figure S4. Standard deviation of diversity estimates across unique locations for each species included in analysis. The standard deviations of the species diversity estimates are colored by broad taxonomic group and sorted by value. Values are plotted on a symlog scale.

Figure S5. Standard deviation of diversity estimates across unique locations for each species (symlog scale) plotted against the total diversity estimate for that species (log scale). Points colored by broad taxonomic group.

Figure S6. Phylogenetic correlogram showing the estimated correlation between phylogenetic relatedness and species-level genetic diversity at different temporal depths.

Figure S7. Relationship between range extent and species-level genetic diversity. Points are colored by broad taxonomic group as in Figure 1. Vertical gray lines show the 95% equal-tailed quantile of the posterior predictive distribution of estimates for diversity for each species.

Figure S8. Trait coverage for each of the nine biotic traits included in this study; coverage calculated as a percentage across all 93 species in the study. Note the *x*-axis starts at 70% (rather than 0). The vertical blue dashed line denotes 80%, a commonly accepted conservative threshold for trait coverage. (Note the three abiotic predictors were available for all datasets in this study.)

Figure S9. Biotic trait coverage for each species in the study (*N* = 93) as a percentage. There are nine biotic traits total. Note the *x*-axis starts at 70% (rather than 0). Species are ordered alphabetically by Latin name. The vertical blue dashed line denotes 80%, a commonly accepted conservative threshold for trait coverage.

Appendix S1: Search strategy used to locate relevant genomic datasets in the INSDC

This appendix details the search strategy that we used to locate relevant genomic datasets in the International Nucleotide Sequence Database Collaboration (INSDC). Note that we accessed the INSDC from the North American portal – the National Center for Biotechnology Information (NCBI). We list the general types of filters that we applied, in the order we applied them, and the number (and type) of records remaining after each filtering step in our pipeline. We also detail the exact NCBI variable names and values that we used in our search and filtering pipeline to locate genomic datasets. NCBI uses BioProjects to denote a study or dataset, BioSamples to refer to individual organisms sampled, and SRA sequence records to refer to the genetic sequence data derived from a BioSample.

After the final filtering step listed below – passing species names to the World Register of Marine Species – we identified and dropped BioProjects (datasets) with fewer than 10 BioSamples (individuals) and 2 collection locations using R. We then curated the remaining potential datasets by hand. We dropped known species hybrids, salmonids, freshwater/inland samples, samples from locations not part of the native species' range, and samples from ancient time. We also dropped studies that were focused on specific organelles (e.g., chloroplast) and quantitative genetic studies. Final sequence samples were included if they had collection latitude and longitude, the raw FASTA file downloaded without errors, the FASTA sequence files were at the level of sampled individual (not pooled sequencing), and were not outliers with respect to read count.

Overview of filters applied and number of records retained after each filtering step:

$\Rightarrow$ 184,155 total BioProjects in INSDC on 10/19/2020

Search BioProject database for all BioProjects that are:
   Non-human, non-viral, non-metagenomic, and non-bacterial
   $\Rightarrow$ 32162 BioProjects returned

Remove BioProjects related to:
   Gene expression, RefSeq, targeted loci, metagenomic, metabarcoding, exome, microbial
   $\Rightarrow$ 19,495 BioProjects retained

Download metadata from the Sequence Read Archive (SRA) for all sequence records in each of the above
BioProjects.
   $\Rightarrow$ 948,446 SRA sequence records returned

Retain SRA records with library strategies:
   WGS, WCS, WGA, RAD-Seq, OTHER
   (removing: AMPLICON, ATAC-seq, Bisulfite-Seq, ChIA-PET, ChIP-Seq, CLONE, CLONEEND,
   CTS, DNase-Hypersensitivity, EST, FAIRE-seq, Hi-C, FINISHING, FL-cDNA, MBD, MeDIP,
   miRNA-Seq, MNase-Seq, MRE-Seq, ncRNA-Seq, POOLCLONE, RIP-Seq, RNA-Seq, SELEX,
   Synthetic-Long-Read, Targeted-Capture, TetheredChromatinConformationCapture, Tn-Seq, WXS)
   $\Rightarrow$ 523,537 SRA sequence records retained

Merge retained SRA records to the BioProject records using PRJ BioProject accession numbers.
Remove BioProjects that did not return any SRA records matching the above specified library

 strategies.
⇒ 7,314 BioProjects retained

Use the NCBI common tree online tool to view all taxonomic IDs assigned to the current list of sequence records. Remove sequences with a taxonomic identification of:
Synthetic, metagenome, unidentified, bacteria, environmental, viral

Download metadata from the BioSample database associated with each of the above sequence records.
⇒ 505,932 SRA sequences from
⇒ 432,400 BioSamples from
⇒ 6,755 BioProjects retained representing
⇒ 18,556 purported species

Retain records with species names present in the World Register of Marine Species (WoRMS).
⇒ 114,865 SRA sequences from
⇒ 100,118 BioSamples from
⇒ 1,336 BioProjects retained representing
⇒ 1,738 purported species

Exact variables and terms used to locate and filter INSDC datasets:

The exact search terms and filters that we used to identify the INSDC datasets analyzed in this publication follow. We use "!=" below to denote "does not equal". Bold headers refer to the NCBI databases, quoted terms denote variables/fields of metadata defined by NCBI within said database, and text beneath quoted terms denotes controlled vocabulary values of the specified variable.

Initial BioProject search:

```
"scope multiisolate"[Filter] OR "scope multispecies"[Filter] OR
"scope other"[Filter] NOT "org human"[Filter] NOT "org archaea"[Filter]
NOT "org viruses"[Filter] NOT "org bacteria"[Filter] NOT "metagenome"[Filter]
NOT "targeted locus loci"[Filter] NOT "clone ends"[Filter]
NOT "metagenomic assembly"[Filter] NOT "transcriptome gene expression"[Filter]
NOT "proteome"[Filter] NOT "epigenomics"[Filter] NOT "exome"[Filter]
NOT "bioproject protein"[Filter] NOT "bioproject gds"[Filter]
NOT "phenotype genotype"[Filter] NOT "metagenome"[All Fields] NOT "map"[Filter]
```

BioProject filters:

```
"project_data_type" !=

RefSeq Transcriptome or Gene expression
Targeted loci cultured
RefSeq Genome
RefSeq Genome sequencing and assembly
RefSeq Other
RefSeq Targeted Locus (Loci)

"project_target_material" !=
```

```
Phenotype
Proteome
Purified Chromosome
Reagent
Transcriptome
Targeted loci environmental
```

```
"project_target_capture" !=

Clone Ends
Exome
```

```
"organism_name" does not contain:

metagenome
```

```
"project_title" does not contain:

metagenome, Metagenome, Metagenomic, metagenomic, metabarcoding
transcriptome, Transcriptome, Transcriptomic
RNA
16S, 16s, 18S, 18s
CRISPR
gene expression, Gene Expression, Gene expression"
viral
microbiome, Microbiome, microbial, microbiota
bacteria, Bacteria
methylation
Plasmodium falciparum
```

SRA filters:

```
"library_strat" =
RAD-Seq
WCS
WGA
WGS
OTHER
```

```
"scientific_name" !=
metagenome, Metagenome
```

The below taxonomic IDs represent synthetic DNA, environmental DNA, metagenomic DNA, and bacterial DNA that we identified using the NCBI phylogenetic common tree online tool.

```
"taxid" !=
```

```
2293429
1440148
32630
1427524
81077
32644
496923
77133
669196
2282120
668369
727
2590021
28448
272943
2021969
527802
477819
562
227
83333
316407
703612
224308
1229511
1126252
1126251
```

Taxonomy filters:

```
"division" !=
Environmental samples
Bacteria
Viruses
```

BioSample filters:

```
"organism" =
# species returned by
worms::wormsbynames(match = FALSE, marine_only = TRUE)
```

We filtered remaining records to retain BioProjects with at least 10 BioSample IDs and at least two unique georeferenced sampling locations per species/BioProject combination (dataset). Finally, we manually dropped samples (or entire datasets) that were located in non-native areas of the species' ranges, were sampled from captive or lab individuals, and those derived from quantitative genetic studies, ancient time, or pooled genetic sequences. We also dropped known species hybrids, salmonids, and samples collected from freshwater or inland locations.

Appendix S2: Citations for genomic datasets

This appendix contains citations for the scientific papers associated with the genetic sequence data utilized in this study. The BioProject accession number where the genetic data are archived in the National Center for Biotechnology Information (NCBI) is also included with each citation. Note some papers include multiple BioProjects and some genomic datasets (BioProjects) have multiple paper citations.

Citations

1. Kelly S Andrews, Krista M Nichols, Anna Elz, Nick Tolimieri, Chris J Harvey, Robert Pacunski, Dayv Lowry, K Lynne Yamanaka, and Daniel M Tonnes. Cooperative research sheds light on population structure and listing status of threatened and endangered rockfish species. *Conservation Genetics*, 19:865–878, 2018. *Genetic sequence data archived under NCBI BioProject Accession:* PRJNA451040.

2. Emily S. Bellis, Reid. B. Edlund, Hazel K. Berrios, Harilaos A. Lessios, and Dee R. Denver. Molecular signatures of host specificity linked to habitat specialization in *Exaiptasia* sea anemones. *Ecology and Evolution*, 8(11):5413–5426, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA394157.

3. Yacine Ben Chehida, Julie Thumloup, Cassie Schumacher, Timothy Harkins, Alex Aguilar, Asunción Borrell, Marisa Ferreira, Lorenzo Rojas-Bracho, Kelly M Robertson, Barbara L Taylor, et al. Mitochondrial genomics reveals the evolutionary history of the porpoises (Phocoenidae) across the speciation continuum. *Scientific Reports*, 10(1):15190, 2020. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA659918.

4. Laura Benestan, Thierry Gosselin, Charles Perrier, Bernard Sainte-Marie, Rémy Rochette, and Louis Bernatchez. RAD genotyping reveals fine-scale genetic structuring and provides powerful population assignment in a widely distributed marine species, the American lobster (*Homarus americanus*). *Molecular Ecology*, 24(13):3299–3315, 2015. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA281764.

5. Laura Benestan, Brady K. Quinn, Halim Maaroufi, Martin Laporte, Fraser K. Clark, Spencer J. Greenwood, Rémy Rochette, and Louis Bernatchez. Seascape genomics provides evidence for thermal adaptation and current-mediated population structure in American lobster (*Homarus americanus*). *Molecular Ecology*, 25(20):5073–5092, 2016. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA281764.

6. Giacomo Bernardi, Peter Nelson, Michelle Paddack, John Rulmal Jr, and Nicole Crane. Genomic islands of divergence in the yellow tang and the brushtail tang surgeonfishes. *Ecology and Evolution*, 8(17):8676–8685, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA449377.

7. Pim Bongaerts, Cynthia Riginos, Ramona Brunner, Norbert Englebert, Struan R Smith, and Ove Hoegh-Guldberg. Deep reefs are not universal refuges: reseeding potential varies among

coral species. *Science Advances*, 3(2):e1602373, 2017. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA361144.

8. Elisa Boscari, Marco Abbiati, Fabio Badalamenti, Giorgio Bavestrello, Lisandro Benedetti-Cecchi, Rita Cannas, Angelo Cau, Carlo Cerrano, Giovanni Chimienti, Federica Costantini, Simonetta Fraschetti, Gianmarco Ingrosso, Ilaria A. M. Marino, Francesco Mastrototaro, Chiara Papetti, Marta Paterno, Massimo Ponti, Lorenzo Zane, and Leonardo Congiu. A population genomics insight by 2b-RAD reveals populations' uniqueness along the Italian coastline in *Leptopsammia pruvoti* (Scleractinia, Dendrophylliidae). *Diversity and Distributions*, 25(7):1101–1117, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA521302.

9. Kristina M. Cammen, Thomas F. Schultz, W. Don Bowen, Michael O. Hammill, Wendy B. Puryear, Jonathan Runstadler, Frederick W. Wenzel, Stephanie A. Wood, and Michael Kinnison. Genomic signatures of population bottleneck and recovery in Northwest Atlantic pinnipeds. *Ecology and Evolution*, 8(13):6599–6614, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA450328.

10. E. Sally Chang, Maria E. Orive, and Paulyn Cartwright. Nonclonal coloniality: Genetically chimeric colonies through fusion of sexually produced polyps in the hydrozoan *Ectopleura larynx*. *Evolution Letters*, 2(4):442–455, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA420851.

11. Julie C Chow, Paul E Anderson, and Andrew M Shedlock. Sea turtle population genomic discovery: Global and locus-specific signatures of polymorphism, selection, and adaptive potential. *Genome Biology and Evolution*, 11(10):2797–2806, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA559677.

12. Gemma V. Clucas, Jane L. Younger, Damian Kao, Louise Emmerson, Colin Southwell, Barbara Wienecke, Alex D. Rogers, Charles-André Bost, Gary D. Miller, Michael J. Polito, Patrick Lelliott, Jonathan Handley, Sarah Crofts, Richard A. Phillips, Michael J. Dunn, Karen J. Miller, and Tom Hart. Comparative population genomics reveals key barriers to dispersal in Southern Ocean penguins. *Molecular Ecology*, 27(23):4680–4697, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA343012, PRJNA493660.

13. Lila Colston-Nepali, Anna Tigano, Brian Boyle, and Vicki Friesen. Hybridization does not currently pose conservation concerns to murres in the Atlantic. *Conservation Genetics*, 20(6):1465–1470, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA564451.

14. David J. Combosch, Sarah Lemer, Peter D. Ward, Neil H. Landman, and Gonzalo Giribet. Genomic signatures of evolution in *Nautilus* - an endangered living fossil. *Molecular Ecology*, 26(21):5923–5938, 2017. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA369717.

15. Brendan H. Cornwell. Gene flow in the anemone *Anthopleura elegantissima* limits signatures of local adaptation across an extensive geographic range. *Molecular Ecology*, 29(14):2550–

2566, 2020. *Genetic sequence data archived under NCBI BioProject Accession*: PR-JNA632874.

16. Nicole L Crane, Juliette Tariel, Jennifer E Caselle, Alan M Friedlander, D Ross Robertson, and Giacomo Bernardi. Clipperton Atoll as a model to study small marine populations: Endemism and the genomic consequences of small population size. *PLoS One*, 13(6):e0198901, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA448430, PRJNA448438.

17. Hayley M. DeHart, Leocadio Blanco-Bercial, Mollie Passacantando, Jennifer M. Questel, and Ann Bucklin. Population connectivity of chaetognath *Eukrohnia hamata*. *National Center for Biotechnology Information*, BioProject Accession: PRJNA576675, Registration date: 09-October-2019.

18. M. Lisette Delgado, Konrad Górski, Evelyn Habit, and Daniel E. Ruzzante. The effects of diadromy and its loss on genomic divergence: The case of amphidromous *Galaxias maculatus* populations. *Molecular Ecology*, 28(24):5217–5231, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA564121.

19. James L. Dimond, Sanoosh K. Gamblewood, and Steven B. Roberts. Genetic and epigenetic insight into morphospecies in a reef coral. *Molecular Ecology*, 26(19):5031–5042, 2017. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA379028.

20. R Fernández, M Schubert, AM Vargas-Velázquez, A Brownlow, GA Víkingsson, U Siebert, LF Jensen, N Øien, D Wall, E Rogan, et al. A genomewide catalogue of single nucleotide polymorphisms in white-beaked and Atlantic white-sided dolphins. *Molecular Ecology Resources*, 16(1):266–276, 2016. *Genetic sequence data archived under NCBI BioProject Accession*: PR-JNA280898.

21. María José Frugone, María Eugenia López, Nicolás I. Segovia, Theresa L. Cole, Andrew Lowther, Pierre Pistorius, Gisele P.M. Dantas, Maria Virginia Petry, Francesco Bonadonna, Phil Trathan, Andrea Polanowski, Barbara Wienecke, Ke Bi, Cynthia Y. Wang-Claypool, Jonathan M. Waters, Rauri C.K. Bowie, Elie Poulin, and Juliana A. Vianna. More than the eye can see: Genomic insights into the drivers of genetic differentiation in Royal/Macaroni penguins across the Southern Ocean. *Molecular Phylogenetics and Evolution*, 139:106563, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA523574.

22. Michelle R. Gaither, Moisés A. Bernal, Richard R. Coleman, Brian W. Bowen, Shelley A. Jones, W. Brian Simison, and Luiz A. Rocha. Genomic signatures of geographic isolation and natural selection in coral reef fishes. *Molecular Ecology*, 24(7):1543–1557, 2015. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA371502.

23. Michelle R. Gaither, Darren J. Coker, Samuel Greaves, Fatih Sarigol, Samuel D. Payet, Veronica Chaidez, Tane H. Sinclair-Taylor, Joseph D. DiBattista, and Michael L. Berumen. Does color matter? Molecular and ecological divergence in four sympatric color morphs of a coral reef fish. *Ecology and Evolution*, 10(18):9663–9681, 2020. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA640135.

24. Michelle R Gaither, Georgios A Gkafas, Menno De Jong, Fatih Sarigol, Francis Neat, Thomas Regnier, Daniel Moore, Darren R Gröcke, Neil Hall, Xuan Liu, et al. Genomics of habitat choice and adaptive evolution in a deep-sea fish. *Nature Ecology & Evolution*, 2(4):680–687, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA430030.

25. Genomic Resources Development Consortium, P. Álvarez, Wolfgang Arthofer, Maria M. Coelho, D. Conklin, A. Estonba, Ana R. Grosso, S. J. Helyar, J. Langa, Miguel P. Machado, I. Montes, Joana Pinho, Alexander Rief, Manfred Schartl, Birgit C. Schlick-Steiner, Julia Seeber, Florian M. Steiner, and C. Vilas. Genomic resources notes accepted 1 June 2015 – 31 July 2015. *Molecular Ecology Resources*, 15(6):1510–1512, 2015. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA253681.

26. Lani U. Gleason and Ronald S. Burton. Genomic evidence for ecological divergence against a background of population homogeneity in the marine snail *Chlorostoma funebralis*. *Molecular Ecology*, 25(15):3557–3573, 2016. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA286949.

27. Fanny L Gonzalez-Zapata, Pim Bongaerts, Catalina Ramirez-Portilla, Boahemaa Adu-Oppong, Gretchen Walljasper, Alejandro Reyes, and Juan A Sanchez. Holobiont diversity in a reef-building coral over its entire depth range in the mesophotic zone. *Frontiers in Marine Science*, 5:29, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA385083.

28. Konrad Górski, EM Habit, MA Pingram, and AJ Manosalva. Variation of the use of marine resources by *Galaxias maculatus* in large Chilean rivers. *Hydrobiologia*, 814:61–73, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA564121.

29. Alison L Gould and Paul V Dunlap. Shedding light on specificity: population genomic structure of a symbiosis between a coral reef fish and luminous bacterium. *Frontiers in Microbiology*, 10:492715, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA385011.

30. Ray Hilborn. Are MPAs effective? *ICES Journal of Marine Science*, 75(3):1160–1162, 08 2017. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA508986.

31. Richard G. J. Hodel, L. Lacey Knowles, Stuart F. McDaniel, Adam C. Payton, Jordan F. Dunaway, Pamela S. Soltis, and Douglas E. Soltis. Terrestrial species adapted to sea dispersal: Differences in propagule dispersal of two Caribbean mangroves. *Molecular Ecology*, 27(22):4612–4626, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA480308, PRJNA480409.

32. Jennifer A. Hoey and Malin L. Pinsky. Genomic signatures of environmental selection despite near-panmixia in summer flounder. *Evolutionary Applications*, 11(9):1732–1747, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA477007.

33. Sho Hosoya, Shotaro Hirase, Kiyoshi Kikuchi, Kusuto Nanjo, Yohei Nakamura, Hiroyoshi Kohno, and Mitsuhiko Sano. Random pcr-based genotyping by sequencing technology GRAS-Di (genotyping by random amplicon sequencing, direct) reveals genetic structure of man-

grove fishes. *Molecular Ecology Resources*, 19(5):1153–1163, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJDB7819.

34. Peter J Hundt, Emma B Liddle, Stuart V Nielsen, Brendan J Pinto, and Tony Gamble. Sex chromosomes and sex-specific molecular markers in Indo-Pacific combtooth blennies (Blenni-idae, Istiblennius). *Marine Ecology Progress Series*, 627:195–200, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA553554.

35. Eeva Jansson, Francois Besnier, Ketil Malde, Carl André, Geir Dahle, and Kevin A Glover. Genome wide analysis reveals genetic divergence between Goldsinny wrasse populations. *BMC Genetics*, 21:1–15, 2020. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA508986.

36. Sheila A Kitchen, Aakrosh Ratan, Oscar C Bedoya-Reina, Richard Burhans, Nicole D Fogarty, Webb Miller, and Iliana B Baums. Genomic variants among threatened *Acropora* corals. *G3 Genes|Genomes|Genetics*, 9(5):1633–1646, 05 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA473816.

37. Ljerka Lah, Daronja Trense, Harald Benke, Per Berggren, Þorvaldur Gunnlaugsson, Christina Lockyer, Ayaka Öztürk, Bayram Öztürk, Iwona Pawliczka, Anna Roos, et al. Spatially explicit analysis of genome-wide SNPs detects subtle population structure in a mobile marine mammal, the harbor porpoise. *PloS One*, 11(10):e0162792, 2016. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA343959.

38. Philip Lavretsky, Jeffrey L. Peters, Kevin Winker, Volker Bahn, Irina Kulikova, Yuri N. Zhu-ravlev, Robert E. Wilson, Chris Barger, Kirsty Gurney, and Kevin G. McCracken. Becoming pure: Identifying generational classes of admixed individuals within lesser and greater scaup populations. *Molecular Ecology*, 25(3):661–674, 2016. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA299372.

39. Sabrina Le Cam, Claire Daguin-Thiébaut, Sarah Bouchemousse, Aschwin H. Engelen, Nova Mieszkowska, and Frédérique Viard. A genome-wide investigation of the worldwide invader *Sargassum muticum* shows high success albeit (almost) no genetic diversity. *Evolutionary Applications*, 13(3):500–514, 2020. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA549138.

40. A. Le Moan, P.-A. Gagnaire, and F. Bonhomme. Parallel genetic divergence among coastal–marine ecotype pairs of European anchovy explained by differential introgression after secondary contact. *Molecular Ecology*, 25(13):3187–3202, 2016. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA311981.

41. Yu-Long Li, Dong-Xiu Xue, Bai-Dong Zhang, and Jin-Xian Liu. Population genomic signatures of genetic structure and environmental selection in the catadromous roughskin sculpin *Trachidermus fasciatus*. *Genome Biology and Evolution*, 11(7):1751–1764, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA360511.

42. N. S. Locatelli and J. A. Drew. Population structure and clonal prevalence of scleractinian corals (*Montipora capitata* and *Porites compressa*) in Kaneohe Bay, Oahu. *bioRxiv*, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA544861.

43. Derek P. Manzello, Mikhail V. Matz, Ian C. Enochs, Lauren Valentino, Renee D. Carlton, Graham Kolodziej, Xaymara Serrano, Erica K. Towle, and Mike Jankulak. Role of host genetics and heat-tolerant algal symbionts in sustaining populations of the endangered coral *Orbicella faveolata* in the Florida Keys with ocean warming. *Global Change Biology*, 25(3):1016–1031, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA508589.

44. Edith Martinez, Vincent Buonaccorsi, John R. Hyde, and Andres Aguilar. Population genomics reveals high gene flow in grass rockfish (*Sebastes rastrelliger*). *Marine Genomics*, 33:57–63, 2017. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA316872.

45. Morten Mattingsdal, Sissel Jentoft, Ole K. Tørresen, Halvor Knutsen, Michael M. Hansen, Joana I. Robalo, Zuzanna Zagrodzka, Carl André, and Enrique Blanco Gonzalez. A continuous genome assembly of the corkwing wrasse (*Symphodus melops*). *Genomics*, 110(6):399–403, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA354496.

46. Morten Mattingsdal, Per Erik Jorde, Halvor Knutsen, Sissel Jentoft, Nils Christian Stenseth, Marte Sodeland, Joana I. Robalo, Michael M. Hansen, Carl André, and Enrique Blanco Gonzalez. Demographic history has shaped the strongly differentiated corkwing wrasse populations in Northern Europe. *Molecular Ecology*, 29(1):160–171, 2020. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA354496.

47. Paolo Momigliano, Henri Jokinen, Antoine Fraimout, Ann-Britt Florin, Alf Norkko, and Juha Merilä. Extraordinarily rapid speciation in a marine fish. *Proceedings of the National Academy of Sciences*, 114(23):6074–6079, 2017. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA382467.

48. Iratxe Montes, Darrell Conklin, Aitor Albaina, Simón Creer, Gary R. Carvalho, María Santos, and Andone Estonba. SNP discovery in European anchovy (*Engraulis encrasicolus*, L) by high-throughput transcriptome and genome sequencing. *PLoS ONE*, 8(8):e70051, 2013. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA253681.

49. Marta Paterno, Marcello Schiavina, Giorgio Aglieri, Jamila Ben Souissi, Elisa Boscari, Renato Casagrandi, Aurore Chassanite, Mariachiara Chiantore, Leonardo Congiu, Giuseppe Guarnieri, Claudia Kruschel, Vesna Macic, Ilaria A. M. Marino, Chiara Papetti, Tomaso Patarnello, Lorenzo Zane, and Paco Melià. Population genomics meet Lagrangian simulations: Oceanographic patterns and long larval duration ensure connectivity among *Paracentrotus lividus* populations in the Adriatic and Ionian seas. *Ecology and Evolution*, 7(8):2463–2479, 2017. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA373825.

50. Joseph B. Pfaller, Adam C. Payton, Karen A. Bjorndal, Alan B. Bolten, and Stuart F. McDaniel. Hitchhiking the high seas: Global genomics of rafting crabs. *Ecology and Evolution*, 9(3):957–974, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA471559.

51. L. V. Plough. Population genomic analysis of the blue crab *Callinectes sapidus* using genotyping-by-sequencing. *Journal of Shellfish Research*, 36(1):249–261, 2017. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA340157.

52. D. S. Portnoy, J. B. Puritz, C. M. Hollenbeck, J. Gelsleichter, D. Chapman, and J. R. Gold. Selection and sex-biased dispersal in a coastal shark: the influence of philopatry on adaptive variation. *Molecular Ecology*, 24(23):5877–5885, 2015. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA286089.

53. Gregory N. Puncher, Alessia Cariani, Gregory E. Maes, Jeroen Van Houdt, Koen Herten, Rita Cannas, Naiara Rodriguez-Ezpeleta, Aitor Albaina, Andone Estonba, Molly Lutcavage, Alex Hanke, Jay Rooker, James S. Franks, Joseph M. Quattro, Gualtiero Basilone, Igaratza Fraile, Urtzi Laconcha, Nicolas Goñi, Ai Kimoto, David Macías, Francisco Alemany, Simeon Deguara, Salem W. Zgozi, Fulvio Garibaldi, Isik K. Oray, Firdes Saadet Karakulak, Noureddine Abid, Miguel N. Santos, Piero Addis, Haritz Arrizabalaga, and Fausto Tinti. Spatial dynamics and mixing of bluefin tuna in the Atlantic Ocean and Mediterranean Sea revealed using next-generation sequencing. *Molecular Ecology Resources*, 18(3):620–638, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA432036.

54. Gregory Neils Puncher, Sherrylynn Rowe, George A. Rose, Nathalie M. Leblanc, Geneviève J. Parent, Yanjun Wang, and Scott A. Pavey. Chromosomal inversions in the Atlantic cod genome: Implications for management of Canada's Northern cod stock. *Fisheries Research*, 216:29–40, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA528403.

55. Catherine M Purcell, Arun S Seetharam, Owyn Snodgrass, Sofia Ortega-García, John R Hyde, and Andrew J Severin. Insights into teleost sex determination from the *Seriola dorsalis* genome assembly. *BMC Genomics*, 19:1–11, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA319656.

56. Jonathan B Puritz, John R Gold, and David S Portnoy. Fine-scale partitioning of genomic variation among recruits in an exploited fishery: Causes and consequences. *Scientific Reports*, 6(1):36095, 2016. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA329407.

57. Hannah G. Reich, Sheila A. Kitchen, Kathryn H. Stankiewicz, Meghann Devlin-Durante, Nicole D. Fogarty, and Iliana B. Baums. Genotypic similarity among algal symbionts corresponds to associations with closely related coral hosts. *bioRxiv*, 2020. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA473816.

58. Pablo Saenz-Agudelo, Joseph D. Dibattista, Marek J. Piatek, Michelle R. Gaither, Hugo B. Harrison, Gerrit B. Nanninga, and Michael L. Berumen. Seascape genetics along environmental gradients in the Arabian Peninsula: Insights from ddRAD sequencing of anemonefishes. *Molecular Ecology*, 24(24):6241–6255, 2015. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA294760.

59. EM Salas, G Bernardi, Michael L Berumen, MR Gaither, and LA Rocha. RADseq analyses reveal concordant Indian Ocean biogeographic and phylogeographic boundaries in the reef fish

Dascyllus trimaculatus. Royal Society Open Science, 6(5):172413, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA590458.

60. Lauren M Schiebelhut, Jonathan B Puritz, and Michael N Dawson. Decimation by sea star wasting disease and rapid genetic change in a keystone species, *Pisaster ochraceus*. *Proceedings of the National Academy of Sciences*, 115(27):7069–7074, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA445895.

61. Aaron B. A. Shafer, Claire R. Peart, Sergio Tusso, Inbar Maayan, Alan Brelsford, Christopher W. Wheat, and Jochen B. W. Wolf. Bioinformatic processing of RAD-seq data dramatically impacts downstream population genetic inference. *Methods in Ecology and Evolution*, 8(8):907–917, 2017. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA348572.

62. Katherine Silliman. Population structure, genetic connectivity, and adaptation in the Olympia oyster (*Ostrea lurida*) along the west coast of North America. *Evolutionary Applications*, 12(5):923–939, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA511386.

63. Ingrid Spies, Kristen M. Gruenthal, Daniel P. Drinan, Anne B. Hollowed, Duane E. Stevenson, Carolyn M. Tarpey, and Lorenz Hauser. Genetic evidence of a northward range expansion in the eastern Bering Sea stock of Pacific cod. *Evolutionary Applications*, 13(2):362–375, 2020. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA558810.

64. Ingrid Spies, James W Orr, Duane E Stevenson, Pamela Goddard, Gerald Hoff, Jared Guthridge, Myles Hollowed, and Christopher Rooper. Skate egg nursery areas support genetic diversity of Alaska and Aleutian skates in the Bering Sea. *Marine Ecology Progress Series*, 669:121–138, 2021. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA646172.

65. William R. Stovall, Helen R. Taylor, Michael Black, Stefanie Grosser, Kim Rutherford, and Neil J. Gemmell. Genetic sex assignment in wild populations using genotyping-by-sequencing data: A statistical threshold approach. *Molecular Ecology Resources*, 18(2):179–190, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA419445.

66. Joshua A. Thia, Katrina McGuigan, Libby Liggins, Will F. Figueira, Christopher E. Bird, Andrew Mather, Jennifer L. Evans, and Cynthia Riginos. Genetic and phenotypic variation exhibit both predictable and stochastic patterns across an intertidal fish metapopulation. *Molecular Ecology*, 30(18):4392–4414, 2021. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA595396.

67. Laura Timm, Joan A Browder, Shaina Simon, Thomas L Jackson, Ian C Zink, and Heather D Bracken-Grissom. A tree money grows on: the first inclusive molecular phylogeny of the economically important pink shrimp (Decapoda: *Farfantepenaeus*) reveals cryptic diversity. *Invertebrate Systematics*, 33(2):488–500, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA554161.

68. Laura E Timm, Lys M Isma, Matthew W Johnston, and Heather D Bracken-Grissom. Comparative population genomics and biophysical modeling of shrimp migration in the Gulf of Mexico reveals current-mediated connectivity. *Frontiers in Marine Science*, 7:19, 2020. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA553831.

69. L.E. Timm, H.D. Bracken-Grissom, A. Sosnowski, M. Breitbart, M. Vecchione, and H. Judkins. Population genomics of three deep-sea cephalopod species reveals connectivity between the Gulf of Mexico and northwestern Atlantic Ocean. *Deep Sea Research Part I: Oceanographic Research Papers*, 158:103222, 2020. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA553831.

70. L.E. Timm, H.D. Bracken-Grissom, A. Sosnowski, M. Breitbart, M. Vecchione, and H. Judkins. Population genomics of three deep-sea cephalopod species reveals connectivity between the Gulf of Mexico and northwestern Atlantic Ocean. *Deep Sea Research Part I: Oceanographic Research Papers*, 158:103222, 2020. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA573759.

71. Benjamin M. Titus, Paul D. Blischak, and Marymegan Daly. Genomic signatures of sympatric speciation with historical and contemporary gene flow in a tropical anthozoan (Hexacorallia: Actiniaria). *Molecular Ecology*, 28(15):3572–3586, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA542967.

72. Felipe Torquato, Pedro Range, Radhouane Ben-Hamadou, Eva E. Sigsgaard, Philip F. Thomsen, Rodrigo Riera, Michael L. Berumen, John A. Burt, David A. Feary, Alyssa Marshell, Daniele D'Agostino, Joseph D. DiBattista, and Peter R. Møller. Consequences of marine barriers for genetic diversity of the coral-specialist yellowbar angelfish from the Northwestern Indian Ocean. *Ecology and Evolution*, 9(19):11215–11226, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA559876.

73. Madeleine J. H. van Oppen, Pim Bongaerts, Pedro Frade, Lesa M. Peplow, Sarah E. Boyd, Hieu T. Nim, and Line K. Bay. Adaptation to reef habitats through selection on the coral animal and its associated microbiome. *Molecular Ecology*, 27(14):2956–2971, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA450871.

74. Felix Vaux, Leif K. Rasmuson, Lisa A. Kautzi, Polly S. Rankin, Matthew T. O. Blume, Kelly A. Lawrence, Sandra Bohn, and Kathleen G. O'Malley. Sex matters: Otolith shape and genomic variation in deacon rockfish (*Sebastes diaconus*). *Ecology and Evolution*, 9(23):13153–13173, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA560239.

75. Patricia H. Wepfer, Yuichi Nakajima, Makamas Sutthacheep, Veronica Z. Radice, Zoe Richards, Put Ang, Tullia Terraneo, Mareike Sudek, Atsushi Fujimura, Robert J. Toonen, Alexander S. Mikheyev, Evan P. Economo, and Satoshi Mitarai. Evolutionary biogeography of the reef-building coral genus *Galaxea* across the Indo-Pacific Ocean. *Molecular Phylogenetics and Evolution*, 151:106905, 2020. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA576132.

76. Shengyong Xu, Na Song, Linlin Zhao, Shanshan Cai, Zhiqiang Han, and Tianxiang Gao. Genomic evidence for local adaptation in the ovoviviparous marine fish *Sebastiscus marmoratus* with a background of population homogeneity. *Scientific Reports*, 7(1):1562, 2017. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA359404.

77. Tyler S Zemlak, Evelyn M Habit, Sandra J Walde, Cecilia Carrea, and Daniel E Ruzzante. Surviving historical Patagonian landscapes and climate: molecular insights from *Galaxias maculatus*. *BMC Evolutionary Biology*, 10:1–18, 2010. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA564121.

78. Bai-Dong Zhang, Dong-Xiu Xue, Yu-Long Li, and Jin-Xian Liu. RAD genotyping reveals fine-scale population structure and provides evidence for adaptive divergence in a commercially important fish from the Northwestern Pacific Ocean. *PeerJ*, 7:e7242, 2019. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA533707.

79. Yunfeng Zhao, Wenzhu Peng, Huayang Guo, Baohua Chen, Zhixiong Zhou, Jian Xu, Dianchang Zhang, and Peng Xu. Population genomics reveals genetic divergence and adaptive differentiation of Chinese sea bass (*Lateolabrax maculatus*). *Marine Biotechnology*, 20:45–59, 2018. *Genetic sequence data archived under NCBI BioProject Accession*: PRJNA356786.

Appendix S3: List of sequencing adapters used in sequence read cleaning

Below is a list of all adapter sequences that we searched for on the 3' end of all raw sequence reads downloaded from the International Nucleotide Sequence Database Collaboration. Lines that start with > denote adapter names and the directly succeeding line contains the adapter sequence.

We generated the below list from:
https://github.com/stephenturner/adapters/blob/master/adapters_combined_152_unique.fasta;
FastQC (https://github.com/s-andrews/FastQC/blob/master/Configuration/contaminant_list.txt);
and BBMap (https://github.com/BioInfoTools/BBMap/blob/master/resources/adapters.fa);
all accessed on April 9, 2020.

We removed duplicate adapters among the above sources and merged adapters with highly similar sequences using IUPAC base pair ambiguity (names of merged adapters begin with "MERGED"). Using the program cutadapt, we searched each FASTQ sequence file in our study for each of the below sequences. Datasets that contained adapters were easy to distinguish from those that did not; datasets with adapters returned hits in >25% of the reads per individual compared to <5% of reads per individual in clean datasets. When present, we removed identified adapter sequences from all reads in the dataset. We then performed a final adapter search to ensure no adapters remained in raw reads (i.e., all sequences below returned hits in $\sim$ 5% or fewer of the sequence reads per individual across the dataset).

```
>Adapter1 / aka PhiX_read1_adapter aka the Illumina common adapter
AGATCGGAAGAGCGGTTCAGCAGGAATGCCGAGACCGATCTCGTATGCCGTCTTCTGCTTG
>MERGED-Primers_and_dimers-GROUP
AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTMRWKMDRMRGMSVSSWYMVVVRBBHWY
VCBVWBWCCSWWCWCGWMKSYCKTCYKMTSYTTG
>MERGED-PCR_Primer1_rc-PhiX_read2_adapter
AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGTAGATCTCGGTGGTCGCCGTATCATTAAAAAA
>MERGED-Reverse_adapter-1-Reverse_adapter-2
MKRTCKSWWRWRCACACGTCTGAACTCCAGTCACATCACGATCTCGTATGCCGTCTTCTGCTTG
>MERGED-TruSeq_Adapter_Index-GROUP1
GATCGGAAGAGCACACGTCTGAACTCCAGTCACVNBNVNNNATCTCGTATGCCGTCTTCTGCTTG
>MERGED-TruSeq_Adapter_Index-GROUP2
GATCGGAAGAGCACACGTCTGAACTCCAGTCACNNNNNNNTCTCGTATGCCGTCTTCTGCTTG
>MERGED-TruSeq3_IndexedAdapter / PE2_rc-Bisulfite_R1
AGATCGGAAGAGCACACGTCTGAACTCCAGTCAC
>MERGED-ABI_Solid3_Adapter_B-ABI_Solid3_5p_AMP_Primer
CCACTACGCCTCCGCTTTCCTCTCTATGGGCAGTCGGTGAT
>MERGED-Illumina_DpnII_and_NlaIII_Adapters-GROUP
ACAGGTTCAGAGTTCTACAGTCCGACATG
>MERGED-I7_Primer_Nextera_XT-GROUP
CCGAGCCCACGAGACNNNNNNNATCTCGTATGCCGTCTTCTGCTTG
>MERGED-Clontech_SMARTer_II_A_Oligonucleotide-Clontech_SMART_CDS_Primer_II_A
AAGCAGTGGTATCAACGCAGAGTACT
>MERGED-ABI_Solid3_Adapter_A-ABI_Solid3_3p_AMP_Primer
CTGCCCCGGGTTCCTCATTCTCTCAGCAGCATG
>MERGED-I5_Primer_Nextera_XT-GROUP
GACGCTGCCGACGANNNNNNNNGTGTAGATCTCGGTGGTCGCCGTATCATT
>MERGED-Bisulfite_R2-PE1_rc / TruSeq3_UniversalAdapter
AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGTA
>MERGED-Nextera_Transposase-GROUP
CTGTCTCTTATACACATCTSHSRBBSSVSDVGVVAGGC
>MERGED-Illumina_PCR_Primer_Index-GROUP
CAAGCAGAAGACGGCATACGAGATNNNNNNGTGACTGGAGTTC
```

```
1501  >PrefixPE_1 / PE1
1502  TACACTCTTTCCCTACACGACGCTCTTCCGATCT
1503  >TruSeq2_PE_r
1504  AGATCGGAAGAGCGGTTCAGCAGGAATGCCGAG
1505  >I5_Adapter_Nextera
1506  CTGATGGCGCGAGGGAGGCGTGTAGATCTCGGTGGTCGCCGTATCATT
1507  >I7_Adapter_Nextera_No_Barcode
1508  CTGAGCGGGCTGGCAAGGCAGACCGATCTCGTATGCCGTCTTCTGCTTG
1509  >Nextera_LMP_Read1_External_Adapter / Illumina Multiplexing Index Sequencing Primer
1510  GATCGGAAGAGCACACGTCTGAACTCCAGTCAC
1511  >Nextera_LMP_Read2_External_Adapter
1512  GATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT
1513  >Stop_Oligo_STP_8
1514  CCACGGGAACGTGGTGGAATTC
1515  >TruSeq2_SE
1516  AGATCGGAAGAGCTCGTATGCCGTCTTCTGCTTG
1517  >PrefixNX_1 / >PrefixNX/2
1518  AGATGTGTATAAGAGACAG
1519  >Illumina_Single_End_Adapter_1
1520  GATCGGAAGAGCTCGTATGCCGTCTTCTGCTTG
1521  >Illumina_Single_End_Adapter_2 / Illumina Single End PCR Primer 2 /
1522  >Illumina DpnII expression Adapter 2
1523  CAAGCAGAAGACGGCATACGAGCTCTTCCGATCT
1524  >Illumina_Single_End_Sequencing_Primer / Illumina Paired End Adapter 1 /
1525  >Illumina Paried End Sequencing Primer 1 / Illumina Multiplexing Adapter 2 /
1526  >Illumina Multiplexing Read1 Sequencing Primer
1527  ACACTCTTTCCCTACACGACGCTCTTCCGATCT
1528  >Illumina_Paired_End_Adapter_2
1529  GATCGGAAGAGCGGTTCAGCAGGAATGCCGAG
1530  >Illumina_Paired_End_PCR_Primer_2 / PrefixPE/2 / PCR_Primer2
1531  CAAGCAGAAGACGGCATACGAGATCGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCT
1532  >Illumina_Paired_End_Sequencing_Primer_2
1533  CGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCT
1534  >Illumina_DpnII_expression_Adapter_2 / Illumina DpnII expression PCR Primer 1 /
1535  >Illumina NlaIII expression Adapter 2 / Illumina NlaIII expression PCR Primer 1 /
1536  >Illumina Small RNA RT Primer / Illumina DpnII Gex Adapter 2 /
1537  >Illumina DpnII Gex PCR Primer 1 / Illumina NlaIII Gex Adapter 2.01 /
1538  >Illumina NlaIII Gex PCR Primer 1 / Illumina Small RNA PCR Primer 1
1539  CAAGCAGAAGACGGCATACGA
1540  >Illumina_DpnII_expression_PCR_Primer_2 /
1541  >Illumina NlaIII expression PCR Primer 2 / Illumina DpnII Gex PCR Primer 2 /
1542  >Illumina NlaIII Gex PCR Primer 2
1543  AATGATACGGCGACCACCGACAGGTTCAGAGTTCTACAGTCCGA
1544  >Illumina_DpnII_expression_Sequencing_Primer / Illumina DpnII Gex Sequencing Primer
1545  CGACAGGTTCAGAGTTCTACAGTCCGACGATC
1546  >Illumina_NlaIII_expression_Sequencing_Primer / Illumina NlaIII Gex Sequencing Primer
1547  CCGACAGGTTCAGAGTTCTACAGTCCGACATG
1548  >Illumina_Multiplexing_Adapter_1
1549  GATCGGAAGAGCACACGTCT
1550  >Illumina_Multiplexing_PCR_Primer_2_01 / Illumina Multiplexing Read2 Sequencing Primer
1551  GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT
1552  >Illumina_DpnII_Gex_Adapter_1
1553  GATCGTCGGACTGTAGAACTCTGAAC
1554  >Illumina_DpnII_Gex_Adapter_2_01 / Illumina NlaIII Gex Adapter 2.02
1555  TCGTATGCCGTCTTCTGCTTG
1556  >Illumina_NlaIII_Gex_Adapter_1_01
1557  TCGGACTGTAGAACTCTGAAC
1558  >ABI_Dynabead_EcoP_Oligo
1559  CTGATCTAGAGGTACCGGATCCCAGCAGT
1560  >ABI_Solid3_EF1_alpha_Sense_Primer
1561  CATGTGTGTTGAGAGCTTC
1562  >ABI_Solid3_EF1_alpha_Antisense_Primer
1563  GAAAACCAAAGTGGTCCAC
1564  >ABI_Solid3_GAPDH_Forward_Primer
1565  TTAGCACCCCTGGCCAAGG
1566  >ABI_Solid3_GAPDH_Reverse_Primer
1567  CTTACTCCTTGGAGGCCATG
1568  >Clontech_Universal_Primer_Mix_Short
```

```
1569  CTAATACGACTCACTATAGGGC
1570  >Clontech_Universal_Primer_Mix_Long
1571  CTAATACGACTCACTATAGGGCAAGCAGTGGTATCAACGCAGAGT
1572  >FlowCell2
1573  TTTTTTTTTTCAAGCAGAAGACGGCATACGA
1574  >FlowCell1
1575  TTTTTTTTTTAATGATACGGCGACCACCGAGATCTACAC
1576  >Trans1
1577  TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG
1578  >Trans2
1579  GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG
```

Appendix S4: Extended definitions and coding of biotic traits

This appendix contains detailed explanations of how we defined the nine biotic traits included in this study. Headers indicate the variable name used in the paper followed parenthetically by the variable name used in archived data files and scripts. Bullet points denote possible values for discrete traits; values after the colon specify the short-hand coding (capital letters) used to record trait data, numerical values represent the dummy numerical values that characters were transformed into for modeling, and bracketed text specifies how discrete traits are shown in Figure 1. See *Supplemental Methods – Collecting biotic, abiotic, and phylogenetic predictors* for a description of how trait values were obtained.

Iteroparity ("Generational_Structure")

This variable captures variability in the number of times an individual can breed in its lifetime. Iteroparous species can breed more than once in their lifetime, whereas semelparous species only breed once.

- Iteroparous: I = 1 [present]

- Semelparous: S = 0 [absent]

Philopatry ("ReturnToSpawningGround")

We defined philopatry as the intentional movement of reproductively mature individuals back to an established spawning or breeding area to reproduce. Species where evidence indicated such migration were coded as TRUE and species lacking such evidence were coded as FALSE. For species that could plausibly have such migratory life histories (i.e., *Bathyraja panthera* and *Pyroteuthis margaritifera*, with large bodied and strong swimming adults with poorly described life histories with related species known to be philopatric), we scored this trait as unknown.

- Philopatric: TRUE = 1 [present]

- Not philopatric: FALSE = 0 [absent]

- Unknown trait state: NA [missing data]

Spawning ("Spawning_mode")

Marine taxa have a wide array of reproductive strategies with respect to fertilization and subsequent embryonic development and parental care. Our primary focus here was on capturing variation in how far embryos disperse from parents. For animals, eggs or developing young can be internally fertilized and then retained in a parents' body or on a parent's body in specialized brooding structures (pouches, chambers, sacs). Alternatively, animals can lay eggs in nests or other benthically attached structures; parents often (but not always) then disassociate from the laid eggs. Finally, many fishes and invertebrates release free floating embryos, typically preceded by broadcast spawning of gametes. We coded all mammals as "I" and all birds as "N"; fishes and invertebrates had a diverse mix of values. For plants and algae, we applied similar logic as for animals and scored species that retain young on the parent as "I".

- Embryos internally fertilized and attached to parents: I = 0 [absent]

- Embryos in a nest or otherwise benthically attached: N = 1 [intermediate]

- Embryos free floating: F = 2 [present]

Benthicity ("isBenthic")

This variable attempts to capture the diversity of relationships between organisms and a fixed location (e.g., for birds this can be nesting sites on land). We considered organisms to be "never" benthic if their entire life cycle was dissociated from any bottom surface (e.g., anchovy that have planktonic larvae and pelagic swimming adults or dolphins that give birth at sea to swimming young). "Sometimes" benthic organisms have a mixture of life stages that are benthic and pelagic, for example many fishes and invertebrates with planktonic larvae and benthic adult stages. "Always" benthic organisms live their entire life with a strong connection to the bottom (such as hydrozoans and anemonefishes that lack planktonic larvae).

- Always: A = 2 [present]

- Sometimes: S = 1 [intermediate]

- Never: N = 0 [absent]

Planktonicity ("isPlanktonic_atanypoint")

We defined planktonicity to capture whether or not species were planktonic – floating or drifting and not able to actively swim against ocean currents – at any point in their lifetime. This variable attempts to distinguish species that have at least one planktonic life stage (larval, adult, or both) from species that entirely lack a planktonic stage. For example, all mammals, birds and elasmobranchs were coded as FALSE.

- Species is planktonic at any point in their lifecycle: TRUE = 1 [present]

- Species is not planktonic at any point in their lifecycle: FALSE = 0 [absent]

Planktotrophy ("Larval_feeding")

Many marine species, especially invertebrates, have a larval stage that differs in morphology from their adult forms. One common way to classify differences in larval life history is based on larval feeding mode, which in turn influences dispersal (as well as development time and survival probabilities). We defined three categories of larval feeding for this trait: species that do not have planktonic larvae, species with non-feeding planktonic larvae, and species with feeding planktonic larvae. Species that do not have planktonic larvae produce progeny that develop directly into young that resemble miniature adults; these young do not spend time drifting or floating in the water column. Species with non-feeding planktonic larvae, often called lecithotrophic, produce energy-rich eggs that facilitate development from larvae into juveniles without the need to feed; as these larvae are planktonic, they drift in the water column while developing, but development is relatively quick given the well-provisioned eggs. Finally, feeding (or planktotrophic) planktonic

larvae derive from eggs that provide only enough energy to produce a larvae; these larvae must then spend time drifting in the water column feeding on particulate food to fuel development into juveniles, which usually takes longer than in (well-provisioned) non-feeding planktonic larvae. All fish with planktonic larvae are feeding (coded "P"). For corals, we coded species that do not have zooxanthellae in their larvae and rely fully on maternal provisioning as "L", and coded corals with larvae that eat at some point and/or are autotrophic (contain zooxanthellae that can photosynthesize and provide nutrition "on the go") as "P".

- No planktonic larvae: N = 0 [absent]

- Non-feeding (lecithotrophic) planktonic larvae: L = 1 [intermediate]

- Feeding (planktotrophic) planktonic larvae: P = 2 [present]

Pelagic larval duration ("PLD_point2")

A continuous variable representing the mean or midpoint of the estimates available for each species for the length of time (in days) that marine larvae spend in the plankton. We assigned this variable a value of 0 for species that have live birth or brood their young that become benthic adults (isBenthic=Always) and for species who are never benthic (isBenthic=Never); thus pelagic larval duration and benthicity are not conflated in our analyses.

Egg size ("Fecundity_EggSize")

A continuous variable in millimeters, representing the largest dimension of the egg or juvenile at birth or hatching (the propagule size, in the sense of (*3*)).

Body size ("BodySize")

A continuous variable in centimeters, representing the largest dimension of an average organism of each species at its full-grown adult size. As corals and seagrasses are colonial, the meaning of body size is unclear. While we acknowledge that genet size (the collective size of a genotype that might include many discrete ramets or colonies) would be most appropriate in terms of scaling to likely reproductive output, genet sizes are largely unknown and therefore we used ramet or colony size. In some instances only colony area was provided, so we assumed a circular shape to derive the diameter.

Appendix S5: Citations for biotic trait data

This appendix contains the citations for all of the biotic trait values used in this study. The citation numbers in this document correspond to the numbers in the columns labeled: "**<trait_name>_citation**" in the data file: "**/data/biotic/final_biotic_traits_for_suppmat-with-citation-numbers.csv**" in the Dryad archive for this study. Note that the trait Benthicity ("isBenthic") does not have a corresponding citation column (in "final_biotic_traits_for_suppmat-with-citation-numbers.csv") because these values reflect common knowledge within our author group. Other individual trait values that have "NA" in the corresponding citation column also reflect common knowledge within our author group.

We obtained as many trait values as possible from public trait databases (e.g., FishLifeBase (*120*) and SeaLifeBase (*121*)). For species/trait combinations not present in centralized databases (which were the majority), we performed Boolean searches in Web of Science or Google Scholar (e.g., <species name> AND egg size).

Citations

1. L.E. Timm, L.M. Isma, M.W. Johnston, and H.D. Bracken-Grissom. Comparative population genomics and biophysical modeling of shrimp migration in the Gulf of Mexico reveals current-mediated connectivity. *Frontiers in Marine Science*, 7, 2020.

2. J.S. Madin, K.D. Anderson, M.H. Andreasen, C.L.T. Bridge, S.D. Cairns, S.R. Connolly, E.S. Darling, M. Diaz, D.S. Falster, E.C. Franklin, R.D. Gates, A.T.M Harmer, M.O. Hoogenboom, D. Huang, S.A. Keith, M.A. Kosnik, . Kuo, C.-Y, J.M. Lough, C.E. Lovelock, O. Luiz, J. Martinelli, T. Mizerek, J.M. Pandolfi, X. Ponchon, M.S. Pratchett, H.M. Putnam, T.E. Roberts, M. Stat, C.C. Wallace, E. Widman, and A.H. Baird. The coral trait database, a curated database of trait information for coral species from the global oceans. *Scientific Data*, 3, 2016.

3. K.P. Sebens. Reproductive ecology of the intertidal sea anemones *Anthopleura xanthogrammica* (Brandt) and *A. elegantissima* (Brandt): Body size, habitat, and sexual reproduction. *Journal of Experimental Marine Biology and Ecology*, 54:225–250, 1981.

4. BirdLife International. *Aptenodytes patagonicus. The IUCN Red List of Threatened Species 2020*, e.T22697748A184637776, 2020.

5. B.L. Chilvers and S.D. Goldsworthy. *Arctocephalus forsteri. The IUCN Red List of Threatened Species 2015*, e.T41664A45230026, 2015.

6. BirdLife International. *Aythya marila. The IUCN Red List of Threatened Species 2018*, e.T22680398A132525108, 2018.

7. E. O'Reilly, B.M. Titus, M.W. Nelsen, S. Ratchford, and N.E. Chadwick. Giant ephemeral anemones? Rapid growth and high mortality of corkscrew sea anemones *Bartholomea annulata* (Le Sueur, 1817) under variable conditions. *Journal of Experimental Marine Biology and Ecology*, 509:44–53, 2018.

8. D.A. Ebert. Reproductive biology of skates, *Bathyraja* (Ishiyama), along the eastern Bering Sea continental slope. *Journal of Fish Biology*, 66:618–649, 2005.

9. M.R. Gaither, G.A. Gkafas, M. de Jong, F. Sarigol, F. Neat, T. Regnier, D. Moore, D.R. Gröcke, N. Hall, X. Liu, J. Kenny, A. Lucaci, M. Hughes, S. Haldenby, and A.R. Hoelzel. Genomics of habitat choice and adaptive evolution in a deep-sea fish. *Nature Ecology & Evolution*, 2:680–687, 2018.

10. H.-J.T. Hoving, V.V. Laptikhovsky, and B.H. Robison. Vampire squid reproductive strategy is unique among coleoid cephalopods. *Current Biology*, 25:R322–R323, 2015.

11. E.M. Salas, G. Bernardi, M.L. Berumen, M.R. Gaither, and L.A. Rocha. RADseq analyses reveal concordant Indian Ocean biogeographic and phylogeographic boundaries in the reef fish *Dascyllus trimaculatus*. *Royal Society Open Science*, 6:172413, 2019.

12. B.A. Fach. Modeling the influence of hydrodynamic processes on anchovy distribution and connectivity in the Black Sea. *Turkish Journal of Fisheries and Aquatic Sciences*, 14, 2014.

13. BirdLife International. *Eudyptes chrysolophus*. *The IUCN Red List of Threatened Species 2020*, e.T22697793A184720991, 2020.

14. D. Grawunder, E.A. Hambleton, M. Bucher, I. Wolfowicz, N. Bechtoldt, and A. Guse. Induction of gametogenesis in the cnidarian endosymbiosis model *Aiptasia sp. Scientific Reports*, 5, 2015.

15. A.J. Edwards. FAO species catalogue. Vol. 10. Gadiform fishes of the world (Order Gadiformes). An annotated and illustrated catalogue of cods, hakes, grenadiers and other gadiform fishes known to date. *Marine Pollution Bulletin*, 24:326–327, 1990.

16. D. Bowen. *Halichoerus grypus*. *The IUCN Red List of Threatened Species 2016*, e.T9660A45226042, 2016.

17. H. Kopack. Animal Diversity Web: *Lagenorhynchus acutus*. https://animaldiversity.org/accounts/Lagenorhynchus_acutus/, 2000. Accessed: July 2021–December 2023.

18. G.C.D. Estrada, C.H. Callado, M.L.G. Soares, and C.S. Lisi. Annual growth rings in the mangrove *Laguncularia racemosa* (Combretaceae). *Trees*, 22:663–670, 2008.

19. B.-D. Zhang, D.-X. Xue, Y.-L. Li, and J.-X. Liu. RAD genotyping reveals fine-scale population structure and provides evidence for adaptive divergence in a commercially important fish from the Northwestern Pacific Ocean. *PeerJ*, 7:e7242, 2019.

20. R. Froese, D. Pauly, Editors. FishBase, 2023. World Wide Web electronic publication. Available from https://www.fishbase.org, Accessed July 2021–December 2023.

21. M.E. Brandt. The effect of species and colony size on the bleaching response of reef-building corals in the Florida Keys during the 2005 mass bleaching event. *Coral Reefs*, 28:911–924, 2009.

22. A.E. Hopkins. Ecological observations on spawning and early larval development in the Olympia oyster (*Ostrea lurida*). *Ecology*, 17:551–566, 1936.

23. D. Vafidis, C. Antoniadou, and K. Kyriakouli. Reproductive cycle of the edible sea urchin *Paracentrotus lividus* (Echinodermata: Echinoidae) in the Aegean Sea. *Water*, 11:1029, 2019.

24. Indian River Species Inventory. *Panaeus duorarum*. https://www.irlspecies.org/taxa/index.php?taxon=9671, N.D. Accessed: July 2021–December 2023.

25. G. Braulik, G. Minton, M. Amano, and A. Bjørge. *Phocoena phocoena. The IUCN Red List of Threatened Species 2020*, e.T17027A50369903, 2020.

26. L. Rojas-Bracho and B.L. Taylor. *Phocoena sinus. The IUCN Red List of Threatened Species 2017*, e.T17028A50370296, 2017.

27. P. Gèlin, C. Fauvelot, V. Mehn, S. Bureau, H. Rouzè, and H. Magalon. Superclone expansion, long-distance clonal dispersal and local genetic structuring in the coral *Pocillopora damicornis* type β in Reunion Island, south western Indian Ocean. *PLOS ONE*, 12:e0169692, 2017.

28. K. Soong. Sexual reproductive patterns of shallow-water reef corals in Panama. *Bulletin of Marine Science*, 49:832–846, 1991.

29. BirdLife International. *Pygoscelis adeliae. The IUCN Red List of Threatened Species 2020*, e.T22697758A157660553, 2020.

30. BirdLife International. *Pygoscelis antarcticus. The IUCN Red List of Threatened Species 2020*, e.T22697761A184807209, 2020.

31. BirdLife International. *Pygoscelis papua. The IUCN Red List of Threatened Species 2020*, e.T22697755A157664581, 2020.

32. N. Pasiec. *Rhizophora mangle* (red mangrove). In *CABI Compendium*. CABI Digital Library, 2015.

33. J.P. Wourms. Reproduction and development of *Sebastes* in the context of the evolution of piscine viviparity. In *Rockfishes of the genus Sebastes: Their reproduction and early life history*, pages 111–126. Springer, 1991.

34. D. Moran, C.K. Smith, B. Gara, and C.W. Poortenaar. Reproductive behaviour and early development in yellowtail kingfish (*Seriola lalandi* Valenciennes 1833). *Aquaculture*, 262:95–104, 2007.

35. A.L. Gould, K.E. Dougan, S.T. Koenigbauer, and P.V. Dunlap. Life history of the symbiotically luminous cardinalfish *Siphamia tubifer* (Perciformes: Apogonidae). *Journal of Fish Biology*, 89:1359–1377, 2016.

36. C. Bartilotti and A. Dos Santos. The secret life of deep-sea shrimps: Ecological and evolutionary clues from the larval description of *Systellaspis debilis* (Caridea: Oplophoridae). *PeerJ*, 7:e7334, 2019.

37. BirdLife International. *Uria aalge. The IUCN Red List of Threatened Species 2018*, e.T22694841A132577296, 2018.

38. F. Trillmich. *Zalophus wollebaeki. The IUCN Red List of Threatened Species 2015*, e.T41668A45230540, 2015.

39. M.E. Bushnell, J.T. Claisse, and C.W. Laidley. Lunar and seasonal patterns in fecundity of an indeterminate, multiple-spawning surgeonfish, the yellow tang *Zebrasoma flavescens*. *Journal of Fish Biology*, 76:1343–1361, 2010.

40. J.A.B. Claydon. *The structure and dynamics of spawning aggregations of coral reef fish*. PhD thesis, James Cook University, 2005.

41. G.V. Clucas, J.L. Younger, D. Kao, L. Emmerson, C. Southwell, B. Wienecke, A.D. Rogers, C. Bost, G.D. Miller, M.J. Polito, et al. Comparative population genomics reveals key barriers to dispersal in Southern Ocean penguins. *Molecular Ecology*, 27:4680–4697, 2018.

42. G.R. Hoff. Identification of multiple nursery habitats of skates in the eastern Bering Sea. *Journal of Fish Biology*, 88:1746–1757, 2016.

43. J.H. Churchill, J.P. Kritzer, M.J. Dean, J.H. Grabowski, and G.D. Sherwood. Patterns of larval-stage connectivity of Atlantic cod (*Gadus morhua*) within the Gulf of Maine in relation to current structure and a proposed fisheries closure. *ICES Journal of Marine Science*, 74:20–30, 2017.

44. J.M. González-Irusta and P.J. Wright. Spawning grounds of Atlantic cod (*Gadus morhua*) in the North Sea. *ICES Journal of Marine Science: Journal du Conseil*, 73:304–315, 2015.

45. A. Ellison, E. Farnsworth, and G. Moore. *Laguncularia racemosa. The IUCN Red List of Threatened Species 2010*, e.T178798A7609219, 2010.

46. B.A. McIntyre, E.E. McPhee-Shaw, M.B.A. Hatch, and S.M. Arellano. Location matters: Passive and active factors affect the vertical distribution of Olympia oyster (*Ostrea lurida*) larvae. *Estuaries and Coasts*, 44:199–213, 2020.

47. L.E. Timm, T.L. Jackson, J.A. Browder, and H.D. Bracken-Grissom. Population genomics of the commercially important Gulf of Mexico pink shrimp *Farfantepenaeus duorarum* (Burkenroad, 1939) support models of juvenile transport around the Florida peninsula. *Frontiers in Ecology and Evolution*, 9, 2021.

48. A.A. Hohn, A.J. Read, S. Fernandez, O. Vidal, and L.T. Findley. Life history of the vaquita, *Phocoena sinus* (Phocoenidae, Cetacea). *Journal of Zoology*, 239:235–251, 1996.

49. T. Brunel, C.J.G. van Damme, M. Samson, and M. Dickey-Collas. Quantifying the influence of geography and environment on the northeast Atlantic mackerel spawning distribution. *Fisheries Oceanography*, 27:159–173, 2017.

50. K. Takano, A. Takemura, M. Furihata, T. Nakanishi, and A. Hara. Annual reproductive and spawning cycles of female *Sebastiscus marmoratus*. In *Developments in Environmental Biology of Fishes*, pages 39–48. Springer Netherlands, 1991.

51. W.B. Driggers, B.S. Frazier, D.H. Adams, G.F. Ulrich, C.M. Jones, E.R. Hoffmayer, and M.D. Campbell. Site fidelity of migratory bonnethead sharks *Sphyrna tiburo* (L. 1758) to specific estuaries in South Carolina, USA. *Journal of Experimental Marine Biology and Ecology*, 459:61–69, 2014.

52. D.E. Lee, C.L. Abraham, P.M. Warzybok, R.W. Bradley, and W.J. Sydeman. Age-specific survival, breeding success, and recruitment in common murres (*Uria aalge*) of the California Current System. *The Auk*, 125:316–325, 2008.

53. Wikipedia contributors. Thick-billed murre. https://en.wikipedia.org/wiki/Thick-billed_murre, 2023. Accessed: July 2021–December 2023.

54. R.K. Cowen and S. Sponaugle. Relationships between early life history traits and recruitment among coral reef fishes. In *Early Life History and Recruitment in Fish Populations*, pages 423–449. Springer, 1997.

55. M.W. Miller, A.J. Bright, R.E. Pausch, and D.E. Williams. Larval longevity and competency patterns of caribbean reef-building corals. *PeerJ*, 8:e9705, 2020.

56. D.E. Morse, N. Hooker, A.N.C. Morse, and R.A. Jensen. Control of larval metamorphosis and recruitment in sympatric agariciid corals. *Journal of Experimental Marine Biology and Ecology*, 116:193–217, 1988.

57. N.K. Dulvy and J.D. Reynolds. Evolutionary transitions among egg–laying, live–bearing and maternal inputs in sharks and rays. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 264:1309–1315, 1997.

58. P. Jivoff, A.H. Hines, and L.S. Quackenbush. Chapter 7: Reproduction biology and embryonic development. In *The Blue Crab: Callinectes sapidus*, pages 255–298. University of Maryland, 2007.

59. A.L. Moran. Spawning and larval development of the black turban snail *Tegula funebralis* (Prosobranchia: Trochidae). *Marine Biology*, 128:107–114, 1997.

60. J.S. Nelson, T.C. Grande, and M.V.H. Wilson. *Fishes of the world*. John Wiley & Sons, Hoboken, New Jersey, 2016.

61. A. Arkhipkin. Age and growth of planktonic squids *Cranchia scabra* and *Liocranchia reinhardti* (Cephalopoda, Cranchiidae) in epipelagic waters of the central-east Atlantic. *Journal of Plankton Research*, 18:1675–1683, 1996.

62. E.S. Chang, M.E. Orive, and P. Cartwright. Nonclonal coloniality: Genetically chimeric colonies through fusion of sexually produced polyps in the hydrozoan *Ectopleura larynx*. *Evolution Letters*, 2:442–455, 2018.

63. J.J. Grigor, M.S. Schmid, and L. Fortier. Growth and reproduction of the chaetognaths *Eukrohnia hamata* and *Parasagitta elegans* in the Canadian Arctic Ocean: Capital breeding versus income breeding. *Journal of Plankton Research*, 39:910–929, 2017.

64. A. Schlesinger, E. Kramarsky-Winter, H. Rosenfeld, R. Armoza-Zvoloni, and Y. Loya. Sexual plasticity and self-fertilization in the sea anemone *Aiptasia diaphana*. *PLOS ONE*, 5:e11874, 2010.

65. Y. Nakajima, Y. Zayasu, C. Shinzato, N. Satoh, and S. Mitarai. Genetic differentiation and connectivity of morphological types of the broadcast-spawning coral *Galaxea fascicularis* in the Nansei Islands, Japan. *Ecology and Evolution*, 6:1457–1469, 2016.

66. M.F. Gomon and D.J. Bray. Fishes of Australia: Common Galaxias, *Galaxias maculatus* (Jenyns 1842). https://fishesofaustralia.net.au/home/species/2129, 2021. Accessed: July 2021–December 2023.

67. E.M. Moore, T.G. Langley, J.S. Goldstein, and W.H. Watson. American lobster, *Homarus americanus*, reproduction and recruitment in a New England estuary. *Estuaries and Coasts*, 43:2141–2151, 2020.

68. R.I. Lonard, F.W. Judd, H.R. DeYoe, and R. Stalter. Biology and ecology of the halophyte *Laguncularia racemosa* (l.) Gaertn. F.: A review. In *Handbook of Halophytes*, pages 1–15. Springer International Publishing, 2020.

69. J.G. Myoung, Y.U. Kim, Y.J. Park, B.G. Kim, J.M. Kim, and H.T. Heo. Embryonic development of larvae and juveniles of the small yellow croaker (*Larimichthys polyactis*) reared in aquarium. *Korean Journal of Fisheries and Aquatic Sciences*, 37:478–484, 2004.

70. S. Goffredo, V. Airi, J. Radeti'c, and F. Zaccanti. Sexual reproduction of the solitary sunset cup coral *Leptopsammia pruvoti* (Scleractinia, Dendrophylliidae) in the Mediterranean. 2. quantitative aspects of the annual reproductive cycle. *Marine Biology*, 148:923–931, 2005.

71. J.L. Padilla-Gamiño, R.R. Bidigare, D.J. Barshis, A. Alamaru, L. H'edouin, X. Hernàndez-Pech, F. Kandel, S. Leon Soon, M.S. Roth, L.J. Rodrigues, A.G. Grottoli, C. Portocarrero, F. Wagenhauser, S.A. Buttler, and R.D. Gates. Are all eggs created equal? a case study from the Hawaiian reef-building coral *Montipora capitata*. *Coral Reefs*, 32:137–152, 2012.

72. R.T. Hanlon and J.B. Messenger. *Cephalopod Behaviour*. Cambridge University Press, 1998.

73. S.W. Davies, M.E. Strader, J.T. Kool, C.D. Kenkel, and M.V. Matz. Modeled differences of coral life-history traits influence the refugium potential of a remote Caribbean reef. *Coral Reefs*, 36:913–925, 2017.

74. J.S. Barber, J.E. Dexter, S.K. Grossman, C.M. Greiner, and J.T. Mcardle. Low temperature brooding of Olympia oysters (*Ostrea lurida*) in northern Puget Sound. *Journal of Shellfish Research*, 35:351–357, 2016.

75. C. Carreras, A. Garcìa-Cisneros, O.S. Wangensteen, V. Ordóñez, C. Palacín, M. Pascual, and X. Turon. East is east and west is west: Population genomics and hierarchical analyses reveal genetic structure and adaptation footprints in the keystone species *Paracentrotus lividus* (Echinoidea). *Diversity and Distributions*, 26:382–398, 2019.

76. J.M. Leis. The pelagic stage of reef fishes: The larval biology of coral reef fishes. In *The Ecology of Fishes on Coral Reefs*, pages 183–230. Springer, 1991.

77. M.F. Strathmann. *Reproduction and Development of Marine Invertebrates of the Northern Pacific Coast*. University of Washington Press, 1987.

78. M.G. Frick, H.R. Martins, A.B. Bolten, K.A. Bjorndal, and K.L. Williams. Diet and fecundity of Columbus crabs, *Planes minutus*, associated with oceanic-stage loggerhead sea turtles, *Caretta caretta*, and inanimate flotsam. *Journal of Crustacean Biology*, 24:350–355, 2004.

79. E.G. Neves. Histological analysis of reproductive trends of three *Porites* species from Kane'ohe Bay, Hawai'i. *Pacific Science*, 54:195–200, 2000.

80. D.D. Turgeon, J.F. Quinn Jr., A.E. Bogan, E.V. Coan, F.G. Hochberg, W.G. Lyons, P.M. Mikkelsen, R.J. Neves, C.F.E. Roper, G. Rosenberg, A. Scheltema, F.G. Thompson, M. Vecchione, and J.D. Williams. *Pyroteuthis margaritifera* (Rüppell, 1844) Jewel enope squid. In *Common and scientific names of aquatic invertebrates from the United States and Canada: Mollusks, 2nd ed.*, page 526. American Fisheries Society, 1998. Accessed: July 2021–December 2023 via SeaLifeBase; https://www.sealifebase.ca/summary/Pyroteuthis-margaritifera.html.

81. P. Kerrison and H.N. Le. Environmental factors on egg liberation and germling production of *Sargassum muticum*. *Journal of Applied Phycology*, 28:481–489, 2015.

82. M.J.A. Vermeij, K.L. Barott, A.E. Johnson, and K.L. Marhaver. Release of eggs from tentacles in a Caribbean coral. *Coral Reefs*, 29:411–411, 2010.

83. B. Johnson. Animal Diversity Web: *Vampyroteuthis infernalis*. https://animaldiversity.org/accounts/Vampyroteuthis_infernalis/, 2000. Accessed: July 2021–December 2023.

84. P. Foxton. Observations on the early development and hatching of the eggs of *Acanthephyra purpurea* a. Milne-edwards. *Crustaceana*, 6:235–237, 1964.

85. E.M. Graham, A.H. Baird, S.R. Connolly, M.A. Sewell, and B.L. Willis. Rapid declines in metabolism explain extended coral larval longevity. *Coral Reefs*, 32:539–549, 2013.

86. B.H. Cornwell. Gene flow in the anemone *Anthopleura elegantissima* limits signatures of local adaptation across an extensive geographic range. *Molecular Ecology*, 29:2550–2566, 2020.

87. T.D. Williams. *The Penguins Spheniscidae. Bird Familes of the World*, volume 2. Oxford University Press, 1995.

88. I. Wolfowicz, S. Baumgarten, P.A. Voss, E.A. Hambleton, C.R. Voolstra, M. Hatta, and A. Guse. *Aiptasia sp.* larvae as a model to reveal mechanisms of symbiont selection in cnidarians. *Scientific Reports*, 6, 2016.

89. D. Macedo, I. Caballero, M. Mateos, R. Leblois, S. McCay, and L.A. Hurtado. Population genetics and historical demographic inferences of the blue crab *Callinectes sapidus* in the US based on microsatellites. *PeerJ*, 7:e7780, 2019.

90. COSEWIC. *COSEWIC assessment and status report on the Roundnose Grenadier Coryphaenoides rupestris in Canada*. Committee on the Status of Endangered Wildlife in Canada, 2008.

91. W. Van Treuren, K.K. Brower, L. Labanieh, D. Hunt, S. Lensch, B. Cruz, H.N. Cartwright, C. Tran, and P.M. Fordyce. Live imaging of *Aiptasia* larvae, a model system for coral and anemone bleaching, using a simple microfluidic device. *Scientific Reports*, 9:9275, 2019.

92. C.P. Marquis, A.H. Baird, R. de Nys, C. Holmström, and N. Koziumi. An evaluation of the antimicrobial properties of the eggs of 11 species of scleractinian corals. *Coral Reefs*, 24:248–253, 2005.

93. E.R. Annis, L.S. Incze, R.S. Steneck, and N. Wolff. Estimates of *in situ* larval development time for the lobster, *Homarus americanus*. *Journal of Crustacean Biology*, 27:454–462, 2007.

94. S.D. Cairns. Species richness of recent Scleractinia. *Atoll Research Bulletin*, 459:1–46, 1999.

95. A.H. Baird, J.R. Guest, and B.L. Willis. Systematic and biogeographical patterns in the reproductive biology of scleractinian corals. *Annual Review of Ecology, Evolution, and Systematics*, 40:551–571, 2009.

96. R. Villanueva, E.A.G. Vidal, F.Á. Fernández-Álvarez, and J. Nabhitabhata. Early mode of life and hatchling size in cephalopod molluscs: Influence on the species distributional ranges. *PLOS ONE*, 11:e0165334, 2016.

97. A. Hettinger, E. Sanford, T.M. Hill, J.D. Hosfelt, A.D. Russell, and B. Gaylord. The influence of food supply on the response of Olympia oyster larvae to ocean acidification. *Biogeosciences*, 10:6629–6638, 2013.

98. L. Fenaux, C. Cellario, and M. Etienne. Croissance de la larve de l'oursin *Paracentrotus lividus*. *Marine Biology*, 86:151–157, 1985.

99. B.L. Jennison. Reproduction in three species of sea anemones from Key West, Florida. *Canadian Journal of Zoology*, 59:1708–1719, 1981.

100. J.B. Pfaller, A.C. Payton, K.A. Bjorndal, A.B. Bolten, and S.F. McDaniel. Hitchhiking the high seas: Global genomics of rafting crabs. *Ecology and Evolution*, 9:957–974, 2019.

101. A. Moreno, A. Dos Santos, U. Piatkowski, A.M.P. Santos, and H. Cabral. Distribution of cephalopod paralarvae in relation to the regional oceanography of the Western Iberia. *Journal of Plankton Research*, 31:73–91, 2008.

102. E.J. Guerra-Castro, J.E. Conde, A. Barcelo, and J.J. Cruz-Motta. Variation in fouling assemblages associated with prop roots of *Rhizophora mangle* L. in the Caribbean: The role of neutral and niche processes. *Austral Ecology*, 46:991–1007, 2021.

103. P. Baweja, S. Kumar, D. Sahoo, and I. Levine. Biology of seaweeds. In *Seaweed in Health and Disease Prevention*, pages 41–106. Elsevier, 2016.

104. K.N. Kosobokova and A.I. Isachenko. The gonad maturation and size structure of the population of abundant planktonic predator *Eukrohnia hamata* (Möbius, 1875) (Chaetognatha) in the Eurasian basin of the Arctic Ocean in summer. *Russian Journal of Marine Biology*, 43:25–33, 2017.

105. M. Bucher, I. Wolfowicz, P.A. Voss, E.A. Hambleton, and A. Guse. Development and symbiosis establishment in the cnidarian endosymbiosis model *Aiptasia sp. Scientific Reports*, 6, 2016.

106. R.C. Babcock and A.J. Heyward. Larval development of certain gamete-spawning scleractinian corals. *Coral Reefs*, 5:111–116, 1986.

107. B.D. Limer, J. Bloomberg, and D.M. Holstein. The influence of eddies on coral larval retention in the Flower Garden banks. *Frontiers in Marine Science*, 7, 2020.

108. S.K. Grossman, E.E. Grossman, J.S. Barber, S.K. Gamblewood, and S.C. Crosby. Distribution and transport of Olympia oyster *Ostrea lurida* larvae in Northern Puget Sound, Washington. *Journal of Shellfish Research*, 39:215, 2020.

109. J.H. Vandermeulen. Studies on reef corals. III. Fine structural changes of calicoblast cells in *Pocillopora damicornis* during settling and calcification. *Marine Biology*, 31:69–77, 1975.

110. H.M. Austin. The eggs and planktonic stages of British marine fishes. *The Quarterly Review of Biology*, 52:216–216, 1977.

111. R. Collin, M.C. Díaz, J. Norenburg, R.M. Rocha, J.A. Sánchez, M. Schulze, A. Schwartz, and A. Valdés. Photographic identification guide to some common marine invertebrates of Bocas Del Toro, Panama. *Caribbean Journal of Science*, 41:638–707, 2005. Accessed: July 2021–December 2023 via SeaLifeBase; https://www.sealifebase.ca/summary/Stephanocoenia-intersepta.html.

112. D.T. Wilson and M.I. McCormick. Microstructure of settlement-marks in the otoliths of tropical reef fishes. *Marine Biology*, 134:29–41, 1999.

113. G.B. Nanninga, P. Saenz-Agudelo, P. Zhan, I. Hoteit, and M.L. Berumen. Not finding Nemo: Limited reef-scale retention in a coral reef fish. *Coral Reefs*, 34:383–392, 2015.

114. E. Macpherson and N. Raventos. Relationship between pelagic larval duration and geographic distribution of Mediterranean littoral fishes. *Marine Ecology Progress Series*, 327:257–265, 2006.

115. Ronald E. Thresher, P.L. Colin, and L.J. Bell. Planktonic duration, distribution and population structure of western and central Pacific damselfishes (Pomacentridae). *Copeia*, 1989:420, 1989.

116. I. Palomera, B. Morales-Nin, and J. Lleonart. Larval growth of anchovy, *Engraulis encrasicolus*, in the western Mediterranean Sea. *Marine Biology*, 99:283–291, 1988.

117. H.G. Hansson. SeaLifeBase: *Eukrohnia hamata* (Möbius, 1875). https://www.sealifebase.ca/summary/Eukrohnia-hamata.html, 1998. Accessed: July 2021–December 2023.

118. S. Campana. Year-class strength and growth rate in young Atlantic cod *Gadus morhua*. *Marine Ecology Progress Series*, 135:21–26, 1996.

119. R. Suwa and M. Nakamura. A precise comparison of developmental series of oocyte growth and oocyte maturation between real-oocytes and pseudo-oocytes in the coral *Galaxea fascicularis*. *Journal of Coral Reef Studies*, 20:1–7, 2018.

120. R.E. Thresher and E.B. Brothers. Reproductive ecology and biogeography of Indo-West Pacific angelfishes (Pisces: Pomacanthidae). *Evolution*, 39:878–887, 1985.

121. N. Raventos and E. Macpherson. Planktonic larval duration and settlement marks on the otoliths of Mediterranean littoral fishes. *Marine Biology*, 138:1115–1120, 2001.

122. Wikipedia contributors. Japanese sea bass. https://en.wikipedia.org/wiki/Japanese_sea_bass, 2023. Accessed: July 2021–December 2023.

123. A. Jackson. Sunset cup coral (*Leptopsammia pruvoti*), 2008. Accessed: July 2021–December 2023 via MarLIN; https://www.marlin.ac.uk/species/detail/1285.

124. J.B. Puritz, J.R. Gold, and D.S. Portnoy. Fine-scale partitioning of genomic variation among recruits in an exploited fishery: Causes and consequences. *Scientific Reports*, 6, 2016.

125. G. Alexander, J.R. Hancock, A.S. Huffmyer, and S.B. Matsuda. Larval thermal conditioning does not improve post-settlement thermal tolerance in the dominant reef-building coral, *Montipora capitata*. *Coral Reefs*, 41:333–342, 2022.

126. R.C. Williams, B.C. Jackson, L. Duvaux, D.A. Dawson, T. Burke, and W. Sinclair. The genetic structure of *Nautilus pompilius* populations surrounding Australia and the Philippines. *Molecular Ecology*, 24:3316–3328, 2015.

127. S. L'opez, X. Turon, E. Montero, C. Palac'in, C. Duarte, and I. Tarjuelo. Larval abundance, recruitment and early mortality in *Paracentrotus lividus* (Echinoidea). interannual variability and plankton-benthos coupling. *Marine Ecology Progress Series*, 172:239–251, 1998.

128. E.B. Brothers and R.E. Thresher. Pelagic duration, dispersal, and the distribution of Indo-Pacific coral-reef fishes. In *The Ecology of Coral Reefs*, pages 53–69. NOAA Publication, 1985.

129. M. Oshima, D. Robert, Y. Kurita, M. Yoneda, O. Tominaga, T. Tomiyama, Y. Yamashita, and S. Uehara. Do early growth dynamics explain recruitment success in Japanese flounder *Paralichthys olivaceus* off the Pacific coast of northern Japan? *Journal of Sea Research*, 64:94–101, 2010.

130. J.B. Pfaller and M.A. Gil. Sea turtle symbiosis facilitates social monogamy in oceanic crabs via refuge size. *Biology Letters*, 12:20160607, 2016.

131. A.L. Primo, A.C. Vaz, D. Crespo, F. Costa, M. Pardal, and F. Martinho. Contrasting links between growth and survival in the early life stages of two flatfish species. *Estuarine, Coastal and Shelf Science*, 254:107314, 2021.

132. V.R. Cumbo, T.Y. Fan, and P.J. Edmunds. Effects of exposure duration on the response of *Pocillopora damicornis* larvae to elevated temperature and high pCO2. *Journal of Experimental Marine Biology and Ecology*, 439:100–107, 2013.

133. P. Edmunds, R. Gates, and D. Gleason. The biology of larvae from the reef coral *Porites astreoides*, and their response to temperature disturbances. *Marine Biology*, 139:981–989, 2001.

134. C.D. Storlazzi, M. van Ormondt, Y.-L. Chen, and E.P.L. Elias. Modeling fine-scale coral larval dispersal and interisland connectivity to help designate mutually-supporting coral reef marine protected areas: Insights from Maui Nui, Hawaii. *Frontiers in Marine Science*, 4, 2017.

135. L.E. Timm, H.D. Bracken-Grissom, A. Sosnowski, M. Breitbart, M. Vecchione, and H. Judkins. Population genomics of three deep-sea cephalopod species reveals connectivity between the Gulf of Mexico and northwestern Atlantic ocean. *Deep Sea Research Part I: Oceanographic Research Papers*, 158:103222, 2020.

136. B. Villamor, M. Bernal, and C. Hernandez. Models describing mackerel (*Scomber scombrus*) early life growth in the north and northwest of the Iberian Peninsula in 2000. *Scientia Marina*, 68:571–583, 2004.

137. M.S. Love, M. Yoklavich, and L.K. Thorsteinson. *The Rockfishes of the Northeast Pacific*. University of California Press, 2002.

138. A.L. Gould and P.V. Dunlap. Shedding light on specificity: Population genomic structure of a symbiosis between a coral reef fish and luminous bacterium. *Frontiers in Microbiology*, 10, 2019.

139. G.M. Wellington and B.C. Victor. Planktonic larval duration of one hundred species of Pacific and Atlantic damselfishes (Pomacentridae). *Marine Biology*, 101:557–567, 1989.

140. R.E. Young, M. Vecchione, P.R. Stephens, and J.Z. Young. Morphological observations on a hatching and a paralarva of the vampire squid, *Vampyroteuthis infernalis* Chun (Mollusca: Cephalopoda). *Proceedings of the Biological Society of Washington*, 112:661–666, 1999.

141. G.E. Pickford. The distribution of the eggs of *Vampyroteuthis infernalis* Chun. *Journal of Marine Research*, 8:73–83, 1949.

142. C.K. Callan, A.I. Burgess, C.R. Rothe, and R. Touse. Development of improved feeding methods in the culture of yellow tang, *Zebrasoma flavescens*. *Journal of the World Aquaculture Society*, 49:493–503, 2018.

143. S.A. Sudnik and T. Falkenhaug. Maturation, fecundity and embryos development in three deep-water shrimps (Decapoda: Caridea: Pasiphaeidae, Oplophoridae) along the mid-Atlantic Ridge from Iceland to the Azores. *Arthropoda Selecta*, 24:401–416, 2015.

144. J.E. Randall. A revision of the surgeonfish genus *Acanthurus*. *Pacific Science*, 10:159–235, 1956.

145. E.A. Agudo-Adriani, J. Cappelletto, F. Cavada-Blanco, and A. Croquer. Colony geometry and structural complexity of the endangered species *Acropora cervicornis* partly explains the structure of their associated fish assemblage. *PeerJ*, 4:e1861, 2016.

146. A.L. Zubillaga, L.M. M'arquez, A. Cr'oquer, and C. Bastidas. Ecological and genetic data indicate recovery of the endangered coral *Acropora palmata* in Los Roques, Southern Caribbean. *Coral Reefs*, 27:63–72, 2007.

147. Wikipedia contributors. Elkhorn coral. https://en.wikipedia.org/wiki/Elkhorn_coral, 2023. Accessed: July 2021–December 2023.

148. J. Voss and L. Richardson. Coral diseases near Lee Stocking Island, Bahamas: patterns and potential drivers. *Diseases of Aquatic Organisms*, 69:33–40, 2006.

149. Coralpedia. *Agaricia undata* (Ellis 1786). https://coralpedia.bio.warwick.ac.uk/en/corals/agaricia_undata, 2023. Accessed: July 2021–December 2023.

150. Wikipedia contributors. *Bartholomea annulata*. https://en.wikipedia.org/wiki/Bartholomea_annulata, 2023. Accessed: July 2021–December 2023.

151. J.A. Thia, K. McGuigan, L. Liggins, W.F. Figueira, C.E. Bird, A. Mather, J.L. Evans, and C. Riginos. Genetic and phenotypic variation exhibit both predictable and stochastic patterns across an intertidal fish metapopulation. *Molecular Ecology*, 30:4392–4414, 2021.

152. E. Beukhof, T.S. Dencker, M.L.D. Palomares, and A. Maureaud. A trait collection of marine fish species from North Atlantic and Northeast Pacific continental shelf seas [dataset]. *Pangaea*, 2019.

153. National Wildlife Federation. Blue crab (*Callinectes sapidus*). https://www.nwf.org/Educational-Resources/Wildlife-Guide/Invertebrates/Blue-Crab, N.D. Accessed: July 2021–December 2023.

154. Wikipedia contributors. *Tegula funebralis*. https://en.wikipedia.org/wiki/Tegula_funebralis, 2023. Accessed: July 2021–December 2023.

155. Wikipedia contributors. Cranchiidae. https://en.wikipedia.org/wiki/Cranchiidae, 2023. Accessed: July 2021–December 2023.

156. E. Jansson, F. Besnier, K. Malde, C. Andr'e, G. Dahle, and K.A. Glover. Genome wide analysis reveals genetic divergence between Goldsinny wrasse populations. *BMC Genetics*, 21:1–15, 2020.

157. Wikipedia contributors. *Ectopleura larynx*. https://en.wikipedia.org/wiki/Ectopleura_larynx, 2023. Accessed: July 2021–December 2023.

158. Wikipedia contributors. Exaiptasia. https://en.wikipedia.org/wiki/Exaiptasia, 2023. Accessed: July 2021–December 2023.

159. J.E.N. Veron and M. Stafford-Smith. *Corals of the World*. Australian Institute of Marine Science, 2000.

160. Wikipedia contributors. American lobster. https://en.wikipedia.org/wiki/American_lobster, 2023. Accessed: July 2021–December 2023.

161. H.K. Lim, M.H. Le, C.M. An, S.Y. Kim, M.S. Park, and Y.J. Chang. Reproductive cycle of yellow croaker *Larimichthys polyactis* in southern waters off Korea. *Fisheries Science*, 76:971–980, 2010.

162. E. Caroselli, F. Zaccanti, G. Mattioli, G. Falini, O. Levy, Z. Dubinsky, and S. Goffredo. Growth and demography of the solitary scleractinian coral *Leptopsammia pruvoti* along a sea surface temperature gradient in the Mediterranean Sea. *PLOS ONE*, 7:e37848, 2012.

163. A. Nishikawa, R.A. Kinzie, and K. Sakai. Fragmentation and genotypic diversity of the scleractinian coral *Montipora capitata* in Kaneohe Bay, Hawaii. *Journal of the Marine Biological Association of the United Kingdom*, 89:101–107, 2008.

164. Wikipedia contributors. Chambered nautilus. https://en.wikipedia.org/wiki/Chambered_nautilus, 2023. Accessed: July 2021–December 2023.

165. R. Bak and E. Meesters. Coral population structure: the hidden information of colony size-frequency distributions. *Marine Ecology Progress Series*, 162:301–306, 1998.

166. P.S.E. zu Ermgassen, M.W. Gray, C.J. Langdon, M.D. Spalding, and R.D. Brumbaugh. Quantifying the historic contribution of Olympia oysters to filtration in Pacific Coast (USA) estuaries and the implications for restoration objectives. *Aquatic Ecology*, 47:149–161, 2013.

167. C.F. Boudouresque and M. Verlaque. *Paracentrotus lividus*. In *Developments in Aquaculture and Fisheries Science*, pages 297–327. Elsevier, 2013.

168. R.L. Brownell, L.T. Findley, O. Vidal, A. Robles, and N.S. Manzanilla. External morphology and pigmentation of the vaquita, *Phocoena sinus* (Cetacea: Mammalia). *Marine Mammal Science*, 3:22–30, 1987.

169. Wikipedia contributors. *Pisaster ochraceus*. https://en.wikipedia.org/wiki/Pisaster_ochraceus, 2023. Accessed: July 2021–December 2023.

170. Wikipedia contributors. *Planes minutus*. https://en.wikipedia.org/wiki/Planes_minutus, 2023. Accessed: July 2021–December 2023.

171. R.H. Richmond. Energetic relationships and biogeographical differences among fecundity, growth and reproduction in the reef coral *Pocillopora damicornis*. *Bulletin of Marine Science*, 41:594–604, 1987.

172. C.L. Hunter. *Genotypic Diversity and Population Structure of the Hawaiian Reef Coral, Porites compressa*. PhD thesis, University of Hawai'i, Honolulu, 1988.

173. M.E. Flock and T.L. Hopkins. Species composition and food habits of the micronektonix cephalopod assemblage in the eastern Gulf of Mexico. *Bulletin of Marine Science*, 49:638–659, 1991.

174. Wikipedia contributors. *Rhizophora mangle*. https://en.wikipedia.org/wiki/Rhizophora_mangle, 2023. Accessed: July 2021–December 2023.

175. A.L. Vereshchaka, J. Olesen, and A.A. Lunina. Global diversity and phylogeny of pelagic shrimps of the former genera *Sergestes* and *Sergia* (Crustacea, Dendrobranchiata, Sergestidae), with definition of eight new genera. *PLOS ONE*, 9:e112057, 2014.

176. F. Vaux, L.K. Rasmuson, L.A. Kautzi, P.S. Rankin, M.T.O. Blume, K.A. Lawrence, S. Bohn, and K.G. O'Malley. Sex matters: Otolith shape and genomic variation in deacon rockfish (*Sebastes diaconus*). *Ecology and Evolution*, 9:13153–13173, 2019.

177. G.W. Baeck, J.M. Jeong, Y.M. Yeo, S.-H. Huh, and J.M. Park. Length-weight and length-length relationships for 10 species of scorpionfishes (Scorpaenidae) on the south coast of Korea. *Journal of Applied Ichthyology*, 28:677–679, 2012.

178. F.B. Griffiths and S.B. Brandt. Distribution of mesopelagic decapod Crustacea in and around a warm-core eddy in the Tasman Sea. *Marine Ecology Progress Series*, 12:175–184, 1983.

179. A.V. Golikov, F.R. Ceia, R.M. Sabirov, J.D. Ablett, I.G. Gleadall, G. Gudmundsson, H.J. Hoving, H. Judkins, J. Pálsson, A.L. Reid, R. Rosas-Luis, E.K. Shea, R. Schwarz, and Xavier J.C. The first global deep-sea stable isotope assessment reveals the unique trophic ecology of vampire squid *Vampyroteuthis infernalis* (Cephalopoda). *Scientific Reports*, 9, 2019.

180. T. Foster and J. Gilmour. Egg size and fecundity of biannually spawning corals at Scott Reef. *Scientific Reports*, 10, 2020.

181. R.M. Coleman. Measuring parental investment in nonspherical eggs. *Copeia*, 1991:1092–1098, 1991.

182. C. E. Ford. Reproduction in the aggregating sea anemone, *Anthopleura elegantissima*. *The Pacific Science*, 18(2):138–145, 1964.

183. Y-H. Kong and I-S. Chen. Reproductive biology of the intertidal Frillfin goby, *Bathygobius fuscus* in Keelung, Taiwan. *Journal of Marine Science and Technology*, 21:213–215, 2013.

184. D.J. Graham, R. Fulford, P. Biesiot, and H. Perry. Fecundity and egg diameter of primiparous and multiparous blue crab *Callinectes sapidus* (Brachyura: Portunidae) in Mississippi waters. *Journal of Crustacean Biology*, 32:49–56, 2012.

185. N.G. Emel'yanova, D.A. Pavlov, and L.T.B. Thuan. Hormonal stimulation of maturation and ovulation, gamete morphology, and raising of larvae in *Dascyllus trimaculatus* (Pomacentridae). *Journal of Ichthyology*, 49:249–263, 2009.

186. N. Okubo, T. Mezaki, Y. Nozawa, Y. Nakano, Y.-T. Lien, H. Fukami, D.C. Hayward, and E.E. Ball. Comparative embryology of eleven species of stony corals (Scleractinia). *PLOS One*, 8:e84115, 2013.

187. P. Ouellet and F. Plante. An investigation of the sources of variability in American lobster (*Homarus americanus*) eggs and larvae: Female size and reproductive status, and interannual and interpopulation comparisons. *Journal of Crustacean Biology*, 24:481–495, 2004.

188. R.H. Young-Gil. Studies on sexual maturation of spotted sea bass, *Lateolabrax maculatus*. *Korean Journal of Fisheries and Aquatic Sciences*, 34:526–535, 2001.

189. K. Kasimatis and C. Riginos. A phylogenetic analysis of egg size, clutch size, spawning mode, adult body size, and latitude in reef fishes. *Coral Reefs*, 35:387–397, 2015.

190. Z. Masood and R.Y. Farooq. Morphology and early life history pattern of some *Lutjanus* species: a review. *International Journal of Biology and Biotechnology*, 8:455–461, 2011.

191. K. De Baets, C. Klug, D. Korn, and N.H. Landman. Data from: Early evolutionary trends in ammonoid embryonic development, 2015. *Dryad*, 10.5061/dryad.k84c3k8j.

192. D.M. Holstein, T.B. Smith, J. Gyory, and C.B. Paris. Fertile fathoms: Deep reproductive refugia for threatened shallow corals. *Scientific Reports*, 5:12407, 2015.

193. M.P. Velazquez and A. Gracia. Fecundity of *Litopenaeus setiferus*, *Farfantepenaeus aztecus*, and *F. duorarum*, in the Southwestern Gulf of Mexico. *Gulf and Caribbean Research*, 12:1–1, 2000.

194. J. Stoddart and R. Black. Cycles of gametogenesis and plantation in the coral *Pocillopora damicornis*. *Marine Ecology Progress Series*, 23:153–164, 1985.

195. M.-Y. Leu, C.-H. Liou, W.-H. Wang, S.-D. Yang, and P.-J. Meng. Natural spawning, early development and first feeding of the semicircle angelfish [*Pomacanthus semicirculatus* (Cuvier, 1831)] in captivity. *Aquaculture Research*, 40:1019–1030, 2009.

196. Animalia. *Pterygioteuthis microlampas*. https://animalia.bio/pterygioteuthis-microlampas, 2023. Accessed: July 2021–December 2023.

197. B. Gibson, L. Gibson, and J. Gibson. *Rhizophora mangle*. https://www.sdnhm.org/oceanoasis/fieldguide/rhiz-man.html, 2000. Accessed: July 2021–December 2023.

198. J.R. Lueg, A.L. Moulding, V.N. Kosmynin, and D.S. Gilliam. Gametogenesis and spawning of *Solenastrea bournoni* and *Stephanocoenia intersepta* in Southeast Florida, USA. *Journal of Marine Biology*, 2012:1–13, 2012.

199. M.E. Bushnell. *Reproduction of Zebrasoma flavescens: oocyte maturation, spawning patterns, and an estimate of reproductive potential for female yellow tang in Hawaii*. PhD thesis, University of Hawaii at Manoa, 2007.

200. D.A. Pavlov and N.G. Emel'yanova. Fertility of ovulated oocytes after their short-term storage in water in several species of marine tropical fishes. *Journal of Ichthyology*, 48:655–664, 2008.

201. B.B. Collette. Hemiramphidae, halfbeaks. In K.E. Carpenter and N. De Angelis, editors, *The living marine resources of the Eastern Central Atlantic. Volume 3: Bony fishes part 1 (Elopiformes to Scorpaeniformes)*. FAO Species Identification Guide for Fishery Purposes, Rome, 2016.

202. B.J. Laurel, T.P. Hurst, L.A. Copeman, and M.W. Davis. The role of temperature on the growth and survival of early and late hatching Pacific cod larvae (*Gadus macrocephalus*). *Journal of Plankton Research*, 30:1051–1060, 2008.

203. S. Hinckley, W.T. Stockhausen, K.O. Coyle, B.J. Laurel, G.A. Gibson, C. Parada, A.J. Hermann, M.J. Doyle, T.P. Hurst, A.E. Punt, and C. Ladd. Connectivity between spawning and nursery areas for Pacific cod (*Gadus macrocephalus*) in the Gulf of Alaska. *Deep Sea Research Part II: Topical Studies in Oceanography*, 165:113–126, 2019.

204. M.D. Guiry and P. Kuipers. *Sargassum muticum* Wireweed. https://www.seaweed.ie/sargassum, 2023. Accessed: July 2021–December 2023.

205. Wikipedia contributors. Common murre. https://en.wikipedia.org/wiki/Common_murre, 2023. Accessed: July 2021–December 2023.

206. J.L. Padilla-Gamiño and R.D. Gates. Spawning dynamics in the Hawaiian reef-building coral *Montipora capitata*. *Marine Ecology Progress Series*, 449:145–160, 2012.

207. J.D. Miller, C.J. Limpus, and M.H. Godfrey. Chapter 8: Nest site selection, oviposition, eggs, development, hatching, and emergence of loggerhead turtles. In *Loggerhead Sea Turtles*, pages 125–143. Smithsonian Books, 2003.

208. N. Takeshita, N. Onikura, S. Matsui, and S. Kimura. Embryonic, larval and juvenile development of the roughskin sculpin, *Trachidermus fasciatus* (Scorpaeniformes: Cottidae). *Ichthyological Research*, 44:257–266, 1997.

209. S. Miyashita, K. Kato, Y. Sawada, O. Murata, Y. Ishitani, K. Shimizu, S. Yamamoto, and H. Kumai. Development of digestive system and digestive enzyme activities of larval and juvenile bluefin tuna, *Thunnus thynnus*, reared in the laboratory. *Suisanzoshoku*, 46:111–120, 1998.

210. NOAA Fisheries. Loggerhead turtle. https://www.fisheries.noaa.gov/species/loggerhead-turtle, 2014. Accessed: July 2021–December 2023.

211. H. Knutsen, P.E. Jorde, O.A. Bergstad, and M. Skogen. Population genetic structure in a deepwater fish *Coryphaenoides rupestris*: patterns and processes. *Marine Ecology Progress Series*, 460:233–246, 2012.

212. P. Casale and A.D. Tucker. *Caretta caretta* (amended version of 2015 assessment). *The IUCN Red List of Threatened Species 2017*, e.T3897A119333622, 2017.

213. Stazione zoologica di Napoli. *Fauna e Flora del Golfo di Napoli, Monografia 38: Uova, Larve e Stadi Giovanili di Teleostei*. Springer-Verlag, 1892. Monographs elaborated with the use of material collected and serialized by S. Lo Bianco from U. D'Ancona, L. Sanzo, A. Sparta, F. Bertolini, G. Montalenti, E. Padoa, S. Ranzi, E. Tortonese, M. Vialli.

Appendix S6: Search parameters and citations for Global Biodiversity Information Facility species occurrence records

This appendix contains the search parameters and citations for the resulting occurrence records that we downloaded from the Global Biodiversity Information Facility (GBIF) for each species included in this study. The search parameters section denotes the exact search and filtering terms that we passed to the `occ_download` function (rgbif R package V3.5.0, Chamberlain and Boettiger 2017) to retrieve species occurrence records from GBIF for each species in this study. The citations capture the specific snapshots of GBIF data used in this study; the DOIs are specific to the exact search parameters that we used and the date that we executed the query.

Search and filtering parameters used to retrieve GBIF records:

```r
library(rgbif)

OurSearchRequest <- occ_download(type='and',
                                 pred('taxonKey',taxonKey),
                                 pred("hasGeospatialIssue", FALSE),
                                 pred("hasCoordinate", TRUE),
                                 pred_not(
                                   pred("basisOfRecord", "FOSSIL_SPECIMEN")
                                 ),
                                 pred_or(
                                   pred_not(pred("establishmentMeans","MANAGED")),
                                   pred_not(pred_notnull("establishmentMeans"))
                                 ),
                                 pred_or(
                                   pred("occurrenceStatus","PRESENT"),
                                   pred_not(pred_notnull("occurrenceStatus"))
                                 ),
                                 format='SIMPLE_CSV',
                                 user=user,pwd=password,email=email)
```

Citations

1. GBIF.org. GBIF occurrence download for species: *Acanthephyra purpurea*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.TFGCY5`.

2. GBIF.org. GBIF occurrence download for species: *Acanthurus olivaceus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.UVAHKV`.

3. GBIF.org. GBIF occurrence download for species: *Acropora cervicornis*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.36NY9X`.

4. GBIF.org. GBIF occurrence download for species: *Acropora palmata*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.AU6DJV`.

5. GBIF.org. GBIF occurrence download for species: *Acropora prolifera*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.WZCUDM`.

6. GBIF.org. GBIF occurrence download for species: *Agaricia fragilis*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.B9X8M6`.

7. GBIF.org. GBIF occurrence download for species: *Agaricia undata*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.T3Y6RZ`.

8. GBIF.org. GBIF occurrence download for species: *Amphiprion bicinctus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.TSQNZD`.

9. GBIF.org. GBIF occurrence download for species: *Anthopleura elegantissima*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.8GXBNZ`.

10. GBIF.org. GBIF occurrence download for species: *Aptenodytes patagonicus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.DCVK9N`.

11. GBIF.org. GBIF occurrence download for species: *Arctocephalus forsteri*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.5AG5D8`.

12. GBIF.org. GBIF occurrence download for species: *Aythya marila*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.QEBGXH`.

13. GBIF.org. GBIF occurrence download for species: *Bartholomea annulata*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.NFGHK2`.

14. GBIF.org. GBIF occurrence download for species: *Bathygobius cocosensis*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.86NGUR`.

15. GBIF.org. GBIF occurrence download for species: *Bathyraja aleutica*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.JAKMBR`.

16. GBIF.org. GBIF occurrence download for species: *Bathyraja panthera*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.C57HRQ`.

17. GBIF.org. GBIF occurrence download for species: *Callinectes sapidus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.2NYJZA`.

18. GBIF.org. GBIF occurrence download for species: *Caretta caretta*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.BH7VFR`.

19. GBIF.org. GBIF occurrence download for species: *Coryphaenoides rupestris*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.ZYK7RM`.

20. GBIF.org. GBIF occurrence download for species: *Cranchia scabra*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.W3P2G7`.

21. GBIF.org. GBIF occurrence download for species: *Ctenolabrus rupestris*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.MNVJY9`.

22. GBIF.org. GBIF occurrence download for species: *Dascyllus trimaculatus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.VMSTXW`.

23. GBIF.org. GBIF occurrence download for species: *Ectopleura larynx*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.GPK9PA`.

24. GBIF.org. GBIF occurrence download for species: *Engraulis encrasicolus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.SC82NB`.

25. GBIF.org. GBIF occurrence download for species: *Eudyptes chrysolophus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.72SJT9`.

26. GBIF.org. GBIF occurrence download for species: *Eukrohnia hamata*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.7FNKN4`.

27. GBIF.org. GBIF occurrence download for species: *Exaiptasia diaphana*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.7DD9HN`.

28. GBIF.org. GBIF occurrence download for species: *Fibramia amboinensis*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.UHFXAZ`.

29. GBIF.org. GBIF occurrence download for species: *Gadus macrocephalus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.A3G3S2`.

30. GBIF.org. GBIF occurrence download for species: *Gadus morhua*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.NMKD65`.

31. GBIF.org. GBIF occurrence download for species: *Galaxea fascicularis*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.DQNHKE`.

32. GBIF.org. GBIF occurrence download for species: *Galaxea horrescens*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.K2MUNY`.

33. GBIF.org. GBIF occurrence download for species: *Galaxias maculatus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.UHQEZM`.

34. GBIF.org. GBIF occurrence download for species: *Holacanthus passer*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.NNSR9J`.

35. GBIF.org. GBIF occurrence download for species: *Homarus americanus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.C5U8NF`.

36. GBIF.org. GBIF occurrence download for species: *Istiblennius lineatus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.Z5CXH6`.

37. GBIF.org. GBIF occurrence download for species: *Lagenorhynchus acutus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.NEWHA4`.

38. GBIF.org. GBIF occurrence download for species: *Laguncularia racemosa*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.RKDKA5`.

39. GBIF.org. GBIF occurrence download for species: *Larimichthys polyactis*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.Y43WRW`.

40. GBIF.org. GBIF occurrence download for species: *Lateolabrax japonicus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.PUB9H8`.

41. GBIF.org. GBIF occurrence download for species: *Leptopsammia pruvoti. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.X5U5WQ`.

42. GBIF.org. GBIF occurrence download for species: *Lutjanus campechanus. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.NHDMZG`.

43. GBIF.org. GBIF occurrence download for species: *Lutjanus fulvus. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.YRXHMU`.

44. GBIF.org. GBIF occurrence download for species: *Montipora capitata. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.53R9JZ`.

45. GBIF.org. GBIF occurrence download for species: *Nautilus pompilius. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.8EBKFQ`.

46. GBIF.org. GBIF occurrence download for species: *Orbicella faveolata. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.85WJZ9`.

47. GBIF.org. GBIF occurrence download for species: *Ostrea lurida. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.TA4X9R`.

48. GBIF.org. GBIF occurrence download for species: *Paracentrotus lividus. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.DCTTTT`.

49. GBIF.org. GBIF occurrence download for species: *Paracirrhites forsteri. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.YR3BTX`.

50. GBIF.org. GBIF occurrence download for species: *Paralichthys dentatus. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.RYM5CK`.

51. GBIF.org. GBIF occurrence download for species: *Penaeus duorarum. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.3TAZ82`.

52. GBIF.org. GBIF occurrence download for species: *Phocoena phocoena. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.W8G4JW`.

53. GBIF.org. GBIF occurrence download for species: *Phocoena sinus. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.545WDZ`.

54. GBIF.org. GBIF occurrence download for species: *Pisaster ochraceus. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.MTVEG5`.

55. GBIF.org. GBIF occurrence download for species: *Planes minutus. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.9VNBHU`.

56. GBIF.org. GBIF occurrence download for species: *Platichthys flesus. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.279XZB`.

57. GBIF.org. GBIF occurrence download for species: *Pocillopora damicornis. The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.VCTSTZ`.

58. GBIF.org. GBIF occurrence download for species: *Pomacanthus maculosus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.95A22Y`.

59. GBIF.org. GBIF occurrence download for species: *Porites astreoides*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.BVRJKB`.

60. GBIF.org. GBIF occurrence download for species: *Porites compressa*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.TMQU9N`.

61. GBIF.org. GBIF occurrence download for species: *Pygoscelis adeliae*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.HEVD2H`.

62. GBIF.org. GBIF occurrence download for species: *Pygoscelis antarcticus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.XSFVMM`.

63. GBIF.org. GBIF occurrence download for species: *Pygoscelis papua*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.F9C8AG`.

64. GBIF.org. GBIF occurrence download for species: *Pyroteuthis margaritifera*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.F5JGYN`.

65. GBIF.org. GBIF occurrence download for species: *Rhizophora mangle*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.RWEYY6`.

66. GBIF.org. GBIF occurrence download for species: *Robustosergia robusta*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.9J5NM3`.

67. GBIF.org. GBIF occurrence download for species: *Sargassum muticum*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.4C6P4V`.

68. GBIF.org. GBIF occurrence download for species: *Scomber scombrus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.G9KUEH`.

69. GBIF.org. GBIF occurrence download for species: *Sebastes diaconus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.SZPU3S`.

70. GBIF.org. GBIF occurrence download for species: *Sebastes paucispinis*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.MJP3JG`.

71. GBIF.org. GBIF occurrence download for species: *Sebastes pinniger*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.PDKCR3`.

72. GBIF.org. GBIF occurrence download for species: *Sebastes rastrelliger*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.XQK8D6`.

73. GBIF.org. GBIF occurrence download for species: *Sebastes ruberrimus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.ZYTDQ9`.

74. GBIF.org. GBIF occurrence download for species: *Sebastiscus marmoratus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.M7G2UC`.

75. GBIF.org. GBIF occurrence download for species: *Seriola lalandi dorsalis*. *The Global Biodiversity Information Facility*, 08 September 2021. URL: https://www.gbif.org/occurrence/download/0013024-210819072339941, `doi:10.15468/DL.S2ZFUB`.

76. GBIF.org. GBIF occurrence download for species: *Siphamia tubifer*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.RVNHSA`.

77. GBIF.org. GBIF occurrence download for species: *Sphyrna tiburo*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.NKFBD9`.

78. GBIF.org. GBIF occurrence download for species: *Stegastes beebei*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.U38T9S`.

79. GBIF.org. GBIF occurrence download for species: *Stegastes leucorus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.X79N5K`.

80. GBIF.org. GBIF occurrence download for species: *Stephanocoenia intersepta*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.Y85A8A`.

81. GBIF.org. GBIF occurrence download for species: *Symphodus melops*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.BRSXN6`.

82. GBIF.org. GBIF occurrence download for species: *Systellaspis debilis*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.Y6SQ7R`.

83. GBIF.org. GBIF occurrence download for species: *Tegula funebralis*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.UTQWT9`.

84. GBIF.org. GBIF occurrence download for species: *Thunnus thynnus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.JMXT2M`.

85. GBIF.org. GBIF occurrence download for species: *Trachidermus fasciatus*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.BERRP6`.

86. GBIF.org. GBIF occurrence download for species: *Uria aalge*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.F5S5AE`.

87. GBIF.org. GBIF occurrence download for species: *Uria lomvia*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.2AHRUG`.

88. GBIF.org. GBIF occurrence download for species: *Vampyroteuthis infernalis*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.EYPAXR`.

89. GBIF.org. GBIF occurrence download for species: *Zalophus wollebaeki*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.77C2WD`.

90. GBIF.org. GBIF occurrence download for species: *Zebrasoma flavescens*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.WGS9RC`.

91. GBIF.org. GBIF occurrence download for species: *Zebrasoma scopas*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.ZJJUCK`.

92. GBIF.org. GBIF occurrence download for species: *Zenarchopterus dunckeri*. *The Global Biodiversity Information Facility*, 08 September 2021. `doi:10.15468/DL.2WC9S7`.

93. GBIF.org. GBIF occurrence download for species: *Halichoerus grypus*. *The Global Biodiversity Information Facility*, 17 January 2022. `doi:10.15468/DL.26TYNY`.