Comparing phenotypic manifolds with Kompot: Detecting differential abundance and gene expression at single-cell resolution

Dominik J. Otto[1,2,3], Erica Arriaga-Gomez[4], Elana Thieme[1,2,3,4,5], Ruijin Yang[1,2,3], Stanley C. Lee[4]+, Manu Setty[1,2,3]+

[1] Basic Sciences Division, Fred Hutchinson Cancer Center, Seattle WA

[2] Computational Biology Program, Public Health Sciences Division, Seattle WA

[3] Translational Data Science IRC, Fred Hutchinson Cancer Center, Seattle WA

[4] Translational Science and Therapeutics Division, Fred Hutchinson Cancer Center, Seattle WA

[5] Molecular and Cellular Biology Program, University of Washington, Seattle WA

+ Corresponding authors:

      Manu Setty (msetty@fredhutch.org)

      Stanley Lee (sclee3@fredhutch.org)

<u>Abstract</u>

Kompot is a statistical framework for holistic comparison of multi-condition single-cell datasets, supporting both differential abundance and differential expression. Differential abundance captures changes in how cells populate the phenotypic manifold across conditions, while differential expression identifies condition-specific changes in gene regulation that may be localized to particular regions of that manifold. Kompot models the distribution of cells and gene expression as continuous functions over a low-dimensional representation of cell states, enabling single-cell resolution inference with calibrated uncertainty estimates. Applying Kompot to aging murine bone marrow, we identified a continuum of shifts in hematopoietic stem cell and mature cell states, transcriptional remodeling of monocytes independent of compositional changes, and divergent regulation of oxidative stress response genes across cell types. By capturing both global and cell-state–specific effects of perturbation, Kompot reveals how aging reshapes cellular identity and regulatory programs across the hematopoietic landscape. This framework is broadly applicable to dissecting condition-specific effects in complex single-cell landscapes.

Introduction

Multi-condition single-cell datasets offer new opportunities to dissect how mutations or other perturbations affect both differentiation trajectories and disease phenotypes[1-4]. Systematic analyses of such datasets require characterization of two key aspects of condition-induced variability: differential abundance and differential expression.

Differential abundance refers to differences in the composition of cell states between two conditions (**Fig. 1A, B**). Even in the absence of overt phenotypic shifts such as loss or gain of broad cell types, changes in the frequency of specific cell states can provide early indicators of disease[5, 6] and response to treatment[7, 8]. These changes are often nuanced and cell-type specific, necessitating statistical approaches that rigorously assess significance while accounting for underlying biological variability.

Perturbations can also reshape the phenotypic landscape itself through differential expression, where gene expression changes that would be unlikely or impossible in unperturbed conditions can emerge[9] (**Fig. 1A, C**). These transcriptional responses reveal condition-specific molecular programs and regulatory networks that often manifest as subtle yet systematic changes across multiple states[9] and can frequently be masked by the inherent variability within biological systems .

Several computational approaches have been developed for comparative analysis of multi-condition single-cell data with differential abundance receiving greater attention[10-15]. A number of these methods rely on discretizations such as clustering cells into discrete groups to stabilize variability estimates. While facilitating interpretation, such discretizations lead to loss in resolution when changes are gradual or occur along continuous trajectories (**Supp. Fig. 1A**). Additionally, many existing methods lack principled significance measures, making it difficult to distinguish genuine biological signals from technical noise.

Here, we introduce Kompot, a novel computational framework that provides a continuous and statistically rigorous approach for differential abundance and differential expression testing using multi-condition single-cell data. Kompot leverages Gaussian Process (GP) modeling to estimate continuous functions across the cell-state space. This approach enables quantification of differential abundance and differential expression at single-cell resolution, while incorporating uncertainty estimates to provide statistically calibrated comparisons. By addressing both aspects of multi-condition analysis within a unified statistical framework, Kompot provides a comprehensive toolkit for extracting biological insights from increasingly complex single-cell datasets across a range of biological systems.

Results

Overview of the Kompot modeling framework

A single-cell RNA-seq dataset provides a snapshot of cellular states in a biological system, with the molecules measured representing a sample of the transcriptome of the cell. To characterize the system and compare it to similar systems under different conditions (such as mutations or other perturbations), we need to understand what this data tells us about the true composition of transcribed genes in a cell and the true distribution of cell states in the system. With Kompot we leverage high-dimensional, continuous cell-state representations and Bayesian Inference to learn both, the composition of the system and transcriptomes of each cell, enabling a holistic comparison of biological systems at single-cell resolution with uncertainty quantification. This comparison is divided into differential cell-state abundance that reflect changes in cell-state composition, and differential-expression that reflect changes in the transcriptome of cells in equivalent states.

The input to Kompot is a latent representation of co-embedded multi-condition single-cell data. This latent representation can incorporate batch effect correction[16, 17] to ensure that cell states considered equivalent share similar locations in that state space (**Supp. Fig. 1B**). From this foundation, Kompot employs

diffusion maps[18] to derive a cell-state representation with biologically meaningful structure (**Fig. 1A**).
Diffusion maps are particularly effective for single-cell analysis as they capture the intrinsic geometry of
the data while providing biologically meaningful measures of cell-to-cell distance[19, 20]. This distance metric
supports both accurate local abundance estimation and principled information sharing when mapping
from cell-state space to gene expression, enabling the statistical comparisons at the core of Kompot's
differential testing framework.

Kompot differential abundance testing

To quantify differences in cell-state composition between conditions, Kompot first constructs condition-
specific density functions on the diffusion map representation. For each condition, Kompot computes a
separate continuous density function (**Fig. 1B, Supp. Fig. 2A-B**) using our Mellon framework[21]. Mellon
leverages nearest-neighbor distances within a Gaussian Process model to estimate continuous density
functions that capture the distribution of cells across the state space[21]. The continuous nature of these
density estimates enables evaluation at any position within the cell-state space, including states
predominantly observed in the alternate condition (**Supp. Fig. 2B**). By computing the ratio of density
estimates at corresponding positions in the cell-state space, Kompot derives a measure of differential
abundance that quantifies the fold-change in cell density between conditions at single-cell resolution
(**Supp. Fig. 2C**).

While the ratio of density estimates provides a measure of fold-change, rigorous comparison between
conditions requires statistical assessment of significance. Kompot addresses this challenge by computing
uncertainty estimates for each density function. These uncertainty estimates account for two critical
factors: limited data availability in regions of the cell-state space and inherent uncertainty through the
sparse function representation. By incorporating these uncertainty estimates into the statistical
framework, Kompot calculates posterior tail probabilities (PTPs), an alternative for p-values, for
differential abundance at each cell state. This enables the generation of volcano plots for differential
abundance testing where each point represents a single cell-state (**Fig. 1B**), allowing for threshold-based
significance assessments. Thus, Kompot can identify subtle cell-state transitions and rare subpopulations
that would likely be undetected by clustering-based approaches, providing insights into the continuous
spectrum of cellular responses to perturbations (**Fig. 1D**).

Kompot differential expression testing

For differential expression testing between conditions, Kompot constructs condition-specific mappings
from cell-states to gene expression space (**Fig. 1A, C**). These mappings are computed using Gaussian
Processes (GPs)[22], by predicting log-transformed gene expression as a continuous function of the
diffusion map coordinates. The non-linear relationship between diffusion maps and gene expression is
effectively captured by GPs[22], essentially serving as a rigorous approach to gene expression imputation.
The continuous nature of these mapping functions allows Kompot to estimate gene expression in one
condition for cell states observed in another, producing counterfactual expression profiles that reflect how
those same cell states would behave under different conditions (**Fig. 1C**). By computing the ratio between
the actual expression in a cell and its counterfactual expression, Kompot quantifies gene expression-fold
changes for every gene at single-cell resolution across the entire phenotypic landscape (**Fig. 1C**).

Given the large number of genes evaluated, we require a robust significance measure that allows us to
identify the most relevant genes, accounting for both uncertainty in the gene expression function and
statistical interdependence of imputed values. Kompot therefore employs the Mahalanobis distance,
which measures the distance of a point from a multivariate normal distribution, considering both variance
and covariance[23]. It effectively measures how many standard deviations a gene expression shift is away

from expected variation, while accounting for the correlation structure between neighboring cell states,
making it an ideal significance measure for differential expression. Kompot therefore uses Mahalanobis
distances as the score in a quasi-volcano plot (**Fig. 1C**) to identify significantly differentially expressed
genes either in the full population of cells or in specific subsets (**Fig. 1D**).

Differential Abundance in Aging Bone Marrow

Aging is a compelling biological context for evaluating differential abundance methods, as it induces
progressive and often subtle shifts in cellular composition rather than dramatic phenotypic
transformations. Hematopoiesis has emerged as a particularly informative system for studying aging[24],
characterized by declining stem cell function[25], reduced lymphocyte production[26], and increased
susceptibility to myeloid malignancies[27]. To investigate these age-related changes, we generated a novel
scRNA-seq dataset of murine bone marrow at 10, 63, and 103 weeks of age, representing young, middle-
aged, and older timepoints (**Supp. Fig. 3A**). After quality control verification (**Supp. Fig. 3B, C**), we
successfully identified most hematopoietic cell types across all ages (**Fig. 2A, B, Supp. Fig. 4**), providing
us with a rich dataset to characterize cell-state changes and gene expression changes in aging
hematopoiesis using Kompot.

We applied Kompot differential abundance testing on our aging hematopoiesis dataset to determine age-
dependent composition changes (**Fig. 2C-E**). We first derived a unified cell-state representation for all
cells using diffusion maps. We then calculated age-specific density functions using the subsets of cells
from each timepoint (**Supp. Fig. 5A-B**). We evaluated these individual density functions at all cell-states
in the dataset to derive condition-specific densities and associated uncertainties (**Supp. Fig. 5C-D**). This
approach enabled pairwise comparisons that quantify age-related changes in cell-type composition (**Fig.
2C-D**). A key advantage of our framework is the ability to set thresholds based on both effect size and
statistical significance at single-cell resolution, allowing for identification of cell states with significant
differential abundance rather than relying solely on fold-change cutoffs (**Fig. 2C-E**).

Our analysis revealed significant expansions and contractions in distinct cell populations over time.
Comparing young to old mice, we observed an expansion of the hematopoietic stem cell (HSC)
compartment and a marked shift from naïve to memory cells in both B-cell and T-cell compartments (**Fig.
2F-G**). The depletion of naïve B-cells and an enrichment of memory B-cells in older mice is a well-
characterized phenomenon of hematopoietic aging[26]. This shift reflects fundamental changes in
differentiation dynamics: naive B-cells rely on a continuous influx from hematopoietic progenitors,
whereas memory B-cells accumulate over time due to antigen exposure. The observed decline in naïve
B-cells reflects the reduced lymphoid differentiation potential of aged HSCs[28], while the expansion of
memory B-cells, including age-associated B-cells, is driven by persistent immune activation[29]. Similarly,
the T-cell compartment displayed a significant reduction in naïve T-cells with a concurrent increase in
memory T-cells, a hallmark of immunosenescence that contributes to diminished adaptive immune
responses in aging[30].

An increased accumulation of hematopoietic stem cells (HSCs) is the most significant change we
observed in the progenitor cells (**Fig. 2F-G**). Progressive enrichment of HSCs with age is a well-
documented phenomenon[31]. HSCs reside at the top of the hematopoietic hierarchy, maintaining blood
production through self-renewal and differentiation into mature immune and blood cells. With aging, HSC
function declines, leading to accumulation in the bone marrow while their ability to differentiate into
different lineages becomes impaired[31]. While the accumulation of aged HSCs has been previously
observed, it has traditionally been quantified using bulk measurements or clustering-based approaches.
We took advantage of the single-cell resolution of Kompot to better characterize the age-induced change
in HSCs. By comparing the differential abundance fold-change along pseudotime ordering[19], we
observed that the expansion of HSC population is in fact a subtle shift in cell-state, rather than a global

increase in a predefined population (**Fig. 2H**). Specifically, HSCs in older mice progressively shift towards
a less differentiation-primed state, creating a continuum of altered stem cell states rather than a uniform
expansion (**Fig. 2H**). This subtle but consequential repositioning along the differentiation trajectory
provides a possible explanation for the functional decline in aged HSCs, revealing how changes in cell
abundance are accompanied by fundamental alterations in cellular function and developmental potential.

Comparing differential abundance results across ages, we observed that changes in mature cell
populations predominantly occurred between young and middle-aged mice (**Fig. 2D, Supp. Fig. 6A**),
while alterations in the HSC compartment were more pronounced between middle-aged and older mice
(**Fig. 2E, Supp. Fig. 6B**). This temporal separation suggests a sequential progression of aging, where
compositional shifts in differentiated cells precede fundamental changes in stem cell states. The early
alterations in mature immune cells may represent initial adaptive responses to aging, while the later HSC
changes potentially indicate more persistent reprogramming of the hematopoietic system.

Our results demonstrate the ability of Kompot to identify subtle but biologically meaningful abundance
changes between conditions. By combining a continuous representation of cell states with robust
statistical measures, Kompot enables precise detection of population-level changes while maintaining
single-cell resolution.

Differential Gene Expression in Aging Hematopoiesis

We applied Kompot differential expression testing on our mouse aging hematopoiesis dataset to
characterize age-related transcriptional changes (**Fig. 3**). We first inferred age-specific mapping functions
from diffusion map coordinates to log-normalized gene expression values, using the subset of cells from
each timepoint. We then used these mappings to impute gene expression and associated uncertainty at
all cell-states across the dataset. This enabled us to compute gene-level fold changes by comparing
observed expression values to their counterfactual predictions under alternative age conditions. To
assess significance, we calculated Mahalanobis distances for each gene across specified groups of cells,
quantifying deviations from expected variation while accounting for the covariate structure among
neighboring cell states. We excluded T and NK cells for this analysis since do not differentiate in the bone
marrow (**Fig. 3H**).

We first applied our approach to characterize gene expression changes in HSCs, a population known to
undergo functional and transcriptional changes with age[31], by comparing HSCs from middle-aged to old-
aged mice (**Fig. 3A-B**). Aging in HSCs has been extensively studied, and the Hematopoietic Aging Atlas
has synthesized results from multiple datasets to identify genes consistently up- or downregulated with
age[32]. For each gene, this atlas assigns a consistency score that reflects the reproducibility of age-
associated expression changes across studies[32]. Kompot's differential expression results strongly
prioritized genes with the highest consistency scores in the Hematopoietic Aging Atlas, with genes
showing large fold changes and Mahalanobis distances highly enriched for reproducible aging signatures
(**Fig. 3C**, **Supp. Fig. 7**).

Our differential abundance analysis revealed that HSCs in older mice shift towards a less differentiated
state along the cell-state continuum (**Fig. 2H**). We next asked whether the genes differentially expressed
in aged HSCs reflected this shift. Indeed, we found that gene expression fold changes were structured
along the same continuum: genes exhibited larger fold changes in cell states that were more differentially
abundant (**Fig. 3D**). These results highlight the transcriptional changes in HSCs that accompany and
may underlie the shift toward a less differentiated state with age,

We next investigated age-related gene expression changes in mature cell types. We focused on changes
from young to old mice since the abundance changes in mature cell types were most pronounced in this
comparison (**Fig. 2D-F**). To distinguish expression changes from abundance-driven effects, we restricted

our analysis to cell types that *did not show significant compositional changes*. Monocytes exhibited clear
transcriptional changes despite stable abundance (**Fig. 3E-F**). Gene ontology analysis revealed that
downregulated genes were enriched in core monocyte/macrophage functions such as defense response
(**Fig. 3G**), consistent with their reduced functional capacity with age[33]. In contrast, upregulated genes
were enriched for antigen processing and presentation pathways, driven by increased expression of MHC
class II genes (**Fig. 3G**), which have been implicated in age-related immune remodeling[34]. Notably, our
results recapitulate findings from a previous study that identified H2-Aa, H2-Ab1, H2-Eb1, Cd74, and
Aw112010 as the top dysregulated genes in aging monocytes across both mice and humans, with only
Igkc surpassing H2-Eb1 in our ranking[34]. The near-perfect agreement with experimentally validated gene
sets underscores the power of our differential expression testing approach to detect biologically
meaningful transcriptional shifts. Gene expression differences in other mature cell types were not as
pronounced but did show a tendency for upregulation of MHC Class II genes similar to monocytes (**Supp.**
Fig. 8).

Our use of Mahalanobis distance as a significance measure enables Kompot to identify genes that exhibit
coordinated but opposing changes across subpopulations. For example, consider a gene that is
upregulated in one cell type and downregulated in another. While the net fold change may be close to
zero, the Mahalanobis distance remains high, reflecting a statistically significant deviation from expected
variation if unaffected. This occurs because such a pattern is unlikely under the local covariance structure
of the cell-state space, where neighboring states are typically expected to exhibit similar expression
patterns. As a result, Kompot can detect structured differential expression that would be missed by
conventional fold-change-based approaches.

To test this capability, we applied Kompot differential expression testing between young and old cells
across all cell-types that differentiate in the bone marrow (**Fig. 3H-I**). The top genes with both high positive
fold change and high Mahalanobis distance were strongly enriched for MHC class II genes, consistent
with our monocyte-specific results and indicating that MHC class II upregulation is a general feature of
hematopoietic aging (**Fig. 3J**). Likewise, genes with high Mahalanobis distance and strong negative fold
change were consistently downregulated across multiple cell types (**Fig. 3J**). In contrast, genes ranked
highly by Mahalanobis distance but with low absolute fold change exhibited divergent expression patterns
across cell types, with upregulation in some and downregulation in others (**Fig. 3J**). To illustrate these
distinctions, we visualized expression levels of representative genes from each category across the cell-
state manifold (**Fig. 3K**). Genes selected by fold change showed broad, monotonic shifts in expression,
whereas Mahalanobis-selected genes revealed distinct, cell-type-specific regulation (**Fig. 3K**).

We next performed gene ontology analysis to assess the biological relevance of genes with high
Mahalanobis distance but low absolute fold change. Antioxidant activity emerged as the most significantly
enriched term (**Fig. 3L**), a pathway known to be impacted by aging[35]. Notably, our results indicate that
while antioxidant activity is broadly impacted across hematopoietic cell types, the direction of expression
changes of the pathway genes varies across genes and cell types, suggesting a context-specific
adaptation to aging (**Fig. 3M**). Oxidative stress is a known driver of HSC dysfunction, promoting DNA
damage, apoptosis, and biased differentiation toward myeloid lineages[35]. The differential expression of
antioxidant pathway genes observed in our study likely represents a compensatory mechanism to
mitigate reactive oxygen species (ROS) accumulation in aged HSCs[35].

These results demonstrate that Mahalanobis distance captures biologically meaningful heterogeneity in
gene expression and provides a powerful approach for aggregating structured expression differences
across cell states without requiring prior clustering, making it especially suited for complex, continuous
single-cell landscapes.

Discussion

Kompot provides a flexible statistical framework for quantifying condition-specific effects in single-cell data, modeling both cell abundance and gene expression as continuous functions over the cell-state manifold. By estimating condition-specific densities and expression profiles using Gaussian processes, Kompot performs differential analysis at single-cell resolution while accounting for uncertainty and spatial structure in the data. This approach supports biologically grounded interpretations of perturbation effects, whether they manifest as global shifts in population structure, localized transcriptional responses, or context-specific coordination between abundance and expression changes.

Applying Kompot to aging murine bone marrow revealed both well-established and previously unrecognized features of hematopoietic aging. Kompot recapitulated known expansions in memory lymphocytes and age-associated B cells, while resolving more subtle shifts in hematopoietic stem cell states that do not correspond to discrete population changes. In the transcriptomic landscape, Kompot uncovered gene expression programs that are either conserved or cell-type specific, including divergent regulation of oxidative stress pathways. These findings demonstrate how modeling continuous phenotypic structure supports more nuanced detection of biological change. Beyond aging, Kompot is broadly applicable to studies of development, disease progression, and response to perturbation, where condition-induced effects often unfold along continuous, high-dimensional trajectories.

Methods

Kompot framework

Kompot is a tool for comparing single-cell phenotypic manifolds across conditions. It provides frameworks for both differential abundance testing and differential gene expression testing at single-cell resolution using multi-condition datasets. Differential abundance testing enables the detection of abundance shifts of cell-states across the phenotypic manifold, without requiring predefined cell-type boundaries. This is achieved by estimating condition-specific cell-state densities using the Bayesian inference scheme implemented in Mellon. In parallel, through differential gene expression testing, Kompot identifies gene expression changes across the same manifold, capturing subtle but structured transcriptional shifts that may vary across cell states and may not follow consistent directionality To do so, Kompot leverages the Gaussian process inference implemented in Mellon to estimate condition-specific gene expression functions for thousands of genes within minutes, while explicitly sharing information across neighboring cell states to address sparsity and enable statistically meaningful comparisons

Differential Abundance

This section explains the methods of Kompot's differential abundance (DA) analysis framework. Our density quantification relies on the Gaussian Process (**GP**) estimation implemented in Mellon. This estimation process is a Bayesian inference that provides us with both a "best estimate" function that maps each point in the cell-state space to a density value $\mu(x_i)$, and a quantification of uncertainty $\sigma(x_i)$. Since we infer these quantities independently for each condition, the posterior distribution for the density change from condition a to condition b at any cell state x_i is given by

$$\delta(x_i) \sim \mathcal{N}\left(\mu_b(x_i) - \mu_a(x_i), \sigma_a^2(x_i) + \sigma_b^2(x_i)\right).$$

Therefore, our density log-fold change is defined by $\mu_b(x_i) - \mu_a(x_i)$, and our uncertainty quantification by $\sigma_a^2(x_i) + \sigma_b^2(x_i)$. Notably, using the correct intrinsic dimensionality, Mellon reports density in the unit log number of cells per volume.

To make density values comparable across samples with different cell counts, we can transform this unit into log fraction of cells per volume by subtracting the log total number of cells. Therefore, we normalize density in Kompot using

$$\mu'_a(x_i) = \mu_a(x_i) - \log N_a$$

where N_a represents the total number of cells measured for condition a. For this normalization to be valid, it is essential, that we assume the correct intrinsic dimensionality of the phenotypic manifold. To that end we use Mellon dimensionality estimation that is assumed to be constant across the dataset.

Finally, we can define the Kompot's local density log-fold change statistic as

$$\Delta(x_i) = \mu_b(x_i) - \mu_a(x_i) - \log {N_a}/{N_b}.$$

Additionally, we provide a p-value-like significance value through the posterior tail probability (**PTP**). For example, if the density change $\Delta(x_i)$ is positive, then we compute $\mathbb{P}(\delta(x_i) < 0)$, and $\mathbb{P}(\delta(x_i) > 0)$ when $\Delta(x_i)$ is negative. The PTP quantifies the probability that the true fold change goes into the opposite direction of our best estimate, given our data and modeling assumptions. Since our posterior distribution is a normal distribution, the local Kompot PTP is given by:

318
$$PTP(x_i) = \Phi\left(\frac{-|\Delta(x_i)|}{\sqrt{\sigma_a^2(x_i) + \sigma_b^2(x_i)}}\right)$$

were Φ is the cumulative distribution function of the standard normal distribution.

Differential Expression

Similar to the density differences, we also leverage Mellon's GP implementation (**Supp. Note 1**) for
differential expression (DE) analysis. In contrast to the density inference algorithm, this DE method
directly conditions the gene-expression functions on the measured data, as classically done in GPs. For
each condition, the goal is to compute a gene-expression function that predicts an expression value for
each gene at any location in the cell-state space, specific to that condition. These functions provide a
smooth and continuous representation of gene expression across the manifold and allow for quantifying
differential expression between conditions at any point in state space, effectively integrating all data
points without requiring predefined cluster boundaries.

The resulting gene-expression functions and uncertainty estimation can be used equivalently to the cell-
state density functions to compute local gene-expression differences, featuring local fold change and
PTP. However, apart from the question in which cell-state a gene may be differentially expressed, we
want to answer if a gene asserts any significant DE within a cell population or across the entire cell-state
space, effectively aggregating measurements across multiple cells.

This comes with an important caveat: The GP based gene-expression functions are an imputation that
already accumulates measurements of multiple cells, and the resulting gene-expression values $f(x_i)$ at
different cell states x_i are not independent measurements that can be aggregated directly. This caveat
can be addressed very directly through Mellon's uncertainty quantification: It covers not only the variance
for a function value $f(x_i)$ but also the covariance to other function values $f(x_j)$ based on the similarity of
x_i and x_j. This covariance can be understood as a statistical connection between the expressions of two
cell states that comes from our assumption that genes change expression smoothly throughout cell-state
space. Formally, under our modelling assumption, the true gene expression values $f(x)$ at cell states
$x = \{x_i\}_{i=1}^m$ are distributed according to a posterior

344
$$f(x) \sim \mathcal{N}\big(\mu(x), \Sigma(x)\big).$$

345   Therefore, the significance for an expressional change between two conditions k and l can be quantified
346   through the Mahalanobis distance (**Supp. Note 2**):

347
$$D(a,b) = \sqrt{(\mu_a - \mu_b)^T (\Sigma_a + \Sigma_b)^{-1} (\mu_a - \mu_b)}$$

Notably, this adequately accounts for redundancy, as adding a nearly identical cell state contribute nearly
nothing to the value of this statistic, while more distinct cell-states have a larger contribution. We leverage
this to our advantage. By producing a fixed set of landmarks, using k-means clustering, we limit the
evaluation of the gene-expression function and the size of the covariance matrix to a fixed size. By using
a fixed number of steps and k++ initialization, both the clustering and the GP only grow linearly in run-
time with the number of cells. Notably, k-means grows with $O(t \times k \times d \times n)$ where t is the number of
iterations, k the number of landmarks, d the dimensionality of the space, and n the number of cells.

Data analysis

Detailed descriptions of data generation and analysis described in the manuscript will be included in a subsequent version of the manuscript.

Data Availability
Raw single-cell data for the murine aging hematopoiesis will be deposited to GEO. Processed anndata object with annotations is available to download from Zenodo[36].

Code Availability
Kompot is available as a Python module at https://github.com/settylab/Kompot. Jupyter notebooks detailing the usage of Kompot differential expression and differential abundance testing are available at https://github.com/settylab/kompot/tree/main/examples and documentation at https://kompot.readthedocs.io/en/latest/.

Acknowledgements
This study was supported by National Institute of General Medical Studies grant no. R35 GM147125 to M.S.; National Cancer Institute grant no. R01CA292932, Mark Foundation for Cancer Research Endeavor Award and Edward P. Evans Foundation Discovery Research Grant to S.C.L; Brotman Baty Institute Pilot Award and Translational Data Science IRC New Collaboration Award to M.S. and S.C.L; National Institute of General Medicine Studies grant no. T32GM136534 to E.T.; National Institutes of Health grant no. ORIP S10OD028685 to support high-performance computing at the Fred Hutchinson Cancer Center.

Competing Interests
The authors declare no competing interests.

Figures

Figure 1: Overview of Kompot's framework for differential abundance and expression analysis.

(A) Illustration of high-dimensional single-cell transcriptomic data from two conditions, projected into a shared low-dimensional cell-state space using diffusion maps.

(B) Differential abundance analysis in Kompot. Condition-specific cell-state densities with associated uncertainty estimates are computed and compared to yield density log-fold changes. Significance is quantified using posterior tail probability (PTP), visualized in a quasi-volcano plot.

(C) Differential expression analysis. Left: schematic of inferred phenotypic manifolds for both conditions with counterfactual cell pairs illustrating expression comparison. Middle: gene-specific differences are illustrated for one example gene, showing condition-specific expression levels, statistical interdependence, and uncertainty. Right: A quasi-volcano plot summarizes gene-level results with the average log-fold change on the x-axis and Mahalanobis distance (accounting for variance and covariance) as a multivariate significance score on the y-axis.

(D) Visualization of differential abundance (left) and differential expression (right) across the cell-state space, colored by log-fold change magnitude.

Figure 2: Kompot differential abundance analysis in aging hematopoiesis.

(A) UMAP projection of aging murine bone marrow scRNA-seq dataset, colored by medium-resolution cell types.

(B) UMAPs highlighting cells from mice grouped by age: Young (10 weeks), Mid (63 weeks), and Old (103 weeks).

(C) Differential abundance results for the Young to Mid (left), Young to Old (middle) and Mid to Old (right) comparisons. Quasi-cell volcano plot with density log-fold change on the x-axis and Posterior Tail Probabilities (PTP) on the y-axis. Significantly different cell-states (absolute log fold change > 1, PTP < 10^{-3}) are highlighted by their cell type annotations from (A).

(D) UMAPs displaying the estimated local density log-fold changes across the full joint cell-state space derived from all samples (Young, Mid, and Old) for comparisons in C.

(E) UMAPs highlighting the subset of cell states with statistically significant abundance changes, as defined by the thresholds shown in the volcano plots in C.

(F) Differential abundance volcano plots showing density log-fold changes across cells, stratified by major cell type and colored by subtype in the Young to Old comparison. y-axis indicates posterior tail probability. Violin plots in the background indicate abundance of cells at the respective density log-fold change in this group.

(G) Direction and magnitude of abundance changes per cell type in the Young to Old comparison. Bars

show the fraction of significantly changing cell-states within each annotated cell type, color-coded by the

direction of change. Cell types are ordered along a progenitor-to-mature axis.

(H) Kompot results for early hematopoietic progenitors. Left: UMAP highlighting HSC, and LMPP subsets.

Right: Scatterplot of cells colored by density log-fold change along pseudotime (x-axis), showing dynamic

abundance shifts during early differentiation.

Figure 3: Kompot differential expression analysis in aging hematopoiesis.

(A) UMAP from Fig. 2A, with HSCs highlighted.

(B) Differential expression results for Mid to Old HSCs. Quasi-cell volcano plot with expression log-fold change on the x-axis and Mahalanobis distance on the y-axis. Significantly different genes (absolute log fold change > 0.15, Mahalanobis distance > 3) are highlighted.

(C) like (B) colored by weighted average of reported "Aging Consistency" using standard Gaussian Kernel after scaling x- and y-axis to standard deviation 1.

(D) Same plot as Fig. 2H, colored by Mid to Old HSC gene expression log fold change for a gene downregulated with age (Cd52, left) and an upregulated gene (Ifitm1, right).

(E) UMAP from Fig. 2A, with monocytes highlighted.

(F) Same as (B), for Young to Old monocytes.

(G) Gene ontology analysis from STRING[37] for downregulated genes (left) and upregulated genes (right) from Fig. 2F.

(H) UMAP for Fig. 2A, highlighting all cell types except T and NK cells

(I) Differential expression results for all cells in Fig. 2H for Young to Old comparisons. Top 10 genes with
positive fold change and high Mahalanobis distance are shown in red, bottom 10 genes with negative
fold change and high Mahalanobis distance are shown in blue, and genes with log absolute fold change
but high Mahalanobis distance are shown in green (absolute log fold change < 0.07, Mahalanobis
distance > 6). 20 genes are labeled.

(J) Heatmap showing Young to Old gene expression log fold changes across different cell types for genes
highlighted in (I).

(K) Top: Imputed expression of gene H2-Q7 using young cells (left), using old cells (right) and fold change
from young to old (middle). Middle, Bottom: Same as the top row for Mef2c and Dynll1. These genes
serve as examples of genes with high positive fold change, high Mahalanobis but low absolute fold
change, and high negative fold change respectively.

(L) Gene ontology analysis using genes with high Mahalanobis distance and low absolute fold change
from (I).

(M) Heatmap showing Young to Old gene expression log fold changes across different cell types for
genes involved in antioxidant activity.

References

1. Tabula Muris C. A single-cell transcriptomic atlas characterizes ageing tissues in the mouse. Nature. 2020;583(7817):590-5, PMC8240505.

2. Konturek-Ciesla A, Dhapola P, Zhang Q, Sawen P, Wan H, Karlsson G, et al. Temporal multimodal single-cell profiling of native hematopoiesis illuminates altered differentiation trajectories with age. Cell Rep. 2023;42(4):112304.

3. Herault L, Poplineau M, Mazuel A, Platet N, Remy E, Duprez E. Single-cell RNA-seq reveals a concomitant delay in differentiation and cell cycle of aged hematopoietic stem cells. BMC Biol. 2021;19(1):19, PMC7851934.

4. Kucinski I, Campos J, Barile M, Severi F, Bohin N, Moreira PN, et al. A time- and single-cell-resolved model of murine bone marrow hematopoiesis. Cell Stem Cell. 2023.

5. Gazestani V, Kamath T, Nadaf NM, Dougalis A, Burris SJ, Rooney B, et al. Early Alzheimer's disease pathology in human cortex involves transient cell states. Cell. 2023;186(20):4438-53 e23, PMC11107481.

6. Mayr CH, Simon LM, Leuschner G, Ansari M, Schniering J, Geyer PE, et al. Integrative analysis of cell state changes in lung fibrosis with peripheral protein biomarkers. EMBO Mol Med. 2021;13(4):e12871, PMC8033531.

7. Rehman SK, Haynes J, Collignon E, Brown KR, Wang Y, Nixon AML, et al. Colorectal Cancer Cells Enter a Diapause-like DTP State to Survive Chemotherapy. Cell. 2021;184(1):226-42 e21, PMC8437243.

8. van der Leun AM, Thommen DS, Schumacher TN. CD8(+) T cell states in human cancer: insights from single-cell analysis. Nat Rev Cancer. 2020;20(4):218-32, PMC7115982.

9. Replogle JM, Saunders RA, Pogson AN, Hussmann JA, Lenail A, Guna A, et al. Mapping information-rich genotype-phenotype landscapes with genome-scale Perturb-seq. Cell. 2022;185(14):2559-75 e28, PMC9380471.

10. Burkhardt DB, Stanley JS, 3rd, Tong A, Perdigoto AL, Gigante SA, Herold KC, et al. Quantifying the effect of experimental perturbations at single-cell resolution. Nat Biotechnol. 2021;39(5):619-29, PMC8122059.

11. Dann E, Henderson NC, Teichmann SA, Morgan MD, Marioni JC. Differential abundance testing on single-cell data using k-nearest neighbor graphs. Nat Biotechnol. 2022;40(2):245-53.

12. Yi H, Plotkin A, Stanley N. Benchmarking differential abundance methods for finding condition-specific prototypical cells in multi-sample single-cell datasets. Genome Biology. 2024;25(1):9.

13. Missarova A, Dann E, Rosen L, Satija R, Marioni J. Leveraging neighborhood representations of single-cell data to achieve sensitive DE testing with miloDE. Genome Biol. 2024;25(1):189, PMC11256449.

14. Boyeau P, Hong J, Gayoso A, Kim M, McFaline-Figueroa JL, Jordan MI, et al. Deep generative modeling of sample-level heterogeneity in single-cell genomics. bioRxiv. 2024:2022.10.04.510898.

15. Ahlmann-Eltze C, Huber W. Analysis of multi-condition single-cell data with latent embedding multivariate regression. Nat Genet. 2025;57(3):659-67, PMC11906359.

16. Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K, et al. Fast, sensitive and accurate integration of single-cell data with Harmony. Nat Methods. 2019;16(12):1289-96, PMC6884693.

17. Luecken MD, Buttner M, Chaichoompu K, Danese A, Interlandi M, Mueller MF, et al. Benchmarking atlas-level data integration in single-cell genomics. Nat Methods. 2022;19(1):41-50, PMC8748196.

18. Coifman RR, Lafon S, Lee AB, Maggioni M, Nadler B, Warner F, et al. Geometric diffusions as a tool for harmonic analysis and structure definition of data: diffusion maps. Proc Natl Acad Sci U S A. 2005;102(21):7426-31, PMC1140422.

19. Setty M, Kiseliovas V, Levine J, Gayoso A, Mazutis L, Pe'er D. Characterization of cell fate probabilities in single-cell data with Palantir. Nat Biotechnol. 2019;37(4):451-60.

20.   Haghverdi L, Buttner M, Wolf FA, Buettner F, Theis FJ. Diffusion pseudotime robustly reconstructs
lineage branching. Nat Methods. 2016;13(10):845-8.

21.   Otto D, Jordan C, Dury B, Dien C, Setty M. Quantifying Cell-State Densities in Single-Cell Phenotypic
Landscapes using Mellon. Nature Methods. 2024, PMC10369887.

22.   Rasmussen CE, Williams CKI. Gaussian Processes for Machine Learning2006.

23.   PC M. Reprint of: Mahalanobis, P.C. (1936) "On the Generalised Distance in Statistics.". Sankhya A.
2018;80(1):1-7.

24.   Jaiswal S, Ebert BL. Clonal hematopoiesis in human aging and disease. Science. 2019;366(6465),
PMC8050831.

25.   Chambers SM, Shaw CA, Gatza C, Fisk CJ, Donehower LA, Goodell MA. Aging hematopoietic stem cells
decline in function and exhibit epigenetic dysregulation. PLoS Biol. 2007;5(8):e201, PMC1925137.

26.   Li H, Cote P, Kuoch M, Ezike J, Frenis K, Afanassiev A, et al. The dynamics of hematopoiesis over the
human lifespan. Nat Methods. 2025;22(2):422-34, PMC11908799.

27.   Appelbaum FR, Gundacker H, Head DR, Slovak ML, Willman CL, Godwin JE, et al. Age and acute
myeloid leukemia. Blood. 2006;107(9):3481-5, PMC1895766.

28.   Florian MC, Klose M, Sacma M, Jablanovic J, Knudson L, Nattamai KJ, et al. Aging alters the epigenetic
asymmetry of HSC division. PLoS Biol. 2018;16(9):e2003389, PMC6168157.

29.   Bulati M, Caruso C, Colonna-Romano G. From lymphopoiesis to plasma cells differentiation, the age-
related modifications of B cell compartment are influenced by "inflamm-ageing". Ageing Res Rev. 2017;36:125-
36.

30.   Goronzy JJ, Weyand CM. Immune aging and autoimmunity. Cell Mol Life Sci. 2012;69(10):1615-23,
PMC4277694.

31.   de Haan G, Lazare SS. Aging of hematopoietic stem cells. Blood. 2018;131(5):479-87.

32.   Flohr Svendsen A, Yang D, Kim K, Lazare S, Skinder N, Zwart E, et al. A comprehensive transcriptome
signature of murine hematopoietic stem cell aging. Blood. 2021;138(6):439-51.

33.   De Maeyer RPH, Chambers ES. The impact of ageing on monocytes and macrophages. Immunol Lett.
2021;230:1-10.

34.   Barman PK, Shin JE, Lewis SA, Kang S, Wu D, Wang Y, et al. Production of MHCII-expressing classical
monocytes increases during aging in mice and humans. Aging Cell. 2022;21(10):e13701, PMC9577948.

35.   Shao L, Li H, Pazhanisamy SK, Meng A, Wang Y, Zhou D. Reactive oxygen species and hematopoietic
stem cell senescence. Int J Hematol. 2011;94(1):24-32, PMC3390185.

36.   Otto D, Arriaga-Gomez E, Thieme E, Yang R, Lee S, Setty M. Single-cell CITE-seq of murine bone
marrow across aging (Young, Mid, Old): Zenodo; 2025 [Available from: https://doi.org/10.5281/zenodo.15587767.

37.   Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, et al. STRING v11: protein-protein
association networks with increased coverage, supporting functional discovery in genome-wide experimental
datasets. Nucleic Acids Res. 2019;47(D1):D607-D13, PMC6323986.