Title:

# TESS: A Forward Simulation Framework for Studying the Role of

# Transposable Elements in Genome Expansion and Contraction

# Authors:

Christopher W. Benson[1][†], Ting-Hsuan Chen[2][†][*], Tianyu Lu[1], Olivia Angelin-Bonnet[3], Susan J.

Thomson[2], Cecilia H. Deng[4], Shujun Ou[1][*]

1 Department of Molecular Genetics, Ohio State University, Columbus, OH, 43210, USA

2 The New Zealand Institute for Plant and Food Research Limited, Lincoln 7608, New Zealand

3 The New Zealand Institute for Plant and Food Research Limited, Palmerston North 4442, New

Zealand

4 The New Zealand Institute for Plant and Food Research Limited, Auckland 1025, New

Zealand

† Christopher W. Benson and Ting-Hsuan Chen contributed equally to this work.

* Correspondence: Ting-Hsuan.chen@plantandfood.co.nz; ou.195@osu.edu

# Abstract

Genome size varies by several magnitudes among eukaryotes. Expansion and contraction are

primarily driven by transposable element activity, but the underlying processes remain enigmatic

due to a lack of historical records tracing these changes. Here, we present the TE Evolution

Simulation Suite, or TESS, for versatile simulation of whole-genome sequences with varying

transposon dynamics. We analyzed the pangenome of *Pucciniomycotina*, a subdivision of fungi

containing the myrtle rust (*Austropuccinia psidii*) genome, which drastically expanded to 1018

Mb. Using TESS, we simulated digital replicas of *Pucciniomycotina* genomes and replicated the

genome size evolution of *A. psidii* driven by LTR retrotransposons.

Keywords: Transposable Element, Genome Size, Forward Simulation, InDels, Solo LTR,

Structural Variation, Speciation, Rust Fungus

# Background

Transposable elements (TEs), or "jumping genes", are dynamic DNA sequences that replicate
and relocate within genomes. TEs constitute 46% of the human (*Homo sapiens*) genome [1], 85%
of the maize (*Zea mays*) genome [2], and over 90% of the genome of the fungal pathogen
*Austropuccinia psidii* that causes the myrtle rust disease [3]. The prevalence of TEs across the
Tree of Life reflects their dual roles as both genomic parasites and drivers of evolutionary
innovation. Beyond inflating genome sizes, TEs drive structural variation, modulate gene
expression, and facilitate chromosomal rearrangements [4]. In both plant and mammalian
genomes, bursts of TE activity correlate with speciation events and adaptive radiations,
enabling rapid responses to environmental stressors like pathogens or climate shifts [5].
Recent studies highlight TE's contribution to the collective genetic repertoire of a
species. Core genomes, conserved across individuals, are often gene-rich and TE-poor [2], while
"variable" or "dispensable" regions are TE-dense and enriched for stress-response genes,
pathogen resistance loci, and structural variants [6]. In *Brassica oleracea*, resistance gene
analogs cluster in TE-rich variable regions [7]. Similarly, tandem clusters of leucine-rich repeat
(LRR) genes in maize are found in genomic regions rich with young long terminal repeat (LTR)
retrotransposons [8], suggesting that TE-driven structural plasticity contributes to adaptive traits.
Related lineages can become karyotypically disparate over generations due to differences in
their ability to repress TEs [9]. The global impact of TEs on macro-chromosomal structure plays
an important role in evolutionary fates by influencing meiotic stability and similarity between
cytotypes [10]. These findings underscore TEs as dynamic regulators of genome evolution,
influencing both microevolutionary adaptation and macroevolutionary diversification.
Genome size varies considerably across taxa. Mammalian and bird genomes appear
more tightly constrained in genome size than those of both fungi and plants [11]. Numerous
studies across the Mammalia class indicate this size constraint is not through a lack of TE
radiation bursts [12] but through mechanistic interplay with purifying selection, where bursts are
followed by TE silencing [13] and DNA loss [12]. Genome size variation in plants facilitates
adaptation to specific environments, primarily carried out by whole genome duplication (WGD)
and TE expansion [14]. Extremes of genome size can be observed from the compact genome of
Arabidopsis (~135 Mb) through to the giant genomes of the Chinese pine (*Pinus tabuliformis*) at
~25 Gb [15], lily (*Lilium sargentiae*) at ~36 Gb [16], and the New Caledonian fork fern (*Tmesipteris
oblanceolata*) at ~160 Gb [17]. These are examples of unconstrained genome expansion, with
extremely high TE content within intergenic regions and also within introns. Conversely, there
are plenty of signs that genome size expansion is not left unchecked in plants: Arabidopsis has
undergone several WGDs, yet remains compact partially through subsequent gene loss that
purges redundant paralogs [18]. The dynamic of genome size change has been difficult to
accurately evaluate primarily because of the lack of high-fidelity tools for simulation and
modeling studies.
Advances in long-read sequencing and computational methods have empowered efforts
in resolving TE dynamics [19], yet reconstructing their evolutionary history remains challenging
due to TEs often being fragmented, nested, and highly identical to each other. Simulations of TE
evolution are thus indispensable for testing hypotheses about genome diversification, evaluating
TE annotation tools, and modeling the interplay between TE activity and selection. By

generating synthetic genomes with known TE architectures, including nested insertions,
sequence divergence, and polymorphism, researchers can disentangle the stochastic and
deterministic forces shaping genome evolution [20].
Existing TE simulators vary widely in scope and range of biological scenarios that they
mimic. Early models, such as those by Clough et al. (1996), introduced foundational concepts
by simulating varying modes of mammalian transposition [21]. Modern tools address specific
niches. SimulaTE [22] generates complex TE landscapes through predefined TE architectures,
limiting its utility for exploring stochastic evolutionary scenarios. Population-genetic frameworks
like SLiM [23] excel at modeling selection and demography but compress TEs into individual units,
precluding modeling of TE sequence mutagenesis, nesting insertion, and the consequential
genome size evolution. TESD [24] incorporates spatial population structure and selection against
deleterious insertions, yet its inaccessibility hinders further application. replicaTE [25] engineers
synthetic genomes by deleting and reinserting TEs into real assemblies but lack dynamic
iterations through evolutionary epochs. Meanwhile, pMEsim [26] focuses on simulating
polymorphic TE variants for pan-genomes, sacrificing long-term evolutionary dynamics, and
simuG [27] serves as a generic genome simulator that can simulate TE insertions. GARLIC [28]
specializes in generating realistic artificial genomic sequences by modeling intergenic
sequences based on kmer profiles, GC contents and local repeat landscape, which serve as
high-quality negative controls for evaluating annotation tools.
While existing simulation tools collectively address diverse needs, few allow flexibility in
genome customization, TE sequence mutagenesis, and dynamic modulation of
insertion/deletion rates across evolutionary epochs; none provides granular family-specific
controls with iterative simulations for hypothesis testing; and most prioritize synthetic data
generation over empirical parameter inference.

# Results

## Introduction of the TE Evolution Simulation Suite (TESS)

Here, we present the TE Evolution Simulation Suite (TESS) that allows dynamic forward
simulation of TE insertions, deletions, truncations, and divergence and speciation events. TESS
is a simulation toolbox comprising three novel components, RandSeqInsert,
TEGenomeSimulator, and PrinTE (**Fig. 1; Table S1**). Each component is capable of executing
independently and interconnecting with the other, allowing for versatile simulation of complex
genome evolution scenarios.
1. RandSeqInsert specializes in the simulation of TIR DNA transposon insertions. It uses a
tree-based algorithm for efficient sequence insertions at random positions, allowing for
benchmarking of TE annotation and structural variant detection tools;
2. TEgenomeSimulator uses single time-point simulation of TE insertion to produce a
genomic evolutionary snapshot. It can model a breadth of TE mobilization scenarios with
deference to the timing and duration of TE burst, and accumulated mutation. These
features allow for flexible approximation of the TE composition of real genomes;

3.  PrinTE enables iterative forward simulations to explore the effects of TE insertion,
mutation, and deletion on genome size evolution, genomic divergence, and the impact of
selection on TE turnover.

**Fig. 1. Programmatic components and functionality of the TE Evolution Simulation Suite**
**(TESS).** TESS is composed of three modules: TEgenomeSimulator, RandSeqInsert, and
PrinTE, represented by distinct colour blocks. The high-level computational workflows of
TEgenomeSimulator and PrinTE are illustrated with rectangles and solid lines. The output from
RandSeqInsert can serve as input for TEgenomeSimulator and PrinTE. See the main text for
detailed descriptions of each module's functionality.

## *RandSeqInsert:* a light-weight simulator for TIR insertions and structural variants

RandSeqInsert was developed to simulate Terminal Inverted Repeat (TIR) transposable
element insertions in genomic sequences. This high-performance Python tool implements an

AVL (named after inventors Adelson-Velsky and Landis) tree-based algorithm for efficient
sequence insertions at random positions in target sequences, enabling precise control over
insertion frequency (**Fig. S1A**). With support for nested insertion, the tool accurately represents
how TIR transposons accumulate in genomes over time. RandSeqInsert can track all insertions,
generating FASTA, BED, and DOT graph files to support further analysis of simulated
transposition patterns (**Fig. S1**).
129            RandSeqInsert simulates target site duplications (TSDs) as biological insertion
signatures with adjustable length and mutation rates, which allow for more accurate detection by
structure-based *de novo* annotation tools (**Fig. S2**). We demonstrated the simulation of TIR
insertions using RandSeqInsert and the detection of these events using minigraph [29],
showcasing the simulation of biologically relevant structural variants and how this simulation
framework can be extended to model TE variation at the population or pan-genome levels (**Fig.
S1B**). Accurate simulation of TE insertions allows for the creation of ground truth sets for
benchmarking and method development purposes (**Fig. S1; Fig. S2**).

***TEgenomeSimulator:*** **simulating diverse real-world TE evolution**

TEgenomeSimulator is a versatile tool that builds on and extends functions from denovoTE-eval
[30], allowing users to define and test a range of TE mutagenesis scenarios that can be reflective
of an evolutionary time point, followed by random and nested insertion into a randomly
synthesized or user-provided genome (**Fig. S3**). It tracks TE locations, sequence identity,
integrity, and breakpoints from nested insertion, providing flexibility to explore individual TE
components and their simulated decay within synthetic genomes (**Table S2**). Key developments
include support for multi-chromosome genome synthesis, flexible TE mutagenesis parameters
incorporating informed TSD length adjustments, and the ability to analyze and reuse annotated
TE composition profiles. Additionally, it streamlines the synthesis of artificial genomes, enabling
efficient simulation of digital replicas of real genomes (**Table S2**).
148            Multiple simulation modes are implemented within TEgenomeSimulator (**Fig. 1**). The
149    Random Synthesized Genome Mode is used to generate a fully synthesized genome adhering
150    to user-defined parameters reflecting genome size and GC content. Users can then modify TE
151    mutation parameters such as TE sequence identity and integrity, as well as copy number.
152            The simulation of TEs within synthetic genomes is often substantially different from what
is observed in real genomes in terms of composition, length, and sequence diversity. To
address this, TEgenomeSimulator implemented the TE Composition Approximation Mode,
whereby TE composition of a provided genome can be used to build a model for inference of TE
mutagenesis parameters. This genome-informed approach can produce a simulated genome
whose TE content closely mirrors that of the original genome, in terms of copy number, genomic
occupancy, sequence divergence, and length ratios relative to consensus sequences. As a
result, it provides a synthetic snapshot of the genome's current TE evolutionary landscape,
closely resembling the observed state of the input genome.
161            Moreover, the Custom Genome Mode implemented in TEgenomeSimulator gives users
the flexibility to utilize a hybrid strategy, combining genome backbones with custom TE libraries.
For example, users can first generate a digital replica of a real genome using the TE
Composition Approximation Mode, then use this simulated genome as an input for the Custom
Genome Mode along with a TE library that contains only LTR retrotransposon families from a

different species. By adjusting the range of mean identity, SD, and TE copy numbers, users can
simulate a TE burst within the digital replica, modelling scenarios such as horizontal TE transfer.
The resulting genome can be used not only to benchmark computational methods but
also as a burn-in phase for forward-time simulations in genome evolution studies, as such
simulated genomes and their TE sequences can be used very effectively in combination with
PrinTE.

## *PrinTE:* forward simulations of TE insertions, deletions, and genome divergence

PrinTE is a forward-time genome simulation tool designed to model the dynamics of TEs within
eukaryotic genomes across evolutionary time and was developed to incorporate as many
biologically relevant parameters into the model as possible (**Fig. S4**). It uses a modular
approach similar to TEgenomeSimulator, beginning with a user-provided (Custom Genome
Mode) or synthetic genome (Random Synthesized Genome Mode) that represents a present-
day or theoretical ancestral genome. Users may generate these inputs from prior PrinTE runs,
import them from TEgenomeSimulator or RandSeqInsert, or construct them entirely from
scratch with PrinTE's built-in parameterization tools.
PrinTE implements two methods for simulation. The variable-rate method aims to
incorporate consequences from some of the biological and selective forces that guide TE
evolution. We established several rules in this method: 1) The host genome survival and TE
proliferation are conflicting forces pitted against each other; 2) TE superfamilies compete with
each other since they occupy the same limited space; 3) The rate of TE insertion is positively
correlated with the number of intact TEs in the previous iteration; 4) TE deletions are guided by
the number of TEs that are deleterious to the host's survival. For this purpose, the simulated
genome consists of three components, all of which are receptive to new TE insertions: existing
TEs, genic space, and intergenic space. If a TE inserts into another TE or intergenic space,
there is no cost to the host and no impact on the TE deletion rate. In contrast, TE insertions into
genic space are costly to the host, resulting in elevated TE deletion rates in the next generation.
Thus, the outcome of the variable-rate method is less predictable, which may lead to
generations of genome size increase followed by subsequent generations of decrease. It is
useful for those who wish to explore the regulation of TEs through time and to understand the
forces that affect TEs' bursts and purges.
The fixed-rate method is designed to generate more controlled and predictable outputs.
In this method, per-generation TE insertion and deletion rates are predefined by the user,
allowing the modeling of known evolutionary trajectories and the generation of results that more
closely match expectations (**Fig. S4**). In practice, we find it most helpful to either start from a
simulated genome that mirrors a present-day version and forecast a few million generations of
TE evolution. Alternatively, it can start from a genome as we expect it may have looked a few
million generations ago and simulate it to the present day. This approach allows users to test
hypotheses relating to TE bloat and purge and to potentially superimpose TE evolutionary
narratives onto the species of interest.
PrinTE can simulate as briefly as a single generation, which can generate subtle
changes to the simulated genome. To allow for simulating millions of generations, PrinTE
employs a stepwise approach in which it can simulate TE evolution of a number of generations
in steps until it reaches the total number of generations specified by the user. To approximate

TE mutation in multiple generations, PrinTE uses an exponential decrease distribution to mimic
DNA substitution of TE sequences over time, in which most TEs have few mutations.
In summary, TESS represents a state-of-the-art simulation of TE dynamics in eukaryotic
genomes, enabling the modeling of TE contributions to genome evolution and the estimation of
past and future genome evolution trajectories.

214
**Fig. 2. Modeling TE insertion and composition for various evolutionary scenarios using**
**TEgenomeSimulator.** (**A**) Impact of altering TE copy number ranges for each TE family on
simulated genome size. (**B**) Distribution of TE sequence divergence under variable ranges of
mean identity across all TE families. A fixed TE copy number range of 5-100 was used in these
simulations. (**C**) Four scenarios of historical TE activity that can be modelled by controlling the
range of mean identity that represents the timeline of TE activities and the standard deviation
(SD) of mean identity that represents the duration of TE activities. (**D**) Various distribution
patterns of TE sequence divergence in association with the settings of mean identity and SD in
**C**.

## Modeling the Time and Duration of TE Activity

The composition of TEs within a genome is shaped by a dynamic interplay of TE bursts, host-
mediated removal processes, latency of TE activity, accumulation of single-nucleotide variants

(SNVs), insertions, deletions, and varying degrees of sequence truncation. The biological
insights from various TE profiles are often found associated with evolutionary milestones [31].
Accurately modeling TE composition under different amplification and duration scenarios is,
therefore, crucial for understanding genome evolution and implementing forward evolutionary
simulations.
The observed TE copy number in a genome represents the net result of TE amplification
and subsequent removal. A history of active TE mobilization, combined with inefficient removal
mechanisms, can lead to a substantial accumulation of TE sequences, often resulting in
genome expansion. TEgenomeSimulator encapsulates this concept through the modification of
TE copy number introduced into simulated genomes. Using TEgenomeSimulator's Custom
Genome Mode, in combination with the TE-depleted *Arabidopsis thaliana* (TAIR10) genome and
the curated *Arabidopsis* TE library (see Methods), we simulated a range of genome size
expansions from under 200 Mb to over 1 Gb. This was achieved by increasing the maximum
copy number per TE family from 10 to 2000 (**Fig. 2A**).
The observed TE divergence distribution is usually the result of multiple TE family bursts
at different times. To simulate this, we can vary the range of mean sequence identity in
TEgenomeSimulator. For simplicity, the minimum and maximum copy number of simulated TE
families was set to 5 and 100 respectively, and we applied a range of identity varying in
distribution sizes (10% - 30%) and sliding between 70% - 100% of mean sequence identity (**Fig.**
**2B**). As the identity range increases, the peak of the divergence distribution becomes
progressively flattened. This broad range of TE identity indicates continuous TE bursts and can
lead to a more gradual and flattened divergence profile.
The duration of TE bursts, sometimes an indicator of the lifespan of a TE family, which
can be measured by the standard deviation (SD) of sequence identity, also influences
divergence distribution. A short-lived TE burst results in a rapid deposition of nearly identical
insertions. These insertions then gradually diverge owing to accumulated mutations. In contrast,
a long-lasting burst over evolutionary time leads to the continuous deposition of similar TE
sequences, causing a less synchronized divergence pattern owing to overlapping degradation
timelines. Moreover, ancient TE bursts tend to show greater sequence deterioration due to
prolonged exposure to mutational decay. We can simulate genomes that represent different
evolutionary phases of TE activity (**Fig. 2C**), offering genomic canvases with distinct base-level
TE composition that can serve as different burn-in phases for more advanced simulations. As
shown in **Fig. 2C**, the representation of TE burst timeline and duration on the x-axis and y-axis,
respectively, allows four distinct TE divergence landscapes to emerge: I) high sequence identity
with high SD, modelling recent TE bursts with ongoing or prolonged activity over time; II) low
sequence identity with high SD, modeling ancient and prolonged activity, or multiple
independent bursts occurring over an extended period; III) low sequence identity with low SD,
reflecting ancient TE bursts that were quickly inactivated, with TE activity remaining silenced
since the initial event; IV) high sequence identity with low SD, inferring very recent and rapid TE
bursts, resulting in a synchronized insertion of highly similar sequences.
We modeled these four scenarios using the Custom Genome mode of
TEgenomeSimulator. The identity ranges and SD ranges were implemented as shown in **Fig.**
**2C**, with identity ranges of 90% to 100% and 70% to 80%, and SD ranges of 15 to 20 and 1 to
5. The TE copy number range was fixed between 5 and 1000. The resulting sequence

divergence profiles demonstrated a narrowing of the peak as the SD range decreased from 15–
20 to 1–5, indicating more instantaneous bursts (**Fig. 2D**). Furthermore, shifting the identity
range from 70–80% to 90–100% led to a leftward shift of the divergence peak (from ~0.25 to
~0.1) and an increase in peak height. These results demonstrate that TEgenomeSimulator is
well-suited for modeling TE composition under diverse evolutionary scenarios, which can be
considered the base-level scenarios before introducing additional evolutionary TE activity, by
supplying the simulated genome as a new input to TEgenomeSimulator or PrinTE.

## Genome Size Evolution Through TE Expansion and Contraction

LTR retrotransposons (LTR-RTs) are the primary contributing TEs to genome size in many
plant, fungi, and animal genomes [32]. To isolate the effect of TEs on genome size across
temporal gradients, we focused on modeling LTR-RT evolution using PrinTE's fixed-rate method
to model predictable patterns of TE expansion and contraction. Here, we simulated two dynamic
scenarios subsequently on the same genome. The first scenario used a TE deletion rate
fourfold higher than that of insertions. We observed a 45% decrease in genome size after five
million generations of forward simulation (**Fig. 3A**). The decrease was initially driven by a swift
reduction in the number of intact TEs (**Fig. S5A**). The number of fragmented TEs increased as
new TE insertions created nesting and fragmentation of TEs in subsequent generations. The
number of intact TEs became limited through generations of purging, and because of the
constant deletion rate, we observed an elevated purging of fragmented TEs in later generations
(**Fig. S5B**).
Following genome size decrease, we continued the simulation but reversed the TE
insertion rate to fourfold higher than that of deletions. The elevated insertion rate caused rapid
genome bloating (**Fig. 3A**) powered by an increase in the numbers of both intact and
fragmented TEs (**Fig. S5C,D**). The sequence lengths of solo, fragmented, and intact LTR-RTs
shifted in response to purging pressures from the host (**Fig. S6**). During periods of genome
expansion, TEs (fragmented, intact, or solo) tend to be longer in length.
Because the two LTRs of an intact LTR-RT were identical at transposition, DNA
mutations in this region are used to estimate the age of the LTR-RT [33,34]. During simulations of
genome size decrease, when the rate of TE deletions exceeds the rate of insertion, we
observed unbiased removal of LTR-RTs through generations. Reduced LTR-RT insertions and
accumulation of DNA substitutions led to a shift in the distribution of LTR-RT ages across time
(**Fig. 3B**). There were two distinct density peaks as simulation proceeds, even though the rate
of TE insertions and deletions remains constant (**Fig. 3B**). The smaller peak represents younger
LTR-RTs that were inserted recently and have had less time to be purged, and the larger peak
that shifts to the right and decreases in density through time was retained from the standing
body of LTR-RTs in the ancestral genome. In cases where the rate of TE insertions exceeds
deletions, young LTR-RTs quickly proliferate in high abundance and dominate the density
distribution of LTR-RTs in the simulated genome, masking the signal of remaining older LTR-
RTs (**Fig. 3C**).
When species become reproductively isolated from each other, their TEs are also
isolated and evolve independently. The two genomes share TEs that derive from their common
ancestor and also contain novel insertions that are unique to the other. With PrinTE, we
simulated a speciation event by evolving a genome for a period of time, then diverted the

genome into two isolated evolutionary paths with shifted TE dynamics to create divergent
evolutionary trajectories of genome expansion or contraction (**Fig. 3D,E**). In the contracting
genome, there are relatively few unique TEs created after speciation (**Fig. 3D; Fig. S7**). In
contrast, the expanding genome had a higher abundance of young and unique TEs (**Fig. 3E;**
**Fig. S7**). In the fixed-rate method, the rate of TE insertion and deletion remains relatively stable,
only deviating by the stochastic variability introduced by sampling a Poisson distribution, while
the number of insertions and deletions introduced scales to the size of the genome (**Fig. 3F**).
Deletion of TEs is mainly achieved by unequal or illegitimate recombination, which both
operate through sequence homology between TE sequences. In PrinTE, we allow users to set
TE deletion bias exponentially to the TE length using the $-k$ parameter. Larger $k$ has a stronger
bias in deleting longer TEs and thus has more constraints on genome size (**Fig. S8**). When $k$ is
set to 0, TE deletion has no bias to TE length, allowing genome size growth through the
accumulation of long TEs (**Fig. S8**). Illegitimate recombination between LTR regions of
individual LTR retrotransposons will partially remove the element and create solo LTRs as
remnants. Given a fixed deletion rate, PrinTE can model the proportion of LTR deletions that
goes through this process by implementing the "$--solo-rate$" parameter. Increased solo rate
can increase the number of solo LTRs in the genome and slow down the purging of LTR
retrotransposons, resulting in a larger genome size (**Fig. S9**, **Fig. S10**). By adjusting parameters
of deletion length bias and LTR solo rate that interplay with TE insertion and deletion rates, we
can create complex scenarios for modeling genome size evolution in different systems.
Recognizing that the rates of TE insertion and deletion can be dynamic through time and
may be modified by varying biological processes, we implemented the variable-rate method in
PrinTE, which can be used to explore specific molecular features that constrain TE evolution
and its impact on genome size. Here, rather than setting fixed TE insertion and deletion rates,
we start with an initial rate and allow the tool to dynamically determine subsequent insertion and
deletion rates based on the genetic makeup of the most recently simulated genome (**Fig. 3J**).
We simulated the genome size and TE evolution of the present-day *Arabidopsis thaliana* and
introduced theoretical parameter conditions to forecast one million generations of TE evolution.
With the tested conditions, there is an initial period of genome bloating followed by a longer
period of TE purging and genome size contraction (**Fig. 3G**). Expectedly, genome size
increased through the accumulation of both intact and fragmented TEs and decreased due to
the reduction of intact TEs (**Fig. 3H**). The divergence of intact LTR-RTs reflects the genome
size expansion and contraction (**Fig. 3I**). These features allow us to simulate TE dynamics close
to what is observed in eukaryotic genomes.

**Fig. 3. PrinTE models TE contribution to genome size evolution using fixed (A-F) and variable (G-J) rate methods.** (**A-F**) Ten million generations of LTR-RT evolution explores their long-term impact on genome size using the fixed rate method in PrinTE. For demonstration, we set the first five million generations of evolution using a rate of TE deletion that exceeds insertion by fourfold. Over the second five million generations, the rates were reversed, and insertion exceeded deletion by fourfold. (**A**) A syntenic ribbon plot illustrates the change in chromosome size. Pie charts corresponding to each time point show the relative contribution of each feature to genome size. (**B,C**) Density distributions of LTR-RT divergence estimated using the Kimura two-parameter (K2P) model during the first five million generations (**B;** genome size decrease) and the second five million generations (**C;** genome size increase). Five million generations is set as a transition time point between genome size decrease and increase and is shown in both plots as dashed red lines. Phylogenetic trees show the evolutionary relationship of LTR-RTs from the *Tekay* family after four million generations of genome size decrease (**D**) and four million geenrations of genome size increase (**E**). (**F**) LTR-RT insertion and deletion counts (solid line) and rates (dashed line) across 10 million generations. (**G,H**) A simulated genome replica of *Arabidopsis thaliana* is forward-evolved one million generations using

366    PrinTE's variable rates method. (**H**) Intact (top) and fragmented (bottom) TE counts per
367    superfamily across one million generations. (**I**) The distribution of LTR-RT divergence estimated
368    using the K2P model. (**J**) TE insertion and deletion counts (solid line) and rates (dashed line)
369    illustrate the self-adjusting functionality of PrinTE's variable rate method.

## Simulating Digital Replicas of Eukaryotic Genomes

TE composition in real genomes varies substantially across TE families and species because of
their complex and diverse evolutionary histories. This variability makes it challenging to simulate
realistic genome-wide TE landscapes. To address this, we used TEgenomeSimulator's TE
Composition Approximation Mode to model the TE profiles of *A. thaliana* (TAIR10) [35] and the
myrtle rust pathogen, *A. psidii* [3]. The sizes of the simulated genomes, 118.2 Mb for *A. thaliana*
and 0.98 Gb for *A. psidii*, closely matched the actual genome sizes (**Fig. 4A**). Similarly, the
proportion of the genome occupied by TEs closely approximated the observed values in both
organisms (**Fig. 4B**). In terms of TE loci and TE bases per superfamily, the simulated TE
composition also aligned well with the real data (**Fig. S11**).
We estimated the sequence divergence distribution of each TE family in real genomes
for simulations of digital replicas. While the simulated divergence profiles resembled the real
data, they were not identical: the fine-scale fluctuations present in real genomes were smoothed
because of the use of normal distributions in the simulator. This approach provides a
generalized distribution pattern that captures the major trends in TE sequence evolution (**Fig.
4C**).
TE fragmentation patterns also vary widely across TE families and genomes. The
Random Synthesized Genome and Custom Genome modes of TEgenomeSimulator use a beta
distribution to model TE integrity across all TE families. When alpha and beta parameters are
set between 0 and 1, with alpha equal to or smaller than beta (**Fig. S12A**), the distribution takes
on a U-shape that approximates general fragmentation patterns previously observed in
grapevine data [36]. However, because of the unique TE evolutionary histories of different
genomes, this approach may fit well for one species but can overestimate the proportion of TE
loci with intermediate degrees of fragmentation in others (**Fig. S12B**). Moreover, fragmentation
patterns can differ substantially between TE families, in particular those identified by homology-
based methods, which often lack members with high structural integrity (**Fig. S12C**). To improve
accuracy, the TE Composition Approximation Mode directly extracts the empirical distribution of
TE length ratios (relative to consensus sequences) for each TE family. This enables the
simulator to reproduce the integrity distributions more authentically in the resulting simulated
genomes (**Fig. 4D**).
These digital replicas of real genomes provide a robust foundation for simulating future
TE-related events, such as new TE bloating or purging episodes, effectively serving as burn-in
phases for modelling genome evolution from the current state.

**Fig. 4**. **TE Composition Approximation for the *Arabidopsis thaliana* and *Austropuccinia psidii* genomes.** (**A**) Size of the original and simulated genomes. (**B**) The proportion of original and simulated genomes occupied by TE and non-TE bases. (**C**) TE sequence diversity distribution colored by TE superfamily. (**D**) The distribution of TE sequence integrity, which is the length ratio relative to the consensus sequence, is colored by TE superfamily as the same in (**C**).

## Genome Size Evolution of Rust Fungus

We further studied the impact of TE dynamics on genome size evolution using rust fungal genomes for their wide range of genome sizes. We downloaded 58 *Pucciniomycotina* fungal

genomes from NCBI and JGI MycoCosm [37] to investigate the expansion and contraction of genome size (**Table S3**). *Puccniomycotina* is a subdivision of fungus that contains 10 classes and over 8400 species [38]. We identified and removed alternative haplotypes from these genomes using Redundans [39]. The resulting haploid genome size varies from 13.6 Mb (*Mixia osmundae*) to 1018.1 Mb (*Austropuccinia psidii*), representing over 70-fold differences (**Table S4**). We performed Benchmarking Universal Single-Copy Orthologs (BUSCOs) evaluations using compleasm [40] and the odb12 basidiomycota database. The complete BUSCOs (single and duplicated) averaged 90.7% (SD = 5.4) and contig N50 averaged 846 kbp (ranging from 111 kbp to 3.8 Mb), suggesting high assembly quality of haploid genomes (**Table S4**).

**Fig. 5.** (**A**) The phylogeny of *Puccniomycotina* fungal genomes used in this study. Protein sequences of shared single-copy BUSCO were used for tree inference with 1000 bootstrap replications. Bootstrap support percentage is shown on each branch. The two classes, *Puccniomycotina* and *Microbotryomycetes,* are labeled beside the phylogeny. The genome size of each species is indicated by color dots. (**B**) Principal Component Analysis (PCA) on LTR metrics in fungus genomes. Each dot represents one genome. PCA loadings of each metric are shown. Genomes selected for further studies are labeled. (**C**) Pearson's correlation (*r*) between genome size and LTR sequence content (LTR_percent), intact LTR-RT count (Intact.count), solo LTR count (Solo.count), and solo:intact ratio. Each dot represents one genome. The correlation *r* and p-value are shown on the panel title. (**D**) Divergence of intact LTR retrotransposons in *Cronartium ribicola* (white pine blister rust), *Puccinia coronata* f. sp. *avenae* (oat crown rust), and *Austropuccinia psidii* (myrtle rust) genomes. The top five most abundant LTR families in each genome are colored, and they are not shared between genomes. A density line is shown as the contour in each plot. The locations of these species on the phylogeny are indicated by dotted lines.

We identified the shared single-copy BUSCOs and used iqtree [41] to construct the
phylogeny of *Pucciniomycotina*. The resulting phylogeny was over-represented by two classes,
*Microbotryomycetes* and *Pucciniomycetes* (**Fig. 5A**). *Microbotryomycetes* genomes of
*Microbotryomycetes* are small, ranging from 17.5 Mb to 50.3 Mb. Correspondingly, on average,
only 0.4% of the genome was represented by LTR sequences with a range of 0% to 3.0%
(**Table S4**). This observation suggests that the best strategy for controlling genome size is to
avoid the invasion of LTR retrotransposons. *Pucciniomycetes* have widely distributed genome
sizes, ranging from 26.8 Mb to 1018.1 Mb, with an LTR sequence content corresponding to
0.4% to 67.4% (**Table S4**).
To further study the dynamic of LTR retrotransposons in these fungal genomes, we *de*
*novo* identified intact LTR elements and solo LTR sequences, and calculated whole-genome
solo:intact ratio using LTR_retriever [33]. By using genomes with LTR sequence content larger
than 1% (n = 22), we conducted Principal Component Analysis (PCA) on LTR metrics and
genome size, showing clustering of smaller genomes and larger genomes as outliers, including
*Cronartium ribicola* (white pine blister rust), *Puccinia coronata* f. sp. *avenae* (oat crown rust),
and *Austropuccinia psidii* (myrtle rust; **Fig. 5B**). PCA loadings suggest that solo LTR number,
intact LTR number, and their ratio were all drivers to the outliers of large genomes (**Fig. 5B**).
Including total LTR content, these LTR metrics showed strong positive correlations with genome

size (Pearson's r ≥ 0.65, p ≤ 0.001; **Fig. 5C**).

Not surprisingly, intact LTR elements as the product of LTR proliferation were strongly
correlated with genome size (r = 0.76; **Fig. 5C**). However, solo LTR sequences as the product
of LTR removal [33] showed an even stronger correlation, almost linearly correlated with genome
size (r = 0.98; **Fig. 5C**). Another counter-intuitive observation is the positive correlation between
genome size and solo:intact LTR ratio, which is a metric used to measure the intensity of LTR
purging [34]. In the traditional understanding, the higher the solo:intact LTR ratio, the more intense
purging of LTR sequences, and thus the smaller the genome [42]. The positive correlation
between solo:intact ratio and genome size suggests that illegitimate recombination may not be
an effective way of escaping genome obesity.
We selected *C. ribicola*, *P. coronata* f. sp. *avenae*, and *A. psidii* as representatives of
three distinct roadmaps to genome size evolution. *C. ribicola* has a genome size of 211.6 Mb
with the LTR divergence distribution showing an extremely quick turnaround time for LTR
element insertions (**Fig. 5D; Fig. S13**). The solo:intact ratio of *C. ribicola* is 27.4 (**Table S4**), the
highest among all genomes, suggesting a quick insertion–quick removal mode of action, and a
relatively stable or slowly increasing genome size. *P. coronata* f. sp. *avenae* has a genome size
of 113.2 Mb and a solo:intact ratio of 1.7 (**Table S4**). The LTR divergence distribution showed a
much slower turnaround time for LTR insertions (**Fig. 5D; Fig. S13**), suggesting a slow
insertion–slow removal mode of action, and a relatively stable genome size. *A. psidii* has an
extremely large genome size of 1018.1 Mb and a solo:intact ratio of 23.5, the second highest
ratio among our data (**Table S4**). However, the LTR divergence of *A. psidii* shows the peak of
LTR insertion passed the current time (**Fig. 5D; Fig. S13**), suggesting a slowdown of insertion
activities [8] and a decreasing genome size.
480
481

**Fig. 6. Simulating genome size evolution in three *Pucciniomycetes* genomes.** (**A-C**)
*Cronartium ribicola* (white pine blister rust), (**D-F**) *Puccinia coronata* f. sp. *avenae* (oat crown
rust), and (**G-I**) *Austropuccinia psidii* (myrtle rust). (**A, D, G**) Sequence divergence compares the
empirical divergence distribution of LTR-RTs (Reference) to simulated genomes. LTR
divergence was estimated using LTR_retriever. Three timepoints were plotted for *C. ribicola* (**A**)
and *P. coronata* f. sp. *avenae* (**D**) to showcase the relatively stable TE dynamic. The three
timepoints of *A. psidii* (**G**) highlight the rapid purging of LTR-RTs during the period of genome
size contraction. (**B, E, H**) Genome size changes across generations. (**C, F, I**) The empirical
solo:intact ratio estimated using LTR_retriever. (**J**) The estimated LTR insertion and deletion
rates of fungal genomes. The expansion and contraction phase of the *A. psidii* simulation was

493     plotted separately. (**K**) Microsynteny between the simulated *P. coronata* f. sp. *avenae* before
494     (top) and after (bottom) the TE evolution that led to genome size increase in the simulated *A.*
495     *psidii* genome. Collinear genes are blue, and LTR-RTs are orange.

## Accessing Genome Evolution with Forward Simulations of LTR Activities

Here, we demonstrate the utility of TESS in modeling genome size evolution under the impact of
TE bloat and purge. Despite their shared ancestry (**Fig. 5A**), the three selected fungal species
(*C. ribicola*, *P. coronata* f. sp. *avenae*, and *A. psidii* ) show immense variation in genome size
and LTR-RT evolution. To test our hypothesis that *C. ribicola* has a stable genome size that it
achieved by rapid but balanced LTR-RT insertion and deletion, we used TEgenomeSimulator to
create a digital replica of the *C. ribicola* genome and forward evolved it using PrinTE with a
variety of insertion and deletion rates in search of the observed LTR-RT divergence distribution.
At each trial, we compared the LTR-RT age distribution with that of the real genome (**Fig. S14)**.
We found that an LTR-RT insertion rate of $2\times10^{-9}$ per bp per generation and a deletion rate of
$2.4\times10^{-9}$ per bp per generation provided a simulated genome that closely matched the *C.*
*ribicola* genome with the genome size increasing moderately through time (**Fig. 6A,J**). Forward-
projecting 200,000 generations of TE evolution indicated that the density distribution of LTR-RT
ages in the simulation closely matched the distribution of the real genome (**Fig. 6A**, **Fig. S14**).
The solo:intact ratio of the simulated genome was 13 (**Fig. 6C**) in comparison to 27.4 in the real
genome (**Table S4**). Although the forward-simulated genome does not share an identical LTR-
RT footprint to that of *C. ribicola*, its proximity to the real data suggests that PrinTE can
effectively infer TE dynamics through forward simulations. Simulation of the *C. ribicola* genome
suggests that the species might have inserted and deleted (either intact or fragmented) one
effective LTR-RT every two generations during the latest 200,000 generations.
We repeated the same process using a digital replica of *P. coronata* f. sp. *avenae*a
simulated by TEgenomeSimulator to test our hypothesis that the species has slow TE insertion
and slow removal, leading to a relatively stable genome size. An LTR-RT insertion rate of $8\times10^{-12}$
per bp per generation and a deletion rate of $1.10\times10^{-11}$ per bp per generation were found to be
optimally adapted for maintaining the LTR-RT characteristics of the real genome for over 20
million generations (**Fig. 6B,J; Fig. S15**). Dating LTR-RTs in forward-simulated and the real
genome showed that the distributions were nearly identical (**Fig. 6D**). The solo:intact ratio in the
forward-simulated genome mimics that of the real genome, at 2.0 and 1.7, respectively (**Fig. 6F**,
**Table S4**). With these rates, we see a gradual increase in average genome size, adding
approximately 1.25 Mb per million generations (**Fig. 6E**). These results suggest that, in contrast
to *C. ribicola, P. coronata* f. sp. *avenae*a appears to have a mostly inactive LTR-RT footprint
with an average of 1090 generations between each effective LTR-RT insertion and an effective
deletion every 790 generations.
For our final simulation, we began with a relatively small genome that is typical in the
*Pucciniomycetes,* and simulated the TE evolution that led to the enormous genome bloating of
*A. psidii*. Based on the genome size and LTR-RT divergence distribution (**Fig. 5A,D**), we
expected that TE insertions exceeded deletions for a long period of time, leading to genome
size increase, followed by a shorter period of rapid LTR-RT deletions. Through evaluation of
various dynamic parameters, we found that an insertion rate of $1.4\times10^{-11}$ per bp per generation
and a deletion rate of $5\times10^{-13}$ per bp per generation for 24 million generations was optimally

tuned to mimic the observed LTR-RT age distribution of the *A. psidii* genome during the long
period of genome size increase (**Fig. 6G,J**; **Fig. S16A,B**). During this period, we aimed to
achieve a genome size that was marginally larger than the real genome so that we could tune
up the rate of deletion to reflect the more recent decline in young LTR-RTs observed in *A. psidii*
(**Fig. 5D**). At peak LTR-RT invasion, the simulated genome was 1,178 Mb in size (**Fig. 6H**).
The drop of recent LTR-RT insertions in *A. psidii* suggests a reduced insertion rate and
an increased deletion rate. Using a substitution rate of $2.0 \times 10^{-9}$ per bp per generation [43], we
estimated that the insertion reduction phase of *A. psidii* had lasted for 1.2 million generations.
We then forward-simulated an additional 1.2 million generations using elevated deletion rates.
Systematic evaluation of TE dynamic parameters revealed that an insertion rate of $1.4 \times 10^{-13}$ per
bp per generation and a deletion rate of $3 \times 10^{-11}$ per bp per generation can achieve a genome
size that is near *A. psidii*, at 1,027 Mb (**Fig. 6H,J**; **Fig. S16C**), and the LTR-RT divergence
distribution mimics the real genome (**Fig. 6G**; **Fig. S16D**). The solo:intact ratio of the simulated
*A. psidii* genome terminated at 6.8 (**Fig. 6I**), which is lower than the observed value at 23.5
(**Table S4**), suggesting other evolutionary processes may play roles in shaping the modern-day
*A. psidii* genome.
Simulation of *A. psidii* suggests two major phases that contributed to the present-day
genome size: a long period of genome size increase in which LTR-RT insertions exceeded
deletions by 28-fold and a short period where deletions exceeded insertions by 214-fold. Our
results support a recent contraction model for genome size evolution in *A. psidii* (**Fig. 6J**). It is
important to note that the estimated parameters are effective insertion and deletion rates
averaged from a long period of evolutionary time, which may not capture short periods of TE
expansion and contraction.
Because these three species were selected to represent the extremes of LTR-RT
dynamics in the sampled *Pucciniomycetes* genomes, we suggest that the amplification and
degradation of LTR-RT can vary widely within the fungal kingdom. In future studies, it will be
interesting to see if such a wide variety of LTR-RT dynamics exists at narrower phylogenetic
levels.

# Discussion

The TE composition profile of a genome serves as an evolutionary archive, preserving
signatures of TE dynamics and host genome responses that collectively shape genome
evolution. Deciphering these patterns from empirical data is challenging. Whether to test
evolutionary hypotheses or to predict future trajectories, simulation is often necessary. To
address this, we built TESS, a simulation framework composed of RandSeqInsert,
TEgenomeSimulator, and PrinTE. RandSeqInsert is a light-weighted simulator that excels in
insertions of TIR DNA transposons with the AVL tree architecture. TEgenomeSimulator and
PrinTE serve as the two central pillars of TESS, each serving distinct but complementary roles.
Used together, they significantly expand the capacity for simulating complex TE dynamics and
testing evolutionary hypotheses.

## TEgenomeSimulator enables flexible modeling of TE composition and evolutionary dynamics

TEgenomeSimulator is a versatile tool for simulating TE composition across diverse evolutionary trajectories, with strong potential for broader applications. It encapsulates the dynamic balance between TE insertions and host-mediated TE removal through its TE copy number parameter, which directly influences the final size of the simulated genome (**Fig. 2A**). It can also simulate genomes with different histories of TE activities by varying the ranges of mean sequence density and its standard deviation representing the duration of the TE burst, allowing flexible modelling of different evolutionary scenarios (**Fig. 2B–D**).

Simulating TE compositions to closely resemble real genomes is a rare capability among existing tools. TEgenomeSimulator stands out by enabling realistic modelling of TE content based on empirical patterns (**Fig. 4**). It integrates three major features of TE composition, a) TE family-specific copy number, b) sequence identity (modeled using normal distribution), and c) TE integrity profiles, into a coherent mutagenesis framework. These features allow the replication of the evolutionary patterns observed in real genomes with additional critical information, enabling a more precise manipulation of TE compositions for hypothesis testing.

In comparison, GARLIC [28] is a genome-wide simulation program prioritized for benchmarking purposes. It begins by extracting coding and non-coding genes and repeats from a user-provided genome, then profiles repeat diversity and analyzes the intergenic space using k-mer and GC content profiles to model genomic context. To simulate TEs, GARLIC first mutates the consensus repeat sequences, then inserts them into a non-repetitive genomic background.

While GARLIC provides more controls in simulating genes and intergenic architecture, TEgenomeSimulator remains more TE-centric, offering flexible, user-driven control of TE-specific parameters without extensive genome preprocessing. This streamlined design lowers the barrier for simulating TE dynamics under various evolutionary assumptions. As such, GARLIC is well-suited for studies focused on genes and their annotations, whereas TEgenomeSimulator is ideal for hypothesis-driven investigations into TE-specific evolutionary scenarios with customizable control over TE dynamic parameters.

## PrinTE allows forward simulation on TE dynamics

Transposable elements are a major genomic component and subject to all evolutionary forces, including purifying selection, drift, and mutation. The ability to trace TEs' responses to these events will greatly facilitate the studies of TE dynamics. Forward simulation is a key approach that allows the artificial evolution of TEs and other genomic features, which provides opportunities to sample the intermediate state of the evolutionary process.

SLiM is a powerful forward simulation program that can simulate the majority of evolutionary scenarios [23]. However, its operation on single nucleotide variants limits the application on TEs with varying lengths (**Table S2**). SLiM provides a work-around, by first compressing the whole TE locus into a presence and absence unit and performing simulations on these units. Sequence content is then filled back to replace the units to generate genomic sequences. This approach cannot model the variation accumulated through generations on the simulated TE sequences, including the deletion and formation of solo LTRs, which are critical

for genome size studies. On the contrary, PrinTE operates on whole-genome sequences,
including invariant sites that allow intuitive modeling of complex scenarios and interpretation of
results. In particular, PrinTE models the insertion and deletion rates of TEs and generates solo
LTRs, which empowers the study of genome size evolution driven by LTR bursts and purges.
The input files of PrinTE are highly flexible (**Fig. 1**). It has a built-in engine to simulate
highly customizable starting genomes, in which users can specify genome size, TE content, TE
superfamily abundance, and TE divergence in their simulations. PrinTE integrates CDS
sequences in the simulation, allowing for modeling purifying selections, tracing large structural
variants, and studying the evolution of gene space. PrinTE also takes the simulated genomes
from TEgenomeSimulator, allowing forward simulation from different diverse backgrounds of
genomes. Most importantly, PrinTE can also take its own output as input, allowing for simulation
of varying TE dynamics and studying the host-TE arms race. This feature also enables the
simulation of genomic divergence and speciation driven by differences in TE activities.

## 630 TE contributes to genome size expansion

The contribution of TE contents to genome size expansion has been confirmed across many
taxa and in our data (**Fig. 5C**). Over time, genomic data have come faster and of a higher
quality and yet, as noted by Osmanski *et al.* (2023) [44], the role of TEs in genome architecture
and evolution is often an afterthought. More in-depth analyses become reliant on domain
specialists for a meaningful evolutionary perspective.
In order to put forward mechanistic hypotheses, authors have developed specialized
computational methods to detect signals of TE radiation and impacts on speciation and genome
size. For example, Ricci *et al.* (2018) [13] developed a measure of TE Density of Insertion and
Relative Rate of Speciation to model TE radiation bursts and silencing. Kapusta *et al.* (2017) [12]
estimated TE-driven DNA gains and losses and proposed the 'accordion' model of TE radiation
followed by DNA loss for tight genome size maintenance in mammals and birds. Osmanski *et al.*
(2023) [44] applied a Redundancy Analysis to study TE composition across mammalian taxa
generated through the Zoonomia genomes [45], and found evidence of multiple recent TE
expansions driving larger genome sizes and wide-spreading TE-silencing events across the
mammalian tree. These authors hypothesize that variations in TE defense mechanisms may
have played a role in whether specific lineages were overcome by a TE burst at a given
evolutionary time.
However, we have limited knowledge about the maintenance of compact genomes
despite evidence of TE radiation bursts and the maintenance and regulation of extremely large
genomes. The Chinese pine attributes 69.4% of its 25.4 Gb genome to TEs, 86.5% of which are
made up of LTR retrotransposons [15]. The Chinese pine genome showed no sign of recent
whole-genome duplication, but 91.2% of its genes were undergoing discrete duplication or
family expansion, strongly associated with TEs. Our results revealed two different genome
dynamics in the *Pucciniomycotina* subdivision of fungi, including the *Microbotryomycetes* class
that harbours compact genome size (18-50 Mb) and the *Pucciniomycetes* class with drastic
differences in genome size (27-1018 Mb) (**Fig. 5A**; **Table S4**). The *Microbotryomycetes* class is
almost devoid of LTR sequences (**Table S4**), suggesting that the most effective way of
maintaining compact genome sizes is to be immune to the invasion of LTR retrotransposons.
Genomes in the *Pucciniomycetes* class have varying richness of LTR retrotransposons (**Table**

**S4**), and the PCA reveals several types of LTR dynamics (**Fig. 5B**) that are likely major drivers
of genome size variation.
The myrtle rust fungus expanded its genome size from an approximate ancestral size of
110 Mb to the current 1,018 Mb, experiencing a ~10 times increase. Our simulation studies
revealed that the genome is experiencing a downsizing effect, with LTR deletions faster than
insertions (**Fig. 6**). Using PrinTE's built-in genome simulation utility, we were able to simulate a
genome that mimic the TE landscape of the ancestral genome of myrtle rust, and further used
PrinTE to simulate two phases of evolution, including a long expansion period and a short
contraction period. This model fits well with the observed composition of the myrtle rust
genome, suggesting a close approximation of its trajectory of genome size evolution.
We recognize that TE activity is tightly linked to host genome regulation, particularly
when it impacts gene function and organismal fitness. It is also important to account for more
complex patterns in sequence identity distributions, which may go beyond simple normal
distributions. In our study, TEgenomeSimulator helped to shape TE composition and sequence
divergence under various evolutionary scenarios, while PrinTE simulated the real-time
mobilization and insertion behavior of TEs. Crucially, the combined use of these two tools
greatly enhanced the simulation capacity, enabling richer and more flexible hypothesis testing
across both evolutionary and mechanistic contexts. Future development would benefit from
incorporating features such as customizable identity distributions, more functional genomic
context, epigenetic regulation mechanisms, and insertion site preferences.

# Conclusions

In conclusion, the TE Evolution Simulation Suite (TESS) is a versatile collection of tools that
allows flexible forward simulation of whole-genome sequences with fine-grain user control of
genome size, TE content, superfamily abundance, TE divergence profile, TE insertion, TE
deletion, TE mutation, solo LTR formation, genome divergence, and speciation. We
demonstrated the use of TESS in resolving the genome size evolution of *Pucciniomycotina*
genomes. In particular, we simulated the genome size evolution of the myrtle rust
(*Austropuccinia psidii*) and other large fungal genomes. Our work transforms the study of
genome size evolution into a highly traceable and explorative simulation framework, and allows
testing of data-driven hypotheses.

# **Methods**

### **Implementation of RandSeqInsert**

RandSeqInsert employs an effective method for randomly inserting genomic fragments into
sequences (either user-provided or randomly generated). The implementation is built around a
balanced binary tree data structure that enables efficient sequence manipulation while
maintaining performance even for long sequences. Curated rice and maize TIR transposon
libraries are included with the tool to serve as donor sequences during the simulation.
***Balanced Sequence Tree***

The backbone of RandSeqInsert is the "SequenceNode" class, which implements an AVL tree
for sequence representation. An AVL tree is a balanced binary tree where the height difference
between left and right branches of any node never exceeds 1 [46]. When this balance is disrupted
during insertions, automatic tree rotations help to restore the balance.
The tree structure represents sequences as connected nodes that can be rearranged
without extensive copying. Each node stores the unique identifier (UID), the actual sequence
fragment (these can be non-TE backbones or inserted TEs), length data, total length of all
sequences in its subtree (essential for position mapping), height of the node, 5' and 3' TSD
sequences, and additional information. After each insertion operation, the balance factor is
recalculated at each node along the path from the insertion site to the root, and rotations are
performed automatically. Eventually, the complete sequence can be retrieved through an in-
order traversal of the tree, which is a depth-first search algorithm that visits nodes in a binary
tree following the Left-Root-Right order.
This structure offers major advantages over alternative approaches. A simple direct
sequence implementation would require rebuilding the entire structure after each operation,
resulting in $O(n)$ time complexity per operation and $O(n^2)$ complexity for $n$ operations. Using
regular binary tree would risk performance degradation to $O(n)$ per operation when insertions
occur in patterns (such as sorted order) that transform the tree structure into essentially a linked
list (which is a linear data structure where each node contains both data and a reference to the
next node), eliminating the logarithmic performance advantage. The AVL tree maintains $O($
$log\ n)$ per-operation time across all operation patterns through a self-balancing mechanism via
rotational operations [46].
**_Event Journal for Nested Insertions_**
RandSeqInsert implements a sophisticated event sourcing architecture through the
"NestingEventJournal" class to track complex nested insertion scenarios. This journal selectively
records only nested insertions (TE-to-TE insertions), captures critical information for each
nested insertion event, including the donor node UID, target node UID, left and right fragment
node UIDs, and TSD information. This selective recording strategy enables the reconstruction of
donor sequences that have been fragmented by subsequent insertions.
**_Insertion Algorithms_**
RandSeqInsert offers both recursive and iterative insertion algorithms, with the iterative method
working better for large sequences by avoiding recursion limits. The insertion process identifies
the appropriate position in the tree based on absolute sequence position, splits the target node
if insertion occurs within it, creates new nodes for the inserted donor sequence and split
fragments, and updates balance information while rebalancing the tree.
Position mapping is crucial to the algorithm. Given an absolute position in the full
sequence, the algorithm determines which node contains this position, the relative position
within that node, and how to split and reorganize nodes to incorporate the new sequence. When
insertion occurs within a node, that node is split into three nodes: a left node containing the
sequence before the insertion site, the current node repurposed for the inserted donor
sequence, and a right node containing the sequence after the insertion site. When a donor
sequence is inserted into another donor sequence (nested insertion), the system automatically
records this event, preserving the original target sequence information for future reconstruction.
**_Target Site Duplication (TSD) Modeling_**

The implementation includes a detailed modeling of Target Site Duplications, a biological
process where short sequences at insertion sites are duplicated during TE integration. The TSD
generation process takes the source TSD sequence at the insertion site, creates both 5' and 3'
TSD sequences with configurable lengths, and applies mutations to simulate biological variation
according to configurable probabilities. TSD sequences are integrated into the node structure
and are handled separately from the core donor sequence data.
Two types of mutations are supported: SNP mutations and InDel mutations. SNP
mutations replace a single nucleotide at a random position with a different randomly selected
base. InDel mutations represent a single random nucleotide insertion or deletion.
***Donor Sequence Tracking***
RandSeqInsert effectively tracks and records information about inserted donor sequences and
their positions. The system identifies donor nodes and calculates their exact positions in the
final sequence through a complete tree traversal. Following this process, it generates a FASTA
file that records each donor sequence with its precise start and end coordinates, and a BED file
that is compliant with PrinTE.
***Sequence Tree Visualization***
For research and analysis purposes, RandSeqInsert can generate two visualizations in
Graphviz DOT format [47]. The sequence tree visualization displays the hierarchical structure of
the AVL tree, with each node color-coded by type and annotated with position, length, and UID
information. The event graph visualization demonstrates the insertion event relationships,
showing the impacts of the nested insertions on sequence integrity. This dual visualization
approach enables researchers to understand both the structural organization of the sequence
data and the history of insertion events.

# Implementation of TEgenomeSimulator

TEgenomeSimulator was built on denovoTE-eval [30] and further developed to incorporate new
functionalities. It comprises three modes: Random Synthesized Genome Mode (`-m 0 or --`
`mode 0`), Custom Genome Mode (`-m 1 or --mode 1`), and TE Composition Approximation
mode (`-m 2 or --mode 2`). These modes all perform random TE insertion and nested TE
insertion sequentially. Depending on the mode chosen, the genome and TE sequences undergo
distinct preparations prior to TE insertion. The implementation and input requirements for each
mode are described in the following section.
***The Three Simulation Modes***
The Random Synthesized Genome mode will synthesize chromosome sequences using a set of
user-defined parameters, including chromosome length and GC content. A TE library can be
provided using the '`--repeat`' flag, which is used to simulate a population of TE sequences
through random mutation, including nucleotide substitutions, 1-base InDel, target site
duplication, and fragmentation. Simulated TE sequences are then randomly inserted into the
synthesized chromosome sequences.
The Custom Genome mode requires users to provide a genome in which TE sequences
have been removed. For convenience, this mode provides an option for users to generate such
a genome using the '`--to_mask`' flag. RepeatMasker [48] is used to mask and remove repeats
from the genome based on a TE library supplied by users. TE sequence mutation and random

784 nested insertions are introduced in the same way as the first mode. Users must also specify a
785 second TE library with '`--repeat2`', which is used for TE masking and removal. Note that the
786 simulator allows users to provide two different TE libraries for masking repeats and for TE
787 sequence mutation, respectively.

788   The TE Composition Approximation mode utilizes the '`--to_mask`' flag implemented in
789 the Custom Genome Mode, with the option of using two different TE libraries. By default,
790 RepeatMasker output is utilized in the configuration of downstream TE mutations. In brief, the
791 simulated TE composition is modeled on that of the original genome in terms of nucleotide
792 substitution rate, InDel rate, integrity, and the absolute and relative abundance of TE families.
793 The TE insertions and nested insertions in this mode are independent of the TE distribution in
794 the original genome.

795 ***Simulating TE Mutagenesis***
796 TE libraries provided by the user must adhere to the RepeatMasker recommended format
797 (">repeat_family_name#class/subclass"), e.g, ">ATCOPIA10#LTR/Copia". It is recommended
798 that the full sequence of each TE family be stored in the TE library file. For example, the LTR
799 and internal domain sequences of an LTR retrotransposon are stitched in the conjunction form,
800 LTR-INT-LTR, rather than stored separately. TEgenomeSimulator will check for split sequences
801 of LTR retrotransposons in the user-provided library and assemble full sequences for the user.

802   TEgenomeSimulator extracts classification information from the TE library for
803 incorporation into a final GFF file. Metadata, including TE family name, superfamily, subclass,
804 and sequence length, are also extracted from the TE library headers and compiled into the
805 output file 'TElib_sim_list.table'. This table also defines the mutagenesis parameters for each TE
806 family, including TE family copy number, mean sequence identity ($I$), standard deviation of
807 mean identity, and the proportion of sequence divergence ($D = 100 - I$) contributed by InDel ($i$).
808 The remaining proportion of divergence is attributed to nucleotide substitution ($s$), such that $i + s$
809 $= 100$. The table also includes values for target site duplication (TSD) length and proportions of
810 fragmented TE loci and nested TE insertions, each relative to the total TE copy number. The
811 assignment of copy number, sequence identity, and sequence integrity are further described
812 below.

813 ***Copy number assignment.*** In both Random Synthesized Genome and Custom Genome
814 modes, the user is required to provide a value range for the random assignment of a copy
815 number to each TE family. The TE Composition Approximation mode, however, uses the copy
816 number information of each family obtained from RepeatMasker as the exact copy number
817 value.

818 ***Sequence identity****.* Both Random Synthesised and Custom Genome modes assign sample
819 sequence identity values from predefined normal distributions. The mean value and the
820 standard deviation of the mean are, by default, randomly selected from ranges between 80 to
821 95 (%) and 1 to 20 (%), respectively. For the Genome Approximation mode, the mean and
822 standard deviation of sequence identity distributions are derived from the RepeatMasker output
823 for each TE family member. For each TE insertion, an integer is then randomly drawn from the
824 generated distribution, regarded as the original identity of the family sequence. This value is
825 further adjusted, following the approach implemented in denovoTE-eval [37], by adding 50% of
826 the difference between the sampled identity value and 100. It is calculated as $I' = I + (100 -$
827 $I)*0.5$, where $I$ is the original identity sampled from the normal distribution, and $I'$ is the adjusted

identity value. This adjustment causes a skew of the identity distribution to the right (i.e., closer
to 100%), reflecting that TE sequences with high divergence from the family sequence are less
likely to be recognized as family members.
***Target Site Duplication (TSD)***. TSD length is implemented based on literature findings [49], while
the sequence of TSD is randomly synthesized. The simulator then randomly assigns the TE
member to forward or reverse strands before performing sequence fragmentation and
integration.
***Sequence integrity***. Both Random Synthesized Genome and Custom Genome modes
implement a beta distribution to model the integrity of TE sequences. The default alpha and
beta values are 0.5 and 0.7, respectively, to create an asymmetric U-shape distribution in which
the probability of low sequence integrity (e.g., <10%) is higher than that of high integrity
(e.g.,>90%). Alpha and beta values can be adjusted using the `--alpha` (or `-a`) and `--beta` (or
`-b`) parameters. The integrity value for a TE member is sampled from the established beta
distribution and then converted to a number representing the bases to be removed from the 5'
end of that sequence. For example, if a TE member is 1 kbp at its full length and is assigned
with 70% integrity, the simulator would remove 300 bases, which accounts for 30% of the intact
length, from its 5' end. By default, only 0.1% of TE copies in each TE family remain untouched,
ensuring the representation of intact TEs in the simulation. For sequence integrity in the TE
Composition Approximation mode, TEgenomeSimulator utilizes a randomly sampled value from
a sequence integrity distribution extracted from the RepeatMasker output.
***Random and Nested TE Insertion***
TEgenomeSimulator randomly samples a value between 0 and 30% as the default proportion of
TE copy number to form nested insertions. This is only applied to LTR retrotransposons and is
implemented across all simulation modes. The simulator performs simple TE insertion before
executing nested TE insertion. Target sites for simple insertion and target TEs for nested
insertion are randomly chosen.
***Output Files of TEgenomeSimulator***
TEgenomeSimulator generates a series of output files that are useful for downstream analysis,
including the fasta file of the simulated genome, the fasta file of all TE sequences inserted in the
simulated genome, and the TE annotation file in GFF format that records the sequence identity,
integrity, and information to link the nested insertion to disrupted TEs.

## Implementation of PrinTE

### Burn-in

PrinTE simulates the forward evolution of genomes. The starting point for simulation is a burn-in
genome in FASTA format and a corresponding PrinTE-formatted BED file. These will serve as
the starting point (0 generation) for which no forward evolution has yet taken place. The burn-in
FASTA and BED can be generated by TEgenomeSimulator, RandSeqInsert, *de novo* by
PrinTE, or previous PrinTE outputs feeding back into PrinTE for additional forward simulations.
PrinTE *de novo* burn-in allows many customizations, including genome size, chromosome
number, percentage of the genome to be genic and TEs, and the ratio of each TE superfamily in
the genome. Each TE insertion creates a TSD with a length determined by the TE superfamily,
which is also customizable. In some instances, it may be beneficial to apply baseline mutation to

TEs in the burn-in such that they do not exactly match elements in the TE library. This is
particularly relevant for self-dating elements, like LTR-RTs, where the default TE library only
contains very young LTR-RTs, with exactly identical LTRs. For baseline mutations in the burn-in
genome, we apply a normalized exponential decay function to skew mutations heavily toward
lower mutation values:

The probability density function is aimed at flexibly and accurately modeling TE age distributions
as they are observed in real data, where aggressive TE purging leads to higher probability of
observing young TEs (**Fig. 5D**). The parameters $M_{max}$ (maximum mutation percentage; '--
`TE_mut_Mmax`') and $k$ (decay constant; '`--TE_mut_k`') are customizable, allowing users to
adjust the shape of the distribution to reflect a wide range of evolutionary and biological
scenarios and to match the observed TE half-life of any species (see **Fig. S4A,B**).

***Two Simulation Modules***

$$f(M) = \frac{k}{M_{max} \cdot (1 - e^{-k})} \cdot e^{-k \cdot \frac{M}{M_{max}}}$$

Forward simulation begins once the burn-in genome is established. Users can specify the total
number of generations ('`--generation_end`') and a smaller number of generations for each
iteration ('`--step`'). There are three sequential modules for forward simulation: 1) genome
mutation, 2) TE insertion, and 3) TE deletion. The pipeline will iterate over these processes for
each step until the target number of generations is simulated. PrinTE can iterate through
modules using constant and **fixed** TE insertion and deletion rates, or it can iterate using self-
adjusting **variable** TE insertion and deletion rates. The fixed and variable methods begin
identically, starting with the mutation of the genome. The expected number of DNA mutations
per sequence ($\lambda m$) is calculated as:

$$\lambda m = DNA\,mutation\,rate \times genome\,size \times generations$$
892
893           Because TE insertions can occur at any generation of a defined evolutionary window
(*i.e.*, a simulation step), they could accumulate DNA mutations less than $\lambda m$. To capture this
dynamic, we used an exponential decay function similar to the decay function used for the burn-
in genome to weight TE mutations toward lower values. We anchored the mutation distribution
here to approximate the complicated evolutionary dynamics of TE activities. The actual number
of mutations for each newly simulated TE is sampled from the probability density function,
$f(mTE)$, to help weight their observed mutations toward lower frequencies:

$$f(m_{TE}) = \frac{1.4 \times 10^{-6} \cdot e^{-1.4 \times 10^{-6} \cdot m_{TE}}}{1 - e^{-1.4 \times 10^{-6} \cdot 3,000,000}}, \quad 0 \leq m_{TE} \leq 3,000,000$$

***Simulation with Fixed Rates***

***TE insertion.*** After DNA mutations, TEs are inserted into the genome at random locations.
They may insert into intergenic regions, into other TEs, resulting in nested TE insertions, or into

genes, resulting in fragmented genes. Information about nesting and fragmentation is recorded
in the BED file. In the fixed rate method, the number of TEs to be inserted ($\lambda_{i-fix}$) is calculated
as:

$$\lambda_{i-fix} = TE\ insertion\ rate\ \times\ genome\ size\ \times\ generations$$

**TE deletion and solo LTR.** Donor TEs are randomly selected from the TE library based on
user-defined abundance ratios per superfamily. A nested insertion will fragment the acceptor TE
into two fragments. For example, if TE1 inserts into TE2, it will generate three sequences for
further operations (e.g., deletion), including two resulting fragments of TE2 and the complete
sequence of TE1. This differs from simple insertions into intergenic regions, where each
insertion results in only one sequence for further operations. During the deletion process, full-
length TEs and TE fragments are all eligible for deletion. The number of TE deletions ($\lambda_{d-fix}$) is
calculated as follows:

$$\lambda_{d-fix} = TE\ deletion\ rate\ \times\ genome\ size\ \times\ generations$$

Illegitimate recombination is a prevalent process for TE deletions and is facilitated by
sequence similarity between nearby homologous sequences [34]. Longer sequences have a
higher probability of causing illegitimate recombination than shorter sequences because there
are more similar base pairs to initiate pairing. As a result of this phenomenon, longer TEs have
a higher probability of facilitating illegitimate recombination and a higher likelihood of being
deleted than shorter TEs. We used an exponential decay function to model an exponentially
increasing probability of TE deletion as TE length increases:

$$f(L) = e^{-k \cdot \left(1 - \frac{L}{L_{95}}\right)}$$

Here, $L_{95}$ represents the TE length at the 95$^{th}$ percentile of the length distribution in the genome,
*L* represents the length of a TE (whether intact or fragmented), and *k* is a user-defined
parameter ('--k') that controls the rate at which deletion probability increases with sequence
length. This modification allows us to favor the deletion of longer TE features by assigning
higher deletion weights to sequences that are closer in length to the longest observed TE.
Except for LTR-RTs, TE deletions span the full length of the TE feature. If a nested TE is
selected for deletion, the adjacent TE fragments it disrupts are reconstituted into an intact TE.
LTR-RTs are an exception to full-length deletions because they can undergo recombination with
themselves owing to the sequence homology between their long terminal repeats, resulting in
the formation of a solo LTR. PrinTE supports deletions that lead to solo LTR formation. If an
LTR is selected for deletion, it will be converted to a solo LTR at a user-specified frequency,
which is set by default to 95% ('--solo_rate').

***Simulation with Variable Rates***

In addition to the fixed-rate TE insertion/deletion method outlined above, in which the number of
insertions and deletions per generation is constant and modeled by user-selected rates, PrinTE
also offers a variable method that self-adjusts the number of insertions and deletions per

generation based on genomic landscapes. This aims to replicate a subset of the biological
forces that affect TE insertions and deletions. The core processes, 1) genome mutation, 2) TE
insertion, and 3) TE deletion, are identical between methods. The genome mutation step is
identical between methods and is described above.

*TE insertions.* In real genomes, the rate of TE transposition is influenced by the number of
intact TEs, since only intact TEs are capable of initiating new insertions. Genomes with a high
number of intact TEs tend to experience more transpositions per generation than those with
few. Fragmented TEs and solo LTRs are no longer viable for transposition. In the variable
method, the number of TE insertions ($\lambda_{i-var}$) is modeled as follows:

949
$$\lambda_{i-var} = TE\ insertion\ rate\ \times\ number\ of\ intact\ TEs\ \times\ generations$$

This simulates insertions via transposition, but some TEs will be acquired vertically and drift into
the genome. We set that the frequency of vertically acquired transpositions is influenced by TE
prevalence within the species. We set that species with TE-rich genomes are more likely to
acquire additional TEs vertically, while species with TE-poor genomes are less likely to do so.
For this calculation, the TE composition of the genome is surveyed before simulations and used
as a proxy for TE abundance in the species. Vertically acquired TEs ($\lambda_{i-ver}$) are then simulated
accordingly:

957
$$\lambda_{i-ver} = TE\ vertical\ rate\ \times\ TE\ composition\ of\ the\ species\ \times\ generations$$

*Insertion bias.* The distribution of TEs across chromosomes is uneven and can be influenced
by chromatin states, with euchromatin generally more accessible for TE insertion. In the variable
method, we model this by treating genes and their surrounding regions as euchromatin, and the
gene-poor genomic backbone as heterochromatin. We can modulate insertion location
probability by chromatin type using a bias factor ('`--chromatin_bias_insert`'). We assign
insertion probability weights to euchromatic and heterochromatic sequences as follows:

$$Euchromatin\ weight = length \times bias\ factor; \quad Heterochromatin\ weight = length$$

It is important to note that PrinTE does not use genes as functional units, but rather as
representative segments of the genome to which cellular and selective attributes can be
assigned. This enables the modeling of global genome features. Unlike the fixed-rate method,
where TEs are inserted at random in proportions defined by the user, the variable-rate method
inserts TEs in proportions that reflect the current genomic composition.

*Fitness-dependent TE deletions.* TEs can trigger extensive and rapid proliferation, resulting in
a sudden increase in their genomic abundance. If left unchecked, this proliferation can
destabilize the genome and impose a selective disadvantage on the host organism. Under the
variable-rate method, the number of TE deletions is dynamically adjusted according to the
extent of gene disruption caused by TE insertions. In this context, genes are modeled as
sequences subject to purifying selection, whereby insertions of TEs into these regions are
deleterious and thus reduce the observed number of TE insertions. The number of TE deletions
($\lambda_{d-var}$) is then determined by the following expression:

In actual genomes, not all genes that confer a selective advantage exert the same influence on
fitness. While most TE insertions into genes are deleterious, some have more severe effects. To
capture this variability, we model the distribution of TE selection pressures across genes using a
log-normal distribution (**Fig. S4C**):

$$\lambda_{d\text{-}var} = TE\ deletion\ rate \times \sum (disrupted\ gene\ weights) \times generations$$

$$f(x;\sigma) = \frac{1}{x\sigma\sqrt{2\pi}}\ e^{-\frac{(\ln x - \sigma^2)^2}{2\sigma^2}}, \quad x > 0$$

In this framework, each disrupted gene contributes a positive value to the total disrupted
gene weight. The mode of the distribution is fixed at 1 to gain better control of the spread.
Because of the long tail of the distribution, a small subset of genes exerts a disproportionately
large influence on fitness. Consequently, TE insertions in these regions lead to a substantial
increase in deletion events. The parameter σ, which is set via the command line ('`--sigma`'),
determines the spread of the distribution. Larger values of σ produce longer tails and increase
the range of selective effects among genes. In reality, TE insertions in some genomic regions
can be lethal. PrinTE does not simulate lethality explicitly. Instead, we assume that simulating
many generations using a stepwise approach effectively filters out lethal insertions. Over time,
such deleterious types are eliminated from the population and do not persist in the simulated
genomic landscape.
***TE deletion bias.*** Once the number of deletions is calculated, the specific TEs to delete are
determined by the same exponential growth function ($f(L)$) used in the fixed-rate method
multiplied by additional genomic attributes that might affect the probability of TE deletion. These
genomic attributes include the selection coefficient parameter ('`--sel_coeff`') and the
chromatin bias parameter ('`--chromatin_bias_delete`'). The selection coefficient
parameter is used to assign additional deletion weight toward the TEs that disrupt genes.
Similarly, we can assign additional weight toward deletions in simulated euchromatin regions.
Longer TEs and TEs nearer to or within impacted genes are more likely to be deleted based on
user-specified thresholds.

## Searching for TE Dynamic Parameters using PrinTE

1002   To identify optimal TE dynamics in a given genome, we employed a systematic parameter
1003   sweep approach using PrinTE. This method involved iteratively testing combinations of key
1004   parameters to approximate empirical TE behavior and genome size observed in the genome of
1005   interest. The '`pipeline.report`' file contains an updating log of TE insertion and deletion
1006   counts per simulated iteration and can also be useful in monitoring PrinTE runs.

### *Parameter Space Exploration*

We conducted a grid search over two critical parameters, insertion and deletion rates, governing
TE insertion dynamics. Users can specify parameter ranges (upper and lower limits) and
increments of parameters. Based on the hypothesis of genome size dynamics, constraints on
the deletion rate can be applied to narrow down the search space. For example, if a genome

evolves with a stable size, a deletion rate between 1.2 to 1.5 folds of the insertion rate can be
used. We also monitor the size of simulated genomes and abort simulations when over or under
the user-set size to ensure strict resource consumption. For example, if the target genome size
is 150 Mb, simulation parameters that create genomes over 180 Mb in any steps are deemed
unfit for the hypothesis and thus aborted to reduce the computation load. PrinTE's built-in '--
max_size' parameter can be used to set the upper limit of genome size and stop the simulation
when exceeding the specified size.

1019

Additional parameters included length bias for deletions ('--k') and solo LTR rates ('--
solo_rate'). In practice, we found k values of 0, 3, or 6 and solo rates from 70% to 95% (with
5% increments) could provide sufficient dynamics on top of the insertion and deletion rates.
Higher k value will lead to slower genome size increase due to the preference for deleting
longer TEs (**Fig. S8**). Higher solo rate will lead to faster genome size increase due to the
creation of solo LTRs instead of full deletions (**Figs. S9, S10**) and increased solo:intact ratio.

### High-Throughput Simulation Framework

PrinTE simulations were executed through an HPC cluster using array jobs (SLURM workload
manager), enabling parallel processing of parameter sets. Users can set the total number of
generations ('-ge') with sampling steps ('-st') for all simulations during parameter sweeps.

The species-specific substitution rate ('--mutation_rate') is also critical to the
simulation outcome. In cases where an accurate substitution rate is not available for the
species, users can try rates in different magnitudes (e.g., $1 \times 10^{-8}$, $1 \times 10^{-9}$, or $1 \times 10^{-10}$ per bp per
generation) in the simulation, then compare the LTR divergence of simulated genomes to the
real genome. Higher mutation rates will introduce more mutations to TEs in a given timeframe,
thus creating gentler slopes of LTR divergence. In reverse, lower mutation rates will create
steeper slopes of LTR divergence.

### Empirical Estimation of LTR Divergence and solo:intact Ratio
Outputs were processed through a standardized pipeline to assess TE dynamics, including
genome size change, LTR divergence distributions, and solo:intact ratio. Intact LTR elements
were extracted from each simulation using the "extract_intact_LTR.py" script; their
divergences were estimated with the K2P model and a mutation rate of $2.0 \times 10^{-9}$ per bp per
generation using the "seq_divergence.py" script, and divergences were plotted using the
"ltr_dens.py" script, all available in PrinTE.

Although the BED file output of PrinTE contains the annotation of all intact and solo LTR
sequences, we perform a *de novo* scan of intact LTR-RTs using LTR_retriever (v3.0.1) [33] on the
simulated genomes, so that these results are comparable to those obtained from the real
genome. The resulting LTR library was used to annotate whole-genome LTR sequences with
RepeatMasker (v4.1.7-p1) [48]. Solo LTRs were then identified by the
"solo_intact_ratio_wrap.sh" script available in the LTR_retriever package, which also
calculated the overall solo:intact ratio of each genome. For the same reason, LTR divergence
distributions of simulated genomes were also plotted using intact LTR-RTs identified by

LTR_retriever. The "`plot_divergence.R`" script available in PrinTE was used to plot the
density of LTR divergence in simulated and real genomes. Parameter sets were ranked by their
ability to reproduce three empirical features: 1) Simulated genome sizes are consistent with
hypothesis predictions; 2) Genome-wide LTR divergence distributions are consistent with
empirical whole-genome data; 3) Solo:intact LTR ratios are close to the empirical value. This
approach identified optimal parameters where simulated TE dynamics showed convergence
with multiple empirical metrics while maintaining computational feasibility.

## Syntenic Ribbon Plot and LTR-RT Phylogeny

To simulate genomes presented in **Fig. 3A**, 19,621 *Arabidopsis thaliana* cds sequences were
inserted into the simulated genome using PrinTE with "`--cds_num 19621`" to guide collinear
anchoring and enable syntenic analysis. The simulated genomes were processed pairwise
using "`jcvi.compara.catalog ortholog --cscore=.99`" in JCVI (v1.5.4) [50]. Syntenic
anchors were visualized using "`matplotlib.pyplot`" in Matplotlib (v3.10.3) to create the
ribbon plot (**Fig. 3A**). The LTR-RT phylogeny (**Fig. 3D,E**) was created using only elements in
the *Tekay* family, which were classified as such by TEsorter (v1.4.7) [51]. *Tekay* LTR-RT headers
were tagged with "`_shared`" and "`_uniq`" to facilitate post-PrinTE parsing. Shared LTR-RTs
were inserted using PrinTE with parameters "`-p 80 -cn 1 -sz 10Mb -tmx 10 --TE_lib`
`Tekay_shared.fa -tk 15`". The outputs ("`burnin.fa`" and "`burnin.bed`") were
duplicated and allowed to continue simulation using either genome expansion parameter ( "`--`
`generation_end 5000000 --step 500000 -F 2e-10,1e-10 --TE_lib`
`Tekay_uniq.fa`") or genome contraction parameters ( "`--generation_end 5000000 --`
`step 500000 -F -F 2e-11,1e-11 --TE_lib Tekay_uniq.fa`"). Intact LTR-RTs were
extracted using "`extract_intact_LTR.py`" and classified using TEsorter with "`-db rexdb-`
`plant -dp2`". Reverse transcriptase domains identified from *Tekay* LTR-RTs were aligned
using MAFFT (v7.525) [52] and provided to VeryFastTree (v4.0.5) [53] for phylogenetic analysis.
The resulting newick trees were plotted using ggtree (v3.1.5) [54] with branches colored by their
ancestral relationship.

## *Pucciniomycotina* Genomes and Annotations

The myrtle rust (*Austropuccinia psidii*) genome was downloaded from NCBI with accession ID
ERS3953144 [3]. Genomes of the *Pucciniomycotina* fungus subdivision were obtained from the
JGI Fungal Genomics Resource MycoCosm
(https://mycocosm.jgi.doe.gov/pucciniomycotina/pucciniomycotina.info.html) (**Table S3**). Rust
fungus genomes with contig N50 values higher than 100 kbp were used, for a total of 47
genomes (**Table S4**).
Benchmarking Universal Single-Copy Orthologs (BUSCOs) evaluations were performed
on each genome by compleasm (v0.2.7) [40] with the odb12 database and the basidiomycota
lineage that has 2409 universal single-copy orthologs. For genomes with duplicated BUSCO
higher than 5%, the Redundans (v2020.01.28) [39] program was used to identify alternative
haplotypes. Haplotype alignments with an identity higher than 90% and coverage of more than
80% were removed using "`--identity 0.9 --overlap 80`" parameters. compleasm was

again used to evaluate the purged genomes with the same parameter. The resulting duplicated
BUSCO were considered biological duplications (**Table S4**). The haplotype genomes (de-
duplicated) were used in the following analyses.
Genomes with single-copy BUSCOs higher than 50% were used to construct the
phylogeny. Melame1 (BUSCO S:1134,D:1147,F:33,I:0,M:95) and MecolCla_1 (BUSCO
S:826,D:1452,F:34,I:0,M:97) were not included in this analysis owing to the high proportion of
duplicated BUSCO. The protein sequences of single-copy BUSCOs present in the remaining 45
genomes were extracted and further aligned using mafft (v7.525) [52] with a focus on local
alignments using the "`--maxiterate 1000 --localpair`" parameters. The phylogeny was
constructed using the aligned BUSCO and iqtree (v2.4.0) [41]. Each BUSCO alignment was
treated as an individual partition [55], and ModelFinder [56] was used to identify the best model for
each partition using the "`-m MFP`" parameter in iqtree. The Shimodaira–Hasegawa-like
approximate likelihood ratio test (SH-aLRT) [57] and bootstrap tests were performed 1000 times
each with parameters "-bb 1000 -alrt 1000" in iqtree. The bootstrap consensus tree was plotted
by ggtree (v3.1.5) [58] in R.
Transposable elements in each fungal genome were annotated using EDTA (v2.2.2) [59]
following the instructions of Benson et al. (2024) [60]. The "`--force 1`" parameter was used to
overcome analysis in genomes with missing types of TEs (*i.e.*, LTR retrotransposons). The "`--
anno 1`" parameter was used to obtain whole-genome TE annotations. The "`--cds`" parameter
was used to provide coding sequences to EDTA to improve TE annotations. Primary coding
sequences of each genome were obtained from JGI MycoCosm and Tobias et al. (2021) [3]. Solo
and intact LTR retrotransposons were identified from whole-genome TE annotations using
LTR_retriever (v3.0.1) [33]. The solo:intact ratio was calculated on the whole-genome level, which
is the number of whole-genome solo LTR divided by the number of whole-genome intact LTR.
"inf" values were given to genomes without intact LTR retrotransposons. The divergence of
intact LTR retrotransposons was estimated by LTR_retriever (v3.0.1) [33] using the one-
parameter Jukes-Cantor model [61].

## TIR Insertion Simulation with RandSeqInsert

To demonstrate RandSeqInsert's capabilities, three targeted simulations were conducted. Two
simulations with 10 rice TIR transposons as donor sequences were inserted into the
*Arabidopsis* Col-CEN mitochondrial chromosome (ChrM) [10] as the acceptor sequence. One
simulation utilized the "`--visual`" parameter to generate DOT visualizations of the balanced
sequence tree structure (**Fig. S1A**), while the second simulation's insertion events were
recaptured using minigraph (v0.21) [29] with default parameters and visualized with Bandage [62]. A
larger-scale simulation was then performed by inserting 30,000 rice TIR transposons into a 350
Mb random acceptor sequence under two conditions: with and without TSD (9 bp). The inserted
elements were subsequently redetected using TIR-Learner (v3.0.7)
(https://github.com/lutianyu2001/TIR-Learner) [63] with the "`--species others`" parameter and
compared against RandSeqInsert's record to calculate the detection rate.

## Simulating Digital Replicas of *Arabidopsis thaliana* and Rust Fungus Genomes

The TAIR10 *A. thaliana* genome was downloaded from NCBI with accession ID GCF_000001735.4. The *Arabidopsis* TE library for simulation and TE removal was obtained from the EDTA GitHub repository (https://github.com/oushujun/EDTA/blob/master/database/) [59]. Three fungal genomes were chosen for simulations, including *Puccinia coronata* f. sp. *avenae* (PuccoNC29_1), *Cronartium ribicola* (Crori1), and *Austropuccinia psidii* (myrtle rust). TE libraries used for TE removal and simulating the fungal genomes were obtained via *de novo* TE annotation using EDTA (as described above). The Genome Approximation Mode of TEgenomeSimulator was used for simulations. The resulting digital replicas of the rust fungus genomes were further used to simulate LTR dynamics using PrinTE (see sections below).

## Forward Simulation of Rust Fungus Genomes

### Simulating LTR Dynamics in Rust Fungus Genomes

***Cronartium ribicola***. We hypothesized that this genome is relatively stable, with high rates of LTR insertions and deletions. We aimed to evolve the genome with a relatively stable genome size. The digital replica of the *C. ribicola* genome was generated by TEGenomeSimulator with a size of 232.6 Mb, which was used as input for PrinTE using the '`--bed`' and '`--fasta`' parameters. Each experiment was simulated with 20,000 generations per step and a total of 200,000 generations (`-ge 200000 -st 20000`). A series of fixed insertion and deletion rates were used in different experiments, including the range of insertion rates from $1.0\times10^{-9}$ to $1.9\times10^{-8}$ per bp per generation and the range of deletion rates from $1.4\times10^{-9}$ to $4.0\times10^{-8}$ per bp per generation specified by the '`--fix`' parameter. Additionally, DNA substitution rates of $1.0\times10^{-8}$, $2.0\times10^{-9}$, and $5.0\times10^{-9}$ per bp per generation, k values of 0, 3, 6, and 10, and solo rates of 60, 70, 75, 80, and 95 were used in the parameter sweep analysis. The simulated genomes were summarized by PrinTE and LTR_retriever and compared with the real *C. ribicola* genome for the determination of the best parameters.

***Puccinia coronata*** **f. sp.** ***avenae.*** We hypothesized that this genome is relatively stable with low rates of LTR insertions and deletions. We aimed to evolve the genome with a relatively stable genome size. The digital replica of the *P. coronata* f. sp. *avenae* genome was generated by TEGenomeSimulator with a size of 112.7 Mb, which was used as input for PrinTE using the '`--bed`' and '`--fasta`' parameters. Each experiment was simulated with 1 million generations per step and a total of 10 million generations (`-ge 10000000 -st 1000000`). A series of fixed insertion and deletion rates were used in different experiments, including the range of insertion rates from $8.0\times10^{-12}$ to $5.0\times10^{-11}$ per bp per generation and the range of deletion rates from $9.6\times10^{-12}$ to $9.0\times10^{-11}$ per generation specified by the '`--fix`' parameter. Additionally, mutation rates of $2.0\times10^{-9}$ and $5.0\times10^{-9}$ per bp per generation, k values of 0, 3, and 6, and solo rates of 70 and 95 were used in the parameter sweep analysis. The simulated genomes were summarized by PrinTE and compared with the real *P. coronata* f. sp. *avenae* genome for the determination of the best parameters.

***Austropuccinia psidii***. We hypothesized that this was once an expanding genome but now a fast downsizing genome with a low rate of LTR insertion and a high rate of LTR deletion. Owing

to the relative phylogenetic closeness to *A. psidii*, we use the genome size and TE composition of *P. coronata* f. sp. *avenae* genome as an approximation of the ancestral state of *A. psidii* before genome expansion. We used PrinTE to simulate such an ancestral genome with parameters '`--cds_percent 20 --TE_num 6000 --chr_number 20 --size 113Mb --TE_mut_k 10 --TE_mut_Mmax 20`'. With a fixed deletion rate of $5.0\times10^{-13}$ per bp per generation, we explored insertion rates from $2.0\times10^{-12}$ to $9.8\times10^{-11}$ per bp per generation and identified that the insertion rate of $1.4\times10^{-11}$ per bp per generation could yield LTR divergence similar to the expansion phase of *A. psidii* (**Fig. S16A,B**). We thus used PrinTE to simulate a fast-expanding phase with parameters "`-k 0 --fix 1.4e-11,5e-13 --mutation_rate 2e-9 --solo_rate 95`" until it reached 1,178 Mb. For the contraction phase, we slowed down the insertion rate to $1.4\times10^{-13}$ per bp per generation and explored deletion rates from $3.0\times10^{-13}$ to $5.0\times10^{-11}$ per bp per generation. We identified that the deletion rate of $3.0\times10^{-11}$ per bp per generation could yield LTR divergence and genome size similar to those of the *A. psidii* genome (**Fig. S16A,B**). Finally, we used the outputs of the last iteration as inputs for PrinTE to simulate a fast-contracting phase with parameters "`-k 0 --fix 1.4e-13,3e-11 --mutation_rate 2e-9 --solo_rate 95`" until it reached 1000 Gb, approximately the size of the *A. psidii* genome.

## Use of AI in this study

ChatGPT 4o, o3-mini, o1, o3, o4-mini, and o4-high, Qwen3-235B-A22B, and DeepSeek R1 were used to help draft and improve scripts during the development of PrinTE and TEgenomeSimulator.

# **Data Availability**

All codes that produced the findings of the study, including all main and supplemental figures, are available at https://github.com/cwb14/TESS, under the GPL-3.0 license. TEgenomeSimulator is available at https://github.com/Plant-Food-Research-Open/TEgenomeSimulator under the GPL-3.0 license.

# **Acknowledgments**

We want to thank Drs. Jason Slot, James Pease, and H. Lisle Gibbs for helpful discussions of this project. We are also grateful to Drs. Chen Wu, Julie Blommaert, and David Chagné for reviewing this manuscript and providing insightful comments.

# **Funding**

This research was supported by the Ohio State University STEM Education Faculty Startup Award (S.O.), Undergraduate Research Access Innovation Seed Grant (S.O.), the United States Department of Agriculture (USDA) National Institute of Food and Agriculture (NIFA) Agriculture and Food Research Initiative (AFRI) Postdoctoral Fellowship (AWD-117797 to C.W.B.), and by

1207 Plant & Food Research through their Kiwifruit Royalty Investment Programme (T.-H.C., O.A.B.,
1208 C.H.D., and S.J.T.).

# Author information

1210 Both Christopher W. Benson and Ting-Hsuan Chen contributed equally to this work and have
1211 the right to list their name first in their CV.

## Authors and Affiliations

1213 Department of Molecular Genetics, Ohio State University, Columbus, OH, 43210, USA
1214 Christopher W. Benson, Tianyu Lu & Shujun Ou
1215
1216 The New Zealand Institute for Plant and Food Research Limited, Lincoln 7608, New Zealand
1217 Ting-Hsuan Chen & Susan J. Thomson
1218
1219 The New Zealand Institute for Plant and Food Research Limited, Palmerston North 4442, New
1220 Zealand
1221 Olivia Angelin-Bonnet
1222
1223 The New Zealand Institute for Plant and Food Research Limited, Auckland 1025, New Zealand
1224 Cecilia H. Deng

## Contributions

1226 S.O., S.J.T., and C.H.D. conceived the project and supervised the work. T.L. developed
1227 RandSeqInsert, T.-H.C. developed TEgenomeSimulator, and C.W.B developed PrinTE and the
1228 TESS framework. C.W.B, T.-H.C., T.L., O.A.B., and S.O. performed the analysis. C.W.B, T.-
1229 H.C., S.J.T., C.H.D., and S.O. wrote the manuscript text. All authors reviewed the manuscript.

## Corresponding author

1231 Correspondence to Ting-Hsuan Chen ([Ting-Hsuan.Chen@plantandfood.co.nz](mailto:Ting-Hsuan.Chen@plantandfood.co.nz)) and Shujun Ou
1232 ([ou.195@osu.edu](mailto:ou.195@osu.edu)).

# Ethics declarations

## Ethics approval and consent to participate

1235 Not applicable.

## Competing interests

1237 The authors declare that they have no competing interest.

# References

1. Lander, E. S. *et al.* Initial sequencing and analysis of the human genome. *Nature* **409**, 860–921 (2001).

2. Hufford, M. B. *et al.* De novo assembly, annotation, and comparative analysis of 26 diverse maize genomes. *Science* **373**, 655–662 (2021).

3. Tobias, P. A. *et al.* Austropuccinia psidii, causing myrtle rust, has a gigabase-sized genome shaped by transposable elements. *G3 (Bethesda)* **11**, (2021).

4. Serrato-Capuchina, A. & Matute, D. R. The Role of Transposable Elements in Speciation. *Genes* **9**, 254 (2018).

5. Beric, A. *et al.* Comparative phylogenetics of repetitive elements in a diverse order of flowering plants (Brassicales). *G3 (Bethesda)* **11**, (2021).

6. Qin, P. *et al.* Pan-genome analysis of 33 genetically diverse rice accessions reveals hidden genomic variations. *Cell* **184**, 3542–3558.e16 (2021).

7. Bayer, P. E. *et al.* Variation in abundance of predicted resistance genes in the Brassica oleracea pangenome. *Plant Biotechnol. J.* **17**, 789–800 (2019).

8. Ou, S. *et al.* Differences in activity and stability drive transposable element variation in tropical and temperate maize. *Genome Res* **34**, 1140–1153 (2024).

9. Hu, Y. *et al.* Gossypium barbadense and Gossypium hirsutum genomes provide insights into the origin and evolution of allotetraploid cotton. *Nat Genet* **51**, 739–748 (2019).

10. Naish, M. *et al.* The genetic and epigenetic landscape of the Arabidopsis centromeres. *Science* **374**, eabi7489 (2021).

11. Blommaert, J. Genome size evolution: towards new model systems for old questions. *Proc. Biol. Sci.* **287**, 20201441 (2020).

12. Kapusta, A., Suh, A. & Feschotte, C. Dynamics of genome size evolution in birds and mammals. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E1460–E1469 (2017).

13. Ricci, M., Peona, V., Guichard, E., Taccioli, C. & Boattini, A. Transposable elements activity

is positively related to rate of speciation in mammals. *J. Mol. Evol.* **86**, 303–310 (2018).

14. Pellicer, J., Hidalgo, O., Dodsworth, S. & Leitch, I. J. Genome size diversity and its impact

on the evolution of land plants. *Genes (Basel)* **9**, (2018).

15. Niu, S. *et al.* The Chinese pine genome and methylome unveil key features of conifer

evolution. *Cell* **185**, 204–217.e14 (2022).

16. Liang, Y. *et al.* The giant genome of lily provides insights into the hybridization of cultivated

lilies. *Nat. Commun.* **16**, 45 (2025).

17. Fernández, P. *et al.* A 160 Gbp fork fern genome shatters size record for eukaryotes.

*iScience* **27**, 109889 (2024).

18. Wright, S. I. & Agren, J. A. Sizing up Arabidopsis genome evolution. *Heredity (Edinb.)* **107**,

509–510 (2011).

19. Kovaka, S., Ou, S., Jenike, K. M. & Schatz, M. C. Approaching complete genomes,

transcriptomes and epi-omes with accurate long-read sequencing. *Nat. Methods* **20**, 12–16

(2023).

20. Hoen, D. R. *et al.* A call for benchmarking transposable element annotation methods. *Mob.*

*DNA* **6**, 13 (2015).

21. Clough, J. E., Foster, J. A., Barnett, M. & Wichman, H. A. Computer simulation of

transposable element evolution: random template and strict master models. *J Mol Evol* **42**,

52–58 (1996).

22. Kofler, R. SimulaTE: simulating complex landscapes of transposable elements of

populations. *Bioinformatics* **34**, 1439 (2018).

23. Bourgeois, Y., Ruggiero, R. P., Hariyani, I. & Boissinot, S. Disentangling the determinants

of transposable elements dynamics in vertebrate genomes using empirical evidences and

simulations. *PLoS Genet* **16**, e1009082 (2020).

24. Deceliere, G., Letrillard, Y., Charles, S. & Biémont, C. TESD: a transposable element

dynamics simulation environment. *Bioinformatics* **22**, 2702–2703 (2006).

25. Verneret, M., Van Anthony, L., Faraut, T., Turpin, J. & Lerat, E. Particular sequence

characteristics induce bias in the detection of polymorphic transposable element insertions.

*bioRxiv* 2024.09.25.614865 (2024) doi:10.1101/2024.09.25.614865.

26. Groza, C., Chen, X., Wheeler, T. J., Bourque, G. & Goubert, C. A unified framework to

analyze transposable element insertion polymorphisms using graph genomes. *Nat*

*Commun* **15**, 8915 (2024).

27. Yue, J.-X. & Liti, G. simuG: a general-purpose genome simulator. *Bioinformatics* **35**, 4442–

4444 (2019).

28. Caballero, J., Smit, A. F. A., Hood, L. & Glusman, G. Realistic artificial DNA sequences as

negative controls for computational genomics. *Nucleic Acids Res* **42**, e99 (2014).

29. Li, H., Feng, X. & Chu, C. The design and construction of reference pangenome graphs

with minigraph. *Genome Biol.* **21**, 265 (2020).

30. Rodriguez, M. & Makałowski, W. Software evaluation for de novo detection of transposons.

*Mobile DNA* **13**, 1–14 (2022).

31. Tusso, S., Suo, F., Liang, Y., Du, L.-L. & Wolf, J. B. W. Reactivation of transposable

elements following hybridization in fission yeast. *Genome Res.* **32**, 324–336 (2022).

32. Nowoshilow, S. *et al.* The axolotl genome and the evolution of key tissue formation

regulators. *Nature* **554**, 50–55 (2018).

33. Ou, S. & Jiang, N. LTR_retriever: A Highly Accurate and Sensitive Program for

Identification of Long Terminal Repeat Retrotransposons. *Plant Physiol.* **176**, 1410–1422

(2018).

34. Ma, J., Devos, K. M. & Bennetzen, J. L. Analyses of LTR-retrotransposon structures reveal

recent and rapid genomic DNA loss in rice. *Genome Res.* **14**, 860–869 (2004).

35. Swarbreck, D. *et al.* The Arabidopsis Information Resource (TAIR): gene structure and

function annotation. *Nucleic Acids Res.* **36**, D1009–14 (2008).

36.  Chen, T.-H. & Winefield, C. Comprehensive analysis of both long and short read

transcriptomes of a clonal and a seed-propagated model species reveal the prerequisites

for transcriptional activation of autonomous and non-autonomous transposons in plants.

*Mob. DNA* **13**, 16 (2022).

37.  Ahrendt, S. R., Mondo, S. J., Haridas, S. & Grigoriev, I. V. MycoCosm, the JGI's fungal

genome portal for comparative genomic and multiomics data analyses. *Methods Mol. Biol.*

**2605**, 271–291 (2023).

38.  *Systematics and Evolution*. (Not Avail, 2014). doi:10.1007/978-3-642-55318-9.

39.  Pryszcz, L. P. & Gabaldón, T. Redundans: an assembly pipeline for highly heterozygous

genomes. *Nucleic Acids Res.* **44**, e113 (2016).

40.  Huang, N. & Li, H. compleasm: a faster and more accurate reimplementation of BUSCO.

*Bioinformatics* **39**, btad595 (2023).

41.  Minh, B. Q. *et al.* IQ-TREE 2: New models and efficient methods for phylogenetic inference

in the genomic era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

42.  Devos, K. M., Brown, J. K. M. & Bennetzen, J. L. Genome size reduction through

illegitimate recombination counteracts genome expansion in Arabidopsis. *Genome Res.* **12**,

1075–1079 (2002).

43.  Edwards, H. M. & Rhodes, J. Accounting for the biological complexity of pathogenic fungi in

phylogenetic dating. *J. Fungi (Basel)* **7**, 661 (2021).

44.  Osmanski, A. B. *et al.* Insights into mammalian TE diversity through the curation of 248

genome assemblies. *Science* **380**, eabn1430 (2023).

45.  Zoonomia Consortium. A comparative genomics multitool for scientific discovery and

conservation. *Nature* **587**, 240–245 (2020).

46.  G. M. Adel'son-Vel'skii, E. M. Landis, 'An algorithm for organization of information', Dokl.

Akad. Nauk SSSR, 146:2 (1962), 263–266.

https://www.mathnet.ru/php/archive.phtml?wshow=paper&jrnid=dan&paperid=26964&optio

n_lang=eng.

47. Gansner, E. R. & North, S. C. An open graph visualization system and its applications to software engineering. *Softw. Pract. Exp.* **30**, 1203–1233 (2000).

48. Smit, A. F. A., Hubley, R. & Green, P. RepeatMasker Open-4.0. 2013--2015. Preprint at (2015).

49. Zhao, D., Ferguson, A. A. & Jiang, N. What makes up plant genomes: The vanishing line between transposable elements and genes. *Biochim. Biophys. Acta* **1859**, 366–380 (2016).

50. Tang, H. *et al.* JCVI: A versatile toolkit for comparative genomics analysis. *Imeta* **3**, e211 (2024).

51. Zhang, R.-G. *et al.* TEsorter: an accurate and fast method to classify LTR-retrotransposons in plant genomes. *Hortic. Res.* **9**, uhac017 (2022).

52. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

53. Piñeiro, C., Abuín, J. M. & Pichel, J. C. Very Fast Tree: speeding up the estimation of phylogenies for large alignments through parallelization and vectorization strategies. *Bioinformatics* **36**, 4658–4659 (2020).

54. Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T.-Y. Ggtree: An r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol. Evol.* **8**, 28–36 (2017).

55. Chernomor, O., von Haeseler, A. & Minh, B. Q. Terrace aware data structure for phylogenomic inference from supermatrices. *Syst. Biol.* **65**, 997–1008 (2016).

56. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**, 587–589 (2017).

57. Anisimova, M. & Gascuel, O. Approximate likelihood-ratio test for branches: A fast, accurate, and powerful alternative. *Syst. Biol.* **55**, 539–552 (2006).

58. Xu, S. *et al.* Ggtree: A serialized data object for visualization of a phylogenetic tree and

annotation data. *Imeta* **1**, e56 (2022).

59. Ou, S. *et al.* Benchmarking transposable element annotation methods for creation of a

streamlined, comprehensive pipeline. *Genome Biol.* **20**, 275 (2019).

60. Benson, C. W., Heringer, P. & Ou, S. Four strategies for whole-genome annotation of

transposable elements and repeats in maize. *Cold Spring Harb. Protoc.* (2024)

doi:10.1101/pdb.prot108578.

61. Jukes, T. H. & Cantor, C. R. Evolution of Protein Molecules. in *Mammalian Protein*

*Metabolism* 21–132 (Elsevier, 1969).

62. Wick, R. R., Schultz, M. B., Zobel, J. & Holt, K. E. Bandage: interactive visualization of de

novo genome assemblies. *Bioinformatics* **31**, 3350–3352 (2015).

63. Su, W., Gu, X. & Peterson, T. TIR-Learner, a New Ensemble Method for TIR Transposable

Element Annotation, Provides Evidence for Abundant New Transposable Elements in the

Maize Genome. *Mol. Plant* **12**, 447–460 (2019).