PrinTE: A Forward Simulation Framework for Studying the Role of Transposable Elements in Genome Expansion and Contraction

Authors:

Christopher W. Benson[1], Ting-Hsuan Chen[2], Susan J. Thomson[2], Cecilia H. Deng[3], Shujun Ou[1]*

1 Department of Molecular Genetics, Ohio State University, Columbus, OH, 43210, USA

2 Plant and Food Research Group, The New Zealand Institute for Bioeconomy Science Limited, Lincoln 7608, New Zealand

3 Plant and Food Research Group, The New Zealand Institute for Bioeconomy Science Limited, Auckland, 1025, New Zealand

* Correspondence: ou.195@osu.edu

Abstract

Genome expansion and contraction are reportedly driven by transposable element (TE) activity, but the underlying dynamics remain enigmatic due to a lack of historical records tracing these changes. Here, we present PrinTE for versatile, forward-time simulation of whole-genome sequences with highly customizable transposon dynamics. Through simulations, we confirm that the distribution of TE sequence divergence reflects their historical insertion and deletion dynamics, which can be used to infer TE dynamic parameters through PrinTE simulations. We analyzed the pangenome of *Pucciniomycotina*, a subdivision of fungi containing myrtle rust (*Austropuccinia psidii*), which drastically expanded its genome size to 1018 Mb. Our analyses reveal that the best strategy for controlling genome size is to avoid the invasion of LTR retrotransposons (LTR-RTs). While illegitimate recombination (IR) is considered the most effective counteraction of LTR-RT invasions leaving only solo LTR remnants, we observed a strong positive correlation between solo:intact LTR ratio (strength of LTR-RT removal) and genome size ($r = 0.65$), and a near-linear correlation between solo LTR count and genome size ($r = 0.98$). This result suggests that IR alone may not effectively prevent genome obesity. Through simulation of *Pucciniomycotina* genomes, we proposed that *A. psidii* might experience a prolonged period of genome expansion followed by a short, potent, and likely ongoing period of contraction. PrinTE is freely available at https://github.com/cwb14/PrinTE.git.

Keywords: Transposable Element, Genome Size, Forward Simulation, InDels, Solo LTR, Speciation, Rust Fungus

Background

Transposable elements (TEs), or "jumping genes", are dynamic DNA sequences that replicate and relocate within genomes. TEs constitute 46% of the human (*Homo sapiens*) genome [1], 85% of the maize (*Zea mays*) genome [2], and over 90% of the genome of the fungal pathogen *Austropuccinia psidii* that causes the myrtle rust disease [3]. The prevalence of TEs across the Tree of Life reflects their dual roles as both genomic parasites and drivers of evolutionary innovation. Beyond inflating genome sizes, TEs drive structural variation, modulate gene expression, and facilitate chromosomal rearrangements [4]. In both plant and mammalian genomes, bursts of TE activity correlate with speciation events and adaptive radiations, enabling rapid responses to environmental stressors like pathogens or climate shifts [5].

Recent studies highlight TE's contribution to the collective genetic repertoire of a species. Core genomes, conserved across individuals, are often gene-rich and TE-poor [2], while "variable" or "dispensable" regions are TE-dense and enriched for stress-response genes, pathogen resistance loci, and structural variants [6]. In *Brassica oleracea*, resistance gene analogs cluster in TE-rich variable regions [7]. Similarly, tandem clusters of leucine-rich repeat (LRR) genes in maize are found in genomic regions rich with young long terminal repeat (LTR) retrotransposons (LTR-RT) [8], suggesting that TE-driven structural plasticity contributes to adaptive traits. Related lineages can become karyotypically disparate over generations due to differences in their ability to repress TEs [9]. The global impact of TEs on macro-chromosomal structure plays an important role in evolutionary fates by influencing meiotic stability and similarity between cytotypes [10]. These findings underscore TEs as dynamic regulators of genome evolution, influencing both microevolutionary adaptation and macroevolutionary diversification.

Genome size varies considerably across taxa. Mammalian and bird genomes appear more tightly constrained in genome size than those of both fungi and plants [11]. Numerous studies across the Mammalia class indicate this size constraint is not through a lack of TE radiation bursts [12] but through mechanistic interplay with purifying selection, where bursts are followed by TE silencing [13] and DNA loss [12]. Genome size variation in plants facilitates adaptation to specific environments, primarily carried out by whole genome duplication (WGD) and TE expansion [14], especially LTR-RTs [15]. Extremes of genome size can be observed from the compact genome of *Arabidopsis thaliana* (~135 Mb) through to the giant genomes of the Chinese pine (*Pinus tabuliformis*) at ~25 Gb [16], lily (*Lilium sargentiae*) at ~36 Gb [17], and the New Caledonian fork fern (*Tmesipteris oblanceolata*) at ~160 Gb [18]. These are examples of unconstrained genome expansion, with extremely high TE content within intergenic regions and introns. Conversely, there are numerous signs that genome size expansion in plants is subject to constraints and regulatory mechanisms. *Arabidopsis* has undergone several WGDs, yet remains compact partially through subsequent gene loss that purges redundant paralogs [19]. The dynamic of genome size change has been difficult to accurately evaluate primarily because of the lack of high-fidelity tools for simulation and modeling studies.

Advances in long-read sequencing and computational methods have empowered efforts in resolving TE dynamics [20], yet reconstructing their evolutionary history remains challenging due to TEs often being fragmented, nested, and highly identical to each other. Simulations of TE evolution are thus indispensable for testing hypotheses about genome diversification, evaluating

TE annotation tools, and modeling the interplay between TE activity and selection. By
generating synthetic genomes with known TE architectures, including nested insertions,
sequence divergence, and polymorphism, researchers can disentangle the stochastic and
deterministic forces shaping genome evolution [21].
Existing TE simulators vary widely in scope and range of biological scenarios that they
mimic. Early models, such as those by Clough et al. (1996), introduced foundational concepts
by simulating varying modes of mammalian transposition [22]. Modern tools address specific
niches. SimulaTE [23] generates complex TE landscapes through predefined TE architectures,
limiting its utility for exploring stochastic evolutionary scenarios. Population-genetic frameworks
like SLiM [24] excel at modeling selection and demography but compress TEs into individual units,
precluding modeling of TE sequence mutagenesis, nested insertion, and the consequential
genome size evolution. TESD [25] incorporates spatial population structure and selection against
deleterious insertions, yet its inaccessibility hinders further application. ReplicaTE [26] engineers
synthetic genomes by deleting and reinserting TEs into real assemblies but lacks dynamic
iterations through evolutionary epochs. Meanwhile, pMEsim [27] focuses on simulating
polymorphic TE variants for pan-genomes, sacrificing long-term evolutionary dynamics, and
simuG [28] serves as a generic genome simulator that can simulate TE insertions. GARLIC [29]
specializes in generating realistic artificial genomic sequences by modeling intergenic
sequences based on kmer profiles, GC contents, and local repeat landscape, which serve as
high-quality negative controls for evaluating annotation tools. TEgenomeSimulator is a single-
timepoint simulator, useful for benchmarking and creating replicas of the TE landscapes of real
genomes, but it does not perform forward simulation for complex evolutionary scenarios.
While existing simulation tools collectively address diverse needs, few allow flexibility in
genome customization, TE sequence mutagenesis, and dynamic modulation of
insertion/deletion rates across evolutionary epochs; none provides granular family-specific
controls with iterative simulations for hypothesis testing; and most prioritize synthetic data
generation over empirical inference of TE dynamic parameters.

104   Results

105   Introduction of PrinTE

106   Here, we present PrinTE, a transposon-focused tool that allows dynamic forward simulation of
107   TE insertions, deletions, truncations, and sequence substitutions. PrinTE enables iterative
108   forward simulations to explore complex genome evolutionary scenarios, including the effects of
109   TE dynamics on genome size evolution, genomic divergence, the impact of selection on TE
110   turnover, and speciation (**Fig. 1**).

Fig. 1. Programmatic components and functionality of PrinTE. The high-level computational
workflow of PrinTE is illustrated with rectangles and solid lines. See the main text for detailed
descriptions of functionality.

Forward simulations of TE insertions, deletions, and genome divergence

PrinTE is a forward-time genome simulation tool designed to model the dynamics of TEs within
eukaryotic genomes across evolutionary timescales and was developed to incorporate as many
biologically relevant parameters into the model as possible. It uses a modular approach,
beginning with a synthetic genome (Synthetic Genome Mode) or a user-provided (Custom
Genome Mode) that represents a present-day or theoretical ancestral genome. Users may
generate these inputs entirely from scratch with PrinTE's built-in parameterization tools, import
prior PrinTE runs for continued simulation, or use TEgenomeSimulator [33] outputs, which are
supported by PrinTE. PrinTE implements two methods for simulation, which are the variable
rate method and the fixed rate method. The variable-rate method aims to incorporate
consequences from some of the biological and selective forces that guide TE evolution. We
established several rules in this method: 1) The host genome survival and TE proliferation are
conflicting forces pitted against each other; 2) TE superfamilies compete with each other since
they occupy the same limited genomic space; 3) The rate of TE insertion is positively correlated
with the number of intact TEs in the previous iteration; and 4) TE deletions are guided by the
number of TEs that are deleterious to the host's survival. For this purpose, the simulated
genome consists of three components, all of which are receptive to new TE insertions, namely
existing TEs, genic space, and intergenic space. If a TE inserts into another TE or intergenic
space, there is no cost to the host and no impact on the TE deletion rate. In contrast, TE
insertions into genic space are costly to the host, resulting in elevated TE deletion rates in the
next generation. Thus, the outcome of the variable-rate method is less predictable, which may
lead to generations of genome size increase followed by subsequent generations of decrease. It
is useful for those who wish to explore the regulation of TE activities through time and to
understand the forces that affect TEs' bursts and purges.

The fixed-rate method is designed to generate more controlled and predictable outputs.
In this method, per-generation TE insertion and deletion rates are predefined by the user,
allowing the modeling of known evolutionary trajectories and the generation of results that more
closely match expectations (**Fig. S1**). In practice, forward simulation should begin from a
simulated genome that mirrors a present-day version and forecast a few million generations of

TE evolution into the future. Alternatively, it can start from a genome as we expect it may have
looked a few million generations ago and simulate it to the present day. This approach allows
users to test hypotheses relating to TE bloat and purge and to potentially superimpose TE
evolutionary narratives onto the species of interest.
PrinTE can simulate as briefly as a single generation, which can generate subtle
changes to the simulated genome. To allow for simulating millions of generations, PrinTE
employs a stepwise approach in which it can simulate TE evolution of a number of generations
in steps until it reaches the total number of generations specified by the user. To approximate
TE mutation in multiple generations, PrinTE uses a normalized exponential decay distribution to
guide DNA substitutions, mimicking the natural turnover of TE sequences that would occur
during iterative steps.
PrinTE represents a state-of-the-art simulation of TE dynamics in eukaryotic genomes,
enabling the modeling of TE contributions to genome evolution and the estimation of past and
future genome evolution trajectories.

Performance and benchmarking

The 5' and 3' LTR regions of an intact LTR-RT are identical upon transposition. Under
the molecular clock hypothesis, mutations to the LTRs reflect the amount of time since
transposition [30,31]. Accurately estimating the sequence divergence of LTR-RTs is critical for the
inference of TE activity over evolutionary epochs. We developed a k-mer-based LTR-RT dating
approach, called Kmer2LTR (https://github.com/cwb14/Kmer2LTR.git), to evaluate PrinTE's
accuracy in LTR-RT modeling and its ability to forward-evolve a genome. PrinTE allows users to
customize the transition/transversion ratio (Ti/Tv), which enables evaluating various nucleotide
substitution models that were incorporated into PrinTE. For comparisons, we used SLiM to
simulate genome divergence based on a simple neutral model. Single-nucleotide
polymorphisms (SNPs) were projected onto the SLiM-simulated genealogies to obtain whole-
genome sequences that were comparable to PrinTE simulations. Divergence of SLiM-simulated
genomes was estimated based on mutations detected using Minimap2 whole-genome
alignments.
In all comparisons, the Kimura two-parameter (K2P) model outperformed p-distance
estimates, regardless of the Ti/Tv ratio (**Fig. 2A**). The one-parameter Jukes-Cantor 69 (JC69)
model performed equally as well as K2P when Ti/Tv=1, but underperformed when Ti/Tv=10
(**Fig. 2A**). PrinTE and SLiM/Minimap2 each show high accuracy in their ability to detect
mutations up to 25% sequence divergence, but the latter quickly dropped out at divergences
exceeding 25%. PrinTE's Kmer2LTR approach benefits from pre-determined LTR boundaries,
which allow it to force a global sequence alignment onto the otherwise highly divergent LTRs
and accurately detect mutation percentages up to 30%. Divergences at >30% are detectable by
PrinTE's LTR-RT approach with a slight reduction in accuracy (**Fig. 2A**). Simulations at higher
generation counts lead to a wider range of sequence divergence in LTR-RTs, but the overall
divergence estimate remains centered on the expected values even at the relatively high
divergence range (**Fig. 2B**). These results demonstrate a robust LTR-RT divergence estimation
by PrinTE's Kmer2LTR method, especially when using the K2P model.

Fig. 2. Benchmarking PrinTE. (**A**) Estimation of LTR retrotransposon divergence simulated by PrinTE with p-distance (p-dist), Jukes-Cantor 69 (JC69), and Kimura two-parameter (K2P) DNA substitution models and transition/transversion ratios (TiTv) of 1 and 10. Single-nucleotide polymorphisms were projected onto SLiM-simulated genealogies and detected using Minimap2 whole-genome alignment. (**B**) Density distribution of LTR sequence divergence simulated under various divergence scenarios and estimated by Kmer2LTR. Vertical grey dashed lines show the cumulative sequence divergence of all LTR-RTs in the simulated genome. Vertical red dashed lines show the expected sequence divergence at each time point. (**C**) LTR-RT recall rates by LTR_retriever using three TE simulators. TSD, target site duplication. (**D**) Performance metrics of LTR-RT identification by RepeatMasker using the PrinTE library and the simulated genome as input. Error bars show 95% confidence intervals (n = 1,000). (**E**) PrinTE runtime and memory utilization using different numbers of CPU threads. Genome size variation with deletion rates (**F**), deletion length preference parameter k (**G**), and solo-LTR frequencies (**H**).

Next, we assessed the simulation fidelity in PrinTE and existing TE simulators, SimulaTE and ReplicaTE. The success of structure-based TE annotation depends on the presence of

known sequence structures and motifs. We reasoned that intact TEs in simulated genomes should be detectable using a structure-based annotation approach. However, existing TE simulation tools often fail to generate target site duplications (TSD), a key signature of LTR-RT integration [30]. Our results show that LTR-RTs simulated by PrinTE have a significantly higher detection rate than those that were simulated using ReplicaTE and SimulaTE (77% vs 15% and 8%, respectively), suggesting that PrinTE was better at reflecting the biological features that compose a TE (**Fig. 2C**). We used RepeatMasker with PrinTE-derived TE library to evaluate detection accuracy of PrinTE simulations and observed very strong quality metrics (F1 = 0.978**),** suggesting that the output TE library and corresponding genome were accurately simulated (**Fig. 2D**).

To evaluate the computational performance of PrinTE, we simulated one million generations of TE evolution into a 100Mb genome using a step size of 100,000 generations. The initial genome was seeded with 6,000 TEs, and each subsequent iterative step resulted in insertion and deletion of ~500 TEs. PrinTE demonstrated efficient memory usage with peak memory less than 2GB. PrinTE runtimes scale well at low thread counts and begin to saturate with increased thread allocation (**Fig. 2E**). These performance metrics demonstrate that PrinTE is highly efficient, optimized for large-scale analysis, and capable of running on personal computers or compute servers.

Genome Size Evolution Through TE Expansion and Contraction

LTR-RTs are the primary contributing TEs to genome size in many plant, fungi, and animal genomes [32]. To isolate the effect of TEs on genome size across temporal gradients, we focused on modeling LTR-RT evolution using PrinTE's fixed-rate method to simulate predictable patterns of TE expansion and contraction. Here, we simulated two dynamic scenarios subsequently on the same genome. The first scenario used a TE deletion rate fourfold higher than that of insertions. We observed a 45% decrease in genome size after five million generations of forward simulation (**Fig. 3A**). The decrease was initially driven by a swift reduction in the number of intact TEs (**Fig. S2A**) followed by an increase in fragmented TEs and solo LTRs (**Fig. 3A**, **Fig. S2B**).

Following the genome size decrease, we continued the simulation but reversed the TE insertion rate to fourfold higher than that of deletions. The elevated insertion rate caused rapid genome bloating (**Fig. 3A**) powered by an increase in the numbers of both intact and fragmented TEs (**Fig. S2C,D**). The sequence lengths of solo, fragmented, and intact LTR-RTs shifted in response to purging pressures from the host (**Fig. S3**). Interestingly, because k>0 leads to a higher probability for deleting longer LTR-RTs, the length of intact LTR-RTs was overall reduced in the genome size decrease phase and slowly increased in the genome size increase phase when purging pressure was abated (**Fig. S3**).

During simulations of genome size decrease, when the rate of TE deletions exceeded the rate of insertion, we observed unbiased removal of LTR families through generations. Accumulation of DNA substitutions and reduced LTR-RT insertions led to elevated sequence divergence of the remaining pool of intact LTR-RTs and a shift in the age distribution across time (**Fig. 3B**). There were two distinct density peaks as simulation proceeds, even though the rate of TE insertions and deletions remains constant (**Fig. 3B**). The smaller peak represents younger LTR-RTs that were inserted recently and have had less time to be purged, while the

larger peak that shifts to the right and decreases in density through time was retained from the
standing body of LTR-RTs in the ancestral genome. In cases where the rate of TE insertions
exceeds deletions, young LTR-RTs quickly proliferate in high abundance and dominate the
density distribution of LTR-RTs in the simulated genome, masking the signal of remaining older
LTR-RTs (**Fig. 3C**). We conclude that the divergence distribution of intact LTR-RTs reflects their
insertion and deletion dynamics through time.
When species become reproductively isolated from each other, their TEs are also
isolated and evolve independently. Genomes share TEs that derive from their common ancestor
and also contain novel insertions that are unique to the other. With PrinTE, we simulated a
speciation event by evolving a genome for a period of time, then diverted the genome into two
isolated evolutionary paths with shifted TE dynamics to create divergent trajectories of genome
expansion or contraction (**Fig. 3D**,**E**). In the contracting genome, there are relatively few unique
TEs created after speciation (**Fig. 3D**; **Fig. S4**). In contrast, the expanding genome had a higher
abundance of young and unique TEs (**Fig. 3E**; **Fig. S4**). In the fixed-rate method, the rate of TE
insertion and deletion remains relatively stable, only deviating by the stochastic variability
introduced by sampling a Poisson distribution, while the number of insertions and deletions
introduced scales with the size of the genome (**Fig. 3F**).
Further, we profiled three deletion-related parameters in PrinTE to contextualize their
impact to genome size, namely deletion rate, TE length bias, and solo LTR rate (**Fig. 2F-H**).
Deletion rate controls the number of TEs that are processed by PrinTE's deletion engine.
Expectedly, we observe decreasing genome size with increasing deletion rates (**Fig. 2F**).
Deletion of TEs is mainly achieved by unequal or illegitimate recombination, which both operate
through sequence homology between TE sequences. In PrinTE, we allow users to set TE
deletion bias exponentially to the TE length using the `-k` parameter, which can approximate the
disproportionate impact of illegitimate recombination on purging longer TEs. Larger k has a
stronger bias in deleting longer TEs and thus has more constraints on genome size (**Fig. 2G**,
Fig. S5). When k is set to 0, TE deletion has no bias to TE length, allowing genome size growth
through the accumulation of long TEs (**Fig. 2G**, **Fig. S5**).
Illegitimate recombination between LTR regions of individual LTR-RTs will partially
remove the element and create solo LTRs as remnants. Given a fixed deletion rate, PrinTE can
model the proportion of LTR deletions that go through this process by implementing the "`--`
`solo-rate`" parameter. Increased solo rate can increase the number of solo LTRs in the
genome and slow down the purging of LTR-RTs, resulting in a larger genome size (**Fig. 2H**, **Fig.**
S6, **Fig. S7**). By adjusting parameters of deletion length bias and LTR solo rate that interplay
with TE insertion and deletion rates, we can create complex scenarios for modeling genome
size evolution in different systems.
Recognizing that the rates of TE insertion and deletion can be dynamic through time and
may be modified by varying biological processes, we implemented the variable-rate method in
PrinTE, which can be used to explore specific molecular features that constrain TE evolution
and its impact on genome size. Here, rather than setting fixed TE insertion and deletion rates,
we start with an initial rate and allow the tool to dynamically determine subsequent insertion and
deletion rates based on the genetic makeup of the most recently simulated genome (**Fig. 3J**).
We simulated the genome size and TE evolution of the present-day *Arabidopsis thaliana* to
forecast one million generations of TE evolution using the variable-rate method in PrinTE. With

the tested conditions, there is an initial period of genome bloating followed by a longer period of
TE purging and genome size contraction (**Fig. 3G**). Expectedly, genome size increased through
the accumulation of both intact and fragmented TEs and decreased due to the reduction of
intact TEs (**Fig. 3H**). The divergence of intact LTR-RTs reflects the genome size expansion and
contraction (**Fig. 3I**). These features allow us to simulate TE dynamics close to what is observed
in eukaryotic genomes.

Fig. 3. Modeling TEs' contribution to genome size evolution using fixed (A-F) and
variable (G-J) rate methods. (**A-F**) Ten million generations of LTR-RT evolution explores their
long-term impact on genome size using the fixed rate method. For demonstration, we set a rate
of TE deletion that exceeded insertion by fourfold in the first five million generations, then
reversed the rates by setting the insertion rate exceeding deletion by fourfold over the next five
million generations. (**A**) A syntenic ribbon plot illustrates the change in chromosome size. Pie
charts corresponding to each time point show the relative contribution of each feature to
genome size. (**B,C**) Density distributions of LTR-RT divergence estimated using the K2P model
during the first five million generations (**B;** genome size decrease) and the second five million
generations (**C;** genome size increase). Five million generations is the transition between
genome size decrease and increase, which is shown in both plots as dashed red lines.
Phylogenetic trees show the evolutionary relationship of LTR-RTs from the *Tekay* family after
four million generations of genome size decrease (**D**) or four million generations of genome size
increase (**E**). (**F**) LTR-RT insertion and deletion counts (solid lines) and rates (dashed lines)

across 10 million generations. (**G,H**) A simulated genome replica of *Arabidopsis thaliana* is
forward-evolved one million generations using PrinTE's variable rates method. (**H**) Intact (top)
and fragmented (bottom) TE counts per superfamily across one million generations. (**I**) The
distribution of LTR-RT divergence estimated using the K2P model. (**J**) TE insertion and deletion
counts (solid lines) and rates (dashed lines) illustrate the self-adjusting functionality of PrinTE's
variable rate method.

Genome Size Evolution of Rust Fungi

We further studied the impact of TE dynamics on genome size evolution using rust fungal
genomes due to their wide range of genome sizes. The *Pucciniomycotina* is a subdivision of
fungi that contains 10 classes and over 8400 species [35]. We downloaded 58 *Pucciniomycotina*
fungal genomes from NCBI and JGI MycoCosm [34] to investigate genome expansion and
contraction (**Table S1**). We identified and removed alternative haplotypes from these genomes
using Redundans [36]. The resulting haploid genome size varies from 13.6 Mb (*Mixia osmundae*)
to 1018.1 Mb (*Austropuccinia psidii*), representing over 70-fold differences (**Table S2**). We
performed Benchmarking Universal Single-Copy Orthologs (BUSCOs) evaluations using
compleasm [37] and the odb12 *Basidiomycota* database. The complete BUSCOs (single and
duplicated) averaged 90.7% (SD = 5.4) and contig N50 averaged 846 kbp (ranging from 111
kbp to 3.8 Mb), suggesting high assembly quality of haploid genomes (**Table S2**).

Fig. 4. (**A**) The phylogeny of *Pucciniomycotina* fungal genomes used in this study. Protein
sequences of shared single-copy BUSCO were used for tree inference with 1000 bootstrap
replications. Bootstrap support percentage is shown on each branch. The two classes,
Pucciniomycotina and *Microbotryomycetes,* are labeled beside the phylogeny. The genome size

of each species is indicated by color dots. (**B**) Principal Component Analysis (PCA) on LTR
metrics in fungal genomes. Each dot represents one genome. PCA loadings of each metric are
shown. Genomes selected for further studies are labeled. (**C**) Pearson's correlation (*r*) between
genome size and LTR sequence content (LTR_percent), intact LTR-RT count (Intact.count),
solo LTR count (Solo.count), and solo:intact ratio. Each dot represents one genome. The
correlation and p-value are shown on the panel title. (**D**) Divergence of intact LTR-RTs in
Cronartium ribicola (white pine blister rust), *Puccinia coronata* f. sp. *avenae* (oat crown rust),
and *Austropuccinia psidii* (myrtle rust) genomes. The top five most abundant LTR families in
each genome are colored, and they are not shared between genomes. A density line is shown
as the contour in each plot. The locations of these species on the phylogeny are indicated by
dotted lines.
346
We identified the shared single-copy BUSCOs and used IQ-TREE [38] to construct the
phylogeny of *Pucciniomycotina*. The resulting phylogeny was over-represented by two classes,
Microbotryomycetes and *Pucciniomycetes* (**Fig. 4A**). *Microbotryomycetes* genomes are small,
ranging from 17.5 Mb to 50.3 Mb. Correspondingly, on average, only 0.4% of the genome was
represented by LTR sequences with a range of 0% to 3.0% (**Table S2**). This observation
suggests that the best strategy for controlling genome size is to avoid the invasion of LTR-RTs.
Pucciniomycetes have widely distributed genome sizes, ranging from 26.8 Mb to 1018.1 Mb,
with an LTR sequence content corresponding to 0.4% to 67.4% (**Table S2**).
To further study the dynamics of LTR-RTs in these fungal genomes, we *de novo*
identified intact LTR elements and solo LTR sequences, and calculated the whole-genome
solo:intact ratio using LTR_retriever [30]. Using genomes with >1% LTR sequence content (n =
22), we performed Principal Component Analysis (PCA) on LTR metrics and genome size. The
analysis revealed clustering among smaller genomes, whereas larger genomes appeared as
outliers, including *Cronartium ribicola* (white pine blister rust), *Puccinia coronata* f. sp. *avenae*
(oat crown rust), and *Austropuccinia psidii* (myrtle rust; **Fig. 4B**). PCA loadings suggested that
solo LTR number, intact LTR number, and their ratio were all drivers to the outliers of large
genomes (**Fig. 4B**). Including total LTR content, these LTR metrics showed strong positive
correlations with genome size (Pearson's $r \geq 0.65$, $p \leq 0.001$; **Fig. 4C**). This observation
supports LTR-RTs' major role in impacting genome size upon successful invasion.
Not surprisingly, intact LTR elements, as the product of LTR-RT proliferation, were
strongly correlated with genome size ($r = 0.76$; **Fig. 4C**). However, solo LTR sequences as the
product of LTR-RT removal [30] showed an even stronger correlation, almost linearly correlated
with genome size ($r = 0.98$; **Fig. 4C**). Another counter-intuitive observation is the positive
correlation between genome size and solo:intact LTR ratio, the latter is a metric used to
measure the intensity of LTR-RT purging [31]. This observation contradicts the conventional
understanding that the higher the solo:intact LTR ratio, the more intense purging of LTR
sequences, and thus the smaller the genome [39]. The positive correlation between solo:intact
ratio and genome size suggests that illegitimate recombination alone may not be an effective
way of escaping genome obesity.
We selected *C. ribicola*, *P. coronata* f. sp. *avenae*, and *A. psidii* as representatives of
three distinct roadmaps to genome size evolution. *C. ribicola* has a genome size of 211.6 Mb
with the LTR divergence distribution showing an extremely quick turnaround time for LTR

element insertions (**Fig. 4D**; **Fig. S8**). The solo:intact ratio of *C. ribicola* is 27.4 (**Table S2**), the
highest among all genomes analyzed in this study, suggesting a quick insertion–quick removal
mode of action, and a relatively stable or slowly increasing genome size. The *P. coronata* f. sp.
avenae has a genome size of 113.2 Mb and a solo:intact ratio of 1.7 (**Table S2**). The LTR
divergence distribution showed a much slower turnover time for LTR-RT insertions (**Fig. 4D**;
Fig. S8), suggesting a slow insertion–slow removal mode of action, and a relatively stable
genome size. *A. psidii* has an extremely large genome size of 1018.1 Mb and a solo:intact ratio
of 23.5, the second highest ratio among our data (**Table S2**). However, the LTR divergence of
A. psidii shows the peak of LTR-RT insertion passed the current time (**Fig. 4D**; **Fig. S8**),
suggesting a slowdown of insertion activities [8] and a decreasing genome size.

Simulating Digital Replicas of Rust Fungal Genomes

TE composition in real genomes varies substantially across TE families and species
because of their complex and diverse evolutionary histories. This variability makes it challenging
to simulate genome-wide TE landscapes that can be used for modeling and hypothesis testing.
To address this, we used TEgenomeSimulator [33] with its TE Composition Approximation Mode
to model the TE profiles of *C. ribicola*, *P. coronata* f. sp. *avenae*, and *A. psidii* genomes within
the *Pucciniomycotina* subdivision (**Fig. 5**, **Fig. S9**, **Fig. S10**). The sizes of these simulated
genomes closely matched their actual genome sizes. Similarly, the TE proportion of the
simulated genomes closely approximated the observed values in terms of TE loci and TE bases
per superfamily (**Fig. 5AB**, **Fig. S9**, **Fig. S10**). TEgenomeSimulator estimated the sequence
divergence and integrity distributions of each TE family in the real genomes for the simulation of
digital replicas. The simulator uses normal distributions to smooth out small-scale fluctuations
present in the real genome, resulting in a simulated divergence profile that broadly resembled
the real data (**Fig. 5C**, **Fig. S9**, **Fig. S10**).
TEs can break down over time due to nested insertions, recombination, small deletions,
and other processes. The whole-genome distribution of TE integrity clearly reflects the turnover
rate of TEs in the three selected fungal genomes (**Fig. 5D**, **Fig. S9D**, **Fig. S10D**). The *C. ribicola*
genome contains mostly high-integrity TEs, suggesting a high rate of TE turnover and active TE
insertions (**Fig. S9D**). In contrast, the *P. coronata* f. sp. *avenae* genome harbors highly
fragmented TEs and very few intact TEs, suggesting a low activity profile (**Fig. S10D**). The
myrtle rust genome, *A. psidii*, presents a bimodal distribution in TE integrity, suggesting both
heavy bloating and purging may have been prominent drivers in the evolutionary history of the
genome (**Fig. 5**). The digital replicas of these genomes have distributions of TE integrity that are
almost identical to the real genomes (**Fig. 5D**, **Fig. S9**, **Fig. S10**). These simulations yield
generalized distribution patterns that capture the major trends in TE sequence evolution of each
genome, providing a robust foundation for PrinTE's simulations of TE bloating or purging
episodes, effectively serving as burn-in phases for modelling genome evolution.

416
417 **Fig. 5**. **TE Composition Approximation for the *Austropuccinia psidii* genome.** (**A**)
418 Comparison of total TE loci between the original and simulated genomes. (**B**) Comparison of
419 total TE bases between original and simulated genomes. (**C**) TE sequence diversity distribution
420 colored by TE superfamily. (**D**) The distribution of TE sequence integrity, which is the length
421 ratio relative to the consensus sequence, is colored by TE superfamily as in (**C**).

Accessing Genome Evolution with Forward Simulations of LTR Activities

Our simulations using PrinTE present an effective capturing of LTR-RT dynamics using the
divergence distribution (**Fig. 2A-C**), which, in reverse, can be used to empirically estimate
parameters of LTR-RT dynamics. Here, we demonstrate the utility of PrinTE in modeling
genome size evolution under the impact of TE bloat and purge. Despite their shared ancestry
(**Fig. 4A**), the three selected fungal species (*C. ribicola*, *P. coronata* f. sp. *avenae*, and *A. psidii*)
show immense variation in genome size and LTR-RT evolution. To test our hypothesis that *C.
ribicola* has a stable genome size that it achieved by rapid but balanced LTR-RT insertion and
deletion, we forward-evolved the *C. ribicola* genome digital replica using PrinTE with a variety of
insertion and deletion rates in search of an LTR-RT divergence distribution that closely matches
the observed distribution. At each trial, we compared the LTR-RT divergence distribution with
that of the real genome (**Fig. S11**). We found that an LTR-RT insertion rate of 2×10^{-9} per bp per
generation and a deletion rate of 2.4×10^{-9} per bp per generation provided a simulated genome
that closely matched the *C. ribicola* genome, with the genome size increased moderately
through time (**Fig. 6A,J**). Forward-projecting 200,000 generations of TE evolution indicated that
the density distribution of LTR-RT ages in the simulation closely matched that of the real
genome (**Fig. 6A**, **Fig. S9**). The solo:intact ratio of the simulated genome was 13 (**Fig. 6C**) in
comparison to 27.4 in the real genome (**Table S2**). Although the forward-simulated genome

does not share an identical LTR-RT footprint to that of *C. ribicola*, its proximity to the real data
suggests that PrinTE can effectively infer TE dynamics through forward simulations. Simulation
of the *C. ribicola* genome suggests that the species might have inserted and deleted (either
intact or fragmented) one effective LTR-RT every two generations during the last 200,000
generations.
We repeated the same process using the genomic digital replica of *P. coronata* f. sp.
*avenae*a to test our hypothesis that the species has slow TE insertion and slow removal, leading
to a relatively stable genome size. An LTR-RT insertion rate of 8×10^{-12} per bp per generation
and a deletion rate of 1.10×10^{-11} per bp per generation were found to be optimally adapted for
maintaining the LTR-RT characteristics of the real genome for over 20 million generations (**Fig.**
6B,J; **Fig. S12**). Dating LTR-RTs in the forward-simulated and the real genome showed that the
distributions were nearly identical (**Fig. 6D**). The solo:intact ratio in the forward-simulated
genome mimics that of the real genome, at 2.0 and 1.7, respectively (**Fig. 6F**, **Table S2**). With
these rates, we see a gradual increase in average genome size, adding approximately 1.25 Mb
per million generations (**Fig. 6E**). These results suggest that, in contrast to *C. ribicola, P.*
coronata f. sp. *avenae*a appears to have an almost inactive LTR-RT footprint with an average of
1090 generations between each effective LTR-RT insertion and an effective deletion every 790
generations.
For our final simulation, we used PrinTE to create an initial burn-in genome that
mimicked the gene and TE landscape of a hypothetical ancestral genome of *A. psidii*,
highlighting PrinTE's utility as a stand-alone TE simulation and evolution tool. We began with a
relatively small genome of 110Mb that is typical in the *Pucciniomycetes,* and simulated the TE
evolution that led to the enormous genome bloating of *A. psidii*. Based on the genome size and
LTR-RT divergence distribution (**Fig. 4A**,**D**), we expected that TE insertions exceeded deletions
for a long period of time, leading to genome size increase, followed by a shorter period of rapid
LTR-RT deletions. Through evaluation of various dynamic parameters, we found that an
insertion rate of 1.4×10^{-11} per bp per generation and a deletion rate of 5×10^{-13} per bp per
generation for 24 million generations was optimally tuned to mimic the observed LTR-RT age
distribution of the *A. psidii* genome during the long period of genome size increase (**Fig. 6G**,**J**;
Fig. S13A,**B**). During this period, we aimed to achieve a genome size that was marginally larger
than the real genome so that we could tune up the rate of deletion to reflect the more recent
decline in young LTR-RTs observed in *A. psidii* (**Fig. 4D**). At peak LTR-RT invasion, the
simulated genome was 1,178 Mb in size (**Fig. 6H**).
The drop of recent LTR-RT insertions in *A. psidii* suggests a reduced insertion rate and
an increased deletion rate. Using a substitution rate of 2.0×10^{-9} per bp per generation [40], we
estimated that the insertion reduction phase of *A. psidii* had lasted for 1.2 million generations.
We then forward-simulated an additional 1.2 million generations using elevated deletion rates
on top of the increasing phase of simulation. Systematic evaluation of TE dynamic parameters
revealed that an insertion rate of 1.4×10^{-13} per bp per generation and a deletion rate of 3×10^{-11}
per bp per generation can achieve a genome size that is near *A. psidii*, at 1,027 Mb (**Fig. 6H**,**J**;
Fig. S13C), and the LTR-RT divergence distribution mimics the real genome (**Fig. 6G**; **Fig.**
S13D). The solo:intact ratio of the simulated *A. psidii* genome terminated at 6.8 (**Fig. 6I**), which
is lower than the observed value at 23.5 (**Table S2**), suggesting other evolutionary processes
also play roles in shaping the modern-day *A. psidii* genome.

Fig. 6. Simulating genome size evolution in three *Pucciniomycetes* genomes. (**A-C**)
Cronartium ribicola (white pine blister rust), (**D-F**) *Puccinia coronata* f. sp. *avenae* (oat crown
rust), and (**G-I**) *Austropuccinia psidii* (myrtle rust). (**A, D, G**) Sequence divergence compares the
empirical divergence distribution of LTR-RTs (Reference) to simulated genomes. LTR
divergence was estimated using LTR_retriever with the K2P model. Three timepoints were
plotted for *C. ribicola* (**A**) and *P. coronata* f. sp. *avenae* (**D**) to showcase the relatively stable TE
dynamic. The three timepoints of *A. psidii* (**G**) highlight the rapid purging of LTR-RTs during the
period of genome size contraction. (**B, E, H**) Genome size changes across generations. (**C, F, I**)
The empirical solo:intact ratio estimated using LTR_retriever. (**J**) The estimated LTR-RT
insertion and deletion rates of fungal genomes. The expansion and contraction phases of the *A.*
psidii simulation were plotted separately. (**K**) Microsynteny between the simulated *P. coronata* f.

sp. *avenae* before (top) and after (bottom) the TE evolution that led to genome size increase in
the simulated *A. psidii* genome. Collinear genes are blue, and LTR-RTs are orange.
499
Simulation of *A. psidii* suggests two major phases that contributed to the present-day
genome size. The first was a long period of genome size increase in which LTR-RT insertions
exceeded deletions by 28-fold, and the second was a short period where deletions exceeded
insertions by 214-fold. Our results support a recent contraction model for genome size evolution
in *A. psidii* (**Fig. 6J**). It is important to note that the estimated parameters are effective insertion
and deletion rates averaged from a long period of evolutionary time, which may not capture
short periods of TE bloat and purge.
Because these three species were selected to represent the extremes of LTR-RT
dynamics in the sampled *Pucciniomycetes* genomes, we suggest that the amplification and
degradation of LTR-RT can vary widely within the fungal kingdom. In future studies, it will be
interesting to see if such a wide variety of LTR-RT dynamics exists at narrower phylogenetic
levels.

Discussion

The TE composition profile of a genome serves as an evolutionary archive, preserving
signatures of TE dynamics and host genome responses that collectively shape genome
evolution. Deciphering these patterns from empirical data is challenging. Whether to test
evolutionary hypotheses or to predict future trajectories, simulation is often necessary. To
address this, we built PrinTE, a flexible, forward simulation framework for transposon evolution.
PrinTE significantly expanded the capacity for simulating complex TE dynamics and testing
evolutionary hypotheses.

PrinTE allows forward simulation on TE dynamics

Transposable elements are a major genomic component and subject to all evolutionary forces,
including purifying selection, drift, and mutation. The ability to trace TEs' responses to these
events will greatly facilitate the studies of TE dynamics. Forward simulation is a key approach
that allows the artificial evolution of TEs and other genomic features, which provides
opportunities to sample the intermediate state of the evolutionary process.
SLiM is a powerful forward simulation program that can simulate the majority of
evolutionary scenarios [24]. However, its operation on single-nucleotide variants limits its
application on TEs with varying lengths and abundance of nestings. SLiM provides a
workaround by first compressing the whole TE locus into a presence and absence unit and
performing simulations on these units. Sequence content is then filled back to replace the units
to generate genomic sequences. This approach makes it difficult to model the variation
accumulated through generations on the simulated TE sequences, including the deletion and
formation of solo LTRs, which are critical for genome evolution studies. On the contrary, PrinTE
operates on whole-genome sequences, including invariant sites that allow intuitive modeling of
complex scenarios and interpretation of results. In particular, PrinTE models the insertion and

deletion rates of TEs and generates solo LTRs, which empowers the study of genome size
evolution driven by TE bloat and purge.
The input files of PrinTE are highly flexible (**Fig. 1**). It has a built-in engine to simulate
highly customizable starting genomes, in which users can specify genome size, TE content, TE
superfamily abundance, and TE divergence in their simulations. PrinTE integrates CDS
sequences in the simulation, allowing for modeling purifying selections, tracing large structural
variants, and studying the evolution of gene space. PrinTE also takes simulated genomes from
TEgenomeSimulator, allowing forward simulation from different diverse backgrounds of
genomes. Most importantly, PrinTE can take its own output as input (e.g., **Fig. 3A-F**), allowing
for the simulation of varying TE dynamics and studying the host-TE arms race. This feature also
enables the simulation of genomic divergence and speciation driven by differences in TE
activities.

TE contributes to genome size expansion

The contribution of TE contents to genome size expansion has been confirmed across many
taxa and in our data (**Fig. 4C**). Over time, genomic data have come faster and of a higher
quality, and yet, as noted by Osmanski *et al.* (2023) [41], the role of TEs in genome architecture
and evolution is often an afterthought. More in-depth analyses are restricted to domain
specialists for a meaningful evolutionary perspective.
In order to put forward mechanistic hypotheses, authors have developed specialized
computational methods to detect signals of TE radiation and impacts on speciation and genome
size. For example, Ricci *et al.* (2018) [13] developed a measure of TE Density of Insertion and
Relative Rate of Speciation to model TE radiation bursts and silencing. Kapusta *et al.* (2017) [12]
estimated TE-driven DNA gains and losses and proposed the "accordion" model of TE radiation
followed by DNA loss for tight genome size maintenance in mammals and birds. Osmanski *et al.*
(2023) [41] applied a Redundancy Analysis to study TE composition across mammalian taxa
generated through the Zoonomia genomes [42], and found evidence of multiple recent TE
expansions driving larger genome sizes and wide-spreading TE-silencing events across the
mammalian tree. These authors hypothesize that variations in TE defense mechanisms may
have played a role in whether specific lineages were overcome by a TE burst at a given
evolutionary time.
However, we have limited knowledge about the maintenance of compact genomes
despite evidence of TE radiation bursts and the maintenance and regulation of extremely large
genomes. The Chinese pine attributes 69.4% of its 25.4 Gb genome to TEs, 86.5% of which are
made up of LTR-RTs [16]. The Chinese pine genome showed no sign of recent whole-genome
duplication, but 91.2% of its genes were undergoing discrete duplication or family expansion,
strongly associated with TEs. Our results revealed two different genome dynamics in the
Pucciniomycotina subdivision of fungi, including the *Microbotryomycetes* class that harbours
compact genome size (18-50 Mb) and the *Pucciniomycetes* class with drastic differences in
genome size (27-1018 Mb) (**Fig. 4A**; **Table S2**). The *Microbotryomycetes* class is almost devoid
of LTR sequences (**Table S2**), suggesting that the most effective way of maintaining compact
genome sizes is to be immune to the invasion of LTR-RTs. Genomes in the *Pucciniomycetes*
class have varying richness of LTR-RTs (**Table S2**), and the PCA reveals several types of LTR
dynamics (**Fig. 4B**) that are likely major drivers of genome size variation.

The myrtle rust fungus expanded its genome size from an approximate ancestral size of
110 Mb to the current 1,018 Mb, experiencing a ~10× increase. Our simulation studies revealed
that the genome is experiencing a downsizing effect, with LTR-RT deletions being >200× faster
than insertions (**Fig. 6**). Using PrinTE's built-in genome simulation utility, we were able to
simulate a genome that mimicked the TE landscape of the ancestral genome of myrtle rust, and
further used PrinTE to simulate two phases of evolution, including a long expansion period and
a short contraction period. This model fits well with the observed composition of the myrtle rust
genome, suggesting a close approximation of its trajectory of genome size evolution.

In this study, we used TEgenomeSimulator and PrinTE to smooth out small-scale TE
activities, then used PrinTE to simulate the real-time mobilization and deletion behavior of TEs
to capture the most dominant expansion or contraction events. We recognize that TE activity is
tightly linked to host genome regulation, particularly when it impacts gene function and
organismal fitness. It is also important to account for more complex patterns in sequence
identity distributions, which may go beyond simple normal distributions. The combined use of
PrinTE and TEgenomeSimulator greatly enhanced the simulation capacity, enabling richer and
more flexible hypothesis testing across both evolutionary and mechanistic contexts. Future
development would benefit from incorporating features such as customizable identity
distributions, more functional genomic context, epigenetic regulation mechanisms, and
additional insertion site preferences.

598 Conclusions

In conclusion, PrinTE is a versatile tool that allows flexible forward simulation of whole-genome
sequences with fine-grained user control of genome size, TE content, superfamily abundance,
TE divergence profile, TE insertion, TE deletion, TE mutation, solo LTR formation, genome
divergence, and speciation. We demonstrated the use of PrinTE in resolving the genome size
evolution of *Pucciniomycotina* genomes. In particular, we inferred the TE dynamic parameters
underlying the genome size evolution of myrtle rust (*Austropuccinia psidii*) and other large
fungal genomes. Our work transforms the study of genome evolution into a highly traceable and
explorative simulation framework, and allows testing of data-driven hypotheses.

607 **Methods**

608 **Implementation of PrinTE**

609 **Burn-in**

PrinTE simulates the forward evolution of genomes. The starting point for simulation is a burn-in
genome in FASTA format and a corresponding PrinTE-formatted BED file. These will serve as
the starting point (0 generation) for which no forward evolution has yet taken place. The burn-in
FASTA and BED can be generated *de novo* by PrinTE, or previous PrinTE outputs feeding back
into PrinTE for additional forward simulations. PrinTE *de novo* burn-in allows many
customizations, including genome size, chromosome number, percentage of the genome to be
genic and TEs, and the ratio of each TE superfamily in the genome. Each TE insertion creates a

TSD with the length determined by the TE superfamily, which was adopted from the
TEgenomeSimulator codebase. In some instances, it may be beneficial to apply baseline
mutation to TEs in the burn-in such that they do not exactly match elements in the TE library.
This is particularly relevant for self-dating elements, like LTR-RTs, where the default TE library
only contains very young LTR-RTs, with exactly identical LTRs. For baseline mutations in the
burn-in genome, we apply a normalized exponential decay distribution to skew mutations
heavily toward lower mutation values:

$$f(M) = \frac{k}{M_{max} \cdot (1 - e^{-k})} \cdot e^{-k \cdot \frac{M}{M_{max}}}$$

The probability density function is aimed at flexibly and accurately modeling TE age distributions
as they are observed in real data, where aggressive TE purging leads to higher probability of
observing young TEs (**Fig. 4D**). The parameters M_{max} (maximum mutation percentage; '--
TE_mut_Mmax') and k (decay constant; '--TE_mut_k') are customizable, allowing users to
adjust the shape of the distribution to reflect a wide range of evolutionary and biological
scenarios and to match the observed TE half-life of any species (see **Fig. S1A**,**B**). The half-
point of sequence divergence among the simulated TE is:

$$M_{1/2} = \frac{M_{\max} \ln(2)}{k}$$

Two Simulation Modules

Forward simulation begins once the burn-in genome is established. Users can specify the total
number of generations ('--generation_end') and a smaller number of generations for each
iteration ('--step'). There are three sequential modules for forward simulation: 1) genome
mutation, 2) TE insertion, and 3) TE deletion. The pipeline will iterate over these processes for
each step until the target number of generations is simulated. PrinTE can iterate through
modules using constant and **fixed** TE insertion and deletion rates, or it can iterate using self-
adjusting **variable** TE insertion and deletion rates. The fixed and variable methods begin
identically, starting with the mutation of the genome. The expected number of DNA mutations
per sequence (λm) is calculated as:

$$\lambda m = DNA\ mutation\ rate\ \times\ genome\ size\ \times\ generations$$
643

Because TE insertions can occur at any generation of a defined evolutionary window
(*i.e.*, a simulation step), they could accumulate DNA mutations less than λm. To capture this
dynamic, we used a normalized exponential decay function similar to the decay function used
for the burn-in genome to weight TE mutations toward lower values. We anchored the mutation
distribution here to approximate the complicated evolutionary dynamics of TE activities. The
actual number of mutations for each newly simulated TE is sampled from the probability density
function, $f(mTE)$, to help weight their observed mutations toward lower frequencies:

$$f(m_{\text{TE}}) = \frac{1.4 \times 10^{-6} \cdot e^{-1.4 \times 10^{-6} \cdot m_{\text{TE}}}}{1 - e^{-1.4 \times 10^{-6} \cdot 3,000,000}}, \quad 0 \leq m_{\text{TE}} \leq 3,000,000$$

651   The half-life of simulated TE insertions is $t_{1/2}$ = ln(2)/k, where k = 1.4×10^{-6}, thus $t_{1/2}$ ≈ 495,000,

652   capturing the observation of TE half-life reported in multiple eukaryotic organisms [30].

Simulation with Fixed Rates

TE insertion. After DNA mutations, TEs are inserted into the genome at random locations.
They may insert into intergenic regions, into other TEs, resulting in nested TE insertions, or into
genes, resulting in fragmented genes. Information about nesting and fragmentation is recorded
in the BED file. In the fixed rate method, the number of TEs to be inserted (λ_{i-fix}) is calculated
as:

$$\lambda_{i-fix} = TE\ insertion\ rate\ \times\ genome\ size\ \times\ generations$$

TE deletion and solo LTR. Donor TEs are randomly selected from the TE library based on
user-defined abundance ratios per superfamily. A nested insertion will fragment the acceptor TE
into two fragments. For example, if TE1 inserts into TE2, it will generate three sequences for
further operations (e.g., deletion), including two resulting fragments of TE2 and the complete
sequence of TE1. This differs from simple insertions into intergenic regions, where each
insertion results in only one sequence for further operations. During the deletion process, full-
length TEs and TE fragments are all eligible for deletion. The number of TE deletions (λ_{d-fix}) is
calculated as follows:

$$\lambda_{d-fix} = TE\ deletion\ rate\ \times\ genome\ size\ \times\ generations$$

Illegitimate recombination is a prevalent process for TE deletions and is facilitated by
sequence similarity between nearby homologous sequences [31]. Longer sequences have a
higher probability of causing illegitimate recombination than shorter sequences because there
are more similar base pairs to initiate pairing. As a result of this phenomenon, longer TEs have
a higher probability of facilitating illegitimate recombination and a higher likelihood of being
deleted than shorter TEs. We used an exponential decay function to model an exponentially
increasing probability of TE deletion as TE length increases:

$$f(L) = e^{-k \cdot \left(1 - \frac{L}{L_{95}}\right)}$$

Here, L_{95} represents the TE length at the 95th percentile of the length distribution in the genome,
L represents the length of a TE (whether intact or fragmented), and *k* is a user-defined
parameter ('--k') that controls the rate at which deletion probability increases with sequence
length. This modification allows us to favor the deletion of longer TE features by assigning
higher deletion weights to sequences that are closer in length to the longest observed TE.
Except for LTR-RTs, TE deletions span the full length of the TE feature. If a nested TE is
selected for deletion, the adjacent TE fragments it disrupts are reconstituted into an intact TE.
LTR-RTs are an exception to full-length deletions because they can undergo recombination with

themselves owing to the sequence homology between their long terminal repeats, resulting in
the formation of a solo LTR. PrinTE supports deletions that lead to solo LTR formation. If an
LTR-RT is selected for deletion, it will be converted to a solo LTR at a user-specified frequency,
which is set by default to 95% ('`--solo_rate`').

Simulation with Variable Rates

In addition to the fixed-rate TE insertion/deletion method outlined above, in which the number of
insertions and deletions per generation is constant and modeled by user-selected rates, PrinTE
also offers a variable method that self-adjusts the number of insertions and deletions per
generation based on genomic landscapes. This aims to replicate a subset of the biological
forces that affect TE insertions and deletions. The core processes, 1) genome mutation, 2) TE
insertion, and 3) TE deletion, are identical between methods. The genome mutation step is
identical between methods and is described above.

TE insertions. In real genomes, the rate of TE transposition is influenced by the number of
intact TEs, since only intact TEs are capable of initiating new insertions. Genomes with a high
number of intact TEs tend to experience more transpositions per generation than those with
few. Fragmented TEs and solo LTRs are no longer viable for transposition. In the variable
method, the number of TE insertions (λ_{i-var}) is modeled as follows:

$$\lambda_{i-var} = TE\ insertion\ rate \times number\ of\ intact\ TEs \times generations$$
This simulates insertions via transposition, but some TEs will be acquired horizontally and drift
into the genome. We set that the frequency of horizontally acquired transpositions is influenced
by TE prevalence within the species. We set that species with TE-rich genomes are more likely
to acquire additional TEs horizontally, while species with TE-poor genomes are less likely to do
so. For this calculation, the TE composition of the genome is surveyed before simulations and
used as a proxy for TE abundance in the species. horizontally acquired TEs (λ_{i-hor}) are then
simulated accordingly:

$$\lambda_{i-hor} = TE\ horizontal\ rate \times TE\ composition\ of\ the\ species \times generations$$
Insertion bias. The distribution of TEs across chromosomes is uneven and can be influenced
by chromatin states, with euchromatin generally more accessible for TE insertion. In the variable
method, we model this by treating genes and their surrounding regions as euchromatin, and the
gene-poor genomic backbone as heterochromatin. We can modulate insertion location
probability by chromatin type using a bias factor ('`--chromatin_bias_insert`'). We assign
insertion probability weights to euchromatic and heterochromatic sequences as follows:

$$Euchromatin\ weight = length \times bias\ factor; \quad Heterochromatin\ weight = length$$

It is important to note that PrinTE does not use genes as functional units, but rather as
representative segments of the genome to which cellular and selective attributes can be
assigned. This enables the modeling of global genome features. Unlike the fixed-rate method,
where TEs are inserted at random in proportions defined by the user, the variable-rate method
inserts TEs in proportions that reflect the current genomic composition.

Fitness-dependent TE deletions. TEs can trigger extensive and rapid proliferation, resulting in
a sudden increase in their genomic abundance. If left unchecked, this proliferation can
destabilize the genome and impose a selective disadvantage on the host organism. Under the
variable-rate method, the number of TE deletions is dynamically adjusted according to the
extent of gene disruption caused by TE insertions. In this context, genes are modeled as
sequences subject to purifying selection, whereby insertions of TEs into these regions are
deleterious and thus reduce the observed number of TE insertions. The number of TE deletions
(λ_{d-var}) is then determined by the following expression:

$$\lambda_{d\text{-}var} = TE\ deletion\ rate \times \sum (disrupted\ gene\ weights) \times generations$$

730 In actual genomes, not all genes that confer a selective advantage exert the same influence on
731 fitness. While most TE insertions into genes are deleterious, some have more severe effects. To
732 capture this variability, we model the distribution of TE selection pressures across genes using a
733 log-normal distribution (**Fig. S1C**):

$$f(x;\sigma) = \frac{1}{x\sigma\sqrt{2\pi}}\ e^{-\frac{(\ln x - \sigma^2)^2}{2\sigma^2}}, \quad x > 0$$

In this framework, each disrupted gene contributes a positive value to the total disrupted
gene weight. The mode of the distribution is fixed at 1 to gain better control of the spread.
Because of the long tail of the distribution, a small subset of genes exerts a disproportionately
large influence on fitness. Consequently, TE insertions in these regions lead to a substantial
increase in deletion events. The parameter σ, which is set via the command line ('`--sigma`'),
determines the spread of the distribution. Larger values of σ produce longer tails and increase
the range of selective effects among genes. In reality, TE insertions in some genomic regions
can be lethal. PrinTE does not simulate lethality explicitly. Instead, we assume that simulating
many generations using a stepwise approach effectively filters out lethal insertions. Over time,
such deleterious types are eliminated from the population and do not persist in the simulated
genomic landscape.
TE deletion bias. Once the number of deletions is calculated, the specific TEs to delete are
determined by the same exponential growth function ($f(L)$) used in the fixed-rate method
multiplied by additional genomic attributes that might affect the probability of TE deletion. These
genomic attributes include the selection coefficient parameter ('`--sel_coeff`') and the
chromatin bias parameter ('`--chromatin_bias_delete`'). The selection coefficient
parameter is used to assign additional deletion weight toward the TEs that disrupt genes.
Similarly, we can assign additional weight toward deletions in simulated euchromatin regions.
Longer TEs and TEs nearer to or within impacted genes are more likely to be deleted based on
user-specified thresholds.

LTR-RT dating. PrinTE implements a novel and accurate (**Fig. 2A,B**) approach called
Kmer2LTR to estimate sequence divergence between the LTR regions of LTR-RTs. Optionally,
Kmer2LTRuses known LTR region boundaries as input to initiate the LTR-RT dating process.
For empirical data where the location of LTRs is unknown, we developed a k-mer approach to

estimate LTR region boundaries. The k-mer approach first uses Jellyfish [43] to identify and count
8-12 mers. K-mers occurring exactly twice in the LTR-RT are mapped back to the LTR-RT. The
resulting k-mer pairs are sorted numerically by position. Pairs that disrupted the monotonic
order or that deviated by more than two standard deviations from the mean distribution of map
position are removed. The map coordinates of the remaining k-mer pairs are used to establish
the putative LTR region. A 65bp extension is applied to each of the putative kmer-determined
LTR regions to ensure that the LTRs are fully captured. MAFFT [44] is used to align LTR regions
within each LTR-RT, and trimAl [44,45] is used to prune overhanging extensions and to remove
small gaps. This approach accurately identifies the boundaries of LTR regions from young to
very old LTR-RTs. The gap-free and extension-cleaned LTRs are then fed to the wavefront
alignment (WFA) algorithm [44–46] for global sequence alignment with a two-piece gap-affine
scoring model to estimate sequence divergence. With these implementations, Kmer2LTR only
requires a fasta of LTR-RTs and identifies the LTR regions *de novo*. This pipeline is freely
available on GitHub (https://github.com/cwb14/Kmer2LTR.git).

772   Searching for TE Dynamic Parameters using PrinTE

773   To identify optimal TE dynamics in a given genome, we employed a systematic parameter
774   sweep approach using PrinTE. This method involved iteratively testing combinations of key
775   parameters to approximate empirical TE behavior and genome size observed in the genome of
776   interest. The `'pipeline.report'` file contains an updated log of TE insertion and deletion
777   counts per simulated iteration and can also be useful in monitoring PrinTE runs.

778   *Parameter Space Exploration*

We conducted a grid search over two critical parameters, insertion and deletion rates, governing
TE insertion dynamics. Users can specify parameter ranges (upper and lower limits) and
increments of parameters. Based on the hypothesis of genome size dynamics, constraints on
the deletion rate can be applied to narrow down the search space. For example, if a genome
evolves with a stable size, a deletion rate between 1.2 to 1.5 folds of the insertion rate can be
used. We also monitor the size of simulated genomes and abort simulations when over or under
the user-set size to ensure strict resource consumption. For example, if the target genome size
is 150 Mb, simulation parameters that create genomes over 180 Mb in any step are deemed
unfit for the hypothesis and thus aborted to reduce the computation load. PrinTE's built-in '`--`
`max_size`' parameter can be used to set the upper limit of genome size and stop the simulation
when exceeding the specified size.

Additional parameters included length bias for deletions ('`--k`') and solo LTR rates ('`--`
`solo_rate`'). In practice, we found k values of 0, 3, or 6 and solo rates from 70% to 95% (with
5% increments) could provide sufficient dynamics on top of the insertion and deletion rates.
Higher k value will lead to slower genome size increase due to the preference for deleting
longer TEs (**Fig. S5**). Higher solo rate will lead to faster genome size increase due to the
creation of solo LTRs instead of full deletions (**Figs. S6, S7**) and increased solo:intact ratio.

796   *High-Throughput Simulation Framework*

PrinTE simulations were executed through an HPC cluster using array jobs (SLURM workload
manager), enabling parallel processing of parameter sets. Users can set the total number of
generations ('-ge') with sampling steps ('-st') for all simulations during parameter sweeps.

The species-specific substitution rate ('--mutation_rate') is also critical to the
simulation outcome. In cases where an accurate substitution rate is not available for the
species, users can try rates in different magnitudes (e.g., 1×10^{-8}, 1×10^{-9}, or 1×10^{-10} per bp
per generation) in the simulation, then compare the LTR divergence of simulated genomes to
the real genome. Higher mutation rates will introduce more mutations to TEs in a given
timeframe, thus creating gentler slopes of LTR divergence. In reverse, lower mutation rates will
create steeper slopes of LTR divergence.

Empirical Estimation of LTR Divergence and solo:intact Ratio
Outputs were processed through a standardized pipeline to assess TE dynamics, including
genome size change, LTR divergence distributions, and solo:intact ratio. Intact LTR elements
were extracted from each simulation using the "extract_intact_LTR.py" script; their
divergences were estimated with the K2P model and a mutation rate of 2.0×10^{-9} per bp per
generation using the "seq_divergence.py" script, and divergences were plotted using the
"ltr_dens.py" script, all available in PrinTE.

Although the BED file output of PrinTE contains the annotation of all intact and solo LTR
sequences, we perform a *de novo* scan of intact LTR-RTs using LTR_retriever (v3.0.1) [30] on the
simulated genomes, so that these results are comparable to those obtained from the real
genome. The resulting LTR library was used to annotate whole-genome LTR sequences with
RepeatMasker (v4.1.7-p1) [47]. Solo LTRs were then identified by the
"solo_intact_ratio_wrap.sh" script available in the LTR_retriever package, which also
calculated the overall solo:intact ratio of each genome. For the same reason, LTR divergence
distributions of simulated genomes were also plotted using intact LTR-RTs identified by
LTR_retriever. The "plot_divergence.R" script available in PrinTE was used to plot the
density of LTR divergence in simulated and real genomes. Parameter sets were ranked by their
ability to reproduce three empirical features: 1) Simulated genome sizes are consistent with
hypothesis predictions; 2) Genome-wide LTR divergence distributions are consistent with
empirical whole-genome data; 3) Solo:intact LTR ratios are close to the empirical value. This
approach identified optimal parameters where simulated TE dynamics showed convergence
with multiple empirical metrics while maintaining computational feasibility.

Benchmarking and performance testing

For benchmarking purposes, we disabled the probability density function that controls the
number of LTR-RT mutations in the burnin ($f(mTE)$) so that LTR-RTs had mutation rates
mirroring those of the genome. This allowed us to compare with other simulators that do not
model such phenomena. We used PrinTE with parameters "-P 10 -itn 150 -cn 1 -sz
100Mb -ge 7500000 -st 500000 -t 20 -k 0 -kt -m 3e-8 -sr 0 -tmx 0 --
TE_lib ./PrinTE/athrep.ltr.full.gz". We used extremely low rates of TE insertion
and deletion ("-F 3.0e-200,5e-200") to effectively disable transpositions and purging so that

the genome and its LTR-RTs are allowed to mutate with only DNA substitutions. Three reps with

"`-TiTv 10`" and "`-TiTv 1`" allowed us to compare the performance of PrinTEs models of

evolution. For comparison, we used SLiM to evolve a single 100 Mb diploid lineage (N=1). SLiM

does not implement an iterative stepwise approach and instead models each generation. To

reduce runtime, we used SLiM to model ancestral trees with no mutations (MU=0), then

projected DNA substitutions onto the mutation-free SLiM genealogies at half-million

generationsteps from 0 - 7.5 million generations using a mutation rate of 3×10^{-8} per bp per

generation and Ti/Tv's of 1 or 10. Mutations in SLiM simulated genomes were detected by

whole-genome alignments using Minimap2 with flags "`-cx asm20 --cs=long --`

`secondary=no`". The PAF alignment files were provided to "`paftools.js`" from the Minimap2

package with the "`stat`" command to calculate p-distance from the reported number of

substitutions and alignment length. Similarly, K2P divergence was calculated using the cs tag in

the PAF file by classifying substitutions as either transitions (purine to purine or pyrimidine to

pyrimidine) or transversions (purine ↔ pyrimidine).

LTR-RT redetection was compared between SimulaTE, ReplicaTE, and PrinTE

simulations. SimulaTE requires a chassis genome as input. PrinTE was used to build the

chassis using "`--burnin_only --cds_percent 0 --TE_percent 0 --chr_number 1`

`--size 1Gb`". The "`define-landscape_random-insertions-freq-range.py`" from

SimulaTE was run using "`--N 3 --insert-count 2000 --min-distance 1000 --`

`min-freq 0.1 --max-freq 0.9`". The resulting pgd file was provided to "`build-`

`population-genome.py`" from SimulaTE to create the final TE-infused genome. ReplicaTE

was used to insert Ty1 and Ty3 elements from *Drosophila melanogaster* chromosome 2 using

the GenBank file from the package. PrinTE was run with parameters "`--burnin_only --`

`cds_percent 0 --TE_num 1000 --chr_number 1 --size 500Mb -s 21 --`

`threads 200`". Three replicates were used for each tool. LTR-RT redetection was conducted

using `LTR_FINDER_parallel` and `LTR_HARVEST_parallel` with parameters "`-w 2 -C -D`

`15000 -d 100 -L 7000 -l 60 -itp 20 -M 0.85 -S 3`" and "`-minlenltr 60 -`

`maxlenltr 7000 -mintsd 4 -maxtsd 8 -mindistltr 300 -maxdistltr 20000 -`

`similar 85 -vic 10 -seed 20 -seqids yes -xdrop 6`", respectively. The resulting

LTR-RT candidates were provided to LTR_retriever with "`-motif [TCCA TGCT TACA TACT`

`TGGA TATA TGTA TGCA TGAA AGTT AGCT]`" to identify LTR-RTs.

PrinTE runtime and peak memory were assessed on an HPC server with 256 CPU

threads of AMD EPYC 7763 2.45GHz and 1024 GB DIMM DDR4 memory at 3200 MHz. Metrics

were collected from the report generated using "`/usr/bin/time -v`". PrinTE was run using "`-`

`cn 20 -sz 100Mb -P 20`" to simulate a 100Mb genome with 20 chromosomes in which 20% of

the genome is genes. "`-itn 6000`" was used to simulate 6000 initial TE insertions. "`-ge`

`1000000 -st 100000`" simulated 1,000,000 generations of TE evolution with step sizes of

100,000 generations (10 iterations total). Insertion and deletion rates, "`-F 3.0e-11,5e-11`", were

used to insert and delete ~500 TE insertions per iteration.

Syntenic Ribbon Plot and LTR-RT Phylogeny

To simulate genomes presented in **Fig. 3A**, 19,621 *Arabidopsis thaliana* cds sequences were inserted into the simulated genome using PrinTE with "`--cds_num 19621`" to guide collinear anchoring and enable syntenic analysis. The burn-in genome consisted of 80% intact LTR-RTs (`-itp 80`). The genome contraction phase was run using "`-ge 5000000 -st 1000000 -F 1e-11,4e-11 -dg -k 3`". The genome size increase phase was run using identical parameters except "`-F 4e-11,1e-11 –continue`" and the contraction phase outputs as the input. The simulated genomes were processed pairwise using "`jcvi.compara.catalog ortholog --cscore=.99`" in JCVI (v1.5.4) [48]. Syntenic anchors were visualized using "`matplotlib.pyplot`" in Matplotlib (v3.10.3) to create the ribbon plot (**Fig. 3A**). The LTR-RT phylogeny (**Fig. 3D,E**) was created using only elements in the *Tekay* family, which were classified as such by TEsorter (v1.4.7) [49]. *Tekay* LTR-RT headers were tagged with "`_shared`" and "`_uniq`" to facilitate post-PrinTE parsing. Shared LTR-RTs were inserted using PrinTE with parameters "`-itp 80 -cn 1 -sz 10Mb -tmx 10 --TE_lib Tekay_shared.fa -tk 15`". The outputs ("`burnin.fa`" and "`burnin.bed`") were duplicated and allowed to continue simulation using either genome expansion parameter ("`--generation_end 5000000 --step 500000 -F 2e-10,1e-10 --TE_lib Tekay_uniq.fa`") or genome contraction parameters ("`--generation_end 5000000 --step 500000 -F -F 2e-11,1e-11 --TE_lib Tekay_uniq.fa`"). Intact LTR-RTs were extracted using "`extract_intact_LTR.py`" and classified using TEsorter with "`-db rexdb-plant -dp2`". Reverse transcriptase domains identified from *Tekay* LTR-RTs were aligned using MAFFT (v7.525) [50] and provided to VeryFastTree (v4.0.5) [51] for phylogenetic analysis. The resulting newick trees were plotted using ggtree (v3.1.5) [52] with branches colored by their ancestral relationship.

Pucciniomycotina Genomes and Annotations

The myrtle rust (*Austropuccinia psidii*) genome was downloaded from NCBI with accession ID ERS3953144 [3]. Genomes of the *Pucciniomycotina* fungus subdivision were obtained from the JGI Fungal Genomics Resource MycoCosm (https://mycocosm.jgi.doe.gov/) (**Table S1**). Rust fungus genomes with contig N50 values higher than 100 kbp were used, for a total of 47 genomes (**Table S2**).

Benchmarking Universal Single-Copy Orthologs (BUSCOs) evaluations were performed on each genome by compleasm (v0.2.7) [37] with the odb12 database and the basidiomycota lineage that has 2409 universal single-copy orthologs. For genomes with duplicated BUSCO higher than 5%, the Redundans (v2020.01.28) [36] program was used to identify alternative haplotypes. Haplotype alignments with an identity higher than 90% and coverage of more than 80% were removed using "`--identity 0.9 --overlap 80`" parameters. compleasm was again used to evaluate the purged genomes with the same parameter. The resulting duplicated BUSCO were considered biological duplications (**Table S2**). The haplotype genomes (de-duplicated) were used in the following analyses.

Genomes with single-copy BUSCOs higher than 50% were used to construct the phylogeny. Melame1 (BUSCO S:1134,D:1147,F:33,I:0,M:95) and MecolCla_1 (BUSCO

S:826,D:1452,F:34,I:0,M:97) were not included in this analysis owing to the high proportion of
duplicated BUSCO. The protein sequences of single-copy BUSCOs present in the remaining 45
genomes were extracted and further aligned using MAFFT (v7.525) [50] with a focus on local
alignments using the "`--maxiterate 1000 --localpair`" parameters. The phylogeny was
constructed using the aligned BUSCO and IQ-TREE (v2.4.0) [38]. Each BUSCO alignment was
treated as an individual partition [53], and ModelFinder [54] was used to identify the best model for
each partition using the "`-m MFP`" parameter in IQ-TREE. The Shimodaira–Hasegawa-like
approximate likelihood ratio test (SH-aLRT) [55] and bootstrap tests were performed 1000 times
each with parameters "`-bb 1000 -alrt 1000`" in IQ-TREE. The bootstrap consensus tree
was plotted by ggtree (v3.1.5) [56] in R.
Transposable elements in each fungal genome were annotated using EDTA (v2.2.2) [57]
following the instructions of Benson et al. (2024) [58]. The "`--force 1`" parameter was used to
overcome analysis in genomes with missing types of TEs (*i.e.*, LTR-RTs). The "`--anno 1`"
parameter was used to obtain whole-genome TE annotations. The "`--cds`" parameter was
used to provide coding sequences to EDTA to improve TE annotations. Primary coding
sequences of each genome were obtained from JGI MycoCosm and Tobias et al. (2021) [3]. Solo
and intact LTR-RTs were identified from whole-genome TE annotations using LTR_retriever
(v3.0.1) [30]. The solo:intact ratio was calculated on the whole-genome level, which is the number
of whole-genome solo LTRs divided by the number of whole-genome intact LTR-RTs. "inf"
values were given to genomes without intact LTR-RTs. The divergence of intact LTR-RTs was
estimated by LTR_retriever (v3.0.1) [30] using the one-parameter Jukes-Cantor 69 model [59].
938

Simulating Digital Replicas of Rust Fungus Genomes

Three fungal genomes were chosen for simulations, including *Puccinia coronata* f. sp. *avenae*
(PuccoNC29_1), *Cronartium ribicola* (Crori1), and *A. psidii* (myrtle rust). TE libraries used for TE
removal and simulating the fungal genomes were obtained via *de novo* TE annotation using
EDTA (as described above). The Genome Approximation Mode of TEgenomeSimulator (v1.0.0)
was used for simulations. The resulting digital replicas of the rust fungus genomes were further
used to simulate LTR dynamics using PrinTE (see sections below).

Forward Simulation of Rust Fungus Genomes

Simulating LTR-RT Dynamics in Rust Fungus Genomes
Cronartium ribicola. We hypothesized that this genome is relatively stable, with high rates of
LTR-RT insertions and deletions. We aimed to evolve the genome with a relatively stable
genome size. The digital replica of the *C. ribicola* genome was generated by
TEGenomeSimulator with a size of 232.6 Mb, which was used as input for PrinTE using the '`--`
`bed`' and '`--fasta`' parameters. Each experiment was simulated with 20,000 generations per
step and a total of 200,000 generations (`-ge 200000 -st 20000`). A series of fixed insertion
and deletion rates was used in different experiments, including the range of insertion rates from
1.0×10^{-9} to 1.9×10^{-8} per bp per generation and the range of deletion rates from 1.4×10^{-9} to
4.0×10^{-8} per bp per generation specified by the '`--fix`' parameter. Additionally, DNA

substitution rates of 1.0×10^{-8}, 2.0×10^{-9}, and 5.0×10^{-9} per bp per generation, k values of 0, 3, 6,
and 10, and solo rates of 60, 70, 75, 80, and 95 were used in the parameter sweep analysis.
The simulated genomes were summarized by PrinTE and LTR_retriever and compared with the
real *C. ribicola* genome for the determination of the best parameters.
Puccinia coronata* f. sp. *avenae. We hypothesized that this genome is relatively stable with
low rates of LTR-RT insertions and deletions. We aimed to evolve the genome with a relatively
stable genome size. The digital replica of the *P. coronata* f. sp. *avenae* genome was generated
by TEGenomeSimulator with a size of 112.7 Mb, which was used as input for PrinTE using the
'`--bed`' and '`--fasta`' parameters. Each experiment was simulated with 1 million generations
per step and a total of 10 million generations (`-ge 10000000 -st 1000000`). A series of
fixed insertion and deletion rates was used in different experiments, including the range of
insertion rates from 8.0×10^{-12} to 5.0×10^{-11} per bp per generation and the range of deletion rates
from 9.6×10^{-12} to 9.0×10^{-11} per generation specified by the '`--fix`' parameter. Additionally,
mutation rates of 2.0×10^{-9} and 5.0×10^{-9} per bp per generation, k values of 0, 3, and 6, and solo
rates of 70 and 95 were used in the parameter sweep analysis. The simulated genomes were
summarized by PrinTE and compared with the real *P. coronata* f. sp. *avenae* genome for the
determination of the best parameters.
Austropuccinia psidii. We hypothesized that this was once an expanding genome but now a
fast downsizing genome with a low rate of LTR-RT insertion and a high rate of LTR-RT deletion.
Owing to the relative phylogenetic closeness to *A. psidii*, we use the genome size and TE
composition of *P. coronata* f. sp. *avenae* genome as an approximation of the ancestral state of
A. psidii before genome expansion. We used PrinTE to simulate such an ancestral genome with
parameters '`--cds_percent 20 --TE_num 6000 --chr_number 20 --size 113Mb -`
`-TE_mut_k 10 --TE_mut_Mmax 20`'. With a fixed deletion rate of 5.0×10^{-13} per bp per
generation, we explored insertion rates from 2.0×10^{-12} to 9.8×10^{-11} per bp per generation and
identified that the insertion rate of 1.4×10^{-11} per bp per generation could yield LTR divergence
similar to the expansion phase of *A. psidii* (**Fig. S13A**,**B**). We thus used PrinTE to simulate a
fast-expanding phase with parameters "`-k 0 --fix 1.4e-11,5e-13 --mutation_rate`
`2e-9 --solo_rate 95`" until it reached 1,178 Mb. For the contraction phase, we slowed
down the insertion rate to 1.4×10^{-13} per bp per generation and explored deletion rates from
3.0×10^{-13} to 5.0×10^{-11} per bp per generation. We identified that the deletion rate of 3.0×10^{-11} per
bp per generation could yield LTR divergence and genome size similar to those of the *A. psidii*
genome (**Fig. S13A**,**B**). Finally, we used the outputs of the last iteration as inputs for PrinTE to
simulate a fast-contracting phase with parameters "`-k 0 --fix 1.4e-13,3e-11 --`
`mutation_rate 2e-9 --solo_rate 95`" until it reached 1000 Gb, approximately the size
of the *A. psidii* genome.
993

Data Availability

995 All codes that produced the findings of the study, including all main and supplemental figures,
996 are available at https://github.com/cwb14/PrinTE.git, under the GPL-3.0 license. The Kmer2LTR
997 code is available at https://github.com/cwb14/Kmer2LTR.git under the GPL-3.0 license.

Acknowledgments

We want to thank Drs. Jason Slot, James Pease, and H. Lisle Gibbs for helpful discussions of this project. We are also grateful to Drs. Chen Wu, Julie Blommaert, Olivia Angelin-Bonnet, and David Chagné for reviewing this manuscript and providing insightful comments.

Funding

This research was supported by the Ohio State University STEM Education Faculty Startup Award (S.O.), the United States Department of Agriculture (USDA) National Institute of Food and Agriculture (NIFA) Agriculture and Food Research Initiative (AFRI) Postdoctoral Fellowship (AWD-117797 to C.W.B.), and by Plant & Food Research Group, Bioeconomy Science Institute through their Kiwifruit Royalty Investment Programme (T.-H.C., C.H.D., and S.J.T.).

Author information

Authors and Affiliations

Department of Molecular Genetics, Ohio State University, Columbus, OH, 43210, USA

Christopher W. Benson & Shujun Ou

Plant and Food Research Group, The New Zealand Institute for Bioeconomy Science Limited, Lincoln 7608, New Zealand

Ting-Hsuan Chen & Susan J. Thomson

Plant and Food Research Group, The New Zealand Institute for Bioeconomy Science Limited, Auckland 1025, New Zealand

Cecilia H. Deng

Contributions

S.O. conceived the project and supervised the work. C.W.B developed PrinTE and Kmer2LTR. C.W.B, T.-H.C., and S.O. performed the analysis. C.W.B, T.-H.C., C.H.D, S.J.T., and S.O. wrote the manuscript text. All authors reviewed and approved the manuscript.

Corresponding author

Correspondence to Shujun Ou (ou.195@osu.edu).

Ethics declarations

Ethics approval and consent to participate

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Lander, E. S. *et al.* Initial sequencing and analysis of the human genome. *Nature* **409**, 860–921 (2001).

2. Hufford, M. B. *et al.* De novo assembly, annotation, and comparative analysis of 26 diverse maize genomes. *Science* **373**, 655–662 (2021).

3. Tobias, P. A. *et al.* Austropuccinia psidii, causing myrtle rust, has a gigabase-sized genome shaped by transposable elements. *G3 (Bethesda)* **11**, (2021).

4. Serrato-Capuchina, A. & Matute, D. R. The Role of Transposable Elements in Speciation. *Genes* **9**, 254 (2018).

5. Beric, A. *et al.* Comparative phylogenetics of repetitive elements in a diverse order of flowering plants (Brassicales). *G3 (Bethesda)* **11**, (2021).

6. Qin, P. *et al.* Pan-genome analysis of 33 genetically diverse rice accessions reveals hidden genomic variations. *Cell* **184**, 3542–3558.e16 (2021).

7. Bayer, P. E. *et al.* Variation in abundance of predicted resistance genes in the Brassica oleracea pangenome. *Plant Biotechnol. J.* **17**, 789–800 (2019).

8. Ou, S. *et al.* Differences in activity and stability drive transposable element variation in tropical and temperate maize. *Genome Res* **34**, 1140–1153 (2024).

9. Hu, Y. *et al.* Gossypium barbadense and Gossypium hirsutum genomes provide insights into the origin and evolution of allotetraploid cotton. *Nat Genet* **51**, 739–748 (2019).

10. Naish, M. *et al.* The genetic and epigenetic landscape of the Arabidopsis centromeres. *Science* **374**, eabi7489 (2021).

11. Blommaert, J. Genome size evolution: towards new model systems for old questions. *Proc. Biol. Sci.* **287**, 20201441 (2020).

12. Kapusta, A., Suh, A. & Feschotte, C. Dynamics of genome size evolution in birds and

mammals. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E1460–E1469 (2017).

13. Ricci, M., Peona, V., Guichard, E., Taccioli, C. & Boattini, A. Transposable elements activity

is positively related to rate of speciation in mammals. *J. Mol. Evol.* **86**, 303–310 (2018).

14. Pellicer, J., Hidalgo, O., Dodsworth, S. & Leitch, I. J. Genome size diversity and its impact

on the evolution of land plants. *Genes (Basel)* **9**, (2018).

15. Galindo-González, L., Mhiri, C., Deyholos, M. K. & Grandbastien, M.-A. LTR-

retrotransposons in plants: Engines of evolution. *Gene* **626**, 14–25 (2017).

16. Niu, S. *et al.* The Chinese pine genome and methylome unveil key features of conifer

evolution. *Cell* **185**, 204–217.e14 (2022).

17. Liang, Y. *et al.* The giant genome of lily provides insights into the hybridization of cultivated

lilies. *Nat. Commun.* **16**, 45 (2025).

18. Fernández, P. *et al.* A 160 Gbp fork fern genome shatters size record for eukaryotes.

iScience **27**, 109889 (2024).

19. Wright, S. I. & Agren, J. A. Sizing up Arabidopsis genome evolution. *Heredity (Edinb.)* **107**,

509–510 (2011).

20. Kovaka, S., Ou, S., Jenike, K. M. & Schatz, M. C. Approaching complete genomes,

transcriptomes and epi-omes with accurate long-read sequencing. *Nat. Methods* **20**, 12–16

(2023).

21. Hoen, D. R. *et al.* A call for benchmarking transposable element annotation methods. *Mob.*

DNA **6**, 13 (2015).

22. Clough, J. E., Foster, J. A., Barnett, M. & Wichman, H. A. Computer simulation of

transposable element evolution: random template and strict master models. *J Mol Evol* **42**,

52–58 (1996).

23. Kofler, R. SimulaTE: simulating complex landscapes of transposable elements of

populations. *Bioinformatics* **34**, 1439 (2018).

24. Bourgeois, Y., Ruggiero, R. P., Hariyani, I. & Boissinot, S. Disentangling the determinants

of transposable elements dynamics in vertebrate genomes using empirical evidences and

simulations. *PLoS Genet* **16**, e1009082 (2020).

25. Deceliere, G., Letrillard, Y., Charles, S. & Biémont, C. TESD: a transposable element

dynamics simulation environment. *Bioinformatics* **22**, 2702–2703 (2006).

26. Verneret, M., Van Anthony, L., Faraut, T., Turpin, J. & Lerat, E. Particular sequence

characteristics induce bias in the detection of polymorphic transposable element insertions.

bioRxiv 2024.09.25.614865 (2024) doi:10.1101/2024.09.25.614865.

27. Groza, C., Chen, X., Wheeler, T. J., Bourque, G. & Goubert, C. A unified framework to

analyze transposable element insertion polymorphisms using graph genomes. *Nat*

Commun **15**, 8915 (2024).

28. Yue, J.-X. & Liti, G. simuG: a general-purpose genome simulator. *Bioinformatics* **35**, 4442–

4444 (2019).

29. Caballero, J., Smit, A. F. A., Hood, L. & Glusman, G. Realistic artificial DNA sequences as

negative controls for computational genomics. *Nucleic Acids Res* **42**, e99 (2014).

30. Ou, S. & Jiang, N. LTR_retriever: A Highly Accurate and Sensitive Program for

Identification of Long Terminal Repeat Retrotransposons. *Plant Physiol.* **176**, 1410–1422

(2018).

31. Ma, J., Devos, K. M. & Bennetzen, J. L. Analyses of LTR-retrotransposon structures reveal

recent and rapid genomic DNA loss in rice. *Genome Res.* **14**, 860–869 (2004).

32. Nowoshilow, S. *et al.* The axolotl genome and the evolution of key tissue formation

regulators. *Nature* **554**, 50–55 (2018).

33. Chen, T.-H., Angelin-Bonnet, O., Deng, C. H. & Thomson, S. *TEgenomeSimulator: A Tool*

to Simulate TE Mutation and Insertion into a Random-Synthesised or User-Provided

Genome. (2025). doi:https://github.com/Plant-Food-Research-

Open/TEgenomeSimulator/releases/tag/v1.0.0.

34. Ahrendt, S. R., Mondo, S. J., Haridas, S. & Grigoriev, I. V. MycoCosm, the JGI's fungal

genome portal for comparative genomic and multiomics data analyses. *Methods Mol. Biol.*

2605, 271–291 (2023).

35. *Systematics and Evolution*. (Not Avail, 2014). doi:10.1007/978-3-642-55318-9.

36. Pryszcz, L. P. & Gabaldón, T. Redundans: an assembly pipeline for highly heterozygous

genomes. *Nucleic Acids Res.* **44**, e113 (2016).

37. Huang, N. & Li, H. compleasm: a faster and more accurate reimplementation of BUSCO.

Bioinformatics **39**, btad595 (2023).

38. Minh, B. Q. *et al.* IQ-TREE 2: New models and efficient methods for phylogenetic inference

in the genomic era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

39. Devos, K. M., Brown, J. K. M. & Bennetzen, J. L. Genome size reduction through

illegitimate recombination counteracts genome expansion in Arabidopsis. *Genome Res.* **12**,

1075–1079 (2002).

40. Edwards, H. M. & Rhodes, J. Accounting for the biological complexity of pathogenic fungi in

phylogenetic dating. *J. Fungi (Basel)* **7**, 661 (2021).

41. Osmanski, A. B. *et al.* Insights into mammalian TE diversity through the curation of 248

genome assemblies. *Science* **380**, eabn1430 (2023).

42. Zoonomia Consortium. A comparative genomics multitool for scientific discovery and

conservation. *Nature* **587**, 240–245 (2020).

43. Marçais, G. & Kingsford, C. A fast, lock-free approach for efficient parallel counting of

occurrences of k-mers. *Bioinformatics* **27**, 764–770 (2011).

44. Katoh, K., Misawa, K., Kuma, K.-I. & Miyata, T. MAFFT: a novel method for rapid multiple

sequence alignment based on fast Fourier transform. *Nucleic Acids Res* **30**, 3059–3066

(2002).

45. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973

(2009).

46. Marco-Sola, S., Moure, J. C., Moreto, M. & Espinosa, A. Fast gap-affine pairwise alignment
    using the wavefront algorithm. *Bioinformatics* **37**, 456–463 (2021).

47. Smit, A. F. A., Hubley, R. & Green, P. RepeatMasker Open-4.0. 2013--2015. Preprint at
    (2015).

48. Tang, H. *et al.* JCVI: A versatile toolkit for comparative genomics analysis. *Imeta* **3**, e211
    (2024).

49. Zhang, R.-G. *et al.* TEsorter: an accurate and fast method to classify LTR-retrotransposons
    in plant genomes. *Hortic. Res.* **9**, uhac017 (2022).

50. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
    improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

51. Piñeiro, C., Abuín, J. M. & Pichel, J. C. Very Fast Tree: speeding up the estimation of
    phylogenies for large alignments through parallelization and vectorization strategies.
    Bioinformatics **36**, 4658–4659 (2020).

52. Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T.-Y. Ggtree: An r package for
    visualization and annotation of phylogenetic trees with their covariates and other associated
    data. *Methods Ecol. Evol.* **8**, 28–36 (2017).

53. Chernomor, O., von Haeseler, A. & Minh, B. Q. Terrace aware data structure for
    phylogenomic inference from supermatrices. *Syst. Biol.* **65**, 997–1008 (2016).

54. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.
    ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**,
    587–589 (2017).

55. Anisimova, M. & Gascuel, O. Approximate likelihood-ratio test for branches: A fast,
    accurate, and powerful alternative. *Syst. Biol.* **55**, 539–552 (2006).

56. Xu, S. *et al.* Ggtree: A serialized data object for visualization of a phylogenetic tree and
    annotation data. *Imeta* **1**, e56 (2022).

57. Ou, S. *et al.* Benchmarking transposable element annotation methods for creation of a

streamlined, comprehensive pipeline. *Genome Biol.* **20**, 275 (2019).

58. Benson, C. W., Heringer, P. & Ou, S. Four strategies for whole-genome annotation of

transposable elements and repeats in maize. *Cold Spring Harb. Protoc.* (2024)

doi:10.1101/pdb.prot108578.

59. Jukes, T. H. & Cantor, C. R. Evolution of Protein Molecules. in *Mammalian Protein*

Metabolism 21–132 (Elsevier, 1969).

1165