Title

Pasta, an age-shift transcriptomic clock, maps the chemical and genetic determinants of aging and rejuvenation

Authors

Jérôme Salignon[1,2#], Maria Tsiokou[3&], Patricia Marqués[3&], Enriqueta Rodríguez-Diaz[3], Hazel Ang[1,†], Federico Pietrocola[3#], Christian G. Riedel[1,2#]

1. Department of Medicine Huddinge, Karolinska Institute, Stockholm, Sweden
2. Department of Molecular Biosciences, The Wenner-Gren Institute, Stockholm University, Stockholm, Sweden
3. Department of Cell and Molecular Biology, Karolinska Institute, Stockholm, Sweden

[#]: Corresponding authors: jerome.salignon@ki.se, federico.pietrocola@ki.se, and christian.riedel@su.se

[&]: Equal contribution

[†] Present address: Institute of Molecular and Cell Biology, Agency of Science, Technology and Research, Singapore

Abstract

As the prevalence of age-related diseases rises, understanding and modulating the aging process is becoming a priority. Transcriptomic aging clocks (TACs) hold great promise for this endeavor, yet most are hampered by platform or tissue specificity and limited accessibility. Here, we introduce Pasta, a robust and broadly applicable TAC based on a novel age-shift learning strategy. Pasta accurately predicts relative age from bulk, single-cell, and microarray data, capturing senescent and stem-like cellular states through signatures enriched in p53 and DNA damage response pathways. Its predictions correlate with tumor grade and patient survival, underscoring clinical relevance. Applied to the CMAP L1000 dataset, Pasta identified known and novel age-modulatory compounds and genetic perturbations, and highlighted mitochondrial translation and mRNA splicing as key determinants of the cellular propensity for aging and rejuvenation, respectively. Supporting Pasta's predictive power, we validated pralatrexate as a potent senescence inducer and piperlongumine as a rejuvenating agent. Strikingly, chemotherapy drugs were highly enriched among pro-aging hits. Taken together, Pasta represents a powerful and generalizable tool for aging research and therapeutic discovery, distributed as an easy-to-use R package on GitHub.

Introduction

The growing prevalence of chronic non-communicable diseases that occur predominantly later in life have positioned aging as a major driver of pathology. This, coupled with advances in understanding the fundamentals of aging and health[1,2], has intensified interest in pharmacological and lifestyle strategies to interfere with aging[3]. Unlike chronological aging, which is inevitably proceeding at a set rate, biological aging is modulated by endogenous and environmental factors that can accelerate, decelerate, or even reverse aging processes[4]. This highlights biological age as a latent but measurable trait that reflects current and future health, captures individual variability in aging rates, and offers a means to evaluate geroprotective interventions.

Over the past decade, a range of computational tools—collectively termed 'aging clocks'—have emerged to quantify biological age and estimate aging rates. These models draw on diverse molecular and clinical hallmarks across multiple 'omics' layers, including epigenetic, transcriptomic, metabolomic, proteomic, and metagenomic data[5]. Among them, epigenetic clocks based on genome-wide CpG methylation patterns have gained prominence due to their high precision in estimating chronological and biological age, strong predictive value for morbidity and mortality, and the stability of DNA methylation (DNAm) changes, which enable the capture of cumulative aging effects[6]. However, DNAm clocks have limitations, including reduced sensitivity to rapid or transient changes[7] and limited biological interpretability, as CpGs are not always clearly linked to gene function[8]. In addition, single-cell DNAm approaches remain costly, technically demanding, and lack standardized protocols, restricting their utility in resolving aging at single-cell resolution[6].

Transcriptomic aging clocks (TACs) address these limitations: they sense transient shifts, yield interpretable gene/pathway outputs, and can be applied to extensive bulk[9], single-cell[10,11], spatial[10,12], genetic[13,14] and chemical[13,15] perturbation datasets. Furthermore, the number of transcriptomic perturbation datasets, such as Perturb-Seq datasets[14] or the Tahoe-100 dataset[15], is rapidly increasing. Applying TACs to such datasets could expose the causal regulators of aging and rejuvenation with high translational potential: compounds that raise the age of cancer cells may boost therapy[16], whereas agents that lower normal-cell age bear regenerative promise, such as improvement in stem cell fitness[17], iPSC protocols[18], or partial reprogramming cocktails[19]. One study matched three tissue signatures to the CMAP L1000 perturbation atlas[13]—a large scale (> 3 million samples, >250 cell lines) bead-based (L1000 technology) genetic and chemical perturbation dataset—and validated novel modulators[20], but its narrow sample base (109 aging samples, two CMAP cell lines, only chemical perturbations) limited scope.

Human TACs have been developed in blood[21], muscle[22], fibroblasts[23], skin[24], and in multi-tissue settings[25–29]. More recently, single-cell TACs have been constructed using blood[30], immune cells[31,32], and PBMCs[33]. Despite this progress, no TAC has achieved widespread adoption comparable to epigenetic clocks. This is due to several limitations, including unreliable performance, lack of user-friendly software, and platform-specific constraints. Gene expression's sensitivity to environmental changes allows detection of transient states but also introduces greater variability and noise than DNA methylation data, demanding robust modeling across multiple datasets. Most TACs—including many single-tissue models and the multi-tissue RNAAgeCalc[26]—are trained on single datasets, leading to overfitting and poor generalization. Indeed, MultiTIMER[27], which was trained on multiple datasets, largely

outperform RNAAgeCalc[26] despite being trained on less samples (~3,000 vs ~9,600). Another
key limitation is the lack of accessible tools. DeepQA and Shokhirev & Johnson's TACs offer
no usable software[25,28], and RAPToR requires users to generate custom references without
default human datasets[29]. Moreover, most multi-tissue TACs[25–28] necessitate steps like raw
data reprocessing, batch correction, and model retraining, limiting their use to bulk RNA-Seq
data and hampering reproducibility.

To address these limitations, we present Pasta (Predicting age-shift from transcriptomic
analysis): a multi-platform, multi-tissue, ready-to-use TAC. Across diverse datasets it
outperformed MultiTIMER[27], largely via p53-centric gene and pathway features. Pasta's
predictions effectively distinguished between senescent, quiescent, and stem cells, in static
and dynamic settings, indicating that Pasta is optimally suited to capture bidirectional cell
state transitions from stemness to senescence. Pasta was a better predictor of tumor grade than
chronological age, reflecting its usefulness for biological age estimation. Applied to CMAP
L1000[13], Pasta revealed new age modulators—including pralatrexate, a potent senescence
inducer—and revealed molecular traits dictating cell-line responsiveness to pro- or anti-aging
perturbations. Pasta thus delivers a robust, interpretable framework for cross-platform
biological-age estimation and systematic discovery of genetic or chemical modulators of
aging and rejuvenation, with translational promise in oncology and regenerative medicine.

Results

Building a general-purpose transcriptomic aging clock using heterogeneous multi-tissue data

We aimed to create a versatile multi-tissue transcriptomic aging clock usable on any gene
expression sample. Heterogeneous raw or normalized matrices from the Genotype-Tissue
Expression Project (GTEx)[34], the Gene Expression Omnibus (GEO)[35], and the Expression
Atlas[36] were rank-transformed, yielding 17,212 healthy profiles from 21 studies (18 bulk
RNA-seq, 3 microarrays; Fig 1a, Table S1). We then compared three different modeling
strategies: regression models (Fig. 1b), young vs old classifiers (Fig. 1c), and age-shift
classifiers (Fig. 1d). Our proposed age-shift classifier mirrors twin neural network
regression[37,38] and pairwise difference regression[39] or classification[40]: it trains on matched-
tissue sample pairs separated by a defined age gap. Classifiers' predictions were converted to
age scores (see Methods).

Using a leave-one-dataset out (LODO) strategy, we found that the 40-years age-shift model
had the highest median Pearson correlation coefficient (PCC) when considering all samples
(Fig. 1e) or only samples from young and old individuals (Fig. S1a), with accurate predictions
across tissues and platforms (Fig S1b-S3). Models trained on un-ranked counts collapsed
(median PCC < 0.1), confirming the need for rank transformation (Fig 1e, S1a). Next, we
tested the best three models on single-cell RNA-Seq datasets. As only bulk transcriptomic
data were used to train the models, we created pseudobulks by aggregating single cells using
chunk sizes from 1 (single-cell) to 1024 to see how increased data reliability improved model
performance. We found that the age-shift model performed best overall in all three datasets
analyzed (Fig. 1f-h)[41–43]. Interestingly, the young vs old classifier failed in datasets without

samples in both the young and old categories (Fig. 1f,h), highlighting issues with fixed age
cutoffs (Fig. 1c). Overall, we found that the 40-years age-shift model had the best
performance for predicting relative age from heterogeneous transcriptomic data and we refer
to it as Pasta in the remainder of this manuscript.

Pasta outperforms alternative approaches and leverages p53-related pathways to
predict transcriptomic age

To evaluate Pasta's performance, we compared it with MultiTIMER, the current state-of-the-
art, available, multi-tissue RNA-Seq aging clock[27]. We found that Pasta outperformed
MultiTIMER in 6 out of 8 datasets from our LODO analysis (Fig. 2a, Table S2) and in 5 out
of 6 independent datasets (Fig. 2b). These datasets represented diverse tissues (Table S2)
which validate Pasta's multi-tissue capabilities.

We then conducted gene set enrichment analysis (GSEA) on the REACTOME, KEGG,
miRDB, and CGP subsets[44] using Pasta's coefficients. The p53 tumor suppressor pathway
was significantly enriched in the positive coefficients of the model in the KEGG, and CGP
gene sets (Fig. 2c, Table S3). Additionally, irradiation stress terms, which are tightly linked to
p53 dynamics[45], were also enriched. The strongest negative enrichments were targets of
miR29, a p53 activator[46] which drives aging phenotypes in mice[47]. Interestingly, no pathway
was significantly enriched when using the baseline regression model, indicating that the age-
shift learning strategy amplified aging-related signal.

Individual coefficients mirrored these trends. RANBP17, carried the largest negative weight,
nearly twice that of any other gene (Fig. 2d). Interestingly, RANBP17 has been shown to
promote cell proliferation[48] and was proposed to be a master regulator of aging which
decreases with age[49]. The strongest positive weights belonged to PCDHB3, a colorectal-
cancer tumour-suppressor[50], CDKN2A (p16), a canonical p53 target and senescence marker
(Fig. 2e), and ZMAT3 another direct p53 regulator.

In conclusion, Pasta operates as multi-tissue, and multi-platform relative age prediction model
which grounds its predictions in tumor-suppressor genes (TP53, PCDHB3) and core ageing
modulators (TP53, miR29, RANBP17).

Pasta accurately tracks cellular age across senescence, pluripotency and differentiation
states

Since Pasta bases its predictions on p53, a key regulator of cell cycle and senescence, we
investigated whether it could be used to identify senescent cells. Pasta effectively separated
senescent from proliferative cells in 19 out of 30 RNA-Seq datasets (AUC-ROC of 1) and
yielded moderate discrimination (AUC-ROC > 0.7) in five additional datasets (Fig. 3a, Table
S4). It likewise robustly distinguished senescent from quiescent cells in 4/5 datasets (Fig. 3b).
In a 450-day serial-passage microarray experiment of primary fibroblasts from a 5-year-old
donor until replicative senescence, Pasta's age scores tracked culture duration almost linearly
(PCC = 0.896, Fig. 3c). We next asked whether Pasta can detect chemical inducers of
senescence in single-cell data. Merging two studies that assayed the same 138 compounds in

A549 cells—one by scRNA-seq[51], the other by flow cytometry using a senescence
reporter[52]—we observed that Pasta's age scores predicted senescence-inducing molecules,
with AUC > 0.8 at the two highest doses (Fig. 3d).

Given Pasta's ability to identify very old cells, we next tested its sensitivity for detecting very
young cells. Also here, Pasta effectively separated embryonic stem cells from fibroblasts in
four bulk RNA-seq datasets (Fig. 3e) and induced Pluripotent Stem Cells (iPSCs) from
fibroblasts in five more (Fig. 3f). During OSKM-mediated fibroblast reprogramming, Pasta's
age scores declined linearly over time (PCC = -0.905, Fig. 3g). Conversely, Pasta's age scores
had strong correlation with time in a differentiation experiment of human embryonic stem
cells into liver cells (PCC = 0.926, Fig. 3h).

Taking all these observations together, we conclude that Pasta can be used to track cellular
age in static and dynamic settings.

Pasta reveals cancer-type specific associations between cellular age, tumor grade, and
patient survival

Studies using transcriptomic stemness prediction models have shown that higher stemness
correlates with advanced tumor grade and poorer overall survival in cancer patients[53,54].
Recently, older cancer cell age has also been linked to adverse outcomes[16]. Indeed, senescent
cells, both normal and cancerous, accumulate in certain tumors, can drive systemic
inflammation, facilitate relapse, and promote immune evasion[16,55]. However, The Cancer
Genome Atlas (TCGA) transcriptomic data have, to our knowledge, not yet been used to
predict older cellular age, leaving it unclear whether poor prognosis is driven by high
stemness alone or also by senescence. We addressed this gap by leveraging Pasta's capacity to
identify cells spanning both youthful/high-stemness and aged/senescent-like states.

In line with previous results, we found that six out of twelve cancer types had a significantly
negative correlation between age scores and tumor grade (Fig. 3i, j, Table S5). Interestingly,
two cancer types, kidney renal clear cell carcinoma and pancreatic adenocarcinoma, had a
significantly positive correlation between tumor age and tumor grade (Fig. 3i, k) suggesting
that both increased and decreased biological age can be associated with tumor grade. While
Pasta was trained to predict age, its predictions exhibited stronger correlation with tumor
grade than donor age itself, strongly suggesting that Pasta's age scores reflect cellular
biological age instead of merely chronological age (Fig. 3i).

We trained Cox proportional hazards regression models to examine associations between age
or Pasta's age scores and all-cause mortality across 33 cancer types. After multiple testing
correction, younger age scores were significantly linked to poorer survival in ACC, SARC,
and SKCM (Fig. 3l,m; Table S6), while older age scores were similarly associated in THYM
and LAML (Fig. 3l,n). Several other cancers showed strong unadjusted associations. In
contrast, chronological age predicted worse survival in 18 cancer types and was not protective
in any. Thus, both younger and older cancer cell age scores were linked to tumor grade and
survival, suggesting Pasta scores may help stratify patient risk in a cancer-specific manner.

Systematic identification of age-promoting and rejuvenating small molecules in the CMAP L1000 dataset

To systematically identify genetic and chemical perturbations influencing cellular age, we applied Pasta to the CMAP L1000 dataset[13], leveraging its compatibility with heterogeneous transcriptomic data. Age scores were plate-normalized using respective negative controls (DMSO for compounds; empty vectors for genetic perturbations). We identified 259 and 59 compounds significantly associated to older and younger age scores—termed 'Aging' and 'Rejuvenating' compounds, respectively (Fig. 4a; Table S7). Top Rejuvenating compounds included pluripotency-promoting molecules, such as RepSox and PD0325901[56]. Top Aging compounds included senescence inducers such as mitoxantrone, gemcitabine, and doxorubicin[57]. To evaluate Pasta's ability to identify compounds with strong age-modulatory effect, we annotated compounds using drug classes promoting senescence[57] and pluripotency[56]. Notably, 18.9% of Aging compounds belonged to senescence-inducing drug classes, versus <0.2% in the Other/Rejuvenating sets (Fig. 4b), while 16.9% of Rejuvenating compounds belonged to reprogramming-inducing drug classes, again versus <0.2% in the other groups (Fig. 4c). Furthermore, 27.1% of Rejuvenating compounds were annotated as both, compared to 1.54% and 0.07% in the Aging and Other groups (Fig. 4d), with all such hits being HDAC inhibitors—agents known to promote reprogramming[58,59] and also induce senescence[60–62]. When repeating the analysis using the base regression model (Fig. 1), only 12.7%/6.5%/19.4% (vs 18.9%/16.9%/27.1%) of Aging/Rejuvenating/Both sets showed these annotations (Fig. S4), confirming that Pasta's age-shift model yields more biologically informative predictions.

We next explored mechanisms underlying these compound effects using the mechanisms of actions (MOAs) annotated in the dataset. Over 26 MOAs were significantly enriched among Aging compounds (Fig. 4e; Table S8), including several pro-senescence pathways such as topoisomerase, CDK, MDM, and IMPDH inhibitors, largely related to growth, cell cycle, and signaling. Only six MOAs were enriched in Rejuvenating compounds (Fig. 4f), with HDAC inhibitors showing the strongest enrichment, consistent with their potent role in accelerating the reprogramming process[58,59]. This set also included MEK and GSK-3 inhibitors, common in reprogramming cocktails[56], as well as cardiac glycosides and proteasome inhibitors—both described as senolytics[63]. Thus, selective elimination of older cells in culture may lead to net rejuvenation (see Discussion).

Analyzing compound effects across cell lines, proteasome inhibitors MG-132 and bortezomib showed consistent rejuvenating effects, while mitoxantrone showed consistently strong aging effects (Fig. 4g; Table S9). Some HDAC inhibitors (e.g., trichostatin-A, vorinostat, panobinostat) were rejuvenating in most lines but aging in neuronal lines (NEU, NPC, SH-SY5Y), consistent with trichostatin-A's cytotoxicity in SH-SY5Y[64]. These findings support the dual role of HDAC inhibitors in promoting reprogramming and senescence[58–62].

To test if age-modulatory compounds affected cancer cell viability, we integrated Cancer Dependency Map (DepMap)[65] data. Both Aging and Rejuvenating sets were broadly toxic to cancer cells, with median dependency scores of –1.86 and –2.97, respectively (Fig. 4h). MOA analyses revealed a significant link between age and dependency scores only for MDM and IMPDH inhibitors (Fig. 4i,k; Table S10). Among all compounds, only two MDM inhibitors—AMG-232 and CGM-097—showed statistically significant effects (Fig. 4j,k; Table S11). These findings indicate that MDM inhibitors may kill cancer cells by accelerating cellular

251 aging, in line with the dual role of the p53–MDM2 axis in promoting both senescence and
252 apoptosis[66].

Identification of potential cancer-specific chemotherapeutic compounds

Chemotherapeutic drugs are thought to exert their effects by increasing the biological age of
both cancer and normal cells[16], thereby reducing tumor stemness and aggressiveness.
Leveraging the scale of our dataset, we systematically tested this hypothesis. We examined
cancers with ≥5 approved chemotherapeutics to assess enrichment within our three age-
association groups (Aging, Rejuvenating, Other) across cell lines. Notably, 17 of 27 cancers
were significantly enriched in the Aging group (Figure 5a; Table S12,13). Of the ten non-
significant cancers, eight had few approved drugs and adjusted p-values <0.2. Only one
cancer showed enrichment in the Rejuvenating group, and none in the Other group. These
results not only support the pro-aging mechanism of chemotherapeutics[16] but also highlight
our method's capacity to identify candidate chemotherapeutics in a data-driven manner—
particularly valuable for poorly treatable cancers. Future work could expand this approach by
using only cell lines relevant for a given cancer with validation in primary clinical material.

We also searched for compounds inducing stronger senescence in cancer cells than in normal
cells, aiming to discover agents with potentially reduced side effects. Pralatrexate, a
dihydrofolate reductase inhibitor, showed the strongest pro-aging effect—four times higher in
cancer cell lines (Fig. 5b-c; Table S14). Additional promising compounds included Exifone
(an HDAC1 activator) and Brequinar (a dihydroorotate dehydrogenase inhibitor), representing
potential candidates for selective, low-toxicity chemotherapies.

Experimental validation of predicted small molecules' age-modulatory effects

To experimentally assess the predictive power of Pasta in identifying compounds capable of
promoting age-modulatory effects in a cell type–dependent manner, we exposed melanoma
A375 cells (responders; Table S15) and breast cancer MDA-MB-231 cells (non-responders)
to pralatrexate, the 4th top Aging drug (Tables S7). In line with Pasta's prediction, we
observed that pralatrexate induced cell enlargement and positivity for the Senescence-
Associated β-Galactosidase (SA-β-Gal) assay (Figure 6a), transcriptional upregulation of
CDKN1A (also known as p21) and the SASP component IL6 (Figure 6b), proliferative arrest
marked by reduced EdU incorporation (Figure 6c), and downregulation of LMNB1 alongside
upregulation of p21 protein levels (Figure 6d)—all consistent with the induction of a *bona*
fide senescence response. Conversely, in MDA-MB-231 cells, pralatrexate induced a stress
response lacking key hallmarks of cellular senescence, as defined by minimal morphological
changes yet detectable SA-β-Gal positivity (Figure 6e), negligible variations in CDKN1A and
IL6 mRNA levels (Figure 6f), increased proliferation and EdU incorporation (Figure 6g), and
modest changes in LMNB1 accompanied by upregulation of p21 protein levels (Figure 6h).

We also examined the capacity of Piperlongumine—the 20th top rejuvenating agent according
to Pasta scoring (Table S7)—to promote stemness features in prostate cancer PC3 cells
(responders; Table S15). We found that Piperlongumine induced transcriptional upregulation

of the pluripotency transcription factors OCT4, SOX2, and NANOG, accompanied by
increased Oct4 protein levels, in PC3 cells (Figure 6i,j).

These findings underscore the robust predictive power of Pasta in capturing cell type–specific
age-modulatory effects, including the ability to distinguish age-neutral stress response from
bona fide senescence. These data strongly support the potential application of Pasta for in-
silico identification of senescence- or reprogramming-inducing compounds, including through
the mining of existing perturbation datasets.

**Systematic identification of age-promoting and rejuvenating genetic perturbations in the
CMAP L1000 dataset**

We next sought to define the genetic determinants of cellular age from the CMAP L100
dataset, including gene knockdown, knockout, and overexpression. Across cell lines, we
identified 841 gene perturbations (GPs) significantly associated with older age scores
('Aging' GPs) and 54 associated with younger scores ('Rejuvenating' GPs). Top aging GPs
included knockout of CCNA2/cyclin and knockdown of the oncogenes ERG and MYC (Fig.
7a; Table S16), consistent with previous observations[67–69]. Overexpression of KRAS, BRAF,
and RIT1 also led to older scores, reflecting the tumor-suppressive mechanism of oncogene-
induced senescence (OIS)[70]. Conversely, MYC overexpression and SH2B3 or SMAD4
knockdown produced younger scores, aligning with earlier reports[71–73]. Both TP53 knockout
and knockdown were among the top 51 youngest GPs (out of 10,479), consistent with its
central tumor-suppressive role and its mutation in 50-60% of human cancers[74].

To characterize these GPs, we performed pathway enrichment using the Open Genes[75] and
REACTOME[76] databases (Fig. 7b–f; S4; Tables S16–S18), inverting overexpression effects
to pool GPs results. The only Aging Hallmark[75] enriched in Rejuvenating GPs was "stem cells
exhaustion" (Fig. 7b), while "senescent cells accumulation" was the most enriched in Aging
GPs (Fig. 7c). Additional enrichments in Aging GPs were related to transcriptional regulation,
insulin/IGF-1 signaling, and telomeres. In REACTOME, Rejuvenating GPs were enriched in
RAF/MEK/ERK and FGFR signaling (Fig. 7d), whereas Aging GPs were enriched in
pathways linked to cell cycle progression, receptor tyrosine kinase activation (KIT, VEGF,
NTRK), PI3K/AKT signaling, and immune activation (TLR4, CD28) (Fig. 7e). Integration
with compound targets revealed Aging GP enrichment in p53-dependent cell cycle regulation
and growth factor signaling (e.g., VEGF, ERBB2, KIT, extranuclear estrogen) (Fig. 7f; S4g).

Next, we identified GPs with cell-line-specific age effects. Some GPs had consistent age
effect across multiple cell lines: MYC overexpression had a consistent rejuvenating impact,
while MYC knockdown, CCNA2 knockout, and BRAF/KRAS overexpression drove aging
(Fig. 7g; Table S19). Some GPs, like SH2B3 knockdown and ASAH1 knockout, showed
opposing effects depending on cell line, highlighting context-dependent roles.

Finally, we assessed whether age-altering GPs affected cancer cell viability. Aging knockout
GPs had negative DepMap dependency scores (median: -1), while other GP sets hovered
around zero (Fig. 7h). REACTOME-based analysis identified five pathways where age scores
had significant association with dependency (Fig. 7i; Table S20), including retinoic acid
biosynthesis (Fig. 7j), OIS (Fig. 7k), G1/S DNA damage checkpoints, and TP53 stabilization.

These findings suggest that age-promoting GPs, particularly related to senescence and TP53,
also simultaneously promote cancer cell death.

Defining the molecular determinants of cellular aging and rejuvenation

Next, we leveraged the comprehensive nature of the CMAP L1000 and DepMap datasets to
define the molecular determinants of cellular propensity for age changes. Aging and
rejuvenation propensities were computed for 19 cell lines by averaging age scores of
significant age-regulatory perturbations across cell lines (see Methods). Results revealed
marked heterogeneity across lines (Fig. 8a), with MDA-MB-231 showing the strongest
rejuvenation propensity (-14.2 vs. -9.6 in MCF10A), consistent with its role as a model for
aggressive, triple-negative breast cancer[77].

Correlating aging propensity with gene mutation burden, TP53 emerged as the only
significantly associated gene (PCC = 0.875; Fig. 8b; Table S21). Cell lines with two TP53
mutations had negative aging-propensity scores (i.e., JURKAT), those with a single mutation
scored below 6, and TP53-wild-type lines scored above 6 (Fig. 8c). These results are in line
with the key role of TP53 as a tumor suppressor that induces senescence in highly
proliferative cells[78,79] and suggests novel potential tumor suppressor genes (Fig. 8b).
Consistent with this idea, top significant genes in Figure 8b included the known tumor
suppressor genes FAT3[80], HERC1[81], and ABR[82,83].

To identify pathways influencing aging and rejuvenation propensities, we performed
REACTOME-based enrichment on molecules (proteins, transcripts, gene copy numbers)
correlated (|PCC| > 0.4) with each trait (Fig. 8d–i; S5; Table S22). Proteins whose abundance
correlated with aging propensity were enriched in 'mitochondrial translation', 'respiratory
electron transport', and 'RNA processing' (Fig. 8d). Similarly, aging propensity-associated
transcripts were enriched in 'mitochondrial translation', 'ABC transporter disorders', and
'TP53-regulates metabolic genes' (Fig. 8e), suggesting mitochondrial activity is a central
regulator of aging. This echoes recent findings linking p53, mitochondrial translation, and the
senescence-associated secretory phenotype (SASP; see Discussion)[84]. Copy number–based
enrichments included interferon-related pathways—e.g., 'TRAF6-mediated IRF7 activation'
and 'IFNA signaling' (Fig. 8f)—consistent with type I interferons' role in SASP[85], DNA
damage–induced[86], and oncogene-induced senescence[87], as well as their pro-aging effects[88].

Rejuvenation propensity correlated with protein abundance in 'mRNA splicing', 'HATs
acetylate histones', and 'chromatin modifying enzymes' (Fig. 8g). Notably, 'HDACs
deacetylate histones' was the top negatively correlated term for rejuvenation-linked copy
numbers (Fig. 8h), followed by PRC2-related and DNA methylation pathways. These findings
underscore the influence of histone-modifying enzymes on reprogramming capacity, aligning
with HDAC inhibitors' roles in reprogramming[58,59] and the key role of PRC2 in
pluripotency[89] and as an aging/rejuvenation biomarker[90–92]. Finally, proteins negatively
correlated with rejuvenation propensity were enriched in apoptosis and checkpoint modules,
including programmed cell death and G2/M control (Fig. 8i). This result is consistent with
evasion of cell death being a hallmark of cancer[93] and p53/p21 acting as barrier to
reprogramming[94].

Discussion

In this study, we introduced Pasta, a versatile multi-tissue transcriptomic aging clock that accurately predicts relative age across diverse transcriptomic datasets. We demonstrated its superior performance over conventional regression and classification approaches, validated its efficacy in both bulk and single-cell data, and showcased its utility in predicting cellular aging, senescence, and stemness, as well as in cataloging age-modulatory perturbations across cell lines. Collectively, these findings establish Pasta as a robust and generalizable tool for aging research and translational applications.

A key strength of Pasta lies in its adaptability to diverse transcriptomic platforms. Trained on raw and normalized, non-preprocessed RNA-Seq and microarray data, its heterogeneous training set enabled the model to learn age-predictive patterns based solely on the relative ordering of gene expression. Consequently, Pasta can be applied directly to bulk, single-cell, or L1000 data without realignment or batch correction. It outperformed MultiTIMER[27], the current state-of-the-art, available, multi-tissue TAC, on bulk RNA-Seq and is distributed as a user-friendly and readily-available R package (https://github.com/jsalignon/pasta), underscoring its value to the aging field.

Pasta employs an innovative age-shift learning approach based on pairwise sample training, conceptually related to twin neural networks[37,38] and pairwise regression/classification methods[39,40], known to improve prediction accuracy. Our analysis confirmed performance gains across bulk and single-cell datasets. Moreover, Pasta relied on more biologically relevant features (Fig. 2c) and generated more meaningful predictions on CMAP L1000 data (Fig. S4) than a conventional regression model, positioning age-shift as a powerful method for predicting relative age.

Strong evidence implicates DNA damage response (DDR), cell cycle, and p53 signaling as core biological drivers of Pasta's predictions. Enrichment analysis revealed p53- and irradiation-related pathways in Pasta's gene signatures (Fig. 2c), and key predictors like CDKN2A/p16 and ZMAT3 are p53 targets (Fig. 2e). Topoisomerase inhibitors, which induce DNA damage, and CDK inhibitors, which inhibit the cell cycle, were the two most significantly enriched MOAs for Aging compounds (Fig. 4e). Finally, G1/S phase transition and regulation of TP53 were among the most significantly enriched pathways for Aging perturbations with consistent chemical and genetic effect (Fig. 7f). DDR activation in response to genotoxic stress triggers p53-mediated senescence[95], and a recent study proposed DDR as a minimal unit of aging clocks[96], where DDR, cell cycle, and senescence represent aging over hours, days, and months. Within this framework, Pasta may capture transcriptomic aging at various timescales by leveraging predictive features from distinct temporal layers.

Pasta showed strong predictive power for identifying cellular senescence, stemness, and age-related dynamics such as reprogramming and differentiation. Notably, it achieved this without being trained on senescent or stem cell data. For example, the REO stemness index[54], a leading bulk transcriptomic stemness predictor, used GTEx data as a negative class during training (non-stem), whereas Pasta was trained directly on GTEx (among other datasets). This suggests that healthy donor transcriptomes might enhance stemness and senescence prediction models, helping to address the scarcity of senescent and stem cell training data. These

findings also raise questions about the true nature of epigenetic clocks' predictions; i.e., are
they predicting biological age at the organismal (i.e., health status) or cellular
(stemness/senescence) level? Supporting the later hypothesis, recent studies reported a partial
overlap between aging and senescence-related CpGs[97] and the control of stemness by DNA
methylation marks in astrocytes[98]. A quantitative assessment of how well epigenetic clocks
predict stemness and senescence could help interpret results from such clocks.

Clinically, Pasta's age scores significantly correlated with tumor grades and survival
outcomes across multiple cancer types (Fig. 3i-n), supporting the potential clinical utility of
Pasta in stratifying patients in distinct risk groups. Furthermore, the fact that Pasta's age
scores were more strongly associated with tumor grade than age itself (Fig. 3i) indicates that
Pasta's age scores reflect biological rather than chronological age.

CMAP L1000 analysis showed marked enrichment of senescence-inducing perturbations in
the Aging group (Fig. 4b, 7c) and of reprogramming- and stemness-associated perturbations
in the Rejuvenating group (Fig. 4c,d, 7b). Notably, Aging perturbations (259 compounds, 841
GPs) greatly outnumbered Rejuvenating ones (59 compounds, 54 GPs), underscoring the
diverse triggers of accelerated aging versus the precision required for rejuvenation.
Perturbation effects were largely consistent across both cancer and normal cell lines (Tables
S9, S19), indicating conservation of aging and rejuvenation pathways in health and disease.

A recent study showed that OSKM expression in cancer cells can induce senescence and
apoptosis[99]. Furthermore, our top Rejuvenating compounds—MG-132 and bortezomib—have
been reported to impair OSKM-mediated reprogramming and reduce stem cell
pluripotency[100]. These discrepancies may be due to CMAP's short exposure time of ~24h for
most samples. Indeed, proteasome inhibitors like MG-101 and MG-132 initially upregulate
Oct4 (day 2) in adipose precursor cells, but suppress it at later stages (days 25–30)[101]. Thus,
our results likely reflect the early stages of reprogramming, which may be consistent across
cancer and normal cells. Therefore, they have particular relevance for partial reprogramming,
aimed at restoring cell function with minimal identity change, to limit cancer risk[102].

Unexpectedly, proteasome inhibitors and cardiac glycosides were among the top Rejuvenating
compounds (Fig. 4f). Interestingly, the cardiac glycoside ouabain, ranked 34th (Table S7),
was recently predicted to exert rejuvenating effects in mice using a transcriptomic mortality
clock[103]. CT-200783 and piperlongumine—structurally related—were the only top-20
Rejuvenating compounds outside HDAC, MEK, or proteasome inhibitor classes. Notably,
proteasome inhibitors, cardiac glycosides, piperlongumine, and HDAC inhibitors have been
described as having senolytics properties[63]. Interestingly, different studies reported either
senolytic or pro-senescence effects of bortezomib[63], indicating that cellular response may
depend on cell state or environmental conditions, such as time of compound exposure. Our
results suggest that the observed rejuvenation may result from selective removal of older
cells, thereby rejuvenating the population as a whole. Furthermore, the surviving cells could
feed from the remnants of dead cells to further boost growth and stemness[104,105]. Supporting
this, HDAC inhibitors have shown rejuvenating properties[106], and proteasome inhibitors can
initially induce OCT4[101]. Future studies should dissect the relative roles of old cell clearance,
growth from dead cells debris, and intrinsic rejuvenation in these compounds' rejuvenating
effects.

Among the chemical perturbations identified, pralatrexate emerged as a novel senescence-
inducing compound. Ranked fourth overall for senescence induction (Table S7) and validated
experimentally (Fig. 6), it aligns with the known effects of other dihydrofolate inhibitors like
methotrexate[107–109] (ranked 98th), pemetrexed[110], and pyrimethamine[111]. Notably, pralatrexate
ranked first for inducing senescence more strongly in cancer cells (Fig. 5b–c; Table S14),
suggesting it may be an alternative to methotrexate for limiting tumor growth with reduced
side effects[109]. FDA-approved in 2009 for relapsed or refractory peripheral T-cell
lymphoma[112], its senescence-inducing effects had not been previously reported. This
underscores the power of transcriptomic aging clocks like Pasta to uncover unrecognized
therapeutic properties of known drugs and may help clarify pralatrexate's risks and benefits,
particularly by leveraging insights into the role of senescence in normal and cancer cells.

Integrating CMAP L1000 with DepMap data revealed key molecular determinants of aging
and rejuvenation. Transcriptomic and proteomic analyses showed that cell responses to pro-
senescence stimuli were strongly influenced by mitochondrial translation activity (Fig. 8d,e).
This aligns with extensive evidence linking mitochondrial metabolism, ROS production, and
senescence[113]. Mitochondrial translation, essential for biogenesis, generates ROS[114], which
are required for p21-mediated senescence[115]. Additionally, p53 has been shown to enhance
mitochondrial translation in senescent cells, supporting cytokine production via SASP[84].
Genetically, cells with higher type I interferon gene copy numbers were more prone to aging
responses (Fig. 8f), consistent with the pro-senescence role of interferon signaling in SASP
and neighboring cell effects[116]. Together, these findings support a model in which enhanced
mitochondrial translation and interferon signaling prime cells for pro-senescence responses
via ROS and inflammatory pathways.

Conversely, rejuvenation was primarily governed by epigenetic factors (Fig. 8g,h), consistent
with known reprogramming barriers[106,117]. Notably, reduced copy numbers in PRC2 and
HDAC genes increased sensitivity to rejuvenating stimuli (Fig. 8h), in line with PRC2's role
as a universal biomarker of aging and rejuvenation[90] and HDAC inhibitors' efficacy in
reprogramming[106]. Altogether, our analysis identifies both established and novel regulators of
aging and rejuvenation, offering mechanistic insights and potential therapeutic targets.

In conclusion, Pasta is a versatile, robust, and biologically informative transcriptomic aging
clock with broad utility—from fundamental research on aging, senescence, and stem cells to
clinical diagnostics and therapeutic discovery (Figure S7). By enhancing the accuracy and
generalizability of transcriptomic aging clocks, we enabled systematic mining of genetic and
chemical perturbations to identify causal regulators of cellular age. This approach holds
strong translational promise, facilitating data-driven identification of pro-aging perturbations
for cancer therapy and rejuvenating interventions for regenerative medicine and future age-
reversal strategies.

Methods

Construction of the aging transcriptomic database

GTEx Protected Access Data version 8 (17,382 samples) was used to obtain precise age
information[34]. 1,674 GTEx samples with a RIN score below 6 were discarded. An additional

1,137 samples originating from gender-specific tissues (cervix, fallopian, uterus, prostate,
ovary, testis, vagina), tissues with low sample sizes (e.g., bladder, 12 samples), or non-
relevant sources (e.g., *cells_ebv_transformed_lymphocytes*) were further discarded. Raw or
normalized count matrices were retrieved in R using the package GEOquery[118] for GEO[35]
datasets and using the package ExpressionAtlas[36] for Expression Atlas[119] datasets. The
biomaRt R package[120] was used to convert gene or probe ids into Ensembl gene ids. Custom
scripts were used to parse health status, age, and tissue metadata. Overall, 21 datasets with
data from healthy donors were identified. These had a total of 17,212 samples, with 14,571
from the GTEx dataset, and contained 8,113 genes in common, which were also included in
the landmark or best inferred genes sets in the CMAP L1000 dataset[13], and which were used
for further analyses. A rank transformation was applied to each sample using base R's *rank*
function with the parameter *ties.method = 'average'*.

Generating sample pairs

For age-shift models, pairs of samples from the same dataset were generated randomly and
then filtered. Sample IDs were shuffled 200 times for the training set and 2000 times for the
testing set within each dataset to generate IDs for both the first and the second samples in the
pairs. Then, pairs with an age difference below the cutoff, involving non-healthy individuals,
or representing duplicate pairs (i.e., sample 1 and sample 2 swapped) were excluded. Finally,
datasets with more than 1.5 times as many pairs as samples were randomly downsampled to
maintain a ratio of 1.5 pairs per sample. Finally, the ranked gene expression values were
subtracted between sample 1 and sample 2 and used as input to train the models. The output
variable was binary, indicating whether sample 1 was younger or older than sample 2.

Training and evaluating models

Ridge-regularized generalized linear models were built using 10-fold cross-validation to
identify the optimal lambda parameters. This was done using the function *cv.glmnet* from the
glmnet package[121], and using the following parameters: *s_lamba = 'lambda.min'*, and
type.measure = 'mse' for regression models or *type.measure = 'deviance'* for classification
models. Models were compared using a leave-one-dataset-out (LODO) cross-validation
strategy. The age-effects of the classification models (traditional and age-shift classifiers)
refer to the log-odds obtained using the *predict* function from the glmnet package with
parameter *type = 'link'*. The Pearson correlation coefficients of the predicted ages (for the
regression model) or age-effects (for the classification models) with the true ages in the left-
out datasets were used as the metric to compare models. To convert the classifiers' predictions
into age differences, referred to as age scores, they were multiplied by a scaling factor specific
to each classifier. These scaling factors were obtained by applying the cross-validation models
to a set of pairs with balanced age differences in the left-out datasets. Specifically, 1000 pairs
were generated per dataset which were then filtered to retain 20 pairs in each of 12 bins
representing 10-year age-differences. Age predictions were made and results across all
datasets were aggregated. Finally, the scaling factor was defined as the beta coefficient from a
linear regression of age_difference on age_effect.

Comparing Pasta with MultiTIMER

The MultiTIMER model cannot be used directly, as it requires query samples to be normalized alongside the training samples[27]. As a result, it only functions with samples available on ARCHS4[9]. We re-ran the MultiTIMER training code using ARCHS4 version 2.1, incorporating 14 test datasets —8 from the LODO analysis and 6 novel datasets—alongside the MultiTIMER's training samples to ensure joint batch correction. LODO datasets were evaluated using Pasta's LODO performance, while the novel datasets were assessed using predictions from the final Pasta model.

Conducting gene set enrichment analysis

GSEA was performed using the GSEA function from the *clusterProfiler* R package[122]. Gene sets were retrieved via the msigdbr function from the *msigdbr* package[123], aggregating sets from the CPG, KEGG, REACTOME, and MIRDB subcategories. Model coefficients, sorted by decreasing magnitude, were used as input. The *TERM2GENE* argument was set to the aggregated gene sets, with all other parameters left at default.

Analyzing senescence and stem cell datasets

The supplementary Table 1 from the SenCID manuscript[124] was used to identify samples from senescent, proliferative, and quiescent cells. Custom scripts were used to identify GSM ids for each sample. This resulted in 308 samples out of 603 which could be downloaded and analyzed. For stem cell datasets, samples were manually annotated. Data was downloaded mainly from GEO using Pasta's functions, *getting_geo_count_mat_for_age_prediction* and *getting_GEO_ES_for_age_model*, as well as with custom scripts. The pasta R package was used to make age predictions. AUC-ROC values were computed using the *auc* function from the pROC R package[125].

Analyzing TCGA data

Histological tumor grades were obtained from Ping et al[126] for samples from BRCA patients and from Liu et al[127] for samples from patients with other cancer types, as described in Zheng et al[54]. Only cancer types with histological tumor grades ranging from G1 and G4 were retained, excluding BLCA samples. The TCGAbiolinks R package was used to download gene expression data and obtain age and overall survival information[128]. Only tumor samples were used for all analyses. The Pearson product-moment correlation coefficient test, implemented in base R's cor.test function, was used to compute p-values and confidence intervals for the correlation between Pasta scores or age in days with tumor grades (from 1 to 4). P-values were adjusted for multiple testing using the False Discovery Rate method implemented via base R's *p.adjust* function with *method = 'fdr'*. Survival analyses were conducted by fitting Cox proportional hazards regression models using the *coxph* function from the R survival package[129]. The predictor variable was either standardized Pasta age scores or standardized age in days. Hazard ratios and their confidence intervals were obtained

588 by exponentiating the model's coefficients and their confidence intervals. Kaplan-Meier
589 survival analyses were performed using the *survfit* function from the R survival package, with
590 Pasta's tertiles as the predictor variable.

Analyzing CMAP L1000 data – Preprocessing data and predicting age scores

Raw level 3 data and metadata files were obtained from the https://clue.io website, and MD5
checksums were verified. Several CMAP plates had incorrect or missing annotations, which
were corrected using custom scripts. Plates with missing treatment information or fewer than
three control samples were excluded from all analyses. The perturbation metadata table was
reduced from 3,027,596 to 3,017,946 entries after filtering. Missing mechanisms of actions
(MOAs) were manually annotated for 60 compounds. Missing compounds names were
manually annotated for seven compounds that had only BRD (Broad) IDs available. All data
were split into individual *.rds* files by plate to facilitate parallel computation with reduced
memory usage. For each sample, its final age score was defined as its raw age score minus the
mean age score of control samples on the same plate. Control samples were defined as
DMSO-treated for compound perturbations and untreated for genetic perturbations.

Analyzing CMAP L1000 data – Annotating senescence- and reprogramming-related
compounds

Senescence-inducing compounds were obtained from the Table 1 of Petrova et al[57].
Reprogramming-inducing compounds were obtained from the Supplementary Table 1 of
Knyazer et al[56]. Due to the large number of compounds listed, only those used in at least 3
studies were selected. The selected compounds were manually checked for their presence in
the CMAP dataset. In total, 31 senescence-inducing compounds and 11 reprogramming-
promoting compounds were identified. The mechanism of actions (MOA) in CMAP for each
of these compounds was identified. MOAs for compounds present in both sets were annotated
as 'both'. This resulted in 5 MOAs for the 'reprogramming' group, 11 MOAs for the
'senescence' group, and 2 MOAs for the 'both' group. All CMAP compounds belonging to
these MOAs were annotated accordingly.

Analyzing CMAP L1000 data – Statistical analysis

Two-sided t-tests were used to assess whether age scores significantly differed from zero for
each perturbation, either across all cell lines or within each individual cell line. All dosages,
incubation times, and BRD IDs for a given perturbation were pooled to increase statistical
power. Bonferroni correction was applied to account for multiple testing. For GPs, a two-
stage approach was employed to mitigate noise associated with variability in genetic construct
quality. In the first stage, significance was tested for each BRD ID. In the second stage,
significance was tested for each GP using only its top two BRD IDs with the highest absolute
average score_pv values. Two metrics were created to combine statistical significance with
effect size: *score_pv = -log10(p-value) * mean(age-score)*, and *score_pa = -log10(adjusted*
*p-value) * mean(age-score)*. Significant results were defined as entries with a score_pa

greater than or equal to -log10(0.05). Over-representation of MOAs among the significant
entries, relative to other entries, was assessed using Fisher's exact test, implemented in base
R's fisher.test function with *alternative = 'greater'*. FDR correction was applied to account
for multiple testing. Analysis of cancer-specific pro-aging compounds was conducted using
one-sided two-sample t-tests to assess whether the mean age in tumor cell lines was greater
than that in normal cell lines. Only compounds with at least two normal and two tumor cell
lines and a mean tumor age effect greater than two were selected. The score_pa and score_pv
metrics were derived as described above but using the difference in mean age effect between
tumor and normal cell lines as the measure of effect size.

Analyzing CMAP L1000 data – Selecting perturbations and cell lines to display in the heatmaps

641 Perturbation and cell line selection was performed separately for the Aging and Rejuvenating
642 groups. The perturbation-by-cell line results tables were sorted by decreasing score_pa for the
643 Rejuvenating group and by increasing score_pa for the Aging group. The top 50 perturbations
644 were selected for each group, which ensured to display the most significant results from our
645 study. For each selected perturbation, the top 5 cell lines were selected.

Analyzing CMAP L1000 data – Pathway enrichment analysis

Significance results from overexpressed genes were inverted to generate a pooled set of GPs
with consistent directional effects per gene. The aging hallmarks gene sets were obtained
from OpenGene[75]. REACTOME gene sets were obtained from the Human MSigDB
Collections[44], using the msigdbr R package[123]. Only REACTOME pathways with 300 genes
or fewer were retained (1,565 out of 1,615). Over-representation analysis was performed
using the *enricher* function from the clusterProfiler R package[122] with the parameter *universe*
set to all the unique genes in the input database.

Analyzing CMAP L1000 data – Analysis of anti-cancer drugs

Anti-cancers drugs were obtained from the CancerDrugs_DB[130] (build date: 28/11/24). We
selected drugs approved from the FDA or the EMA or Euro nationally approved drugs which
kept 311 out of 321 drugs. We then kept 132 drugs that were present in the CMAP L1000
dataset. For each cancer, over-representation of its anti-cancer drugs among the compounds
from a given age set (Aging, Rejuvenating or Other), relative to other compounds, was
assessed using Fisher's exact test, implemented in base R's fisher.test function with
alternative = 'greater'. FDR correction was applied to account for multiple testing.

Analyzing CMAP L1000 data – Identifying consistent compound target / gene perturbation results

To identify significant GPs whose age effects are consistent with the targets of significant
compounds, we computed the mean score_pa for each gene, using the pooled GP significance
table, and for each compound target. Compounds without annotated targets were excluded
from this analysis. For compounds with multiple annotated targets, each target was analyzed
separately. This resulted in mean score_pa values for 7,638 genes and for 868 targets of
compounds. The intersection of these two sets yielded 813 entries, of which 591 had a
consistent directionality of effect and were retained. 47 of these had non-zero score_pa values
in both sets and were retained. Finally, only perturbation pairs with a min score_pa below
0.05 were retained to include only genes significant in both genetic and chemical
perturbations. This resulted in 1 gene for the rejuvenation set (MAP2K1) and 14 genes for the
Aging set.

Analyzing DepMap data – Data sources

DepMap data were obtained using the depmap R package[131]. Cell lines metadata was
retrieved using the *depmap_metadata* function. Drug, CRISPR and RNAi sensitivity data
were obtained using the *drug_sensitivity_21Q2*, *depmap_crispr*, and *depmap_rnai* functions,
respectively. Copy number, gene expression, protein abundance, and mutation data were
obtained using the *depmap_copyNumber*, *depmap_TPM*, *depmap_proteomic*, and
depmap_mutationCalls functions, respectively.

Analyzing DepMap data – Dependency scores

The significance of age scores for compounds was computed for each BRD ID and cell line,
using the same approach as described above (see Statistical analysis). CMAP age score
significance results and DepMap dependency scores were then merged by BRD IDs for
compounds (41,199 overlapping entries), and by gene names for CRISPR (46,752
overlapping entries) and RNAi (27,401 overlapping entries). The associations between
dependency scores and perturbations (compound, CRISPR or RNAi) were assessed by fitting
linear models with dependency score as the dependent variable and mean age score as the
independent variable. The association between dependency scores and gene sets (Mechanism
of actions for compounds and REACTOME pathways for CRISPR) were tested by fitting
linear model with dependency score as the dependent variable and both mean age score and
perturbation name as the independent variables. P-values were adjusted for multiple testing
using the Bonferroni correction. This analysis resulted in the identification of 2 MOAs, 2
compounds, 9 REACTOME pathways and 0 gene perturbations whose mean age scores were
significantly associated with dependency scores at an adjusted p-value significance cutoff of
0.05. The relatively low number of replicates per gene perturbation (median 9, 3^{rd} quartile 12)
could explain the lack of significant entries for this analysis.

Analyzing DepMap data – Computing cell lines' propensity for Aging and Rejuvenation

The compound-by-cell line and GP-by-cell line age score significance tables were merged and
filtered to retain only perturbations with significant age scores across all cell lines. To derive

robust conclusions, the table was further filtered to retain only entries from 19 cell lines in
which at least 25 significant perturbations had been tested for both the Rejuvenation and the
Aging groups. Propensity scores were then computed as the mean age score of all significant
perturbations within both the Rejuvenation and the Aging groups.

713 Analyzing DepMap data – Correlating propensity scores with molecular features

The significance of the association between gene mutation and propensity scores was assessed
using the Pearson product-moment correlation coefficient test, implemented in base R's
cor.test function. P-values were corrected for multiple testing using the Bonferroni method.
Out of 1,724 genes, only TP53 showed a significant association. PCCs between cell line's
propensity for Aging or Rejuvenation and protein expression levels, log-transformed gene
copy numbers, or gene expression (in transcripts per million) were computed for each
molecule (e.g., gene or protein). Molecules with an absolute PCC higher than 0.4 were
analyzed using ORA with the REACTOME database, as described above (see Pathway
enrichment analysis).

723 Cell Culture

A-375 (human melanoma) cells were kindly provided by Dr. Aishe Sarshad (University of
Gothenburg, Sweden) and Dr. Alexander Espinosa (Karolinska Institutet, Sweden). MDA-
MB-231 (human breast cancer) cells were provided by Pr. Kirsty Spalding (Karolinska
Institutet, Sweden). PC3 cells were kindly provided by Dr. Yvonne Ceder (Lund University).
Jurakt cells were purchased from ATCC. Cell lines were maintained in standard DMEM
(Gibco). All media were supplemented with 10% heat-inactivated fetal bovine serum (FBS;
Gibco) and 1% antibiotics (penicillin/streptomycin 100 U/mL; Gibco). Cells were maintained
in a humidified incubator at 37°C and 5% CO_2. Cells were tested monthly for *Mycoplasma*
contamination using MycoAlert[TM] Mycoplasma detection kit (Lonza[TM] LT07-4118-
11630271), and only negative cells were used for experiment. For the assessment of pro-aging
or rejuvenating effects, cells were incubated in the presence of Pralatrexate
(MedChemExpress, #HY-10446, 100 nM in DMSO) or Piperlongumine (MedChemExpress,
#HY-N2329, 10 μM) for 72 hours. The DNA damage-inducing agent doxorubicin (D1515,
Sigma-Aldrich, 100 nM) was used as positive control for senescence induction.

739 SAβG Assay

740 Cells were washed with PBS, fixed with 0.2% glutaraldehyde for 10 minutes, washed with
741 PBS, and incubated overnight at 37°C with a staining solution containing 1 mg/mL X-Gal
742 (Calbiochem, #203782-1GM) prepared in dimethylformamide (DMF; Sigma-Aldrich, D4551)
743 at pH 6. Cells were then washed in PBS and visualized using an Olympus IX73 brightfield
744 microscope.

746 Cell Proliferation analysis

747 Cell proliferation was assessed by EdU incorporation by Click-iT™ Plus EDU Alexa Fluor™
748 647 assay (C10634, Thermo Fisher Scientific) as described by the manufacturer. EdU
749 positivity was assessed by Flow Cytometry using a BD FACSCanto II (BD) at the
750 Biomedicum Flow Core Facility. FACS data were analyzed using the FlowJo software.

Immunoblotting

For immunoblotting, proteins extracted by cellular lysis in RIPA buffer were run on 4–12%
Bis-Tris acrylamide gels (#NP0322, Thermo Fisher Scientific) and electrotransferred to
0.2 μM polyvinylidene fluoride (PVDF) (Biorad,#11704156) using Trans-Blot Turbo
Transfer System (BioRad). Non-specific binding sites were saturated by incubating
membranes for 1 h in 0.05% Tween 20 (#P9416, Sigma Aldrich) v-v in Tris-buffered saline
(TBS) supplemented with 5% BSA (Sigma-Aldrich, #10735078001, w:v in TBS), followed
by an overnight incubation with primary antibodies specific for Waf1/Cip1/CDKN1A p21
(Santa Cruz Biotechnology, #sc-6246), LaminB1 (12987-1-AP, Proteintech) and Oct3/4 (sc-
5279, Santa Cruz Biotechnology). Equal protein loading was monitored with GAPDH (#2118,
Cell Signaling) and alpha-tubulin (#2144, Cell Signaling) specific antibodies). Membranes
were cut in order to allow simultaneous detection of different molecular weight proteins.
Membranes were developed with suitable horseradish peroxidase conjugates followed by
chemiluminescence-based detection with the Cytiva Amersham ECL Prime (#RPN2232, GE
Healthcare) and the Amersham ImageQuant500 software-assisted imager (Cytiva).

Gene Expression Analysis by RT-qPCR

Total RNA was extracted from cell samples using TRIzol reagent (Invitrogen, #15596018)
according to the manufacturer's instructions. Up to 1 μg of total RNA was reverse transcribed
into cDNA using the iScript cDNA Synthesis Kit (Bio-Rad, #1725038) RT-qPCR.
Quantitative real-time PCR was performed using GoTaq PCR Master Mix (Promega,
#A6002) in a QuantStudio 5 Real-Time PCR System (ThermoFisher). The average expression
of ACTB (for sdetection of senescence genes) or GAPDH (for detection of stemness genes)
served as endogenous normalization controls. Primers used in this study are the following:

Gene	Forward	Reverse
CDKN1A	CCGCCCCCTCCTCTAGCTGT	CCCCCATCATATACCCCTAACACA
IL6	TACCCCCAGGAGAAGATTCC	TTTTCTGCCAGTGCCTCTTT
OCT4	CCTGAAGCAGAAGAGGATCACC	AAAGCGGCAGATGGTCGTTTGG
SOX2	GCTACAGCATGATGCAGGACCA	TCTGCGAGCTGGTCATGGAGTT
NANOG	TTTGGAAGCTGCTGGGGAAG	GATGGGAGGAGGGGAGAGGA
GAPDH	AATGAAGGGGTCATTGATGG	AAGGTGAAGGTCGGAGTCAA
ACTB	GTTGTCGACGACGAGCG	GCACAGAGCCTCGCCTT

Data availability

All datasets used to train and evaluate the Pasta model are publicly available from the GTEx Portal (https://gtexportal.org), the Gene Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/), and the Expression Atlas (https://www.ebi.ac.uk/gxa/home). Detailed accession numbers and metadata for each dataset are listed in Supplementary Table S1. The Cancer Cell Line Encyclopedia (CCLE) and DepMap datasets were accessed via the depmap R package (https://depmap.org/portal/download/). The CMAP L1000 perturbation dataset was obtained from the Connectivity Map website (https://clue.io). The Pasta R package, including pre-trained models, functions for data download, processing, and age prediction, is freely available on GitHub at https://github.com/jsalignon/pasta.

Funding

This work was supported by grants from the Swedish Research Council (VR; 2017-06088, 2019-04868, and 2023-04383), the Swedish Cancer Society (Cancerfonden; 20 1034 Pj and 23 2994 Pj), the Novo Nordisk Foundation (NNF21OC0070427 and NNF22OC0078353), Karolinska Institutet (KID2016-00207 and KID2021-00495) and by an ICMC project grant to C.G.R, and by grants from the Swedish Research Council (MH 2023-02156), the Swedish Cancer Society (24 3543 Pj; 23 0690 JIA), Longevity Impetus Grant from Norn Group (First round), Longevity Impetus Grant from Norn Group, Rosenkranz Foundation and Hevolution Foundation (Third Round), Wellcome Leap's Dynamic Resilience Program (jointly funded by Temasek Trust), Ming Wai Lau Centre for Reparative Medicine (MWLC), Karolinska Institutet Senior Researcher Grant to Federico Pietrocola and from Wenner-Gren Stiftelserna (UPD2024-0048) to Federico Pietrocola and Patricia Marques Gonzalez.

Acknowledgements

We thank all members of the Riedel and Pietrocola laboratories for their valuable feedback. We acknowledge the GTEx Consortium, GEO, Expression Atlas, TCGA, CMAP, and DepMap teams for making their transcriptomic and functional genomics datasets publicly available, enabling this research. We thank Sara Hägg (Karolinska Institute) for providing early advice on how to validate our transcriptomic aging clock. We thank Bader Zarrouki (AstraZeneca) for scientific discussions.

Conflicts of interest

The authors declare no conflict of interests.

Author contributions

J.S. performed all computational analyses, developed the R package, generated the computational figures and tables in the manuscript, and wrote the first draft of the manuscript. M.T., P.M.G., E.R.D. conducted wet lab experiments. F.P., M.T. and P.M.G generated wet lab figures and associated methods. J.S., H.A., F.P. and C.G.R. contributed to subsequent editing and revision of the manuscript. J.S., F.P. and C.G.R. conceived and supervised the study.

References

1. López-Otín, C., and Kroemer, G. (2021). Hallmarks of Health. Cell *184*, 33–63. https://doi.org/10.1016/j.cell.2020.11.034.

2. López-Otín, C., Blasco, M.A., Partridge, L., Serrano, M., and Kroemer, G. (2023). Hallmarks of aging: An expanding universe. Cell *186*, 243–278. https://doi.org/10.1016/j.cell.2022.11.001.

3. Rosen, R.S., and Yarmush, M.L. (2023). Current Trends in Anti-Aging Strategies. Annual Review of Biomedical Engineering *25*, 363–385. https://doi.org/10.1146/annurev-bioeng-120122-123054.

4. Min, M., Egli, C., Dulai, A.S., and Sivamani, R.K. (2024). Critical review of aging clocks and factors that may influence the pace of aging. Front. Aging *5*. https://doi.org/10.3389/fragi.2024.1487260.

5. Rutledge, J., Oh, H., and Wyss-Coray, T. (2022). Measuring biological age using omics data. Nat Rev Genet, 1–13. https://doi.org/10.1038/s41576-022-00511-7.

6. Teschendorff, A.E., and Horvath, S. (2025). Epigenetic ageing clocks: statistical methods and emerging computational challenges. Nat Rev Genet, 1–19. https://doi.org/10.1038/s41576-024-00807-w.

7. Bird, A. (2002). DNA methylation patterns and epigenetic memory. Genes Dev. *16*, 6–21. https://doi.org/10.1101/gad.947102.

8. Horvath, S., and Raj, K. (2018). DNA methylation-based biomarkers and the epigenetic clock theory of ageing. Nature Reviews Genetics *19*, 371–384. https://doi.org/10.1038/s41576-018-0004-3.

9. Lachmann, A., Torre, D., Keenan, A.B., Jagodnik, K.M., Lee, H.J., Wang, L., Silverstein, M.C., and Ma'ayan, A. (2018). Massive mining of publicly available RNA-seq data from human and mouse. Nat Commun *9*, 1366. https://doi.org/10.1038/s41467-018-03751-6.

10. CZI Cell Science Program, Abdulla, S., Aevermann, B., Assis, P., Badajoz, S., Bell, S.M., Bezzi, E., Cakir, B., Chaffer, J., Chambers, S., et al. (2025). CZ CELLxGENE Discover: a single-cell data platform for scalable exploration, analysis and modeling of

aggregated data. Nucleic Acids Research *53*, D886–D900.
https://doi.org/10.1093/nar/gkae1142.

11. Youngblut, N.D., Carpenter, C., Prashar, J., Ricci-Tam, C., Ilango, R., Teyssier, N.,
Konermann, S., Hsu, P.D., Dobin, A., Burke, D.P., et al. (2025). scBaseCamp: an AI agent-
curated, uniformly processed, and continually expanding single cell data repository.
Preprint at bioRxiv, https://doi.org/10.1101/2025.02.27.640494
https://doi.org/10.1101/2025.02.27.640494.

12. Xu, Z., Wang, W., Yang, T., Li, L., Ma, X., Chen, J., Wang, J., Huang, Y., Gould, J.,
Lu, H., et al. (2024). STOmicsDB: a comprehensive database for spatial transcriptomics
data sharing, analysis and visualization. Nucleic Acids Research *52*, D1053–D1061.
https://doi.org/10.1093/nar/gkad933.

13. Subramanian, A., Narayan, R., Corsello, S.M., Peck, D.D., Natoli, T.E., Lu, X., Gould,
J., Davis, J.F., Tubelli, A.A., Asiedu, J.K., et al. (2017). A Next Generation Connectivity
Map: L1000 Platform and the First 1,000,000 Profiles. Cell *171*, 1437-1452.e17.
https://doi.org/10.1016/j.cell.2017.10.049.

14. Wei, Z., Si, D., Duan, B., Gao, Y., Yu, Q., Guo, L., and Liu, Q. (2024). PerturBase: a
comprehensive database for single-cell perturbation data analysis and visualization.
Preprint at bioRxiv, https://doi.org/10.1101/2024.02.03.578767
https://doi.org/10.1101/2024.02.03.578767.

15. Zhang, J., Ubas, A.A., Borja, R. de, Svensson, V., Thomas, N., Thakar, N., Lai, I.,
Winters, A., Khan, U., Jones, M.G., et al. (2025). Tahoe-100M: A Giga-Scale Single-Cell
Perturbation Atlas for Context-Dependent Gene Function and Cellular Modeling. Preprint
at bioRxiv, https://doi.org/10.1101/2025.02.20.639398
https://doi.org/10.1101/2025.02.20.639398.

16. Demaria, M. (2025). Cancer treatments accelerate ageing. Nat Rev Cancer, 1–2.
https://doi.org/10.1038/s41568-025-00801-2.

17. Abbott, A. (2024). Stem cells head to the clinic: treatments for cancer, diabetes and
Parkinson's disease could soon be here. Nature *637*, 18–20.
https://doi.org/10.1038/d41586-024-04160-0.

18. Cao, S., and Chen, J. (2025). A speedy cocktail. Nat Chem Biol, 1–2.
https://doi.org/10.1038/s41589-024-01804-0.

19. Paine, P.T., Nguyen, A., and Ocampo, A. (2024). Partial cellular reprogramming: A
deep dive into an emerging rejuvenation technology. Aging Cell *23*, e14039.
https://doi.org/10.1111/acel.14039.

20. Zhang, C., Saurat, N., Cornacchia, D., Chung, S.Y., Sikder, T., Nemchik, A., Minotti,
A., Studer, L., and Betel, D. Identifying Age-Modulating Compounds Using a Novel
Computational Framework for Evaluating Transcriptional Age. Aging Cell *n/a*, e70075.
https://doi.org/10.1111/acel.70075.

21.   Peters, M.J., Joehanes, R., Pilling, L.C., Schurmann, C., Conneely, K.N., Powell, J.,
Reinmaa, E., Sutphin, G.L., Zhernakova, A., Schramm, K., et al. (2015). The
transcriptional landscape of age in human peripheral blood. Nature Communications 6,
8570. https://doi.org/10.1038/ncomms9570.

22.   Mamoshina, P., Volosnikova, M., Ozerov, I.V., Putin, E., Skibina, E., Cortese, F., and
Zhavoronkov, A. (2018). Machine Learning on Human Muscle Transcriptomic Data for
Biomarker Discovery and Tissue-Specific Drug Target Identification. Frontiers in Genetics
9.

23.   Fleischer, J.G., Schulte, R., Tsai, H.H., Tyagi, S., Ibarra, A., Shokhirev, M.N., Huang,
L., Hetzer, M.W., and Navlakha, S. (2018). Predicting age from the transcriptome of
human dermal fibroblasts. Genome Biology 19, 221. https://doi.org/10.1186/s13059-018-
1599-6.

24.   Holzscheck, N., Falckenhayn, C., Söhle, J., Kristof, B., Siegner, R., Werner, A.,
Schössow, J., Jürgens, C., Völzke, H., Wenck, H., et al. (2021). Modeling transcriptomic
age using knowledge-primed artificial neural networks. npj Aging Mech Dis 7, 1–13.
https://doi.org/10.1038/s41514-021-00068-5.

25.   Shokhirev, M.N., and Johnson, A.A. Modeling the human aging transcriptome across
tissues, health status, and sex. Aging Cell n/a, e13280. https://doi.org/10.1111/acel.13280.

26.   Ren, X., and Kuan, P.F. (2020). RNAAgeCalc: A multi-tissue transcriptional age
calculator. PLOS ONE 15, e0237006. https://doi.org/10.1371/journal.pone.0237006.

27.   Jung, S., Arcos Hodar, J., and del Sol, A. (2023). Measuring biological age using a
functionally interpretable multi-tissue RNA clock. Aging Cell 22, e13799.
https://doi.org/10.1111/acel.13799.

28.   Qi, H., Zhao, H., Li, E., Lu, X., Yu, N., Liu, J., and Han, J. DeepQA: A Unified
Transcriptome-Based Aging Clock Using Deep Neural Networks. Aging Cell n/a, e14471.
https://doi.org/10.1111/acel.14471.

29.   Bulteau, R., and Francesconi, M. (2022). Real age prediction from the transcriptome
with RAPToR. Nat Methods 19, 969–975. https://doi.org/10.1038/s41592-022-01540-0.

30.   Zakar-Polyák, E., Csordas, A., Pálovics, R., and Kerepesi, C. (2024). Profiling the
transcriptomic age of single-cells in humans. Commun Biol 7, 1–15.
https://doi.org/10.1038/s42003-024-07094-5.

31.   Schmidt, Y.G., Wu, D., Madan, S., Sinha, S., Sahni, S., Gopalan, V., Wang, B.,
Dhruba, S.R., Schäffer, A.A., Weng, N., et al. (2024). IMMClock reveals immune aging
and T cell function at single-cell resolution. Preprint at bioRxiv,
https://doi.org/10.1101/2024.11.13.623449 https://doi.org/10.1101/2024.11.13.623449.

32.   Li, W., Zhang, Z., Kumar, S., Botey-Bataller, J., Zoodsma, M., Ehsani, A., Zhan, Q.,
Alaswad, A., Zhou, L., Grondman, I., et al. (2025). Single-cell immune aging clocks reveal

924 inter-individual heterogeneity during infection and vaccination. Nat Aging, 1–15.
925 https://doi.org/10.1038/s43587-025-00819-z.

926 33. Zhu, H., Chen, J., Liu, K., Gao, L., Wu, H., Ma, L., Zhou, J., Liu, Z., and Han, J.-D.J.
927 (2023). Human PBMC scRNA-seq–based aging clocks reveal ribosome to inflammation
928 balance as a single-cell aging hallmark and super longevity. Science Advances *9*,
929 eabq7599. https://doi.org/10.1126/sciadv.abq7599.

930 34. THE GTEX CONSORTIUM (2020). The GTEx Consortium atlas of genetic
931 regulatory effects across human tissues. Science *369*, 1318–1330.
932 https://doi.org/10.1126/science.aaz1776.

933 35. Barrett, T., Wilhite, S.E., Ledoux, P., Evangelista, C., Kim, I.F., Tomashevsky, M.,
934 Marshall, K.A., Phillippy, K.H., Sherman, P.M., Holko, M., et al. (2013). NCBI GEO:
935 archive for functional genomics data sets—update. Nucleic Acids Research *41*, D991–
936 D995. https://doi.org/10.1093/nar/gks1193.

937 36. Keays, M. (2023). ExpressionAtlas: Download datasets from EMBL-EBI Expression
938 Atlas https://doi.org/10.18129/B9.bioc.ExpressionAtlas.

939 37. Wetzel, S.J., Ryczko, K., Melko, R.G., and Tamblyn, I. (2022). Twin Neural Network
940 Regression. Applied AI Letters *3*, e78. https://doi.org/10.1002/ail2.78.

941 38. Wetzel, S.J., Melko, R.G., and Tamblyn, I. (2022). Twin neural network regression is
942 a semi-supervised regression algorithm. Mach. Learn.: Sci. Technol. *3*, 045007.
943 https://doi.org/10.1088/2632-2153/ac9885.

944 39. Tynes, M., Gao, W., Burrill, D.J., Batista, E.R., Perez, D., Yang, P., and Lubbers, N.
945 (2021). Pairwise Difference Regression: A Machine Learning Meta-algorithm for
946 Improved Prediction and Uncertainty Quantification in Chemical Search. J. Chem. Inf.
947 Model. *61*, 3846–3857. https://doi.org/10.1021/acs.jcim.1c00670.

948 40. Belaid, M.K., Rabus, M., and Hüllermeier, E. (2024). Pairwise Difference Learning
949 for Classification. Preprint at arXiv, https://doi.org/10.48550/arXiv.2406.20031
950 https://doi.org/10.48550/arXiv.2406.20031.

951 41. Wang, A., Chiou, J., Poirion, O.B., Buchanan, J., Valdez, M.J., Verheyden, J.M., Hou,
952 X., Kudtarkar, P., Narendra, S., Newsome, J.M., et al. (2020). Single-cell multiomic
953 profiling of human lungs reveals cell-type-specific and age-dynamic control of SARS-
954 CoV2 host genes. eLife *9*, e62522. https://doi.org/10.7554/eLife.62522.

955 42. Lukowski, S.W., Lo, C.Y., Sharov, A.A., Nguyen, Q., Fang, L., Hung, S.S., Zhu, L.,
956 Zhang, T., Grünert, U., Nguyen, T., et al. (2019). A single☐cell transcriptome atlas of the
957 adult human retina. The EMBO Journal *38*, e100811.
958 https://doi.org/10.15252/embj.2018100811.

959 43. Solé-Boldo, L., Raddatz, G., Schütz, S., Mallm, J.-P., Rippe, K., Lonsdorf, A.S.,
960 Rodríguez-Paredes, M., and Lyko, F. (2020). Single-cell transcriptomes of the human skin

reveal age-related loss of fibroblast priming. Commun Biol *3*, 1–12.
https://doi.org/10.1038/s42003-020-0922-4.

44.　Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette,
M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set
enrichment analysis: A knowledge-based approach for interpreting genome-wide
expression profiles. Proceedings of the National Academy of Sciences *102*, 15545–15550.
https://doi.org/10.1073/pnas.0506580102.

45.　Stewart-Ornstein, J., Iwamoto, Y., Miller, M.A., Prytyskach, M.A., Ferretti, S.,
Holzer, P., Kallen, J., Furet, P., Jambhekar, A., Forrester, W.C., et al. (2021). p53
dynamics vary between tissues and are linked with radiation sensitivity. Nat Commun *12*,
898. https://doi.org/10.1038/s41467-021-21145-z.

46.　Park, S.-Y., Lee, J.H., Ha, M., Nam, J.-W., and Kim, V.N. (2009). miR-29 miRNAs
activate p53 by targeting p85α and CDC42. Nat Struct Mol Biol *16*, 23–29.
https://doi.org/10.1038/nsmb.1533.

47.　Swahari, V., Nakamura, A., Hollville, E., Hung, Y.-H., Kanke, M., Kurtz, C.L.,
Caravia, X.M., Roiz-Valle, D., He, S., Krishnamurthy, J., et al. (2024). miR-29 is an
important driver of aging-related phenotypes. Commun Biol *7*, 1–11.
https://doi.org/10.1038/s42003-024-06735-z.

48.　Mandic, R., Marquardt, A., Terhorst, P., Ali, U., Nowak-Rossmann, A., Cai, C.,
Rodepeter, F.R., Stiewe, T., Wezorke, B., Wanzel, M., et al. (2022). The importin beta
superfamily member RanBP17 exhibits a role in cell proliferation and is associated with
improved survival of patients with HPV+☐HNSCC. BMC Cancer *22*, 785.
https://doi.org/10.1186/s12885-022-09854-0.

49.　Mertens, J., Paquola, A.C.M., Ku, M., Hatch, E., Böhnke, L., Ladjevardi, S., McGrath,
S., Campbell, B., Lee, H., Herdy, J.R., et al. (2015). Directly Reprogrammed Human
Neurons Retain Aging-Associated Transcriptomic Signatures and Reveal Age-Related
Nucleocytoplasmic Defects. Cell Stem Cell *17*, 705–718.
https://doi.org/10.1016/j.stem.2015.09.001.

50.　Ye, W., Ling, S., Liu, R.-Y., Pan, Z.-Z., Wang, G., Gao, S., Wu, J., Cao, L., Dong, L.,
Li, Y., et al. (2018). Exome sequencing reveals the genetic landscape and frequent
inactivation of PCDHB3 in Chinese rectal cancers. The Journal of Pathology *245*, 222–
234. https://doi.org/10.1002/path.5073.

51.　Srivatsan, S.R., McFaline-Figueroa, J.L., Ramani, V., Saunders, L., Cao, J., Packer, J.,
Pliner, H.A., Jackson, D.L., Daza, R.M., Christiansen, L., et al. (2020). Massively
multiplex chemical transcriptomics at single-cell resolution. Science *367*, 45–51.
https://doi.org/10.1126/science.aax6234.

52.　Wang, L., Leite de Oliveira, R., Wang, C., Fernandes Neto, J.M., Mainardi, S., Evers,
B., Lieftink, C., Morris, B., Jochems, F., Willemsen, L., et al. (2017). High-Throughput
Functional Genetic and Compound Screens Identify Targets for Senescence Induction in
Cancer. Cell Reports *21*, 773–783. https://doi.org/10.1016/j.celrep.2017.09.085.

53.   Malta, T.M., Sokolov, A., Gentles, A.J., Burzykowski, T., Poisson, L., Weinstein,
J.N., Kamińska, B., Huelsken, J., Omberg, L., Gevaert, O., et al. (2018). Machine Learning
Identifies Stemness Features Associated with Oncogenic Dedifferentiation. Cell *173*, 338-
354.e15. https://doi.org/10.1016/j.cell.2018.03.034.

54.   Zheng, H., Song, K., Fu, Y., You, T., Yang, J., Guo, W., Wang, K., Jin, L., Gu, Y., Qi,
L., et al. (2021). An absolute human stemness index associated with oncogenic
dedifferentiation.   Briefings   in   Bioinformatics   *22*,   2151–2160.
https://doi.org/10.1093/bib/bbz174.

55.   Park, S.S., Choi, Y.W., Kim, J.-H., Kim, H.S., and Park, T.J. (2021). Senescent tumor
cells: an overlooked adversary in the battle against cancer. Exp Mol Med *53*, 1834–1841.
https://doi.org/10.1038/s12276-021-00717-5.

56.   Knyazer, A., Bunu, G., Toren, D., Mracica, T.B., Segev, Y., Wolfson, M., Muradian,
K.K., Tacutu, R., and Fraifeld, V.E. (2021). Small molecules for cell reprogramming: a
systems   biology   analysis.   Aging   (Albany   NY)   *13*,   25739–25762.
https://doi.org/10.18632/aging.203791.

57.   Petrova, N.V., Velichko, A.K., Razin, S.V., and Kantidze, O.L. (2016). Small
molecule compounds that induce cellular senescence. Aging Cell *15*, 999–1017.
https://doi.org/10.1111/acel.12518.

58.   Huangfu, D., Maehr, R., Guo, W., Eijkelenboom, A., Snitow, M., Chen, A.E., and
Melton, D.A. (2008). Induction of pluripotent stem cells by defined factors is greatly
improved   by   small-molecule   compounds.   Nat   Biotechnol   *26*,   795–797.
https://doi.org/10.1038/nbt1418.

59.   Anokye-Danso, F., Trivedi, C.M., Juhr, D., Gupta, M., Cui, Z., Tian, Y., Zhang, Y.,
Yang, W., Gruber, P.J., Epstein, J.A., et al. (2011). Highly Efficient miRNA-Mediated
Reprogramming of Mouse and Human Somatic Cells to Pluripotency. Cell Stem Cell *8*,
376–388. https://doi.org/10.1016/j.stem.2011.03.001.

60.   Di Bernardo, G., Squillaro, T., Dell'Aversana, C., Miceli, M., Cipollaro, M., Cascino,
A., Altucci, L., and Galderisi, U. (2009). Histone Deacetylase Inhibitors Promote
Apoptosis and Senescence in Human Mesenchymal Stem Cells. Stem Cells and
Development *18*, 573–582. https://doi.org/10.1089/scd.2008.0172.

61.   Li, X.-N., Shu, Q., Su, J.M.-F., Perlaky, L., Blaney, S.M., and Lau, C.C. (2005).
Valproic acid induces growth arrest, apoptosis, and senescence in medulloblastomas by
increasing histone hyperacetylation and regulating expression of p21Cip1, CDK4, and
CMYC. Molecular Cancer Therapeutics *4*, 1912–1922. https://doi.org/10.1158/1535-
7163.MCT-05-0184.

62.   An, H.-M., Xue, Y.-F., Shen, Y.-L., Du, Q., and Hu, B. (2013). Sodium Valproate
Induces Cell Senescence in Human Hepatocarcinoma Cells. Molecules *18*, 14935–14947.
https://doi.org/10.3390/molecules181214935.

63.   Rad, A.N., and Grillari, J. (2024). Current senolytics: Mode of action, efficacy and
limitations, and their future. Mechanisms of Ageing and Development *217*, 111888.
https://doi.org/10.1016/j.mad.2023.111888.

64.   Wang, Y., Wang, X., Liu, L., and Wang, X. (2009). HDAC inhibitor trichostatin A-
inhibited survival of dopaminergic neuronal cells. Neuroscience Letters *467*, 212–216.
https://doi.org/10.1016/j.neulet.2009.10.037.

65.   Tsherniak, A., Vazquez, F., Montgomery, P.G., Weir, B.A., Kryukov, G., Cowley,
G.S., Gill, S., Harrington, W.F., Pantel, S., Krill-Burger, J.M., et al. (2017). Defining a
Cancer   Dependency   Map.   Cell   *170*,   564-576.e16.
https://doi.org/10.1016/j.cell.2017.06.010.

66.   Huang, Y., Che, X., Wang, P.W., and Qu, X. (2024). p53/MDM2 signaling pathway in
aging, senescence and tumorigenesis. Seminars in Cancer Biology *101*, 44–57.
https://doi.org/10.1016/j.semcancer.2024.05.001.

67.   Xu, S., Wu, W., Huang, H., Huang, R., Xie, L., Su, A., Liu, S., Zheng, R., Yuan, Y.,
Zheng, H., et al. (2019). The p53/miRNAs/Ccna2 pathway serves as a novel regulator of
cellular senescence: Complement of the canonical p53/p21 pathway. Aging Cell *18*,
e12918. https://doi.org/10.1111/acel.12918.

68.   Guney, I., Wu, S., and Sedivy, J.M. (2006). Reduced c-Myc signaling triggers
telomere-independent senescence by regulating Bmi-1 and p16INK4a. Proceedings of the
National Academy of Sciences *103*, 3645–3650. https://doi.org/10.1073/pnas.0600069103.

69.   Fang, L., Li, D., Yin, J., Pan, H., Ye, H., Bowman, J., Capaldo, B., and Kelly, K.
(2022). TMPRSS2-ERG promotes the initiation of prostate cancer by suppressing
oncogene-induced   senescence.   Cancer   Gene   Ther   *29*,   1463–1476.
https://doi.org/10.1038/s41417-022-00454-5.

70.   Zhu, H., Blake, S., Kusuma, F.K., Pearson, R.B., Kang, J., and Chan, K.T. (2020).
Oncogene-induced senescence: From biology to therapy. Mechanisms of Ageing and
Development *187*, 111229. https://doi.org/10.1016/j.mad.2020.111229.

71.   Zhuang, D., Mannava, S., Grachtchouk, V., Tang, W.-H., Patil, S., Wawrzyniak, J.A.,
Berman, A.E., Giordano, T.J., Prochownik, E.V., Soengas, M.S., et al. (2008). C-MYC
overexpression is required for continuous suppression of oncogene-induced senescence in
melanoma cells. Oncogene *27*, 6623–6634. https://doi.org/10.1038/onc.2008.258.

72.   Balcerek, J., Jiang, J., Li, Y., Jiang, Q., Holdreith, N., Singh, B., Chandra, V., Lv, K.,
Ren, J., Rozenova, K., et al. (2018). Lnk/Sh2b3 deficiency restores hematopoietic stem cell
function and genome integrity in Fancd2 deficient Fanconi anemia. Nat Commun *9*, 3915.
https://doi.org/10.1038/s41467-018-06380-1.

73.   Song, H., Wang, J., Wang, X., Yuan, B., Li, D., Hu, A., Guo, Y., Cai, S., Jin, S., Zhou,
Y., et al. (2022). HNF4A-AS1-encoded small peptide promotes self-renewal and
aggressiveness of neuroblastoma stem cells via eEF1A1-repressed SMAD4 transactivation.
Oncogene *41*, 2505–2519. https://doi.org/10.1038/s41388-022-02271-4.

74.   Baugh, E.H., Ke, H., Levine, A.J., Bonneau, R.A., and Chan, C.S. (2018). Why are
there hotspot mutations in the TP53 gene in human cancers? Cell Death Differ 25, 154–
160. https://doi.org/10.1038/cdd.2017.180.

75.   Rafikova, E., Nemirovich-Danchenko, N., Ogmen, A., Parfenenkova, A., Velikanova,
A., Tikhonov, S., Peshkin, L., Rafikov, K., Spiridonova, O., Belova, Y., et al. (2024). Open
Genes—a new comprehensive database of human genes associated with aging and
longevity. Nucleic Acids Research 52, D950–D962. https://doi.org/10.1093/nar/gkad712.

76.   Milacic, M., Beavers, D., Conley, P., Gong, C., Gillespie, M., Griss, J., Haw, R.,
Jassal, B., Matthews, L., May, B., et al. (2024). The Reactome Pathway Knowledgebase
2024. Nucleic Acids Research 52, D672–D678. https://doi.org/10.1093/nar/gkad1025.

77.   Sagara, A., Miura, S., Kobinata, A., Naganawa, R., Yaginuma, S., Saito, S., Saito, R.,
Kominato, H., Yumoto, T., and Sato, F. (2023). COL8A1 enhances the invasion/metastasis
in MDA-MB-231 cells via the induction of *IL1B* and *MMP1* expression. Biochemical and
Biophysical   Research   Communications   *642*,   145–153.
https://doi.org/10.1016/j.bbrc.2022.12.046.

78.   Sugrue, M.M., Shin, D.Y., Lee, S.W., and Aaronson, S.A. (1997). Wild-type p53
triggers a rapid senescence program in human tumor cells lacking☐functional☐p53.
Proceedings   of   the   National   Academy   of   Sciences   *94*,   9648–9653.
https://doi.org/10.1073/pnas.94.18.9648.

79.   Rufini, A., Tucci, P., Celardo, I., and Melino, G. (2013). Senescence and aging: the
critical roles of p53. Oncogene *32*, 5129–5143. https://doi.org/10.1038/onc.2012.640.

80.   Jiang, Z., Zhu, M., Zhang, L., Cui, H., Jiang, R., and Yang, Y. (2024). Antitumor
immunity and prognosis value elicited by FAT3 and LRP1B co-mutation in endometrial
cancer. Gynecologic Oncology *187*, 1–11. https://doi.org/10.1016/j.ygyno.2024.04.023.

81.   Schneider, T., Martinez-Martinez, A., Cubillos-Rojas, M., Bartrons, R., Ventura, F.,
and Rosa, J.L. (2019). Large HERCs Function as Tumor Suppressors. Front. Oncol. *9*.
https://doi.org/10.3389/fonc.2019.00524.

82.   Namasu, C.Y., Katzerke, C., Bräuer-Hartmann, D., Wurm, A.A., Gerloff, D.,
Hartmann, J.-U., Schwind, S., Müller-Tidow, C., Hilger, N., Fricke, S., et al. (2017). ABR,
a novel inducer of transcription factor C/EBPα, contributes to myeloid differentiation and
is a favorable prognostic factor in acute myeloid leukemia. Oncotarget 8, 103626–103639.
https://doi.org/10.18632/oncotarget.22093.

83.   Full article: Role of Rho guanine nucleotide exchange factors in non-small cell lung
cancer
https://www.tandfonline.com/doi/full/10.1080/21655979.2021.2006519?rfr_dat=cr_pub++
0pubmed&url_ver=Z39.88-2003&rfr_id=ori%3Arid%3Acrossref.org.

84.   Jiang, X., Baig, A.H., Palazzo, G., Del Pizzo, R., Bortecen, T., Groessl, S., Zaal, E.A.,
Amaya Ramirez, C.C., Kowar, A., Aviles-Huerta, D., et al. (2024). P53-dependent

hypusination of eIF5A affects mitochondrial translation and senescence immune surveillance. Nat Commun *15*, 7458. https://doi.org/10.1038/s41467-024-51901-w.

85. Wang, B., Han, J., Elisseeff, J.H., and Demaria, M. (2024). The senescence-associated secretory phenotype and its physiological and pathological implications. Nat Rev Mol Cell Biol *25*, 958–978. https://doi.org/10.1038/s41580-024-00727-x.

86. Yu, Q., Katlinskaya, Y.V., Carbone, C.J., Zhao, B., Katlinski, K.V., Zheng, H., Guha, M., Li, N., Chen, Q., Yang, T., et al. (2015). DNA-Damage-Induced Type I Interferon Promotes Senescence and Inhibits Stem Cell Function. Cell Reports *11*, 785–797. https://doi.org/10.1016/j.celrep.2015.03.069.

87. Katlinskaya, Y.V., Katlinski, K.V., Yu, Q., Ortiz, A., Beiting, D.P., Brice, A., Davar, D., Sanders, C., Kirkwood, J.M., Rui, H., et al. (2016). Suppression of Type I Interferon Signaling Overcomes Oncogene-Induced Senescence and Mediates Melanoma Development and Progression. Cell Reports *15*, 171–180. https://doi.org/10.1016/j.celrep.2016.03.006.

88. Sun, E.D., Zhou, O.Y., Hauptschein, M., Rappoport, N., Xu, L., Navarro Negredo, P., Liu, L., Rando, T.A., Zou, J., and Brunet, A. (2024). Spatial transcriptomic clocks reveal cell proximity effects in brain ageing. Nature, 1–12. https://doi.org/10.1038/s41586-024-08334-8.

89. Pereira, C.F., Piccolo, F.M., Tsubouchi, T., Sauer, S., Ryan, N.K., Bruno, L., Landeira, D., Santos, J., Banito, A., Gil, J., et al. (2010). ESCs Require PRC2 to Direct the Successful Reprogramming of Differentiated Cells toward Pluripotency. Cell Stem Cell *6*, 547–556. https://doi.org/10.1016/j.stem.2010.04.013.

90. Moqri, M., Cipriano, A., Simpson, D.J., Rasouli, S., Murty, T., de Jong, T.A., Nachun, D., de Sena Brandine, G., Ying, K., Tarkhov, A., et al. (2024). PRC2-AgeIndex as a universal biomarker of aging and rejuvenation. Nat Commun *15*, 5956. https://doi.org/10.1038/s41467-024-50098-2.

91. Koldobskiy, M.A., Camacho, O., Reddy, P., Belmonte, J.C.I., and Feinberg, A.P. (2023). Convergence of aging- and rejuvenation-related epigenetic alterations on PRC2 targets. Preprint at bioRxiv, https://doi.org/10.1101/2023.06.08.544045 https://doi.org/10.1101/2023.06.08.544045.

92. Salignon, J., Millan-Ariño, L., Garcia, M.U., and Riedel, C.G. (2024). Cactus: A user-friendly and reproducible ATAC-Seq and mRNA-Seq analysis pipeline for data preprocessing, differential analysis, and enrichment analysis. Genomics *116*, 110858. https://doi.org/10.1016/j.ygeno.2024.110858.

93. Seidel, E., and von Karstedt, S. (2022). Extrinsic cell death pathway plasticity: a driver of clonal evolution in cancer? Cell Death Discov. *8*, 1–6. https://doi.org/10.1038/s41420-022-01251-7.

94. Li, H., Collado, M., Villasante, A., Strati, K., Ortega, S., Cañamero, M., Blasco, M.A., and Serrano, M. (2009). The Ink4/Arf locus is a barrier for iPS cell reprogramming. Nature *460*, 1136–1139. https://doi.org/10.1038/nature08290.

95. Rufini, A., Tucci, P., Celardo, I., and Melino, G. (2013). Senescence and aging: the critical roles of p53. Oncogene *32*, 5129–5143. https://doi.org/10.1038/onc.2012.640.

96. Han, J.-D.J. (2024). The ticking of aging clocks. Trends in Endocrinology & Metabolism *35*, 11–22. https://doi.org/10.1016/j.tem.2023.09.007.

97. Wagner, W. (2019). The Link Between Epigenetic Clocks for Aging and Senescence. Front Genet *10*, 303. https://doi.org/10.3389/fgene.2019.00303.

98. Kremer, L.P.M., Cerrizuela, S., El-Sammak, H., Al Shukairi, M.E., Ellinger, T., Straub, J., Korkmaz, A., Volk, K., Brunken, J., Kleber, S., et al. (2024). DNA methylation controls stemness of astrocytes in health and ischaemia. Nature *634*, 415–423. https://doi.org/10.1038/s41586-024-07898-9.

99. Pedrosa, P., Zhang, Z., Nuñez-Quintela, V., Macias, D., Ge, J., Denholm, M., Dyas, A., Estevez-Souto, V., Lado-Fernandez, P., Gonzalez, P., et al. (2024). Inhibition of lung tumorigenesis by transient reprogramming in cancer cells. Cell Death Dis *15*, 1–16. https://doi.org/10.1038/s41419-024-07207-2.

100. Selenina, A.V., Tsimokha, A.S., and Tomilin, A.N. (2017). Proteasomes in Protein Homeostasis of Pluripotent Stem Cells. Acta Naturae *9*, 39–47.

101. Floyd, E.Z., Staszkiewicz, J., Power, R.A., Kilroy, G., Kirk-Ballard, H., Barnes, C.W., Strickler, K.L., Rim, J.S., Harkins, L.L., Gao, R., et al. (2015). Prolonged Proteasome Inhibition Cyclically Upregulates Oct3/4 and Nanog Gene Expression, but Reduces Induced Pluripotent Stem Cell Colony Formation. Cell Reprogram *17*, 95–105. https://doi.org/10.1089/cell.2014.0030.

102. Singh, P.B., and Zhakupova, A. (2022). Age reprogramming: cell rejuvenation by partial reprogramming. Development *149*, dev200755. https://doi.org/10.1242/dev.200755.

103. Tyshkovskiy, A., Kholdina, D., Ying, K., Davitadze, M., Molière, A., Tongu, Y., Kasahara, T., Kats, L.M., Vladimirova, A., Moldakozhayev, A., et al. (2024). Transcriptomic Hallmarks of Mortality Reveal Universal and Specific Mechanisms of Aging, Chronic Disease, and Rejuvenation. Preprint at bioRxiv, https://doi.org/10.1101/2024.07.04.601982 https://doi.org/10.1101/2024.07.04.601982.

104. Huang, Q., Li, F., Liu, X., Li, W., Shi, W., Liu, F.-F., O'Sullivan, B., He, Z., Peng, Y., Tan, A.-C., et al. (2011). Caspase 3–mediated stimulation of tumor cell repopulation during cancer radiotherapy. Nat Med *17*, 860–866. https://doi.org/10.1038/nm.2385.

105. Haak, V.M., Huang, S., and Panigrahy, D. (2021). Debris-stimulated tumor growth: a Pandora's box? Cancer Metastasis Rev *40*, 791–801. https://doi.org/10.1007/s10555-021-09998-8.

1190   106.   Pereira, B., Correia, F.P., Alves, I.A., Costa, M., Gameiro, M., Martins, A.P., and
1191   Saraiva, J.A. (2024). Epigenetic reprogramming as a key to reverse ageing and increase
1192   longevity.   Ageing   Research   Reviews   *95*,   102204.
1193   https://doi.org/10.1016/j.arr.2024.102204.

1194   107.   Hattangadi, D.K., DeMasters, G.A., Walker, T.D., Jones, K.R., Di, X., Newsham, I.F.,
1195   and Gewirtz, D.A. (2004). Influence of p53 and caspase 3 activity on cell death and
1196   senescence in response to methotrexate in the breast tumor cell. Biochemical
1197   Pharmacology *68*, 1699–1708. https://doi.org/10.1016/j.bcp.2004.06.033.

1198   108.   Dabrowska, M., Mosieniak, G., Skierski, J., Sikora, E., and Rode, W. (2009).
1199   Methotrexate-induced senescence in human adenocarcinoma cells is accompanied by
1200   induction of p21waf1/cip1 expression and lack of polyploidy. Cancer Letters *284*, 95–101.
1201   https://doi.org/10.1016/j.canlet.2009.04.015.

1202   109.   Csik, B., Vali Kordestan, K., Gulej, R., Patai, R., Nyul-Toth, A., Shanmugarama, S.,
1203   Mukli, P., Ungvari, A., Balsara, K.E., McNall, R.Y., et al. (2025). Cisplatin and
1204   methotrexate induce brain microvascular endothelial and microglial senescence in mouse
1205   models   of   chemotherapy-associated   cognitive   impairment.   GeroScience.
1206   https://doi.org/10.1007/s11357-025-01560-6.

1207   110.   Chen, Y., Zhang, C., Jin, S., Li, J., Dai, J., Zhang, Z., and Guo, R. (2023). Pemetrexed
1208   induces ROS generation and cellular senescence by attenuating TS-mediated thymidylate
1209   metabolism to reverse gefitinib resistance in NSCLC. Journal of Cellular and Molecular
1210   Medicine *27*, 2032–2044. https://doi.org/10.1111/jcmm.17799.

1211   111.   Dong, H., Hu, L., Li, W., Shi, M., He, L., Wang, C., Hu, Y., Wang, H., Wen, C., Liu,
1212   H., et al. (2022). Pyrimethamine inhibits cell growth by inducing cell senescence and
1213   boosting CD8+ T-cell mediated cytotoxicity in colorectal cancer. Mol Biol Rep *49*, 4281–
1214   4292. https://doi.org/10.1007/s11033-022-07262-y.

1215   112.   O'Connor, O.A., Amengual ,Jennifer, Colbourn ,Donald, Deng ,Changchun, and and
1216   Sawas, A. (2017). Pralatrexate: a comprehensive update on pharmacology, clinical activity
1217   and   strategies   to   optimize   use.   Leukemia   &   Lymphoma   *58*,   2548–2557.
1218   https://doi.org/10.1080/10428194.2017.1306642.

1219   113.   Sun, N., Youle, R.J., and Finkel, T. (2016). The Mitochondrial Basis of Aging.
1220   Molecular Cell *61*, 654–666. https://doi.org/10.1016/j.molcel.2016.01.028.

1221   114.   Abu Shelbayeh, O., Arroum, T., Morris, S., and Busch, K.B. (2023). PGC-1α Is a
1222   Master Regulator of Mitochondrial Lifecycle and ROS Stress Response. Antioxidants
1223   (Basel) *12*, 1075. https://doi.org/10.3390/antiox12051075.

1224   115.   Macip, S., Igarashi, M., Fang, L., Chen, A., Pan, Z., Lee, S.W., and Aaronson, S.A.
1225   (2002). Inhibition of p21□mediated ROS accumulation can rescue p21□induced
1226   senescence. The EMBO Journal *21*, 2180–2188. https://doi.org/10.1093/emboj/21.9.2180.

1227   116.   Frisch, S.M., and MacFawn, I.P. (2020). Type I interferons and related pathways in
1228   cell senescence. Aging Cell *19*, e13234. https://doi.org/10.1111/acel.13234.

117. Ji, S., Xiong, M., Chen, H., Liu, Y., Zhou, L., Hong, Y., Wang, M., Wang, C., Fu, X.,
and Sun, X. (2023). Cellular rejuvenation: molecular mechanisms and potential therapeutic
interventions for diseases. Sig Transduct Target Ther *8*, 1–39.
https://doi.org/10.1038/s41392-023-01343-5.

118. Davis, S., and Meltzer, P.S. (2007). GEOquery: a bridge between the Gene Expression
Omnibus (GEO) and BioConductor. Bioinformatics *23*, 1846–1847.
https://doi.org/10.1093/bioinformatics/btm254.

119. George, N., Fexova, S., Fuentes, A.M., Madrigal, P., Bi, Y., Iqbal, H., Kumbham, U.,
Nolte, N.F., Zhao, L., Thanki, A.S., et al. (2024). Expression Atlas update: insights from
sequencing data at both bulk and single cell level. Nucleic Acids Research *52*, D107–
D114. https://doi.org/10.1093/nar/gkad1021.

120. Durinck, S., Spellman, P.T., Birney, E., and Huber, W. (2009). Mapping identifiers for
the integration of genomic datasets with the R/Bioconductor package biomaRt. Nature
Protocols *4*, 1184–1191.

121. Friedman, J.H., Hastie, T., and Tibshirani, R. (2010). Regularization Paths for
Generalized Linear Models via Coordinate Descent. Journal of Statistical Software *33*, 1–
22. https://doi.org/10.18637/jss.v033.i01.

122. Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W.,
Zhan, L., et al. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting
omics data. The Innovation *2*, 100141. https://doi.org/10.1016/j.xinn.2021.100141.

123. Dolgalev, I. (2022). msigdbr: MSigDB Gene Sets for Multiple Organisms in a Tidy
Data Format.

124. Tao, W., Yu, Z., and Han, J.-D.J. (2024). Single-cell senescence identification reveals
senescence heterogeneity, trajectory, and modulators. Cell Metabolism *36*, 1126-1143.e5.
https://doi.org/10.1016/j.cmet.2024.03.009.

125. Robin, X., Turck, N., Hainard, A., Tiberti, N., Lisacek, F., Sanchez, J.-C., and Müller,
M. (2011). pROC: an open-source package for R and S+ to analyze and compare ROC
curves. BMC Bioinformatics *12*, 77. https://doi.org/10.1186/1471-2105-12-77.

126. Ping, Z., Xia, Y., Shen, T., Parekh, V., Siegal, G.P., Eltoum, I.-E., He, J., Chen, D.,
Deng, M., Xi, R., et al. (2016). A microscopic landscape of the invasive breast cancer
genome. Sci Rep *6*, 27545. https://doi.org/10.1038/srep27545.

127. Liu, J., Lichtenberg, T., Hoadley, K.A., Poisson, L.M., Lazar, A.J., Cherniack, A.D.,
Kovatich, A.J., Benz, C.C., Levine, D.A., Lee, A.V., et al. (2018). An Integrated TCGA
Pan-Cancer Clinical Data Resource to Drive High-Quality Survival Outcome Analytics.
Cell *173*, 400-416.e11. https://doi.org/10.1016/j.cell.2018.02.052.

128. Colaprico, A., Silva, T.C., Olsen, C., Garofano, L., Cava, C., Garolini, D., Sabedot, T.,
Malta, T.M., Pagnotta, S.M., Castiglioni, I., et al. (2015). TCGAbiolinks: An

R/Bioconductor package for integrative analysis of TCGA data. Nucleic Acids Research.
https://doi.org/10.1093/nar/gkv1507.

129.   Therneau, T.M. (2024). A Package for Survival Analysis in R.

130.   Pantziarka, P., Capistrano I, R., De Potter, A., Vandeborne, L., and Bouche, G. (2021).
An Open Access Database of Licensed Cancer Drugs. Front Pharmacol *12*, 627574.
https://doi.org/10.3389/fphar.2021.627574.

131.   Gatto, L., and Killian, T. (2023). depmap: Cancer Dependency Map Data Package
https://doi.org/10.18129/B9.bioc.depmap.

132.   Kim, Y.-M., Byun, H.-O., Jee, B.A., Cho, H., Seo, Y.-H., Kim, Y.-S., Park, M.H.,
Chung, H.-Y., Woo, H.G., and Yoon, G. (2013). Implications of time-series gene
expression profiles of replicative senescence. Aging Cell *12*, 622–634.
https://doi.org/10.1111/acel.12087.

133.   Liu, X., Ouyang, J.F., Rossello, F.J., Tan, J.P., Davidson, K.C., Valdes, D.S.,
Schröder, J., Sun, Y.B.Y., Chen, J., Knaupp, A.S., et al. (2020). Reprogramming roadmap
reveals route to human induced trophoblast stem cells. Nature *586*, 101–107.
https://doi.org/10.1038/s41586-020-2734-6.

134.   Li, Q., Hutchins, A.P., Chen, Y., Li, S., Shan, Y., Liao, B., Zheng, D., Shi, X., Li, Y.,
Chan, W.-Y., et al. (2017). A sequential EMT-MET mechanism drives the differentiation
of human embryonic stem cells towards hepatocytes. Nat Commun *8*, 15166.
https://doi.org/10.1038/ncomms15166.

1286

1287

1288

Figure legends

Figure 1. The age-shift classifier is a novel modeling approach that enables cross-platform aging clocks.

(**a**) Overview of the evaluated modelling strategies. Models were trained and tested exclusively on samples from healthy donors. (**b-d**) Schematic diagrams of the different modeling strategies. (**e**) Comparative performance in predicting relative age, assessed with leave-one-dataset-out cross-validation. All samples were rank-transformed except for REG_RC and AS40_RC where raw counts ('RC') were used. The y-axis shows the Pearson Correlation Coefficient (PCC) between chronological ages and predicted age scores in the held-out datasets. (**f-h**) Age predictions for scRNA-Seq aging atlases of the retina[42] (**f**), skin[43] (**g**), and lung[41] (**h**). Abbreviations: REG, regression; AS30/40/50, age-shift classifier with 30/40/50 years of age difference, TC_30_70|TC_40_60, traditional classifiers with young cutoffs at 30/40 years and old cutoffs at 70/60 years, RC: raw counts.

1302

Figure 2. Pasta is a state-of-the-art, multi-tissue relative age-prediction model whose output is predominantly driven by p53-linked signals.

(**a-b**) Pearson Correlation Coefficient (PCC) of age scores for Pasta and MultiTIMER vs true ages for healthy individuals. (**a**) Results from the leave-one-dataset-out analysis. (**b**) Results on external datasets. (**c**) Gene-Set Enrichment Analysis (GSEA) of the coefficients of the Pasta model, using the REACTOME, CPG, KEGG and MIRDB gene sets from the Human MSigDB Collections. (**d-e**) Top 10 most positive (**d**) and negative (**e**) coefficients.

1310

Figure 3. Pasta identifies senescent and stem cells and stratifies tumor grade and patient overall survival.

(**a-b**) Pasta's ability to discriminate between senescent and proliferative (**a**) or quiescent (**b**) cells, assessed using AUC-ROC (Area Under the Curve of the Receiver Operating Characteristic), across multiple RNA-Seq datasets. (**c**) Pasta's predictions on a microarray senescence time course dataset[132]. (**d**) Pasta was applied to pseudobulks from a SciPlex dataset generated by exposing the A549[51] cell line to 138 drugs, and the resulting age scores were evaluated for their capacity to recover senescence-inducing drugs, as reported in Wang et al 2017[52]. (**e-f**) Pasta's ability to discriminate between stem cells (**e**) or induced-pluripotent stem cells (iPSC) (**f**) and fibroblasts, assessed by AUC-ROC. (**g**) Pasta's age scores on a bulk RNA-Seq reprogramming time course dataset[133]. (**h**) Pasta's age scores on a bulk RNA-Seq liver differentiation time course dataset[134]. (**i**) Correlation of patient ages (red) and Pasta's age scores (blue) with tumor grade. (**j-k**) Boxplots of Pasta's age scores for breast invasive carcinoma (**j**) and kidney renal clear cell carcinoma (**k**) patients, grouped by tumor grade. (**l**) Hazard ratios from Cox proportional-hazards regression model fitted for scaled ages (red) and scaled age scores (blue). (**m-n**) Kaplan-Meier survival estimates for adrenocortical carcinoma (**m**) and thymoma (**n**) patients, grouped by age score tertiles.

1328

Figure 4. Pasta identifies age-modulatory compounds.

(**a**) Volcano plot showing the most significant Rejuvenating (negative age score) and Aging (positive age score) compounds. Points in yellow, blue, and purple indicate compounds from drug classes that induce senescence[57], promote reprogramming[56], or do both, respectively. For display purposes, p-values of zero were set to 10^{-300}. (**b-d**) Proportion of compounds annotated as senescence-inducing (**b**), pro-reprogramming (**c**), or both (**d**) for significantly age-increasing ('Aging'), age-decreasing ('Rejuvenating') compounds, and other compounds ('Other'). (**e-f**) All significantly enriched Mechanisms Of Actions (MOAs) for the Aging (**e**) and Rejuvenating (**f**) compounds. Colors are defined as in (a). The grey lines show the 0.05 adjusted p-value cutoff. (**g**) Heatmap of compounds by cell line with the most significant age scores (see Methods). Colors indicate the square root of a metric balancing significance and effect size: *-log10(adjusted p-value) * mean(age score)*. Missing values are shown in grey. (**h**) Boxplots and violin plots of dependency scores for compounds by cell line in the Aging (175 entries), Rejuvenating (75 entries) and Other (40,906 entries) groups. (**i-j**) MOAs (**i**) and compounds (**j**) with the strongest association between age score and dependency. The yellow

1344 and blue colors indicate a negative and positive Pearson correlation coefficient between age
1345 score and dependency, respectively. (**k**) Mean age score vs dependency for MDM inhibitors.
1346 Red and blue points show results for AMG-232 and CGM-097, respectively, across different
1347 cell lines.

1348

Figure 5. Pasta can prioritize novel anti-cancer drugs with potentially diminished side-
effects.

(**a**) Cancers whose anti-cancer drugs are significantly enriched in age-modulatory compounds
across all cell lines, with yellow and blue colors indicating enrichment in the Aging and
Rejuvenating sets, respectively. Only cancers with at least 5 approved anti-cancer drugs
according to CancerDrugs_DB[130] were selected. 17 out 27 cancers were significant for the
Aging group. (**b**) Top 10 compounds whose age score is statistically higher in cancer cell
lines than in normal cell lines (see Methods). (**c**) Mean age score in cancer and normal cell
lines for the 10 compounds shown in (b).

1358

Figure 6. Experimental validation of pralatrexate as a senescence-inducing compound
and piperlongumine as a rejuvenating compound.

(**a**) Representative images from Senescence Associated Beta Galactosidase assay performed
on A-375 cells exposed to Pralatrexate (Prltx). Doxorubicin (Doxo) was used as positive
control for senescence induction. Scale Bar: 10 µm. (**b**) qPCR analysis of CDKN1A and IL6
mRNA levels in A-375 cells in response to Prtlx and Doxo.; N=3 biological replicates; *** p<
0.001 (one-way ANOVA). (**c**) Cell proliferation assay by Edu incorporation performed in A-
375 cells exposed to Prltx and Doxo (one representative experiment with n=3
replicates/condition). *** p< 0.001 (one-way ANOVA, calculated on Edu$^+$ cells). (**d**)
Immunoblotting showing LMNB1, CDKN1A and GAPDH protein levels in A-375 cells
exposed to Prltx and Doxo (N=3 biological replicates). (**e**) Representative images from
Senescence Associated Beta Galactosidase assay performed on MDA-231 cells exposed to
Pralatrexate (Prltx). Doxorubicin (Doxo) was used as positive control for senescence
induction. Scale Bar: 10 µm. (**f**) qPCR analysis of CDKN1A and IL6 mRNA levels in MDA-
231 cells in response to Prltx and Doxo.; N=3 biological replicates; *p< 0.05 (one-way
ANOVA). (**g**) Cell proliferation assay by Edu incorporation performed in MDA-231 cells
exposed to Prltx and Doxo (one representative experiment with n=3 replicates/condition). **
p< 0.01; *** p< 0.001 (one-way ANOVA, calculated on Edu + cells). (**h**) Immunoblotting
showing LMNB1, CDKN1A and GAPDH protein levels in A-375 cells exposed to Prltx and
Doxo (N=3 biological replicates). (i) qPCR analysis of Oct4, Sox2 and Nanog levels in PC3
cells exposed to Piperlongumine (Piplng). * p< 0.05 Student's t test. (j) Immunoblotting
showing OCT3/4 and alpha-tubulin protein levels in PC3 cells incubated with Piplng..

1381

1382 **Figure 7. Pasta identifies age-modulatory gene perturbations.**

1383 (**a**) Volcano plot displaying the most significant Rejuvenating (negative age score) and Aging
1384 (positive age score) gene perturbations (GPs). (**b-f**) Enrichment analysis using the Hallmarks

of Aging[75] (**b-c**) and REACTOME (**d-f**) gene sets for Rejuvenating GPs (**b, d**), Aging GPs (**c, e**), and Aging GPs that are also compound targets of significant Aging compounds (**f**). Significant results from the three types of GPs (oe, sh, and xpr) were pooled, with overexpression results inverted (i.e., MYC_oe was included in the Aging GP set). Note that the Hallmarks of Aging gene sets are not directional (i.e., genes in the set "stem cell exhaustion" are related to stem cell in general; the last nouns were not used when compiling these gene sets). (**g**) Heatmap of GPs by cell lines with the most significant age scores (see Methods). Colors indicate the square root of a metric balancing significance and effect size: -*log10(adjusted p-value) * mean(age score)*. Missing values are shown in grey. (**h**) Boxplots and violin plots of dependency scores for GPs by cell line in the Aging, Rejuvenating and Other groups. The knockdown (KD) sets contain RNAi data for DepMap and shRNA data from CMAP, while the knockout (KO) sets contain CRISPR data from both databases. Number of entries per set: Other (KD: 27,243, KO: 46,684 entries), Aging (KD: 152, KO: 55), Rejuvenation (KD: 6, KO: 13). (**i**) REACTOME pathways whose PCCs are most significantly associated with dependency scores in the knockout set. All terms were negatively correlated with dependency scores. The grey line shows the 0.05 adjusted p-value cutoff. (**j**) Mean age score vs dependency for GPs in the retinoic acid (RA) biosynthesis pathway. Red and blue points show results for RDH13 and ALDH1A1, respectively, across different cell lines. (**k**) Mean age score vs dependency for GPs in the oncogene-induced senescence pathway. Red and blue points show results for MDM2 and TP53 knockout, respectively, across different cell lines. Abbreviations: oe, over-expression; sh, shRNA; xpr, CRISPR; KD, knockdown; KO, knockout.

Figure 8. Pasta uncovers the molecular determinants of cell lines' propensity for aging and rejuvenation.

(**a**) Propensity of various cell lines to become older or younger following exposure to age-regulatory perturbations (see Methods). (**b**) Gene mutations most strongly negatively correlated with the propensity for Aging. (**c**) Number of TP53 mutations versus the propensity of cells for Aging. (**d-i**) REACTOME pathway enrichment for genes whose protein levels (**d, g, i**), expression levels (**e**), and copy numbers (**f, h**) exhibit an absolute PCC higher than 0.4 with the propensity of cells for Rejuvenation (**d, e, f**) and Aging (**g, h, i**).

Supplementary Figures legends

Figure S1. Results of leave-one-dataset-out (LODO) analysis.

(**a**) Comparative performance in predicting relative age, assessed with LODO, when considering only young (< 40 years old) and old (> 60 years old) individuals. (**b**) Detailed LODO results for the AS40/Pasta model when considering all individauls. The x-axis shows the AUC-ROC for predicting whether the first sample is younger than the second in each evaluation pair.

1426 **Figure S2. Predicted versus true age differences for sample pairs using the AS40/Pasta**
1427 **LODO models.**

1428 For each dataset, predictions were generated by a model trained on all other datasets. Each
1429 point corresponds to a sample pair with an age difference of at least 40 years.

1430

1431 **Figure S3. Predicted age scores versus true ages for all samples using the AS40/Pasta**
1432 **LODO models.**

1433 For each dataset, predictions were generated by a model trained on all other datasets. Each
1434 point corresponds to one sample.

1435

Figure S4. Analysis of L1000 data with the baseline regression model recovers fewer
known age-modulatory compounds.

Proportion of compounds annotated as senescence-inducing (**a**), pro-reprogramming (**b**), or
both (**c**) for significantly age-increasing ('Aging'), age-decreasing ('Rejuvenating')
compounds, and other compounds ('Other'). Analyses were performed identically to Figure
4b-d but employed the baseline regression model from Figure 1 rather than the age-shift
model.

1443

1444 **Figure S5. REACTOME enrichment analysis of gene perturbations.**

1445 Results for Rejuvenating knockdown (**a**), knockout (**b**), and overexpression (**c**), and for Aging
1446 knockdown (**d**), knockout (**e**), and overexpression (**f**). (**g**) Genes with non-zero adjusted p-
1447 value scores (score_pa) across all cell types that showed consistency between the pooled gene
1448 perturbations and the compound target analyses.

1449

Figure S6. REACTOME enrichment analysis of pathways significantly associated with
the propensity for Aging and Rejuvenation.

Results for genes whose copy number positively correlates with the propensity for
Rejuvenation (**a**), proteins whose abundance negatively correlates with the propensity for
Aging (**b**), and genes whose transcript levels positively correlate with the propensity for
Rejuvenation (**c**) or negatively correlate with the propensity for Rejuvenation (**d**).

1456
1457 **Figure S7. Graphical summary of the study.**

1458

Supplementary Tables legends

Table S1. Datasets used to train and evaluate the age models.

The *dataset* column includes GEO accession numbers, Expression Atlas IDs, or project names (GTEx and TCGA). The *n_samples* column indicates how many samples from healthy individuals were used to train and evaluate the models. The *n_pairs* column indicates how many sample pairs were used to train the final AS40/Pasta model. The *age_Q1/Q2/Q3* columns represent the first, second and third quartile of donor age.

Table S2. Benchmarking results comparing Pasta and MultiTIMER.

This table contains the Pearson Correlation Coefficients (PCCs) between true age and predicted age score for Pasta and MultiTIMER in the datasets shown in panels 2a and 2b, and the tissue type for each dataset.

Table S3. GSEA pathway enrichment results based on Pasta's coefficients.

Gene-Set Enrichment Analysis (GSEA) of the coefficients of the Pasta model, using the REACTOME, CPG, KEGG and MIRDB gene sets from the Human MSigDB Collections.

Table S4. AUC-ROC results for senescence and stem cell datasets.

This table reports the AUC-ROC results for datasets shown in panels 3a, 3d, 3e, and 3f. It includes the number of samples (or pseudobulks for panel 3d) for each group as well as the tissue type for each comparison, and compound dose for panel 3d.

Table S5. Association between age scores or age and tumor grade.

Results from correlation analyses between tumor grade and either chronological age or Pasta age scores. Abbreviations: padj – adjusted p-value, signif – significant result.

Table S6. Association between age scores, chronological age, and all-cause survival.

Results from Cox proportional-hazards regression analyses using scaled Pasta age predictions and scaled chronological ages. Abbreviations: padj – adjusted p-value.

Table S7. Compounds with significant age-associations across all cell types.

1490 Abbreviations: padj – adjusted p-value, n – number of samples, n_cell – number of cell lines,
1491 med_age – median age scores.

1492

1493 **Table S8. Mechanisms of actions (MOAs) with significant age association across all cell**
1494 **types.**

1495 *bg* indicates the background and lists the MOA of all compounds included in the analyses.
1496 Abbreviations: padj – adjusted p-value, tot – total.

1497

1498 **Table S9. Compounds with significant age associations in specific cell types.**

1499 Abbreviations: padj – adjusted p-value, n – number of samples, med_age – median age score.

1500

Table S10. MOAs with a significant association between age score and dependency.

This table contains the significance of the association between mean age score and
dependency score across cell lines and compounds for each MOA, adjusted by compound
effects. Only entries with an unadjusted p-value below 0.05 are included in this table. The
column *n_cellXdrug* indicates the number of data points (i.e., compound-cell line pair) used to
fit the regression model. Abbreviations: PCC – Pearson Correlation Coefficient, padj –
adjusted p-value.

1508

Table S11. Compounds with a significant association between age score and dependency.

This table contains the significance of the association between mean age score and
dependency for each compound. Only entries with a non-adjusted p-value below 0.05 are
included in this table. The column *n_cell* indicates the number of data points (i.e., cell lines)
used to fit the regression model. Abbreviations: PCC – Pearson Correlation Coefficient, padj
– adjusted p-value.

1515

1516 **Table S12. Drugs used to treat cancers.**

1517 Drugs used to treat the various cancers analyzed in Figure 5. The *age_group* column indicates
1518 if a drug has a pro-aging effect, pro-rejuvenating effect, or no effect (group Other) across cell
1519 lines.

1520

Table S13. Enrichment of anti-cancer drugs in age-modulatory compound groups.

Abbreviations: in_cat_yes/in_cat_no – number of drugs that are used to treat the indicated
cancer that are in the given age-modulatory compound group (in_cat_yes) or that are not in

this group (in_cat_no), out_cat_yes/out_cat_no – number of drugs that are used to treat the
indicated cancer that are in the other age-modulatory compound groups (out_cat_yes) or that
are not in these groups (out_cat_no), padj – adjusted p-value.

1527

1528 **Table S14. Cancer-specific pro-aging compounds.**

1529 Abbreviations: sd – standard deviation, n_cell – number of cell lines, n_rep – number of
1530 entries, padj – adjusted p-value, age_diff – difference in mean age effect between tumor and
1531 normal cell lines.

1532

1533 **Table S15. Results for pralatrexate and piperlongumine in all cell types.**

1534 Abbreviations: padj – adjusted p-value, n – number of samples, med_age – median age score.

1535

1536 **Table S16. Gene perturbations with significant age associations across all cell types.**

1537 Abbreviations: padj – adjusted p-value, n – number of samples, n_cell – number of cell lines,
1538 med_age – median age score.

1539

1540 **Table S17. Hallmarks of Aging significantly enriched in Aging and Rejuvenating GPs.**

1541

1542 **Table S18. REACTOME terms significantly enriched in Aging and Rejuvenating GPs.**

1543

1544 **Table S19. Gene perturbations with significant age associations in specific cell types.**

1545 Abbreviations: padj – adjusted p-value, n – number of samples, med_age – median age score.

1546

Table S20. REACTOME terms with a significant association between age score and
dependency.

This table contains the significance of the association between mean age score and
dependency score across cell lines and gene knockout for each REACTOME term, adjusted
for the effect of gene knockouts. Only entries with a non-adjusted p-value below 0.05 are
included in this table. The column *n_cellXgeneKO* indicates the number of data points (i.e.,
GP-cell line pair) used to fit the regression model. Abbreviations: PCC – Pearson Correlation
Coefficient, padj – adjusted p-value, REAC_term – REACTOME term.

1555

1556 **Table S21. Genes whose mutation count across cell lines is associated with cellular**
1557 **propensity for Aging.**

1558 Only entries with a non-adjusted p-value below 0.05 are included in this table. Abbreviations:
1559 PCC – Pearson Correlation Coefficient, padj – adjusted p-value, n_cell – number of cell lines,
1560 n_mut – number of mutations, n_cell_mut – number of cell lines with at least one mutation.

1561

1562 **Table S22. REACTOME terms significantly associated with the propensity for Aging**
1563 **and Rejuvenation.**

1564 Abbreviations: PCC_dir – direction of the Pearson Correlation Coefficient.

1565

a Database. Heterogeneous count matrices from18 bulk RNA-Seq and 3 microarrays. → Modeling. Compare age modeling strategies through a leave-one-dataset-out cross-validation strategy.

b Regression (REG)

Predicted age / True age

c Traditional classifier (TC)

Dataset 1: Dataset 2: Age: 0 20 40 60 80 100

Class: Young (< 40) Old (> 60)

d Age−shift classifier (AS)

Dataset 1: Dataset 2: Age: 0 20 40 60 80 100

Class: Younger (< −40) Older (> 40)

e Leave−one−dataset−out cross−validation results

PCC all samples

REG_RC AS40_RC AS50 TC_30_70 AS30 REG TC_40_60 AS40|PASTA

Modeling strategy

f Retina scRNA−Seq

PCC / log2(chunk_size)

g Skin scRNA−Seq

PCC / log2(chunk_size)

h Lung scRNA−Seq

PCC / log2(chunk_size)

Modeling_strategy ● REGR ● TC46 ● AS40

a Anti-cancer drugs enriched in age-modulatory compounds

b Top cancer-specific senescence-inducing compounds

c Mean age effect for cancer and normal cell lines

Senescence induction in responder A375

a
Ctrl | Prltx | Doxo

b
CDKN1A mRNA (F.C) — Ctrl, Prltx, Doxo; *** ***

IL6 mRNA (F.C) — Ctrl, Prltx, Doxo; *** .075

c
Cell proliferation (%) — Ctrl, Prltx, Doxo; *** ***
cells (%): EdU⁺, EdU⁻

d
LMNB1 — 67 kDa
CDKN1A — 21 kDa
GAPDH — 36 kDa
Ctrl | Prltx | Doxo

Senescence induction in non-responder MDA-MB-231

e
Ctrl | Prltx | Doxo

f
CDKN1A mRNA (F.C) — Ctrl, Prltx, Doxo; ns *

IL6 mRNA (F.C) — Ctrl, Prltx, Doxo; ns *

g
Cell proliferation (%) — Ctrl, Prltx, Doxo; ** ***
cells (%): EdU⁺, EdU⁻

h
LMNB1 — 67 kDa
CDKN1A — 21 kDa
GAPDH — 36 kDa
Ctrl | Prltx | Doxo

Stemness induction in responder PC3

i
OCT4 mRNA (F.C) — Ctrl, Piplng; *
SOX2 mRNA (F.C) — Ctrl, Piplng; *
NANOG mRNA (F.C) — Ctrl, Piplng; *

j
OCT 3/4 — 38 kDa
TUBA1A — 50 kDa
Ctrl | Piplng

a Propensity of cells for age-changes

b Gene mutations most negatively correlated with propensity for Aging

c # of mutations in TP53 vs propensity for Aging

d REAC. terms for proteins whose levels positively correlate with propensity for Aging

e REAC. terms for genes whose mrna levels positively correlate with propensity for Aging

f REAC. terms for genes whose copy number positively correlate with propensity for Aging

g REAC. terms for proteins whose levels positively correlate with propensity for Rejuv.

h REAC. terms for genes whose copy number negatively correlate with propensity for Rejuv.

i REAC. terms for proteins whose levels negatively correlate with propensity for Rejuv.