**Molecular characterisation of neurons across animals identify conserved gene expression.**

Rivera-Rivera Carlos J.*[1], Feuda R.[2,3], Pisani D.*[1]

[1] Bristol Palaeobiology Group, School of Biological Sciences and School of Earth Sciences, University of Bristol (UK).

[2] Neurogenetics Group, University of Leicester (UK)

[3] Department of Genetics, Genomics and Cancer Sciences, University of Leicester (UK)

* Corresponding authors. Contact details: CJRR; carlos.riverarivera@bristol.ac.uk, Life Sciences Building, 24 Tyndall Ave, BS8 1TG Bristol, United Kingdom. DP; davide.pisani@bristol.ac.uk, Life Sciences Building, 24 Tyndall Ave, BS8 1TG Bristol, United Kindgom.

Running Title: *Conserved nerve cell gene batteries*

**Abstract**

The evolution of the nervous system has been shrouded in controversy since the onset of the genomics era. A large part of this controversy stems from the lack of phylogenetic consensus for the main branches of the animal tree, where often animals with nervous systems do not form a monophyletic group. However, this question can be informed from other non-phylogenetic perspectives, such as comparative genomics. Here we ask how similar are genes differentially expressed in neurons across a representative set of eight animals with a nervous system, and tally the presence or absence of their homologs in 10 animals and two choanoflagellates. We show that proteins from 39 families are differentially expressed in all neurons, regardless of the phylogenetic placement of their lineage, and that the majority of these gene families are present in the (unicellular) closest relatives of animals, choanoflagellates. We found that the members of these 39 gene families are enriched in domains for ion transport and juxtacrine signalling, and that there is one gene family of zinc-dependent extracellular matrix-remodelling proteins which is only found in animals bearing a nervous system. Our results show that common genetic toolkits are in place for the function of nervous systems. We identify a large number of potential new genomic markers linked to the nervous system and hope they can complement ongoing research efforts to better understand this quintessentially animal system.

**Introduction**

The nervous system, most commonly defined as a collection of neuronal cell types organised into a system via synaptic or anastomosed cell contacts (Sachkova et al. 2021), provided animals a discrete tissue system for integrating sensory information thus unlocking a free-living lifestyle in large multicellular organisms (Liebeskind et al. 2017). It can be argued that the nervous system is a hallmark of animal life and distinguishes the lineage in contrast to the other kingdoms of multicellular life.

Traditionally, it has been assumed that the nervous system is homologous across all animals. However, this hypothesis has recently been challenged (Moroz 2015). Uncertainty in our understanding of

nervous system evolution follows from the emergence of incongruence at the root of the animal phylogeny (Dunn et al. 2008a; Philippe et al. 2009; Schierwater et al. 2009; Philippe et al. 2011; Ryan et al. 2013; Whelan et al. 2015; Feuda et al. 2017; Steenwyk and King 2024). Traditionally, sponges (the Porifera), were considered the sister of all the other animals followed by the Placozoa (a small lineage of diploblastic marine organisms), and the Eumetazoa, traditionally identified as the lineage including all animals with a nervous system (Fig 1A, B, and E). Within Eumetazoa, two phyla; Cnidaria (corals and jellyfishes) and Ctenophora (comb jellies) were identified as sisters to each other in a lineage named Coelenterata. In turn, Coelenterata was identified as the sister to Bilateria (the lineage including all animals with bilateral body symmetry - acoel flatworms, arthropods and vertebrates - Fig. 1E). However, genomic data failed to corroborate this phylogeny, with some studies identifying the comb jellies as the sister of all the other animals (Fig. 1C and D) and others still identifying the sponges as the sister of all the other animals, but in a context where the comb jellies are the second lineage emerging along the animal tree (the tree in Fig. 1A and B). In these new scenarios, a single origin of the nervous system with no subsequent losses cannot be reconciled, suggesting more complex evolutionary histories where the nervous system might have evolved multiple times independently (Moroz 2021;    Moroz and Romanova 2022; Moroz and Romanova 2023) or, if it evolved only once, it was then secondarily lost multiple times (Fig. 1 summarises the possible scenarios), irrespective of the relative relationships of sponges and comb jellies.

**Figure 1**. Proposed hypotheses for the emergence(s) of the nervous system(s). A - Assuming that sponges are the first metazoan lineage to branch out (Porifera-sister hypothesis), the nervous system emerged in the ancestor to all animals minus sponges and was lost in the ancestor to placozoans. B - Assuming Porifera-sister, the nervous system emerged in the ancestor to all animals and was independently lost in sponges and in placozoans. C - Assuming ctenophores are the first metazoan lineage to branch out (Ctenophora-sister hypothesis), at least two independent emergences occurred, at the ancestor of all ctenophores, and again in the ancestor to cnidarians and bilaterians. D - Again assuming Ctenophora-sister, nervous systems emerged in the ancestor to all animals and were subsequently lost in sponges and placozoans. E - Assuming the Coelenterata hypothesis (sponges as the first branching metazoan and cnidarians + ctenophores sister to bilaterians), the nervous system emerged in the ancestor to all animals with a nervous system.

Here we address the problem of the origin of the nervous system with reference to gene content. We investigate the evolutionary history of gene families with a special focus on gene families that are expressed in the nervous system. We test how many such families are shared across different animal lineages and whether we could expect such numbers of shared families to be consistent with multiple origins of the nervous system. We used readily accessible whole-organism single-cell transcriptomes (scRNA-seq) for a variety of key taxa and the highly curated orthogroup database eggNOG (Huerta-Cepas et al. 2019). We use scRNA-seq data from a ctenophore, three cnidarians, and four bilaterians to determine which gene orthogroups are differentially expressed in different nervous systems, how similar they are to each other, and what their functional profiles are. We contrasted those results with the total coding sequence (CDS) data for 12 additional species spanning all main branches of animals and their closest relatives, the choanoflagellates, to determine at which taxonomic level each nervous system-expressed orthogroup emerged. Finally, we test whether the number of shared orthogroups is what we would expect if the nervous systems of key organisms evolved independently.

We found that irrespective of the relative relationships of ctenophores and sponges, all animals with a nervous system differentially express a core set of orthogroups in their neuronal cells. Pairwise tests showed that these sets are more similar to each other than expected by chance alone. We interpret

these results to suggest that the nervous system most likely evolved only once, and was subsequently lost, in Placozoa (if sponges are the sister of all the other animals), or in sponges and Placozoa (if ctenophores are the sister of all the other animals). If it evolved twice independently, this underpinning similarity suggests that the same genes were independently co-opted in the evolving nervous systems. This interpretation is possible but less parsimonious.

**Results**

*Nervous systems share 39 orthogroups*

We identified the following numbers of nsOGs for the eight core species in our dataset: *C. gigas* (3,227), *D. melanogaster* (8,498), *D. rerio* (9,854), *H. vulgaris* (6,552), *M. leidyi* (1,125), *M. musculus* (7,023), *N. vectensis* (1,826), *S. pistillata* (3,889) (Table 1, nervous system cell barcodes in Supplementary Table S1). Of these, a total of 39 nsOGs were differentially expressed in all eight species' neuronal clusters (mouse gene family names on Supplementary Table S2). We found that the most common biological process GO term among the 39 families was 'transmembrane transport - GO:0055085'; for molecular function, 'protein binding - GO:0005515'; and for cell compartment, 'membrane - GO:0016020' (Full data in Supplementary Table S3). The most common protein domains in these 39 orthogroups include a large number of transmembrane proteins domains including protein, amino acid, carbohydrate, and ion transporters (mainly $Ca^{2+}$, although in one case, an $Na^+$ transporter, and in a different case $H^+$ transporter) as well as GTPases, chaperones, cytoskeleton-associated proteins, kinases, and zinc fingers (Supplementary Table S3).

**Table 1**. eggNOG search results overview. For all species that had scRNA-seq data analysed in this work, the total of sequences in the (CDS) reference dataset, how many of those were found in the eggNOG 5.0.2 database, how many of those had Opisthokonta-level orthogroup annotation. Then, how many genes were found to be differentially expressed in neurons, how many of those were found in the eggNOG 5.0.2 database, how may had an Opisthokonta-level orthogroup annotation, and finally, the total number of opisthokont orthogroups represented in the set of neuronally differentially-expressed sequences. The reference for *S. pistillata* included sequence data from several different sources and some sequences may have been repeated, which explains the large number of sequences in the reference. When tallying the total numbers of orthogroups (both for the CDS and neuronal datasets), repeated orthogroups were not allowed, which effectively removed repeats.

| Species | Total seqs. in reference | Reference eggNOG hits | Reference Opis. OGs | Neuronal sequences (NS) | Annotated NSs | NSs with Opis. OG | Total Opis. NS OGs |
|---|---|---|---|---|---|---|---|
| *Danio rerio* | 32,057 | 28,037 (87%) | 27,465 (86%) | 19,129 | 18,742 (98%) | 18,478 (97%) | 9,854 |
| *Mus musculus* | 28,862 | 25,981 (90%) | 25,566 (89%) | 9,767 | 9,564 (98%) | 9,564 (98%) | 7,023 |
| *Drosophila melanogaster* | 17,825 | 14,287 (80%) | 13,959 (78%) | 13,399 | 11,832 (88%) | 11,651 (87%) | 8,498 |
| *Crassostrea gigas* | 36,012 | 24,165 (67%) | 21,964 (61%) | 5,362 | 4,216 (79%) | 4,012 (75%) | 3,227 |
| *Mnemiopsis leidyi* | 16,548 | 12,666 (77%) | 9,578 (58%) | 1,680 | 1,402 (83%) | 1,214 (72%) | 1,125 |
| *Nematostella vectensis* | 32,280 | 24,632 (76%) | 18,918 (59%) | 3,044 | 2,547 (83%) | 2,071 (68%) | 1,719 |
| *Stylophora pistitllata* | 50,024* | 41,926 (84%) | 35,834 (72%) | 6,918 | 5,851 (90%) | 5,244 (76%) | 3,889 |
| *Hydra vulgaris* | 38,749 | 24,084 (62%) | 16,419 (42%) | 12,279 | 10,179 (83%) | 8,773 (71%) | 6,552 |

We found that the species with the highest percentage of unique nsOGs was *D. melanogaster*, with 35.81% of its 8,498 nsOGs unique to its set (similar to the uniqueness of its totOGs set). The smallest set of unique nsOGs (7.20%) was that of *M. leidyi* (Fig. 2). The largest difference in uniqueness between the totOG and nsOG was found in *D. rerio*, for which 11.31% of its totOGs are lineage-specific, while 28.65% of its nsOGs are lineage-specific. *M. musculus* showed the smallest difference in number

of unique OGs with 13.69% unique totOGs and 15.63% unique nsOGs (Fig. 2).

**Figure 2.** Genomic uniqueness for all species used in this study. Bars are shown for each species studied, and the colours represent the following: orange - percent of orthogroups in full CDS reference datasets which were unique to that species; green - percent of nervous system orthogroups which were unique to a particular species; and blue - percent of sequences (not orthogroups) which did not have a hit in the eggNOG 5.0 database. Species name codes are as follow: Sr - *Salpingoeca rosetta* (Choanoflagellata) Mb - *Monosiga brevicollis* (Choanoflagellata), Sl - *Spongilla lacustris* (Porifera), Em - *Ephydatia muelleri* (Porifera), Aq - *Amphimedon queenslandica* (Porifera), Pb - *Pleurobrachia bacheii* (Ctenophora), Ml - *Mnemiopsis leidyi* (Ctenophora), Sp - *Stylophora pistilliata* (Cnidaria), Hv - *Hydra vulgaris* (Cnidaria), Nv - *Nematostella vectensis* (Cnidaria), Ta - *Trichoplax adhaerens* (Placozoa), Cg - *Crassostrea gigas* (Mollusca - bilaterian), Dm - *Drosophila melanogaster* (Arthropoda – bilaterian), Sm - *Schimidtea mediterranea* (Platyhelminthes – bilaterian), Sa - *Schistosoma mansoni* (Platyhelminthes – bilaterian), Su – *Strongylocentrotus purpuratus* (Echinodermata – bilaterian) Cc - *Carcharadon carcharias* (Chordata – bilaterian), Dr - *Danio rerio* (Chordata – bilaterian), Xt - *Xenopus tropicalis* (Chordata – bilaterian), Mm - *Mus musculus* (Chordata – bilaterian).

*Orthogroups in common in CDS data studied*

CDS sequences from the expanded dataset of 20 species with no hits in the eggNOG database ranged from 3.0% (the white shark) to 75.6% (the placozoan), ($\bar{x} = 25.2\%, \sigma = 18.6\%, N = 20$, for

average, standard deviation, and sample size, respectively – relative to all other species; see Supplementary Table S4a). The organism with the least number of totOGs is *T. adhaerens* (3,830), and the one with the most, *X. tropicalis* (12,514), ($\bar{x} = 7,944, \sigma = 2,719.1, N = 20$). All 20 totOG sets (including two choanoflagellates and the placozoan) shared 437 orthogroups, whereas when constrained to only the species with a nervous system (14 in total, including all eight with scRNA-seq data, two flatworms, one more ctenophore, the white shark, and the purple sea urchin) this number is 1,258. In contrast, the totOG sets of the eight species with scRNA-seq data show 2,049 shared orthogroups. We also calculated the percent of the total set of orthogroups which was unique to the species when compared to all the other 19 totOG sets. The species with the highest percentage of unique totOGs was *D. melanogaster*, with 33.67% of its totOGs not found in any of the other 19 totOG sets. The species with the least percentage of unique totOG sets was *P. bacheii*, with 0.79% of its totOGs as lineage-specific. *D. melanogaster*'s percent of unique CDSs is also reflected in the percent of unique nsOGs, 35.81%. *M. leidyi*'s CDS reference had 2.50% of its totOGs as unique (see Fig. 2, and Supplementary Table S4a for fully detailed numbers).

*Taxonomic distribution of global nsOGs*

Of the 39 nsOGs found in common among all of the eight species of eumetazoans we studied, 84.6% were present in the choanoflagellate *S. rosetta* CDS data, and 76.9% in the choanoflagellate *M. brevicollis*. Sponges also showed expression of this set: *S. lacustris* (97.4%), *A. queenslandica* (89.7%), and *E. muelleri* (94.9%). Finally, the placozoan *T. adhaerens*, which has a secondarily reduced body

plan and no known nervous system (Najle et al. 2023) only 69.2% of the nsOGs were present in its CDS reference (Supplementary Table S4b). All deuterostomes in our dataset had 100% of the NeurMet-nsOGs in their CDS datasets (Supplementary Table S4c).

According to the eggNOG taxonomic profiles, 29 of the 39 (74%) nsOGs shared by our core set of eight species, were present in the common ancestor of animals and Fungi. Six (15%) emerged in the holozoan MRCA, three (7.5%) in the metazoan MRCA, and one (~2.6%) in the ancestor to Eumetazoa (as defined in eggNOG database: i.e. Bilateria plus cnidaria and ctenophores – Supplementary Table S3). This eumetazoan specific nsOG is a member of the matrix metalloprotease (MMP – eggNOG code 38CRR@33154), a family of zinc-dependent

**Figure 3.** Presence/absence matrix of all 39 NeurMet-nsOGs in the CDS datasets of all the 20 species used in this study, using the following colour coding: Blue - orthogroup is present in the CDS dataset of that species; Green – orthogroup is absent in CDS dataset of that species. The species which are highlighted are species in our dataset which lack a nervous system and the OG highlighted (MMP) is one found only in organisms with a nervous system. OG names are based on mouse protein families.

extracellular matrix-remodelling proteins which belong to the metzincin protease superfamily (de Almeida et al. 2022), and we could confirm its absence in all three sponge CDS datasets as well as in *Trichoplax*'s (Fig. 3).

*NsOGs set are more similar to each other than expected by chance*

All pairwise comparisons among the nsOGs showed that the nsOGs of different species are more similar to each other than expected by chance, with all the observed values being larger than those from the permuted distributions (Supplementary Figure S1).

Our validatory test showed that there is usually greater similarity between nsOGs sets than between nsOGs and digestive system OGs (dsOGs). Indeed, for the cases of *D. melanogaster* and *H. vulgaris*, the number of OGs shared between the nsOG set and the dsOG set is random (Fig. 4). However, for the other cases it is not random. In the comparisons against the nsOGs of the remaining bilaterian taxa, the proportion of OGs shared between different nsOGs is higher than the proportion of OGs shared between dsOGs and nsOGs (Fig. 4). However, comparisons between the dsOGs of *M. leidyi* against the nsOGs of *N. vectensis* and *S. pistillata*, did not conform to the expected pattern, finding greater similarity than when comparing nsOGs among themselves. (Fig. 4).

**Figure 4.** Overlap between the nervous and digestive system orthogroup sets of *M. leidyi* and the nervous system orthogroup sets of all other animals with a nervous system. Red distributions show the overlap between nervous systems expected by chance, and the red vertical line shows the observed overlap. Green distributions show the overlap expected by chance between the digestive system orthogroup set of *M. leidyi* and the second species' nervous system orthogroup set, and the green vertical line is the observed overlap between both sets.

**Discussion**

By combining scRNA-seq data and comparative genomics we show that 39 gene orthogroups have a common pattern of differential expression in neural cells. Furthermore, we showed that these orthogroups (four of which are Metazoan innovations) are enriched in protein domains that mediate cell-cell signalling as well as signalling within cells, but crucially, most of the orthogroups are present in choanoflagellates, the sister group to animals (84.6% and 76.9%, respectively). Finally, we showed that the set of nervous system orthogroups (nsOG) of the ctenophore *Mnemiopsis leidyi*, which has been previously hypothesised to have independently evolved its nervous system (Moroz 2015), is more similar to the set of the nsOGs of all other animals than is expected by chance. We compared these results with the total CDS data of 10 other animals and showed that most of these 39 nsOGs present in all nervous system-bearing animals are also found in animals without a nervous system (sponges and placozoans).

*Conservation/co-option of functional and developmental cues in animal nervous systems*

Whether the nervous system emerged once at the root of the animal tree and was lost in sponges and placozoans, emerged independently in ctenophores and in the ancestor to cnidarians and bilaterians, or emerged once in an ancestor to ctenophores, cnidarians, and bilaterians after this lineage separated from sponges and placozoans, remains a fundamental but as of yet, unanswered question (Marlow and Arendt 2014; Ryan et al. 2015; Feuda et al. 2017; Schultz et al. 2023). While part of this question can only be addressed phylogenetically, we can gain important clues about the origin of the nervous system also performing analyses which do not rely on the relative placement of ctenophores, sponges and placozoans in the animal phylogeny, but identifies strongly differentially expressed gene families across nervous systems. We focus here on characterising the genomic component of nervous systems in general, with the intention of adding evidence to this question. Our results show that nervous systems in general share an enriched expression of domains for protein traffic and import/export, and ion channels. This does not come as a surprise, given the known role of

neuropeptides and of paracrine signalling in whole-organism intercommunication (Jékely 2021; Moroz 2021), as well as the role of ion channels in creating actions potentials for neuronal firing (Hodgkin and Huxley 1952). It stands out that the neurons we studied here also share an enrichment in juxtacrine signalling domains. These domains are known to be essential mediators of cell differentiation during neurogenesis (Notch/Jagged, EGF-like, semaphorin), as well as crucial signals in axon guidance (Yaron and Sprinzak 2012) and are present in all nsOGs. As more non-model organisms are studied using single-cell sequencing methods, this core set of genes can be revised and vetoed, and we look forward to a more comprehensive understanding of gene orthogroups involved in the nervous system development and function.

*One nsOG is present in all nervous systems and absent from the CDS sets of animals lacking a nervous system*

The taxonomic profile of one orthogroup, MMP (38CRR@33154) shows it to be unique to cnidarians, ctenophores, and bilaterians (both in the eggNOG database and in our own CDS data), suggesting a nervous system-specific gain that clusters all animals with a nervous system (Fig. 4). A second orthogroup, KANSL1 (38MCJ@33154) showed a similar pattern in the eggNOG database (unique to cnidarians, ctenophores, bilaterians, and placozoans), but we found it in two of our three sponge CDS datasets (*E. muelleri, S. lacustris,* and *A. queenslandica*). The MMP-orthogroup sequences from *M. leidyi* are assigned by eggnog-mapper the KEGG pathway of 'Parathyroid hormone synthesis, secretion, and action' and the enriched Pfam domains for this OG across all animals were 'Peptidase_M10', 'PG_binding_1', 'Hemopexin', and 'C1q', domains related to peptide cleaving, bacterial cell wall degradation, haem binding, and activation of serum complement system, respectively (Supplementary Table S3).

*All nsOGs are more similar to each other than expected by chance*

In pairwise comparisons, all nsOG sets were more similar to each other than expected by chance (Supplementary Figure S1). This is not surprising for the species which are hypothesised to have inherited the nervous system from a common ancestor, but it is interesting when considering the possibility that ctenophores independently evolved neurons and the nervous system. Assuming an independent emergence of the nervous system in ctenophores, one can envision that a rudimentary neuronal components were already in place in the ancestor to all metazoans, and thus was inherited by all clades (as discussed by Jékely et al. 2015). These rudimentary neuronal components then evolved in parallel to form nervous systems which employ the same biochemical mechanisms If this is the case, ctenophores, a group that in any case deserves much attention, will become a fundamental model organism for asking questions about the parallel evolution of complex systems in multicellular animals. On the other hand, if ctenophores inherited a fully integrated nervous system from an eumetazoan ancestor, they would still remain a fundamental group to study lineage-specific specialisation of nervous systems (Moroz 2015; Sachkova et al. 2021; Burkhardt et al. 2023). However, these lines of inquiry will be contingent on a consensus on the phylogenetic placement of ctenophores in the animal tree of life.

*The evolution of a nervous system in an uncertain phylogenetic context*

If we contextualise the results from this work into the most common phylogenetic hypotheses set forth for the early branching patterns of metazoans, we have the following possible scenarios. 1. Assuming the Ctenophora-sister hypothesis, the ancestor to all animals had a rudimentary signalling system that led to the parallel evolution of nervous systems with 39 orthogroups in common in ctenophores and then in all other animals, the ancestor of sponges lost its nervous system, and alongside it, the MMP orthogroup, and both the nervous system and the MMP orthogroup were again lost independently in the ancestor to placozoans (Fig. 5A). 2. Assuming the Ctenophora-sister hypothesis, the ancestral ctenophore evolved a nervous system that, due to functional constraints ended up sharing 39 orthogroups (including MMP, a group found in all other metazoans with a nervous

system) with that of an independently evolved nervous system on all other metazoans barring sponges, and this nervous system, as well as the MMP orthogroup was lost in placozoans (Fig. 5B). 3. Assuming the Porifera-sister hypothesis, the ancestor to all non-poriferan metazoans developed a nervous system which used a new clade-specific orthogroup, MMP, which the ancestor to placozoans subsequently lost alongside its nervous system (Fig. 5C). 4. Assuming the Porifera-sister hypothesis, the ancestor to all animals developed a rudimentary signalling system which was lost in the ancestor to sponges, then the ancestor to all animals except sponges specialised this signalling system into an integrated nervous system which used a new orthogroup, MMP, which the ancestor to placozoans lost in its morphological reduction (Fig. 5D). 5. Assuming the Coelenterata hypothesis, the ancestor to all animals with a nervous system evolved an integrated nervous system, which included the MMP orthogroup (Fig. 5E). A recent study in Placozoan peptidergic neurons showed support for the hypothesis that the earliest instance of a nervous system was a secretory network of cells, which lends support to the hypotheses presented in Fig. 5A and 5C (Najle et al. 2023).

**Figure 5.** The five phylogenetic hypotheses discussed in this paper with the data from this study included in them. The hypotheses are as follows: A. Assuming Ctenophora-sister, a rudimentary paracrine signalling nervous proto-system which included the MMP OG was present in the ancestor to all animals. This nervous proto-system evolved into the nervous system we observe in ctenophores nowadays, and on its sister branch, it was lost independently (at least as a purely nervous signalling system), alongside the MMP OG, two times – in Porifera and in Placozoa. Finally, in the ancestor to Bilateria and Cnidaria, the nervous proto-system evolved into the nervous system we see in these group. B. Assuming Ctenophora-sister, the ancestor to all animals already had a fully integrated nervous system which included the MMP OG, and this system, along with the MMP gene family was lost independently in Porifera and Placozoa. C. Assuming Porifera-sister, a nervous proto-system and the MMP OG were present in the ancestor to all non-poriferan animals, which then independently evolved a fully-integrated nervous system in Ctenophora and Bilateria + Cnidaria, and Placozoans lost the MMP OG as well as the nervous proto-system. D. Assuming Porifera-sister, the ancestor to all non-sponge animals evolved a fully-integrated nervous system that used the MMP OG in its development/operation, and this system and the OG were lost in Placozoa. E. Assuming the Coelenterata grouping, the ancestor of Bilateria + Coelenterata evolved a fully integrated nervous system which used the MMP OG in its development and/or operation.

Here we completed the first detailed description of the orthogroup composition of metazoan nervous systems, using scRNA-seq data from representative lineages of eight main metazoan branches. We showed how, regardless of phylogeny, animal nervous systems share important components of their

biochemical machinery both for development and function, and we show that a single orthogroup which is differentially expressed in all nervous systems is present only in the genomes of animals with a nervous system and absent otherwise. This work represents an essential first step into reconstructing the regulatory networks conserved/co-opted in nervous systems across animals, which can be a key non-phylogenomic source of information to clarify the contentious evolutionary path of this quintessential animal system.

**Materials and Methods**

*Data acquisition and curation*

We obtained post-processed unique molecular identifier (UMI) count tables from previously published scRNA-seq studies for eight metazoans with nervous systems: *Mnemiopsis leidyi* (Ctenophora) (Sébé-Pedrós et al. 2018a)*, Hydra vulgaris* (Cnidaria) (Siebert et al. 2019)*, Nematostella vectensis* (Cnidaria) (Sébé-Pedrós et al. 2018b)*, Stylophora pistillata* (Cnidaria) (Levy et al. 2021)*, Crassostrea gigas* (Protostomia: Lophotrochozoa) (Piovani et al. 2023)*, Drosophila melanogaster* (Protostomia: Ecdysozoa) (Li & Janssens et al. 2022)*, Mus musculus* (Vertebrata) (Pijuan-Sala et al. 2019; Griffiths and Lun 2023), and *Danio rerio* (Vertebrata) (Lange et al. 2023). For each, we also obtained the reference genomic CDS datasets used to annotate the scRNA-seq reads, as well as cell annotations where available (see Supplementary Table S5 for dataset and original references details). The cell annotation for nervous system cells of *M. leidyi* was produced by Sachkova et al. (2021), a different study from the one that initially produced the whole-organism scRNA-seq datasets. We also obtained full-organism CDS data for 12 other holozoan lineages: Choanoflagellata: *Monosiga brevicollis*, *Salpingoeca rosetta;* Porifera: *Spongilla lacustris, Amphimedon queenslandica, Ephydatia muelleri*; Ctenophora: *Pleurobrachia bacheii*; Placozoa: *Trichoplax adhaerens*; Platyhelminthes: *Schmidtea mediterranea, Schistosoma mansoni*; Echinodermata: *Strongylocentrotus purpuratus*; and Vertebrata: *Carcharadon carcharias, Xenopus tropicalis* (source dataset details in Supplementary Table S5).

Combined with the eight species whose scRNA-seq data we analysed, we worked with a total of 20 species: two choanoflagellates and 18 metazoans. We analysed all 20 genome-level CDS datasets with eggnog-mapper v2.1.8 (Cantalapiedra et al. 2021) and queried the eggNOG v5.0.2 (Huerta-Cepas et al. 2019) database with all the reference sequence sets using the diamond v2.0.14 (Buchfink et al. 2021) search algorithm. We then extracted, for each species, the list of orthogroups in the whole CDS set (totOGs from now on) based on their Opisthokonta-level annotation. Focusing on the Opisthokonta annotation allowed us to directly compare gene families across all species in our dataset. We also tallied the number of CDS without an eggNOG database hit, number of sequences without an Opisthokont orthogroup assignment, and the total number of orthogroups represented in each CDS dataset. These values allowed us to gauge the quality and completeness of our OG assignments (see Fig. 6 for a visual representation of the pipeline).

**Figure 6.** Schematic representation of the pipeline for obtaining the list of eggNOG orthogroups differentially expressed in nervous systems across animals. ScRNA-seq datasets for eight animal species were combined with the lists of neuronal cell barcodes or a list of neuropeptide markers (for *Crassostrea gigas*) in order to produce species-specific lists of genes differentially expressed in neurons. In parallel, the CDS reference datasets for all eight species plus 10 additional animals and two choanoflagellates were analysed with eggnog-mapper which provided the Opisthokonta-level orthogroup assignment for most CDS (totOGs). We gauged the quality and completeness of the eggnog-mapper results (see Fig. 3 below) to detect biases stemming from eggNOG 5.0.2's database exhaustiveness. Next, each neuronal marker was queried in its corresponding CDS eggnog-mapper result, and the opisthokont-level orthogroup classification recorded. The resulting lists of nervous system orthogroups (nsOGs) were then compared with all sets against each other using in order to obtain the final list of 39 pan-metazoan orthogroups differentially expressed in neurons of the eight animals studied ('NeurMet-nsOGs').

*Pipeline for identifying nervous system cells*

To detect differentially expressed genes in the scRNA-seq datasets we developed a bespoke script

(available on https://github.com/carlosj-rr/scrnaseq_wrangle/DEtector.py). First, for each dataset

and species, all genes which showed a stable expression pattern across cells (and as such were non-

informative) were removed. The genes have a distribution statistically indistinguishable (using the

Kolmogorov-Smirnov test) from a monotone dataset of the same size based on the median value.

Next, we ranked differentially expressed genes and all UMI counts were recoded to '1' for any cell in

which the gene was expressed below the 5$^{th}$ percentile of all cells (downregulation), '2' for any cell in

which the gene was expressed between the 5th and 95th percentiles of all cells (standard expression), and '3' for any cell in which the gene was expressed above the 95th percentile of all cells (upregulation). We used a 5% cutoff to determine down- and upregulation because we wanted to focus on genes showing strong differential expression. These recoded tables (one per species) are available as Tables S6-S13 in data.bris). We used two different approaches to identify nervous system genes from the table of differentially expressed genes. For all taxa but the mollusc *C. gigas,* and the cnidarian *N. vectensis,* we extracted strongly differentially expressed genes in cell clusters identified as belonging to the nervous system in their original publication. For *N. vectensis*, the authors had identified a set of neural markers which we used to subselect the nervous system cells. For *C. gigas*, Piovani et al. (2023) identified 44 neuropeptide markers in their dataset, and based on that list, we were able to classify cells as nervous system cells if they strongly differentially expressed more than 95% of the 44 neuropeptides.

*Detecting genes differentially expressed in nervous system cells*

Next, we developed an algorithm and a metric to identify genes that are differentially expressed in neurons. First we turned the recoded expression matrices S2 to S9 into Boolean tables by changing all '1's and '3's (down-/up-regulation) to 'true' and all '2's (standard expression) to false. To detect genes with high differential expression in neuronal cells, we defined a metric that we named the proportional expression preference (PEP). This metric defines in what proportion of neuronal cells ('foreground' population) a gene is expressed in comparison to non-neuronal cells (the 'background' population).

$$PEP_i = P(FG_i = True) - P(BG_i = True),$$

where $P(FG_i = True)$ represents the proportion of neuronal cells in which gene $i$ is differentially expressed, and $P(BG_i = True)$ the proportion of the non-neuronal cells in which the same gene is differentially expressed. This metric ranges from -1, if a gene is differentially expressed in all non-

neuronal and zero neuronal cells, to 1, for a gene that is preferentially expressed in all neuronal cells and no non-neuronal cells.

Next, a permutation test was performed to define whether the observed PEP value for each gene was above what was expected purely by random expression noise. For each species, we counted the number of cells classified as nervous system cells. After that we randomly sampled the same number of cells from across the full dataset (nervous system cells included), 100 times. We then calculated the PEP values for each gene across the 100 random samples of cell populations. If the observed PEP value for a gene was above the 95th percentile of the distribution of PEP values from random foreground/background cell combinations, the gene was identified as differentially expressed in nervous system cells. This resulted in the generation of species-specific lists of genes showing strong differential expression in nervous system cells.

To find common neuronal orthogroups among all species, we extracted the eggNOG opisthokont orthogroup annotation for each of the genes differentially expressed in nerve cells. In the eggNOG database, a gene family is equivalent to an orthogroup, as they encompass all orthologous and paralogous genes which descend from a common ancestor. Using the Opisthokonta-level orthogroup (OG) ID enabled us to compare OGs between all lineages in our dataset and allows for the results of our work to be extended to Fungi and other opisthokont clades that were not included in our analyses. This is important because, while Fungi and other opisthokonts do not have a nervous system, gene families with nervous system expression have deep evolutionary roots (McCoy and Fire 2024).

*Measuring similarities of nervous system orthogroup and total genomic orthogroup sets at different taxonomic levels*

We calculated the intersection of total OG sets (totOGs) across all 20 CDS datasets. We determined, for our 20-species datasets, how many orthogroups were in common among vertebrates, protostomes, bilaterians, ctenophores, cnidarians, "coelenterates" (assuming their monophyly),

eumetazoans, eumetazoans excluding the ctenophore, metazoans excluding the ctenophores, all metazoans (including ctenophores), and holozoans (metazoans plus choanoflagellates). We did the same analysis with the eight sets of OGs differentially expressed in neural cells (nsOGs), although the taxonomic group were limited to our eight species with scRNA-seq data.

Once we obtained a set of shared OGs across all metazoan nsOGs (NeurMet-nsOG), we defined at which point each neural cell OG emerged by cross-referencing with the much more complete eggNOG 5.0.2 database. We searched each NeurMet-nsOG's taxonomic profile in the database's website (http://eggnog5.embl.de/#/app/home; Huerta-Cepas et al. 2019), which shows the species in which a given OG is found, thus giving a good idea of when the OG may have emerged. We specifically asked which orthogroups were present in the last common ancestor of Fungi and animals, the last common ancestor of holozoans and animals, and within animals, the last common ancestor of all eumetazoans. Note that in the eggNOG 5.0.2 database Eumetazoa includes ctenophores by default. For all comparisons, we cross-referenced with our own datasets, to make sure our data corroborated or expanded those of the eggNOG database. This also enabled us to count the number of totOGs in common between all metazoans excluding ctenophores, a grouping that would emerge if the Ctenophora-sister hypothesis is true. Due to the way the eggNOG web interface is set up for the exploration of taxonomic profiles of gene families, we could not leverage the much larger taxonomic sample of that database for detecting totOGs which could have emerged in the 'Metazoa minus ctenophora' clade. In addition, we checked whether any of the NeurMet-nsOGs were present in the CDS sets of both choanoflagellates of this study.

*Functional profiling of NeurMet-nsOGs*

To clarify the protein functional domains present in each NeurMet-nsOGs we used Pfam (Mistry et al. 2021) and the InterPro database (Paysan-Lafosse et al 2022) for the gene ontology (GO) annotation (where available).

*Quantifying difference between nsOG sets*

We developed a permutation-based approach to test whether two nsOG sets were more similar to each other than expected by chance. Briefly, we extracted random subsamples of totOG of the same size as the nsOG set for each species and calculated the proportion of OGs found on both subsamples, relative to the size of the nsOG of a species of interest (100 permutations). We then tested whether the number of nsOGs shared between two species was significantly larger than that expected when randomly sampling OGs from these two species. To make sure the signal picked up in our analyses did not identify background similarity shared among cell types, we performed a validatory test using *M. leidyi*. For this test, we compared the OGs from the digestive cell cluster (dsOGs) of *M. leidyi* against the nsOGs of every other species in our dataset. We reasoned that if our results represented background similarity, rather than cell-specific homologous similarity, pairwise comparison between dsOGs and nsOGS should return results that are indistinguishable from those obtained when comparing nsOGs among each other.

**Data access**

All scRNA-seq and CDS data were obtained from public databases (see Supplementary Table S1 for sources). All supplementary figures, tables and data are available at the University of Bristol data repository, data.bris, at https://data.bris.ac.uk/data/dataset/20n4tk7g5vjgd2fxagwcqmghqx. All code to replicate the analysis, with detailed explanations on how each function works is available in CJRR's GitHub repository: github.com/carlosj-rr/scrnaseq_wrangle.

**Competing interests**

The authors do not have any competing interests.

**Acknowledgements**

This work was funded by a Marie Skłodowska-Curie Action to CJRR (project code RipGEESE), RF is supported by the Royal Society (UF160226 and URF/R/221011) and DP and RF are supported by a Leverhulme Trust Grant RPG 2024-030. We thank all the members of the Bristol Palaeobiology group

**References**

Buchfink B, Reuter K, Drost HG. 2021. Sensitive protein alignments at tree-of-life scale using DIAMOND. *Nat Methods* **18**: 366–368. http://dx.doi.org/10.1038/s41592-021-01101-x.

Burkhardt P, Colgren J, Medhus A, Digel L, Naumann B, Soto-Angel JJ, Nordmann E-L, Sachkova MY, Kittleman M. 2023. Syncytial nerve net in a ctenophore sheds new light on the early evolution of nervous systems. *Science (80- )* 293–297. https://www.science.org/doi/epdf/10.1126/science.ade5645.

Cantalapiedra CP, Herņandez-Plaza A, Letunic I, Bork P, Huerta-Cepas J. 2021. eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the Metagenomic Scale. *Mol Biol Evol* **38**: 5825–5829.

de Almeida LGN, Thode H, Eslambolchi Y, Chopra S, Young D, Gill S, Devel L, Dufour A. 2022. Matrix Metalloproteinases: From Molecular Mechanisms to Physiology, Pathophysiology, and Pharmacology. *Pharmacol Rev* **74**: 712–768.

Dunn CW, Hejnol A, Matus DQ, Pang K, Browne WE, Smith SA, Seaver E, Rouse GW, Obst M, Edgecombe GD, et al. 2008. Broad phylogenomic sampling improves resolution of the animal tree of life. *Nature* **452**: 745–749.

Feuda R, Dohrmann M, Pett W, Philippe H, Rota-Stabelli O, Lartillot N, Wörheide G, Pisani D. 2017. Improved Modeling of Compositional Heterogeneity Supports Sponges as Sister to All Other Animals. *Curr Biol* **27**: 3864-3870.e4.

Griffiths J, Lun A. 2023. BioconductoR's MouseGastrulationData.

Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, Mende DR, Letunic I, Rattei T, Jensen LJ, et al. 2019. EggNOG 5.0: A hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Res* **47**: D309–D314.

Jékely G. 2021. The chemical brain hypothesis for the origin of nervous systems. *Philos Trans R Soc B Biol Sci* **376**.

Jékely G, Paps J, Nielsen C. 2015. The phylogenetic position of ctenophores and the origin(s) of nervous systems. *Evodevo* **6**.

Lange M, Granados A, VijayKumar S, Bragantini J, Ancheta S, Santhosh S, Borja M, Kobayashi H, McGeever E, Solak AC, et al. 2023. Zebrahub – Multimodal Zebrafish Developmental Atlas Reveals the State Transition Dynamics of Late Vertebrate Pluripotent Axial Progenitors.

*bioRxiv* 2023.03.06.531398. http://biorxiv.org/content/early/2023/03/07/2023.03.06.531398.abstract.

Levy S, Elek A, Grau-Bové X, Menéndez-Bravo S, Iglesias M, Tanay A, Mass T, Sebé-Pedrós A. 2021. A stony coral cell atlas illuminates the molecular and cellular basis of coral symbiosis, calcification, and immunity. *Cell* **184**: 2973-2987.e18.

Li H, Janssens J, de Waegeneer M, Kolluru SS, Davie K, Gardeux V, Saelens W, David FPA, Brbić M, Spanier K, et al. 2022. Fly Cell Atlas: A single-nucleus transcriptomic atlas of the adult fruit fly. *Science (80- )* **375**.

Liebeskind BJ, Hofmann HA, Hillis DM, Zakon HH. 2017. Evolution of Animal Neural Systems. *Annu Rev Ecol Evol Syst* **48**: 377–398.

Marlow H, Arendt D. 2014. Evolution: Ctenophore genomes and the origin of neurons. *Curr Biol* **24**: 757–761.

Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, Tosatto SCE, Paladin L, Raj S, Richardson LJ, et al. 2021. Pfam: The protein families database in 2021. *Nucleic Acids Res* **49**: D412–D419.

Moroz LL, Romanova DY. 2023. Homologous vs. homocratic neurons: revisiting complex evolutionary trajectories. *Front Cell Dev Biol* **11**: 1–6.

Moroz LL, Romanova DY. 2022. Alternative neural systems: What is a neuron? (Ctenophores, sponges and placozoans). *Front Cell Dev Biol* **10**: 1–19.

Moroz LL. 2021. Multiple Origins of Neurons From Secretory Cells. *Front Cell Dev Biol* **9**: 1–9.

Moroz LL. 2015. Convergent evolution of neural systems in ctenophores. *J Exp Biol* **218**: 598–611.

Najle SR, Grau-Bové X, Elek A, Navarrete C, Cianferoni D, Chiva C, Cañas-Armenteros D, Mallabiabarrena A, Kamm K, Sabidó E, et al. 2023. Stepwise emergence of the neuronal gene expression program in early animal evolution. *Cell* **186**: 4676-4693.e29.

Paysan-Lafosse T, Blum M, Chuguransky S, Grego T, Pinto BL, Salazar GA, Bileschi ML, Bork P, Bridge A, Colwell L, et al. 2023. InterPro in 2022. *Nucleic Acids Res* **51**: D418–D427.

Philippe H, Brinkmann H, Copley RR, Moroz LL, Nakano H, Poustka AJ, Wallberg A, Peterson KJ, Telford MJ. 2011. Acoelomorph flatworms are deuterostomes related to Xenoturbella. *Nature* **470**: 255–258.

Pijuan-Sala B, Griffiths JA, Guibentif C, Hiscock TW, Jawaid W, Calero-Nieto FJ, Mulas C, Ibarra-Soria X, Tyser RCV, Ho DLL, et al. 2019. A single-cell molecular map of mouse gastrulation and early organogenesis. *Nature* **566**: 490–495.

Piovani L, Leite DJ, Guerra LAY, Simpson F, Musser JM, Salvador-Martínez I, Marlétaz F, Jékely G, Telford MJ. 2023. Single-cell atlases of two lophotrochozoan larvae highlight their complex evolutionary histories. *Sci Adv* **9**: 1–15.

Pisani D, Pett W, Dohrmann M, Feuda R, Rota-Stabelli O, Philippe H, Lartillot N, Wörheide G. 2015. Genomic data do not support comb jellies as the sister group to all other animals. *Proc Natl Acad Sci* **112**: 15402–15407.

Ryan JF, Chiodin M. 2015. Where is my mind? How sponges and placozoans may have lost neural cell types. *Philos Trans R Soc B Biol Sci* **370**.

Ryan JF, Pang K, Schnitzler CE, Nguyen AD, Moreland RT, Simmons DK, Koch BJ, Francis WR, Havlak P, Smith SA, et al. 2013. The genome of the ctenophore Mnemiopsis leidyi and its implications for cell type evolution. *Science (80- )* **342**: 1336.

Sachkova MY, Nordmann EL, Soto-Àngel JJ, Meeda Y, Górski B, Naumann B, Dondorp D, Chatzigeorgiou M, Kittelmann M, Burkhardt P. 2021. Neuropeptide repertoire and 3D anatomy of the ctenophore nervous system. *Curr Biol* **31**: 5274-5285.e6.

Schierwater B, Eitel M, Jakob W, Osigus HJ, Hadrys H, Dellaporta SL, Kolokotronis SO, DeSalle R. 2009. Concatenated analysis sheds light on early metazoan evolution and fuels a modern "urmetazoon" hypothesis. *PLoS Biol* **7**.

Schultz DT, Haddock SHD, Bredeson J V, Green RE, Simakov O. 2023. Ancient gene linkages support ctenophores as sister to other animals. *Nature*.

Sebé-Pedrós A, Chomsky E, Pang K, Lara-Astiaso D, Gaiti F, Mukamel Z, Amit I, Hejnol A, Degnan BM, Tanay A. 2018. Early metazoan cell type diversity and the evolution of multicellular gene regulation. *Nat Ecol Evol* **2**: 1176–1188. http://dx.doi.org/10.1038/s41559-018-0575-6.

Sebé-Pedrós A, Saudemont B, Chomsky E, Plessier F, Mailhé MP, Renno J, Loe-Mie Y, Lifshitz A, Mukamel Z, Schmutz S, et al. 2018. Cnidarian Cell Type Diversity and Regulation Revealed by Whole-Organism Single-Cell RNA-Seq. *Cell* **173**: 1520-1534.e20.

Siebert S, Farrell JA, Cazet JF, Abeykoon Y, Primack AS, Schnitzler CE, Juliano CE. 2019. Stem cell differentiation trajectories in Hydra resolved at single-cell resolution. *Science (80- )* **365**.

Simion P, Philippe H, Baurain D, Jager M, Richter DJ, Di Franco A, Roure B, Satoh N, Quéinnec É, Ereskovsky A, et al. 2017. A Large and Consistent Phylogenomic Dataset Supports Sponges as the Sister Group to All Other Animals. *Curr Biol* **27**: 958–967.

Steenwyk JL, King N. 2024. The promise and pitfalls of synteny in phylogenomics. *PLoS Biol* **22**: 1–23. http://dx.doi.org/10.1371/journal.pbio.3002632.

Whelan N V, Kocot KM, Moroz LL, Halanych KM. 2015. Error, signal, and the placement of Ctenophora sister to all other animals. *Proc Natl Acad Sci U S A* **112**: 5773–8. http://www.ncbi.nlm.nih.gov/pubmed/25902535%5Cnhttp://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4426464.

Yaron A, Sprinzak D. 2012. The cis side of juxtacrine signaling: A new role in the development of the nervous system. *Trends Neurosci* **35**: 230–239. http://dx.doi.org/10.1016/j.tins.2011.12.003.