Advancing Ligand Binding Affinity Prediction with Cartesian Tensor-Based Deep Learning

Jie Yu,[∘,1,2,3] Xia Sheng,[∘,1,4] Zhehuan Fan,[∘,1,4] Zhaokun Wang,[∘,1,4] Duanhua Cao,[∘,5] Yongxin Hao,[1,6] Yingying Zhang,[1,6] Panpan Shao,[1,3,7] Huicong Ma,[8,9] Tian Cao,[9,10] JingXin Rao,[1,4] Mingan Chen,[1,3,11] Kaixian Chen,[1,4] Xutong Li,[1,4] Dan Teng,[1,4] Xiaomin Luo,[1,4] Mingliang Wang,[*,8,9,12] Sulin Zhang[*,1,4], Mingyue Zheng[*,1,2,4]

[1]Drug Discovery and Design Center, State Key Laboratory of Drug Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, 555 Zuchongzhi Road, Shanghai 201203, China

[2]School of Information Science and Technology, Shanghai Tech University, Shanghai, 201210, China

[3]Lingang Laboratory, Shanghai 200031, China

[4]University of Chinese Academy of Sciences, No. 19A Yuquan Road, Beijing 100049, China

[5]School of Life Sciences and Technology, Tongji University, Shanghai, 200092, China

[6]Division of Life Science and Medicine, University of Science and Technology of China, Hefei, 230026, Anhui, China.

[7]School of Chinese Materia Medica, Nanjing University of Chinese Medicine, Nanjing 210023, China

[8]School of Pharmacy, Zunyi Medical University, Zunyi, 563000, China

[9]Zhongshan Institute for Drug Discovery, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Zhongshan Tsuihang New District, Guangdong 528400, China

[10]State Key Laboratory of Discovery and Utilization of Functional Components in Traditional Chinese Medicine, School of Pharmaceutical Sciences, Guizhou Medical University, Guiyang 550014, China

[11]School of Physical Science and Technology, Shanghai Tech University, Shanghai, 201210, China

[12]Department of Medicinal Chemistry，Shanghai Institute of Materia Medica, Chinese Academy of Sciences 555 Zu Chong Zhi Road, Shanghai 201203, China

Corresponding Author

[*](Mingliang Wang) E-mail: wangmingliang@simm.ac.cn

[*](Sulin Zhang) E-mail: slzhang@simm.ac.cn

[*](Mingyue Zheng) E-mail: myzheng@simm.ac.cn

Author Contributions

[∘]J.Y., X.S., Z.H.F., Z.K.W., and D.H.C. contributed equally to this work. M.Y.Z. and J.Y. designed the research study. J.Y. and X.S. developed the method and wrote the code. M.L.W., H.C.M, and T.C. conducted chemical synthesis. All authors contributed to the analysis of the results. J.Y., M.Y.Z. and X.S. wrote the paper. All authors read and approved the manuscript.

Note

The authors declare no competing financial interest.

0. Abstract

We present PBCNet2.0, a cartesian tensor-based Siamese Neural Network for protein-ligand relative binding affinity prediction. Trained on 8.6 million protein-ligand complex structure pairs, PBCNet2.0 achieves zero-shot performance comparable to computationally intensive physics-based simulations. Our prioritization experiments show that PBCNet2.0 speeds up binding affinity optimization by 718% while reducing resource use by 41%. Through extensive retrospective experiments, we demonstrate that PBCNet2.0 intrinsically comprehends protein-ligand interactions, showing high sensitivity to intermolecular interactions and exceptional perception of spatial geometric information. Strikingly, PBCNet2.0 exhibits an emergent capability to predict affinity changes induced by binding residue variations, highlighting its potential for identifying resistance mutation. We prospectively validated these capabilities on two targets ENPP1 and ALDH1B1, where PBCNet2.0 successfully identified affinity shifts arising from subtle molecular interactions and conformational differences, and pinpointed critical binding residues with an 83% hit rate. This combination of computational efficiency, spatial geometric perception of binding site, and generalizable affinity prediction establishes PBCNet2.0 as a transformative tool for developing pharmacological probes for all human proteins.

1. Introduction

The human genome harbors a vast number of proteins that remain underexplored despite their potential therapeutic value[1,2]. Many of these proteins are poorly understood and have not been adequately studied, yet they hold significant promise for drug development. A key factor contributing to this underutilization is the lack of high-quality, well-characterized molecular probes[1]. Recognizing this gap, the global initiative Target 2035 has been launched with the ambitious goal of identifying pharmacological probes for all human proteins[3,4].

Within this context, protein-ligand binding affinity prediction[5–7] plays crucial roles in accelerating the development of molecular probes. While physics-based methods such as free energy perturbation (FEP, e.g., Schrödinger's FEP+[8]) have achieved chemical accuracy (root-mean-square error $\sim 1.1 \, \text{kcal} \cdot \text{mol}^{-1}$) in predicting relative binding affinities, their substantial computational cost, complexity in system setup, and dependency on expert intervention limit their broad applicability[9]. Consequently, artificial intelligence (AI), especially deep learning, presents a compelling opportunity for high-throughput and accurate affinity predictions by learning directly from structural data.

Recent advances in deep learning models that incorporate three-dimensional (3D) structural information have demonstrated promising results in protein-ligand binding prediction[5,10]. Our group developed EquiScore, an equivariant graph neural network that integrates physical principles of ligand-protein interactions, showing robust performance in virtual screening tasks[5]. In parallel, Moon et al. developed PIGNet2.0, a physics-informed graph neural network that explicitly models various protein-ligand interactions to predict binding free energies, proving effective in both virtual screening and lead optimization[11]. We previously introduced the Pairwise Binding Comparison Network (PBCNet), which employs a Siamese Neural Network (SNN) architecture enhanced with physics-informed graph attention mechanisms to rank binding affinities among structurally similar ligands[6]. PBCNet processes pairs of protein-ligand complexes to predict relative binding affinities, with affinity labels derived from identical assays to minimize systematic errors common in large-scale affinity databases[12]. While PBCNet showed significant advantages over existing high-throughput methods and achieved comparable performance to Schrödinger's FEP+ after fine-tuning,

there remains a performance gap in zero-shot predictions.

We identified two primary limitations. First, PBCNet's training dataset, though carefully

curated, encompasses only ~20,000 unique small molecules and 960 protein targets, representing a

limited chemical and biological space. Recent research on scaling laws[13,14], validated across

multiple domains including protein structure prediction[15] and large language models[16], indicates

that model performance correlates positively with data volume and model complexity. The restricted

scale of PBCNet's training set may therefore limit its generalization across diverse chemical

structures and protein families. Second, PBCNet's geometric encoding approach introduces inherent

constraints. The model utilizes atom pairwise statistical potentials[17] (APSP) and angular encoding

modules to guide attention weights during message passing. While these components implement

physical principles such as distance-dependent interactions and directional bonding, they rely on

predetermined geometric features from known atom pairs and fixed angle bins. This methodology

potentially limits the model's ability to learn context-specific interactions from structural data and

may be suboptimal for novel atom types not included in the predefined APSP set[18]. These geometric

constraints could impede the detection of unusual binding patterns, particularly in systems with non-

standard ligand chemistry[19].

To address these limitations and advance the field of protein-ligand binding affinity prediction,

we present PBCNet2.0, a next-generation model leveraging equivariant graph neural networks (E-

GNNs)[20]. Our approach is guided by two key insights from recent advances in artificial intelligence.

First, inspired by scaling laws, we expanded our training dataset to encompass 8.6 million protein-

ligand complex pairs from BindingDB[21], representing a 14-fold increase over PBCNet's original

dataset. This expanded dataset covers a broader chemical and biological landscape, with 0.28

million unique small molecules and 1,122 distinct protein targets, enabling more comprehensive

learning of diverse interaction patterns. Second, we fundamentally redesigned our model

architecture by implementing a Cartesian-based E-GNN[22] framework for message passing. This

approach naturally encodes three-dimensional structural relationships through equivariant

transformations, moving beyond PBCNet's reliance on predefined potential functions to learn

geometric constraints directly from data. This study presents our investigation into these innovations

through several key aspects:

First, we evaluate PBCNet2.0's performance in lead optimization tasks, comparing it with

existing computational methods including physics-based approaches and AI models. We then

analyze the model's ability to capture and interpret protein-ligand interactions through detailed

ablation studies and geometric perturbation experiments. The discussion extends to an unexpected

emergent capability: the prediction of binding affinity changes induced by pocket mutations, which

has notable implications for understanding drug resistance[23–26]. Finally, we validate our findings

through prospective experiments on two drug targets, ectonucleotide

pyrophosphatase/phosphodiesterase 1 (ENPP1)[27] and aldehyde dehydrogenase 1B1 (ALDH1B1)[28],

demonstrating the practical utility of our approach in real-world molecular probe development and

lead optimization scenarios.

2. Results

2.1 PBCNet2.0: a structure-based $\Delta pAct$ predictor

Building upon the Siamese Neural Network (SNN) architecture of PBCNet, PBCNet2.0 introduces an equivariant message passing module to analyze protein-ligand binding interfaces and predict relative binding affinity changes. Here, we describe the model's training methodology, architectural design, and intended applications. Throughout our analysis, we use $\Delta pAct$ to denote the negative logarithm of relative binding affinity, applicable across multiple experimental measurements (IC_{50}, EC_{50}, K_i, and K_d).

Recent studies of scaling laws in machine learning have demonstrated that model performance consistently improves with increased training data volume and model complexity, providing empirical guidance for AI system optimization. Based on this principle, we assembled a comprehensive training dataset of 8.6 million protein-ligand complex pairs (Fig. 1a), a 14-fold increase from PBCNet's original dataset (0.6 million). The ***Training set*** section details the data processing methodology.

The PBCNet2.0 architecture consists of three primary components: (1) a message-passing module, (2) a readout module, and (3) a prediction module, as illustrated in Figure 1b. The message-passing module facilitates information transfer across protein-ligand interaction graphs to generate node-level representations. The key advancement in PBCNet2.0 lies in the integration of an equivariant graph neural network[20] with Cartesian tensors[22], enabling comprehensive analysis of distance-dependent and angular geometric information in protein-ligand interactions and molecular conformations.

In the implementation, each node's hidden representation is encoded as a Cartesian rank-2 tensor, represented by a 3×3 matrix. This matrix is decomposed into three independent 3×3 matrices with 1, 3, and 5 degrees of freedom (detailed in Equation 1, ***Methods***), corresponding to scalar, vector, and tensor features. These features undergo parallel updates through 3×3 matrix operations, maintaining model equivariance efficiently. During message passing, protein pocket and ligand atom representations are updated through their interactions, after which pocket atoms are removed while ligand atoms are retained.

The readout module generates graph- and pair-level representations. Graph-level

representations (x_{ref} and x_{query}) are computed by averaging the invariant node representations

of reference and query ligands. The pair-level representation is defined as $\Delta x = x_{ref} - x_{query}$,

which differs from PBCNet's concatenation of x_{ref}, x_{query}, and Δx. This modification enhances

the model's focus on protein-ligand interactions, as structural similarity between analogous ligands

would otherwise result in Δx approaching zero, creating computational challenges. The pair-level

representation is processed through a three-layer feedforward neural network in the prediction

module to estimate relative binding affinity ($\Delta \hat{y}$). To ensure anti-symmetry ($f(\Delta x) = -f(-\Delta x)$),

we incorporate $-\Delta x = x_{query} - x_{ref}$ during training and compute the combined loss. Further

technical details are provided in the ***Methods*** section.

For inference, PBCNet2.0 requires only the binding poses of two structurally similar molecules

bound to the same protein target to predict their relative binding affinity (Fig. 1c). This enables

systematic ranking of compound series based on predicted binding affinities, facilitating rational

lead optimization and probe development. Notably, PBCNet2.0 exhibits an additional capability:

when presented with structures of an identical ligand bound to protein pockets containing specific

amino acid mutations, the model can predict the resulting changes in binding affinity (Fig. 1c). This

capability will be thoroughly examined in subsequent sections.

Figure 1   a. Processing of training data for PBCNet2.0.   The raw data was collected from BindingDB (2023.12 version), and input binding poses were generated through docking and rule-based filtering. PBCNet2.0 is trained on 8.6 million protein-ligand pairs. The training pairs share the same protein pocket but differ in ligand structures. **b. The framework of PBCNet2.0.** The complexes are processed through a Cartesian rank-2 Tensor-based message-passing module, designed to facilitate mutual information exchange between the ligands and the protein pocket. This module yields node-level

representations. In this process, each hidden dimension of each node-level representation is represented as a 3×3 matrix, which can be decomposed into three matrices with degrees of freedom 1, 3, and 5, respectively, using the irreducible representation of tensors. This decomposition enables efficient equivariant information propagation through matrix multiplication. In the Readout module, molecular representations (graph-level) are obtained by averaging the scalar node-level representations of the ligand atoms. The difference between the representations of the reference and query molecules (x^{ref} and x^{que}) serves as the final pair-wise representation (Δx), which is then fed into the Prediction module to output the predicted relative binding affinity $\Delta \hat{y}$. A computationally efficient sign-flip augmentation strategy is also implemented here. **c. The application of PBCNet2.0.** During inference, the PBCNet2.0 framework requires only the binding poses of two structurally similar small molecules bound to the same protein to predict their relative binding affinity. Additionally, when given structures with identical ligand molecules bound to protein pockets containing specific residue mutations, the model can predict the resulting changes in binding affinity.

2.2 PBCNet2.0 Achieves Substantial Improvement in Lead Optimization Tasks with Perception of Protein - Ligand Interaction Information

Following our previous work on PBCNet[6], we first systematically evaluated the performance of PBCNet2.0 in lead optimization tasks through three experimental paradigms: (1) **Zero-shot prediction**, which assess guidance capability during early-stage optimization when no structure-activity relationship (SAR) data is available; (2) **Few-shot prediction**, which evaluates performance in mid-to-late-stage optimization where a modest amount of SAR data enables model fine-tuning; and (3) **Prioritization experiment**, which validates the model's ability to identify high-activity compounds in real-world lead optimization scenarios. All metrics and methodologies matched our previous work[6], as summarized them in the Supplementary Section 1.

The zero-shot prediction results, presented in Figure 2a and Supplementary Figure 1, show the average performance of various methods across 16 chemical series referred to as the FEP test set (combining FEP1 and FEP2 sets). We assessed ranking performance using Pearson's correlation coefficient (R) and Spearman's rank correlation coefficient (ρ) (refer to ***Evaluation metrics*** section). We compared against twelve baseline methods (detailed in ***Baseline models*** section): a deep learning based co-folding model (Boltz-2[29]), three traditional structure-based approaches (Glide SP[30], MM-GB/SA[31], and Schrödinger's FEP+[8]), three sequence-based AI models (PSICHIC[7], BIND[32], and PLAPT[33]), and five structure-based AI models (PIGNet2[11], RTMScore[34], GenScore[35], OnionNet-2[36], and PBCNet[6]). The detailed prediction results of each method on each chemical

series are summarized in the Supplementary Data 1.

Benchmarking results establish PBCNet2.0's superior predictive capability, achieving $\rho = $

0.67 and $R = 0.66$ on the FEP set, an 18% enhancement over its predecessor PBCNet ($\rho = 0.57$,

$R = 0.56$). Notably, PBCNet2.0 approaches the performance of the gold-standard Schrödinger

FEP+ ($\rho = 0.70$ and $R = 0.70$, $\Delta\rho = 0.03$ and $\Delta R = 0.04$), with statistically indistinguishable

ranking accuracy ($p = 0.39 > 0.05$). This result makes PBCNet2.0 the first AI approach to achieve

parity with industrial-strength FEP simulations in critical evaluation metrics.

To visualize PBCNet2.0's improvements, we compared predicted ΔpAct values from both

PBCNet2.0 and PBCNet for test systems from the FEP set (Supplementary Fig. 2). PBCNet2.0's

predictions exhibit a stronger correlation with experimental values and cover a broader value range.

For example, in the pfkfb3 system, PBCNet2.0's predictions span [-2, 2], closely matching the

experimental value range, while PBCNet's predictions remain limited to [-1, 1].

Figure 2b illustrates the few-shot prediction results, with detailed data summarized in

Supplementary Data 2. The x-axis indicates the number of compounds used for fine-tuning, while

the y-axis shows the ranking capability metrics. Cyan dots represent PBCNet's performance, red

dots show PBCNet2.0's performance, and a blue dashed line indicates the Schrödinger's FEP+

baseline. Fine-tuning proves highly effective for PBCNet2.0: its performance improves steadily as

more fine-tuning ligands are added, consistently outperforming both PBCNet and Schrödinger's

FEP+. This data-efficient refinement capability makes PBCNet2.0 valuable when limited SAR data

is available for a target of interest.

The real-world utility of PBCNet2.0 is demonstrated through retrospective prioritization

experiments across 8 pharmaceutically relevant targets (Fig. 2c). The detailed data and evaluation

process are summarized in Supplementary Data 3 and the Supplementary Section 1. Benchmarking

results show that PBCNet2.0 outperforms both PBCNet and MM-GB/SA on all metrics. Specifically,

PBCNet2.0 achieved an **advantage order** of 13.57, while compared to PBCNet's 11.18 and MM-

GB/SA's 5.89. The **advantage ratio**, which measures theoretical resource savings in model-guided

lead optimization, shows PBCNet2.0 reduced resource investment by 40.51%, surpassing PBCNet

(30.18%) and MM-GB/SA (14.93%). The **efficiency improvement ratio**, reflecting enhanced

efficiency in compound optimization, shows PBCNet2.0 accelerated lead-optimization projects by

717.81%, surpassing both PBCNet (473.00%) and MM-GB/SA (281.31%). Statistical analysis

confirms that the performance of PBCNet2.0 in the prioritization experiment is significantly (p ≤
0.05) superior to MM-GB/SA.

These comprehensive results demonstrate that PBCNet2.0 exhibits significant advances in
relative binding affinity prediction, with notable improvements in both accuracy and reliability. This
enhanced predictive capability positions PBCNet2.0 as a valuable tool for molecular probe
development and lead optimization. We next examine the underlying mechanisms contributing to
its predictive performance.

Mastropietro et al. reported that many machine learning-based scoring models for protein-
ligand binding suffer from a bias where predictions are predominantly driven by ligand
memorization rather than understanding protein-ligand interactions[37]. To rigorously test whether
PBCNet2.0 overcomes this bias. First, we conducted a strategic ablation study using the FEP dataset.
We systematically removed all protein-ligand interaction edges from the original molecular graphs
(**Complete graphs**), generating the **Incomplete graphs** (Supplementary Fig. 3a). We then
monitored the performance dynamics of PBCNet2.0 on both graphs over the course of training
(Supplementary Fig. 3b). During the initial phase of training (approximately up to 12.5K steps),
both curves exhibit a similar upward trend, suggesting that the model initially mainly relies on
molecular structure information for predictions. However, as training progresses beyond 12.5K
steps, a marked divergence emerges. The model performance on the **Incomplete graphs** declines
steadily, whereas that on the **Complete graphs** continues to improve. This pattern indicates that
protein-ligand interaction information has emerged as the primary determinant of PBCNet2.0's
predictions, implying that PBCNet2.0 may progressively develop an understanding of the
interaction information.

To further verify that PBCNet2.0 learns protein-ligand interaction patterns rather than
memorizing molecular features or structural variations, we evaluated its performance on the SAR-
Diff set, which was specifically designed to decouple structural modifications from binding affinity
changes (refer to ***SAR-Diff held-out*** test set section). Specifically, the SAR-Diff set comprises eight
pairs of chemical series (16 series total) where each pair share identical R-group modification but
exhibit distinct structure-activity relationships[38–40]. As illustrated in Figure 2d with an example from
the first series pair, and further demonstrated in Supplementary Figure 3c showing the absence of
activity correlation between series, this dataset design necessitates models to identify interaction-

dependent activity variations instead of relying on structural memorization. Therefore, successful

predictions on this benchmark require accurate modeling of the interactions between modified

chemical groups and their corresponding protein binding sites.

Figure 2e presents the comprehensive evaluation results (Supplementary Data 4), with heatmap

organized to show performance of different methods (y-axis) across series pairs (x-axis). Spearman

correlation coefficients (ρ) quantify prediction quality. The two numbers in each heatmap cell stand

for the ρ of a model on a given series pair. Each column indicates the ρ of different models on

the same chemical series, as illustrated by the arrows between Figure 2d and Figure 2e. When the

$\rho \geq 0.4$, the corresponding model is considered to have predictive power on the series. The darkest

green cells denote that models successfully predicted both series within a pair, lighter green indicate

models only own predictive ability for one series, and pale cells represent models have no predictive

ability for either series. PBCNet2.0 exhibited consistent predictive capability across all series pairs,

distinguishing itself from other methods, including Boltz-2 which succeeded in only four pairs.

Notably, both PBCNet2.0 and PBCNet were unique in avoiding complete prediction failures for any

series pair.

Figure 2f provides a detailed comparison of average model performance on these 16 series

(ranking performance measured by ρ and prediction accuracy measured by root-mean-square error,

r.m.s.e.). PBCNet2.0 demonstrated superior performance in both metrics. These results highlight

that PBCNet2.0 successfully masters the interactions between modified groups and protein pockets,

with protein-ligand interaction patterns serving as key determinants of prediction accuracy. A

comparison between Figure 2f and 2a reveals that the SAR-Diff set presented a more challenging

evaluation scenario than the FEP set for most models (with the exception of RTMScore), further

validating PBCNet2.0's robust performance under demanding conditions.

These collective results demonstrate that PBCNet2.0 serves as an effective computational tool

for lead optimization and molecular probe development. The model shows measurable

improvements over its predecessor PBCNet across multiple performance metrics, including ranking

accuracy and prediction capability. The mechanistic analysis reveals that PBCNet2.0 effectively

incorporates protein-ligand interaction information to enable accurate relative binding affinity

predictions.

304
305 **Figure 2 a. Direct prediction results.** The x-axis represents various binding affinity prediction
306 methods, while the y-axis denotes their average of their ranking capability (Spearman) on the FEP set. A
307 two-sided Wilcoxon signed-rank test compared each model against PBCNet2.0, with significant levels:

*p ≤ 0.05, **p ≤ 0.01, ***p ≤ 0.001, ****p ≤ 0.0001, and ns (not significant) for p > 0.05. Different model types are shown in different colors. **b. Finetuned prediction results.** The x-axis represents the number of compounds used for fine-tuning, while the y-axis shows the ranking capability metrics. Cyan dots represent PBCNet performance, pink dots show PBCNet2.0 performance, and a blue dashed line indicates the Schrödinger's FEP+ baseline. **c. Selection experiment results.** The x-axis represents different methods, and the y-axis shows performance across eight systems. Three panels display distinct evaluation metrics. A one-sided Wilcoxon signed-rank test evaluated whether PBCNet2.0 and PBCNet significantly outperformed MM-GB/SA. **d. Illustration of the partial modification process for a chemical series pair from the SAR-Diff Set.** A schematic representation of structural changes within a chemical series pair. **e. The predictive performance of the models on the SAR-Diff set.** The x-axis lists 8 series pairs, and the y-axis represents 11 prediction methods. Each cell in the heatmap displays the Spearman correlation coefficient for a model on a given series pair. Dark green cells indicate models successfully predicted both series within a pair, light green cells show models with predictive ability for only one series, and pale cells represent models without predictive ability for either series. **f-g. Average performance of methods on SAR-Diff set.** The x-axis represents various binding affinity prediction methods, while the y-axis denotes the average of their ranking capability (Spearman) and prediction accuracy (r.m.s.e with ΔpAct) on the SAR-Diff set. The one-sided Wilcoxon signed-rank test was used to analyze whether the ranking performance of PBCNet2.0 was significantly superior to others and whether its prediction error was significantly lower. *p ⩽ 0.05, **p ⩽ 0.01, ***p ⩽ 0.001, ****p ⩽ 0.0001, and ns (not significant) for p > 0.5. Different types of models are distinguished using bars of different colors.

     2.3 PBCNet2.0 Captures Subtle Intermolecular Interactions with Strict Geometric

Constraints

Following our demonstration that protein-ligand interaction information plays a crucial role in

PBCNet2.0's predictions, we sought to evaluate the model's capability in identifying specific

protein-ligand interactions and its utilization of geometric information. While our previous analysis

of PBCNet focused primarily on hydrogen bond (H-bond) interaction recognition[41], we extended

this investigation to PBCNet2.0. Through examination of the same cases analyzed in the PBCNet

study, we found that PBCNet2.0 maintains comparable effectiveness in H-bond interaction

identification, as evidenced in Supplementary Figure 4.

Beyond H-bonds, we investigated fluorine orthogonal multipolar interactions[42], an important

but frequently overlooked protein-ligand interaction type[43–45]. Traditional empirical expressions for

intermolecular interactions, including many general-purpose force fields, often inadequately

represent this interaction[43,46]. We investigated whether PBCNet2.0, utilizing a data-driven approach,

could learn and identify this interaction from comprehensive structural data. As illustrated in Figure

3a, fluorine's high electronegativity creates a nucleophilic center, while the carbonyl oxygen's

electron-withdrawing effect makes the carbonyl carbon electrophilic. This electronic

complementarity enables orthogonal multipolar interactions between fluorine and carbonyl carbon.

These interactions are characterized by specific geometric requirements: the fluorine-carbon

distance must be under 3.8Å, with the $F \cdots C = O$ angle approaching $90°$. Accurate detection of

such interactions requires precise modeling of both 3D spatial relationships and angular

dependencies.

To evaluate this capability systematically, we developed the F-Opt benchmark test set (see ***F-***

Opt test set section), comprising 10 selected compound pairs where fluorine substitution creates

orthogonal multipolar interactions that enhance binding affinity (Fig. 3b, Supplementary Fig. 5).

We assessed various binding affinity prediction methods using this benchmark. Figure 3c

(Supplementary Data 5) displays the absolute prediction errors for each model across compound

pairs (identified by PDB IDs), with darker shading indicating larger errors. White numbers denote

cases   where   predicted   and   experimental ΔpAct   values   show   opposite   signs

(Sign(Predicted ΔpAct) $\neq$ Sign(Experimental ΔpAct)). PBCNet2.0 demonstrated the highest

accuracy with the lowest mean absolute error ($\mathrm{m.a.e.} = 0.33$) among all methods. PBCNet

achieved the second-best performance ($\mathrm{m.a.e.} = 0.52$). Notably, only PBCNet2.0 and PBCNet

achieved 100% sign concordance, while other methods showed error rates between 20% and

50%, highlighting the reliability of both approaches.

To gain mechanistic insights, we conducted interpretability analyses on two systems from the

F-Opt benchmark (5G4O and 4E3N, detailed in ***Interpretability analysis*** section) to understand how

fluorine orthogonal multipolar interactions are recognized. As shown in Figure 3d and

Supplementary Figure 6, these interactions are represented by purple dashed lines. We annotated

protein atoms within 5Å of the fluorine atoms (central atoms) with weight values ($0.00 - 1.00$),

with boldface numbers indicating the interaction weights of participating carbon atoms. The

consistently high weights ($0.98 - 1.00$) assigned to these carbon atoms in both systems demonstrate

PBCNet2.0's ability to identify geometrically constrained fluorine interactions. To evaluate

PBCNet2.0's sensitivity to angular parameters, we conducted a perturbation study on the 5G4O

system. Using Maestro, we modified two key interaction angles by moving the central fluorine atom:

reducing one angle from $79.2°$ to $52.9°$ and increasing another from $95.6°$ to $133.7°$ (both

angles falling outside the acceptable range). Throughout this process, the ligand pose was adjusted

to maintain structural integrity. Figure 3e shows that the weight values for the interacting carbon

atoms decreased substantially (from 0.98 to 0.50 and from 1.00 to 0.56), demonstrating that the

equivariant framework enables PBCNet2.0 to recognize angular constraints. This geometric

sensitivity is essential for accurately modeling subtle intermolecular interactions.

Despite PBCNet's strong predictive performance on the F-Opt test set, its interpretability

analysis reveals limitations in mechanistic understanding (Supplementary Fig. 7a). Analysis of

PBCNet's statistical atom-pair potential function shows that this empirical approach assigns

unexpectedly low interaction probabilities to fluorine-carbonyl carbon pairs at relevant distances

(Supplementary Fig. 7c), indicating inadequate modeling of fluorine orthogonal multipolar

interactions. Additionally, PBCNet shows limited sensitivity to angular parameters (Supplementary

Fig. 7b). These limitations stem from its relatively coarse 3D modeling approach, which constrains

its ability to capture precise physicochemical interactions. In comparison, PBCNet2.0's geometry-

aware equivariant architecture, combined with extensive training data, enables more sophisticated

modeling of intermolecular interactions. This architectural advancement not only improves

predictive performance but also establishes a robust framework for learning complex interaction

patterns through data-driven approaches. The clear correlation between predictions and underlying

physicochemical principles enhances PBCNet2.0's utility for both probe development and lead

optimization.

	4NBL	4OG6	4E3N	5A4L	5G4O	4I80	3FLN	3OQK	4DKO	1OYT	Mean
Boltz-2	0.47	0.40	0.79	0.75	0.30	0.42	1.21	0.01	0.37	0.79	0.55
Glide SP	0.63	0.20	1.65	0.74	0.07	1.43	0.98	1.27	1.31	0.44	0.87
MM-GB/SA	1.78	3.23	1.99	1.59	0.18	1.45	3.25	2.56	0.47	1.66	1.82
PSICHIC	0.79	0.19	0.31	0.97	0.12	0.62	0.65	1.09	0.38	0.58	0.57
BIND	0.92	0.19	1.41	0.56	0.94	0.50	0.72	0.98	0.65	0.90	0.78
Plapt	0.38	0.34	2.04	0.89	0.66	0.46	0.01	0.43	1.49	0.20	0.69
PIGNet2.0	0.34	0.33	0.93	0.31	0.68	0.02	0.33	1.75	0.44	0.40	0.55
RTMScore	1.58	0.81	0.49	1.90	3.20	0.50	2.66	1.53	0.95	3.66	1.73
GenScore	4.67	2.91	0.84	1.35	1.43	3.76	3.10	2.19	1.44	0.12	2.18
Onionnet2.0	1.29	0.22	1.15	1.20	0.39	0.46	1.01	1.27	0.49	1.34	0.88
PBCNet	0.42	0.01	0.98	0.73	0.29	0.39	0.50	0.88	0.29	0.70	0.52
PBCNet2.0	0.38	0.24	0.62	0.10	0.72	0.33	0.24	0.10	0.19	0.38	0.33

Figure 3   a. Illustration of fluorine orthogonal multipolar interaction. A schematic demonstrating
its formation mechanism and geometric constraints. **b. Examples from the F-Opt Set.** Representative
cases showcasing fluorine orthogonal multipolar interactions, including key distances and angles. **c.
Absolute prediction errors of models on each compound pair:** The y-axis represents various methods,
and the x-axis corresponds to compound pairs (denoted by PDB IDs). The numbers in the table indicate

absolute errors, with white numbers highlighting incorrect predictions. **d. Interpretability analysis of the 5G4O system**. For a molecule with a trifluoromethyl R-group, two fluorine orthogonal multipolar interactions with the protein are identified. The binding mode and geometric constraints are illustrated, with fluorine orthogonal multipolar interactions represented by purple dashed lines. Protein atoms within 5 Å of the central fluorine atoms are annotated with weights, and bold numbers denote the weights of carbon atoms involved in the interactions. **e. Disruption of geometric constraints in fluorine orthogonal multipolar interaction**. The angles of fluorine orthogonal multipolar interactions were disrupted, changing one angle from 79.2° to 52.9° and another from 95.6° to 133.7° (both being unreasonable angles). Consequently, the weight distribution of surrounding atoms was altered.

2.4 PBCNet2.0 Demonstrates Emergent Abilities for Identifying Drug-Resistant Mutations

Residue mutations can directly impact protein structure, stability, function, and molecular interactions[24–26] (Fig. 4a). Residue mutations may alter binding site properties and drug binding affinities, potentially leading to drug resistance[23]. For instance, in non-small cell lung cancer (NSCLC) treated with first-generation EGFR inhibitors (e.g., Gefitinib[47]), the acquired T790M mutation in EGFR kinase domain emerges in $> 50\%$ of resistant cases within 9-14 months[48,49], sterically hindering drug binding and causing therapeutic failure. This rapid resistance crisis necessitated urgent development of next-generation inhibitors, exemplified by Osimertinib[50]—a third-generation EGFR inhibitor specifically engineered to overcome T790M-mediated resistance. Critically, the stronger the drug selection pressure, the more rapidly resistance develops because drug target has been proved to be very plastic[51]. Hence, an early prediction and identification of mutations that may potentially impact ligand binding is important for lead optimization and probe development. Although PBCNet2.0 was not explicitly trained on data involving amino acid mutations (the training set only contained data for different ligands binding to the same protein site), we were curious whether PBCNet2.0 could have automatically learned to predict mutation-induced binding affinity changes. This hypothesis was motivated by the shared nature of both tasks— modeling changes in protein-ligand interactions—and by PBCNet2.0's ability to capture subtle molecular interactions as demonstrated above. To rigorously test this hypothesis, we constructed a Mutation Benchmark Set (see ***Mutation test set*** section) containing 8 pharmaceutically relevant targets with clinically observed binding pocket mutations. As shown in Figure 4b (Supplementary

Data 6), PBCNet2.0 demonstrated relative solid predictive performance, achieving an average Spearman correlation of 0.53 ($\rho = 0.53$). This is a stark contrast to methods like Boltz-2, Glide SP, BIND, PLAPT, RTMScore, and GenScore, which even exhibited negative correlations. The failure of Boltz-2 on this task may be attributable to the co-folding method's difficulty in accurately predicting the structural changes caused by mutations in the amino acid sequence, thereby causing its affinity prediction module to completely fail.[52] A deeper target-level look into PBCNet2.0's predictions exposed varying performances across different targets. While PBCNet2.0 exhibited strong ranking performance for tACE[53] ($\rho = 0.76$) and HIV-1 PR[54] ($\rho = 0.75$) targets (Supplementary Fig. 8a), it showed less satisfactory results for AKR1B1[55] ($\rho = 0.28$) and PfDFHR-TS[56] ($\rho = 0.26$) targets (Supplementary Fig. 8b).

To further investigate the rationale behind PBCNet2.0's predictions, we conducted an interpretability analysis of **Lactobacillus casei Thymidylate Synthase** (LTS)[57] target. The N229C mutation abolished a key hydrogen bond between the ASN229 amide oxygen and ligand pyridine nitrogen (Fig. 4c). This mutation decreased the binding affinity ($\Delta pK_d = -0.62$)[57], closely matching PBCNet2.0's prediction ($\Delta pK_d = -0.79$). Additionally, we visualized the weights (details in ***Interpretability analysis*** section) assigned to neighboring atoms of the pyridine nitrogen during the model's message-passing phase (Fig. 4d). In Figure 4d, the hydrogen bond is depicted as green dashed lines, and protein atoms within 5 Å of the central nitrogen atom were annotated with weights, where boldface numeral indicates the weight of the oxygen atom involved in the interaction. We found that the involved oxygen atom was assigned a relative high weight (0.99) in wild-type simulations, with complete signal loss post-mutation. This phenomenon demonstrates the validity of PBCNet2.0's predictions. Through its acquired ability to capture protein-ligand interactions, the model has successfully generalized to mutation prediction tasks.

Thereafter, we retrospectively analyzed three cases of mutant-selective inhibitors that are either in clinical trials or already in the market: (1) Olutasidenib, approved by the Food and Drug Administration (FDA) in December 2022, targets the mutant **Isocitrate Dehydrogenase 1** (IDH1) (with $>$1000-fold selectivity) to treat acute myeloid leukemia (AML)[58]; (2) MRTX1133, a potent, noncovalent, and selective **Kirsten Rats Arcomaviral Oncogene Homolog** (KRAS) G12D inhibitor, entered clinical trials in March 2023 to address manifold tumors harboring KRAS[G12D] mutations[59]; and (3) Pirtobrutinib, the first-in-class noncovalent **Bruton's tyrosine kinase** (BTK)

inhibitor approved by the FDA[60]. The structures, binding modes, and experimental and predicted

relative binding affinities of these inhibitors are depicted in Figure 4e. As shown in Figure 4e,

PBCNet2.0 accurately predicted relative binding affinities across these structurally diverse

inhibitors, despite never being explicitly trained on mutation-specific data. This emergent

capability[61,62] likely stems from the model's generalized understanding of protein-ligand

recognition patterns, developed through training on a large-scale dataset containing 8.6 million

entries.

The above experimental results demonstrate that PBCNet2.0 can identify critical mutations,

positioning it as a versatile tool for both probe development and molecular optimization.

492
Figure 4   a. Illustration of amino acid mutation. A schematic representation of a mutation at position
113 in the protein pocket, where THR is mutated to TYR. This alters the protein pocket environment,
affecting small molecule binding and resulting in changes to binding affinity. **b. Average performance**
of methods on the Mutation set. The x-axis represents various binding affinity prediction methods,
while the y-axis denotes the average of their ranking capability (Spearmen) on the Mutation set. The one-
sided Wilcoxon signed-rank test was used to analyze whether the ranking ability of PBCNet2.0 was
significantly superior to others. *$p \leq 0.05$ and **$p \leq 0.01$. Different types of models are distinguished
using bars of different colors. **c. Binding Mode of the Mutation Case.** The mutation of ASN at position
229 to CYS causes the protein to lose a hydrogen bond with the small molecule, leading to a decrease in
pK_d by 0.62. PBCNet2.0 predicts a decrease of 0.79. **d. Interpretability analysis.** The hydrogen bond
is depicted as green dashed lines, and protein atoms within 5 Å of the central nitrogen atom were
annotated with weights, where the bold number indicates the weights associated with the oxygen atom

involved in the hydrogen bond. **e. Retrospective analysis of mutant-selective inhibitors.** Three mutant-
selective inhibitors that are either in clinical trials or already in the market are shown: Olutasidenib (target:
IDH1), MRTX1133 (target: KRAS), and Pirtobrutinib (target: BTK). For each compound, the figure
shows chemical structures on the left, binding modes in wild-type and mutant proteins in the middle
panels, and quantitative data on the right. The right section presents wild-type and mutant residues
alongside experimental and predicted ΔpIC_{50} values for each protein-ligand pair.

2.5 Experimental Validation

513 Building upon our retrospective evaluation and analysis of PBCNet2.0, we operationalized it
514 in a prospective experimental validation study to verify its capacity to capture subtle intermolecular
515 interactions and predict resistance-associated mutations that govern ligand binding.

2.5.1 Experimental Validation of Subtle Molecular Interaction Capture in ENPP1
Inhibitors

In our previous work, we identified compound **A1** (Fig. 5a) as a promising ENPP1 inhibitor[27],
and the binding mode analysis suggested it might form a fluorine orthogonal multipolar
interaction with the target protein. Here, we further investigate **A1** with detailed analysis. First, we
conducted an interpretability analysis on the fluorine of **A1** (Fig. 5b). Results showed a relative high
weight (0.94) for the potential dipole interaction between this fluorine and the target protein's
carbonyl oxygen. Then, based on **A1**, we designed an analog **A2** (Fig. 5a), replacing the fluorine
with hydrogen to eliminate the potential dipole effect while minimizing other structural changes.
Using PBCNet2.0, we predicted **A2**'s binding affinity for ENPP1 with **A1** as the reference.
PBCNet2.0 predicted a substantial difference in binding affinity between **A1** and **A2**, specifically
forecasting a 40-fold ($\Delta pAct = 1.61$) reduction in binding affinity upon fluorine-to-hydrogen
substitution. This marked disparity in predicted binding affinities provides strong computational
evidence supporting the existence of the fluorine orthogonal multipolar interaction.

To validate the accuracy of these computational results, we synthesized **A1** and **A2**
(Supplementary Section 3) and conducted corresponding experimental validations (Supplementary
Section 2). The protein thermal shift (PTS) assay revealed that both **A1** and **A2** increased the thermal
stability of ENPP1 protein (Fig. 5c), suggesting direct binding of these compounds to ENPP1. As

expected, **A1** indeed exhibited a stronger thermostabilizing effect on ENPP1 ($\Delta T_m = 3.8°C$) than

A2 ($\Delta T_m = 2.7°C$). *In vitro* enzyme activity assays showed that **A1** inhibited ENPP1 enzymatic

activity ($IC_{50} = 13.3nM$, Fig. 5d) obviously more potently than **A2** ($IC_{50} = 2284nM$, Fig. 5d).

These data suggested that **A1** possessed stronger binding affinity for ENPP1 protein. This was

confirmed by subsequent surface plasmon resonance (SPR) studies (Fig. 5e-f), which showed K_D

values of 35.3nM and 1030nM for **A1** and **A2** binding to ENPP1, respectively, indicating a

substantially higher binding affinity of **A1**. The $\Delta p K_D$ measured by SPR assay was 1.46, closely

aligning with PBCNet2.0's prediction of 1.61, with a minor error of 0.15. These experimental

observations collectively demonstrate the establishment of a robust fluorine-mediated orthogonal

multipolar interaction between compound **A1** and ENPP1.

This analysis provides compelling validation of PBCNet2.0's ability to accurately predict the

impact of subtle intermolecular interactions on protein-ligand binding affinity. The strong

correlation between computational predictions and experimental measurements supports the utility

of PBCNet2.0 as a reliable tool for rational structure-based probe development and lead

optimization.

2.5.2 Experimental Validation of Subtle Molecular Interaction Capture and Drug-

Resistant Mutation Prediction in ALDH1B1 Inhibitors

In our previous investigation, we identified a racemic compound exhibiting inhibitory activity

against ALDH1B1. Two critical questions remained to be addressed: (1) The potential differential

biological activities between stereoisomers, and (2) The identification of key amino acid residues

that could potentially confer drug resistance. Here, we employed PBCNet2.0 to systematically

investigate these aspects.

To investigate the first question, we performed computational analysis of the enantiomeric pair

(**B1** and **B2**, Fig. 5g). Molecular docking revealed that despite overall similar binding modes, a key

distinction arose from their stereochemical differences: **B2**'s pyridine ring could not establish the

favorable $\pi - \pi$ stacking interaction[63,64] with PHE296 that was observed with **B1** (Fig. 5h).

PBCNet2.0 predicted a significant difference in binding affinity, suggesting that **B1** would exhibit

approximately 10-fold higher potency compared to **B2** (predicted $\Delta pAct = 1.07$). Model

interpretability analysis supported this docking observation by highlighting the significance of the

$\pi - \pi$ stacking interaction between **B1**'s pyridine ring and PHE296 (Supplementary Fig. 9).

To validate these computational predictions experimentally, we synthesized the racemate and

separated the enantiomers **B1** and **B2** through chiral chromatography. Their absolute configurations

were determined by X-ray crystallography (Fig. 5i, Supplementary Section 4). PTS assays (Fig. 5j)

demonstrated that both compounds stabilized ALDH1B1, with **B1** showing notably higher

thermostabilization ($\Delta T_m = 8.8°C$) compared to **B2** ($\Delta T_m = 1.1°C$). Enzymatic assays (Fig. 5k)

revealed that **B1** inhibited ALDH1B1 ($IC_{50} = 20nM$) substantially more potently than **B2** ($IC_{50} = $

$5574nM$). SPR measurements (Fig. 5l-m) further confirmed this differential binding, yielding K_D

values of $15.2nM$ and $1210nM$ for **B1** and **B2**, respectively. The experimental binding affinity

differences aligned with PBCNet2.0's predictions, demonstrating the model's capability to identify

stereochemistry-dependent activity variations and capture subtle three-dimensional interaction

features.

To address the second question, we performed a systematic pocket-directed *in silico* alanine

scanning[65,66] by computationally mutating each binding pocket residue to alanine. PBCNet2.0 was

then employed to predict changes in binding affinity between each mutated ALDH1B1-**B1** complex

and the wild-type complex. Based on the predictions shown in Figure 5n and Supplementary Figure

10, six residues (PHE296, ASN457, LEU477, ILE458, PHE170, and GLU124) were selected for

experimental validation through site-directed mutagenesis and binding assays (Supplementary

Section 2). PTS assays and SPR analyses demonstrated that alanine substitutions at five of these

positions (all except GLU124) significantly impaired **B1** binding (Fig. 5o-t and Supplementary Fig.

11). This high success rate (5/6) in identifying functionally important residues from the selected

candidates demonstrates PBCNet2.0's potential utility in predicting resistance-conferring mutations.

For comparison, we also analyzed these residues using the conventional MM-GB/SA method, which

ranked PHE296 and PHE170 among its top predictions, while placing ASN457, ILE458, and

LEU477 at lower positions (Supplementary Fig. 10b).

The application of PBCNet2.0 led to three key findings: (1) Accurate binding mode prediction

through the integration of molecular docking and interpretable AI analysis; (2) Identification of a

stereo-chemically pure ALDH1B1 inhibitor, **B1**, with improved potency compared to its racemic

predecessor; (3) Successful prediction of potential resistance mutations via computational alanine

scanning. These results demonstrate PBCNet2.0's utility in providing structural and

pharmacophoric insights to guide rational probe development and structure-based lead optimization.

Figure 5. **a. Chemical structures of ENPP1 inhibitors.** Reference ligand **A1** (R = F) and query

ligand **A2** (R = H) where the fluorine atom was replaced with hydrogen. **b. Interpretability analysis.**

The fluorine orthogonal multipolar interaction between **A1** and TRP322 of ENPP1, with interaction

distance (d = 3.22Å) and angle (θ = 109.2°). Fluorine orthogonal multipolar interaction represented by

purple dashed lines. Protein atoms within 5 Å of the central fluorine atoms are annotated with weights,

and bold numbers denote the weights of carbon atoms involved in the interactions. **c. Protein thermal**

shift assay. Impact of **A1** (50 μM) and **A2** (50 μM) on the thermal stability of ENPP1 protein (2 μM)

as determined by PTS assay. Temperature (°C) on the x-axis and normalized signal (%) on the y-axis. **d.**

***In vitro* enzyme activity assay.** Dose-dependent inhibition of **A1** and **A2** against ENPP1. The substrate

for the ENPP1 enzymatic reaction is thymidine 5'-monophosphate p-nitrophenyl ester (TMP). Error bars

represent the mean ± SEM of three independent experiments. **e-f. Surface plasmon resonance binding**
assay. The binding kinetics measurement of **A1** and **A2** to ENPP1 protein were using SPR assay. Graphs
depicting equilibrium response units versus **A1** and **A2** concentrations were plotted. **g. Chemical**
structures of ALDH1B1 inhibitors. Chiral centers of Reference ligand **B1** and query ligand **B2** were
different. **h. Binding modes of B1 and B2.** While **B1** and **B2** adopt largely similar binding modes, a
critical difference emerged due to their distinct stereochemistry: the pyridine ring of **B2** is unable to form
the favorable $\pi - \pi$ stacking interaction with PHE296 that is observed for **B1**. **i. X-Ray Structural**
Data. Absolute configurations for **B1** and **B2**. **j. Protein thermal shift assay.** Impact of **B1** (50μM)
and **B2** (50μM) on the thermal stability of ALDH1B1 protein (5μM) Temperature (°C) on the x-axis and
normalized signal (%) on the y-axis. **k. *In vitro* enzyme activity assay.** Dose-dependent inhibition of **B1**
and **B2** against ALDH1B1. Error bars represent the mean ± SEM of three independent experiments. **l-m.**
Surface plasmon resonance binding assay. The binding kinetics measurement of **B1** and **B2** to
ALDH1B1 protein were measured using SPR assay. Graphs depicting equilibrium response units versus
B1 and **B2** concentrations were plotted. **n. Mutation prediction results of PBCNet2.0.** The x-axis
represents the value of the decrease in binding affinity predicted by the model after the mutation, and the
y-axis represents the mutated residue. The 15residues prioritized by the model (out of 43) are shown. **o-**
t. Protein thermal shift assay. Impact of **B1** on the thermal stability of various ALDH1B1 mutants.
623

3. Discussion

We present PBCNet2.0, an advanced deep learning framework designed to predict protein-ligand relative binding affinities. The model leverages an extensive training dataset of 8.6 million protein-ligand pairs and employs a Cartesian tensor-based equivariant network architecture. Our benchmarking demonstrates that PBCNet2.0 achieves comparable accuracy to Schrödinger's FEP+ simulations in zero-shot prediction scenarios, while maintaining computational efficiency. Through systematic evaluation, we show that PBCNet2.0 accurately captures protein-ligand interaction patterns and geometric features at the atomic level. Notably, the model exhibits emergent ability to predict binding affinity changes resulting from pocket residue mutations, despite not being specifically trained for this task. Both retrospective and prospective validation studies support PBCNet2.0's utility in expediting lead optimization and probe development processes.

While demonstrating significant capabilities, PBCNet2.0 faces certain methodological limitations common to contemporary structure-based deep learning approaches. A primary constraint lies in its reliance on static protein-ligand binding conformations, which precludes explicit consideration of entropic effects and conformational dynamics during the binding process. Given recent advances in protein-ligand co-folding methods[15,67], incorporating protein flexibility through integrated structure prediction approaches represents a promising avenue for future development.

Additionally, PBCNet2.0 shows limitations in predicting complete activity loss cases where R-group modifications lead to altered binding modes. This challenge arises from two main factors: First, the scarcity of experimentally validated negative data in public databases, especially regarding binding mode verification and target engagement confirmation. Second, the technical difficulties in accurately measuring binding affinities for compounds with very low activity. Since PBCNet2.0 is designed to compare binding affinity differences between compounds sharing similar binding modes, preliminary conformational analysis and pose validation can help filter out compounds with potentially disrupted binding modes, thereby avoiding this type of inaccurate predictions. Several promising research directions warrant further investigation. First, incorporating high-quality protein mutation datasets into PBCNet2.0's training could enhance the model's ability to capture specific

protein-ligand interactions. This integration may improve the model's generalization capabilities

across different protein targets and binding scenarios. Second, the current training framework of

PBCNet2.0, which focuses on learning generalizable protein-ligand interaction patterns, could serve

as an effective pretraining strategy. The pretrained model could then be fine-tuned for specific

downstream applications such as virtual screening, potentially addressing the common challenge of

ligand structure memorization in traditional training approaches. Third, rigorous validation through

real-world drug optimization projects would be essential to assess PBCNet2.0's practical utility.

Such applications would not only evaluate the model's performance but also identify specific areas

requiring improvement, thereby guiding its refinement for drug discovery applications.

4. Methods and Data

4.1 Datasets

4.1.1 Training set

The training dataset of PBCNet2.0 consists of 8.6 million protein-ligand complex pairs, processed through a systematic pipeline (Supplementary Fig. 12) comprising three sequential stages: (1) Data Collection, (2) Data Preprocessing, and (3) Three-Dimensional Structure Generation. In the initial data collection phase, we extracted approximately 2.81 million protein-ligand binding affinity measurements (IC_{50}, EC_{50}, K_i, and K_d) from the BindingDB Database (2023.12 version)[21].

In the data preprocessing phase, we systematically filtered and refined the dataset. Molecules with invalid SMILES[68] structures during RDKit[69] preparation were excluded. We removed affinity data entries marked as NaN, 0, or containing inequality symbols ('<' or '>'). The data were then organized into groups based on three key identifiers: 'Entry DOI', 'Target name', and 'Target source', ensuring that affinity measurements within each group corresponded to the same target and publication source. Within each group, molecules were clustered using a Tanimoto similarity threshold of 0.5 to establish chemical series of structural analogs. For molecules with multiple reported affinity values, we implemented a quality control protocol: when the maximum-to-minimum affinity ratio was below 3-fold, we used the arithmetic mean; otherwise, the molecule was excluded. For each compound series, we gathered relevant PDB entries based on target information. To select appropriate PDB structures for docking, we analyzed structural similarities between co-crystallized ligands and compounds in each series by calculating Tanimoto coefficients. The PDB structure containing the co-crystallized ligand with the highest similarity was selected for docking. To maintain data quality, chemical series without co-crystallized ligands showing similarity above 0.5 were excluded from further analysis.

Following the confirmation of ligands and PDB structures, we employed Glide[30] for generating 3D docking conformations. The ligands underwent preprocessing via the Schrödinger LigPrep module using default parameters. The protein structures were prepared using the Protein Preparation Wizard in Schrödinger suite according to standard protocols. Water molecules forming more than

three hydrogen bonds with either ligand or receptor atoms were preserved, and receptor grids were

generated centered on the co-crystallized ligand. Subsequently, docking was performed using the

Glide module with default parameters, generating up to 200 poses per ligand.

Given that structurally similar ligands from the same chemical series typically exhibit

comparable binding poses[6], a fundamental assumption in our modeling approach, pose selection

was carefully executed. For each series, we first identified the maximum common substructure

(MCS) between individual ligands and their corresponding co-crystallized reference. The root-

mean-square deviation (r. m. s. d.) was calculated between each docked pose and the experimentally

determined reference pose within the MCS region. Poses with $r. m. s. d. \leq 2.0$ Å were deemed

acceptable, indicating preservation of the reference binding mode. For ligands with multiple

acceptable poses, the one with the highest Glide score was selected. Ligands failing to yield

acceptable poses were excluded to maintain data integrity. All operations were conducted using

Schrödinger version 2022-3 via the Schrödinger Python API.

Following preprocessing, we systematically paired molecules within each chemical series,

implementing a maximum threshold of 15,000 pairs per series. To ensure comprehensive coverage,

priority was assigned to previously unpaired molecules during the pairing process. Analysis of the

resulting label distribution revealed a normal distribution pattern, with a notable concentration of

values within the $[-1, 1]$ range (Supplementary Fig. 13a, left). This concentration presented a

potential risk of model overfitting, as the model could achieve artificially low training error by

defaulting to predictions near the mean value (Supplementary Fig. 13b). To mitigate this risk, we

implemented a data balancing strategy, employing under-sampling in regions of high data density

and over-sampling in regions of low density. The resulting balanced dataset exhibited a more

uniform label distribution, as illustrated in Supplementary Figure 13a (right).

To ensure data independence, we excluded protein-ligand pairs that overlapped with the test

sets. A comprehensive comparison between the training sets of PBCNet2.0 and its predecessor

reveals substantial improvements in both target coverage and chemical diversity, as illustrated in

Supplementary Figure 13c.

We also addressed potential modification bias, where systematic changes in binding affinity

might correlate with specific R-group modifications (e.g., methyl to ethyl conversions). Such bias

could lead to model overfitting through simple structural pattern recognition rather than learning

meaningful protein-ligand interactions. Our analysis of binding affinity changes across pairs with

identical R-group modifications, presented in Supplementary Figure 13d, demonstrates normally

distributed activity changes centered around zero, confirming the absence of systematic

modification bias.

4.1.2 FEP held-out test set

For model evaluation, we utilized two established FEP benchmark datasets. The first dataset

(FEP1) from Wang et al.[8] contains eight chemical series with experimentally determined binding

free energies (ΔG) and corresponding FEP+ predictions. The binding free energies were converted

to pIC_{50} values under non-competitive binding assumptions using the equation:

$$pIC_{50} \approx -\log_{10}(e^{\frac{\Delta G}{RT}})$$

where R represents the gas constant (1.987×10^{-3} kcal $\cdot$ K^{-1} $\cdot$ mol^{-1}), T is the thermodynamic

temperature at 297 K, and e denotes Euler's number (2.718). The second dataset (FEP2) from

Schindler et al.[70] comprises eight congeneric series targeting pharmaceutically relevant proteins,

with experimental IC_{50} measurements. FEP2 exhibits enhanced molecular diversity compared to

FEP1, featuring variations in charge properties, ring systems, and core structures. Both datasets

underwent logarithmic transformation of binding metrics and standardized ligand pairing protocols

consistent with our training methodology. These datasets were combined to form the comprehensive

FEP set, with detailed information available in Supplementary Table 1.

4.1.3 SAR-Diff held-out test set

The SAR-Diff test set comprises eight pairs of chemical series, where each pair contains two

distinct series. While both series within a pair undergo identical R-group modifications, they exhibit

different structure-activity relationships and core scaffolds. This test set specifically evaluates the

model's ability to recognize and interpret protein-ligand interactions, as accurate predictions require

analysis of binding pocket interactions rather than simple molecular structure pattern recognition.

The dataset was curated from the ChEMBL database[71], with series selected from identical

assays (as identified by their ChEMBL assay ID) to minimize potential ranking errors within each

series. Following data acquisition and ligand structure extraction, we identified appropriate PDB

structures and performed docking procedures consistent with our training set protocol to generate

3D structural data. The complete dataset specifications and metadata are provided in Supplementary

Table 2.

4.1.4 Mutation test set

To evaluate the model's capacity for capturing protein-ligand interactions, we assembled a

mutation dataset consisting of eight targets from the mutation-induced drug resistance database

(MdrDB)[72]. The dataset was filtered to include only mutations within the protein-ligand binding

pocket that preserved the original interaction patterns (hydrogen bonds, hydrophobic interactions,

salt bridges, π-stacking, π-cation interactions, and halogen bonds). For each target system, we

examined a series of mutations while maintaining an identical binding ligand. Each series

incorporated a wild-type (WT) complex and multiple mutant (MT) complexes, with corresponding

relative binding energy measurements ($\Delta\Delta G$). Protein and ligand structures were prepared using

Schrödinger Maestro, followed by docking simulations to generate binding conformations. The

most probable conformations were selected based on their binding modes to establish the mutation

test set. The final dataset encompassed mutation data from 8 distinct targets, with comprehensive

details provided in Supplementary Table 3.

4.1.5 F-Opt test set

To evaluate the model's interpretability, we analyzed its performance using protein-ligand

fluorine orthogonal multipolar interactions as a model system. We curated a fluorine-focused

optimization (F-Opt) dataset from published studies, comprising 10 pairs of protein-ligand

complexes. In these pairs, ligand modifications from -H or -CH3 to -F or -CF3 groups led to

enhanced binding affinities through the formation of fluorine orthogonal multipolar interactions.

Two key criteria guided the dataset selection: (1) explicit literature documentation of fluorine

orthogonal multipolar interactions as the primary mechanism for improved activity, and (2)

availability of co-crystal structures for structural validation. Complete details of the protein-ligand

systems are provided in Supplementary Table 4.

4.2 PBCNet2.0 implement

PBCNet2.0 predicts relative binding affinities between pairs of protein-ligand complexes. At

its core, the model employs an equivariant graph neural network architecture with Cartesian tensor

representations, building upon TensorNet[22]. The model represents atomic interactions in Cartesian

coordinates using rank-2 tensors (3×3 matrices) as atomic descriptors. These Cartesian rank-2

tensors can be efficiently decomposed into irreducible components under rotational transformations

(Equation 1). This decomposition enables straightforward 3×3 matrix operations during the

message-passing phase, enhancing the computational efficiency of the equivariant network. The

following sections detail the irreducible decomposition of rank-2 Cartesian tensors and the

architectural components of PBCNet2.0.

4.2.1 Irreducible Cartesian rank-2 tensor decomposition

785   Here, X denotes a rank-2 tensor (3×3 matrix) defined in Cartesian coordinates, with X_{ij}

786 representing the value in the $i-$th row and $j-$th column. Any tensor X can be decomposed in

787 the following manner:

$$X = \frac{1}{3}\text{Tr}(X)\text{Id} + \frac{1}{2}(X - X^T) + \frac{1}{2}\left(X + X^T - \frac{2}{3}\text{Tr}(X)\text{Id}\right), \tag{1}$$

where $\text{Tr}(X) = \sum_i X_{ii}$ is the trace operator, X^T denotes the transpose of X and Id is the identity

matrix. The first term is proportional to the identity matrix, possessing only one degree of freedom.

This term is invariant under rotations and represents a scalar. The second term is a skew-symmetric

matrix and contains three independent components that transform as a vector under rotations. The

third term is a symmetric and traceless, with five independent components that represent a higher-

order tensor. From a representation theory perspective, Equation 1 shows that the 9-dimensional

Cartesian rank-2 tensor (X) can be decomposed into components of dimensions 1 (scalar), 3 (vector),

and 5 (tensor), respectively. We refer to X as a "full tensor", while its components are referred to

as scalar (I), vector (A), and tensor (S) features.

The tensor representation can be constructed from relative position vectors in neural networks.

For example, given a vector $\boldsymbol{r} = (r_x, r_y, r_z)$ (e.g., the difference between the coordinates of two

nodes in a graph), we can build a well-behaved tensor $X = I + A + S$ as follows:

801
$$I = f(\|\boldsymbol{r}\|)\text{Id}, A = \begin{pmatrix} 0 & r_z & -r_y \\ -r_z & 0 & r_x \\ r_y & -r_x & 0 \end{pmatrix}, S = \boldsymbol{r}\boldsymbol{r}^{\text{T}} - \frac{1}{3}\text{Tr}(\boldsymbol{r}\boldsymbol{r}^{\text{T}})\text{Id}, \tag{2}$$

where $\|\boldsymbol{r}\|$ is the length of $\boldsymbol{r}$, $f(\cdot)$ is mapping function, and $\boldsymbol{r}\boldsymbol{r}^{\text{T}}$ denotes the outer product of $\boldsymbol{r}$

with itself. We can also modify the full tensor by multiplying invariant quantities to each component,

$X' = f_I I + f_A A + f_S S$, where f_I, f_A, and f_S can be constants or learnable invariant functions (e.g.,

functions of distances). These modifications preserve the transformation properties of the tensor

under rotations.

4.2.2 Graph preprocessing and feature initialization

In our protein-ligand interaction graph construction, we defined the protein pocket as residues

within 8Å of the ligand. The graph representation treats each heavy atom in the pocket-ligand

complex as a node and each covalent bond as an edge. Additional distance edges were established

between protein-ligand and ligand-ligand node pairs with spatial distances ≤ 5.0Å. For protein-

protein node pairs, we implemented a dual-threshold approach: if either protein atom was connected

to a ligand atom via a distance edge, a distance edge was created when their spatial distance was

≤ 5Å; otherwise, a more stringent threshold of ≤ 3Å was applied. This hierarchical graph

construction methodology optimizes the representation of protein-ligand interactions while

efficiently reducing computational complexity by limiting distant protein-protein connections.

For feature initialization, while the original TensorNet architecture was limited to atom types

and edge lengths as node and edge initial features, we expanded the feature set to include

comprehensive chemical descriptors. The enhanced model incorporates nine distinct atomic features

(detailed in Supplementary Table 5) and various bond types (listed in Supplementary Table 6),

providing a more complete representation of atoms and their local chemical environments.

4.2.3 Initial tensor representations

PBCNet2.0 learns a Cartesian full tensor representation $X_i = \{X_i^0, X_i^1, \cdots X_i^d\}$ (d represents

the number of hidden dimensions) for an atom a_i. In the following, we describe the computational

process using X_i as an example. Atomic representation X_i can be decomposed into scalar, vector

and tensor contributions I_i, A_i, S_i via Equation 1. Conversely, these components can be merged

to form X_i.

We treat a_i as the central atom, $a_j \in \{a_{j_1}, a_{j_2}, \cdots, a_{j_n}\}$ (n is the number of neighbor atoms

of a_i) represents a neighbor atom of a_i, and $e_{ij} \in \{e_{ij_1}, e_{ij_2}, \cdots, e_{ij_n}\}$ is an edge between a_i and

a_j. The coordinates of a_i and a_j are presented as $\boldsymbol{r}_i$ and $\boldsymbol{r}_j$, the edge vector $\boldsymbol{r}_{ij} = \boldsymbol{r}_j - \boldsymbol{r}_i$ can

be obtained, and $\|\boldsymbol{r}_{ij}\|$ presents the length of e_{ij}. We initialize each edge scalar features by the

identity matrix $I_{ij}^{initial} = Id$, and each vector ($A_{ij}^{initial}$) and tensor ($S_{ij}^{initial}$) feature are initialized

using the normalized edge vectors $\hat{\boldsymbol{r}}_{ij} = \frac{\boldsymbol{r}_{ij}}{\|\boldsymbol{r}_{ij}\|} = (\hat{r}_{ij}^x, \hat{r}_{ij}^y, \hat{r}_{ij}^z)$ through Equation 2. Thus, the initial

full tensor representation of e_{ij} is $X_{ij}^{initial} = I_{ij}^{initial} + A_{ij}^{initial} + S_{ij}^{initial}$. Next, we aim to encode

bond- and atom-invariant information, including edge distances, edge and atom chemical features

(Supplementary Table 5 and 6), into the initial full tensor $X_{ij}^{initial}$. This information is encoded

using linear layers, with the distance information extended by a radial basis function (RBF) before

encoding:

839
$$\boldsymbol{Z}_i = Linear_1(\boldsymbol{z}_i), \boldsymbol{Z}_j = Linear_1(\boldsymbol{z}_j), \boldsymbol{Z}_{ij}^{atom} = \boldsymbol{Z}_i \oplus \boldsymbol{Z}_j,$$

840
$$\boldsymbol{Z}_{ij}^{chem} = Linear_2(\boldsymbol{z}_{ij}), \boldsymbol{Z}_{ij}^{dist} = \mathrm{RBF}(\|\boldsymbol{r}_{ij}\|),$$

841
$$\boldsymbol{Z}_{ij}^{bond} = \boldsymbol{Z}_{ij}^{chem} \oplus \boldsymbol{Z}_{ij}^{dist}, \tag{3}$$

842   where $\boldsymbol{z}_i$, $\boldsymbol{z}_j$ and $\boldsymbol{z}_{ij}$ are chemical features of a_i, a_j and e_{ij}, respectively. $\|\boldsymbol{r}_{ij}\|$ is the length

843   of e_{ij}. RBF($\cdot$) and $\oplus$ are the radial basis function and concatenate operation. $\boldsymbol{Z}_i$, $\boldsymbol{Z}_j$ and

844    $\boldsymbol{Z}_{ij}^{chem}$ are chemical embeddings of a_i, a_j and e_{ij}, while $\boldsymbol{Z}_{ij}^{dist}$ is distance embedding of e_{ij}.

845    $\boldsymbol{Z}_{ij}^{atom}$ and $\boldsymbol{Z}_{ij}^{bond}$ are atom pair-wise and bond-invariant representation of e_{ij}. Then $\boldsymbol{Z}_{ij}^{atom}$ and

846    $\boldsymbol{Z}_{ij}^{bond}$ are combined with $I_{ij}^{initial}$, $A_{ij}^{initial}$ and $S_{ij}^{initial}$ according to:

847
$$f_I^{initial} = Linear_3(\boldsymbol{Z}_{ij}^{bond}), f_A^{initial} = Linear_4(\boldsymbol{Z}_{ij}^{bond}), f_s^{initial} = Linear_5(\boldsymbol{Z}_{ij}^{bond}),$$

848
$$X_{ij} = \boldsymbol{Z}_{ij}^{atom} \times (f_I^{initial} I_{ij}^{initial} + f_A^{initial} A_{ij}^{initial} + f_s^{initial} S_{ij}^{initial}), \tag{4}$$

where X_{ij} is the invariant information-merged initial tensor representation of e_{ij}. It should be

noted that the original TensorNet[22] includes an additional distance decay function in Equation 4.

The assumption of distance decay function is that atoms that are farther apart have less influence on

each other, which obviously affects the identification of long-range interaction. Moreover, this

function also inevitably introduces a bias, where the correlation weights between atom pairs become

larger as their distance decreases, and vice versa. In PBCNet2.0, we have eliminated this function.

We then get atom-wise initial tensor representation of a_i by aggregating all neighboring edge

tensor representations:

857
$$X_i = \sum_{j \in \mathcal{N}(i)} X_{ij},$$

858   where $\mathcal{N}(i)$ means the neighbor set of the central atom a^i. At this point, Frobenius norm (an

859   invariant norm) $\|X\|_F = \text{Tr}(X^T X)$ of X_i is computed and fed to a series of layers:

860
$$f_i^I, f_i^A, f_i^S = SiLU(Linear_7(SiLU(Linear_6(LN(\|X_i\|_F))))),$$

where $LN(\cdot)$ and $SiLU(\cdot)$ are layer norm and SiLU activation. After the decomposition of this

atom-wise tensor representation X_i into its irreducible representations via Equation 1, f_i^I, f_i^A and

f_i^S are then used to modify component-wise linear combinations to obtain the final initial atomic

tensor representations $X_i = f_i^I \times Linear_8(I_i) + f_i^A \times Linear_9(A_i) + f_i^S \times Linear_{10}(S_i)$.

4.2.4 Tensor representations updating

PBCNet2.0 consists of three message passing layers. We denote the input of the first layer as

X_i^0 (i.e., the initial atomic tensor representation), the output of the first layer (which becomes the

input of the second layer) as X_i^1, and so on. To introduce the message passing process, we focus on

the transition from X_i^0 to X_i^1 as an example. Firstly, X_i^0 is normalized by

870
$$X_i^0 = \frac{X_i^0}{\|X_i^0\|_F + 1},$$

871   Then X_i^0 is decomposed into scalar, vector and tensor features $X_i^0 = I_i^0 + A_i^0 + S_i^0$. The obtained

872   features are transformed by independent linear combinations to get an updated tensor embedding

873    $X_i^{0\_ud}$ according to

874
$$X_i^{0\_ud} = Linear_{11}(I_i^0) + Linear_{12}(A_i^0) + Linear_{13}(A_i^0).$$

875   We then aim to merge the information from neighboring atoms used to further update $X_i^{0\_ud}$. Firstly,

876   the bond-invariant embeddings $\mathbf{Z}_{ij}^{bond}$ are concatenated with the $O(3)$-invariant scalar embedding

877   of a_i and a_j ($\mathbf{x}_i^{0\_scalar}$ and $\mathbf{x}_j^{0\_scalar}$, see *Scalar embedding extracting* section for more details),

878   and then are transformed into three invariant terms as follows:

879
$$\mathbf{x}_i^{0\_scalar} = SiLU(Linear_{14}(LN(\|I_i^0\|_F \oplus \|A_i^0\|_F \oplus \|S_i^0\|_F))),$$

880
$$\mathbf{x}_j^{0\_scalar} = SiLU(Linear_{14}(LN(\|I_j^0\|_F \oplus \|A_j^0\|_F \oplus \|S_j^0\|_F))),$$

$$x_{ij}^{0\_bond} = x_i^{0\_scalar} \oplus x_j^{0\_scalar} \oplus Z_{ij}^{bond}$$

$$f_{ij}^I, f_{ij}^A, f_{ij}^S = SiLU(Linear_{17}(SiLU(Linear_{16}(SiLU\left(Linear_{15}(x_{ij}^{0\_bond})\right)))))). \qquad (5)$$

Note that we also eliminated the application of distance decay function compared to the original

TensorNet here. Next, the messages M_{ij} sent from neighbors a_j to atom a_i are computed as:

$$X_j^{0\_ud} = I_j^{0\_ud} + A_j^{0\_ud} + S_j^{0\_ud},$$

$$M_{ij} = f_{ij}^I \times I_j^{0\_ud} + f_{ij}^A \times A_j^{0\_ud} + f_{ij}^S \times S_j^{0\_ud}$$

These messages are then aggregated:

$$M_i = \sum_{j \in \mathcal{N}(i)} M_{ij}$$

where $\mathcal{N}(i)$ means the neighbor set of the central atom a^i. Then updated tensor embedding $X_i^{0\_ud}$

is further updated by M_i with a parity-preserved form:

$$X_i^{0\_ud} = X_i^{0\_ud} \times M_i + M_i \times X_i^{0\_ud}.$$

Next, $X_i^{0\_ud}$ is further decomposed into scalar, vector, and tensor features ($I_i^{0\_ud}$, $A_i^{0\_ud}$, and $S_i^{0\_ud}$)

with Equation 1. These features are individually normalized according to:

$$I_i^{0\_ud} = \frac{I_i^{0\_ud}}{\left\|X_i^{0\_ud}\right\|_F + 1}, A_i^0 = \frac{A_i^{0\_ud}}{\left\|X_i^{0\_ud}\right\|_F + 1}, S_i^0 = \frac{S_i^{0\_ud}}{\left\|X_i^{0\_ud}\right\|_F + 1}.$$

These normalized features are then used to compute independent linear combinations:

$$X_i^{0\_ud} = Linear_{18}(I_i^{0\_ud}) + Linear_{19}(A_i^{0\_ud}) + Linear_{20}(S_i^{0\_ud}).$$

Finally, a residual message ΔX_i^0 is computed in a parity-preserving manner and used to update X_i^0

to get X_i^1:

$$\Delta X_i^0 = X_i^{0\_ud} + (X_i^{0\_ud})^2,$$

$$X_i^1 = X_i^0 + \Delta X_i^0.$$

Repeating the above process, we can get X_i^2 and X_i^3 for the subsequent layers.

4.2.5 Scalar embedding extracting

Since the protein-ligand binding affinity is a $O(3)$-invariant property, the Frobenius norms of

each component of X_i^3, which is also $O(3)$-invariant, are used to get final atomic scalar embedding

x_i^{scalar},

$$x_i^{scalar} = SiLU(Linear_{21}(LN(\|I_i^3\|_F \oplus \|A_i^3\|_F \oplus \|S_i^3\|_F))) \qquad (6)$$

4.2.6 Readout

After the above process, we obtain the final scalar embedding x_i^{scalar} of each atom. We then summed the scalar embeddings from the ligand atoms as the final complex embedding x^{scalar},

$$x^{scalar} = \sum_{i \in \mathcal{L}(lig)} x_i^{scalar}$$

where $\mathcal{L}(lig)$ is the ligand atom set of a pocket-ligand complex. Finally, we utilize the embedding of the reference complex x_{ref}^{scalar} to subtract that of the query complex x_{query}^{scalar}, thereby obtaining the difference embedding $\Delta x^{scalar} = x_{ref}^{scalar} - x_{query}^{scalar}$, which then be fed into a 3-layer perceptron to obtain the final prediction $\Delta \hat{y}$. If the binding affinity of the reference complex y_{ref} is known, we can get the predicted binding affinity of tested complex $\hat{y}_{query} = y_{ref} - \Delta \hat{y}$. In the training process, to ensure, to the extent possible, that $Model(\Delta x^{scalar}) = -Model(-\Delta x^{scalar})$, we input $-\Delta x^{scalar} = x_{query}^{scalar} - x_{ref}^{scalar}$ into the fully connected layer as well and calculate the loss together as a data augmentation strategy with a low computational overhead.

4.3 Baseline models

To thoroughly evaluate our method, we selected a range of representative baseline models. For traditional approaches, we chose Schrödinger's **Glide-SP**[30], **MM-GBSA**[31], and Schrödinger's **FEP+**[8]. The Glide-SP and MM-GBSA modules were used with Schrödinger's default parameters, while the performance of FEP+ is based on the reported results[8,70].

Boltz-2[29] is a recently published deep learning-based co-folding model. This model takes sequence representations of both the protein and the ligand as input and is capable of predicting the three-dimensional structure of the protein-ligand complex along with its binding affinity. The weight parameters and source code for Boltz-2 were obtained from its original publication. Boltz-2 reports the binding affinity value as $\log(IC_{50})$, derived from an IC_{50} measurement in micromolar units (μM). Therefore, we adopt '6 − predicted values' as the final prediction results.

For deep learning scoring methods, we considered both sequence-based and structure-based

approaches. The sequence-based methods include:

• **PSICHIC**[7]. It directly decodes protein-ligand interaction fingerprints from sequence data by

incorporating physicochemical constraints. The weights and code of PSICHIC is from its

original publication. The binding affinity values output by the model are pK_d or pK_i, and we

directly evaluate the model using its outputs.

• **BIND**[32]. It is a graph neural network that uses Evolutionary Scale Modeling 2[73] (ESM-2)

embeddings for proteins and graph-based representations for ligands to perform virtual

screening. We ran the model with the default parameters provided in the paper and evaluated

it using the IC_{50} values from the output.

• **PLAPT**[33]. It utilizes transfer learning from pretrained transformers like ProtBERT[74], requiring

only 1D protein and ligand string data for predictions. The model's weights are set to their

default values, and the predicted IC_{50} values are directly used for performance calculation.

The structure-based methods differ in terms of input forms (ranging from 2D contact features

to 3D graph structures), learning objectives (from direct binding affinity prediction to fine-grained

interaction modeling), and model architectures, providing a comprehensive evaluation benchmark.

These models comprise:

• **RTMScore**[34]. It predicts binding affinity by fitting a protein-ligand distance likelihood

potential, using a Mixture Density Network (MDN) framework. In the pretrained models

provided in the paper, we selected the best-performing model, rtmscore_model3. The paper

defines a custom affinity score, where higher values indicate stronger binding affinity.

• **GenScore**[35]. Building on RTMScore, the model adds a ranking loss term to the training loss

and incorporates a fine-tuning strategy. We evaluated the model using GatedGCN_ft_1.0,

which showed the strongest ranking ability according to the results. The output affinity values

are also custom-defined.

• **PIGNet2**[11]. The model focuses on modeling the interactions between protein-ligand pairs,

refining the types of interactions into hydrogen bonds, metal bonds, van der Waals forces, and

hydrophobic interactions. During replication of the paper's results, we found that the model

pda_3 had the performance most closely aligned with the reported results on the FEP test set,

so we selected this model for comparison. The predicted binding free energy (ΔG) is converted

into pIC$_{50}$.

● **OnionNet-2**[36]. It is a scoring model based on convolutional neural network to predict the

binding affinity pKd, where the protein-ligand interactions are characterized by the number of

contacts between protein residues and ligand atoms in multiple distance shells. We used the

default parameter weights for the model.

4.4 Evaluation metrics

Model performance was evaluated using four metrics: the Pearson correlation coefficient (r),

Spearman's rank correlation coefficient (ρ), root-mean-square error (r. m. s. e.), and mean-absolute

error (m. a. e.). The correlation coefficients range from 0 to 1, where 1 indicates perfect correlation.

The Pearson correlation measures the linear relationship between predicted and experimental

binding affinities, while Spearman's correlation assesses the monotonic relationship between

predicted and experimental rankings. Root-mean-square error measures the average magnitude of

prediction errors, where lower values indicate better prediction accuracy. Mean absolute error

represents the average absolute difference between predicted and experimental values, where lower

values indicate better performance.

4.5 Interpretability analysis

In the message passing module of PBCNet2.0, the pocket-ligand complex is deemed a whole

graph and information on atoms is transmitted across the molecule graph. This module provide

atom–atom pairwise interpretation of key inter-molecular interactions, which is helpful for guiding

the design a drug for a specific target. The invariant terms f_{ij}^I, f_{ij}^A, and f_{ij}^S obtained from the

Equation 5 can be seen as a measure of importance between atoms a_i and a_j. We extracted these

invariant terms from the first layer of the message-passing model and computed their average as

relevance weights f_{ij} using the formula:

986
$$f_{ij} = \frac{f_{ij}^I + f_{ij}^A + f_{ij}^S}{3}.$$

987 Next, we performed min-max normalization on the weights of all neighboring atoms to obtain the

988 final weights w_{ij} used in visualization:

$$f_{i\_min} = \min_{j \in \mathcal{N}(i)} f_{ij},$$

$$f_{i\_max} = \max_{j \in \mathcal{N}(i)} f_{ij},$$

$$w_{ij} = \frac{f_{ij} - f_{i\_min}}{f_{i\_max} - f_{i\_min}}.$$

The normalized weights w_{ij} have a minimum value of 0 and a maximum value of 1, facilitating their use in visualization. An ideal model will tend to assign a higher weight to the pair of atoms which may form a key inter-molecular interaction because they usually have a larger influence on each other's chemical environment and on ligand binding affinity.

Notably, because the positions of the hydrogen atoms depended heavily on the corresponding program used to add hydrogens, and due to the sheer number of hydrogen atoms which would substantially increase the computational load, we did not take hydrogen atoms into account in the modeling process. Here, for hydrogen-bond donors, the heavy atoms covalently linked with hydrogen atoms were chosen for further analysis, consistent with previous practices in PBCNet.

5. Data availability

The training data is from BindingDB source and can be found at Zendo (https://zenodo.org/records/15656365).

The test data are available under an MIT License via GitHub (https://github.com/YuJie-0202/PBCNet2.0).

6. Coda availability

The code of this work is available under an MIT License via GitHub at (https://github.com/YuJie-0202/PBCNet2.0).

7. Acknowledgments

We gratefully acknowledge financial support from Strategic Priority Research Program of the Chinese Academy of sciences (XDB0830200), National Natural Science Foundation of China (T2225002, 82273855, and 82474143), National Key Research and Development Program of China (2022YFC3400504 and 2023YFC2305904), LingangLaboratory (LGL-8888), Youth Innovation Promotion Association CAS (2023296), and Young Elite Scientists Sponsorship Program by CAST (2023ONRC001). We thank the staff members of the Large-scale Protein Preparation System at the National Facility for Protein Science in Shanghai (NFPS), Shanghai Advanced Research Institute, Chinese Academy of Science, China for providing technical support and assistance in data collection and analysis.

References

1. Oprea, T. I. *et al.* Unexplored therapeutic opportunities in the human genome. *Nat. Rev. Drug Discov.* **17**, 317–332 (2018).

2. Dewson, G., Eichhorn, P. J. A. & Komander, D. Deubiquitinases in cancer. *Nat. Rev. Cancer* **23**, 842–862 (2023).

3. Ackloo, S. *et al.* Target 2035 – an update on private sector contributions. *RSC Med. Chem.* **14**, 1002–1011 (2023).

4. Ackloo, S. *et al.* CACHE (Critical Assessment of Computational Hit-finding Experiments): A public–private partnership benchmarking initiative to enable the development of computational methods for hit-finding. *Nat. Rev. Chem.* **6**, 287–295 (2022).

5. Cao, D. *et al.* Generic protein–ligand interaction scoring by integrating physical prior knowledge and data augmentation modelling. *Nat. Mach. Intell.* **6**, 688–700 (2024).

6.   Yu, J. *et al.* Computing the relative binding affinity of ligands based on a pairwise

binding comparison network. *Nat. Comput. Sci.* **3**, 860–872 (2023).

7.   Koh, H. Y., Nguyen, A. T. N., Pan, S., May, L. T. & Webb, G. I. Physicochemical

graph neural network for learning protein–ligand interaction fingerprints from sequence data.

Nat. Mach. Intell. **6**, 673–687 (2024).

8.   Wang, L. *et al.* Accurate and Reliable Prediction of Relative Ligand Binding Potency

in Prospective Drug Discovery by Way of a Modern Free-Energy Calculation Protocol and Force

Field. *J. Am. Chem. Soc.* **137**, 2695–2703 (2015).

9.   Cournia, Z., Allen, B. & Sherman, W. Relative Binding Free Energy Calculations in

Drug Discovery: Recent Advances and Practical Considerations. *J. Chem. Inf. Model.* **57**, 2911–

2937 (2017).

10.   Cao, D. *et al.* SurfDock is a surface-informed diffusion generative model for reliable

and accurate protein–ligand complex prediction. *Nat. Methods* **22**, 310–322 (2025).

11.   Moon, S., Hwang, S.-Y., Lim, J. & Kim, W. Y. PIGNet2: a versatile deep learning-

based protein–ligand interaction prediction model for binding affinity scoring and virtual

screening. *Digit. Discov.* **3**, 287–299 (2024).

12.   Landrum, G. A. & Riniker, S. Combining IC50 or Ki Values from Different Sources

Is a Source of Significant Noise. *J. Chem. Inf. Model.* **64**, 1560–1567 (2024).

13.   Sorscher, B., Geirhos, R., Shekhar, S., Ganguli, S. & Morcos, A. Beyond neural

scaling laws: beating power law scaling via data pruning. *Adv. Neural Inf. Process. Syst.* **35**,

19523–19536 (2022).

14.   Bahri, Y., Dyer, E., Kaplan, J., Lee, J. & Sharma, U. Explaining neural scaling laws.

Proc. Natl. Acad. Sci. **121**, e2311878121 (2024).

15.   Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with

AlphaFold 3. *Nature* **630**, 493–500 (2024).

16.   OpenAI   *et   al.*   GPT-4   Technical   Report.   Preprint   at

https://doi.org/10.48550/arXiv.2303.08774 (2024).

17.   Muegge, I. & Martin, Y. C. A General and Fast Scoring Function for Protein−Ligand

Interactions: A Simplified Potential Approach. *J. Med. Chem.* **42**, 791–804 (1999).

18.   Shen, M. & Sali, A. Statistical potential for assessment and prediction of protein

structures. *Protein Sci.* **15**, 2507–2524 (2006).

19.   Ferrada, E. & Melo, F. Effective knowledge-based potentials. *Protein Sci.* **18**, 1469–

1485 (2009).

20.   Nehil-Puleo, K., Quach, C. D., Craven, N. C., McCabe, C. & Cummings, P. T. E(n)

Equivariant Graph Neural Network for Learning Interactional Properties of Molecules. *J. Phys.*

Chem. B **128**, 1108–1117 (2024).

21.   Liu, T., Lin, Y., Wen, X., Jorissen, R. N. & Gilson, M. K. BindingDB: a web-

accessible database of experimentally determined protein–ligand binding affinities. *Nucleic*

Acids Res. **35**, D198–D201 (2007).

22.   Simeon, G. & De Fabritiis, G. TensorNet: Cartesian Tensor Representations for

Efficient Learning of Molecular Potentials. *Adv. Neural Inf. Process. Syst.* **36**, 37334–37353

(2023).

23.   Cohen, M. L. Epidemiology of Drug Resistance: Implications for a Post—

Antimicrobial Era. *Science* **257**, 1050–1055 (1992).

24.   Alfalah, M., Keiser, M., Leeb, T., Zimmer, K. & Naim, H. Y. Compound

Heterozygous Mutations Affect Protein Folding and Function in Patients With Congenital

Sucrase-Isomaltase Deficiency. *Gastroenterology* **136**, 883–892 (2009).

25.   Teng, S., Madej, T., Panchenko, A. & Alexov, E. Modeling Effects of Human Single

Nucleotide Polymorphisms on Protein-Protein Interactions. *Biophys. J.* **96**, 2178–2188 (2009).

26.   Hashimoto, K. & Panchenko, A. R. Mechanisms of protein oligomerization, the

critical role of insertions and deletions in maintaining different oligomeric states. *Proc. Natl.*

Acad. Sci. **107**, 20352–20357 (2010).

27.   Zhan, S. *et al.* Discovery of Imidazo[1,2-a]pyrazine Derivatives as Potent ENPP1

Inhibitors. *J. Med. Chem.* **67**, 18317–18333 (2024).

28.   Chen, Y. *et al.* Aldehyde dehydrogenase 1B1 (ALDH1B1) is a potential biomarker

for human colon cancer. *Biochem. Biophys. Res. Commun.* **405**, 173–179 (2011).

29.   Passaro, S. *et al.* Boltz-2: Towards Accurate and Efficient Binding Affinity

Prediction. Preprint at https://doi.org/10.1101/2025.06.14.659707 (2025).

30.   Friesner, R. A. *et al.* Glide: A New Approach for Rapid, Accurate Docking and

Scoring. 1. Method and Assessment of Docking Accuracy. *J. Med. Chem.* **47**, 1739–1749 (2004).

31.   Kollman, P. A. *et al.* Calculating Structures and Free Energies of Complex

Molecules: Combining Molecular Mechanics and Continuum Models. *Acc. Chem. Res.* **33**, 889–

897 (2000).

32.   Lam, H. Y. I., Guan, J. S., Ong, X. E., Pincket, R. & Mu, Y. Protein language models

are performant in structure-free virtual screening. *Brief. Bioinform.* **25**, bbae480 (2024).

33.   Rose, T., Monti, N., Anand, N. & Shen, T. PLAPT: Protein-Ligand Binding Affinity

Prediction   Using   Pretrained   Transformers.   Preprint   at

https://www.biorxiv.org/content/10.1101/2024.02.08.575577v3 (2024).

34.   Shen, C. *et al.* Boosting Protein–Ligand Binding Pose Prediction and Virtual

Screening Based on Residue–Atom Distance Likelihood Potential and Graph Transformer. *J.*

Med. Chem. **65**, 10691–10706 (2022).

35.   Shen, C. *et al.* A generalized protein–ligand scoring framework with balanced

scoring, docking, ranking and screening powers. *Chem. Sci.* **14**, 8129–8146 (2023).

36.   Wang, Z. *et al.* OnionNet-2: A Convolutional Neural Network Model for Predicting

Protein-Ligand Binding Affinity Based on Residue-Atom Contacting Shells. *Front. Chem.* **9**,

753002 (2021).

37.   Mastropietro, A., Pasculli, G. & Bajorath, J. Learning characteristics of graph neural

networks predicting protein–ligand affinities. *Nat. Mach. Intell.* **5**, 1427–1436 (2023).

38.   Marshall, G. R. & Cramer, R. D. Three-dimensional structure-activity relationships.

Trends Pharmacol. Sci. **9**, 285–289 (1988).

39.   Peltason, L. & Bajorath, J. SAR Index: Quantifying the Nature of Structure−Activity

Relationships. *J. Med. Chem.* **50**, 5571–5578 (2007).

40.   Guha, R. On exploring structure-activity relationships. *Methods Mol. Biol.* **993**, 81–

94 (2013).

41.   Kenny, P. W. Hydrogen-Bond Donors in Drug Design. *J. Med. Chem.* **65**, 14261–

14275 (2022).

42.   Paulini, R., Müller, K. & Diederich, F. Orthogonal Multipolar Interactions in

Structural Chemistry and Biology. *Angew. Chem. Int. Ed.* **44**, 1788–1805 (2005).

43.   Bissantz, C., Kuhn, B. & Stahl, M. A Medicinal Chemist's Guide to Molecular

Interactions. *J. Med. Chem.* **53**, 5061–5084 (2010).

44.   Schneider, H.-J. Hydrogen bonds with fluorine. Studies in solution, in gas phase and

by computations, conflicting conclusions from crystallographic analyses. *Chem. Sci.* **3**, 1381–

1394 (2012).

45.   Pollock, J. *et al.* Rational Design of Orthogonal Multipolar Interactions with Fluorine

in Protein–Ligand Complexes. *J. Med. Chem.* **58**, 7465–7474 (2015).

46.   Zhou, P., Zou, J., Tian, F. & Shang, Z. Fluorine Bonding — How Does It Work In

Protein−Ligand Interactions? *J. Chem. Inf. Model.* **49**, 2344–2355 (2009).

47.   Herbst, R. S., Fukuoka, M. & Baselga, J. Gefitinib — a novel targeted approach to

treating cancer. *Nat. Rev. Cancer* **4**, 956–965 (2004).

48.   Yu, H. A. *et al.* Local therapy with continued EGFR tyrosine kinase inhibitor therapy

as a treatment strategy in EGFR-mutant advanced lung cancers that have developed acquired

resistance to EGFR tyrosine kinase inhibitors. *J. Thorac. Oncol.* **8**, 346–351 (2013).

49.   Yu, H. A. *et al.* Analysis of tumor specimens at the time of acquired resistance to

EGFR-TKI therapy in 155 patients with EGFR-mutant lung cancers. *Clin. Cancer Res.* **19**, 2240–

2247 (2013).

50.   Schmid, S., Li, J. J. N. & Leighl, N. B. Mechanisms of osimertinib resistance and

emerging treatment options. *Lung Cancer* **147**, 123–129 (2020).

51.   Hao, G.-F., Yang, G.-F. & Zhan, C.-G. Structure-based Methods for Predicting Target

Mutation-induced Drug Resistance and Rational Drug Design to Overcome the Problem. *Drug*

Discov. Today **17**, 1121–1126 (2012).

52.   Cheng, J. *et al.* Accurate proteome-wide missense variant effect prediction with

AlphaMissense. *Science* **381**, eadg7492.

53.   Watermeyer, J. M., Kröger, W. L., O'Neill, H. G., Sewell, B. T. & Sturrock, E. D.

Probing the Basis of Domain-Dependent Inhibition Using Novel Ketone Inhibitors of

Angiotensin-Converting Enzyme. *Biochemistry* **47**, 5942–5950 (2008).

54.   Kim, E. E. *et al.* Crystal structure of HIV-1 protease in complex with VX-478, a

potent and orally bioavailable inhibitor of the enzyme. *J. Am. Chem. Soc.* **117**, 1181–1182 (1995).

55.   Steuber, H., Heine, A., Podjarny, A. & Klebe, G. Merging the Binding Sites of Aldose

and Aldehyde Reductase for Detection of Inhibitor Selectivity-determining Features. *J. Mol. Biol.*

379, 991–1016 (2008).

56.   Vanichtanankul, J. *et al.* Combined Spatial Limitation around Residues 16 and 108

of Plasmodium falciparum Dihydrofolate Reductase Explains Resistance to Cycloguanil.

Antimicrob. Agents Chemother. **56**, 3928–3935 (2012).

57.   Finer-Moore, J. S. *et al.* Partitioning Roles of Side Chains in Affinity, Orientation,

and Catalysis with Structures for Mutant Complexes: Asparagine-229 in Thymidylate Synthase.

Biochemistry **35**, 5125–5136 (1996).

58.   Caravella, J. A. *et al.* Structure-Based Design and Identification of FT-2102

(Olutasidenib), a Potent Mutant-Selective IDH1 Inhibitor. *J. Med. Chem.* **63**, 1612–1623 (2020).

59.   Hallin, J. *et al.* Anti-tumor efficacy of a potent and selective non-covalent

KRASG12D inhibitor. *Nat. Med.* **28**, 2171–2182 (2022).

60.   Wang, E. *et al.* Mechanisms of Resistance to Noncovalent Bruton's Tyrosine Kinase

Inhibitors. *New Engl. J. Med.* **386**, 735–743 (2022).

61.   Anderson, P. W. More Is Different. *Science* **177**, 393–396 (1972).

62.   Wei, J. *et al.* Emergent Abilities of Large Language Models. Preprint at

https://doi.org/10.48550/arXiv.2206.07682 (2022).

63.   Chen, T., Li, M. & Liu, J. π–π Stacking Interaction: A Nondestructive and Facile

Means in Material Engineering for Bioapplications. *Cryst. Growth Des.* **18**, 2765–2783 (2018).

64.   Xu, X., Jia, K., Qi, Q., Tian, G. & Xiang, D. Regulation of π–π interactions between

single aromatic molecules by bias voltage. *Phys. Chem. Chem. Phys.* **26**, 14607–14612 (2024).

65.   Heredia, V. V., Cooper, W. C., Kruger, R. G., Jin, Y. & Penning, T. M. Alanine

Scanning Mutagenesis of the Testosterone Binding Site of Rat 3α-Hydroxysteroid

Dehydrogenase Demonstrates Contact Residues Influence the Rate-Determining Step.

Biochemistry **43**, 5832–5841 (2004).

66.   Zou, H. *et al.* Molecular Insight into the Interaction between IFABP and PA by Using

MM−PBSA and Alanine Scanning Methods. *J. Phys. Chem. B* **111**, 9104–9113 (2007).

67.   Krishna, R. *et al.* Generalized biomolecular modeling and design with RoseTTAFold

All-Atom. *Science* **384**, eadl2528 (2024).

68.   Weininger, D. SMILES, a chemical language and information system. 1. Introduction

to methodology and encoding rules. *J. Chem. Inf. Comput. Sci.* **28**, 31–36 (1988).

69.   Landrum, G. RDKit: Open-source cheminformatics from machine learning to

chemical registration. https://www.rdkit.org (2019).

70.   Schindler, C. E. M. *et al.* Large-Scale Assessment of Binding Free Energy

Calculations in Active Drug Discovery Projects. *J. Chem. Inf. Model.* **60**, 5457–5474 (2020).

71.   Gaulton, A. *et al.* ChEMBL: a large-scale bioactivity database for drug discovery.

Nucleic Acids Research **40**, D1100–D1107 (2012).

72.   Yang, Z. *et al.* A mutation-induced drug resistance database (MdrDB). *Commun.*

Chem. **6**, 123 (2023).

73.   Lin, Z. *et al.* Evolutionary-scale prediction of atomic-level protein structure with a

language model. *Science* **379**, 1123–1130 (2023).

74.   Elnaggar, A. *et al.* ProtTrans: Toward Understanding the Language of Life Through

Self-Supervised Learning. *IEEE Trans. Pattern Anal. Mach. Intell.* **44**, 7112–7127 (2022).

1193