Alpha cell MHC-I expression recruits cytotoxic CD8[+] T cells during human aging and type 2 diabetes.

Michael Schleh[1], Jean-Philippe Cartailler[2], Shristi Shrestha[2], Amanda Cambraia[1], Aliyah Habashy[1], Cara Ellis[3], Melanie Cutler[1], Gabriel Ferguson[1], Ricki Colman[4,5], Alvin Powers[6], Rafael Arrojo e Drigo[1,6,7,8*]

[1] Department of Molecular Physiology and Biophysics, Vanderbilt University School of Medicine, Nashville, TN
[2] Creative Data Solutions, Vanderbilt University School of Medicine, Nashville, TN
[3] Alberta Diabetes Institute and Department of Pharmacology, University of Alberta, Edmonton, Canada
[4] University of Wisconsin, Cell and Regenerative Biology, School of Medicine and Public Health, Madison, Wisconsin
[5] Wisconsin National Primate Research Center (WNPRC), Madison, Wisconsin
[6] Vanderbilt Diabetes Research and Training Center, Vanderbilt University Medical Center, Nashville, TN
[7] Center for Cell Systems Biology, Vanderbilt University School of Medicine, Nashville, TN
[8] Mass Spectrometry Research Center, Vanderbilt University School of Medicine, Nashville, TN
*Corresponding author

Keywords: aging, alpha cells, antigen presentation, calorie restriction, T cells

Address of Correspondence:
Rafael Arrojo e Drigo
718 Light Hall
2215 Garland Ave
Nashville, Tennessee 37232
r.drigo@vanderbilt.edu

Running Title: Alpha cell inflammation and calorie restriction in aging

Abstract

Aging is a major risk factor for type 2 diabetes (T2D), driven in part by loss of beta cell identity and function due to stress and islet inflammation. Calorie restriction (CR) improves metabolic health by promoting beta cell longevity in young animals. It remains unknown whether CR reverses cellular aging and dampens islet inflammation in aged animals. We show that aging alpha cells have an active inflammatory phenotype with increased IFNα/γ signaling that activates major histocompatibility complex (MHC-I) expression to promote CD8$^+$ T-cell accumulation in mouse and human islets. During human T2D, aging islet-resident CD8$^+$ T cells transition from memory states to activated effector phenotypes. Moreover, CR dampens alpha cell MHC-I expression to suppress CD8$^+$ cytotoxicity, which is sufficient to prevent islet inflammation and fibrosis in mice and non-human primates. Together, these findings highlight alpha cell MHC-I within the islet endocrine–immune axis in aging that may underlie fibrosis and disease development.

Introduction

Aging is a major risk factor for type 2 diabetes (T2D), affecting 29.2% of adults 65 and older in the United States[1]. Age-related deterioration in glycemic control has been traditionally attributed to declining peripheral insulin sensitivity, reduced beta cell insulin secretion, and increased hepatic insulin clearance[2-4]. However, older individuals also exhibit elevated circulating glucagon levels at basal and after a mixed meal stimulation[2], thus supporting the idea that dysregulation of alpha cell function may contribute to impaired glucose homeostasis during aging. Notably, hyperglucagonemia is a hallmark of adult T2D due to blunted glucose-dependent suppression of alpha cell glucose secretion[5] and elevated amino acid induced-glucagon release[6,7]. Notably, this phenotype is prevalent in older adults with impaired glucose tolerance and may be absent in T2D children[8], which suggests that aging alpha cells could be more likely to failure during T2D. Alpha and beta cells are long-lived endocrine cells that secrete insulin and glucagon, and constitute the majority of the islet[9], and are essential regulators of glucose homeostasis that must maintain function and support throughout the lifetime[10-12]. Aging beta cells have compromised structure-function due to loss of cellular identity markers, increased DNA damage and senescence, endoplasmic reticulum (ER) stress, compromised calcium homeostasis, impaired autophagy, increased transcriptional noise, amyloid deposition, and islet fibrosis[2,13-16]. It remains unknown if and how aging impacts alpha cell structure-function and glucagon-regulatory mechanisms.

Calorie restriction (CR) is a well-established dietary intervention that extends life- and health-spans across species, by delaying the onset of age-related cell dysfunction and disease[17-20]. Moderate reductions in energy intake (10–40%) improve metabolic homeostasis in both young and aged animal, whereas more severe restriction can lead to muscle wasting and immunosuppression[21]. At the molecular level, CR modulates major nutrient-sensing pathways including suppression of insulin-IGF1 signaling[22], inhibition of mammalian target of rapamycin complex (mTORC1) activity and activation of AMPK[23], reduction of oxidative stress and inflammatory tone[24], and enhanced autophagy[25,26]. Pharmacological strategies targeting these pathways, such as rapamycin (mTORC1 inhibition)[27], metformin (AMPK activation)[28], or resveratrol[29] have shown promises, but are limited by off-target effects, including immunosuppression and beta cell failure[30,31], which restricts their translational potential. CR and CR-mimicking interventions (i.e., intermittent fasting and time restricted feeding) in young adult mice promotes beta cell health, function, and longevity by enhancing peripheral insulin sensitivity that in turn reduces the metabolic demand for beta cell insulin secretion[32-35]. These pro-longevity effects are associated with increased expression of beta cell identity and circadian gene expression regulators, improved autophagy and protein homeostasis, mitochondrial homeostasis and ATP production, and reduced beta cell DNA damage and senescence[32-35]. Whether CR can rescue pancreatic inflammation and beta cell dysfunction in aging animals, and whether CR has any significant impact on aging alpha cells remains unclear.

To address these gaps, we integrated large-scale meta-analyses of publicly available human islet proteomic, transcriptomic, and spatial proteomic datasets to define molecular signatures of aging in non-diabetic (ND) donors, and to determine whether these cellular signatures translate to T2D. We then used CR in aged mice to translate whether dietary restriction can reverse age-associated cellular and molecular alterations in old islets. As expected, the aging human islet proteome exhibited increased ER stress, senescence, and beta cell immaturity signatures. Surprisingly, we also observed heightened inflammatory markers linked to innate immunity, coupled with adaptive immunity in antigen presentation markers. Single-cell transcriptomics revealed these proteomic signatures were largely explained by upregulation of hallmark inflammation genes in aging alpha cells, including activation of MHC-I genes and interferon (IFN) α/γ pathways. Using transcription factor inference algorithms, we found activation of alpha cell

inflammatory signatures is attributed to activation of pro-inflammatory transcription factors (TFs)
KLF6, IRF1, and NFKB1. Multiplexed spatial mass spectrometry confirmed higher MHC-I (HLA-
ABC) expression in aging alpha cells, but not beta cells, and revealed an age-related increase
in central memory (CD27$^+$CD45RA$^-$) CD8$^+$ T cells (CD8$_{CM}$) that shifted toward a terminally differ-
entiated effector memory phenotype (CD27$^-$CD45RA$^+$; CD8$_{TEM}$) in T2D islets. Exposure of aged
lean mice to 20% CR for two months significantly reduced alpha cell pro-inflammatory gene ex-
pression and disrupted ligand-receptor inference of alpha-to-T-cell communication, suggesting
that CR broadly attenuates immune activation through the suppression of alpha cell inflamma-
tory signaling. Together, our findings indicate that aging alpha cells exhibit pro-inflammatory sig-
natures that underlie age-related pancreas inflammation and fibrosis via infiltration of cytotoxic
CD8$^+$ T cells. These aspects are conserved in mice and humans and is suppressed by CR in
mouse and non-human primates (NHPs). Therefore, CR may offer the dual benefit of improving
overall metabolic health in lean aging organisms while reversing islet inflammation signatures
associated with impaired beta cell function and compromised glucose homeostasis.

Results

Aging ND islets heighten inflammatory and T-cell receptor activation pathways with
mTORC1 and heightened unfolded protein response. Islet inflammation and fibrosis are rec-
ognized hallmarks of human pancreas aging and diabetes, contributing to beta cell functional
decline[36-38]. To define human proteomic aging signatures, we leveraged a comprehensive hu-
man islet proteomics dataset generated by the HumanIslets consortium (**Figure 1A, Supple-**
mentary Table 1)[39,40]. Comparisons of young (28±4 years, n=26) and old (71±3 years, n=19)
ND adults identified 86 proteins significantly increased by aging and 17 proteins decreased. Re-
actome pathway enrichment revealed increased immune regulatory and innate immune path-
ways, amino acid metabolism and mTORC1-related metabolic factors, and ER stress in old is-
lets (**Figure 1B**). These age-associated changes included elevated beta cell immaturity markers
(RBP4[41]; **Figure 1C**), senescence (CDKN1A/p21[42], **Figure 1D**), ER stress signaling (ATF6,
CHOP, and PERK; **Figure 1E**), and increased mTORC1 activation proteins (LAMP2, LAMTOR,
RAGA, RAGC; **Figure 1F**). Moreover, we observed a robust immune activation signature char-
acterized by increased expression of pro-inflammatory and antigen presentation proteins, in-
cluding the MHC-1 antigen, HLA-A, T-cell receptor (TCR) signaling and memory components
such as CD28 and CD44, and additional antigen processing and immune-regulatory signaling
(TGF-β1, TOM1, and SLC15A4; **Figure 1G**). Together, these findings reinforce previous reports
to support that aging human islets have an active cellular stress response accompanied by en-
hanced inflammatory signaling to upregulate immune cell activation and recruitment, although
the cell-type–specific distinctions to these processes remain to be defined.

Single cell transcriptomics reveal activation of inflammatory signaling in aging human
alpha cells. Over the past decade, several groups have applied diverse single cell tran-
scriptomic approaches to investigate how islet cell gene expression and chromatin accessibility
are regulated during development, aging, and diabetes[41,43-52]. We recently demonstrated that
aging ND beta cells have an active ER stress response[43], which explains the higher level of ER
stress markers in bulk islet proteome (**Figure 1**). In mice, a recent bulk RNA-seq study of sorted
alpha cells, beta cells, and delta cells associated aging with the activation of immune responses
and MHC-I complex gene expression[53]. To identify the specific islet cell types contributing to the
increased expression of pro-inflammatory markers identified in the human islet proteomics (in-
cluding MHC-I), we performed meta-analysis of single cell mRNA sequencing (scRNA-seq) tran-
scriptomics deposited in the HPAP PANC-DB database[54-57]. We initially curated scRNA-seq data
from a total of n=132 donor samples and applied a comprehensive bioinformatics approach to
filter samples and perform robust quality control and covariate correction for sex, BMI, ethnicity,
sequencing chemistry, and sequencing batch (see methods for details) (**Figure 2A**,

Supplementary Figure 1A-B, **Supplementary Table 2**). This resulted in a final dataset con-
taining scRNA-seq profiles from n=32 donors (131,610 annotated islet cells, including 56,851
alpha cells), stratified into young adults (20-40 years of age) and old adults (≥50 years of age)
(**Figure 2B**).

First, we investigated which ND human islet cell types express immune signaling genes
and found that alpha cells exhibit the highest expression of HLA-ABC antigens, which encode
the MHC-1 complex, among all major endocrine cell types – which is followed by endothelial
and actual immune cell types (**Figure 2C**). Pseudobulk differential gene expression and princi-
pal components analysis (PCA) resolved groups into distinct clusters (**Supplementary Figure
1B**). Gene Set Enrichment Analysis (GSEA) identified activation of both IFNα and IFNγ signaling
(*STAT1*, *SOCS1*, *SOCS3*, *IRF7*, *IRF9*, *IL7*, and *IL15*; **Figure 2D-E**), and MHC-I-mediated anti-
gen presentation pathways (*HLA-E*, *HLA-F*, and *TAP1*: **Figure 2F**) in old alpha cells. In addition,
aging alpha cells had increased expression of genes associated with glucose, fatty acid, and
amino acid catabolism (**Figure 2G-I**). These signatures correlated with activation of a "beta cell-
like" senescence program marked (CDKN1A/p21, CDKN1B/p27, CDKN2A/p16, and IGF1R),
and a lack of an active ER stress signature (**Supplementary Figure 1C**). To determine if these
transcriptional changes correlate with alterations in alpha cell glucagon secretion, we analyzed
HPAP human islet perfusion data from young and old donors from two different human islet
analysis centers (UPenn and Vanderbilt), which used complementary stimulation protocols to
test for the effect of several insulin and glucagon secretagogues. This revealed that alpha cells
from older donors have increased glucagon secretion in response to stimulation with a mixed
amino acid (AA) cocktail and to de-polarization with KCL (**Figure 2J**). No changes were ob-
served when alpha cells were stimulated with low glucose, high glucose, and epinephrine (**Sup-
plementary Figure 1D**). Together, these results indicate that aging human alpha cells acquire
senescence-associated features, display heightened AA catabolism and AA-dependent gluca-
gon secretion, and an active IFN signaling and MHC-I antigen-presentation programs.

**Re-organization of gene regulatory networks activates inflammatory gene expression in
aging alpha cells.** We previously applied the transcription factor (TF) inference algorithm
pySCENIC[58] to show that aging beta cell stress arises from reconfigured gene regulatory net-
works (GRNs) and activation of stress TFs[43]. Here, we used pySCENIC and graph network
analysis to characterize GRNs in aging alpha cells and identify TFs driving inflammation. Re-
construction of GRNs in young and old alpha cells revealed canonical alpha cell TFs (ARX,
PAX6, and MAFB) (**Figure 3A-B**). Young alpha cells presented few TFs forming small regulons,
composed of a limited number of TFs, including regulators previously implicated in beta cell
GRNs (MAFA, EGR1, and BCLAF1; **Figure 3A**). In contrast, aging alpha cells exhibited ex-
panded GRNs connected to pro-inflammatory TFs such as KLF6, NFKB1, IRF1, and MXD1,
consistent with enrichment of their downstream targets in older alpha cells (**Figure 3B-D**).

We previously found that TFs maintaining beta cell identity in young adults can be "redi-
rected" to activate cell stress and beta cell dedifferentiation programs during aging[43]. To investi-
gate if a similar mechanism was in place in aging alpha cells, we analyzed the identity of TF
regulons linked to NFKB1, KLF6, IRF1, and MAFB in young versus old alpha cells. This ap-
proach revealed substantial shifts in TF regulon identity, including the activation of a KLF6 regu-
lon with 226 target genes not expressed in young alpha cells, and expansion of NFKB1, IRF1,
and MAFB regulons (**Figure 3C-D**). Reactome Pathway analysis[59] mapped these regulons to
inflammatory pathways, including immune activation and cytokine signaling (i.e., IFN*, IL20*, IL4,
and IL13), caspase-mediated cell death, MAPK and TGF-β signaling, and response to mechani-
cal stress (**Figure 3E-F, Supplementary Figure 2A**). Notably, HNF1A and MAFB regulons in
aging alpha cells were associated with ERBB4 signaling, which mediates epidermal growth fac-
tor (EGF) receptor signaling cascades[60].

To identify potential activators of these inflammatory pathways, we applied CellChat[61] to
infer cell-cell communication from scRNA-seq data across young and old islet cell types (**Figure**
2G-I). This approach identified 74 signaling pathways that are significantly up regulated in aging
human alpha cells (**Supplementary Figure 2B, Supplementary Table 3**). These enriched path-
ways were further categorized into major molecular classes such as extracellular matrix (ECM)
adhesion (Laminin, collagen, Vcam/Icam) and remodeling (Mmp), growth factors (Fgf, Egf,
Pdgf), GPCR signaling (Adgrl, Prostaglandins), immune checkpoints (Lair1, MHC-I, Cd46, IL6),
neurotransmitters and other hormones (Gaba-b, Insulin, ApoE), Semaphorins (Sema3), mem-
brane receptor signaling (Wnt, Tgfb, App), tight junctions (Ocln, Clnd), and "other" (Ptprm, an-
nexin). Approximately 30% of enriched pathways involved ECM biology, reflecting the fibrotic
aging islet microenvironment[37], (**Figure 3G**, **Supplementary Figure 2B-C**). Growth factor path-
ways, including pro-fibrotic factor TGFB1 were the second most represented (**Figure 3H, Sup-**
plementary Figure 2D-E and 3A). Beta cells were primary TGFB1 secretory source in both
young and old islets, followed by quiescent stellate and ductal cells. In young islets, TGFB1 sig-
naling from both alpha and beta cells targeted macrophages, T cells, endothelial cells, and acti-
vated stellate cells; in older islets, signaling expanded to acinar and quiescent microglia-like
cells (**Figure 3H**). Strikingly, alpha and beta cells established WNT signaling networks and acti-
vated stellate and acinar cells communicated via LIF – two molecules that are involved in
TGFB1-mediated fibrosis (**Supplementary Figure 3A**)[62,63]. FGF signaling from ductal, activated
stellate, and microglia cells primarily targeted aging alpha cells, whereas EGF and GDF signal-
ing were mostly restricted to the extra-islet compartment (**Figure 3H, Supplementary Figure**
2D-3 and 3A). Older alpha cells were enriched in immune signaling, including CD46 (amplifying
CD8[+] T-cell signaling[64] and CD4/8[+] IFNγ production[65]), prostaglandin and IL6 signaling (which
regulates alpha cell secretory function[66]), and protein tyrosine phosphatase receptor (PRPRM),
a modulatory of IL6 signaling during stress responses[67] (**Figure 3I, Supplementary Figure 2F**
and 3A). Together, these studies highlight aging human alpha cells acquire a pro-inflammatory
identity through extensive rewiring of their GRNs, marked by activation of stress- and cytokine-
responsive TFs (including IRF1, NFKB1, and KLF6) and reduced MAFB activity. These regulons
appear to be amplified by TGFB1-, WNT-, and T-cell- and cytokine-mediated signaling from
neighboring islet and stromal cells, processes that may contribute to the fibrotic remodeling of
the aging islet microenvironment.

Spatial analysis of aging human pancreases identifies alpha-T-cell interactions *in situ*.
Aging ND human pancreases exhibit multiple features of chronic low-grade inflammation, in-
cluding increased fibrosis, higher incidence of acinar-to-ductal metaplasia (ADM), angiopathy,
and pancreatic adiposity[38]. To validate our bulk proteomic and scRNA-seq findings *in situ*, we
analyzed high-dimensional imaging mass cytometry (IMC) and CO-Detection by indexing (CO-
DEX) datasets from HPAP[56,57]. Human ND pancreas samples were stratified into young adult
(22-40 years) and old adults (47-58 years) groups (**Figure 4A and 4D; Supplementary Figure**
4B and 5B; Supplementary Tables 4 and 5). Individual cells from each donor image were seg-
mented using a custom QuPath-based and spatial analysis pipeline (see methods), enabling un-
supervised clustering into major endocrine and immune cell populations based on marker ex-
pression ((**Supplementary Figure 4A, C-G and 5A, C-G**), including alpha (GCG[+]), beta (C-
peptide[+]), delta (SST[+]), pancreatic polypeptide (PP[+]), endothelial cells (CD31[+]), Epithelial/Acinar
cells (CK19[+]/CA2[+]), M1-like macrophages (CD45[+]CD68[+]CD163[-]) , M2-like macrophages
(CD45[+]CD68[+]CD163[+]), and T cells (CD45[+]CD3[+]CD4[+] and CD45[+]CD3[+]CD8[+]) (**Figure 4B and**
4E, Supplementary Tables 6 and 7). This framework enabled spatial mapping of all major pan-
creatic and immune cell types in either IMC (**Figure 4C**) and CODEX (**Figure 4F**) platforms, as
well as quantification of immune and endocrine cell phenotyping (**Figure 4G-H**; **Supplementary**
Figure 6A and 6C). Notably, aging pancreases showed significant expansion of CD45[+]CD3[+] T

cells, predominantly driven by CD8$^+$ T-cell infiltration (**Figure 4G-H, Supplementary Figure**
6C).

To investigate endocrine-immune interactions underlying CD8$^+$ T-cell expansion, we performed
nearest-neighbor spatial analyses to quantify the proximity of CD8$^+$ T cells to their nearest alpha
and beta cells. Aged pancreases displayed a higher proportion of alpha cells localized within
close proximity (<50µm) to CD8$^+$ T cells (**Figure 4I**), a relationship not observed for beta cells
(**Figure 4J**). No age-associated differences were detected between CD4$^+$ T cells to alpha cells
or in total macrophage density (**Figure 4G-H**). Notably, the increased spatial proximity between
alpha cells and CD8$^+$ T cells was accompanied with increased expression of HLA-ABC abun-
dance (encoding MHC-I complex) and attenuated abundance of HLA-DR abundance (encoding
MHC-II) in alpha cells, whereas beta cells showed no such change (**Figure 4K-M**). Importantly,
we detected expected age-associated decreases in beta cell identity markers PDX-1, NKX6-1,
and PAX-6[43] (**Supplementary Figure 6B**), confirming the classical aging beta cell pheno-
types[38,43]. Together, these findings extend our isolated islet analyses by demonstrating that hu-
man islet and pancreatic aging is marked by increased pro-inflammatory signaling from alpha
cells, which correlates with elevated CD8$^+$ T-cell density *in situ*. Given that MHC molecules pre-
sent antigenic peptides to T cells via TCR engagement, these results suggest that alpha cell-
intrinsic inflammatory pathways contribute to the immune activation and pancreatic inflammation
that emerge with aging.

T2D expands pancreatic CD8$^+$ T cells that shift towards terminal effector-memory pheno-
types. To determine whether whole-pancreas immune features associated with immune aging
overlap with T2D, we analyzed additional CODEX data from T2D donors profiled through HPAP
(**Supplementary Table 8**). These cohorts were matched for age and BMI; however, T2D donors
displayed expected impairments in glycemia, evidenced by higher HbA1c and reduced C-pep-
tide (**Figure 5A**). Integration of 942,724 single-cell profiles identified targeted endocrine (99,744
cells) and immune (195,140 cells) compartments (**Figure 5B**). Endocrine cell clustering identi-
fied two distinct beta cell populations: a CPEP$^+$CHGA$^+$ population and a
CPEP$^+$CHGA$^+$COL4A1$^+$COL6$^+$ "fibrotic" beta cell state enriched in T2D (**Figure 5C**). These cells
exhibited higher interaction with fibrotic ECM and thus were detected as being COL4A1$^+$COL6$^+$
by our cell segmentation algorithms. T2D beta cells were marked by lower CPEP, CHGA, NKX6-
1, and Ki67 expression (**Figure 5D**), while alpha cells had attenuated SOX9 and Ki67 (**Figure**
5E), consistent with diminished endocrine cell identity and proliferative capacity. Within immune
compartments, we observed a trend towards increased CD8$^+$ T-cell representation and de-
creased CD68$^+$CD163$^+$ M2-like macrophages (**Figure 5F**), accompanied by a significant ~2-fold
increase in pancreas CD8$^+$ T-cell density (**Figure 5G**). Together, these findings demonstrate that
T2D amplifies CD8$^+$ T-cell density, mirroring patterns observed in aging and establishes height-
ened cytotoxic pressure within the pancreas.

To resolve whether the increased adaptive immune phenotype observed in the aging
and T2D pancreas was localized within islets, we analyzed HPAP cytometry by time-of-flight
(CyTOF) data from isolated islets of young (26 ± 8 years, n=13), old (54 ± 5 years, n=14), and
T2D (50 ± 8 years, n=13) human donors (**Figure 5H, Supplementary Table 9**). Both aging and
T2D presented greater CD45$^+$ live immune cell abundance compared with young adults, corre-
sponding with to higher frequencies of islet CD3$^+$CD8$^+$ T cells within the CD45$^+$ immune com-
partment (**Figure 5I-J**). CD8$^+$ T cells differentiate along a spectrum from naïve (CD8$_{naïve}$) to
CD8$_{CM}$ and CD8$_{TEM}$ states in response to TCR activation by antigen and cytokine stimulation. In
T2D islets, we observed a reduction in CD45RA$^+$CD27$^+$ CD8$_{naïve}$ T cells compared with young
adults, whereas aging increased CD45RA$^-$CD27$^+$ CD8$_{CM}$ T cells (**Figure 5K-L**). T2D islets fur-
ther shifted CD8$^+$ T-cell populations towards a higher CD45RA$^+$CD27$^-$ CD8$_{TEM}$ phenotype, re-
sulting in elevated effector-to-central memory ratio that was not observed in aging ND samples

(**Figure 5L-M**). Together, these data show that T2D intensifies pancreas and islet CD8[+] T-cell
infiltration and drives their progression toward terminal effector-memory states, establishing a
cytotoxic immune milieu that is distinct from, but overlaps with immune aging.

Calorie restriction improves glucose tolerance and dampens alpha cell inflammation in
mice. Dieting through calorie restriction (CR) and weight loss interventions is commonly pre-
scribed as an initial strategy to normalize glycemia in individuals with prediabetes and T2D. We
and others have shown that CR enhances glucose homeostasis and delays the onset of inflam-
mation and aging hallmarks associated with metabolic dysfunction, including in beta cells[68-72].
Therefore, we tested whether CR could reverse aging signatures in alpha and beta cells *in vivo*.
To achieve this, 72-week-old C57BL/6J male mice were subjected to a 2-month, 20% CR feed-
ing regimen, while age-matched controls remained on ad-libitum (AL) diet (**Figure 6A**). As ex-
pected, CR mice consumed 80% of total kcal intake compared to AL-fed controls (**Supplemen-**
tary Figure 7A-B). This intervention induced a steady decline in body weight over 2 months, re-
sulting in ~12% weight loss (**Figure 6B, Supplementary Figure 7C-D**), which was due to a pre-
dominant loss in fat mass. Tissue-specific weight loss was also observed in the liver and epidid-
ymal white adipose tissue (eWAT) of CR mice (**Supplementary Figure 7E-H**). Metabolic pheno-
typing and behavioral analysis revealed comparable overall energy expenditure and oxygen
consumption between AL and CR mice (**Supplementary Figure 7I-L**). However, carbon dioxide
production and carbohydrate oxidation were higher in CR mice during the dark period, when
food was available (**Supplementary Figure 7M-P**), indicating a higher preference for carbohy-
drate metabolism in CR mice during active feeding periods and a rapid shift towards fatty acid
metabolism during inactivity and food withdrawal. To assess how CR influences glucose homeo-
stasis in aged mice, we performed mixed meal tolerance tests (MTTs; 2 g glucose/kg body
weight, Ensure Original). Baseline MTTs were conducted at 72 weeks, prior to CR. After 2
months of CR, old mice exhibited significantly improved glucose tolerance at 80 weeks, at-
tributed to enhanced insulin sensitivity (**Figure 6C**). This improvement was accompanied by
~50% reductions in fasting and glucose-stimulated beta cell insulin secretion (**Supplementary**
Figure 7Q). Similar to younger CR-treated mice[32], CR preserved beta cell function in aged mice
by reducing the relative insulin secretory demand required to maintain normoglycemia, demon-
strating that CR-mediated functional beta cell adaptation remains intact in aging (**Supplemen-**
tary Figure 7R).

We previously showed that CR modulates beta cell heterogeneity by activating pro-lon-
gevity and homeostasis pathways that enhance beta cell protein and organelle quality-control
mechanisms[32]. This transcriptional response is driven by the activation of TF-directed GRNs
that increase the expression of beta cell identity and longevity genes[32]. To determine whether
CR elicits similar adaptative programs to old beta cells during CR, we performed single nucleus
RNA sequencing (snRNA-seq, 10X genomics) on islet nuclei isolated from AL and CR-fed mice
(**Figure 6A**). A total of 32,380 cells were sequenced (AL=17,344, CR=15,306) to identify all ma-
jor islet cell types, including beta cells (*Ins1*, *Ins2*, and *Ero1b*), alpha cells (*Gcg*), delta cells
(*Sst*), endothelial cells (*Pecam1*), and pancreatic polypeptide cells (*Ppy*) (**Figure 6D-E, Supple-**
mentary Figure 8A). Approximately 65% of the sequenced cells were beta cells, and CR did
not impact the relative composition of the heterogeneous beta cell population or islet cytoarchi-
tecture composition (**Supplementary Figure 8B-H**). Using pseudobulk analysis of beta cells fol-
lowed by KEGG and STRING pathways analysis, we observed a narrow range of CR-mediated
effects on beta cell gene expression, limited to downregulation of few ER protein processing
and folding (*Hspa1a/b/HSP70, Hspa8, Hsp90ab1, Dnaja1/HSP40, and Calr)*, cell stress (*Ypel2*,
Txnip), and insulin synthesis (*Ins1*) genes (**Figure 6F and Supplementary Figure 6D-J**). These
data indicate that beta cell functional adaptation to CR in aging mice primarily reflects compen-
sation for age-related proteostatic stress from insulin production.

Remarkably, CR exerted a stronger transcriptional effect on alpha cells, modulating
nearly five times more genes than beta cells (184 vs. 36 differentially expressed genes; **Figure**
6F). Differential gene expression and pathway enrichment analyses of old CR alpha cell gene
signatures revealed upregulation of pathways associated with glucagon, glucose and fatty acid
metabolism (*Gcg, Gys2, and Acacb*) and Pgc1a (*Ppargc1a*) (**Figure 6G**). In contrast, CR sup-
pressed 159 genes involved in immune signaling, including pathways related to TCR signaling,
antigen presentation, and immune cell recruitment pathways (*Cd44, Cd28, Il7r, Il18r1, and Ccr7;*
Figure 6H). This included genes involved in cell-cell signaling via MHC class I (*H2-Q7*, *H2-Q6*,
H2-K1) and MHC class II (*H2-Aa*, *H2-Ab1*, *H2-Ob*) antigens **(Figure 6H-I)**. These findings
demonstrate that 2 months of mild CR reduces age-associated pro-inflammatory signaling in ag-
ing alpha cells, thus further supporting an immunoregulatory role for alpha cells during aging[53]
(Figures 2-4) and type 1 diabetes[73].

CR suppresses alpha cell immune signaling to reduce islet inflammation. To further
investigate how CR-mediated modulation of aging alpha cell inflammatory gene expression in-
fluences the cellular and molecular profile of islet immune cells, we quantified the relative com-
position and transcriptional signatures immune cell types identified by snRNA-seq (**Figure 7A**).
This revealed a significant decrease in B cells, T cells, proliferating B cells, and antigen present-
ing cells (APCs) in CR mouse islets, whereas the relative density and transcriptional profile of
islet-resident macrophages were unchanged (**Figure 7B**). Furthermore, old CR mouse islets
had a marked reduction in islet T cells, including a near total ablation of cytotoxic CD8[+] effector
T cells (**Figure 7C-D**). Here, the remaining CD8[+] T cells in CR islets had reduced expression of
effector markers granzyme K (*Gzmk*), IL-2 receptor (Il2ra/*Cd25*), Ccr7, and *Txk*; all markers of
activated T cells (**Figure 7E**). We validated these changes *in situ* using immunohistochemistry
and confocal microscopy of AL and CR old pancreases, followed by spatial mapping of alpha
cells, beta cells, and CD3[+] T cells (**Figure 7F)**. This approach revealed a significant reduction in
T-cell density (**Figure 7G**), and increased distance between T cells and alpha cells (**Figure 7H**).
These data support recent findings of immune cell infiltration in the aging mouse islet[61] and
demonstrate that CR modulates the recruitment of Cd8[+] T cells to the islet by reducing their
preferential proximity to old alpha cells.

To identify communication pathways linking aged alpha cells with CD8[+] T cells and their
modulation by CR, we performed receptor-ligand inference analysis using CellChat[74,75] to recon-
struct cell-cell communication networks in AL and CR aging mouse islets (**Figure 7I**)[76]. This ap-
proach revealed an overall downregulation of intercellular communication between islet immune
cells and alpha cells, beta cells, and delta cells (**Figure 7I**). This was underscored by a signifi-
cant decrease in Cd45-based interactions between old alpha cells and T- and B-cells in CR
mice (**Figure 7J**). These data further supports the idea that old alpha cells are important signal-
ing hubs that communicate with effector adaptive immune cells, including cytotoxic CD8[+] T cells
that are associated with islet inflammation in aging and T2D (**Figures 2-5**).

Finally, we investigated whether CR had similar anti-inflammatory effects in the pancreas
of NHPs on AL or 30% CR for at least 17 years[77] (**Figure 7K**). Importantly, like our CR mouse
models (Figure 6-7 and[78]), CR NHPs were more insulin sensitive and had reduced beta cell in-
sulin release (**Figure 7K**). In humans, islet inflammation during aging is associated with in-
creased lipid droplet accumulation, islet and vascular fibrosis, and islet amyloid accumula-
tion[37,38]. Therefore, we ran a double-blind analysis to quantify the impact of CR on these tissue
signatures in the pancreases of age-matched AL and CR NHPs (see methods for details). We
found that CR significantly decreased pancreatic lipid droplet accumulation and islet fibrosis
(**Figure 7L-N**), suggesting an underlying inflammatory may contribute to fibrosis in aged NHPs
and can be rescued by CR.

These results indicate that CR profoundly reshapes the aging islet immune environment
by suppressing alpha cell inflammatory and MHC-I signaling to reduce CD8$^+$ T-cell recruitment
and cytotoxicity, while diminishing intercellular communication between alpha cells and adaptive
immune cells. Moreover, the observation that long-term CR similarly attenuates inflammatory
and fibrotic features in NHPs suggests that CR promotes healthier islet and pancreatic aging
across species.

Discussion

Aging is associated with an active and low-grade inflammatory process in different tis-
sues, where inflammation is a key signature of several age-associated degenerative and meta-
bolic diseases, including Alzheimer's[79], and obesity and diabetes[80]. In the pancreas, aging is as-
sociated with increased islet macrophage infiltration and vascular fibrosis[37], and share other key
hallmarks of T1D and T2D islet pathophysiology such as beta cell ER stress, compromised beta
cell identity and function, and beta cell senescence[15,37,38,42-44,53,81-88]; however, recent studies
have implicated the activation of inflammatory responses in human alpha cells during T1D and
T2D[73,89].

In this manuscript, we identify activation of immune signaling and inflammation as a key
signature of aging human islets from non-diabetic individuals. This phenotype is defined by ac-
tive IFNγ signaling and MHC-I expression in old alpha cells, which is due to activation of IFNγ-
responsive TFs IRF1, NFKB1, and KLF6. In addition, old human alpha cells appear to be more
sensitive to amino-acid-dependent glucagon release, which could be explained by the observed
increase in the expression of several AA catabolism enzymes. Moreover, our data shows that
old alpha cells have increased immune signaling pathways (IL6, CD46), and that aging islets
have increased TGFB1 levels (and of other growth factors like FGF and LIFR) that signal mostly
from beta cells and that may act to promote and increase islet fibrosis. Importantly, activation of
alpha cell immune signaling during aging translates into increased physical interactions between
old alpha cells with cytotoxic CD8$^+$ T cells, which accumulate within the aging human islet micro-
environment. Strikingly, most old islet CD8$^+$ cells have a "memory-like" phenotype (CD27$^+$), thus
indicating that these cells have been exposed to T cell antigens. While our study does not iden-
tify which antigen is presented by alpha cell MHC-Is, a recent study where human islets were
exposed to cytokine stress (e.g., IFNγ) identified several new MHC-I antigens, many of which
are enriched in aging alpha cells (i.e., NFKB1, IRF9, TAP1, IFIT3, PSMB8, and STAT1)[81]. Im-
portantly, while density of islet CD8+ cells is similar between non-diabetic and T2D islets, most
T2D islet CD8+ cells have an effector-like signature (CD45RA$^+$ CD8$^+$). This data indicates that
infiltration of CD8+ cells in aging human islets occurs independently of T2D and further sug-
gests that islet inflammation in individuals with T2D onset during old age involves activation of
alpha cell-interacting and "resident" CD8+ cells.

Calorie restriction (CR) prolongs the lifespan and health span of multiple organisms,
from yeast to mice to non-human primates to humans[70-72,90-92]. The benefits of CR depend on
CR-dependent improvements in glucose homeostasis mechanisms, including enhanced insulin
sensitivity that reduces the metabolic demand for insulin release from beta cells[32,71,93,94]. Our ex-
periments with 80-week-old mice on CR for 2 months demonstrate that the insulin-sensitizing
and beta cell functional adaptation pathways promoted by CR are not impaired during aging, not
that these involve significant transcriptional reorganization in the beta cell. This is likely ex-
plained by the limited plasticity of aging beta cells in response to metabolic challenges[95]. In con-
trast, CR acted on old alpha cells and significantly suppressed the expression of pro-inflamma-
tory genes, including MHC-I and TCR signaling pathways. This CR-dependent modulation
dampened CD45-mediated communication between immune cells and alpha cells, which was
sufficient to reduce the density of T cells in aging islet periphery. Strikingly, a similar anti-

inflammatory effect was observed in NHPs kept on CR for most of their adult life, where CR pre-
vented the accumulation of pancreatic lipid droplets, and of islet fibrosis amyloid deposition.

In conclusion, our findings identify the relationship between alpha cells and immune cells
within the islet of Langerhans during aging that may underlie the onset of islet fibrosis and in-
flammation to promote T2D development during old age.

Data availability: IMC, CODEX, and human islet single cell RNAseq are available via PANC-
DB (Zenodo: 10.5281/zenodo.15588155). Mouse snRNA-seq is readily and freely available at
Zenodo and raw FASTQ data deposited at Gene Expression Omnibus.

Acknowledgements: We acknowledge the following Vanderbilt University (VU) and Vanderbilt
University Medical Center (VUMC) core facilities: VUMC Hormone Assay & Analytical Services
Core (NIH DK020593), VU Metabolic Mouse Phenotyping Center [VMMPC (NIH DK059637;
www.vmmpc.org)], Translational Pathology Shared Resource (NCI/NIH Cancer Center Support
Grant 5P30 CA68485-19), and Vanderbilt Technologies for Advanced Genomics (VANTAGE)
core laboratory. VANTAGE is supported in part by Clinical and Translational Science Award
Grant 5UL1 RR024975-03, Vanderbilt Ingram Cancer Center Grant P30 CA68485, Vanderbilt
Vision Center Grant P30 EY08126, and National Institutes of Health/National Center for Re-
search Resources Grant G20 RR030956. This manuscript used data acquired from the data-
base (https://hpap.pmacs.upenn.edu/) of the Human Pancreas Analysis Program (HPAP;
RRID:SCR_016202; and[56,57]). HPAP is part of a Human Islet Research Network
(RRID:SCR_014393) consortium (UC4-DK112217, U01-DK123594, UC4-DK112232, and U01-
DK123716). We would also like to acknowledge NIH grant R01AG040178. The WNPRC is sup-
ported by research facilities improvement programs NCRR/ORIP grants
P51RR000167/P51OD011106. This work includes data and/or analyses from HumanIslets.com
funded by the Canadian Institutes of Health Research, JDRF Canada, and Diabetes Canada (5-
SRA-2021-1149-S-B/TG 179092) with data from islets isolated by the Alberta Diabetes Institute
IsletCore with the support of the Human Organ Procurement and Exchange program, Trillium
Gift of Life Network, BC Transplant, Quebec Transplant, and other Canadian organ procurement
organizations with written informed donor consent as approved by the Human Research Ethics
Board at the University of Alberta (Pro00013094). Finally, we would like to acknowledge Dr Pat-
rick E MacDonald from ADI – Edmonton (Canada) for critical comments and inputs during data
collection and manuscript writing stages of this project.

Funding: This work was support by the National Institutes of Health grants 1R01DK138141
awarded to R.A.D. M.S. is supported by the National Institutes of Health Molecular Endocrinol-
ogy Training Program (T32DK007563).

Author Contributions: M.S., S.S., JP.C., R.C., and R.A.D designed experiments. M.S., A.C.,
M.C., G.F., S.S., JP.C., R.C., A.H., C.E., A.P., and R.A.D. collected data and assisted with data
analysis and project management. M.S., S.S., JP.C., and R.A.D. completed data analysis. M.S.
and R.A.D. interpreted data and wrote the manuscript.

Disclosures: The authors have nothing to disclose.

REFERENCES

1. National Diabetes Statistics Report. *Centers for Disease Control and Prevention* (2024).
2. Basu, R.*, et al.* Mechanisms of the age-associated deterioration in glucose tolerance: contribution of alterations in insulin secretion, action, and clearance. *Diabetes* **52**, 1738-1748 (2003).
3. Petersen, K.F.*, et al.* Mitochondrial dysfunction in the elderly: possible role in insulin resistance. *Science* **300**, 1140-1142 (2003).
4. Cowie, C.C.*, et al.* Prevalence of diabetes and impaired fasting glucose in adults in the U.S. population: National Health And Nutrition Examination Survey 1999-2002. *Diabetes Care* **29**, 1263-1268 (2006).
5. Unger, R.H., Aguilar-Parada, E., Müller, W.A. & Eisentraut, A.M. Studies of pancreatic alpha cell function in normal and diabetic subjects. *J Clin Invest* **49**, 837-848 (1970).
6. Raskin, P., Aydin, I., Yamamoto, T. & Unger, R.H. Abnormal alpha cell function in human diabetes: the response to oral protein. *Am J Med* **64**, 988-997 (1978).
7. Raskin, P., Aydin, I. & Unger, R.H. Effect of insulin on the exaggerated glucagon response to arginine stimulation in diabetes mellitus. *Diabetes* **25**, 227-229 (1976).
8. D'Alessio, D. The role of dysregulated glucagon secretion in type 2 diabetes. *Diabetes Obes Metab* **13 Suppl 1**, 126-132 (2011).
9. Cabrera, O.*, et al.* The unique cytoarchitecture of human pancreatic islets has implications for islet cell function. *Proceedings of the National Academy of Sciences* **103**, 2334-2339 (2006).
10. Cnop, M.*, et al.* The long lifespan and low turnover of human islet beta cells estimated by mathematical modelling of lipofuscin accumulation. *Diabetologia* **53**, 321-330 (2010).
11. Drigo, R.A.E.*, et al.* Age Mosaicism across Multiple Scales in Adult Tissues. *Cell Metabolism* **30**, 343-+ (2019).
12. Perl, S.*, et al.* Significant human beta-cell turnover is limited to the first three decades of life as determined by in vivo thymidine analog incorporation and radiocarbon dating. *J Clin Endocrinol Metab* **95**, E234-239 (2010).
13. López-Otín, C., Blasco, M.A., Partridge, L., Serrano, M. & Kroemer, G. Hallmarks of aging: An expanding universe. *Cell* **186**, 243-278 (2023).
14. Shrestha, S.*, et al.* Aging compromises human islet beta cell function and identity by decreasing transcription factor activity and inducing ER stress. *Science Advances* **8**, eabo3932 (2022).
15. Sandovici, I.*, et al.* Ageing is associated with molecular signatures of inflammation and type 2 diabetes in rat pancreatic islets. *Diabetologia* **59**, 502-511 (2016).
16. Li, L.*, et al.* Defects in β-cell Ca2+ dynamics in age-induced diabetes. *Diabetes* **63**, 4100-4114 (2014).
17. Fontana, L., Partridge, L. & Longo, V.D. Extending healthy life span--from yeast to humans. *Science* **328**, 321-326 (2010).
18. Mattison, J.A.*, et al.* Caloric restriction improves health and survival of rhesus monkeys. *Nat Commun* **8**, 14063 (2017).
19. Green, C.L., Lamming, D.W. & Fontana, L. Molecular mechanisms of dietary restriction promoting health and longevity. *Nat Rev Mol Cell Biol* **23**, 56-73 (2022).
20. Anderson, R.M., Le Couteur, D.G. & de Cabo, R. Caloric Restriction Research: New Perspectives on the Biology of Aging. *J Gerontol A Biol Sci Med Sci* **73**, 1-3 (2017).
21. Speakman, J.R. & Mitchell, S.E. Caloric restriction. *Mol Aspects Med* **32**, 159-221 (2011).

22.   Mao, K.*, et al.* Late-life targeting of the IGF-1 receptor improves healthspan and lifespan in
female mice. *Nat Commun* **9**, 2394 (2018).

23.   Liu, G.Y. & Sabatini, D.M. mTOR at the nexus of nutrition, growth, ageing and disease. *Nat*
Rev Mol Cell Biol **21**, 183-203 (2020).

24.   Il'yasova, D.*, et al.* Effects of 2 years of caloric restriction on oxidative status assessed by
urinary F2-isoprostanes: The CALERIE 2 randomized clinical trial. *Aging Cell* **17**(2018).

25.   Yang, L., Li, P., Fu, S., Calay, E.S. & Hotamisligil, G.S. Defective hepatic autophagy in obesity
promotes ER stress and causes insulin resistance. *Cell Metab* **11**, 467-478 (2010).

26.   Pyo, J.O.*, et al.* Overexpression of Atg5 in mice activates autophagy and extends lifespan.
Nat Commun **4**, 2300 (2013).

27.   Harrison, D.E.*, et al.* Rapamycin fed late in life extends lifespan in genetically
heterogeneous mice. *Nature* **460**, 392-395 (2009).

28.   Martin-Montalvo, A.*, et al.* Metformin improves healthspan and lifespan in mice. *Nat*
Commun **4**, 2192 (2013).

29.   Baur, J.A.*, et al.* Resveratrol improves health and survival of mice on a high-calorie diet.
Nature **444**, 337-342 (2006).

30.   Chen, C., Liu, Y., Liu, Y. & Zheng, P. mTOR regulation and therapeutic rejuvenation of aging
hematopoietic stem cells. *Sci Signal* **2**, ra75 (2009).

31.   Fraenkel, M.*, et al.* mTOR inhibition by rapamycin prevents beta-cell adaptation to
hyperglycemia and exacerbates the metabolic state in type 2 diabetes. *Diabetes* **57**, 945-
957 (2008).

32.   dos Santos, C.*, et al.* Calorie restriction increases insulin sensitivity to promote beta cell
homeostasis and longevity in mice. *Nature Communications* **15**, 9063 (2024).

33.   Matta, L.*, et al.* Chronic intermittent fasting impairs β cell maturation and function in
adolescent mice. *Cell Rep* **44**, 115225 (2025).

34.   Patel, S., Yan, Z. & Remedi, M.S. Intermittent fasting protects β-cell identity and function in a
type-2 diabetes model. *Metabolism* **153**, 155813 (2024).

35.   Brown, M.R.*, et al.* Time-restricted feeding prevents deleterious metabolic effects of
circadian disruption through epigenetic control of β cell function. *Sci Adv* **7**, eabg6856
(2021).

36.   Mateus Gonçalves, L., Pereira, E., Werneck de Castro, J.P., Bernal-Mizrachi, E. & Almaça, J.
Islet pericytes convert into profibrotic myofibroblasts in a mouse model of islet vascular
fibrosis. *Diabetologia* **63**, 1564-1575 (2020).

37.   Almaca, J.*, et al.* Young capillary vessels rejuvenate aged pancreatic islets. *Proceedings of*
the National Academy of Sciences of the United States of America **111**, 17612-17617
(2014).

38.   Wright, J.J.*, et al.* Exocrine Pancreas in Type 1 and Type 2 Diabetes: Different Patterns of
Fibrosis, Metaplasia, Angiopathy, and Adiposity. *Diabetes* **73**, 1140-1152 (2024).

39.   Ewald, J.D.*, et al.* HumanIslets.com: Improving accessibility, integration, and usability of
human research islet data. *Cell Metab* **37**, 7-11 (2025).

40.   Kolic, J.*, et al.* Proteomic predictors of individualized nutrient-specific insulin secretion in
health and disease. *Cell Metab* **36**, 1619-1633.e1615 (2024).

41.   Camunas-Soler, J.*, et al.* Patch-Seq Links Single-Cell Transcriptomes to Human Islet
Dysfunction in Diabetes. *Cell Metabolism* **31**, 1017-+ (2020).

42.   Aguayo-Mazzucato, C.*, et al.* Acceleration of beta Cell Aging Determines Diabetes and
Senolysis Improves Disease Outcomes. *Cell Metabolism* **30**, 129-+ (2019).

43.   Shrestha, S.*, et al.* Aging compromises human islet beta cell function and identity by
585   decreasing transcription factor activity and inducing ER stress. *Science Advances* 8, 17
586   (2022).
587   44.   Enge, M.*, et al.* Single-Cell Analysis of Human Pancreas Reveals Transcriptional Signatures
588   of Aging and Somatic Mutation Patterns. *Cell* 171, 321-330.e314 (2017).
589   45.   Tritschler, S., Theis, F.J., Licked, H. & Bottcher, A. Systematic single-cell analysis provides
590   new insights into heterogeneity and plasticity of the pancreas. *Molecular Metabolism* 6,
591   974-990 (2017).
592   46.   Augsornworawat, P., Maxwell, K.G., Velazco-Cruz, L. & Millman, J.R. Single-Cell
593   Transcriptome Profiling Reveals β Cell Maturation in Stem Cell-Derived Islets after
594   Transplantation. *Cell Rep* 32, 108067 (2020).
595   47.   Chiou, J.*, et al.* Single-cell chromatin accessibility identifies pancreatic islet cell type- and
596   state-specific regulatory programs of diabetes risk. *Nature Genetics* 53, 455-+ (2021).
597   48.   Bramswig, N.C.*, et al.* Epigenomic plasticity enables human pancreatic α to β cell
598   reprogramming. *J Clin Invest* 123, 1275-1284 (2013).
599   49.   Avrahami, D.*, et al.* Aging-Dependent Demethylation of Regulatory Elements Correlates
600   with Chromatin State and Improved β Cell Function. *Cell Metabolism* 22, 619-632 (2015).
601   50.   Avrahami, D.*, et al.* Single-cell transcriptomics of human islet ontogeny defines the
602   molecular basis of β-cell dedifferentiation in T2D. *Mol Metab* 42, 101057 (2020).
603   51.   Su, C.*, et al.* 3D chromatin maps of the human pancreas reveal lineage-specific regulatory
604   architecture of T2D risk. *Cell Metabolism* 34, 1394-+ (2022).
605   52.   Arda, H.E.*, et al.* Age-Dependent Pancreatic Gene Regulation Reveals Mechanisms
606   Governing Human beta Cell Function. *Cell Metabolism* 23, 909-920 (2016).
607   53.   Staels, W.*, et al.* Comprehensive alpha, beta, and delta cell transcriptomics reveal an
608   association of cellular aging with MHC class I upregulation. *Mol Metab* 87, 101990 (2024).
609   54.   Fasolino, M.*, et al.* Single-cell multi-omics analysis of human pancreatic islets reveals novel
610   cellular states in type 1 diabetes. *Nature Metabolism* 4, 284-+ (2022).
611   55.   Patil, A.R.*, et al.* Single-cell expression profiling of islets generated by the Human Pancreas
612   Analysis Program. *Nat Metab* 5, 713-715 (2023).
613   56.   Shapira, S.N., Naji, A., Atkinson, M.A., Powers, A.C. & Kaestner, K.H. Understanding islet
614   dysfunction in type 2 diabetes through multidimensional pancreatic phenotyping: The
615   Human Pancreas Analysis Program. *Cell Metab* 34, 1906-1913 (2022).
616   57.   Kaestner, K.H., Powers, A.C., Naji, A. & Atkinson, M.A. NIH Initiative to Improve
617   Understanding of the Pancreas, Islet, and Autoimmunity in Type 1 Diabetes: The Human
618   Pancreas Analysis Program (HPAP). *Diabetes* 68, 1394-1402 (2019).
619   58.   van de Sande, B.*, et al.* A scalable SCENIC workflow for single-cell gene regulatory network
620   analysis. *Nature Protocols* 15, 2247-2276 (2020).
621   59.   Wu, G. & Haw, R. Functional Interaction Network Construction and Analysis for Disease
622   Discovery. *Methods Mol Biol* 1558, 235-253 (2017).
623   60.   Zeng, F., Zhang, M.Z., Singh, A.B., Zent, R. & Harris, R.C. ErbB4 isoforms selectively regulate
624   growth factor induced Madin-Darby canine kidney cell tubulogenesis. *Mol Biol Cell* 18,
625   4446-4456 (2007).
626   61.   Jin, S.*, et al.* Inference and analysis of cell-cell communication using CellChat. *Nat
627   Commun* 12, 1088 (2021).
628   62.   Akhmetshina, A.*, et al.* Activation of canonical Wnt signalling is required for TGF-β-
629   mediated fibrosis. *Nat Commun* 3, 735 (2012).
630   63.   Nguyen, H.N.*, et al.* Leukemia inhibitory factor (LIF) receptor amplifies pathogenic
631   activation of fibroblasts in lung fibrosis. *Proc Natl Acad Sci U S A* 121, e2401899121 (2024).

64.   Arbore, G.*, et al.* Complement receptor CD46 co-stimulates optimal human CD8. *Nat*
Commun **9**, 4186 (2018).
65.   Hansen, A.S.*, et al.* CD46 is a potent co-stimulatory receptor for expansion of human IFN-γ-
producing CD8. *Immunol Lett* **200**, 26-32 (2018).
66.   Ellingsgaard, H.*, et al.* Interleukin-6 enhances insulin secretion by increasing glucagon-like
peptide-1 secretion from L cells and alpha cells. *Nat Med* **17**, 1481-1489 (2011).
67.   Im, J.Y.*, et al.* DDIAS promotes STAT3 activation by preventing STAT3 recruitment to PTPRM in
lung cancer cells. *Oncogenesis* **9**, 1 (2020).
68.   Dos Santos, C.*, et al.* Caloric restriction promotes beta cell longevity and delays aging and
senescence by enhancing cell identity and homeostasis mechanisms. *Res Sq* (2023).
69.   Duregon, E.*, et al.* Prolonged fasting times reap greater geroprotective effects when
combined with caloric restriction in adult female mice. *Cell Metab* **35**, 1179-1194.e1175
(2023).
70.   Mitchell, S.J.*, et al.* Effects of Sex, Strain, and Energy Intake on Hallmarks of Aging in Mice.
Cell Metabolism **23**, 1093-1112 (2016).
71.   Pak, H.H.*, et al.* Fasting drives the metabolic, molecular and geroprotective effects of a
calorie-restricted diet in mice. *Nature Metabolism* **3**, 1327-+ (2021).
72.   Spadaro, O.*, et al.* Caloric restriction in humans reveals immunometabolic regulators of
health span. *Science* **375**, 671-+ (2022).
73.   Dos Santos, T.*, et al.* Altered immune and metabolic molecular pathways drive islet cell
dysfunction in human type 1 diabetes. *J Clin Invest* (2025).
74.   Coomans de Brachène, A.*, et al.* Exercise as a non-pharmacological intervention to protect
pancreatic beta cells in individuals with type 1 and type 2 diabetes. *Diabetologia* **66**, 450-
460 (2023).
75.   Carapeto, P.*, et al.* Exercise activates AMPK in mouse and human pancreatic islets to
decrease senescence. *Nature Metabolism* (2024).
76.   Al Jobori, H.*, et al.* Empagliflozin Treatment Is Associated With Improved β-Cell Function in
Type 2 Diabetes Mellitus. *J Clin Endocrinol Metab* **103**, 1402-1407 (2018).
77.   Lee, Y.S., Lee, C., Choung, J.S., Jung, H.S. & Jun, H.S. Glucagon-Like Peptide 1 Increases β-
Cell Regeneration by Promoting α- to β-Cell Transdifferentiation. *Diabetes* **67**, 2601-2614
(2018).
78.   Hahn, O.*, et al.* A nutritional memory effect counteracts benefits of dietary restriction in old
mice. *Nat Metab* **1**, 1059-1073 (2019).
79.   Kellogg, C.M.*, et al.* Microglial MHC-I induction with aging and Alzheimer's is conserved in
mouse models and humans. *Geroscience* **45**, 3019-3043 (2023).
80.   SantaCruz-Calvo, S.*, et al.* Adaptive immune cells shape obesity-associated type 2
diabetes mellitus and less prominent comorbidities. *Nat Rev Endocrinol* **18**, 23-42 (2022).
81.   Nanaware, P.P.*, et al.* The antigen presentation landscape of cytokine-stressed human
pancreatic islets. *Cell Rep* **44**, 115927 (2025).
82.   Marselli, L.*, et al.* beta-Cell inflammation in human type 2 diabetes and the role of
autophagy. *Diabetes Obesity & Metabolism* **15**, 130-136 (2013).
83.   Jelleschitz, J.*, et al.* Insulitis and aging: Immune cell dynamics in Langerhans islets. *Redox*
Biol **82**, 103587 (2025).
84.   De Burghgrave, M.*, et al.* Pancreatic Islet Cells Response to IFNγ Relies on Their Spatial
Location within an Islet. *Cells* **12**(2022).
85.   Avrahami, D.*, et al.* Single-cell transcriptomics of human islet ontogeny defines the
molecular basis of beta-cell dedifferentiation in T2D. *Molecular Metabolism* **42**, 14 (2020).

86.   Horwitz, E.*, et al.* β-Cell DNA Damage Response Promotes Islet Inflammation in Type 1
680   Diabetes. *Diabetes* **67**, 2305-2318 (2018).

681   87.   Aguayo-Mazzucato, C.*, et al.* beta Cell Aging Markers Have Heterogeneous Distribution and
682   Are Induced by Insulin Resistance. *Cell Metabolism* **25**, 898-+ (2017).

683   88.   Thompson, P.J.*, et al.* Targeted Elimination of Senescent Beta Cells Prevents Type 1
684   Diabetes. *Cell Metab* **29**, 1045-1060.e1010 (2019).

685   89.   Dai, X.Q.*, et al.* Heterogenous impairment of alpha cell function in type 2 diabetes is linked
686   to cell maturation state. *Cell Metabolism* **34**, 256-+ (2022).

687   90.   Longo, V.D. & Anderson, R.M. Nutrition, longevity and disease: From molecular
688   mechanisms to interventions. *Cell* **185**, 1455-1470 (2022).

689   91.   Mattison, J.A.*, et al.* Impact of caloric restriction on health and survival in rhesus monkeys
690   from the NIA study. *Nature* **489**, 318-321 (2012).

691   92.   Civitarese, A.E.*, et al.* Calorie restriction increases muscle mitochondrial biogenesis in
692   healthy humans. *PLoS Med* **4**, e76 (2007).

693   93.   da Amaral, M.E.C.*, et al.* Caloric restriction recovers impaired beta-cell-beta-cell gap
694   junction coupling, calcium oscillation coordination, and insulin secretion in prediabetic
695   mice. *American Journal of Physiology-Endocrinology and Metabolism* **319**, E709-E720
696   (2020).

697   94.   Sheng, C.J.*, et al.* Reversibility of beta-Cell-Specific Transcript Factors Expression by Long-
698   Term Caloric Restriction in db/db Mouse. *Journal of Diabetes Research* **2016**, 11 (2016).

699   95.   Tschen, S.I., Dhawan, S., Gurlo, T. & Bhushan, A. Age-dependent decline in beta-cell
700   proliferation restricts the capacity of beta-cell regeneration in mice. *Diabetes* **58**, 1312-1320
701   (2009).

702   96.   Ritchie, M.E.*, et al.* limma powers differential expression analyses for RNA-sequencing and
703   microarray studies. *Nucleic acids research* **43**, e47-e47 (2015).

704   97.   Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
705   RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

706   98.   Jin, S., Plikus, M.V. & Nie, Q. CellChat for systematic analysis of cell-cell communication
707   from single-cell transcriptomics. *Nat Protoc* **20**, 180-219 (2025).

708   99.   Hao, Y.*, et al.* Dictionary learning for integrative, multimodal and scalable single-cell
709   analysis. *Nat Biotechnol* **42**, 293-304 (2024).

710   100.   Wang, Y.J.*, et al.* Single-Cell Mass Cytometry Analysis of the Human Endocrine Pancreas.
711   *Cell Metabolism* **24**, 616-626 (2016).

712   101.   Colman, R.J.*, et al.* Caloric restriction delays disease onset and mortality in rhesus
713   monkeys. *Science* **325**, 201-204 (2009).

Figure Legends

Figure 1 - Aging human islets have an inflammatory activation phenotype combined with
mTORC1 signaling and elevated unfolded protein response stress. (A) Islets from young
(28±4 years, n=26), and old (71±3 years, n=19) ND donors from the HumanIslets consortium
were compared by proteomics. **(B)** Reactome pathways present from significantly abundant pro-
teins between young and old ND donors. **(C-G)** Protein expression levels for beta cell immatu-
rity marker, RBP4 (C); senescence, CDKN1A/p21 (D); unfolded protein response, CHOP
(DDIT3), ATF6, PERK (EIF2AK3) (E); mTORC1 activation, LAMP2, LAMTOR, RAGA, RAGC
(F); and immune regulatory components, HLA-A, TGF-β1, TOM1, SLC15A4, CD44, and CD28,
respectively (G). HumanIslets, Human Islet Data Analysis and Sharing consortium; HLA, human
leukocyte antigen; TCR, T cell receptor; UPR, unfolded protein response. *$p<0.05$, young vs. old
donors. Error bars are mean ± SD.

Figure 2 - Aging increases human alpha cell inflammatory and metabolic signatures. (A)
Age and sex stratification for young (n=11; F=6, M=5) and old (n=21; F=11, M=10) ND donors
for scRNA-seq integration, analysis, and SCENIC TF activity. **(B)** Cell type UMAP of integrated
scRNA-seq data obtained from HPAP database. **(C)** HLA-ABC expression across endocrine and
non-endocrine cell types in human islets. **(D-E)** GSEA hallmarks for gene sets Hallmark Inter-
feron Alpha Response (M5911) and Hallmark Interferon GAMMA Response (M5913). **(F)** KEGG
hallmarks for Antigen Processing and Presentation by MHC Class I Molecules (M47529). **(G-I)**
Significantly expressed Hallmark genes for metabolic processes including Glycolysis (M5937;
G), Fatty acid metabolism (M5935; H), and gene ontology pathways for Amino Acid Metabolic
Process (GO:0006520; I), respectively. **(J)** Isolated human islet perifusion assay including stim-
ulation with amino-acids, glucose, IBMX, and KCL. Dotted line boxes (a', b', and c') highlight
specific stimuli that provoked significant glucagon secretion from old islets versus young islets.
P-values are shown. Data are mean ± SD with boxplots representing mean ± IQR. Welch's two
sample *t-test* was used for panel J, a'-c'. scRNA-seq, single-cell RNA sequencing. TF, transcrip-
tion factor; ND, non-diabetic. HPAP, Human Pancreas Analysis Program; GSEA, gene set en-
richment analysis.

Figure 3 - Re-wiring of gene regulatory networks activates an inflammatory phenotype in
aging alpha cells. (A-B) Gene Regulatory Network (GRN) maps of young (A) and old (B) hu-
man alpha cells. Transcription factors (TFs) are represented by white hexagons and inferred TF
target genes are shown in solid grey circles. Only the top 15% TF regulons after 100X
pySCENIC inference simulations are shown. TF hexagon size represents the TF importance
metric within the GRN structure, and TF-target gene nodes are connected by line edges colored
using a *viridis* hue to represent predicted connection strength. Disconnected and smaller nodes
are shown at the bottom of each GRN network. **(C)** Venn diagrams representing unique and
shared genes found in the regulons associated with IRF1, NFKB1, KLF6, MAFB, and HNF1A
TFs. **(D)** Regulatory network map displaying age-dependent and age-independent regulatory
connections between inflammatory TFs and canonical alpha cell identity TFs. In this map, TFs
are represented by solid grey circles and regulatory relationships are shown by arrows. Green
arrows, connections found in both young and old alpha cells; dark blue and beige arrows repre-
sent connections found uniquely in old and young alpha cells, respectively. **(E-F)** Close up of a
Reactome Pathway map displaying significantly enriched pathways associated with genes
found in aging alpha cell IRF1, NFKB1, and KLF6 regulons. Different tones of magenta hue rep-
resent a false discovery rate (FDR) of at least 0.05 or lower. **(G)** Heatmaps displaying age-de-
pendent changes in outgoing ("senders") and incoming ("receivers") cell-cell communication
pathways significantly modulated in old islet cells versus young islet cells. Pathway targets were
clustered into functional categories (shown x-axis) and their enrichment value for each old cell
type (in the y-axis) is shown. Positive values indicate pathways enriched in old islet cells. **(H-I)**

Circle plots illustrating growth factor (TGFB1 and FGF) and immune signaling pathways (IL6,
CD46) activated in old islets cell types. FGF, IL6, and CD46 were identified as significantly en-
riched pathways in old alpha cells. Each cell type is represented by a color and color legend is
shown at the bottom.

Figure 4 - Aging increases CD8$^+$ islet infiltration via alpha cell MHC-I signaling. (A) Imag-
ing IMC workflow. FFPE pancreatic tissue sections (4-8µm) were labeled simultaneously with 37
metal-conjugated antibodies. Laser ablation was conducted for 1,000 x 1,000 µm ROIs selected
around islet clusters, and signal quantification was measured by CyTOF. **(B)** UMAPs of subset
pancreas cells from all donors, highlighting the islet-endocrine cells (top) and immune cells (bot-
tom). **(C)** Representative IMC images and spatial reconstructions of islet-endocrine (top) and
islet-immune (bottom) fractions. Scale bar = 100µm **(D)** CODEX workflow and schematic for im-
age acquisition and processing of multiplexed datasets generated from 54 markers. **(E)** UMAPs
of pancreas cell subsets obtained through CODEX imaging, highlighting endocrine cells (top)
and immune subsets (bottom). **(F)** Representative CODEX images and spatial reconstructions
for islet endocrine (top) and immune (bottom) subsets. Scale bar = 2000µm. **(G-H)** Immune cell
proportions within the total pancreatic immune pool, collected by both IMC (G) and CODEX (H)
imaging platforms. **(I)** Density plots illustrating networking distance between CD8$^+$ T cells dis-
tance and alpha cells, median CD8$^+$-alpha cell distance, and the percentage of alpha cells local-
ized within ≤50µm of CD8$^+$ T cells. **(J)** Density plots showing network distance between CD8$^+$ T
cells and beta cells, median CD8$^+$-beta cell distance, and the percentage of beta cells localized
within ≤50µm of CD8$^+$ T cells. **(K)** Representative IMC images illustration alpha cells (GCG, red)
beta cells (C-peptide, white), and scaled HLA-ABC expression. Scale bar = 100µm. **(L-M)**
Quantification of HLA-ABC and HLA-DR intensity from segmented alpha cells (L) and beta cells
(M). IMC, imaging mass cytometry; CyTOF, Cytometry by time-of-flight; ROI, region of interest.
Welch's two-sample *t-test* was used to compare young and old donors in panels G-J and L-M.
Data are mean ± SEM (panels G-J), median ± SEM (L-M).

Figure 5 - T2D expands pancreatic CD8$^+$ T-cell populations and shifts islet CD8$^+$ T-cell
states from towards terminal effector-memory phenotypes. (A) Clinical characteristics ND
and T2D adults included for HPAP CODEX imaging. **(B)** UMAP of integrated pancreas samples
(942,724 cells) were segmented into immune cells (195,140 cells) and islet endocrine (99,744
cells) populations. **(C)** Identified islet endocrine cell populations, including alpha (GCG$^+$), beta
(C-PEP$^+$), "fibrotic" beta cells (C-PEP$^+$, COL4A1$^+$, COL6$^+$), delta (SST$^+$), pancreatic polypeptide
(PP$^+$), and epsilon (ghrelin$^+$) cells. **(D)** Scaled beta cell expression of C-PEP, CHGA, NKX6-1
and Ki67. **(E)** Scaled alpha cell expression of SOX9 and Ki67. **(F)** Proportions of major pancre-
atic immune subsets, including M2-like macrophages (CD163$^+$CD68$^+$), M1-like macrophages
(CD68$^+$CD163$^-$), granulocytes/monocytes (CD66b$^+$), CD8$^+$ T cells (CD3$^+$CD8$^+$), and CD4$^+$ T cells
(CD3$^+$CD4$^+$). **(G)** Immune cell density expressed as cells per mm^2. **(H)** CyTOF analysis of islets
isolated from young ND (n=12), old ND(n=15), and T2D (n=13) donors. **(I)** Representative gat-
ing of viable CD45$^+$ immune cells (left) and CD3$^+$CD8$^+$ T cells (right), sourced from HPAP-164.
(J) Proportion of CD45$^+$ immune cells among viable islet cells (left) and CD3$^+$CD8$^+$ within the
CD45$^+$ compartment (right). **(K)** Representative two-parameter (CD27 x CD45RA) plot illustrat-
ing CD8$^+$ T-cell differentiation states; (1) naïve CD27$^+$CD45RA$^+$, (2) central memory
(CD27$^+$CD45RA$^-$), 3) and terminal effector memory (CD27$^-$CD45RA$^+$). **(L)** Distribution of CD8$^+$ T
cells across naïve CD27$^+$CD45RA$^+$; CD8$_{naïve}$), central memory (CD27$^+$CD45RA$^-$; CD8$_{CM}$), and
terminal effector memory (CD27$^-$CD45RA$^+$; CD8$_{TEM}$) phenotypes. **(M)** CD8$^+$ T$_{EM}$/T$_{CM}$ ratio, repre-
senting the relative expansion of terminal effector-memory CD8$^+$ T cells compared with the cen-
tral memory pool. Welch's two sample *t-test* was used for comparisons between pool ND and
T2D donors in panels A and C-G. One-way *ANOVA* compared young, old and T2D donors in
panels J and L-M. DC, dendritic cell; cDC1, conventional cell; PP, pancreatic polypeptide; NK,

natural killer. *p< 0.05, **p<0.01 for age (old vs. young) and ND vs. T2D groups. Data are mean
± SEM.

Figure 6 - Late-life CR improves glucose tolerance while downregulating alpha cell TCR
activation and antigen presentation and processing. (A) 72-week-old C57Bl/6J mice were
fed ad-libitum (AL, control) or subjected to 20% calorie restriction (CR) for 2 months. Islets were
collected at 80 weeks and processed for IHC and snRNA-seq. **(B)** Percent change in body
weight relative to baseline. **(C)** MTTs were performed at baseline (72 weeks) and after the inter-
vention (80 weeks). AUC is presented for each test before and after dietary intervention. **(D)**
UMAP of 32,380 islet cells. **(E)** Selected marker genes for endocrine islet subpopulations by
scaled gene expression and percent cell expression. **(F)** Number of DEGs affected by CR in en-
docrine/endothelial cell populations. **(G)** KEGG pathways significantly modified by CR in alpha
cells. **(H)** Protein-protein interaction networks predicted from genes downregulated by CR in al-
pha cells, and subset by 'leukocyte activation' subcluster in red. **(I)** Scaled single-cell expression
of MHC class I and II genes in alpha cells. AUC, area under curve; IHC, immunohistochemistry;
MTT, mixed meal tolerance test; snRNA-seq, single nucleus RNA sequencing. #p< 0.05 for die-
tary groups (AL vs. CR); ****p<0.0001 for intervention (Pre- vs. Post-CR).

Figure 7 - Late-life CR attenuates islet adaptive CD8$^+$ effector phenotypes and reduces al-
pha cell-mediated T-cell recruitment. (A) UMAP of subset immune cell populations (B cells, T
cells, macrophages, proliferating B cells, and APCs [dendritic cells] from AL and CR groups. **(B)**
Proportion of immune cell populations relative to total islet cells. **(C)** UMAP of T-cell populations
from Al and CR groups. **(D)** Proportion of total T cells; * denotes effector CD8$^+$ T cells. **(E)** Ex-
pression of Th1-associated genes: *Gzmk*, *Cd25/Il2ra*, *CCr7*, and *Txk*. **(F)** Representative IHC
staining for alpha cells (glucagon, red), beta cells (insulin, green) and T cells (CD3, purple).
Scale bar = 100µm. **(G)** Pancreatic T-cell density (#CD3$^+$ cells per mm^2). **(H)** T-cell-endocrine
cell networks, quantifying the average distance of CD3$^+$ T cells to alpha cells (left), and beta
cells (right). **(I)** Intercellular communication networks showing total interaction number relative to
interaction signal strength; edge thickness represents relative signal strength. **(J)** Network plot
of inferred CD45 signaling interactions. **(K)** Twelve age-matched non-human primates (*Macaca*
mulatta) were maintained on AL or 30% CR diets, starting at 7-10 years of age and assessed
longitudinally for metabolic outcomes, including peripheral insulin sensitivity via FSIVGTT, until
natural death. **(L)** Pancreatic sections (5µm) were stained with H&E and Gomori's Trichrome;
islets were manually subset and quantified per image. **(M)** Lipid droplet area, expressed in ab-
solute (#/pancreas) and relative density (#/mm2). **(N)** Trichrome-positive area per islet, subset
per primate pancreas and per islet analyzed. H&E, hematoxylin and eosin; FSIVGTT, fasting
state intravenous glucose tolerance test. Welch's two sample *t-test* was used for comparisons
between AL and CR groups. *p<0.05, ***p<0.001, AL vs. CR. Data are mean ± SEM.

Imaging Mass Cytometry

A. Young (n=18) 20-40 years / Old (n=6) 40-70 years

B. Endocrine cells
- Beta
- Alpha
- Delta
- PP/Ghrelin
- Endothelial
- Mix

Immune cells
- Mφ
- Monocyte
- CD4 T
- CD8 T
- B Cell
- NK

C. DNA CPEP GCG SST — Young HPAP-080 / Old HPAP-066

DNA CD163 CD14 CD4 CD8 CD20 ISLET — Young HPAP-080 / Old HPAP-066

CODEX

D. Young (n=11) 20-40 years / Old (n=9) 40-70 years

E. Endocrine cells
- Beta
- Alpha
- Delta
- Mix
- PP
- Ghrelin
- Stromal

Immune cells
- M2 Mφ
- M1 Mφ
- Mono./Granulo
- cDC1
- DC
- CD4 T
- CD8 T
- B cell

F. DNA CPEP GCG SST Ghrelin PP — Young HPAP-013 / Old HPAP-174

DNA CD163 CD66b CD4 CD8 CD19 ISLET — Young HPAP-013 / Old HPAP-174

G. Imaging Mass Cytometry — % immune cells
- Mono.
- Mφ
- CD8 T
- NK
- CD4 T
- B Cell

young / old

CD8+ T cells per mm² — *
CD4 T cells per mm²

H. CODEX — % immune cells
- M1 Mφ
- M2 Mφ
- Gran./Mono
- CD8 T
- CD4 T
- B Cell
- DC
- cDC1

young / old

%CD8+ of CD3+ — *
%CD4+ of CD3+ — *

I. Alpha cell distance to CD8+ T cells
Density / distance (μm)
median distance (μm) — p=0.10 — Young Old
percent <50μm — * — % Alpha Cells — Young Old

J. Beta cell distance to CD8+ T cells
Density / distance (μm)
median distance (μm) — Young Old
Percent <50μm — % Beta Cells — Young Old

K. CPEP GCG — HLA-ABC
Young HPAP-019 / Inset
Old HPAP-066 / Inset

L. Alpha cells
HLA-ABC — * — young old
HLA-DR — young old
Intensity

M. Beta cells
HLA-ABC — * — young old
HLA-DR — young old
Intensity

A Immune cells

Macrophages
T-cell
APC
prol. B-cell
B-cell

UMAP 2 / UMAP 1

B Proportion of islet cells

% islet cells

- B cell
- T cell
- Macrophage
- Prol. B cell
- Dendritic cell

AL / CR

C T cells

UMAP 2 / UMAP 1

D Proportion of T cells

% T cells

- Exhausted CD 4/8+
- Effector CD8+
- Naive/memory CD4+
- Effector CD8+
- Stromal/Other
- Effector CD4+

AL * * / CR *

E T cell cytotoxicity

Gzmk   *Il2ra*

Ccr7   *Txk*

UMAP 2 / UMAP 1

F DAPI GCG Insulin CD3

AL — *Inset*
CR — *Inset*
100 μm

G T cell density

CD3+ per mm²

AL / CR   *

H CD3+ T cell distances

T cell - Alpha cell
CD3+ distance to alpha cells (μm)
AL / CR   *

T cell - Beta cell
CD3+ distance to beta cells (μm)
AL / CR   ns

I Intercellular communication
Interaction weights/strength

B-cell EC Delta
Ppy Alpha
T-cell APC
SMA prol. B-cell
Macrophages

AL / CR

J CD45 Signaling

AL / CR

K

Control or 30% CR diet

Age

Insulin Sensitivity Index: S_i
FSIVGTT

10^{-4}min^{-1} per μU/mL

15 20 25
Age (years)
* * * p=0.08

L   Control   CR

H&E   *Inset*   *Inset*

Trichrome *Inset*   *Inset*

M Lipid droplet area

Lipid droplet number
Control / CR   *

per mm²
Control / CR   *

N Islet trichrome+ area

% positive
Control / CR   p=0.088

Control / CR   ***