1 **Alpha cell inflammation during human pancreas aging and type 2 dia-**
2 **betes and its reversal by calorie restriction in mice.**
3

4 Michael Schleh[1], Jean-Philippe Cartailler[2], Shristi Shrestha[2], Amanda Cambraia[1], Aliyah Ha-
5 bashy[1], Cara Ellis[3], Melanie Cutler[1], Gabriel Ferguson[1], Alvin Powers[4], Rafael Arrojo e
6 Drigo[1,4,5,6*]
7
[1] Department of Molecular Physiology and Biophysics, Vanderbilt University School of Medicine,
Nashville, TN
[2] Creative Data Solutions, Center for Stem Cell Biology, Vanderbilt University School of Medi-
cine, Nashville, TN
[3] Alberta Diabetes Institute and Department of Pharmacology, University of Alberta, Edmonton,
Canada
[4] Vanderbilt Diabetes Research and Training Center, Vanderbilt University Medical Center,
Nashville, TN
[5] Center for Cell Systems Biology, Vanderbilt University School of Medicine, Nashville, TN
[6] Mass Spectrometry Research Center, Vanderbilt University School of Medicine, Nashville, TN
*Corresponding author

Keywords: aging, alpha cells, antigen presentation, calorie restriction, T cells

Address of Correspondence:
Rafael Arrojo e Drigo
718 Light Hall
2215 Garland Ave
Nashville, Tennessee 37232
r.drigo@vanderbilt.edu

Abstract **(145/150)**
Aging increases risk for type 2 diabetes (T2D), partly through loss of beta cell identity and func-
tion driven by metabolic stress and islet inflammation. While calorie restriction (CR) promotes
beta cell longevity in young animals, its impact on cellular aging and inflammatory burden in
older individuals is unclear. Using SCENIC regulon and multiomic analyses, we find that aging
human alpha cells adopt a coordinated inflammatory phenotype marked by IFN-γ signaling to
increase major histocompatibility complex (MHC) class I presentation, and CD8[+] T cell recruit-
ment and activation towards islets. In T2D, CD8[+] T cells further progress towards an effector
memory state. CR reduces alpha cell MHC-I expression while subsequently suppressing CD8[+]
effector status and accompanied by reduced islet inflammation and immune cell infiltration in
mice. Together, these findings highlight an alpha cell–immune signaling axis in aging and T2D
that may underlie fibrosis and disease pathophysiology.

Introduction

Aging is a major risk factor for type 2 diabetes (T2D), with prevalence reaching 28.8%
among adults aged ≥65 years in the United States (*1*). Age-related deterioration in glycemic
control has largely been attributed to progressive insulin resistance, impaired beta cell insulin
secretion, and altered hepatic insulin clearance (*2-4*). However, aging is also characterized by
tissue loss of homeostatic resilience and increased immune heterogeneity, that may shape met-
abolic health decline (*5*). A central feature of adaptive immune aging is thymic involution, which
reduces naïve T cell output and shifts the T cell compartment towards peripheral expansion and
memory biased populations (*6-8*). Whether such age-associated immune remodeling arises pri-
marily through intrinsic programs or occurs secondary to extrinsic cues in the microenvironment
and whether these aspects are present during diabetes, remains largely unknown.

Pancreatic islets are composed primarily of long-lived endocrine alpha- and beta cells,
which maintain glucose homeostasis through coordinated secretion of glucagon and insulin (*9-*
12). During aging, beta cells exhibit structural and functional decline, including loss of cellular
identity, DNA damage accumulation, endoplasmic reticulum (ER) stress, impaired autophagy,
increased transcriptional noise, amyloid deposition, senescence, and accumulate islet fibrosis
(*2, 13-16*). In contrast, far less is known how aging affects alpha cell biology despite well-known
disruptions in alpha cell glucagon secretion during aging and along the natural history of T2D (*2,*
17-20) that are accompanied by exaggerated amino acid-induced glucagon release (*21, 22*).
Moreover, it remains unknown whether aging promotes intrinsic inflammatory programs within
specific endocrine cell types, or whether islet inflammation is driven predominantly by extrinsic
immune cues. Defining the cellular origins of age-associated islet inflammation is essential to
understanding the intersection of immune and metabolic aging and how these aspects may con-
tribute to diabetes pathogenesis.

Calorie restriction (CR) is a well-established dietary intervention that extends lifespan
and health span across species by delaying several hallmarks of cellular dysfunction during ag-
ing (*23-26*). Moderate energy restriction (10–40%) improves metabolic homeostasis in both
young and aged animals, in part through modulation of nutrient-sensing pathways including in-
sulin-IGF1, mTORC1, and AMPK (*27, 28*). CR also reduces oxidative stress and inflammatory
tone while enhancing autophagy and proteostasis (*29-31*), and it can promote anti-inflammatory
responses in lean adult humans (*32*). In young adult mice, CR and CR-mimicking interventions
improve beta cell function and longevity, and which are thought to be explained by enhanced
peripheral insulin sensitivity (*33-36*). However, whether CR can reverse established age-associ-
ated islet inflammation, particularly within specific endocrine cell populations, remains unre-
solved.

Here, we integrated multimodal and multiomic analyses of publicly available human islet
datasets – including bulk proteomics, singe-cell RNA sequencing (scRNA-seq) and epigenome
profiling, and spatial proteomics – and performed a large-scale meta-analysis to define the mo-
lecular signatures of alpha cells during aging in non-diabetic (ND) donors and to determine their
relevance to T2D. At the islet proteome level, human islets exhibited hallmarks of ER stress, se-
nescence, and inflammatory activation linked to innate and adaptive signaling compartments.
Single cell transcriptomics revealed these inflammatory signatures were significantly enriched in
alpha cells and these resulted from IFN-γ-induced pro-inflammatory signaling via activation of
pro-inflammatory transcription factors (TFs) STAT1, IRF1, and NFKB1. Notably, this pro-inflam-
matory alpha cell state is linked to increased amino-acid induced glucagon release in vitro,
which may explain previous observations of hyperglucagonemia in ND humans in vivo (*2*).
Moreover, spatial proteomics confirmed this pro-inflammatory signaling in alpha cells correlates
with increased abundance of major histocompatibility complex class I (MHC-I; HLA-ABC) in al-
pha cells from aged donors, and which associated with enrichment of CD8[+] T cells within the

islet microenvironment and closer to alpha cells, specifically. In T2D, these islet CD8+ T cells
suffer a shift towards central memory and terminally differentiated CD8+ phenotypes. Similarly,
we show in old mice that this alpha cell-immune association is active, and it can be successfully
reversed by short term CR, which is sufficient to reduce alpha cell inflammation and attenuate
CD8+ T cell effector islet accumulation.

Together, our study identifies alpha cell inflammation as a previously unrecognized hall-
mark of human pancreas aging that is molecularly and functionally distinct from aging beta cells.
These findings establish a link between alpha cell pro-inflammatory programs and adaptive im-
mune activation during aging and demonstrate that CR can reverse age-associated islet im-
mune dysfunction by modulating alpha cells and to preserve systemic metabolic homeostasis.

<u>Results</u>

Aging ND islets exhibit elevated immune activity and unfolded protein response signal-
ing. Islet inflammation and fibrosis are hallmarks of human pancreas aging and diabetes, con-
tributing to beta cell dysfunction (*37-39*). To define proteomic signatures of aging, we leveraged
the HumanIslets consortium dataset comparing young (28±4 years, n=26) and old (71±3 years,
n=19) ND donors (**Figure 1A, Supplementary Table S1A-S1B**) (*40, 41*). Aging was associated
with 86 increased and 17 decreased proteins (**Supplementary Table S2**). Pathway analysis
(Reactome) revealed enrichment of innate immune signaling, alongside elevated amino acid
metabolism, mTORC1 activity, and ER stress pathways in old adults (**Figure 1B, Supplemen-**
tary Table S3). Consistent with these changes, aged islets exhibited increased markers of beta
cell immaturity (RBP4 (*42*); **Figure 1C**), senescence (CDKN1A/p21 (*43*), **Figure 1D**), ER stress
signaling (ATF6, CHOP, PERK; **Figure 1E**), and mTORC1 activation (LAMP2, LAMTOR, RAGA,
RAGC; **Figure 1F**). A robust immune signature was also evident by upregulation of antigen
presentation (HLA-A), T-cell receptor (TCR) signaling and memory proteins (CD28, CD44), and
immune-regulatory signaling (TGF-β1, TOM1, CD63; **Figure 1G**). Together, these findings rein-
force aging human islets exhibit coordinated ER stress and metabolic remodeling alongside
heightened inflammatory and immune engagement within the islet microenvironment.

Single cell transcriptomics reveal inflammatory signaling in aging human alpha cells.
Over the past decade, single-cell approaches have defined how islet gene regulation changes
during development, aging, and diabetes (*42, 44-53*). We previously showed that aging ND beta
cells exhibit chronic ER stress and loss of identity and insulin secretion (*44*), and further sup-
ported by our proteomic findings (**Figure 1**). To identify specific cell types contributing to the in-
flammatory signatures observed in aging islets, we analyzed scRNA-seq data from a uniformly
re-processed dataset created by PanKbase based on previously published studies (*54, 55*). Af-
ter quality control and cohort selection, the final dataset annotated 131,610 islet cells from 32
donors (56,851 alpha cells), stratified into young (20-40 years) and old (≥50 years) groups
(**Supplementary Figure S1A-C, Supplementary Table S4A-S4B).**

Among ND islet endocrine cell types, alpha cells exhibited the highest expression of
MHC-I genes (HLA-ABC), followed by immune, ductal, and endothelial cells (**Supplementary**
Figure S1D). Pseudobulk differential expression and gene set enrichment analysis (GSEA)
across alpha, beta, acinar, and MUC5B ductal cells identified age-dependent transcriptomic
changes for major islet and pancreas cell types (**Figure 2B, Supplementary Figure S1E, Sup-**
plementary Table S5). This approach revealed activation of type II interferon (IFN-γ) signaling
in aging alpha, beta, and acinar cells (**Figure 2C and Supplementary Figure S1F**). To decon-
volve the impact of IFN-γ within the human islet cytoarchitecture, we performed weighted analy-
sis of IFN-γ responsive genes enriched in old alpha cells and beta cells and discovered that ca-
nonical IFN-γ signaling is strongly enriched in old alpha cells (**Figure 2D, Supplementary Fig-**
ure S2A-S2B). Old alpha cells displayed canonical IFNγ signatures, including activation of JAK-

STAT (*STAT1, SOCS1, SOCS3, IRF7/9, IL7*, and *IL15*) and antigen processing pathways
(*TAP1, B2M*), whereas old beta cells display TNF-signaling pathway signatures (**Figure 2D-E,**
Supplementary Figure S2C-S2D). As expected, this analysis also found activation of ER stress
and DNA damage, and impaired oxidative phosphorylation in old beta cells (*44*). None of these
are found in old alpha cells (**Supplementary Figure S1F, S2A-S2B, Supplementary Table 6**).
Old alpha cells also increased expression of senescence-associated genes (*IGF1R, CDKN2A,*
CDKN1A/B) and of alpha cell identity markers (*ALDH1A1, TM4SF4, LOXL4)*, while displaying
reduced expression of ER stress-related genes (HERPUD1, ATF4, JUND). This data hints that
aging alpha cells may develop senescence next to old beta cells however they do not suffer
from significant impairments to identity or activation of ER stress (**Supplementary Figure S1G**)
(*56, 57*). Importantly, unlike old acinar cells, old alpha cells and beta cells did not display signifi-
cant activation of pro-apoptosis pathways (**Supplementary Figure S1H**), thus suggesting that
this pro-inflammatory environment does not lead to loss of alpha or beta cells.

Aging alpha cells also had increased expression of genes across several metabolic pro-
cesses including glucose metabolism and oxidative phosphorylation, fatty acid metabolism, and
amino acid catabolism (**Figure 2F-G and Supplementary Figure S1I**). To determine if these
transcriptional changes correlate with alterations in alpha cell glucagon secretion, we analyzed
HPAP human islet perifusion data from young and old adult donors from two different human is-
let analysis centers (UPenn and Vanderbilt), which used complementary stimulation protocols to
test for the effect of several insulin and glucagon secretagogues. This revealed that alpha cells
from older donors have increased glucagon secretion in response to stimulation with a mixed
amino acid (AA) cocktail and to de-polarization with KCL (**Figure 2H-K**). No changes were ob-
served when alpha cells were stimulated with low glucose, high glucose, and epinephrine (**Sup-**
plementary Figure S3).

Together, these results indicate that aging human alpha cells acquire senescence-asso-
ciated features, display heightened AA-dependent glucagon secretion, and an active IFN-γ sig-
naling coupled with MHC-I antigen-presentation programs.

Re-organization of gene regulatory networks activates inflammatory gene expression in
aging alpha cells. We have previously applied the transcription factor (TF) inference algorithm
pySCENIC (*58*) to show that aging beta cell stress arises from reconfigured gene regulatory net-
works (GRNs) and activation of stress TFs (*44*). Here, we used pySCENIC and graph network
analysis to characterize GRNs in aging alpha cells and identify the TFs linked to activation of
the observed inflammatory response. This approach identified TFs enriched in both age groups
(**Figure 3A**), including canonical alpha cell TFs (NKX6-1,FOXA2), and inflammation-associated
TFs that were significantly enriched in old alpha cells (IRFs, NFKB1, and STAT1) (**Figure 3A,**
Supplementary Figure S4).

We reconstructed enriched GRNs of young and old alpha cells and observed a stronger
inferred activity of pro-inflammatory TFs (IRF1, NFKB1, and STAT1 signaling) and reduced ac-
tivity of MAFB during aging (**Figure 3B**). This data suggests that although inflammation-associ-
ated TFs are found in both age groups, these TFs play a relatively small role in the overall tran-
scriptome landscape in younger alpha cells. To test this hypothesis, we performed analysis of
TF-associated regulons to reveal that inflammatory TFs do not target the same set of genes in
young versus old alpha cells, suggesting a significant re-wiring of alpha cell GRNs during aging.
This is characterized by a low Jaccard similarity index between conserved TFs in both age
groups, and this includes loss of TF-regulon similarity for alpha cell identity TFs (ARX, HNF1A,
HNF4A) as well as inflammation-associated TFs (STAT1, IRF2/7/8/9). In fact, these TFs are re-
directed towards new targets that include several genes identified as part of the pro-inflamma-
tory signaling phenotype of old alpha cells (**Figure 3C**, **Supplementary Table S7**). Moreover,
several of these downstream gene targets were pro-inflammatory TFs themselves, which

created a pro-inflammatory TF network in old alpha cells centered around NFKB1 and IRF1
(**Figure 3D**). In contrast, old alpha cells lose the link between NKX6.1, MAFB, and ISL1 that is
only observed in young cells (**Figure 3D**).

To further validate our SCENIC results, we integrated our analysis with publicly available
chromatin accessibility data from HPAP (snATAC-seq). Here, we performed differential analysis
of peak accessibility using ChromVAR (*59*) to identify TF motifs enriched in chromatin regions
significantly upregulated in old alpha cells. Next, we crossed referenced this list with our old al-
pha cell SCENIC data to identify n=269 shared TFs, including canonical alpha cell TFs (FOXA2,
HNF1A/B) and several inflammatory TFs (IRFs, KLF6, NFKB1/RELs, STAT1) (**Figure 3E**).

These data indicate that aging promotes re-wiring of alpha cell GRNs to activate inter-
feron-stimulated NFKB1, IRF1, and STAT1 gene expression programs and this likely explains
the origins of pro-inflammatory activation in aging alpha cells.

Spatial analysis of aging human pancreases identifies alpha cell-T-cell interactions *in*
***situ*.** Aging ND human pancreases exhibit multiple features of chronic low-grade inflammation,
including increased fibrosis, higher incidence of acinar-to-ductal metaplasia (ADM), angiopathy,
and pancreatic adiposity (*38, 39*). To validate our proteomic and transcriptomics meta-analysis
findings *in situ*, we analyzed high-dimensional imaging mass cytometry (IMC) and CO-Detection
by indexing (CODEX) datasets from HPAP (*60, 61*). Human ND pancreas samples were strati-
fied into younger adult (22-40 years) and older adults (47-58 years) groups (**Supplementary**
Figure S5B and S6B; Supplementary Tables S8 and S9). Individual cells from each donor im-
age were segmented using standard QuPath- and Seurat-based workflows for cell identification
and spatial annotation, cell marker quantification, and unsupervised clustering (**Supplementary**
Figure S5A and S6A). This allowed us to identify and map the spatial landscape of major pan-
creatic cells including alpha cells (GCG$^+$), beta cells (C-peptide$^+$), delta cells (SST$^+$), pancreatic
polypeptide cells (PP$^+$), endothelial cells (CD31$^+$), Epithelial/Acinar cells (CK19$^+$/CA2$^+$), M1-like
macrophages (CD45$^+$CD68$^+$CD163$^-$), M2-like macrophages (CD45$^+$CD68$^+$CD163$^+$), and CD4$^+$
and CD8$^+$ T cell subpopulations (CD45$^+$CD3$^+$CD4$^+$ and CD45$^+$CD3$^+$CD8$^+$, respectively) (**Figure**
4A and 4C, Supplementary Figures S5C-G and S6C-G, Supplementary Tables S10-S11).

This framework enabled spatial mapping of all major pancreatic and immune cell types
in both IMC (**Figure 4A-B**) and CODEX (**Figure 4C-D, Supplementary Figure S7A**) platforms
to quantify immune and endocrine cell composition and their molecular phenotypes. Using this
approach, we found no differences in endocrine islet cell composition (**Supplementary Figure**
S7B) and expected age-associated decreases in beta cell identity markers PDX-1, NKX6-1, and
PAX6 (**Supplementary Figure S7C**) (*44*). This data confirms loss of beta cell identity during ag-
ing without the activation of cell death (*39, 44*). Importantly, we found that ND aging pancreases
have significant expansion of CD45$^+$CD3$^+$ T cells predominantly driven by accumulation of CD8$^+$
T-cells that can be found inside the islet (**Figure 4B, 4D-E, and Supplementary Figure S7D**
and S8A). No age-associated differences were detected for CD4$^+$ T cells or in total macrophage
density (**Figure 4F-G**).

To further investigate the organization of endocrine-immune interactions underlying
CD8$^+$ T-cell expansion, we utilized spatial coordinates to identify nearest-neighbor cell endocrine
beta and alpha cells within a 50µm radius of CD8 T cells, as established previously by other
groups (*62-64*). This revealed that older pancreases have a higher fraction of CD8$^+$ T cells in the
vicinity or in direct contact with alpha cells (**Figure 4E and 4H, Supplementary Figure S8A**),
which was not observed for beta cells (**Figure 4I**).

During an inflammatory event, CD8$^+$ T cells engage neighboring cells by interaction be-
tween the T cell receptor (TCR) and major histocompatibility complex class 1 (MHC-1) mole-
cules on the interacting cell. Accordingly, the increased spatial proximity between alpha cells

and CD8[+] T cells correlated with a significant increase in the expression of MHC-I complex HLA-
ABC molecules in alpha cells (**Figure 4J**). In fact, older alpha cells express ~5x more HLA-ABC
than neighboring beta cells, and both cell types express near background levels of MHC-II HLA-
DR molecules (**Figure 4L**). Because alpha cell mass was unchanged with aging, this suggests
that despite significant interactions between alpha cells and CD8[+] T cells, aging is not associ-
ated with alpha cell loss (as indicated by the single-cell transcriptomics data (**Supplementary
Figure S1H**).

Together, these findings demonstrate that human islet and pancreatic aging landscapes
are defined by increased pro-inflammatory signaling from alpha cells that correlates with ele-
vated CD8[+] T-cell density *in situ*. Given that MHC molecules present antigenic peptides to T
cells via TCR engagement, these results suggest that alpha cell-intrinsic inflammatory pathways
may orchestrate immune activation and islet inflammation during aging.

**T2D expands pancreatic CD8[+] T cells that shift towards terminal effector-memory pheno-
types.** To determine how these pancreas and immune aging signatures overlap with T2D patho-
genesis, we analyzed CODEX data from T2D donors profiled through HPAP (**Supplementary
Table S12**). Cohorts were matched for age and BMI; however, T2D donors displayed expected
impairments in glycemia, evidenced by higher HbA1c and reduced C-peptide (**Figure 5A**). Inte-
gration of 942,724 single-cell profiles identified and characterized endocrine (99,744 cells) and
immune (195,140 cells) compartments (**Figure 5B**). Our endocrine cell clustering approach
identified no changes to major islet cell populations and highlighted two distinct beta cell popula-
tions: a CPEP[+]CHGA[+] population and a CPEP[+]CHGA[+]COL4A1[+]COL6[+] population that was en-
riched in T2D (**Figure 5C**). These cells exhibited higher interaction with fibrotic extra-cellular
matrix (ECM) and thus were detected as being "fibrotic" by our expansion-based cell segmenta-
tion algorithms that captured ECM signals from neighboring regions (**Supplementary Figure
S8A**). Importantly, T2D beta cells were marked by lower CPEP, CHGA, NKX6-1 expression
(**Figure 5D**), while alpha cells had attenuated SOX9 (**Figure 5E**), consistent with compromised
endocrine cell identity during T2D (*65*). Within immune compartments, we observed no changes
in macrophages densities (**Figure 5F**), and a significant ~2-fold increase in pancreas CD8[+] T-
cell density (**Figure 5G**). Together, these findings demonstrate that T2D amplifies CD8[+] T-cell
density to mirror patterns observed during aging in ND pancreases.

To resolve whether the increased adaptive immune phenotype observed in both aging
and T2D pancreases was localized within islets, we analyzed HPAP cytometry by time-of-flight
(CyTOF) data from isolated islets of young (26±8 years, n=13), old (54±5 years, n=14), and T2D
(50±8 years, n=13) human donors (**Figure 5H, Supplementary Table S13-14**). Both aging and
T2D presented greater CD45[+] live immune cell abundance compared with young adults, corre-
sponding with higher frequencies of islet CD8[+] T cells within the CD45[+] immune compartment
(**Figure 5I-J, Supplementary Figure S9**). CD8[+] T cells differentiate along a spectrum from na-
ïve (CD8$_{naïve}$) to CD8$_{CM}$ and CD8$_{TEM}$ states in response to TCR activation by antigen and cyto-
kine stimulation. In T2D islets, we observed a reduction in CD45RA[+]CD27[+] CD8$_{naïve}$ T cells com-
pared to young adults, whereas aging increased CD45RA[-]CD27[+] CD8$_{CM}$ T cells (**Figure 5K-L**).
T2D islets further shifted CD8[+] T-cell populations towards a higher CD45RA[+]CD27[-] CD8$_{TEM}$ phe-
notype, resulting in elevated effector-to-central memory ratio that was not observed in aging ND
samples (**Figure 5L-M**).

Together, these data show that T2D intensifies pancreas and islet CD8[+] T-cell infiltration
patterns found during aging to drive their progression towards terminal effector-memory states.

**Calorie restriction improves glucose tolerance and dampens alpha cell inflammation in
mice.** CR is a dietary intervention approach that can normalize glycemia during prediabetes and
T2D, likely by delaying the onset of workload stress and aging hallmarks in beta cells (*32, 66-*

69). We therefore tested whether CR could reverse aging signatures in alpha and beta cells *in*
vivo. To achieve this, 72-week-old C57BL/6J male mice were subjected to a 2-month, 20% CR
feeding regimen, while age-matched controls remained on ad-libitum (AL) diet (**Figure 6A**). As
expected, CR mice consumed 80% of total kcal intake compared to AL-fed controls (**Supple-**
mentary Figure S10A-B). This intervention induced a steady decline in body weight over 2
months, resulting in ~12% weight loss (**Figure 6B, Supplementary Figure S10C-D**) predomi-
nantly due to a loss in fat mass (**Supplementary Figure S10E-F**). Of note, CR mice also expe-
rienced reductions in liver and epididymal white adipose tissue (eWAT) mass due to reduced li-
pid droplet size (**Supplementary Figure S10G-J**).

Metabolic phenotyping and behavioral analysis revealed comparable overall energy ex-
penditure and oxygen consumption between AL and CR mice (**Supplementary Figure S10K-**
N). However, carbon dioxide production and carbohydrate oxidation were higher in CR mice
during the dark period when food was available (**Supplementary Figure S10O-R**), indicating a
higher preference for carbohydrate metabolism in CR mice during active feeding periods and a
rapid shift towards fatty acid metabolism during inactivity and food withdrawal. To assess how
CR influences glucose homeostasis in aged mice, we performed mixed meal tolerance tests
(MTTs; 2 g glucose/kg body weight). Baseline MTTs were conducted at 72 weeks, prior to CR.
After 2 months of CR, old mice exhibited significantly improved glucose tolerance at 80 weeks,
attributed to enhanced insulin sensitivity (**Figure 6C**). This improvement was accompanied by
~50% reductions in fasting and glucose-stimulated beta cell insulin secretion (**Supplementary**
Figure S10S). This indicates that CR preserves aging beta cell function by reducing the relative
insulin secretory demand required to maintain normoglycemia, demonstrating that known CR-
mediated functional beta cell adaptation (*33*) remains intact in aging (**Supplementary Figure**
S10T).

We previously showed that CR modulates beta cell heterogeneity by activating pro-lon-
gevity and pro-homeostasis transcriptional mechanisms that enhance beta cell protein and or-
ganelle quality-control mechanisms (*33*). This transcriptional response is driven by the activation
of TF-directed GRNs that increase the expression of beta cell identity and longevity genes (*33*).
Therefore, to determine whether CR elicits similar adaptative programs to old beta cells during
CR, we performed single nucleus RNA sequencing (snRNA-seq, 10X genomics) on islet nuclei
isolated from AL and CR-fed mice (**Figure 6A**). A total of 32,380 cells were sequenced
(AL=17,344, CR=15,306) to identify all major islet cell types, including beta cells (*Ins1*, *Ins2*, and
Ero1b), alpha cells (*Gcg*), delta cells (*Sst*), endothelial cells (*Pecam1*), and pancreatic polypep-
tide cells (*Ppy*) (**Figure 6D-E, Supplementary Figure S11A**). Approximately 65% of the se-
quenced cells were beta cells, and CR did not impact the relative composition of the heteroge-
neous beta cell population or islet cytoarchitecture composition (**Supplementary Figure S11B-**
D). Using pseudobulk analysis of beta cells followed by KEGG and STRING databases, we ob-
served a narrow range of CR-mediated effects on beta cell gene expression, limited to downreg-
ulation of few ER protein processing and folding (*Hspa1a/b/HSP70, Hspa8, Hsp90ab1,*
Dnaja1/HSP40, and Calr), cell stress (*Ypel2*, *Txnip*), and insulin synthesis (*Ins1*) genes (**Figure**
6F and Supplementary Figure S11E-J). These data indicate that CR has a very limited impact
on beta cell transcriptional programs in aged mice despite a significant downregulation of insulin
release *in vivo*.

Remarkably, CR exerted a stronger transcriptional effect on alpha cells, modulating
nearly five times more genes than beta cells (184 vs. 36 differentially expressed genes; **Figure**
6F, Supplementary Table S15). Differential gene expression and pathway enrichment analyses
of old CR alpha cell gene signatures revealed upregulation of pathways associated with gluca-
gon, glucose and fatty acid metabolism (*Gcg, Gys2, Acacb*, and *Pgc1a*/*Ppargc1a)* (**Figure 6G**).
In contrast, CR suppressed 159 genes involved in TCR signaling, antigen presentation, and

348 immune cell recruitment pathways (**Figure 6H, Supplement Figure S12**). This included genes
349 involved in cell-cell signaling via MHC class I (H2-D1, *H2-Q7*, *H2-Q6*, *H2-K1*) and some MHC
350 class II (*H2-Aa*, *H2-Ab1*) antigens (**Figure 6H-I**). Because MHC-I antigens is more prominent in
351 islet endocrine cells than MHC-II, we constructed an MHC-I module score based on heavy and
352 light chain genes (*H2-K1, H2-D1, H2-Q2, H2-Q5,H2-Q6,H2-Q7,H2-Q12*) detected in our
353 scRNA-seq data, and found the module score was reduced by CR (**Figure 6J**).

354   Taken together, these findings demonstrate that 2 months of CR reduces pro-inflamma-
355 tory signaling in aging alpha cells in mice, and it induces a state of beta cell rest that does not
356 involve significant changes to beta cell transcriptional mechanisms.

357 **CR suppresses alpha cell immune signaling to reduce islet inflammation.** To further inves-
358 tigate how CR-mediated modulation of aging alpha cell inflammatory gene expression influ-
359 ences the cellular and molecular profile of islet immune cells, we quantified the relative compo-
360 sition and transcriptional signatures immune cell types identified by snRNA-seq (**Figure 7A**).
361 This revealed a significant decrease in B cells, T cells, proliferating B cells, and antigen present-
362 ing cells (APCs) in CR mouse islets, whereas the relative density and transcriptional profile of
363 islet-resident macrophages were unchanged (**Figure 7B**). Old CR mouse islets had a marked
364 reduction in islet T cells, including a near total ablation of cytotoxic CD8[+] effector T cells (**Figure**
365 **7C-D**). In addition, the remaining CD8[+] T cells in CR islets had reduced expression of effector
366 markers granzyme K (*Gzmk*), IL-2 receptor (Il2ra/*Cd25*), Ccr7, and *Txk*; all markers of activated
367 CD8[+] T cells (**Figure 7E**). We validated these changes *in situ* using immunohistochemistry and
368 confocal microscopy of AL and CR old pancreases, followed by spatial mapping of alpha cells,
369 beta cells, and CD3[+] T cells (**Figure 7F**). This approach revealed a significant reduction in T-cell
370 density (**Figure 7G**), and increased distance between T cells and alpha cells (**Figure 7H**). The
371 proportion of T cells in proximity (within 6μm) to alpha cells trended higher (p=0.07), whereas
372 this enrichment was not observed for beta cells (**Supplementary Figure S13A-B**). Total macro-
373 phage content in proximity to islets was not altered by CR (**Supplementary Figure 13C-E**), and
374 the relative proportions of pro-inflammatory (M1-like) and anti-inflammatory (M2-like) macro-
375 phages were likewise unchanged (**Supplementary Figure S13F-K**). These data support recent
376 findings of immune cell infiltration in the aging mouse islet (*70*) and demonstrate that CR modu-
377 lates the recruitment of CD8[+] T cells to the islet by reducing their preferential proximity to old al-
378 pha cells.

379   To identify communication pathways linking aged alpha cells with CD8[+] T cells and their
380 modulation by CR, we performed receptor-ligand inference analysis using CellChat (*71, 72*) to
381 reconstruct cell-cell communication networks in islets from AL and CR mice (*73*). This analysis
382 revealed a downregulation of intercellular signaling between islet immune cells and endocrine
383 cell types, including alpha, beta, and delta cells (**Supplementary Figure S14A**). This was un-
384 derscored by a significant decrease in CD45-based interactions between old alpha cells and T-
385 and B-cells in CR mice (**Figure 7I**), accompanied by reduced complement signaling, particularly
386 toward T cells (**Figure 7J**). Moreover, CR attenuated T cell co-stimulation, as evidenced by re-
387 duced CD80 signaling from APCs and reduced CD86 signaling from B cells to T cells (**Supple-**
388 **mentary Figure S14B–C**). Chemokine signaling via CCL and CXCL pathways towards T cells
389 was similarly reduced under CR (**Supplementary Figure S14D-E**).

390   Overall, these results indicate that CR profoundly remodels the aging islet immune envi-
391 ronment by suppressing alpha cell inflammatory and MHC-I signaling, thereby reducing CD8[+] T
392 cell recruitment and cytotoxicity, and diminishing intercellular communication between alpha
393 cells and adaptive immune populations.

394 **<u>Discussion</u>**

Aging is associated with an active and low-grade inflammatory process in different tis-
sues, where inflammation is a key signature of several age-associated degenerative and meta-
bolic diseases, including Alzheimer's (*74*), and obesity and diabetes (*75*). In the pancreas, aging
is associated with increased islet macrophage infiltration and vascular fibrosis (*38*), and share
other key hallmarks of T1D and T2D islet pathophysiology such as beta cell ER stress, compro-
mised beta cell identity and function, and beta cell senescence (*15, 38, 39, 43-45, 56, 57, 76-*
82). Importantly, recent studies have implicated the activation of inflammatory responses in hu-
man alpha cells during T1D and T2D (*83, 84*).

Here, we identified the activation of immune signaling and inflammation as a key signa-
ture of aging human islets from non-diabetic individuals. This phenotype is defined by active
IFNγ signaling and MHC-I expression in old alpha cells due to activation of IFNγ-responsive TFs
IRF1, NFKB1, and STAT1. Activation of alpha cell immune signaling during aging translates into
increased physical interactions between old alpha cells with pancreas CD8$^+$ T cells, which accu-
mulate in the human islet microenvironment during aging. Strikingly, most old islet CD8$^+$ cells
have a "memory-like" phenotype (CD27$^+$), thus indicating that these cells have been exposed to
T cell antigens. While our study does not identify which antigen is presented by alpha cell MHC-
Is, a recent study where human islets were exposed to cytokine stress (e.g., IFNγ) identified
several new MHC-I antigens, many of which are enriched in aging alpha cells (i.e., NFKB1,
IRF9, TAP1, IFIT3, PSMB8, and STAT1)(*76*).

This aging alpha cell inflammatory phenotype correlates with an increase in the expres-
sion of several AA catabolism enzymes. This could explain the observed increase in amino-acid-
dependent glucagon release from isolated human islets and the elevated levels of glucagon in
old ND humans after a meal (*2*). Notably, elevated glucagon secretion after a mixed meal and
AA infusions is also observed in adults with T2D (*19-21*), thus suggesting that alpha cell inflam-
mation could also play a role during T2D pathogenesis. In fact, while we did not observe
changes in the density of islet CD8+ cells between ND and T2D islets, most T2D islet CD8$^+$
cells have an effector-like signature (CD45RA$^+$CD8$^+$). This suggests that infiltration of CD8$^+$ cells
in aging human islets occurs independently of T2D and further suggests that islet inflammation
in individuals with T2D onset during old age involves activation of alpha cell-interacting and "res-
ident" CD8$^+$ cells and broad islet inflammation signatures (*38, 80*).

In species spanning from yeast to humans, CR has been shown to extend lifespan and
health span (*32, 68, 69, 85-87*). The benefits of CR depend on improvements in glucose home-
ostasis, enhanced insulin sensitivity, and reduced metabolic demand for insulin release from
beta cells leading to beta cell rest (*33, 69, 88, 89*). In addition, CR can reduce overall inflamma-
tion signatures in different models, including humans (*32*). Our experiments with 80-week-old
mice on CR for 2 months demonstrate that the insulin-sensitizing and beta cell functional adap-
tation pathways promoted by CR are not impaired during aging and that these adaptations do
not require significant transcriptional reorganization. This is likely explained by the limited plas-
ticity of aging beta cells to respond to metabolic challenges (*90*), however the mechanisms un-
derlying this limited plasticity remain unknown. Notably, a recent study has shown that aging of
mouse endocrine islet cells is linked to immune activation and MHC-I gene expression (*82*).
While we did not observe this signature in aging beta cells and delta cells, CR acted on old al-
pha cells to significantly suppress the expression of pro-inflammatory genes, including MHC-I
signaling pathways. This CR-dependent modulation dampened CD45-mediated communication
between immune cells and alpha cells, which was sufficient to reduce the density of T cells in
aging islet periphery.

In conclusion, our findings identify the relationship between alpha cells and immune cells
within the islet of Langerhans during aging that may underlie pro-inflammatory injury and pro-
mote T2D development during old age.

Data availability: All source data and metadata, including IMC, CODEX, snRNA-seq, and
source code data are publicly available and can be downloaded from Zenodo (10.5281/ze-
nodo.17874622). Human islet single cell RNAseq uniformly reprocessed is available via PanK-
base Zenodo (https://zenodo.org/records/15596314). Raw FASTQ data from mouse islet se-
quencing data are deposited at Gene Expression Omnibus and processed data is available on
Github and Zenodo.

Acknowledgements: We acknowledge the following Vanderbilt University (VU) and Vanderbilt
University Medical Center (VUMC) core facilities: VUMC Hormone Assay & Analytical Services
Core (NIH DK020593), VU Metabolic Mouse Phenotyping Center [VMMPC (NIH DK059637;
www.vmmpc.org)], Translational Pathology Shared Resource (NCI/NIH Cancer Center Support
Grant 5P30 CA68485-19), and Vanderbilt Technologies for Advanced Genomics (VANTAGE)
core laboratory. VANTAGE is supported in part by Clinical and Translational Science Award
Grant 5UL1 RR024975-03, Vanderbilt Ingram Cancer Center Grant P30 CA68485, Vanderbilt
Vision Center Grant P30 EY08126, and National Institutes of Health/National Center for Re-
search Resources Grant G20 RR030956. This manuscript used data acquired from the data-
base (https://hpap.pmacs.upenn.edu/) of the Human Pancreas Analysis Program (HPAP;
RRID:SCR_016202)(*60, 61*). HPAP is part of a Human Islet Research Network
(RRID:SCR_014393) consortium (UC4-DK112217, U01-DK123594, UC4-DK112232, and U01-
DK123716). This work includes data and/or analyses from HumanIslets.com funded by the Ca-
nadian Institutes of Health Research, JDRF Canada, and Diabetes Canada (5-SRA-2021-1149-
S-B/TG 179092) with data from islets isolated by the Alberta Diabetes Institute IsletCore with the
support of the Human Organ Procurement and Exchange program, Trillium Gift of Life Network,
BC Transplant, Quebec Transplant, and other Canadian organ procurement organizations with
written informed donor consent as approved by the Human Research Ethics Board at the Uni-
versity of Alberta (Pro00013094). we would like to acknowledge Dr Patrick E MacDonald from
ADI – Edmonton (Canada) for critical comments and inputs during data collection and manu-
script writing stages of this project. We would also like to acknowledge NIH grant
R01AG040178.

Funding: This work was support by the National Institutes of Health (NIH) grant R01DK138141
to R.A.D. M.S. is supported by the NIH Molecular Endocrinology Training Program
(T32DK007563) and the American Diabetes Association (1-26-PDF-1109). A.C. was supported
by an American Heart Association Postdoctoral Fellowship (26POST1571444).

Author Contributions: M.S., S.S., JP.C., and R.A.D designed experiments. M.S., A.C., M.C.,
G.F., S.S., JP.C., A.H., C.E., A.P., and R.A.D. collected data and assisted with data analysis and
project management. M.S., S.S., JP.C., and R.A.D. completed data analysis. M.S. and R.A.D.
interpreted data and wrote the manuscript.

Disclosures: The authors have nothing to disclose.

REFERENCES

1. National Diabetes Statistics Report. *Centers for Disease Control and Prevention*, (2024).

2. R. Basu *et al.*, Mechanisms of the age-associated deterioration in glucose tolerance:
contribution of alterations in insulin secretion, action, and clearance. *Diabetes* **52**, 1738-
1748 (2003).

3. K. F. Petersen *et al.*, Mitochondrial dysfunction in the elderly: possible role in insulin
resistance. *Science* **300**, 1140-1142 (2003).

4. C. C. Cowie *et al.*, Prevalence of diabetes and impaired fasting glucose in adults in the U.S.
population: National Health And Nutrition Examination Survey 1999-2002. *Diabetes Care*
29, 1263-1268 (2006).

5. A. Alpert *et al.*, A clinically meaningful metric of immune age derived from high-dimensional
longitudinal monitoring. *Nat Med* **25**, 487-495 (2019).

6. C. Sportès *et al.*, Administration of rhIL-7 in humans increases in vivo TCR repertoire
diversity by preferential expansion of naive T cell subsets. *J Exp Med* **205**, 1701-1714 (2008).

7. M. L. Pekalski *et al.*, Postthymic expansion in human CD4 naive T cells defined by
expression of functional high-affinity IL-2 receptors. *J Immunol* **190**, 2554-2566 (2013).

8. X. Sun *et al.*, Longitudinal analysis reveals age-related changes in the T cell receptor
repertoire of human T cell subsets. *J Clin Invest* **132**, (2022).

9. O. Cabrera *et al.*, The unique cytoarchitecture of human pancreatic islets has implications
for islet cell function. *Proceedings of the National Academy of Sciences* **103**, 2334-2339
(2006).

10. M. Cnop *et al.*, The long lifespan and low turnover of human islet beta cells estimated by
mathematical modelling of lipofuscin accumulation. *Diabetologia* **53**, 321-330 (2010).

11. R. A. E. Drigo *et al.*, Age Mosaicism across Multiple Scales in Adult Tissues. *Cell Metabolism*
30, 343-+ (2019).

12. S. Perl *et al.*, Significant human beta-cell turnover is limited to the first three decades of life
as determined by in vivo thymidine analog incorporation and radiocarbon dating. *J Clin*
Endocrinol Metab **95**, E234-239 (2010).

13. C. López-Otín, M. A. Blasco, L. Partridge, M. Serrano, G. Kroemer, Hallmarks of aging: An
expanding universe. *Cell* **186**, 243-278 (2023).

14. S. Shrestha *et al.*, Aging compromises human islet beta cell function and identity by
decreasing transcription factor activity and inducing ER stress. *Science Advances* **8**,
eabo3932 (2022).

15. I. Sandovici *et al.*, Ageing is associated with molecular signatures of inflammation and type
2 diabetes in rat pancreatic islets. *Diabetologia* **59**, 502-511 (2016).

16. L. Li *et al.*, Defects in β-cell Ca2+ dynamics in age-induced diabetes. *Diabetes* **63**, 4100-
4114 (2014).

17. G. M. Reaven, Y. D. Chen, A. Golay, A. L. Swislocki, J. B. Jaspan, Documentation of
hyperglucagonemia throughout the day in nonobese and obese patients with noninsulin-
dependent diabetes mellitus. *J Clin Endocrinol Metab* **64**, 106-110 (1987).

18. K. Færch *et al.*, Insulin Resistance Is Accompanied by Increased Fasting Glucagon and
Delayed Glucagon Suppression in Individuals With Normal and Impaired Glucose
Regulation. *Diabetes* **65**, 3473-3481 (2016).

19. R. H. Unger, E. Aguilar-Parada, W. A. Müller, A. M. Eisentraut, Studies of pancreatic alpha
cell function in normal and diabetic subjects. *J Clin Invest* **49**, 837-848 (1970).

20. D. D'Alessio, The role of dysregulated glucagon secretion in type 2 diabetes. *Diabetes Obes*
Metab **13 Suppl 1**, 126-132 (2011).

21.   P. Raskin, I. Aydin, T. Yamamoto, R. H. Unger, Abnormal alpha cell function in human
diabetes: the response to oral protein. *Am J Med* **64**, 988-997 (1978).
22.   P. Raskin, I. Aydin, R. H. Unger, Effect of insulin on the exaggerated glucagon response to
arginine stimulation in diabetes mellitus. *Diabetes* **25**, 227-229 (1976).
23.   L. Fontana, L. Partridge, V. D. Longo, Extending healthy life span--from yeast to humans.
Science **328**, 321-326 (2010).
24.   J. A. Mattison *et al.*, Caloric restriction improves health and survival of rhesus monkeys. *Nat*
Commun **8**, 14063 (2017).
25.   C. L. Green, D. W. Lamming, L. Fontana, Molecular mechanisms of dietary restriction
promoting health and longevity. *Nat Rev Mol Cell Biol* **23**, 56-73 (2022).
26.   R. M. Anderson, D. G. Le Couteur, R. de Cabo, Caloric Restriction Research: New
Perspectives on the Biology of Aging. *J Gerontol A Biol Sci Med Sci* **73**, 1-3 (2017).
27.   K. Mao *et al.*, Late-life targeting of the IGF-1 receptor improves healthspan and lifespan in
female mice. *Nat Commun* **9**, 2394 (2018).
28.   G. Y. Liu, D. M. Sabatini, mTOR at the nexus of nutrition, growth, ageing and disease. *Nat Rev*
Mol Cell Biol **21**, 183-203 (2020).
29.   D. Il'yasova *et al.*, Effects of 2 years of caloric restriction on oxidative status assessed by
urinary F2-isoprostanes: The CALERIE 2 randomized clinical trial. *Aging Cell* **17**, (2018).
30.   L. Yang, P. Li, S. Fu, E. S. Calay, G. S. Hotamisligil, Defective hepatic autophagy in obesity
promotes ER stress and causes insulin resistance. *Cell Metab* **11**, 467-478 (2010).
31.   J. O. Pyo *et al.*, Overexpression of Atg5 in mice activates autophagy and extends lifespan.
Nat Commun **4**, 2300 (2013).
32.   O. Spadaro *et al.*, Caloric restriction in humans reveals immunometabolic regulators of
health span. *Science* **375**, 671-+ (2022).
33.   C. dos Santos *et al.*, Calorie restriction increases insulin sensitivity to promote beta cell
homeostasis and longevity in mice. *Nature Communications* **15**, 9063 (2024).
34.   L. Matta *et al.*, Chronic intermittent fasting impairs β cell maturation and function in
adolescent mice. *Cell Rep* **44**, 115225 (2025).
35.   S. Patel, Z. Yan, M. S. Remedi, Intermittent fasting protects β-cell identity and function in a
type-2 diabetes model. *Metabolism* **153**, 155813 (2024).
36.   M. R. Brown *et al.*, Time-restricted feeding prevents deleterious metabolic effects of
circadian disruption through epigenetic control of β cell function. *Sci Adv* **7**, eabg6856
(2021).
37.   L. Mateus Gonçalves, E. Pereira, J. P. Werneck de Castro, E. Bernal-Mizrachi, J. Almaça, Islet
pericytes convert into profibrotic myofibroblasts in a mouse model of islet vascular fibrosis.
Diabetologia **63**, 1564-1575 (2020).
38.   J. Almaca *et al.*, Young capillary vessels rejuvenate aged pancreatic islets. *Proceedings of*
the National Academy of Sciences of the United States of America **111**, 17612-17617
(2014).
39.   J. J. Wright *et al.*, Exocrine Pancreas in Type 1 and Type 2 Diabetes: Different Patterns of
Fibrosis, Metaplasia, Angiopathy, and Adiposity. *Diabetes* **73**, 1140-1152 (2024).
40.   J. D. Ewald *et al.*, HumanIslets.com: Improving accessibility, integration, and usability of
human research islet data. *Cell Metab* **37**, 7-11 (2025).
41.   J. Kolic *et al.*, Proteomic predictors of individualized nutrient-specific insulin secretion in
health and disease. *Cell Metab* **36**, 1619-1633.e1615 (2024).
42.   J. Camunas-Soler *et al.*, Patch-Seq Links Single-Cell Transcriptomes to Human Islet
Dysfunction in Diabetes. *Cell Metabolism* **31**, 1017-+ (2020).

43. C. Aguayo-Mazzucato *et al.*, Acceleration of beta Cell Aging Determines Diabetes and
Senolysis Improves Disease Outcomes. *Cell Metabolism* **30**, 129-+ (2019).
44. S. Shrestha *et al.*, Aging compromises human islet beta cell function and identity by
decreasing transcription factor activity and inducing ER stress. *Science Advances* **8**, 17
(2022).
45. M. Enge *et al.*, Single-Cell Analysis of Human Pancreas Reveals Transcriptional Signatures
of Aging and Somatic Mutation Patterns. *Cell* **171**, 321-330.e314 (2017).
46. S. Tritschler, F. J. Theis, H. Licked, A. Bottcher, Systematic single-cell analysis provides new
insights into heterogeneity and plasticity of the pancreas. *Molecular Metabolism* **6**, 974-990
(2017).
47. P. Augsornworawat, K. G. Maxwell, L. Velazco-Cruz, J. R. Millman, Single-Cell Transcriptome
Profiling Reveals β Cell Maturation in Stem Cell-Derived Islets after Transplantation. *Cell*
Rep **32**, 108067 (2020).
48. J. Chiou *et al.*, Single-cell chromatin accessibility identifies pancreatic islet cell type- and
state-specific regulatory programs of diabetes risk. *Nature Genetics* **53**, 455-+ (2021).
49. N. C. Bramswig *et al.*, Epigenomic plasticity enables human pancreatic α to β cell
reprogramming. *J Clin Invest* **123**, 1275-1284 (2013).
50. D. Avrahami *et al.*, Aging-Dependent Demethylation of Regulatory Elements Correlates with
Chromatin State and Improved β Cell Function. *Cell Metabolism* **22**, 619-632 (2015).
51. D. Avrahami *et al.*, Single-cell transcriptomics of human islet ontogeny defines the
molecular basis of β-cell dedifferentiation in T2D. *Mol Metab* **42**, 101057 (2020).
52. C. Su *et al.*, 3D chromatin maps of the human pancreas reveal lineage-specific regulatory
architecture of T2D risk. *Cell Metabolism* **34**, 1394-+ (2022).
53. H. E. Arda *et al.*, Age-Dependent Pancreatic Gene Regulation Reveals Mechanisms
Governing Human beta Cell Function. *Cell Metabolism* **23**, 909-920 (2016).
54. M. Fasolino *et al.*, Single-cell multi-omics analysis of human pancreatic islets reveals novel
cellular states in type 1 diabetes. *Nature Metabolism* **4**, 284-+ (2022).
55. A. R. Patil *et al.*, Single-cell expression profiling of islets generated by the Human Pancreas
Analysis Program. *Nat Metab* **5**, 713-715 (2023).
56. C. Aguayo-Mazzucato *et al.*, beta Cell Aging Markers Have Heterogeneous Distribution and
Are Induced by Insulin Resistance. *Cell Metabolism* **25**, 898-+ (2017).
57. P. J. Thompson *et al.*, Targeted Elimination of Senescent Beta Cells Prevents Type 1
Diabetes. *Cell Metab* **29**, 1045-1060.e1010 (2019).
58. B. van de Sande *et al.*, A scalable SCENIC workflow for single-cell gene regulatory network
analysis. *Nature Protocols* **15**, 2247-2276 (2020).
59. A. N. Schep, B. Wu, J. D. Buenrostro, W. J. Greenleaf, chromVAR: inferring transcription-
factor-associated accessibility from single-cell epigenomic data. *Nat Methods* **14**, 975-978
(2017).
60. K. H. Kaestner, A. C. Powers, A. Naji, M. A. Atkinson, NIH Initiative to Improve Understanding
of the Pancreas, Islet, and Autoimmunity in Type 1 Diabetes: The Human Pancreas Analysis
Program (HPAP). *Diabetes* **68**, 1394-1402 (2019).
61. S. N. Shapira, A. Naji, M. A. Atkinson, A. C. Powers, K. H. Kaestner, Understanding islet
dysfunction in type 2 diabetes through multidimensional pancreatic phenotyping: The
Human Pancreas Analysis Program. *Cell Metab* **34**, 1906-1913 (2022).
62. É. Korpos *et al.*, Identification and characterisation of tertiary lymphoid organs in human
type 1 diabetes. *Diabetologia* **64**, 1626-1641 (2021).
63. M. H. Wu *et al.*, Single-cell analysis of the human pancreas in type 2 diabetes using multi-
spectral imaging mass cytometry. *Cell Reports* **37**, 15 (2021).

64.   Y. J. Wang *et al.*, Multiplexed In Situ Imaging Mass Cytometry Analysis of the Human
626   Endocrine Pancreas and Immune System in Type 1 Diabetes. *Cell Metab* **29**, 769-783.e764
627   (2019).
628   65.   C. Talchai, S. H. Xuan, H. V. Lin, L. Sussel, D. Accili, Pancreatic beta Cell Dedifferentiation as
629   a Mechanism of Diabetic beta Cell Failure. *Cell* **150**, 1223-1234 (2012).
630   66.   C. Dos Santos *et al.*, Caloric restriction promotes beta cell longevity and delays aging and
631   senescence by enhancing cell identity and homeostasis mechanisms. *Res Sq*, (2023).
632   67.   E. Duregon *et al.*, Prolonged fasting times reap greater geroprotective effects when
633   combined with caloric restriction in adult female mice. *Cell Metab* **35**, 1179-1194.e1175
634   (2023).
635   68.   S. J. Mitchell *et al.*, Effects of Sex, Strain, and Energy Intake on Hallmarks of Aging in Mice.
636   *Cell Metabolism* **23**, 1093-1112 (2016).
637   69.   H. H. Pak *et al.*, Fasting drives the metabolic, molecular and geroprotective effects of a
638   calorie-restricted diet in mice. *Nature Metabolism* **3**, 1327-+ (2021).
639   70.   S. Jin *et al.*, Inference and analysis of cell-cell communication using CellChat. *Nat Commun*
640   **12**, 1088 (2021).
641   71.   A. Coomans de Brachène *et al.*, Exercise as a non-pharmacological intervention to protect
642   pancreatic beta cells in individuals with type 1 and type 2 diabetes. *Diabetologia* **66**, 450-
643   460 (2023).
644   72.   P. Carapeto *et al.*, Exercise activates AMPK in mouse and human pancreatic islets to
645   decrease senescence. *Nature Metabolism*, (2024).
646   73.   H. Al Jobori *et al.*, Empagliflozin Treatment Is Associated With Improved β-Cell Function in
647   Type 2 Diabetes Mellitus. *J Clin Endocrinol Metab* **103**, 1402-1407 (2018).
648   74.   C. M. Kellogg *et al.*, Microglial MHC-I induction with aging and Alzheimer's is conserved in
649   mouse models and humans. *Geroscience* **45**, 3019-3043 (2023).
650   75.   S. SantaCruz-Calvo *et al.*, Adaptive immune cells shape obesity-associated type 2 diabetes
651   mellitus and less prominent comorbidities. *Nat Rev Endocrinol* **18**, 23-42 (2022).
652   76.   P. P. Nanaware *et al.*, The antigen presentation landscape of cytokine-stressed human
653   pancreatic islets. *Cell Rep* **44**, 115927 (2025).
654   77.   L. Marselli *et al.*, beta-Cell inflammation in human type 2 diabetes and the role of
655   autophagy. *Diabetes Obesity & Metabolism* **15**, 130-136 (2013).
656   78.   J. Jelleschitz *et al.*, Insulitis and aging: Immune cell dynamics in Langerhans islets. *Redox
657   Biol* **82**, 103587 (2025).
658   79.   M. De Burghgrave *et al.*, Pancreatic Islet Cells Response to IFNγ Relies on Their Spatial
659   Location within an Islet. *Cells* **12**, (2022).
660   80.   D. Avrahami *et al.*, Single-cell transcriptomics of human islet ontogeny defines the
661   molecular basis of beta-cell dedifferentiation in T2D. *Molecular Metabolism* **42**, 14 (2020).
662   81.   E. Horwitz *et al.*, β-Cell DNA Damage Response Promotes Islet Inflammation in Type 1
663   Diabetes. *Diabetes* **67**, 2305-2318 (2018).
664   82.   W. Staels *et al.*, Comprehensive alpha, beta, and delta cell transcriptomics reveal an
665   association of cellular aging with MHC class I upregulation. *Mol Metab* **87**, 101990 (2024).
666   83.   T. Dos Santos *et al.*, Altered immune and metabolic molecular pathways drive islet cell
667   dysfunction in human type 1 diabetes. *J Clin Invest*, (2025).
668   84.   X. Q. Dai *et al.*, Heterogenous impairment of alpha cell function in type 2 diabetes is linked
669   to cell maturation state. *Cell Metabolism* **34**, 256-+ (2022).
670   85.   V. D. Longo, R. M. Anderson, Nutrition, longevity and disease: From molecular mechanisms
671   to interventions. *Cell* **185**, 1455-1470 (2022).

86.   J. A. Mattison *et al.*, Impact of caloric restriction on health and survival in rhesus monkeys
from the NIA study. *Nature* **489**, 318-321 (2012).
87.   A. E. Civitarese *et al.*, Calorie restriction increases muscle mitochondrial biogenesis in
healthy humans. *PLoS Med* **4**, e76 (2007).
88.   M. E. C. da Amaral *et al.*, Caloric restriction recovers impaired beta-cell-beta-cell gap
junction coupling, calcium oscillation coordination, and insulin secretion in prediabetic
mice. *American Journal of Physiology-Endocrinology and Metabolism* **319**, E709-E720
(2020).
89.   C. J. Sheng *et al.*, Reversibility of beta-Cell-Specific Transcript Factors Expression by Long-
Term Caloric Restriction in db/db Mouse. *Journal of Diabetes Research* **2016**, 11 (2016).
90.   S. I. Tschen, S. Dhawan, T. Gurlo, A. Bhushan, Age-dependent decline in beta-cell
proliferation restricts the capacity of beta-cell regeneration in mice. *Diabetes* **58**, 1312-1320
(2009).
91.   M. E. Ritchie *et al.*, limma powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic acids research* **43**, e47-e47 (2015).
92.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
93.   Y. Hao *et al.*, Dictionary learning for integrative, multimodal and scalable single-cell
analysis. *Nat Biotechnol* **42**, 293-304 (2024).
94.   Y. J. Wang *et al.*, Single-Cell Mass Cytometry Analysis of the Human Endocrine Pancreas.
Cell Metabolism **24**, 616-626 (2016).
95.   R. J. Colman *et al.*, Caloric restriction delays disease onset and mortality in rhesus
monkeys. *Science* **325**, 201-204 (2009).

Figure Legends

Figure 1 - Aging human islets have an inflammatory activation phenotype combined with
mTORC1 signaling and elevated unfolded protein response stress. (A) Islets from young
(28±4 years, n=26), and old (71±3 years, n=19) ND donors from the HumanIslets consortium
were compared by proteomics meta-analysis. **(B)** Reactome pathways present from significantly
abundant proteins between young and old ND donors. **(C-G)** Protein expression levels for beta
cell immaturity marker, RBP4 (C); senescence, CDKN1A/p21 (D); unfolded protein response,
CHOP (DDIT3), ATF6, PERK (EIF2AK3) (E); mTORC1 activation, LAMP2, LAMTOR, RAGA,
RAGC (F); and immune regulatory components, HLA-A, TGF-β1, TOM1, CD63, CD44, and
CD28, respectively (G). HLA, human leukocyte antigen. *p<0.05, young vs. old donors. Error
bars are mean ± SD.

Figure 2 - Aging increases human alpha cell interferon signaling and metabolic signa-
tures. (A) Cell type UMAP of integrated scRNA-seq data obtained from HPAP database. **(B)**
Heatmap with hierarchical clustering showing top 25 genes differentially regulated by age in hu-
man alpha cells. **(C)** GSEA hallmarks for gene sets associated with the Hallmark Interferon
GAMMA Response (M5913). **(D)** Relative gene expression levels of genes in the Hallmark Inter-
feron GAMMA Response (M5913) collection in alpha cells versus beta cells. Asterisks mark
genes significantly upregulated in either cell type. **(E)** Weighted KEGG pathway analysis
demonstrating the enrichment of IFN-γ associated genes in old alpha cells versus old beta cells.
(F-G) Significantly expressed Hallmark genes for metabolic processes including Glycolysis
(M5937; G) and Amino Acid Metabolic Process (GO:0006520), respectively. **(H-K)** Isolated hu-
man islet perifusion assay including stimulation with amino-acids, glucose, IBMX, and KCL. Dot-
ted line boxes and represented quantification (**I**, **J**, and **K**) highlight specific stimuli that provoked
significant glucagon secretion from old islets versus young islets. P-values are shown. Data are
mean ± SD with boxplots representing mean ± IQR. ND, non-diabetic. HPAP, Human Pancreas
Analysis Program; GSEA, gene set enrichment analysis.

Figure 3 - Re-wiring of gene regulatory networks activates an interferon-stimulated in-
flammatory phenotype in aging alpha cells. (A) Regulon specificity scores (RSS) ratios be-
tween old and young alpha cells, with enrichment defined as the top 10% of RSS ratios within
each group. **(B)** Annotated Gene Regulatory Network (GRN) maps of young (top) and old (bot-
tom) human alpha cells. Transcription factors (TFs) are represented by white hexagons and in-
ferred TF target genes are shown in solid grey circles. Only the top 15% TF regulons after 100X
pySCENIC inference simulations are shown. TF hexagon size represents the TF importance
metric within the GRN structure, and TF-target gene nodes are connected by line edges colored
using a *viridis* hue to represent predicted connection strength. Shaded blue boxes are anno-
tated with pathway enrichment terms associated with their respective regulon network. **(C)** Venn
diagrams representing unique and shared genes found in the regulons associated with IRF1,
NFKB1, STAT1, AND MAFB TFs. **(D)** Regulatory network map displaying age-dependent and
age-independent regulatory connections between inflammatory TFs and canonical alpha cell
identity TFs. In this map, TFs are represented by solid grey circles and regulatory relationships
are shown by arrows. Green arrows, connections found in both young and old alpha cells; dark
blue and beige arrows represent connections found uniquely in old and young alpha cells, re-
spectively. **(E)** Venn diagram illustrating the overlap between SCENIC and chromVAR associ-
ated TFs detected in old alpha cells.

Figure 4 - Aging increases CD8[+] islet infiltration via alpha cell MHC-I signaling. (A) Imag-
ing Mass Cytometry (IMC) UMAPs displaying subset pancreas cells from all donors, highlighting
the islet-endocrine cells (top) and immune cells (bottom). **(B)** Representative IMC images and
spatial reconstructions of islet-endocrine (top) and islet-immune (bottom) fractions. Scale bar =
100μm. **(C)** UMAPs of pancreas cell subpopulations, highlighting endocrine cells (top) and

immune subsets (bottom). **(D)** Representative CODEX images and spatial reconstructions for
islet immune subpopulations. Scale bar = 2000μm. **(E)** Representative islet for an aged donor,
displaying C-peptide (yellow), glucagon (green), CD45 (white) and CD8 (magenta), with inset
illustrating alpha-CD8 T cell interaction. Scale bar; image = 100μm, inset = 10μm. **(F-G)** Im-
mune cell proportions within the total pancreatic immune pool, collected by both IMC (F) and
CODEX (H) imaging platforms. **(H)** Density plots illustrating networking distance between CD8$^+$
T cells distance and alpha cells, median CD8$^+$-alpha cell distance, and the percentage of alpha
cells localized within ≤50μm of CD8$^+$ T cells. **(I)** Density plots showing network distance between
CD8$^+$ T cells and beta cells, median CD8$^+$-beta cell distance, and the percentage of beta cells
localized within ≤50μm of CD8$^+$ T cells. **(J)** Representative IMC images illustration alpha cells
(GCG, red) beta cells (C-peptide, white), and scaled HLA-ABC expression. Scale bar = 100μm.
(K-L). Quantification of HLA-ABC (K) and HLA-DR (L) intensity from segmented alpha cells and
beta cells. IMC, imaging mass cytometry; CyTOF, Cytometry by time-of-flight; ROI, region of in-
terest. Welch's two-sample *t-test* was used to compare young and old donors in panels F-G, H-I,
and K-L. Data are mean ± SEM (panels F-I), median ± SEM (K-L).

Figure 5 - T2D expands pancreatic CD8$^+$ T-cell populations and shifts islet CD8$^+$ T-cell
states from towards terminal effector-memory phenotypes. (A) Clinical characteristics ND
and T2D adults included for HPAP CODEX imaging. **(B)** UMAP of integrated pancreas samples
(942,724 cells) were segmented into immune cells (195,140 cells) and islet endocrine (99,744
cells) populations. **(C)** Identified islet endocrine cell populations, including alpha (GCG$^+$), beta
(C-PEP$^+$), "fibrotic" beta cells (C-PEP$^+$, COL4A1$^+$, COL6$^+$), delta (SST$^+$), pancreatic polypeptide
(PP$^+$), and epsilon (ghrelin$^+$) cells. **(D)** Scaled beta cell expression of C-PEP, CHGA, and NKX6-
1. **(E)** Scaled alpha cell expression of SOX9. **(F)** Proportions of major pancreatic immune sub-
sets, including M2-like macrophages (CD163$^+$CD68$^+$), M1-like macrophages (CD68$^+$CD163$^-$),
granulocytes/monocytes (CD66b$^+$), CD8$^+$ T cells (CD3$^+$CD8$^+$), and CD4$^+$ T cells (CD3$^+$CD4$^+$).
(G) Immune cell density expressed as cells per mm^2. **(H)** CyTOF analysis of islets isolated from
young ND (n=12), old ND(n=15), and T2D (n=13) donors. **(I)** Representative gating of viable
CD45$^+$ immune cells (left) and CD3$^+$CD8$^+$ T cells (right), sourced from HPAP-164. **(J)** Proportion
of CD45$^+$ immune cells among viable islet cells (left) and CD3$^+$CD8$^+$ within the CD45$^+$ compart-
ment (right). **(K)** Representative two-parameter (CD27 x CD45RA) plot illustrating CD8$^+$ T-cell
differentiation states; (1) naïve CD27$^+$CD45RA$^+$, (2) central memory (CD27$^+$CD45RA$^-$), 3) and
terminal effector memory (CD27$^-$CD45RA$^+$). **(L)** Distribution of CD8$^+$ T cells across naïve
CD27$^+$CD45RA$^+$; CD8$_{naïve}$), central memory (CD27$^+$CD45RA$^-$; CD8$_{CM}$), and terminal effector
memory (CD27$^-$CD45RA$^+$; CD8$_{TEM}$) phenotypes. **(M)** CD8$^+$ T$_{EM}$/T$_{CM}$ ratio, representing the rela-
tive expansion of terminal effector-memory CD8$^+$ T cells compared with the central memory
pool. Welch's two sample *t-test* was used for comparisons between pool ND and T2D donors in
panels A and C-G. One-way *ANOVA* compared young, old and T2D donors in panels J and L-M.
DC, dendritic cell; cDC1, conventional cell; PP, pancreatic polypeptide; NK, natural killer. *p<
0.05, **p<0.01 for age (old vs. young) and ND vs. T2D groups. Data are mean ± SEM.

Figure 6 - Late-life CR improves glucose tolerance while downregulating alpha cell TCR
activation and antigen presentation and processing. (A) 72-week-old C57Bl/6J mice were
fed ad-libitum (AL, control) or subjected to 20% calorie restriction (CR) for 2 months. Islets were
collected at 80 weeks and processed for IHC and snRNA-seq. **(B)** Percent change in body
weight relative to baseline. **(C)** MTTs were performed at baseline (72 weeks) and after the inter-
vention (80 weeks). AUC is presented for each test before and after dietary intervention. **(D)**
UMAP of 32,380 islet cells. **(E)** Selected marker genes for endocrine islet subpopulations by
scaled gene expression and percent cell expression. **(F)** Number of DEGs affected by CR in en-
docrine/endothelial cell populations. **(G)** KEGG pathways significantly modified by CR in alpha
cells. **(H)** Protein-protein interaction networks predicted from genes downregulated by CR in al-
pha cells, and subset by 'leukocyte activation' subcluster in red. **(I)** Scaled single-cell expression

of MHC class I in alpha cells. **(J)** Scaled expression UMAP and box plot showing relative MHC-I
module scores, derived from *H2-K1, H2-D1, H2-Q2, H2-Q5, H2-Q6, H2-Q7, and H2-Q12* genes,
in AL and CR alpha cells. AUC, area under curve; IHC, immunohistochemistry; MTT, mixed meal
tolerance test; snRNA-seq, single nucleus RNA sequencing. #p< 0.05 for dietary groups (AL vs.
CR); ****p<0.0001 for intervention (Pre- vs. Post-CR).

Figure 7 - Late-life CR attenuates islet adaptive CD8$^+$ effector phenotypes and reduces al-
pha cell-mediated T-cell recruitment. (A) UMAP of subset immune cell populations (B cells, T
cells, macrophages, proliferating B cells, and APCs [dendritic cells] from AL and CR groups. **(B)**
Proportion of immune cell populations relative to total islet cells. **(C)** UMAP of T-cell populations
from AL and CR groups. **(D)** Proportion of total T cells; * denotes effector CD8$^+$ T cells. **(E)** Ex-
pression of Th1-associated genes: *Gzmk, Cd25/Il2ra, CCr7,* and *Txk*. **(F)** Representative IHC
staining for alpha cells (glucagon, red), beta cells (insulin, green) and T cells (CD3, purple).
Scale bar = 100µm. **(G)** Pancreatic T-cell density (CD3$^+$ T-cell counts per mm^2). **(H)** T-cell-endo-
crine cell networks, quantifying the average distance of CD3$^+$ T cells to alpha cells (left), and
beta cells (right). **(I-J)** Intercellular communication networks showing inferred CD45 signaling
and complement signaling interactions, respectively.

A Young Old

Nondiabetic Human Donors
Young (n=26) Old (n=19)
20-35 years 60-85 years

Islet Isolation Proteomics

Age
years

Young Old

B Fold enrichment
Count 5 10 15 20

Enriched young ← → Enriched old

Innate Immune System
Platelet degranulation
TP53 Regulates Metabolic Genes
Cellular responses to stimuli
Immune System
Cellular responses to stress
mTORC1-mediated signalling
Energy dependent regulation of mTOR by LKB1-AMPK
MTOR signalling

Old

Translation
Metabolism of proteins
SRP-dependent cotranslational protein targeting to membrane

Young

$-\log_{10}(FDR)$

C Beta cell immaturity
RBP4
scaled intensity
Young Old

D Senescence
CDKN1A/p21
scaled intensity
Young Old

E Unfolded protein response
DDIT3/CHOP ATF6 EIF2AK3/PERK
scaled intensity
Young Old Young Old Young Old

F mTORC1 activation
LAMP2 LAMTOR
scaled intensity
RAGA RAGC
scaled intensity
Young Old Young Old

G Immune
HLA-A TGFβ1 TOM1 CD63 CD44 CD28
scaled intensity
Young Old Young Old Young Old Young Old Young Old Young Old

Imaging Mass Cytometry

A Endocrine cells

Beta, Alpha, Delta, PP/Ghrelin, Endothelial, Mix

Immune cells

Mθ, Monocyte, CD4 T, CD8 T, B Cell, NK

B DNA CPEP GCG SST

Young HPAP-080 / Old HPAP-066

DNA CD163 CD14 CD4 CD8 CD20 ISLET

Young HPAP-080 / Old HPAP-066

CODEX

C Endocrine cells

Beta, Alpha, Delta, Mix, PP, Ghrelin, Stromal

Immune cells

M2 Mθ, M1 Mθ, Mono./Granulo., cDC1, DC, CD4 T, CD8 T, B cell

D DNA CD163 CD66b CD4 CD8 CD19 ISLET

Young HPAP-013 / Old HPAP-174

E C-PEP GCG CD45 CD8

F Imaging Mass Cytometry

Mono., Mθ, CD8 T, NK, CD4 T, B Cell

CD8+ T cells per mm2

CD4+ T cells per mm2

young / old

G CODEX

M1 Mθ, M2 Mθ, Gran./Mono., CD8 T, CD4 T, B Cell, DC, cDC1

%CD8+ of CD3+

%CD4+ of CD3+

young / old

H Alpha cell distance to CD8+ T cells

Density / distance (µm)

median distance (µm) p=0.10

percent <50um

% Alpha Cells

Young / Old

I Beta cell distance to CD8+ T cells

Density / distance (µm)

median distance (um)

Percent <50µm

% Beta Cells

Young / Old

J Islet / HLA-ABC

DAPI CPEP GCG

Young HPAP-019

Old HPAP-066

Inset

200 µm

K HLA-ABC

Alpha / Beta

Intensity

young / old

L HLA-DR

Alpha / Beta

Intensity

young / old

A Immune cells

Macrophages
T-cell
APC
prol_B-cell
B-cell

UMAP 2
UMAP 1

B Proportion of islet cells

% islet cells

- B cell
- T cell
- Macrophage
- Prol. B cell
- Dendritic cell

10%
5%
0%

AL   CR

C T cells

UMAP 2
UMAP 1

D Proportion of T cells

% T cells

100%
75%
50%
25%
0%

AL   CR

- Exhausted CD 4/8+
- Effector CD8+
- Naïve/memory CD4+
- Effector CD8+
- Stromal/Other
- Effector CD4+

E T cell cytotoxicity

Gzmk
AL   CR

Il2ra
AL   CR

Ccr7
AL   CR

Txk
AL   CR

1
0.5
0

F DAPI GCG Insulin CD3

AL
Inset

CR
Inset

100 μm

G T cell density

CD3+ per mm²

*

AL   CR

H CD3+ T cell distances

T cell - Alpha cell

CD3+ distance to alpha cells (μm)

100
50
0

*

AL   CR

T cell - Beta cell

CD3+ distance to beta cells (μm)

100
50
0

ns

AL   CR

I CD45 Signaling

AL   CR

J Complement Signaling

AL   CR