Calorie restriction suppresses aging alpha cell pro-inflammatory signaling.

Michael Schleh[1], Amanda Cambraia[1], Melanie Cutler[1], Gabriel Ferguson[1], Rafael Arrojo e Drigo[1,2,3,4]

[1] Department of Molecular Physiology and Biophysics, Vanderbilt University School of Medicine, Nashville, TN
[2] Vanderbilt Diabetes Research and Training Center, Vanderbilt University Medical Center, Nashville, TN
[3] Center for Cell Systems Biology, Vanderbilt University School of Medicine, Nashville, TN
[4] Mass Spectrometry Research Center, Vanderbilt University School of Medicine, Nashville, TN

Address of Correspondence:
Rafael Arrojo e Drigo
718 Light Hall
2215 Garland Ave
Nashville, Tennessee 37232
r.drigo@vanderbilt.edu

Running Title: Alpha cell autoimmunity and calorie restriction in aging

<u>Abstract</u>

Aging is a major risk factor for insulin resistance and type 2 diabetes, driven in part by declining

pancreatic beta cell function. While calorie restriction (CR) initiated early in life improves

metabolic health and preserves beta cell function, its impact when initiated later in life remains

unclear. We implemented a two-month, 20% CR intervention in 72-week-old male mice and

assessed *in vivo* glucose homeostasis and beta cell function using meal tolerance tests, single

cell RNA-sequencing, and confocal microscopy. We found that old mice exposed to CR have

significantly improved glucose tolerance due to higher insulin sensitivity, which drives a two-fold

reduction in glucose-stimulated beta cell insulin secretion. At the molecular level, CR enhanced

beta cell proteostasis by downregulating ER stress pathways. Remarkably, CR reprogrammed

the alpha cell transcriptome to suppress antigen presentation pathways and cytokine signaling

pathways, including major histocompatibility complex class I (MHC-I) signaling. These

transcriptional changes correlated with profound reduction in the density of adaptive immune

cells in aging islets, including a near-complete loss of cytotoxic CD8$^+$ T cells due to weakened

intercellular communication between antigen-presenting cells with T and B lymphocytes.

Importantly, aging human alpha cells from non-diabetic donors display similar proinflammatory

phenotypes, including higher expression of MHC-I antigens that correlates with higher density of

Cd8$^+$ cells in aging human pancreases. Together, these findings demonstrate that CR remodels

aging alpha and beta cell structure-function to modulate immune cell interactions in aging

pancreases to mitigate age-related immune activation and islet inflammation.

-243 words

Keywords: aging, alpha cells, antigen presentation, caloric restriction, T cells

Introduction

Aging is a major risk factor for type 2 diabetes (T2D), affecting 29.2% of adults 65 and older in the United States[1]. Aging is associated with poor glycemic control due to progressive loss of peripheral insulin sensitivity and pancreatic beta cell function[2-4]. Beta cells are located in the islet of Langerhans next to glucagon-secreting alpha cells and somatostatin-secreting delta cells; together, these major islet cell types regulate glucose homeostasis[5]. These major islet cell types remain largely post-mitotic throughout most of their host organism lifespan[6, 7], and therefore are under constant pressure to maintain function that supports normal glucose homeostasis. Moreover, postmitotic cells are continuously exposed to age-associated stressors that can increase accumulation of molecular and cellular damage, which causes stress that impairs cell structure-function and contribute to age-associated disease. Accordingly, aging beta cells suffer from loss of cellular identity markers, increased DNA damage and senescence, up regulation of ER stress, compromised calcium homeostasis, impaired autophagy, and increased transcriptional noise, amyloid deposition, and islet fibrosis[8-11]. These molecular signatures are linked changes to beta cell function in aging mice and humans[9]. It remains unknown how aging impacts alpha cell and delta cell function.

Calorie restriction (CR) is a well-established dietary intervention that extends life- and health-spans by delaying the onset of age-related cell dysfunction and disease[12-15]. CR regimens of typically 10–40% restriction of energy intake enhance metabolic homeostasis in young and aged animal models, where higher restriction amounts can lead to adverse effects such as muscle wasting and immunosuppression[16]. At the molecular level, CR acts via modulation of major nutrient-sensing pathway nodes, namely the suppression of insulin and IGF1 signaling[17], inhibition of mammalian target of rapamycin complex (mTORC1) activity and activation of AMPK[18], reduction of oxidative stress and inflammatory tone[19], and activation of autophagy[20, 21]. While many of these pathways have been explored for their therapeutic potential via pharmacological targeting using rapamycin (mTORC1 inhibition)[22], metformin

(AMPK activation)[23], or resveratrol[24]. However, many of these interventions exhibit significant

off-target effects, including immunosuppression and beta cell failure[25, 26], that complicate

translation of these approaches to delay or treat age-related human disease.

CR and CR-mimicking interventions (i.e., intermitted fasting and time restricted feeding)

in young adult mice promotes beta cell health, function, and longevity by enhancing peripheral

insulin sensitivity that in turn reduces the metabolic demand for beta cell insulin secretion[27-30].

These pro-longevity effects are associated with increased expression of beta cell identity and

circadian gene expression regulators, improved autophagy and protein homeostasis,

mitochondrial homeostasis and ATP production, and reduced beta cell DNA damage and

senescence[27-30]. Similar adaptive changes to beta cell function are also observed in other

insulin-sensitizing strategies, such as exercise[31, 32], SGLT2 inhibition[33], and incretin agonist

treatments[34]. In contrast, CR has blunted benefits when initiated in mid- to late-life[35], and as

beta cells age they lose their relative plasticity to adapt to changes to whole body metabolism

and dietary interventions may be limited due to irreversible epigenetic changes[36]. Whether CR

and other pro-longevity approaches can rescue beta cell aging signatures in mature and aged

organisms remains unclear.

Here, we investigated whether short-term CR would improve systemic glucose

homeostasis and modulate beta cell structure-function to reverse the hallmarks of beta cell

aging in old mice. Two months of 20% CR significantly improved glucose homeostasis and

triggered a 2-fold reduction in beta cell insulin release. Single-nucleus RNA sequencing

(snRNA-seq) of isolated islets revealed that CR has a limited effect on aging beta cell gene

expression and marked by downregulation of a small set of genes mainly associated with heat-

shock protein (HSP) folding machinery. Strikingly, CR triggered a significant re-organization of

the aging alpha cell transcriptome with lower expression of several pro-inflammatory genes,

including key mediators of immune signaling via enhanced MHC class I-dependent antigen

presentation. Importantly, aging human islets and alpha cells from non-diabetic donors have

significantly higher expression of several pro-inflammatory genes, which correlates with

increased density of activated $CD8^+$ T cells in aging human pancreases.

Together, our findings suggest that old alpha cells could be involved in the regulation of

immune cell infiltration in the aging pancreas and contribute to age-associated pancreas and

islet inflammation phenotypes. These aspects are suppressed by CR, which remodels the

structure-function of aging islet cells and immune cells to reduce organ inflammation.

Therefore, CR may offer a dual mechanism for improving metabolic health and reverse islet

inflammation in aging pancreases.

Results and Discussion

Calorie restriction improves glucose tolerance and reduces insulin secretion in old mice.

To investigate the potential of CR to reverse beta cell aging signatures in old mice, we exposed

72-week-old C57/BL6 male mice to a 2-month 20% CR feeding regimen, while age-matched

controls remained on ad libitum (AL) diet (**Figure 1A**). As expected, CR mice consumed 80% of

total kcal intake compared to AL-fed controls (**Figure 1B, S1A**). This intervention led to a steady

decline in body weight over 2 months, with an average of ~12% body weight loss (**Figure 1C-D,**

S1B-C) due to a predominant loss of body fat mass (**Figure 1E-F**). Tissue-specific weight loss

was also observed in the liver and epididymal white adipose tissue (eWAT) of CR mice (**Figure**

1G, S1-G).

To investigate the impact of CR on whole body metabolism, we placed old AL and CR

mice in metabolic cages for the last week of the diet intervention. While overall energy

expenditure and oxygen consumption were comparable between CR or AL mice (**Figure S2A-**

D), carbon dioxide production and carbohydrate oxidation were significantly increased in CR

mice during the dark period (**Figure S2E-H**). These results indicate that CR mice have a higher

preference for carbohydrate metabolism during their active phase and quickly shift towards fatty

acid metabolism in phases of inactivity and when food is unavailable. Next, we performed mixed

meal tolerance tests (MTTs; 2 g glucose/kg body weight using, Ensure Original) to investigate

the impact of CR on old mouse glucose homeostasis and basal and glucose-stimulated beta cell

insulin release mechanisms. We performed a baseline MTT before starting the CR intervention

at 72 weeks. After 2 months of CR, old mice had significantly improved glucose tolerance post-

intervention at 80 weeks (**Figure 1H**). This phenotype was accompanied by increased insulin

sensitivity, and a ~50% reduction in fasting and in glucose-stimulated insulin secretion (**Figure**

1I-J). Like in younger CR mice[27], CR in old mice preserved beta cell function by significantly

reducing the relative amount of beta cell insulin secretion required to support normoglycemia

(**Figure 1K-L**).

Together, these data demonstrate that CR in old mice leads to significant weight loss

and improvement of systemic glucose homeostasis mechanisms via enhanced peripheral

insulin sensitivity. This also reduces the amount of basal and glucose-stimulated beta cell insulin

secretion, thus suggesting that the potential of old beta cells to adapt to significant

improvements in insulin sensitivity are not impacted during aging.

CR downregulates ER-dependent protein and antigen processing in aging islet cells. In

young adult mice, CR re-wires the landscape of beta cell gene regulatory networks to increase

the expression of beta cell identity and longevity genes. To investigate how CR impacted old

beta cell transcriptional heterogeneity and gene expression, we applied single nucleus RNA

sequencing (snRNA-seq, 10X genomics) of frozen islet pellets extracted from AL or CR mice

(**Figure 2A**). A total of 32,380 cells were sequenced (AL=17,344, CR=15,306) to identify all

major islet cell types, including beta cells (*Ins1*, *Ins2*, and *Ero1b*), alpha cells (*Gcg*), delta cells

(*Sst*), endothelial cells (*Pecam1*), and pancreatic polypeptide cells (*Ppy*) (**Figure 2B-C**).

Approximately 65% of the sequenced cells were beta cells, and CR did not impact the relative

composition of the heterogeneous beta cell population (**Figure 2D**), as well as the ratio of beta

cells to alpha and delta cells in islets (**Figure S3A**).

Pseudobulk analysis of beta cells, followed by KEGG and STRING pathways analysis,

revealed downregulation of genes related to ER protein processing and folding

(*Hspa1a/b/HSP70, Hspa8, Hsp90ab1, Dnaja1/HSP40, and Calr)*, as well as cell stress (*Ypel2*,

Txnip) and insulin synthesis (*Ins1*) genes (**Figure 2E-H**). CR also up-regulated a small number

of genes involved in AKT signaling cascade regulation (*Kctd20, Fkbp5*), and the transcription

factor *Stat3* and the transcription co-activator Pgc1a (*Ppargc1a*) (**Figure 2E**). Remarkably, CR

induced a stronger effect on alpha cells, where it modulated ~5-times more genes than beta

cells (184 vs. 36 differentially expressed genes; **Table S1-2**). Differential gene expression and

pathway enrichment analysis of old CR alpha cell gene signatures revealed upregulation of

pathways associated with glucagon, glucose and fatty acid metabolism (i.e., *Gcg, Gys2, and*

Acacb) and Pgc1a (*Ppargc1a)* (**Figure 3A-B**). In addition, CR suppressed the expression of

n=159 genes involved in T cell receptor (TCR) signaling, antigen presentation, and immune cell

recruitment pathways (i.e., *Cd44, Cd28, Il7r, Il18r1, and Ccr7;* **Figure 3B-C**), as well as genes

involved in cell-cell signaling and MHC class I (*H2-Q7*, *H2-Q6*, *H2-K1*) and class II (*H2-Aa*, *H2-*

Ab1, *H2-Ob*) antigens **(Figure 3D–E)**.

These findings indicate that CR has a limited effect on the transcriptome of aging beta

cells, which contrasts with our previous findings in young adult mice exposed to CR[27]. These

results could be explained to age-dependent changes on the beta cell epigenome that limit the

plasticity of aging beta cell gene expression patterns[36]. Moreover, our data suggests that CR-

mediated remodeling of the beta cell transcriptome may not be required for the functional beta

cell adaptations observed *in vivo* in young[27] and old mice (**Figure 1**). More importantly, CR-

mediated suppression of age-associated pro-inflammatory responses in aging alpha cells

indicates that this cell type could play a role in mediating islet and pancreas immune responses,

as recently suggested in spatial mapping experiments in NOD mice[37].

CR alpha cells have reduced adaptive immunity and TCR signaling. To further investigate

the impact of CR-mediated changes to alpha cell pro-inflammatory gene expression, we

analyzed the relative composition and transcriptional signatures of immune cell types identified

in our snRNA-seq dataset (**Figure 4A**). This revealed a significant decrease in B cells, T cells,

proliferating B cells, and antigen presenting cells (APCs) in CR mice (**Figure 4B**). Notably, the

number and transcriptional profile of islet-resident macrophages remained relatively constant

between AL or CR diets (**Figure 4B**), thus indicating that CR could modulate the recruitment of

immune cells to the aging pancreas via adaptive immune signatures and does not impact

resident immune cells. Accordingly, CR mice had a marked reduction in islet-associated T cells,

including cytotoxic CD8$^+$ T cells that were nearly ablated (**Figure 4C-D**). The remaining CD8+

cells had lower expression of granzyme K (*Gzk*), IL-2 receptor (*Cd25*), and *Txk*; all markers of

activated T cells (**Figure 4E**). Similar changes were observed in B cells **(Figure 4F-G)**.

213   Next, we reconstructed cell-cell communication networks to investigate how CR impacts

214   intercellular communication between islet endocrine and immune cells in AL and CR mice

215   (**Figure 4H**)[38]. This approach revealed an overall downregulation of intercellular communication

216   between immune cells and alpha cells, beta cells, and delta cells. This was underscored by a

217   significant reduction in Cd45- and MHC-based communication between old alpha cells and T-

218   and B-cells in CR mice (**Figure 4I-J**).

219   These findings suggest that suppression of age-related pro-inflammatory gene

expression in alpha cells by CR alters receptor-ligand interactions between alpha cells and

immune cells. This phenotype is marked by reduced CD8$^+$ cytotoxic T cells and reduced

expression of granzyme K, which suggests that CR could reduce age-associated islet

inflammation phenotypes. It is unclear whether alpha and beta cells respond to CR through cell-

autonomous mechanisms or whether intercellular crosstalk between neighboring endocrine cells

contributes to this response.

Upregulation of TCR activation pathways in aging non-diabetic human islets. To translate

our findings to the human setting and to determine if aging non-diabetic human islets have

similar pro-inflammatory signatures, we leveraged a comprehensive human islet proteomics

dataset recently made public by the Human Islet Data Analysis and Sharing (HI-DAS)

consortium (**Figure 5A**)[39, 40]. Here, data from non-diabetic young adults (27±5 years) and old

adults (71±4 years) was analyzed to identify proteins significantly enriched in aged samples

(**Table S2**). Pathway enrichment analysis (KEGG) uncovered increased signatures of amino

acid metabolism, senescence, and ER stress in old islets (**Figure 5B**), as expected. Specifically,

these changes mapped to age-associated increase of beta cell immaturity markers (RBP4 and

upregulated mTORC1 activity regulators, **Figure 5C**), senescence (CDKN1A/p21, **Figure 5D**),

impaired proteostasis and/or protein misfolding by the UPR (ATF6, CHOP, and PERK; **Figure**

5E), and increased amino acid metabolism (BCAT1) and mTORC1 activation proteins (**Figure**

5F). In addition, we also identified an activated immune system response in aging islets, which

had significantly higher expression of MHC-I and TCR proteins, including HLA-A, CD28, and

CD44 (**Figure 5G**), which were also enriched in aging AL mice (**Figure 3C-E**). These results

demonstrate that aging human islets experience an active cellular stress response associated

with increased amino acid metabolism signatures and links this phenotype with increased

expression of pro-inflammatory signatures of TCR activation and immune cell recruitment.

Aging non-diabetic human pancreases have several signs associated with active pro-

inflammatory signaling, including increased fibrosis, higher incidence of acinar-to-ductal

metaplasia (ADM), angiopathy, and pancreatic adiposity[41]. To validate our previous isolated

mouse and human islet analysis *in situ*, we analyzed high-dimensional imaging mass cytometry

(IMC) data from the human pancreas analysis program (HPAP)[42, 43]. Here, non-diabetic human

pancreases were stratified by age to divide samples into two age groups: young adult (22-40

years) and old adult (47-58 years). Individual cells were clustered into unique cell types based

on marker classification; these included alpha (GCG, 7.8%), beta (C-peptide, 6.8%), delta (SST,

0.8%), pancreatic polypeptide (PP; 12%), endothelial cells (CD31, 6%), Epithelial/Acinar cells

(CK19/CA2, 48%), Macrophages (CD68, 9%), and T cells (CD4=0.1%, CD8=1.2%). As

expected, we observed an age-associated decrease in beta cell identity markers (PDX-1 and

NKX6-1; **Figure 5H**), and up-regulation of islet amyloid polypeptide (IAPP) in both alpha and

beta cells (**Figure 5I**). In alpha cells, aging was associated with increased expression of MHC-I

proteins HLA-ABC and attenuated abundance of MHC-2 HLA-DR protein, whereas beta cells

had only attenuated HLA-ABC abundance during aging (**Figure 5J**). Next, we subset the IMC

dataset to identify CD45$^+$ immune cells and found no changes to overall macrophage density

(CD68$^+$ cells expressing CD45) or in the expression of the anti-inflammatory macrophage

marker CD163 (**Figure 5K**). We also observed an age-induced increase in the proportion of

CD8$^+$ T cells specifically, whereas the CD4$^+$ T cell population density was not affected in aging

pancreases (**Figure 5L**).

These findings support our isolated islet analysis to demonstrate that human islet and

pancreas aging is associated with increased expression of pro-inflammatory markers in alpha

cells, which correlates with increased density of CD8$^+$ T cells in the aging pancreas. Since

MHCs present antigenic peptides to CD4$^+$ and CD8$^+$ T cells via TCR complexes, age-associated

increases in alpha cell pro-inflammatory signatures in mice and humans suggests that alpha

cells may be involved in the process of human islet and pancreas inflammation that occurs

during aging.

Acknowledgements: We acknowledge the following Vanderbilt University (VU) and Vanderbilt

University Medical Center (VUMC) core facilities: VUMC Hormone Assay & Analytical Services

Core (NIH DK020593), VU Metabolic Mouse Phenotyping Center [VMMPC (NIH DK059637;

www.vmmpc.org)], Translational Pathology Shared Resource (NCI/NIH Cancer Center Support

Grant 5P30 CA68485-19), and Vanderbilt Technologies for Advanced Genomics (VANTAGE)

core laboratory. VANTAGE is supported in part by Clinical and Translational Science Award

Grant 5UL1 RR024975-03, Vanderbilt Ingram Cancer Center Grant P30 CA68485, Vanderbilt

Vision Center Grant P30 EY08126, and National Institutes of Health/National Center for

Research Resources Grant G20 RR030956. This manuscript used data acquired from the

database (https://hpap.pmacs.upenn.edu/) of the Human Pancreas Analysis Program (HPAP;

RRID:SCR_016202; PMID: 31127054; PMID: 36206763). HPAP is part of a Human Islet

Research Network (RRID:SCR_014393) consortium (UC4-DK112217, U01-DK123594, UC4-

DK112232, and U01-DK123716).

Funding: This work was support by the National Institutes of Health grants 1R01DK138141

awarded to R.A.D. M.S. is supported by the National Institutes of Health Molecular

Endocrinology Training Program (T32DK007563).

Author Contributions: M.S. and R.A.D designed experiments. M.S., A.C., M.C., G.F., and

R.A.D. collected data and assisted with data analysis and management. M.S. and R.A.D.

completed data analysis. M.S. and R.A.D. interpreted data and wrote the manuscript.

Disclosures: The authors have nothing to disclose.

REFERENCES

1.   National Diabetes Statistics Report. Centers for Disease Control and Prevention. 2024. doi: https://www.cdc.gov/diabetes/php/data-research/index.html.

2.   Basu R, Breda E, Oberg AL, Powell CC, Dalla Man C, Basu A, Vittone JL, Klee GG, Arora P, Jensen MD, Toffolo G, Cobelli C, Rizza RA. Mechanisms of the age-associated deterioration in glucose tolerance: contribution of alterations in insulin secretion, action, and clearance. Diabetes. 2003;52(7):1738-48. doi: 10.2337/diabetes.52.7.1738. PubMed PMID: 12829641.

3.   Petersen KF, Befroy D, Dufour S, Dziura J, Ariyan C, Rothman DL, DiPietro L, Cline GW, Shulman GI. Mitochondrial dysfunction in the elderly: possible role in insulin resistance. Science. 2003;300(5622):1140-2. doi: 10.1126/science.1082889. PubMed PMID: 12750520; PMCID: PMC3004429.

4.   Cowie CC, Rust KF, Byrd-Holt DD, Eberhardt MS, Flegal KM, Engelgau MM, Saydah SH, Williams DE, Geiss LS, Gregg EW. Prevalence of diabetes and impaired fasting glucose in adults in the U.S. population: National Health And Nutrition Examination Survey 1999-2002. Diabetes Care. 2006;29(6):1263-8. doi: 10.2337/dc06-0062. PubMed PMID: 16732006.

5.   Cabrera O, Berman DM, Kenyon NS, Ricordi C, Berggren P-O, Caicedo A. The unique cytoarchitecture of human pancreatic islets has implications for islet cell function. Proceedings of the National Academy of Sciences. 2006;103(7):2334-9. doi: doi:10.1073/pnas.0510790103.

6.   Arrojo e Drigo R, Lev-Ram V, Tyagi S, Ramachandra R, Deerinck T, Bushong E, Phan S, Orphan V, Lechene C, Ellisman MH, Hetzer MW. Age Mosaicism across Multiple Scales in Adult Tissues. Cell Metabolism. 2019;30(2):343-51.e3. doi: https://doi.org/10.1016/j.cmet.2019.05.010.

7.   Cnop M, Hughes SJ, Igoillo-Esteve M, Hoppa MB, Sayyed F, van de Laar L, Gunter JH, de Koning EJ, Walls GV, Gray DW, Johnson PR, Hansen BC, Morris JF, Pipeleers-Marichal M, Cnop I, Clark A. The long lifespan and low turnover of human islet beta cells estimated by mathematical modelling of lipofuscin accumulation. Diabetologia. 2010;53(2):321-30. Epub 20091024. doi: 10.1007/s00125-009-1562-x. PubMed PMID: 19855953.

8.   López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. Hallmarks of aging: An expanding universe. Cell. 2023;186(2):243-78. Epub 20230103. doi: 10.1016/j.cell.2022.11.001. PubMed PMID: 36599349.

9.   Shrestha S, Erikson G, Lyon J, Spigelman AF, Bautista A, Manning Fox JE, dos Santos C, Shokhirev M, Cartailler J-P, Hetzer MW, MacDonald PE, Arrojo e Drigo R. Aging compromises human islet beta cell function and identity by decreasing transcription factor activity and inducing ER stress. Science Advances. 2022;8(40):eabo3932. doi: doi:10.1126/sciadv.abo3932.

10.   Sandovici I, Hammerle CM, Cooper WN, Smith NH, Tarry-Adkins JL, Dunmore BJ, Bauer J, Andrews SR, Yeo GS, Ozanne SE, Constância M. Ageing is associated with molecular signatures of inflammation and type 2 diabetes in rat pancreatic islets. Diabetologia. 2016;59(3):502-11. Epub 20151223. doi: 10.1007/s00125-015-3837-8. PubMed PMID: 26699651; PMCID: PMC4742511.

11.   Li L, Trifunovic A, Köhler M, Wang Y, Petrovic Berglund J, Illies C, Juntti-Berggren L, Larsson NG, Berggren PO. Defects in β-cell Ca2+ dynamics in age-induced diabetes. Diabetes. 2014;63(12):4100-14. Epub 20140701. doi: 10.2337/db13-1855. PubMed PMID: 24985350.

12.   Fontana L, Partridge L, Longo VD. Extending healthy life span--from yeast to humans.
Science. 2010;328(5976):321-6. doi: 10.1126/science.1172539. PubMed PMID: 20395504;
PMCID: PMC3607354.
13.   Mattison JA, Colman RJ, Beasley TM, Allison DB, Kemnitz JW, Roth GS, Ingram DK,
Weindruch R, de Cabo R, Anderson RM. Caloric restriction improves health and survival of
rhesus monkeys. Nat Commun. 2017;8:14063. Epub 20170117. doi: 10.1038/ncomms14063.
PubMed PMID: 28094793; PMCID: PMC5247583.
14.   Green CL, Lamming DW, Fontana L. Molecular mechanisms of dietary restriction
promoting health and longevity. Nat Rev Mol Cell Biol. 2022;23(1):56-73. Epub 20210913. doi:
10.1038/s41580-021-00411-4. PubMed PMID: 34518687; PMCID: PMC8692439.
15.   Anderson RM, Le Couteur DG, de Cabo R. Caloric Restriction Research: New Perspectives
on the Biology of Aging. J Gerontol A Biol Sci Med Sci. 2017;73(1):1-3. doi:
10.1093/gerona/glx212. PubMed PMID: 29240911; PMCID: PMC6279158.
16.   Speakman JR, Mitchell SE. Caloric restriction. Mol Aspects Med. 2011;32(3):159-221.
Epub 20110810. doi: 10.1016/j.mam.2011.07.001. PubMed PMID: 21840335.
17.   Mao K, Quipildor GF, Tabrizian T, Novaj A, Guan F, Walters RO, Delahaye F, Hubbard GB,
Ikeno Y, Ejima K, Li P, Allison DB, Salimi-Moosavi H, Beltran PJ, Cohen P, Barzilai N, Huffman DM.
Late-life targeting of the IGF-1 receptor improves healthspan and lifespan in female mice. Nat
Commun. 2018;9(1):2394. Epub 20180619. doi: 10.1038/s41467-018-04805-5. PubMed PMID:
29921922; PMCID: PMC6008442.
18.   Liu GY, Sabatini DM. mTOR at the nexus of nutrition, growth, ageing and disease. Nat Rev
Mol Cell Biol. 2020;21(4):183-203. Epub 20200114. doi: 10.1038/s41580-019-0199-y. PubMed
PMID: 31937935; PMCID: PMC7102936.
19.   Il'yasova D, Fontana L, Bhapkar M, Pieper CF, Spasojevic I, Redman LM, Das SK, Huffman
KM, Kraus WE. Effects of 2 years of caloric restriction on oxidative status assessed by urinary F2-
isoprostanes: The CALERIE 2 randomized clinical trial. Aging Cell. 2018;17(2). Epub 20180209.
doi: 10.1111/acel.12719. PubMed PMID: 29424490; PMCID: PMC5847862.
20.   Yang L, Li P, Fu S, Calay ES, Hotamisligil GS. Defective hepatic autophagy in obesity
promotes ER stress and causes insulin resistance. Cell Metab. 2010;11(6):467-78. doi:
10.1016/j.cmet.2010.04.005. PubMed PMID: 20519119; PMCID: PMC2881480.
21.   Pyo JO, Yoo SM, Ahn HH, Nah J, Hong SH, Kam TI, Jung S, Jung YK. Overexpression of Atg5
in mice activates autophagy and extends lifespan. Nat Commun. 2013;4:2300. doi:
10.1038/ncomms3300. PubMed PMID: 23939249; PMCID: PMC3753544.
22.   Harrison DE, Strong R, Sharp ZD, Nelson JF, Astle CM, Flurkey K, Nadon NL, Wilkinson JE,
Frenkel K, Carter CS, Pahor M, Javors MA, Fernandez E, Miller RA. Rapamycin fed late in life
extends lifespan in genetically heterogeneous mice. Nature. 2009;460(7253):392-5. Epub
20090708. doi: 10.1038/nature08221. PubMed PMID: 19587680; PMCID: PMC2786175.
23.   Martin-Montalvo A, Mercken EM, Mitchell SJ, Palacios HH, Mote PL, Scheibye-Knudsen
M, Gomes AP, Ward TM, Minor RK, Blouin MJ, Schwab M, Pollak M, Zhang Y, Yu Y, Becker KG,
Bohr VA, Ingram DK, Sinclair DA, Wolf NS, Spindler SR, Bernier M, de Cabo R. Metformin
improves healthspan and lifespan in mice. Nat Commun. 2013;4:2192. doi:
10.1038/ncomms3192. PubMed PMID: 23900241; PMCID: PMC3736576.
24.   Baur JA, Pearson KJ, Price NL, Jamieson HA, Lerin C, Kalra A, Prabhu VV, Allard JS, Lopez-
Lluch G, Lewis K, Pistell PJ, Poosala S, Becker KG, Boss O, Gwinn D, Wang M, Ramaswamy S,

Fishbein KW, Spencer RG, Lakatta EG, Le Couteur D, Shaw RJ, Navas P, Puigserver P, Ingram DK,
de Cabo R, Sinclair DA. Resveratrol improves health and survival of mice on a high-calorie diet.
Nature. 2006;444(7117):337-42. Epub 20061101. doi: 10.1038/nature05354. PubMed PMID:
17086191; PMCID: PMC4990206.
25.   Chen C, Liu Y, Liu Y, Zheng P. mTOR regulation and therapeutic rejuvenation of aging
hematopoietic stem cells. Sci Signal. 2009;2(98):ra75. Epub 20091124. doi:
10.1126/scisignal.2000559. PubMed PMID: 19934433; PMCID: PMC4020596.
26.   Fraenkel M, Ketzinel-Gilad M, Ariav Y, Pappo O, Karaca M, Castel J, Berthault MF, Magnan
C, Cerasi E, Kaiser N, Leibowitz G. mTOR inhibition by rapamycin prevents beta-cell adaptation
to hyperglycemia and exacerbates the metabolic state in type 2 diabetes. Diabetes.
2008;57(4):945-57. Epub 20080103. doi: 10.2337/db07-0922. PubMed PMID: 18174523.
27.   dos Santos C, Cambraia A, Shrestha S, Cutler M, Cottam M, Perkins G, Lev-Ram V, Roy B,
Acree C, Kim K-Y, Deerinck T, Dean D, Cartailler JP, MacDonald PE, Hetzer M, Ellisman M, Arrojo e
Drigo R. Calorie restriction increases insulin sensitivity to promote beta cell homeostasis and
longevity in mice. Nature Communications. 2024;15(1):9063. doi: 10.1038/s41467-024-53127-2.
28.   Matta L, Weber P, Erener S, Walth-Hummel A, Hass D, Bühler LK, Klepac K, Szendroedi J,
Guerra J, Rohm M, Sterr M, Lickert H, Bartelt A, Herzig S. Chronic intermittent fasting impairs β
cell maturation and function in adolescent mice. Cell Rep. 2025;44(2):115225. Epub 20250118.
doi: 10.1016/j.celrep.2024.115225. PubMed PMID: 39827461.
29.   Patel S, Yan Z, Remedi MS. Intermittent fasting protects β-cell identity and function in a
type-2 diabetes model. Metabolism. 2024;153:155813. Epub 20240201. doi:
10.1016/j.metabol.2024.155813. PubMed PMID: 38307325; PMCID: PMC10985623.
30.   Brown MR, Sen SK, Mazzone A, Her TK, Xiong Y, Lee JH, Javeed N, Colwell CS, Rakshit K,
LeBrasseur NK, Gaspar-Maia A, Ordog T, Matveyenko AV. Time-restricted feeding prevents
deleterious metabolic effects of circadian disruption through epigenetic control of β cell
function. Sci Adv. 2021;7(51):eabg6856. Epub 20211215. doi: 10.1126/sciadv.abg6856. PubMed
PMID: 34910509; PMCID: PMC8673777.
31.   Coomans de Brachène A, Scoubeau C, Musuaya AE, Costa-Junior JM, Castela A,
Carpentier J, Faoro V, Klass M, Cnop M, Eizirik DL. Exercise as a non-pharmacological
intervention to protect pancreatic beta cells in individuals with type 1 and type 2 diabetes.
Diabetologia. 2023;66(3):450-60. Epub 20221119. doi: 10.1007/s00125-022-05837-9. PubMed
PMID: 36401627; PMCID: PMC9676790.
32.   Carapeto P, Iwasaki K, Hela F, Kahng J, Alves-Wagner AB, Middelbeek RJW, Hirshman MF,
Rutter GA, Goodyear LJ, Aguayo-Mazzucato C. Exercise activates AMPK in mouse and human
pancreatic islets to decrease senescence. Nature Metabolism. 2024. doi: 10.1038/s42255-024-
01130-8.
33.   Al Jobori H, Daniele G, Adams J, Cersosimo E, Solis-Herrera C, Triplitt C, DeFronzo RA,
Abdul-Ghani M. Empagliflozin Treatment Is Associated With Improved β-Cell Function in Type 2
Diabetes Mellitus. J Clin Endocrinol Metab. 2018;103(4):1402-7. doi: 10.1210/jc.2017-01838.
PubMed PMID: 29342295; PMCID: PMC7328850.
34.   Lee YS, Lee C, Choung JS, Jung HS, Jun HS. Glucagon-Like Peptide 1 Increases β-Cell
Regeneration by Promoting α- to β-Cell Transdifferentiation. Diabetes. 2018;67(12):2601-14.
Epub 20180926. doi: 10.2337/db18-0155. PubMed PMID: 30257975.

35.   Hahn O, Drews LF, Nguyen A, Tatsuta T, Gkioni L, Hendrich O, Zhang Q, Langer T, Pletcher
S, Wakelam MJO, Beyer A, Grönke S, Partridge L. A nutritional memory effect counteracts
benefits of dietary restriction in old mice. Nat Metab. 2019;1(11):1059-73. Epub 20191021. doi:
10.1038/s42255-019-0121-0. PubMed PMID: 31742247; PMCID: PMC6861129.
36.   Avrahami D, Li C, Zhang J, Schug J, Avrahami R, Rao S, Stadler MB, Burger L, Schübeler D,
Glaser B, Kaestner KH. Aging-Dependent Demethylation of Regulatory Elements Correlates with
Chromatin State and Improved β Cell Function. Cell Metab. 2015;22(4):619-32. Epub 20150827.
doi: 10.1016/j.cmet.2015.07.025. PubMed PMID: 26321660; PMCID: PMC4598285.
37.   Balasenthilkumaran NV, Whitesell JC, Pyle L, Friedman RS, Kravets V. Network approach
reveals preferential T-cell and macrophage association with α-linked β-cells in early stage of
insulitis in NOD mice. Front Netw Physiol. 2024;4:1393397. Epub 20240624. doi:
10.3389/fnetp.2024.1393397. PubMed PMID: 38979061; PMCID: PMC11228247.
38.   Jin S, Guerrero-Juarez CF, Zhang L, Chang I, Ramos R, Kuan CH, Myung P, Plikus MV, Nie
Q. Inference and analysis of cell-cell communication using CellChat. Nat Commun.
2021;12(1):1088. Epub 20210217. doi: 10.1038/s41467-021-21246-9. PubMed PMID:
33597522; PMCID: PMC7889871.
39.   Ewald JD, Lu Y, Ellis CE, Worton J, Kolic J, Sasaki S, Zhang D, Dos Santos T, Spigelman AF,
Bautista A, Dai XQ, Lyon JG, Smith NP, Wong JM, Rajesh V, Sun H, Sharp SA, Rogalski JC,
Moravcova R, Cen HH, Manning Fox JE, Atlas E, Bruin JE, Mulvihill EE, Verchere CB, Foster LJ,
Gloyn AL, Johnson JD, Pepper AR, Lynn FC, Xia J, MacDonald PE. HumanIslets.com: Improving
accessibility, integration, and usability of human research islet data. Cell Metab. 2025;37(1):7-
11. Epub 20241001. doi: 10.1016/j.cmet.2024.09.001. PubMed PMID: 39357523.
40.   Kolic J, Sun WG, Cen HH, Ewald JD, Rogalski JC, Sasaki S, Sun H, Rajesh V, Xia YH,
Moravcova R, Skovsø S, Spigelman AF, Manning Fox JE, Lyon J, Beet L, Xia J, Lynn FC, Gloyn AL,
Foster LJ, MacDonald PE, Johnson JD. Proteomic predictors of individualized nutrient-specific
insulin secretion in health and disease. Cell Metab. 2024;36(7):1619-33.e5. doi:
10.1016/j.cmet.2024.06.001. PubMed PMID: 38959864; PMCID: PMC11250105.
41.   Wright JJ, Eskaros A, Windon A, Bottino R, Jenkins R, Bradley AM, Aramandla R, Philips S,
Kang H, Saunders DC, Brissova M, Powers AC. Exocrine Pancreas in Type 1 and Type 2 Diabetes:
Different Patterns of Fibrosis, Metaplasia, Angiopathy, and Adiposity. Diabetes.
2024;73(7):1140-52. doi: 10.2337/db23-0009. PubMed PMID: 37881846; PMCID:
PMC11189834.
42.   Kaestner KH, Powers AC, Naji A, Atkinson MA. NIH Initiative to Improve Understanding of
the Pancreas, Islet, and Autoimmunity in Type 1 Diabetes: The Human Pancreas Analysis
Program (HPAP). Diabetes. 2019;68(7):1394-402. Epub 20190524. doi: 10.2337/db19-0058.
PubMed PMID: 31127054; PMCID: PMC6609987.
43.   Shapira SN, Naji A, Atkinson MA, Powers AC, Kaestner KH. Understanding islet
dysfunction in type 2 diabetes through multidimensional pancreatic phenotyping: The Human
Pancreas Analysis Program. Cell Metab. 2022;34(12):1906-13. Epub 20221006. doi:
10.1016/j.cmet.2022.09.013. PubMed PMID: 36206763; PMCID: PMC9742126.

Figure Legends

Figure 1: Late-life calorie restriction improves glucose tolerance and lowers insulin

secretory demand. A) 72-week-old C57Bl/6J mice were fed ad libitum (AL, control) or

subjected to 20% calorie restriction (CR) for 2 months. B) Cumulative energy intake over the

intervention period. C) Body weight measured before (Pre) and after (Post) 2 months CR. D)

Percent change in body weight relative to baseline. E-F) Fat mass and lean body mass (E), and

fat-to-lean mass ratio after 2 months on diet. G) Pancreas, liver, and epididymal white adipose

tissue (eWAT) weight expressed relative to body mass. H) Mixed meal tolerance tests (MTTs)

were performed at baseline (72 weeks) and after the intervention (80 weeks); Area under curve

(AUC) is presented for each test. I) HOMA-IR, calculated as [glucose (mM) × insulin

(µU/mL)]/405. J) Basal insulin levels. K) Insulin levels measured during the MTT. L) Insulin to

glucose ratio. Student's *t-test* was used when comparing two groups (panels B,D,G,H,K-L).

Two-way ANOVA with diet and intervention as main effects was used in panels C,E-F,I-J). n=13

per group. Data are mean ± SE. * $p < 0.05$ for dietary intervention (pre vs. post); # $p < 0.05$ for

group effect (AL vs. CR). Panel A was created by BioRender.com.

Figure 2: Two months calorie restriction downregulates ER-dependent protein

processing in beta cells and antigen presentation in alpha and beta cells. A) Single-

nucleus RNA sequencing (SnRNA-seq) was performed on islets from 80-week-old mice fed

either AL or CR for 2 months. B) Selected marker genes for non-immune islet subsets based on

gene expression. C) UMAP of 32,380 islet nuclei with beta cells segmented from the population.

D) Beta cell subclusters and the proportion within each subcluster. E) Differential gene

expression in beta cells by pseudobulk analysis. F) KEGG pathways significantly downregulated

by CR in beta cells. G) Protein-protein interaction networks shows predicted and known

associations among genes downregulated by CR (STRING). H) Expression of beta cell genes

related to protein processing and antigen presentation. KEGG, Kyoto encyclopedia of genes

and genomes; UMAP, Uniform Manifold Approximation and Projection. * p< 0.05 for dietary

intervention (AL vs. CR). Panel A was created by BioRender.com.

Figure 3: Calorie restriction downregulates TCR activation and antigen presentation in

alpha cells. A) Alpha cell subpopulations were analyzed from snRNA-seq dataset. B) KEGG

pathways significantly modified by CR in alpha cells. C) Protein-protein interaction networks

predicted from genes downregulated by CR in alpha cells. D) Gene ontology networks subset

from the 'Leukocyte activation' subcluster in panel C.   E) Expression of MHC class I and II

genes in alpha cells.   * p< 0.05 for dietary intervention (AL vs. CR). Panel A was created by

BioRender.com.

Figure 4: Two months calorie restriction dampens islet adaptive immunity, and alpha cell-

specific TCR signaling. A) UMAP of immune cell populations, including B cells, T cells,

macrophages, proliferating B cells, and antigen presenting cells (dendritic cells). B) Proportion

of immune cell types shown relative to total immune cells (left) and total islet cells (right). C)

UMAP of T cell populations. D) Proportion of total T cells. E) Expression of Th1 associated

genes; *Gzmk*, *Cd25*, and *Txk*. F-G) UMAP of B cell clusters and the proportion of B cells within

each cluster. H) Intercellular communication networks visualized by total number of interactions

(absolute), and interaction strength (relative). Edge thickness represents interaction strength. I)

Network plot of inferred CD45 signaling interactions. J) Network plot of inferred MHC signaling,

which was only present in AL-fed control mice.   UMAP, Uniform Manifold Approximation and

Projection.

Figure 5: Aging independently increases beta cell immaturity, protein catabolism ER

stress, mTORC1, and TCR activation pathways in human islets. A) Islets from young (27±5

years, n=30), and old (71±4 years, n=27) nondiabetic donors in the Human Islet Data Analysis

and Sharing (HI-DAS) Consortium and were compared by proteomics. B) Reactome pathways

present from significantly abundant proteins between young and old nondiabetic donors. C-D)

Protein expression for RBP4 (C) and CDKN1A [p21] (D). E) Significantly abundant proteins for

UPR sensors, ATF6, CHOP, and PERK. F) Significantly abundant proteins for BCAA catabolism

(BCAT1) and mTORC1 activation (LAMTOR, LAMP2, RAGA, RAGC). F) Significantly abundant

proteins for TCR activation proteins (HLA-A, CD28, and CD44). G) Antigen presenting and TCR

activation proteins HLA-A, CD28, and CD44. Imaging mass cytometry from the Human

Pancreas Analysis Program (HPAP) database were stratified into young 22-40 year old and old

47-58 year old adults, and spatial segmentation was completed. H) Beta cell identity markers,

PDX-1 and NKX6-1. I) Islet amyloid polypeptide (IAPP) abundance in alpha cells and beta cells.

J) Alpha cell and beta cell MHC-I (HLA-ABC) and MHC-II (HLA-DR) abundance. K) Relative

abundance of macrophages within Cd45$^+$ immune cells, and anti-inflammatory macrophage,

CD163, abundance on macrophages. L) Relative abundance of CD4$^+$ and CD8$^+$ T cells within

Cd45$^+$ immune cells. Error bars are mean±SD. * p<0.05, young vs. old donors.

A

AL
17344 cells

CR
15036 cells

UMAP 2
UMAP 1

B

□ AL ■ CR

KEGG Pathways in Alpha Cells

Down by CR ← → Up by CR

Glucagon signaling pathway
Insulin signaling pathway
AMPK signaling pathway
Insulin resistance
ECM-receptor interaction
Antigen processing and presentation
Type I diabetes mellitus
Th1 and Th2 cell differentiation
Allograft rejection
Hematopoietic cell lineage
T cell receptor signaling pathway
Cell adhesion molecules

− log₁₀ FDR

Count
○ 3
○ 6
○ 9
○ 12

C Alpha cell gene networks downregulated by CR

● Leukocyte activation
● V(D)J recombination
● T cell activation
● LPS signaling

D

● Leukocyte cell-cell adhesion (GO:0007159)
● Antigen receptor signaling (GO:00050851)
● Positive regulation of T cell activation (GO:00050870)

E Alpha Cell Features

H2-Aa H2-Ab1 H2-Ob H2-Q7 H2-Q6 H2-K1

Expression Level