Biliverdin Reductase A is a major determinant of neuroprotective Nrf2 signaling

Chirag Vasavda[a,1], Ruchita Kothari[a,1], Navneet Ammal Kaidery[b,c], Suwarna Chakraborty[d], Sunil Jamuna Tripathi[d], Ryan S. Dhindsa[e,f,g], Samaneh Saberi[h], Julia E. Lefler[h], Priyanka Kothari[i], Kalyani Chaubey[j,k,l,m], Cristina Ricco[a], Adele M. Snowman[a], Michael C. Ostrowski[h], Eugenio Barone[n], Lakshminarayan M. Iyer[o], L. Aravind[o], Sudarshana M. Sharma[h], Andrew A. Pieper[j,k,l,m,p,q], Bobby Thomas[b,c,r,s], Solomon H. Snyder[a,d,t,2], Bindu D. Paul[a,d,t,u,2]

[a]The Solomon H. Snyder Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, MD, USA.
[b]Darby Children's Research Institute, Medical University of South Carolina, Charleston, SC, USA.
[c]Department of Pediatrics, Medical University of South Carolina, Charleston, SC, USA.
[d]Department of Pharmacology and Molecular Sciences, Johns Hopkins University School of Medicine, Baltimore, MD, USA.
[e]Department of Pathology and Immunology, Baylor College of Medicine, Houston, TX, USA
[f]Jan and Dan Duncan Neurological Research Institute, Texas Children's Hospital, Houston, TX, USA.
[g]Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, USA.
[h]Department of Biochemistry and Molecular Biology and Hollings Cancer Center, Medical University of South Carolina, Charleston, SC, USA.
[i]Department of Cell Biology, Johns Hopkins University School of Medicine, Baltimore, MD, USA.
[j]Department of Psychiatry, Case Western Reserve University School of Medicine, Cleveland, OH, USA
[k]Brain Health Medicines Center, Harrington Discovery Institute, University Hospitals Cleveland Medical Center, Cleveland, OH, USA.
[l]Department of Pathology, Case Western Reserve University, School of Medicine, Cleveland, OH, USA.
[m]Department of Neuroscience, Case Western Reserve University, School of Medicine, Cleveland, OH, USA.
[n]Department of Biochemical Sciences "A. Rossi-Fanelli", Sapienza University of Rome, Rome, Italy.
[o]Computational Biology Branch, Division of Intramural Research, National Library of Medicine, National Institutes of Health, Bethesda, MD, USA.
[p]Geriatric Psychiatry, GRECC, Louis Stokes Cleveland VA Medical Center, Cleveland, OH, USA.
[q]Institute for Transformative Molecular Medicine, School of Medicine, Case Western Reserve University, Cleveland, OH, USA.
[r]Department of Neuroscience, Medical University of South Carolina, Charleston, SC, USA.
[s]Department of Drug Discovery, Medical University of South Carolina, Charleston, SC, USA.
[t]Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, MD, USA.
[u]Lieber Institute for Brain Development, Baltimore, MD, USA.
[1]These authors contributed equally.

[2]Correspondence to

Bindu D. Paul, Solomon H. Snyder

Email: bpaul8@jhmi.edu; ssnyder@jhmi.edu

Author Contributions: CV, RK, AAP, BT, SHS and BDP designed the study. CV, RT, NAK, SC, SJT, PK, KC, SS, JL, JEL, NAM, AMS, MCO, EB, LMI, LA, AAP, BT, SHS and BDP conducted experiments and/or analysis. NAK, SS, JL, SMS, MCO and BT collaborated on the ChIP-seq

analysis. LMI and LA collaborated on the phylogenetic and sequence of analysis of BVR and HO
proteins. BT, AAP, SHS, and BDP provided resources and funding. CV, RT, LMI, LA, BT, AAP,
SHS, and BDP wrote the paper with input from all authors.

Competing Interest Statement: The authors do not have any competing interests.

Keywords: Biliverdin reductase A, oxidative stress, Nrf2 signaling, neuroprotection, gene
regulation

Abstract

Biliverdin reductase A (BVRA), the terminal enzyme in heme catabolism, generates the
neuroprotective and lipophilic antioxidant bilirubin. Here, we identify a novel non-enzymatic role for
BVRA in redox regulation. We show that BVRA directly interacts with nuclear factor erythroid-
derived factor-like 2 (Nrf2), the master regulator of redox homeostasis, to modulate target signaling
pathways. ChIP-seq and RNA-seq analyses reveal that this interaction coordinates the expression
of neuroprotective genes that are typically dysregulated in Alzheimer's disease and other
neurodegenerative conditions. Thus, this previously unknown BVRA-Nrf2 axis controls an essential
pathway of redox signaling in neuroprotection. Our findings establish BVRA as a dual-function
integrator of antioxidant defenses in both the lipophilic and hydrophilic subcellular compartments,
bridging these two distinct and critical cellular protection mechanisms in the brain. This
advancement in understanding the endogenous antioxidant system of the brain positions the
BVRA-Nrf2 axis as a promising therapeutic target for neurodegenerative disease.

Significance Statement

We show a non-canonical role for biliverdin reductase A (BVRA), classically known as the
biosynthetic enzyme for bilirubin, in nonenzymatic modulation of antioxidant neuroprotective
nuclear factor erythroid-derived factor-like 2 (Nrf2) signaling in the brain. Both BVRA and Nrf2
signaling are compromised in neurodegenerative diseases such as Alzheimer's disease, and the
BVRA-Nrf2 axis offers a new direction for developing neuroprotective therapies.

20
21   **Main Text**
22
23   **Introduction**
24
Biliverdin reductase A (BVRA), encoded by the *Blvra* gene, and the terminal enzyme in heme
catabolism and the major biosynthetic enzyme for bilirubin in adults (Fig. 1*A*) (1, 2). Bilirubin is a
potent lipophilic antioxidant with protective efficacy in advanced cardiovascular disease associated
with hyperbilirubinemia (3-9), as well as in malaria (10) and metabolic disorders (11, 12). Beyond
its canonical role, BVRA exhibits additional diverse enzymatic and non-enzymatic functions,
modulating key signaling pathways such as focal adhesion kinase signaling in the brain, insulin
signaling, phosphoinositide 3-kinase (PI3K)/Protein kinase B (Akt) signaling, and mitogen-activated
protein kinase (MAPK) cascades (11, 13-21). Interestingly, at near-neutral pH BVRA utilizes NADH
as a cofactor, whereas it uses NADPH at alkaline pH, enabling flexibility of function under different
cellular conditions (22-24). Importantly, while BVRA's roles in peripheral tissues are well-
documented, its functions in the central nervous system are not as well understood, despite its
significant expression in brain (2).

We previously showed that both bilirubin and BVRA are highly abundant in the lipid-rich

brain (2). Bilirubin, being lipophilic, functions in a manner complementary to glutathione (GSH), the

abundant hydrophilic antioxidant to confer beneficial effects (2, 7-9, 25). Bilirubin more effectively

prevents lipid peroxidation than soluble cellular components, such as protecting cytosolic proteins

from oxidation (2, 8). Bilirubin is also a potent scavenger of superoxide ($O_2^{\bullet-}$) radicals, with mice

lacking BVRA displaying elevated susceptibility to oxidative stress (2, 9). We have shown that

bilirubin is a potent scavenger of $O_2^{\bullet-}$ and cannot neutralize other forms of reactive oxygen species

(ROS) such as H_2O_2 and the hydroxyl radical (OH$^{\bullet}$)(2).

Despite bilirubin's apparent lack of reactivity towards these ROS, *Blvra*[-/-] mice are still

hypersensitive to H_2O_2 and complementing cells with BVRA is protective, suggesting that these

mice may be deficient in a mechanism dependent on BVRA, but not bilirubin to counteract oxidative

stress (2). Here, we reveal a redox-regulatory role of BVRA that entails a non-enzymatic interaction

with the nuclear factor erythroid-derived factor-like 2 (Nrf2) transcription factor, which is well known

as the master regulator of antioxidant responses (26). We demonstrate that direct binding of BVRA

to Nrf2 enhances transcriptional regulation of neuroprotective gene networks implicated in

Alzheimer's disease (AD) and neurodegeneration. This BVRA-Nrf2 axis establishes a novel link

between heme metabolism and physiologic redox homeostasis, revealing opportunities to

therapeutically bolster endogenous neuroprotection.

56
57
58 **Results**
59

Heme-independent expression patterns and functional roles of brain BVRA

The first enzyme in heme catabolism in the brain, heme oxygenase 2 (HO-2) accounts for

most of the HO activity in the nervous system and, like HO-1, regulates redox balance (27, 28).

HO-2 deletion aggravates oxidative stress induced by seizures, glutamate, and inflammation, and

also causes cerebral vascular injury (29). To determine whether BVRA operates independently of

the heme oxygenase (HO) system, we first identified brain cell types in which heme metabolism is

crucial. Using single-cell RNA sequencing (scRNA-seq), we compared *Blvra* (which encodes

BVRA) and *Hmox2* (which encodes HO-2) expression profiles in whole-brain lysates. Both *Blvra*

and *Hmox2* were detected in numerous cell types in the brain (Fig. 1*B*). On the other hand, *Hmox1*

mRNA (which encodes HO-1) is inducible (30) and not expressed at baseline, rendering it

undetectable in most cells analyzed. Notably, *Blvra* exhibited more restricted cellular expression

than *Hmox2*, with HO-2 mRNA levels nearly tripling those of *Blvra* transcripts. Strikingly, 23% of

brain cells expressed *Blvra* mRNA without detectable *Hmox2* mRNA and only 10% expressed both

Blvra and *Hmox2* (Fig. 1*C*), a pattern observed broadly across cell types and brain regions (Fig.

1*D*). This surprisingly broad presence of BVRA in the absence of HO-2, is puzzling and suggests

a potential function independent of heme metabolism. This dissociation, corroborated by
independent datasets (Allen Brain Atlas, Protein Atlas) (31), implies that BVRA operates in contexts
unrelated to bilirubin generation.

We next examined the evolutionary histories of these principal enzymes involved in heme
degradation: the BVRs (BVRA, BVRB), and HO. BVRA and BVRB are distinct enzymes with BVRA
being the main bilirubin producing enzyme in adults and BVRB in the fetal stages (32). BVRA
belongs to the expansive glyceraldehyde 3-phosphate dehydrogenase (GAPDH)-like clade of
redox enzymes, whose members feature an N-terminal NADPH/NADH-binding Rossmann fold
catalytic domain and a C-terminal GADC domain responsible for substrate selection and binding.
Structural studies of the related *Synechocystis* BVRA revealed that its C-terminal domain
accommodates two biliverdin molecules via π-π stacking, both of which are believed to participate
in the redox reaction. Based on the *Synechocystis* BVRA structure and its interactions with NADPH
and biliverdin, and the sequence conservation pattern, we identified several residues for further
mutational studies (G17, E97, R172, R46, R227, K219). We also noticed a cysteine residue (C292)
that is conserved across vertebrates and some cyanobacteria, which may play a role in BVRA-
mediated redox homeostasis (Fig. S1). In contrast, BVRB lacks the GADC domain and is
comprised of only the NADPH-binding Rossmann fold catalytic domain, which is also present in
BVRA (Fig. 1*F*). Consistent with the absence of a dedicated substrate-binding domain, it exhibits
greater substrate promiscuity. HO, on the other hand, is a heme-dependent all–α-helical oxygenase
that is structurally unrelated to BVRA and BVRB.

Given that homologs of these enzymes are reported in prokaryotes, we explored their
phylogenetic affinities and genomic contexts to gain insights into the evolution of this pathway.
Phylogenetic analyses show that both BVRA and HO are ultimately of cyanobacterial origin.
However, their patterns of acquisition by eukaryotes differ, with the gene for BVRA appearing to
have been transferred from cyanobacteria to an early metazoan (Fig. 1*E*) and HO having been
acquired earlier in eukaryotic evolution, as evidenced by its presence in diverse eukaryotic lineages
such as oomycetes and haptophytes. BVRB, by contrast, appears to represent a lateral transfer
from a firmicute lineage again into an early metazoan. Notably, homologs related to all three
enzymes were transferred on multiple independent occasions from prokaryotes to eukaryotes. For
example, one such paralog of BVRA in eukaryotes, the diphenol detoxifying enzyme dihydrodiol
dehydrogenase (DHDH), was independently acquired from bacteria.

Furthermore, even within metazoans, the phyletic patterns of BVRA and BVRB enzymes
differ considerably: for example, BVRB is widely present in insects, where HO and BVRA are largely
absent. However, HO and BVRA, display a degree of concordance in their distribution among other
metazoans. These patterns support the hypothesis that the heme catabolic pathway combining the
roles of BVRA and HO likely evolved near the base of the animal lineage, whereas BVRB, given
its broader substrate range, including flavins and ferric ions, may have been secondarily

incorporated into heme catabolism in vertebrates, perhaps as an extension of its more generalized
role in redox modification of exogenous toxic small molecules. The cyanobacterial versions of HO
and BVRA likely functioned originally in the metabolism of phycobilins associated with proteins of
their light-harvesting systems. Their early incorporation and shared distribution in animals imply
that they were co-opted for the emerging importance of heme metabolism in the context of oxygen
transport in multicellular animals.

To further characterize the potentially novel function of BVRA decoupled from HO, we
utilized *Blvra*[-/-] mice lacking exon 3 (encoding the NAD(P)H-binding domain) abolishing bilirubin
production (2). Despite originally creating this model to interrogate bilirubin's antioxidant effects,
our earliest observations implicated BVRA itself in redox regulation, including protection against
the hydrophilic H_2O_2 oxidant. We therefore generated a series of mutants in the NADPH and
biliverdin binding domains of BVRA, including the G17A mutant described earlier (21, 33) to assess
their roles in BVRA function (Fig. S1). As previously reported, a glycine to alanine mutation of
residue 17 (G17A) disrupted NADPH binding and significantly diminished BVRA-mediated bilirubin
production (Fig. 1*G*). Specifically, G17A conferred a 5-fold reduction in Vmax and a 300-fold
increase in Km (Fig. 1*H*) of the enzyme. Furthermore, several other mutants derived from residues
that are predicted to bind NADPH/NADH or biliverdin also showed diminished activities (Fig. 1*G*).

Next, we transfected BVRA[-/-] mouse embryonic fibroblasts (MEFs) with WT or G17A-BVRA
to test whether this enzymatically dead mutant could rescue cells exposed to H_2O_2 despite its
inability to produce bilirubin (Fig. 1*I*). G17A-BVRA rescues *Blvra*[-/-] cells to a similar extent as WT-
BVRA, hinting further at a redox function of BVRA independent of bilirubin production and heme
catabolism (Fig. 1*J*).

Depleting BVRA blunts Nrf2 signaling
136

To assess how BVR may exert antioxidant activity independent of its role in bilirubin production,
we monitored the gene expression of several antioxidant genes via unbiased whole transcriptome
sequencing of WT and BVRA[-/-] hippocampal neurons at baseline and under oxidative stress
induced by H_2O_2. After this toxic exposure, several genes were differentially expressed in both cell
types (Fig. 2*A-C,* (*SI Data Table 1*). We focused on genes that show a fold difference of at least 2
in upregulation or downregulation between the WT and *Blvra*[-/-] genotypes. Among the genes
upregulated, the cholesterol biosynthetic pathway was most prominently represented, with
upregulation in BVRA[-/-] cells under stress BVRA[-/-] cells. Most significant among the downregulated
pathways was the nuclear factor erythroid-derived factor-like 2 (Nrf2) signaling pathway (Fig. 2*D,*
right panel), with Nrf2 ranking significantly high amongst the transcription factors implicated in the
differential gene expression between the WT and BVRA[-/-] cells (Fig. 2*E*). We therefore chose to
focus on Nrf2 signaling for the remainder of our study to determine the specific role of BVRA in the
response to oxidative stress.

Nrf2 is a transcription factor that regulates the expression of several antioxidant response
proteins. Under basal conditions, Nrf2 is sequestered by the Kelch-like ECH-associated protein 1
(Keap1) in the cytosol and ubiquitinated for targeted degradation by the 26S proteasome(34). In
response to oxidative stress, Keap1 is modified on its reactive cysteine residues and dissociates
from Nrf2, preventing its ubiquitination and proteasomal degradation (35-39). Nrf2 then
translocates to the nucleus, where it binds the antioxidant response element (ARE) to orchestrate
transcription of a battery of vital antioxidant genes necessary for cytoprotection. To evaluate
whether Nrf2 signaling is disrupted in BVRA$^{-/-}$ cells, we analyzed the transcript levels of Nrf2 target
genes such as *Hmox1, Gclc*, *Gsta2* and *Cxcl2* (Fig. 2*F-I*). The expression of *Hmox1*, *Gclc* (which
encodes the catalytic subunit of glutamate–cysteine ligase, a key enzyme in glutathione
biosynthesis) and *Gsta2* (which encodes glutathione S-transferase A2, an enzyme involved in
detoxification of electrophilic compounds) were blunted in the BVRA$^{-/-}$ cells, whereas the proteins
involved in the immune response and inflammation such as *Cxcl2* (encoding the chemokine (C-X-
C motif) ligand 2) were upregulated.

Next, we compared the activity of the Nrf2 promoter in WT and BVRA$^{-/-}$ cells using a
luciferase reporter assay. In this dual-luciferase firefly reporter system, where the luciferase gene
is regulated by the promoter for glutathione S-transferase A4 (GSTA4), a target of Nrf2. Thus, Nrf2
activity at the *Gsta4* promoter initiates luciferase transcription, which can be measured via luciferin
luminescence. We quantified the activation of the *Gsta4* promoter in WT and BVRA$^{-/-}$ MEFs in
response to three Nrf2 activators: H_2O_2, sulforaphane (SFN), and tert-butyl hydroquinone (*t*-BHQ).
SFN is a compound known to modify the reactive cysteines on Keap1, thus releasing Nrf2 before
ubiquitination and increasing overall expression of intracellular Nrf2 (40). *t*-BHQ is also known to
activate Nrf2, although its mechanism not well-established (41). H_2O_2 treatment increases luciferin
reporter detection in WT cells, indicating Nrf2 activation by proxy of GST gene transcription.
However, this increase is not seen in BVRA$^{-/-}$ cells, which are unable to elicit Nrf2 activity and GST
luciferase expression. The same pattern was also noted in the SFN and tBHQ treatment conditions
(Fig. 2*J*).

To determine whether enzymatically dead BVRA also potentiates Nrf2 transcriptional
activity, we introduced WT and G17A BVRA in a HEK293 cell line harboring an Nrf2 regulated
luciferase reporter. In this HEK293 cell line, the promoter controlling luciferase expression is directly
downstream of the antioxidant response element promoter that Nrf2 binds, allowing us to use
luciferase expression as a readout for Nrf2 transcriptional activity in general and not limited to
GSTA4 expression as in the previous experiment. WT- and G17A-BVRA both increase luciferase
activity significantly and to the same degree in each treatment condition (Fig. 2*K*). Enzymatically
dead BVRA was equally efficacious as WT BVRA in activating Nrf2 *in vitro*, underscoring BVRA's
novel non-catalytic Nrf2-inducing role. We next analyzed the effect of erastin, an inhibitor of the
cystine transporter, one of whose subunits is encoded by *Slc7a11*, a target of Nrf2, on the viability

of these cells. BVRA$^{-/-}$ MEFs were more susceptible to increasing concentrations of erastin as
compared to WT cells, further confirming that depletion of BVRA compromises Nrf2 signaling (Fig.
2L).

BVRA physically and genetically interacts with Nrf2
To characterize the relationship between BVRA and Nrf2, we aimed to breed BVRA and Nrf2
double-knockout mice. In our attempts to create such mice, no *Blvra$^{-/-}$/ Nfe2l2$^{-/-}$* pups were ever
obtained, indicating that this interaction is crucial for survival and function. In the crosses
conducted, we expected roughly 9% of the pups to present with the double-null genotype but were
unable to obtain double knockout pups (Fig. 3A, B). All other possible genotypes were generated
at near their expected frequencies. This finding suggests that BVRA and Nrf2 are genetically linked
and that deleting both genes is embryonic lethal, confirming a genetic interaction between *Blvra*
and *Nfe2l2* that is crucial for survival.
To determine whether BVRA interacts with Nrf2, we conducted a GST pull down assay in
HEK293 transfected with constructs harboring GFP-tagged Nrf2 and either GST vector alone or a
construct encoding GST-BVRA. GST-BVRA robustly bound Nrf2 unlike GST alone, as revealed by
western blot analysis (Fig. 3C). Given that BVRA and BVRB are the two enzymes involved in
bilirubin production, but have completely different evolutionary histories and are only remotely
related, we analyzed whether BVRB has an Nrf2-interacting role. We simultaneously co-
expressing GFP-Nrf2 with either GST-BVRB, GST-BVRA or GST alone in HEK293 cells and
conducted GST pulldown assays. We observed that BVRB does not associate with Nrf2 and this
function of BVR remains specific to BVRA (Fig. 3D). To determine whether this interaction is
dependent on the BVRA's bilirubin-producing activity, we transfected cells with Nrf2 and
enzymatically dead G17A BVRA, which cannot bind the cofactors NADPH or NADH. G17A BVRA
not only pulls down Nrf2 but does so more robustly than WT BVRA, indicating that this interaction
is independent of its catalytic function (Fig. 3E).
As both BVRA and Nrf2 modulate redox homeostasis, we also analyzed whether the
formation of the BVRA-Nrf2 complex increases in response to oxidative stress. We therefore
simulated pro-oxidant conditions to monitor changes in the BVR-Nrf2 interaction. HEK293 cells
exposed to H_2O_2 and pyrogallol, a superoxide generator (42), exhibit increased Nrf2 expression
compared to vehicle conditions, which was associated with increased BVRA-Nrf2 complex
formation via co-immunoprecipitation. (Fig. 3F). Notably, there was no affinity difference in
interaction under different redox conditions.
We also analyzed whether BVRA influences stability of Nrf2. As Nrf2 stability and levels
are regulated by ubiquitination, we first examined whether BVRA increases Nrf2 expression by
binding and hindering ubiquitin association. Through co-immunoprecipitation, we find that BVRA
does not bind or associate with ubiquitin (Fig. S2A). Additionally, we observed no changes in levels

of Nrf2 in the presence or absence of BVRA both under basal conditions or in response to H_2O_2,
sulforaphane or *t*-BHQ exposure (Fig. S2B). Next, we monitored Nrf2 levels in the nucleus of WT
and BVRA$^{-/-}$ cells upon induction of oxidative stress or with treatment with Nrf2 activators such as
sulforaphane. While whole cell levels of Nrf2 are higher in WT cells with oxidative stress compared
to cells lacking BVRA, the levels of Nrf2 in the nucleus are similar in both WT and BVRA$^{-/-}$ MEFs
under the conditions analyzed (Fig. S3C).

Lastly, we investigated whether binding of BVRA to Nrf2 is modulated by its substrate,
biliverdin. We transfected HEK293 cells with GFP-Nrf2 and either GST-BVRA or GST vector,
treated cells with biliverdin and conducted a pulldown assay. In the presence of biliverdin,
interaction of BVRA with Nrf2 decreased as compared to cells which were not treated (Fig. 3*G*).
Thus it appears that the presence of substrates for BVRA can influence its non-canonical functions.
235

236   **BVRA is a component of the transcriptional program of Nrf2-regulated promoters *in vivo***
237

BVRA has been reported to be a transcription factor regulating the expression of activating
transcription factor 2 (ATF2) and HO-1 by binding to the AP-1 sites in their promoters (43, 44).
Considering the physical and genetic interaction of Nrf2 and BVRA, we examined whether BVRA
might potentially coregulate Nrf2 target genes. We performed a genome-wide ChIP-seq analysis
of MEFs generated from previously described FLAG-BVRA transgenic mouse line (21) using
FLAG and Nrf2 antibodies (Fig. 4*A-D*). The peak distribution demonstrated that approximately 10
% of the peaks were located in promoter regions with majority (approximately 40%) harbored in
distal intergenic regions (10 to 50kb; Fig. 4*B*). Motif analysis of both FLAG-BVRA and Nrf2 ChIP-
seq peaks identified bZIP motifs such as Bach1::Mafk motif and Nrf2 motif respectively (Fig. 4*B*).
ChIP-seq peaks were annotated by nearest neighbor method to identify genes that both Nrf2 and
BVRA potentially regulate. Approximately 43% of the genes in the annotated ChIP-seq data had
both BVRA and Nrf2 peaks (*SI Data Table 2*). We further overlaid the BVRA-Nrf2 gene set with
RNAseq-data from WT and BVRA$^{-/-}$ cells. K-means clustering of the Nrf2-BVRA gene expression
identified two clusters that are predominantly enriched for pathways that are dysregulated in
neurodegenerative diseases (Fig. S3). Further, Gene Set Enrichment Analysis (GSEA) analysis of
BVR-Nrf2 associated genes revealed inflammatory pathways, heme metabolism and redox
signaling (Fig. 4*E*).
255

256   Thus, interaction of BVRA with Nrf2 is physiologically relevant and fits in well with the observation
257   that this association is increased in response to oxidative stress to increase the transcription of
258   Nrf2 targets to maintain redox balance in cells

259
260   **Discussion**
261

In this study, we show that BVRA, the biosynthetic enzyme for bilirubin, is an essential
regulator of neuroprotective signaling by the master regulator of redox homeostasis, Nrf2. We
showed previously that bilirubin is a potent antioxidant capable of scavenging superoxide (2).
However, the role of BVRA in oxidative stress independent of its catalytic activity has not been
explored (2). Using advanced genomic techniques, as well as evolutionary, mutational, genetic,
and biochemical studies, we examined the non-canonical role of BVRA in the antioxidant stress
response. sc-RNA-seq analysis of the brain revealed that a significant proportion of brain cell
populations expressed BVRA but not HO-2, the precursor to BVRA in heme catabolism in the brain,
indicating additional functions for BVRA distinct from its classical role in bilirubin production.
Phylogenetic analysis revealed that although HO-2 and BVRA originated in the cyanobacteria in
the context of phycobilisome synthesis in the light-harvesting complex, they were acquired at
different points in time in the eukaryotes. Thus, they were co-opted and combined in a common
heme catabolism pathway only in Metazoa. Using sequence conservation information from a
multiple sequence alignment, we mutated key residues in BVRA and tested these mutants for
BVRA activity. Further, the presence of a single exposed conserved cysteine within a CC motif at
the extreme C-terminus in vertebrates suggests that this residue might be involved in the sensing
of the redox status of the cell.

Amongst the mutants generated, the G17A mutant was severely compromised in the ability
to produce bilirubin as this mutation affects its ability to bind the cofactor, NAD(P)H as reported
earlier (21, 33). Interestingly, G17A BVRA rescued H_2O_2-mediated cell death, demonstrating a
protective role of BVRA, distinct from its catalytic activity. Furthermore, unbiased RNA-seq analysis
of hippocampal neurons derived from BVRA[-/-] mice, identified Nrf2 signaling as a top downregulated
pathway. In support of these findings, both WT and G17A BVRA bind Nrf2 and stimulate
transcription of target genes. We also noted that in the presence of biliverdin, the substrate for
production of bilirubin, the binding of BVRA to Nrf2 is diminished. This suggests a dual yet distinct
role of BVRA with its additional ability to regulate redox metabolism via Nrf2. Although BVRA does
not influence the nuclear translocation of Nrf2, it is essential for optimal function of Nrf2 as revealed
by our RNA-seq and ChIP-seq data. This role of BVRA as a regulator that interacts with a
transcription factor is closely paralleled by another catalytically inactive Rossman fold
oxidoreductase harboring a from the same clade as BVRA, Gal80, which has been co-opted as a
regulator that binds transcription factors (45).

Our studies revealed that multiple pathways are co-regulated by the BVRA-Nrf2-bound
genes, which encompass a range of functions, including mitochondrial function, heme metabolism
to cytokine signaling and oxidative stress response. Delineating this function of BVRA allows us to
understand heme metabolism in the context of oxidant-induced aging and disease. While heme
oxygenases, which catalyze the first step of heme degradation, are known targets for redox
regulation, their overexpression leads to deleterious off-target effects including inflammation and

cognitive decline (46) reducing its therapeutic efficacy (47). Likewise, while bilirubin itself acts as

an endogenous and exogenous ROS suppressor, its effects are limited to distinct forms of ROS

(2). While exogenous bilirubin may only be effective in curtailing excess $O_2^{\bullet-}$, activators of BVRA

expression may be able to regulate ROS in a more all-encompassing manner. The BVRA/Nrf2

interaction is relevant to a wide array of ROS-mediated pathologies. Suboptimal Nrf2 activity has

been observed in neurodegenerative diseases such as PD and AD (48, 49). However, currently

known activators of Nrf2 are predominantly electrophilic compounds that exert adverse side effects.

In such a scenario, molecules that activate BVRA expression or stimulate its transcriptional activity

are of significant therapeutic value. BVRA's role in activating Nrf2 provides a new therapeutic

signaling pathway that mitigates a broader array of oxidative species via multiple downstream

effectors. Augmenting BVRA therefore promises to a more reliable and efficacious approach for

diseases involving imbalanced Nrf2 activity. Additionally, modulating the Nrf2 pathway through

BVRA may also mitigate ferroptosis, an iron-mediated form of cell death, which is observed in

several neurodegenerative diseases as both BVRA and Nrf2 regulate iron disposition, through its

metabolism or transport, and loss of either of these proteins result in accumulation of excess iron

and vulnerability to oxidative stress (50-53).

Collectively, our findings reveal a previously unrecognized and crucial function of BVRA,

which bridges two distinct and critical pathways, namely heme metabolism and Nrf2 signaling, to

afford cytoprotective effects. Furthermore, BVRA orchestrates antioxidant stress responses in both

the lipophilic and hydrophilic subcellular compartments of the cell through not only the production

of bilirubin which directly scavenges $O_2^{\bullet-}$ radicals, but also through its role as a transcriptional

regulator, respectively (Fig. 4F). Augmenting this latter pathway may offer therapeutic benefits in

diseases characterized by redox imbalance. Both the enzymatic and non-enzymatic roles of BVRA

participate in diverse aspects of normal physiology, and characterizing these activities and their

cell-type-specific roles will likely reveal new paradigms for treating neurodegenerative diseases.

326
327 **Materials and Methods**

328
Plasmids/cDNA

Plasmids encoding GST-BVR (GST-BVRA) and GST-BVRB were generated by cloning cDNA

encoding human BVRA and BVRB into pCMV-GST vector (Clontech/TaKaRa). Likewise, the myc-

BVRA (myc-BVRA) plasmid was generated by cloning cDNA of human BLVRA into pCMV-myc

vector. The GFP-Nrf2 plasmid was generated by cloning cDNA of Nrf2 into GFP-vector.

Cell Culture

Mouse Embryonic Fibroblasts (MEFs)

WT, BVRA$^{-/-}$, and Flag BVRA MEFs were isolated as previously described (2). In brief, embryonic

day 14.5 (E14.5) embryos were obtained from timed BVRA$^{+/-}$ matings. The pups were decapitated

and eviscerated, after which the remaining portion was trypsinized and sheared. Isolated MEFs

were plated in 2 wells of a 6-well plate and cultured overnight in DMEM (Gibco) supplemented with

10% FBS, 100 U/mL penicillin and streptomycin, and 2 mM glutamine at 37°C with 5% CO_2. MEFs

were then expanded to 6-well plates and transiently transfected with SV40T antigen (Addgene) and

maintained until stably proliferative. DNA was isolated from the heads of the pups and genotyped

to confirm the genotype of the corresponding MEF cell line. Data for WT and BVRA$^{-/-}$ cells were

drawn from experiments on at least two independent cell lines of each genotype.

Human Embryonic Kidney (HEK) 293 Cells

HEK293 cells were obtained from the American Type Culture Collection. HEK293 cells were

cultured in Dulbecco's Modified Eagle Medium (DMEM), 10% fetal bovine serum,

penicillin/streptomycin (100 U/ml), and glutamine (2 mM) in an atmosphere of 5% CO_2 at 37°C.

Primary neurons

Primary WT and BVRA$^{-/-}$ neurons were isolated as previously described (2). In brief, hippocampi

from embryonic day 18 (E18) embryos were isolated and washed with HBSS. Tissue was then

incubated with 0.25% trypsin in HBSS for 18 min at 37°C in a conical tube, after which trypsin was

inactivated with the addition of FBS. Tissue were then pelleted at 2000*g* for 5 min at 25°C, after

which the supernatant trypsin/FBS was aspirated off. Tissue were then rinsed twice with HBSS,

resuspended in HBSS, and then gently triturated through a fire-polished glass pipette to promote

additional dissociation. Tissue was then filtered through a 70 μm nylon cell strainer to remove

debris. The dissociated cells were pelleted at 2000*g* for 5 min at 25°C and then resuspended in

neuronal culture media (composed of Neurobasal (Gibco) supplemented with 2% B-27 supplement

and 2 mM glutamine). Cells were filtered once more through a 70 μm nylon cell strainer. Dissociated

cells were then plated onto rinsed, poly-D-lysine-coated dishes. Four days later, the conditioned

media was supplemented with fresh neuronal culture media. The day before dissections, plates

were prepared and coated in 0.25 mg/mL poly-D-lysine overnight at 37°C. Just prior to plating,

plates were rinsed three times with sterile water and once with neuronal culture media.

Transfection

HEK293 cells were transfected with Lipofectamine 3000 (ThermoFisher) as per the manufacturer's

instructions. Briefly, HEK293 cells were seeded in either 10 cm dishes ($3x10^6$ cells) or in 60 mm

dishes ($1x10^6$ cells). The next day, transfection complexes were added dropwise to the plated cells.

For cells in 10 cm dishes, 25 μg of DNA, 23.7 μL of Lipofectamine, and 50 μL of P3000 were

brought to a volume of 1 mL in serum-free Optimem media (Thermo). For cells in 60 mm dishes, 5

ug of DNA, 8.25 μL Lipofectamine, and 10 μL P300 were brought to a volume of 500 μL in Optimem.

Cells were harvested 48 hours after transfection.

Additional details of reagents and methods are available in the *Supplementary Section.*

378
379 **Acknowledgments**
380

BDP, SHS and AAP were supported by the American Heart Association and Paul Allen Foundation

Initiative in Brain Health and Cognitive Impairment (19PABH134580006). BDP and AAP by

NIH/NIA 1R01AG071512. BDP also acknowledges support from NIH/NIA 1R21AG073684, NIH

NIDA, grant P50 DA044123, funding from the Solve-ME foundation and the Catalyst Award from

Johns Hopkins University. B.T. was supported by NIH AG077396, NS101967, NS133688, and the

Department of Defense HT94252310443. M.C.O. and S.M.S. were supported by NIH

P01CA236778. AAP was supported by The Valour Foundation, as the Rebecca E. Barchas, MD,

Professor in Translational Psychiatry of Case Western Reserve University and the Morley-Mather

Chair in Neuropsychiatry of University Hospitals of Cleveland Medical Center, and by the Wick

Foundation, Department of Veterans Affairs Merit Award I01BX005976, NIH/NIA RO1AGs066707,

NIH/NIA 1 U01 AG073323, the Louis Stokes VA Medical Center resources and facilities, the Lincoln

Neurotherapeutics Research Fund, the Leonard Krieger Fund of the Cleveland Foundation, the

Meisel & Pesses Family Foundation, and an anonymous donor. LMI and LA are supported by the

funds of the Division of Intramural Research of the National Library of Medicine at the National

Institutes of Health. We also acknowledge biorender.com, a service we used to design our

schematic figures and graphical abstract, and Intelligenomica LLC for genomics support.

References

399
400

1. Paul BD & Pieper AA (2024) Neuroprotective Roles of the Biliverdin Reductase-A/Bilirubin
Axis in the Brain. *Biomolecules* 14(2).
2. Vasavda C*, et al.* (2019) Bilirubin Links Heme Metabolism to Neuroprotection by
Scavenging Superoxide. *Cell Chem Biol* 26(10):1450-1460 e1457.
3. Vitek L*, et al.* (2002) Gilbert syndrome and ischemic heart disease: a protective effect of
elevated bilirubin levels. *Atherosclerosis* 160(2):449-456.
4. Woronyczova J*, et al.* (2022) Serum Bilirubin Concentrations and the Prevalence of Gilbert
Syndrome in Elite Athletes. *Sports Med Open* 8(1):84.
5. Schwertner HA, Jackson WG, & Tolan G (1994) Association of low serum concentration of
bilirubin with increased risk of coronary artery disease. *Clin Chem* 40(1):18-23.
6. Djousse L*, et al.* (2001) Total serum bilirubin and risk of cardiovascular disease in the
Framingham offspring study. *Am J Cardiol* 87(10):1196-1200; A1194, 1197.
7. Stocker R, Yamamoto Y, McDonagh AF, Glazer AN, & Ames BN (1987) Bilirubin is an
antioxidant of possible physiological importance. *Science* 235(4792):1043-1046.
8. Sedlak TW*, et al.* (2009) Bilirubin and glutathione have complementary antioxidant and
cytoprotective roles. *Proc Natl Acad Sci U S A* 106(13):5171-5176.

9. Vasavda C, Snyder SH, & Paul BD (2021) Quantitative measurement of reactive oxygen
species in ex vivo mouse brain slices. *STAR Protoc* 2(1):100332.
10. Figueiredo A*, et al.* (2025) A Metabolite-Based Resistance Mechanism Against Malaria.
bioRxiv.
11. Lanzillotta C*, et al.* (2024) Biliverdin Reductase-A integrates insulin signaling with
mitochondrial metabolism through phosphorylation of GSK3beta. *Redox Biol* 73:103221.
12. Cimini FA*, et al.* (2019) Reduced biliverdin reductase-A levels are associated with early
alterations of insulin signaling in obesity. *Biochim Biophys Acta Mol Basis Dis*
1865(6):1490-1501.
13. Cimini FA, Perluigi M, Barchetta I, Cavallo MG, & Barone E (2022) Role of Biliverdin
Reductase A in the Regulation of Insulin Signaling in Metabolic and Neurodegenerative
Diseases: An Update. *Int J Mol Sci* 23(10).
14. Kapitulnik J & Maines MD (2009) Pleiotropic functions of biliverdin reductase: cellular
signaling and generation of cytoprotective and cytotoxic bilirubin. *Trends Pharmacol Sci*
30(3):129-137.
15. Salim M, Brown-Kipphut BA, & Maines MD (2001) Human biliverdin reductase is
autophosphorylated, and phosphorylation is required for bilirubin formation. *J Biol Chem*
276(14):10929-10934.
16. Lerner-Marmarosh N*, et al.* (2005) Human biliverdin reductase: a member of the insulin
receptor substrate family with serine/threonine/tyrosine kinase activity. *Proc Natl Acad Sci*
U S A 102(20):7109-7114.
17. Wegiel B*, et al.* (2009) Cell surface biliverdin reductase mediates biliverdin-induced anti-
inflammatory effects via phosphatidylinositol 3-kinase and Akt. *J Biol Chem*
284(32):21369-21378.
18. Hinds TD, Jr.*, et al.* (2016) Biliverdin Reductase A Attenuates Hepatic Steatosis by
Inhibition of Glycogen Synthase Kinase (GSK) 3beta Phosphorylation of Serine 73 of
Peroxisome Proliferator-activated Receptor (PPAR) alpha. *J Biol Chem* 291(48):25179-
25191.
19. Stec DE*, et al.* (2020) Biliverdin Reductase A (BVRA) Knockout in Adipocytes Induces
Hypertrophy and Reduces Mitochondria in White Fat of Obese Mice. *Biomolecules* 10(3).
20. Chen W*, et al.* (2021) Bilirubin deficiency renders mice susceptible to hepatic steatosis in
the absence of insulin resistance. *Redox Biol* 47:102152.
21. Vasavda C*, et al.* (2022) Biliverdin reductase bridges focal adhesion kinase to Src to
modulate synaptic signaling. *Sci Signal* 15(733):eabh3066.
22. Kutty RK & Maines MD (1981) Purification and characterization of biliverdin reductase from
rat liver. *J Biol Chem* 256(8):3956-3962.
23. Maines MD & Trakshel GM (1993) Purification and characterization of human biliverdin
reductase. *Arch Biochem Biophys* 300(1):320-326.
24. Maines MD, Mayer RD, Erturk E, Huang TJ, & Disantagnese A (1999) The oxidoreductase,
biliverdin reductase, is induced in human renal carcinoma--pH and cofactor-specific
increase in activity. *J Urol* 162(4):1467-1472.
25. Dore S*, et al.* (1999) Bilirubin, formed by activation of heme oxygenase-2, protects neurons
against oxidative stress injury. *Proc Natl Acad Sci U S A* 96(5):2445-2450.
26. Baird L & Dinkova-Kostova AT (2011) The cytoprotective role of the Keap1-Nrf2 pathway.
Arch Toxicol 85(4):241-272.
27. Verma A, Hirsch DJ, Glatt CE, Ronnett GV, & Snyder SH (1993) Carbon monoxide: a
putative neural messenger. *Science* 259(5093):381-384.
28. Dore S*, et al.* (2000) Heme oxygenase-2 acts to prevent neuronal death in brain cultures
and following transient cerebral ischemia. *Neuroscience* 99(4):587-592.
29. Parfenova H & Leffler CW (2008) Cerebroprotective functions of HO-2. *Curr Pharm Des*
14(5):443-453.
30. Maines MD, Trakshel GM, & Kutty RK (1986) Characterization of two constitutive forms of
rat liver microsomal heme oxygenase. Only one molecular species of the enzyme is
inducible. *J Biol Chem* 261(1):411-419.
31. Zhang Y*, et al.* (2018) Bliverdin reductase-A improves neurological function in a germinal
matrix hemorrhage rat model. *Neurobiol Dis* 110:122-132.

32. O'Brien L, Hosick PA, John K, Stec DE, & Hinds TD, Jr. (2015) Biliverdin reductase
isozymes in metabolism. *Trends Endocrinol Metab* 26(4):212-220.

33. Lerner-Marmarosh N, Miralem T, Gibbs PE, & Maines MD (2007) Regulation of TNF-alpha-
activated PKC-zeta signaling by the human biliverdin reductase: identification of activating
and inhibitory domains of the reductase. *FASEB J* 21(14):3949-3962.

34. Itoh K*, et al.* (1999) Keap1 represses nuclear activation of antioxidant responsive elements
by Nrf2 through binding to the amino-terminal Neh2 domain. *Genes Dev* 13(1):76-86.

35. Baird L & Yamamoto M (2020) The Molecular Mechanisms Regulating the KEAP1-NRF2
Pathway. *Mol Cell Biol* 40(13).

36. Dinkova-Kostova AT, Kostov RV, & Canning P (2017) Keap1, the cysteine-based
mammalian intracellular sensor for electrophiles and oxidants. *Arch Biochem Biophys*
617:84-93.

37. Uruno A & Motohashi H (2011) The Keap1-Nrf2 system as an in vivo sensor for
electrophiles. *Nitric Oxide* 25(2):153-160.

38. Kobayashi A*, et al.* (2006) Oxidative and electrophilic stresses activate Nrf2 through
inhibition of ubiquitination activity of Keap1. *Mol Cell Biol* 26(1):221-229.

39. Yamamoto M, Kensler TW, & Motohashi H (2018) The KEAP1-NRF2 System: a Thiol-
Based Sensor-Effector Apparatus for Maintaining Redox Homeostasis. *Physiol Rev*
98(3):1169-1203.

40. Hu C, Eggler AL, Mesecar AD, & van Breemen RB (2011) Modification of keap1 cysteine
residues by sulforaphane. *Chem Res Toxicol* 24(4):515-521.

41. Imhoff BR & Hansen JM (2010) Tert-butylhydroquinone induces mitochondrial oxidative
stress causing Nrf2 activation. *Cell Biol Toxicol* 26(6):541-551.

42. Yin XJ, Ma JY, Antonini JM, Castranova V, & Ma JK (2004) Roles of reactive oxygen
species and heme oxygenase-1 in modulation of alveolar macrophage-mediated
pulmonary immune responses to Listeria monocytogenes by diesel exhaust particles.
Toxicol Sci 82(1):143-153.

43. Ahmad Z, Salim M, & Maines MD (2002) Human biliverdin reductase is a leucine zipper-
like DNA-binding protein and functions in transcriptional activation of heme oxygenase-1
by oxidative stress. *J Biol Chem* 277(11):9226-9232.

44. Kravets A, Hu Z, Miralem T, Torno MD, & Maines MD (2004) Biliverdin reductase, a novel
regulator for induction of activating transcription factor-2 and heme oxygenase-1. *J Biol*
Chem 279(19):19916-19923.

45. Aravind L & Koonin EV (1998) Eukaryotic transcription regulators derive from ancient
enzymatic domains. *Curr Biol* 8(4):R111-113.

46. Wang D*, et al.* (2015) Overexpression of heme oxygenase 1 causes cognitive decline and
affects pathways for tauopathy in mice. *J Alzheimers Dis* 43(2):519-534.

47. Uttara B, Singh AV, Zamboni P, & Mahajan RT (2009) Oxidative stress and
neurodegenerative diseases: a review of upstream and downstream antioxidant
therapeutic options. *Curr Neuropharmacol* 7(1):65-74.

48. Wang Q*, et al.* (2017) Meta-Analysis of Parkinson's Disease and Alzheimer's Disease
Revealed Commonly Impaired Pathways and Dysregulation of NRF2-Dependent Genes.
J Alzheimers Dis 56(4):1525-1539.

49. Soni P, Sharma SM, Pieper AA, Paul BD, & Thomas B (2025) Nrf2/Bach1 signaling axis:
A promising multifaceted therapeutic strategy for Alzheimer's disease. *Neurotherapeutics*
22(3):e00586.

50. Stockwell BR (2022) Ferroptosis turns 10: Emerging mechanisms, physiological functions,
and therapeutic applications. *Cell* 185(14):2401-2421.

51. Mezzanotte M & Stanga S (2024) Brain Iron Dyshomeostasis and Ferroptosis in
Alzheimer's Disease Pathophysiology: Two Faces of the Same Coin. *Aging Dis.*

52. Berndt C*, et al.* (2024) Ferroptosis in health and disease. *Redox Biol* 75:103211.

53. Wang ZL, Yuan L, Li W, & Li JY (2022) Ferroptosis in Parkinson's disease: glia-neuron
crosstalk. *Trends Mol Med* 28(4):258-269.

54. Pereira PJ*, et al.* (2001) Structure of human biliverdin IXbeta reductase, an early fetal
bilirubin IXbeta producing enzyme. *Nat Struct Biol* 8(3):215-220.

528
529
530 **Figure Legends**
531
Fig. 1. BVRA exhibits expression profiles and redox homeostatic activity unlinked to heme

catabolism. (*A*) Schematic representation of the heme catabolic pathway. Heme derived

from hemoglobin from senescent red blood cells is metabolized by heme oxygenases (HO)

to produce the green pigment, biliverdin. Biliverdin reductase A (BVRA) then reduces

biliverdin to produce the yellow pigment, bilirubin, which is a potent antioxidant. (*B*) T-

distributed stochastic neighbor embedding (t-SNE) analysis of single-cell RNA expression in

primary neuronal cultures from WT C57BL/6J mice. Each point represents a single cell. Left

panel: Cells are clustered by gene expression into populations with similar cell type and

function. Right panel: Expression distribution of *Blvra*, *Hmox2*, both individually and together,

across cell population clusters. (*C*) Quantified percentage of total cells expressing *Blvra*,

Hmox2, and a combination of both genes. (*D*) Scaled Pearson residuals of *Blvra* and *Hmox2*

gene expression in different cell populations. A residual >1 indicates a higher observed to

expected frequency ratio. Each point represents a single cell. (*E*) Phylogenetic trees of BVRA,

HO, and BVRB respectively. (*F*) Protein structures of BVRA and BVRB obtained from Kavanagh

et. al. (PDB Entry - 2H63 (pdb_00002h63) and Periera et. al., respectively (54). Structural

comparison of the two BVR proteins aligned at the Rossmann fold of the NADPH-binding domain.

(*G*) Activity of WT BVRA and BVRA mutants measured at varying concentrations of NADPH and

10 μM biliverdin. Activity measured by absorbance reading of bilirubin production at 442 nM with

V_o calculated according to the method by Michaelis and Menten. n = 3 in triplicate. (*H*) Quantified

V_{max} and K_m of WT BVRA and BVRA mutants. (*I*) Viability of WT and BVRA MEFs transfected

with empty vector, WT BVRA, and G17A- BVRA after 8 h of exposure to serial dosing of H_2O_2.

Data is normalized to viability of vehicle condition. (*J*) Quantified LD_{50}s (I) of H_2O_2 exposure in

WT and BVRA MEFS. n = 3 in triplicate.

555
556 **Fig.2. Nrf2 activity is disrupted in BVRA$^{-/-}$ neurons and fibroblasts**
557
(*A*) Volcano plot analysis of genes from WT and *Blvra$^{-/-}$* primary hippocampal neurons. Each plot

displays the fold change of gene expression on a log scale with their corresponding P values plotted

on a log scale. Each dot represents a gene of interest. Change in gene expression between WT

from vehicle to H_2O_2 condition are shown (*B*) Change in gene expression in *Blvra$^{-/-}$* neurons from

vehicle to H_2O_2 condition. (*C*) Change in gene expression between WT and *Blvra$^{-/-}$* neurons after

H_2O_2 treatment. Red dots indicate genes that are upregulated with H_2O_2 condition in *Blvra$^{-/-}$* neurons.

Blue dots indicate genes that are downregulated with H_2O_2 condition in *Blvra$^{-/-}$* neurons. (*D*)

Logarithmically-scaled *p* values of molecular pathways that are upregulated (left panel) and

downregulated (right panel) in *Blvra$^{-/-}$* neurons compared to WT controls under the H_2O_2 condition.

(*E*) Transcription factors ranked by log-scaled *p* values. A higher *p* value indicates transcription

factors that are more significantly dysregulated in BVRA$^{-/-}$ compared to WT neurons in the H$_2$O$_2$

condition. Transcription factors ranked from 1, most significant, to 1279, least significant. (*F-I*)

Validation of expression of genes regulated by Nrf2 in WT and BVRA$^{-/-}$ neurons. Wild-type (WT)

and BVRA$^{-/-}$ primary hippocampal neurons were treated with or without 200 μM H$_2$O$_2$ for 6 h and

expression of Nrf2 targets were analyzed by qPCR. n=3, SEM, *p* <0.01, Student's t test. (*J*)

Luciferase reporter assays in WT and BVRA$^{-/-}$ MEFs transfected with Gsta4 luciferase reporter

construct after exposure to vehicle, 50 μM H$_2$O$_2$, 10 μM SFN, and 25 μM TBHQ. n = 4 in triplicate.

Sidak's multiple comparison (*K*) Luciferase reporter assays in stable cell HEK293 cell lines harboring

a 5xARE reporter constructs transfected with either WT BVRA or G17A- BVRA and after exposure

to either vehicle, 200 μM H$_2$O$_2$, 10 μM SFN, and 25 μM TBHQ. n = 4 in triplicate. Tukey's multiple

comparison. (*L*) BVRA$^{-/-}$ MEFs are more sensitive to erastin, an inhibitor of the XcT transporter. WT

and BVRA$^{-/-}$ MEFs were treated overnight for 18 hours with increasing concentrations of erastin, a

xCT inhibitor and ferroptosis inducer, and cell viability was assessed by MTT assay. n=7, SEM, *p

< 0.05, **p <0.01, ***p <0.001*, Student's t test.

583 Fig. 3. BVRA genetically and physically interacts with Nrf2

(*A*) Expected versus observed frequency of pups from the *Blvra*$^{-/-}$ and *Nfe2l2*$^{-/-}$ (*Nrf2*$^{-/-}$) cross. (*B*)

Number of observed pups of each genotype. n = 114 pups total. (*C*) GST pulldown with glutathione

sepharose beads of GST-BVRA and GFP-Nrf2 transfected HEK293 cells. GST-only vector control

was used as a negative control. Blots were probed for GFP-Nrf2 and GST-BVRA with their

respective tags. (*D*) Co-immunoprecipitation of GFP-Nrf2 with GST- BVRA and GST-BVRB. Inputs

at the right indicate samples taken before incubation in beads to monitor whole cell expression of

transfected proteins. Blots were probed for GFP-NRF2 and GST-BVRA with their respective tags.

(*E*) GFP-Nrf2 co-immunoprecipitation with GST-WT BVRA and GST-G17A-BVRA. Shorter

exposure was used for G17A IP lanes and longer exposure was used for the remaining lanes. Blots

were probed for GFP-Nrf2 and GST-BVR with their respective tags. (*F*) Co-immunoprecipitation of

GST-BVRA and GFP-Nrf2 with varying oxidative stress conditions, including vehicle, 200 μM H$_2$O$_2$,

and 100 μM pyrogallol. Blots were probed for GFP-Nrf2 and GST-BVRA with their respective tags.

(*G*) Interaction of GST-BVRA and GFP-Nrf2 in the presence of 10 μM biliverdin in HEK293 cells.

Blots were probed for GFP-Nrf2 and GST-BVRA with their respective tags.

600 Fig. 4. ChIP-Seq analysis reveals genes regulated by both BVRA and Nrf2. (*A*) Schematic

representation of methods used for genomic analysis. (The data obtained from ChIP-seq was

overlaid onto the RNA-seq data from the primary hippocampal neurons in vehicle treated (V) or

H$_2$O$_2$-treated conditions (T) (*B*) Global representation of Flag-BVRA and Nrf2 peak distribution and

Motifs. (*C*) Venn diagram showing overlap of genes associated with both BVRA and Nrf2 peaks.

(*D*) MA plot showing differentially expressed readcounts between WT and BVRA$^{-/-}$. (E) GSEA
analysis of BVRA-Nrf2 associated genes from C. (*F*) Summary of canonical and non-canonical
roles of BVRA. BVRA acts on heme-derived biliverdin to produce the antioxidant-cytoprotectant
bilirubin, which scavenges superoxide generated from mitochondrial activity and other processes
to maintain redox balance. In addition to this role in heme catabolism, BVR also functions as a
serine-threonine and tyrosine kinase that modulates several signaling pathways. We show here
that BVRA also regulates Nrf2 signaling, independent of its enzymatic roles.
612
613
614

A

Heme →(*Heme oxygenase*)→ Biliverdin →(*Biliverdin reductase*)→ Bilirubin → *Mitigates oxidative stress*

Fe²⁺   CO

NAD(P)H → NAD⁺/NADP

B

C

10%   23%   67%

Blvra+
Hmox2+
Blvra+/Hmox2+

D

Blvra

Hmox2

E

BVRA   HO   BVRB

cyanobacteria
metazoans

firmicutes
metazoans
metazoans
fungi
cyanobacteria
metazoans

0.5

F

BVRA
BVRB

BVRA unaligned structure

NAD(P)H-binding
(Rossmann fold)

G

K219A
WT
R227A
R46A
R45A
R172A
E97A
G17A

[NADPH] (µM)

H

WT G17A R45A R46A E97A R172A K219A R227A

µM

WT G17A R45A R46A E97A R172A K219A R227A

I

WT
BVRA⁻/⁻
BVRA⁻/⁻ + WT BVRA
BVRA⁻/⁻ + G17A BVRA

[H₂O₂] (M)

J

H₂O₂

WT   WT BVRA   G17A BVRA

BVRA⁻/⁻

LD₅₀ (µM)

Figure 1

Figure 2

Figure 3

Figure 4