Intra-subtype heterogeneity shapes treatment response in *KMT2A*-rearranged ALL across all age groups

Short title: KMT2Ar B-ALL across age

Alina M. Hartmann[1,3,4], Lorenz Bastian[1,3,4], Malwine J. Barz[1,3,4], Johannes Haas[5], Eric Amelunxen[1], Patrick Ehm[2,3,4], Lennart Lenk[2,3,4], Michaela Kotrova[1,3], Thomas Beder[1,3], Fabio Steffen[5], Kerstin Rauwolf[5], Nadine Wolgast[1,3,4], Sonja Bendig[1,3,4], Cecilia Bozzetti[1,3], Julia Alten[2,3], Mayukh Mondal[4,6], Annika Rademacher[2,3], Julia Heymann[2,3], Wencke Walter[7], Claudia Haferlach[7], Aeint-Steffen Ströh[1,3], Anke K. Bergmann[8], Thomas Burmeister[9], Nicola Gökbuget[10], Beat Bornhauser[5], Jean-Pierre Bourquin[5], Monika Brüggemann[1,3,4], Martin Schrappe[2,3,4], Gunnar Cario[2,3,4] *, Claudia D. Baldus[1,3,4] *

[1] Medical Department II, Hematology and Oncology, University Medical Center Schleswig-Holstein, Campus Kiel, Kiel, Germany

[2] Department of Pediatrics I, Pediatric Hematology/Oncology, University Medical Center Schleswig-Holstein, Campus Kiel, Kiel, Germany

[3] University Cancer Center Schleswig-Holstein (UCCSH), University Medical Center Schleswig-Holstein, Germany

[4] Clinical Research Unit CATCH ALL (KFO 5010) funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), Kiel, Germany

[5] University Children Hospital Zurich, Division of Oncology, Comprehensive Cancer Center Zurich, Zurich, Switzerland

[6] Institute for Clinical Molecular Biology, University Medical Center Schleswig-Holstein, Campus Kiel, Kiel, Germany

[7] MLL Munich Leukemia Laboratory, Munich, Germany

[8] University Hospital Würzburg, Clinical Genetics and Genomic Medicine, Josef-Schneider-Str. 2, 97080, Würzburg, Germany

[9] Department of Hematology, Oncology and Tumor Immunology, CVK, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany

[10] Department of Medicine II, Hematology/Oncology, Goethe University Hospital, Frankfurt/M., Frankfurt/M., Germany

* These Authors contributed equally

Corresponding author: Alina M Hartmann, Rosalind Franklin Str. 12, 24105 Kiel Germany; alina.hartmann@uksh.de; ORCID https://orcid.org/0009-0002-0638-0000

Keywords

KMT2A-rearranged B-ALL, Minimal residual disease (MRD), Age-overriding ALL biology, Biological Heterogeneity, B-cell developmental state, Multiomics profiling, *Ex-vivo* drug response profiling, prognostic gene signature

Abstract

Background

KMT2A-rearranged B-cell acute lymphoblastic leukemia (*KMT2A*r B-ALL) exhibits significant heterogeneity in age of onset, developmental origins, and clinical outcomes. The interplay of individual factors influencing early treatment response within this high-risk molecular subtype remains poorly elucidated. We aimed to comprehensively assess how leukemic developmental state, fusion partner, and patient age jointly influence early treatment response and drug sensitivity.

Methods

To identify determinants of early treatment response to induction chemotherapy, we analysed 465 *KMT2A*r B-ALL cases spanning a wide age range (1 month to 89 years) by integrating transcriptomic and genomic profiling with functional drug response and measurable residual disease (MRD) kinetics. Transcriptomic profiling was used to derive a developmental maturity score based on proximity to physiological B-cell differentiation from a normal B-lymphopoiesis reference. Using an ordinal regression model we identify gene expression programs associated with early MRD response.

Results

We observed a strong inverse correlation between MRD clearance with advancing age (p=2.1E-04), proximity to early B-cell-precursor developmental state (low maturity score, p=1.3E-03) and *AFF1* as fusion partner (p=7.0E-04). A combined model confirmed the predominate impact of both maturity and *KMT2A* fusion partner on MRD response, supporting the concept that the cell's developmental state defines therapy response. Gene expression analysis identified cellular traits that relate to MRD response (e.g. chromatin organization, immune modulation and proliferation). This gene expression classifier grouped cases by MRD response but also by *ex-vivo* induction drug sensitivity. Notably, good responders to *ex-vivo* induction drugs were characterized by a higher maturity score

(p=1.8E-03), whereas for less mature *KMT2A*r B-ALL cases response profiles suggested

higher Venetoclax sensitivity.

Conclusions

Our study provides an integrative framework linking developmental phenotype, fusion

partner, and MRD kinetics across the full age spectrum of *KMT2A*r B-ALL. The maturity

score, derived from bulk transcriptome data, offers a biologically relevant predictor of early

treatment response and drug sensitivity. These insights may support future risk-adapted

strategies and therapeutic targeting, particularly in immature *KMT2A*r B-ALL.

Introduction

KMT2A-rearranged B-cell acute lymphoblastic leukemia (*KMT2A*r B-ALL) is considered a

high-risk molecular subtype due to poor treatment response and high risk of relapse. *KMT2A*

is the most prevalent genomic driver in infant ALL, accounting for ~70% of the cases(1,2),

yet also occurs in older children (~2-5%(2,3)) and adult patients (~15%(4)). The prognosis is

especially poor for infant patients and older adults, suggesting intra-subtype molecular

heterogeneity to account for the differences in treatment response. Over 90 *KMT2A* fusion

partners have been identified with over 80% of *KMT2A*r cases having an *AFF1*, *MLLT1* or

MLLT3 rearrangement(5). Adult cases predominantly carry *KMT2A::AFF1* rearrangements

(*AFF1*r); pediatric and infant cases display a greater variety of fusion partners. *KMT2A*

fusion partners have shown to impact therapy response and outcome with *MLLT3*r cases

showing superior outcome compared to *AFF1*r in infant patients, but not in pediatric patients

(age >1 year)(2,6). Molecular characterization of an infant *KMT2A*r B-ALL cohort revealed

subclusters associated with different driver fusions(7), suggesting different mechanisms

based on the underlying gene fusion partner.

In addition to the heterogeneity of molecular drivers, molecular B-ALL subtypes have

differential proximity to the physiological developmental stages of normal B-cell

development, from hematopoietic stem cells (HSCs) to committed pre-B cells. Age-

overriding immunoglobulin rearrangement profiling across molecular subtypes revealed a

higher immunogenetic maturity in pediatric patients, compared to adults(8,9) and single-cell-

sequencing studies of ALL found increasing cell plasticity with age(10). However, molecular

characterization of *KMT2A*r B-ALL cases have unravelled an immature leukemic origin in

younger patients(10–12) associated with increased cell plasticity, retained latent myeloid

potential and co-expression of myeloid marker genes(10). This immature, high plasticity

phenotype has been shown to impact drug sensitivity(13),(14), necessitating alternative

treatment strategies. As the bispecific T-cell engager monoclonal antibody, blinatumomab,

chimeric antigen receptor T-cell therapy and novel agents such as BH3-mimetic Venetoclax

and Menin inhibitors(15,16) are being incorporated into novel treatment algorithms, futures

investigations will have to assess their efficacy in the light of the underlying leukemic origin

of the ALL.

To address the multi-faceted intra-subtype heterogeneity in an age-overarching cohort and

to explore the impact of the underlying cellular context in disease and therapy response, we

conducted a multi-omics analysis in a large cohort of 465 *KMT2A*r diagnostic B-ALL cases

spanning a unique age range upon diagnosis (1 month to 89 years), and including common

KMT2A fusions with representative frequencies(5). We demonstrate that the maturity score

(reflecting proximity to B-cell developmental stages), the *KMT2A* fusion partner and the age

at diagnosis impact measurable residual disease (MRD) response in *KMT2A*r ALL. A

machine learning model revealed maturity and fusion-partners as stronger predictors for

MRD than age. Importantly, distinct leukemic regulatory programs are linked to treatment

response and thus may provide the basis for understanding and targeting molecular

mechanisms of drug resistance in *KMT2A*r B-ALL.

Methods

Patients and transcriptome data set

We have aggregated a unique cohort of n=465 *KMT2A*r ALL cases spanning the whole age range (1 month to 89 years, Figure 1A) including most common *KMT2A* fusion partners with representative frequencies(5) (Figure 1B). The cohort is well characterized with available omics layers including RNAseq (n=325), Sanger sequencing/RNAseq for gene fusion detection (n=436), SNP arrays (n=135), lymphoid capture panel DNA-sequencing(17) (n=82) and drug response profiling(18,19) (DRP, n=61). Moreover, data on clinical annotations including age (n=422), genomic *IG-R* rearrangement profile (n=47), white blood cell count at diagnosis (WBC, n=223) and longitudinal MRD monitoring (n=214) were available. For the n=325 RNAseq cases, n=148 were uniformly sequenced in our facility and defined as discovery cohort. The remaining n=177 cases (validation cohort) were from publicly available sources(20) (n=133) and from a collaborating laboratory(21) (n=41). (Figure 1A; Supplementary Table S1)

Multi-omics data

Transcriptomic data analysis, fusion gene detection and DNA capture panel sequencing were performed by common standard (Supplementary Methods). DRP was performed in an established co-culture system as previously described(18,19,22) (Supplementary Methods).

Clinical response data

For patients treated according to the AIEOP-BFM ALL/INTERFANT and GMALL study group protocols, MRD measurements were performed in the central reference laboratories. Pediatric-inspired induction therapy protocols in adults with comparable MRD timepoints in pediatric regimens allowed for common kinetic definitions of MRD clearance (Supplementary Methods, Supplementary Figure S1).

Data analysis and statistics

We identified *KMT2A* fusion specific gene sets with multi-comparison ANOVA and LASSO feature selection to define genes uniquely expressed in patients with *AFF1*r, *MLLT1*r or *MLLT3*r. The MRD gene expression signature was defined using an ordinal regression model to account for the structured nature of MRD categories (Supplementary Methods).

To validate gene sets, we performed unsupervised clustering of normalized gene counts across our discovery cohort and, where applicable, the validation cohort, with the R pheatmap package (Version 1.0.12). Biological function was annotated using GO-term enrichment analysis (Supplementary Methods).

To define the patient-specific maturity score, representing proximity to physiological B-cell developmental states, we used the machine learning classifier ALLCatchR[23] to define developmental trajectories and fitted linear slopes on the enrichment across B-cell developmental stages. High maturity score values indicate proximity to mature B-cell stages, low maturity scores correspond to more immature B-cell stages. To validate our maturity score we mapped our cohort to two recently published single-cell atlases of physiological B-cell development[10,24] and additionally correlated it to the B-cell map author's multipotency score[10] (Supplementary Methods).

To quantify how age, maturity and gene fusion effect MRD response, we trained a machine learning decision tree (Supplementary methods). The model did not have high enough predictive capacity to be used as an MRD-clearance classification tool (combined accuracy 56.7%, compared to 27.9% random assignment). However, feature importance values could be extracted to learn which features contribute most to the decision process. We validated these findings in a univariate odds ratio analysis (Supplementary Methods).

Data Availability

RNAseq count data are accessible from zenodo at doi 10.5281/zenodo.15638437 for our cohort or through the original publications (St.Jude[20] and MLL[21]). DRP normalized AUC

167 data are provided in Supplementary Table S7. RNAseq raw fastq files will be provided upon

168 acceptance.

Results

To disentangle the molecular underpinnings of treatment response in *KMT2A*r B-ALL, we

aggregated a cohort of n=465 initial diagnosis samples, including n=325 RNAseq samples

from own (n=148) and external (n=177)(20,21) sequencing, representing three clinical

cohorts and an age spectrum from 1 month to 89 years (Figure 1A). Patients were

categorized for subsequent analyses as infants (<1 year; n=63), pediatric (1-17 years,

n=96), adult (18-54 years, n=207) and elderly patients (>=55 years, n=79). To avoid batch

effects, we processed n=148 samples homogeneously in-house (patient age: 2 months to 79

years, median 13 years), serving as a discovery cohort. Integrative data analysis comprised

genomic profiling for karyotypes (SNParray), single nucleotide variants (capture panel

sequencing(17)), and IG/TR rearrangement status, transcriptomic profiling for gene

expression and gene fusions and functional DRP.

KMT2A fusion partners are associated with age and characterized by few cooperating

genomic aberrations

Among B-ALL molecular subtypes, *KMT2A*r ALL is characterized by the widest age range

and holds a diverse repertoire of *KMT2A* fusion partner genes. We observed a significant

correlation between fusion partners and patient's age (Figure 1B; n=305, Chi-sq. test

P<0.0001). While *AFF1* was the most frequent fusion partner gene (n=221/321; 69%), it was

strongly enriched in adult/elderly patients (71% vs infant/pediatric 48%; p=7.7E-12).

Pediatric patients showed a more diverse fusion partners, with *MLLT3*r exclusively found in

infants/pediatric cases, and *MLLT1*r and *MLLT10*r enriched in infant/pediatric cases (61%

and 92% vs adult/elderly 8% and 0.6%, p=0.008 and p=1.5E-4, respectively). Rare *KMT2A*

fusion partners were also included in our cohort [*EPS15* (n=5), *USP2* (n=2), *DCPS* (n=2),

one of each: *MLLT4*, *UBASH3B*, *NRP2*, *ARSA*, *CBL*, *ELL,* Figure 1C] providing a

comprehensive representation of *KMT2A*r ALL.

To investigate the landscape of cooperating genomic events, we performed capture panel

sequencing for genes recurrently involved in ALL (n=82) and SNParrays (n=104). We

detected pathogenic mutations in 32 of the 82 analysed samples [in 8/26 (31%) pediatric

patients and in 24/56 (43%) adult patients]. In total, 21 of 82 patients (26%) carried either a

KRAS or NRAS mutation, the remaining mutations were found in TP53 (6%), ATM (5%),

CDKN2B (2%) and CXCR4, FAT1, RUNX1, TCF3 (<2%, Supplementary Figure S2A).

Mutations were neither enriched in age groups nor associated with driver fusions. The SNP

arrays revealed diploid karyotypes in nearly all cases, without patterns of recurrent

chromosomal gains and losses (Supplementary Figure S2B).

Gene regulatory landscape of *KMT2A*r fusion partners

Multi-comparison ANOVA and LASSO feature selection were used to define genes uniquely

expressed in *AFF1*-, *MLLT1*- or *MLLT3* fusion positive cases. We unravelled a 237-gene

signature in our discovery cohort. The gene signature separated *AFF1*r cases from other

KMT2A fusions in our discovery cohort (Figure 2A) and in the two independent validation

cohorts(20,21) (Supplementary Figure S3A and S3B respectively). *MLLT1* and *MLLT3*

fusions were separated only at the second clustering level, indicating that these cases are

more similar to each other than to *AFF1*r cases (Figure 2A).

Functionally, this gene signature differentiated *AFF1*r cases from non-*AFF1*r cases.

Upregulated genes in *AFF1*r cases included leukemia-associated proliferation genes *MYC*,

AKT3 and *KMT2A* itself, other known cancer genes (e.g. *HMGA2*, HOXC6, *PLAG1*) and

early stem-cell markers such as *DNTT* and *LRG6* (Figure 2B – above diagonal). This

signature indicates the activation of highly proliferative, stem-cell-near cellular programs with

globally activated gene expression and somatic recombination as well as silenced apoptosis

and differentiation signals suggestive of an immature gene regulation in *AFF1*r cases.

In contrast, genes upregulated in non-*AFF1*r cases included B-cell functional genes like fc-

like receptor protein *FCRLA, IGFBP4* and *ZNF521* and proliferation activation genes like

BMI1 and *IL12RB2,* revealing a more mature, differentiated and functionally active B-cell

program that is nonetheless proliferating (Figure 2B – below diagonal).

***KMT2A*r ALL in adults is enriched for immature transcriptional developmental states**

To systematically characterize the developmental underpinnings of *KMT2A*r ALL, we made

use of our immuno-genomic defined RNAseq reference of human B-lymphopoiesis(23).

Single sample gene set enrichment analysis (ssGSEA) for gene sets defining normal B-

lymphopoiesis was used to define the proximity of individual *KMT2A*r ALL samples to their

physiological B-cell counterparts (Figure 3A). Regression analysis across enrichment scores

was used to condense values into a single maturity score (Figure 3B, Supplementary Figure

S4A, Material & Methods).

We validated the stage of differentiation by correlating *IGH* rearrangement patterns on a

genomic level(9) in n=47 patients (n=14 infant, n=13 pediatric, n=20 adult): most patients

displayed complete *IGH*-rearrangements, indicative of late pro-B to pre-B cells of origin. In

line with previous findings(9), the majority of cases revealed clonally evolving IGH-stems,

and the absence of clonal evolution correlated with higher maturity scores (Supplementary

Figure S4B). In addition, we mapped bulk-RNAseq data to two single cell atlases of human

B-cell development(10,24) confirming the prediction that samples with low maturity scores

corresponded to either the pro-B V(D)J compartment or pro-B cycling cells, whereas

samples with high maturity scores aligned with more mature B-cell stages. (large pre-B,

small pre-B, immature B, Figure 3C and Supplementary Figure S5A). We calculated the

"multipotency score" postulated by Iacobucci et al.(10) for our cohort and found a strong

correlation with our maturity score (R^2=0.54, P<0.001, Supplementary Figure 5B) and

accordingly, an increase in the maturity score and a decrease in the multipotency score with

predicted B-developmental stages based on the B-cell developmental map (Supplementary

Figures S5C and S5D respectively).

Increasing maturity correlated with younger age (spearman rank correlation, R=0.45,

p<2.2e-15). To separate the effect of age on maturity from the effect of fusion gene, we

assessed the maturity scores separately for every driver fusion (Figure 3D). *AFF1*r cases

were most immature (n=221, median maturity score=-0.41) compared to samples carrying

other fusions. Within *AFF1*r cases, we observed increasing immaturity with increasing age

(R=0.16, p=0.024). *MLLT1*r cases had a higher median maturity score than *AFF1*r cases

(n=43, median maturity score=-0.21; p=5.3e-06), but again increasing age correlated with

increasing immaturity (R=0.6, p=4.8e-05). *MLLT3*r and *MLLT10*r had the highest maturity

scores (*MLLT3*r*:* n=28, median maturity score=0.33; *MLLT10*r*:* n=14, median maturity

score=0.34) and were only observed in young patients (median age 0.95 years and 0.9

years, respectively).

Age, driver fusion and maturity modulate MRD response

To define the molecular underpinnings of treatment response, we made use of our

integrated age-overring cohort including infants (14.5%), pediatric (22.9%), adult (52.8%)

and elderly (9.8%) patients treated according to BFM-based pediatric or adult treatment

protocols with corresponding MRD measurements available. MRD was measured at two

timepoints during induction therapy and after induction/before consolidation using clonal

IG/TR or *KMT2A* rearrangements as previously described(25). Of the 214 MRD evaluable

patients, 18.2%, 24.8%, and 57.0% had a fast, intermediate, and slow MRD clearance,

respectively. Age significantly correlated with the initial MRD clearance (n=214, multi-

comparison ANOVA p=0.0002), with the median age of patients in fast MRD responders

being 9.6 years in contrast to 37 years in slow responders (Figure 4A). Overall, 51% of infant

(n=16/31), 31% of pediatric (n=15/49), 65% of adult (n=73/113) and 86% of elderly (n=18/21)

patients showed slow MRD clearance.

Considering driver fusion partner and maturity as factors correlated with age and each other,

we next evaluated MRD clearance with respect to age within each driver fusion group

(Figure S6A) and with respect to maturity score (Figure S6B). Considering the most frequent

fusion partners *AFF1* (n=144), *MLLT1* (n=23), and *MLLT3* (n=18), we observed a skewed

distribution in the three MRD categories fast/intermediate/slow. *AFF1*r cases predominantly

showed an intermediate or slow MRD clearance (92%) compared to *MLLT1* (57%) and

MLLT3 (50%) cases with a faster MRD response (total n=185, Chi-squared test with Monte

Carlo Simulation, p=0.0005; Supplementary Figure S6A). Correspondingly, infants showed

faster MRD response if they were *MLLT1*r or *MLLT3*r, but slower in case of *AFF1*r

(*MLLT1*r*/MLLT3*r: 6/13 slow/intermediate; *AFF1*: 17/17 slow/intermediate, Fishers exact test,

p=0.0008).

In addition, cases with slow or intermediate MRD clearance had significantly lower maturity

scores than patients with fast MRD clearance (Wilcox-test, p=0.0025, Figure 4B). This

pattern persisted when considering maturity score, age group and fusion partner separately

(Supplementary Figure S6B).

To quantify the respective effects of these factors in a combined analysis, we trained a

decision tree (Figure 4C, Material & Methods). The tree grouped patients first by maturity

score (>-0.04 to fast), next by fusion (*MLLT1*r to fast) and the remaining cases by maturity

score (≤-0.28 to slow, >-0.28 to intermediate), suggesting that the combination of maturity

and driver fusion have a high predictive value for MRD response and should be considered

together with patient age. We confirmed the impact of maturity, driver fusion and age on

MRD clearance in a univariate analysis considering age, sex, fusion partner, maturity score

and WBC (Figure 4D).

MRD driven gene expression signatures

To characterize the molecular programs underlying MRD response, we used RNAseq

expression data and built an ordinal regression model, with age and gene fusion as

covariates for MRD-associated genes independent of these factors. This resulted in a 448-

gene signature (Supplementary Table S6) which grouped patients into three main clusters in

an unsupervised clustering approach. Two clusters distinguished patients with fast and slow

MRD clearance (Cluster 1; n=15/23 (65% fast clearance, Cluster 3; n=26/27 (96%)

intermediate/slow clearance, Fisher's exact test p=0.007; Figure 5A). We condensed the

gene signature into a single "resistance score" using ssGSEA and confirmed that this score

was highly predictive for MRD response (Figure 5B).

There was only a minimal overlap of this MRD-gene expression signature with gene

expression profiles associated with *KMT2A* fusion partner or defining different B-cell stages

(Supplementary Figure S6C) underscoring that the MRD-signature represented specific

MRD-associated molecular programs irrespective of the fusion signature.

To functionally annotate the signature of MRD-slow-responders, we applied gene set

enrichment analysis for genes upregulated in MRD-slow-responders (Figure 5C) and found

enrichment of GO-terms associated with chromatin organization, driven by genes coding for

core histones [H1 linker histones *H1-3* and *H1-4*; H2B histones (e.g. *H2BC6, H2BC11*); H2A

histones (e.g. *H2AC16*), and H3 histones]. Group 2 and 4 were functionality associated with

cell differentiation and proliferation, respectively. The fifth GO-term group was associated

with immune response and leukocyte activation and accordingly was driven by immune-

modulator genes like *IL5RA, IL1RL1* and *AZU1* and inflammatory response genes like

ANXA1. Overall, the MRD-associated gene regulation appears not be driven by one

prominent signaling pathway but rather reflects a complex interplay of cellular functions

related to proliferation and differentiation, which in turn points to the developmental

underpinnings (maturity) as a modulator of treatment sensitivity.

Age, maturity and gene fusion reflect *ex-vivo* drug response

To functionally characterize the MRD response we performed *ex-vivo* DRP on diagnostic primary patient material. We screened 61 *KMT2A*r cases (n=21 infant, n=22 pediatric, n=16 adult, n=1 elderly) with a library of n=27 drugs, including both standard-of-care chemotherapeutics and targeted therapy agents such as BCL2-family inhibitors Venetoclax and Navitoclax, proteasome inhibitor Bortezomib or HDAC-inhibitor Panobinostat (Figure 6A).

We split samples into quartiles based on their mean drug response to induction phase drugs (Asparaginase, Cytarabine, Dexamethasone, Daunorubicin, Doxorubicin, Vincristine) quantified by the logarithmic area under the dose response curve (logAUC). We observed high concordance with MRD response: samples with a good *ex-vivo* drug response (Q1 and Q2) were significantly enriched for patients with fast MRD clearance (Q1/2 sensitive DRP group included 13/26 fast MRD clearance patients compared to the Q3/4 DRP resistant group with only 2/22 fast MRD clearance patients; Fisher's exact test p=0.004; Figure 6A, Supplementary Figure S7). This difference in response was also significant for the single drugs Dexamethasone, Daunorubicin, Asparaginase and Vincristine individually (Wilcox-test slow vs. fast MRD clearance; p-value cut-off 0.1; Dexamethasone p=0.0027; Daunorubicin p=0.084, Vincristine p=0.012, Asparaginase p=0.02, Figure 6B).

Q1 and Q2 (most DRP sensitive) were slightly enriched for infant and pediatric patients, although not significantly (Fisher's exact test p=0.16), significantly enriched for non-*AFF1* gene fusion partners (Fisher's exact test p=0.01) and had significantly higher maturity scores (Wilcox-test, p=0.016; Figure 6C and Supplementary Figure S7).

In line with these results, our gene expression-based resistance score (Figure 5B), reflecting high expression of MRD-slow-clearance associated genes, was significantly higher in cases with poor response to induction phase drugs (Q1/2: median resistance score -0.13, Q3/4: median resistance score -0.03, Wilcox-test p=0.002). Thus, the activation of MRD-

associated gene regulation at diagnosis can be captured by gene expression and is

reflected in functional *ex-vivo* drug response profiling (Figure 6A).

We also evaluated three Menin inhibitors (MI-503, SNDX-5613, and VTP-50469). In contrast

to our main DRP panel, these compounds were assessed after six rather than three days of

incubation to account for their slower kinetics. Within the KMT2A-r cohort, we observed no

significant variation in Menin inhibitor sensitivity with respect to developmental maturity, age,

or MRD clearance (Supplementary Figure□S8). For MI-503, AFF1-rearranged cases

showed a slightly reduced response compared to other fusions (ANOVA $P□=□0.02$),

although the effect size was modest and not observed for SNDX-5613 or VTP-50469.

Because the assay used a different incubation time, these data could not be integrated with

the rest of the DRP panel, and we therefore cannot compare overall Menin inhibitor

response to other drugs or to different molecular subtypes.

Venetoclax responders are more frequently immature and characterized by AFF1

fusions

In addition to standard induction chemotherapy drugs, we further analyzed targeted

compounds revealing several drugs with notable activities in our cohort including the

proteasome-inhibitor Bortezomib (n=45), the HDAC-inhibitor Panobinostat (n=57) and the

BCL2-inhibitors Navitoclax (n=46) and Venetoclax (n=58). We matched the sensitivity

profiles for these targeted compounds with MRD clearance and maturity to find potential

vulnerabilities for MRD-slow responding patients.

By comparing drug response with maturity scores [based on the cutoffs defined in the

decision tree in Figure 4C: maturity score <=-0.28 (immature); <=-0.04 and >-0.28, >-0.04

(mature)], we did not observe decreasing sensitivity to Navitoclax, Bortezomib or Venetoclax

with increasing immaturity, as opposed to induction therapy drugs (Figure 6D). Interestingly,

for Venetoclax we noticed a slight inverse trend showing highest sensitivity in the most

immature group, although this trend was not significant (Wilcox-test, p=0.18). Consistently,

we did not observe a correlation of MRD clearance with Venetoclax response (Figure 6B). In

fact, cases sensitive to Venetoclax mirror the MRD-slow-response profile, with low maturity

scores, predominantly *AFF1*r and older age (Supplementary Figure S7), highlighting the

potential of Venetoclax in the treatment of cases with MRD-slow response profile, especially

those with low B-cell maturity.

Discussion

Prognosis of *KMT2A*r ALL varies dramatically across age groups, suggesting an intra-

subtype molecular heterogeneity to account for the differences in treatment response. To

address the underlying biology, we have aggregated the - to our knowledge - largest cohort

of *KMT2A*r patients extensively characterized by multi-level data analysis and clinical

annotations. This is the first study to integrate transcriptomic developmental state, driver

fusion partner, longitudinal MRD response, and *ex-vivo* drug profiling across the full age

spectrum of *KMT2A*r B-ALL. By linking these molecular layers to treatment outcome, we

identify cellular vulnerabilities and provide a framework for risk stratification. Previous

studies(10,7,11–14) on *KMT2A*r ALL have mostly focused on single aspects like age(13),

infant patients(7,11,12), or included only small cohorts of *KMT2A*r patients(11,14). Because

KMT2A fusion partners are unevenly distributed across age groups and influence molecular

heterogeneity and outcome, large data sets are necessary to account for these contributing

factors(1,2,6). In addition, as only few cooperating genomic events characterize the *KMT2A*r

subgroup, it is likely that cell intrinsic factors drive its intra-subtype heterogeneity.

Dissecting the gene expression signatures of *KMT2A*r B-ALLs with respect to fusion partner

revealed distinct cellular programs activated by the different driver fusions, indicating a

highly proliferative, cell-cycle active gene expression program in *AFF1*r cases and a more B-

cell functional, mature signature in non-*AFF1*r cases. Using our newly developed maturity

score we observed that older patients exhibited more transcriptionally immature

developmental states compared to younger patients, even when controlling for fusion

partner. By analyzing maturity within each fusion group and age group, we show that these

patterns are consistent across the full age range, suggesting that chronological age and

leukemic B-cell maturity are inversely correlated. Age specific difference in *KMT2A*r ALL

therefore most likely reflect individual characteristics of a continuous disease spectrum

rather than defined sub-entities.

Consistent with prior reports(1,2,4–6), age correlated with early MRD clearance. However,

our fine-scale dissection demonstrates that fusion partner and maturity correlate with age

and thus a complex interplay of different factors predicts MRD response. Patients with slow

or intermediate MRD clearance had significantly lower maturity scores than fast MRD

responders, and this association held when maturity, age, and fusion were modeled

independently. Integrating these factors as covariates, we derived a novel gene expression

signature separating patients by MRD response. Patients with slow MRD clearance

displayed activation of cellular programs at diagnosis involved in chromatin organisation,

proliferation and immune modulators. Recent studies have begun to link transcriptional

maturity to therapy sensitivity in B-ALL. Yoshimura et al.(13) detected a gene-expression

based "adult-like" phenotype in pediatric *KMT2A*r, *DUX4*r and *CRLF2*r patients. This

phenotype, defined by molecular "adult-likeness" rather than chronological age was linked to

decreased Mercaptopurine-sensitivity. Huang et al(14) analysed 203 *ex-vivo* DRPs across

molecular subtypes and linked Asparaginase-resistance to pre-pro-B cell of origin,

characterized by *BCL2* expression, and suggest Asparaginase and Venetoclax to exhibit

synergistic cytotoxicity. Building on these findings, our study expands the analysis to the

largest *KMT2A*r cohort, integrating fusion-specific effects, full age range, and direct MRD

and *ex-vivo* response correlations.

Functional *ex-vivo* drug profiling mirrored clinical MRD response for drugs administered

during initial therapy protocol in line with previous studies(26,27), further advocating the use

of *ex-vivo* induction therapy response for risk stratification(28). Integrating our drug response

profiling data with the transcriptional profiling revealed that poor response to standard-of-

care chemotherapeutics is associated with immature transcriptional states. Accordingly, the

MRD-slow-response cases were characterized by an immature developmental stage, *AFF1*r

and poorer *ex-vivo* response to standard of care chemotherapeutics. Intriguingly, patients

that were responding best to Venetoclax matched the profile of MRD-poor response, with

immature developmental stage and predominantly *AFF1*r. Thus, the efficacy of targeted

therapies such as Venetoclax was independent of the B-cell developmental states in our

functional DRP, suggesting these compounds as a therapeutic alternative in patients with

increased risk of poor chemotherapy response.

Using single-cell RNAseq of 89 B-ALL cases (including only nine with *KMT2A*r), Iacobucci et

al.(10) developed a "multipotency score", characterizing patients with an enrichment of early

lymphoid progenitors, variation in transcriptional programs and proximity to early B-cell

developmental states. The *KMT2A*r cases grouped into a *KMT2A*-early and *KMT2A*-

commited cluster, however in-depth characterization of the multipotency of these cases with

respect to gene fusion and age is missing. We have reproduced this multipotency score

across our cohort and could show that it highly correlates with our maturity score. We extend

these insights into a bulk-transcriptome context suitable for clinical application. Together, our

findings indicate that leukemic B-cell developmental state and fusion partner jointly shape

therapy response and drug sensitivity, exceeding the predictive value of chronological age.

We provide the analytical groundwork to explore interdependencies between transcriptional

developmental B-cell state, selection of genomic drivers, gene regulation and biological and

clinical phenotypes in B-ALL across molecular subtypes. The inherent correlation of MRD

clearance, our MRD-related gene expression signature and the corresponding drug

response profile show that molecular programs activated already at diagnosis are essential

for and can predict MRD-response as well as are reflected in the functional *ex-vivo* DRP in

*KMT2A*r ALL. Our findings highlight the importance of the leukemic B-cell developmental

states in determining outcome and drug sensitivity of *KMT2A* rearranged ALL. In future, a

consensus definition of B-ALL proximity to normal B-lymphopoiesis might be included in

novel risk stratification models, putatively extending beyond *KMT2A*r B-ALL.

Acknowledgements

RNAseq was performed in cooperation with the Competence Center for Genomic Analysis

Kiel (CCGA).

Special thanks to Dr. Rolf Koehler, Institute for Humangenetics, Heidelberg for providing

parts of the MRD-data in older children and Dr. Claus Meyer, DCAL, Frankfurt for providing

the KMT2A-sequence for MRD analysis.

This study was funded in part by the Deutsche Forschungsgemeinschaft (DFG; German

Research Foundation) project number 444949889 (KFO 5010 Clinical Research Unit

"CATCH ALL" to A.M.H., L.B., M.J.B, P.E., L.L., N.W., S.B., M.M., M.Br., M.S., G.C, and

C.D.B.)

Authorship

<u>Contribution:</u>

A.M.H., L.B., M.J.B., G.C., C.D.B. designed research;

A.M.H., L.B., M.J.B., J.Ha., E.A., P.E., M.K., T.Be., F.S., K.R., N.W., M.M., B.B. established

bioinformatic workflows and performed research;

A.M.H., L.B., M.J.B., L.L., F.S., K.R., B.B., G.C., C.D.B analyzed and interpreted data;

A.M.H., L.B., N.G., A.K.B., M.Br., M.S., G.C., C.D.B. supervised the study;

A.M.H. developed statistical models and produced visualizations.

P.E., S.B contributed RNAseq data;

J.A., A.R., J.He., M.S., G.C., contributed pediatric MRD data;

A.-S.S., T.Bu., N.G. and M.Br. contributed adult MRD data;

J.Ha., E.A., F.S., K.R., B.B., J.-P.B. performed and analyzed drug response profiling;

B.B., J.-P.B. supervised drug response profiling;

C.B. contributed DNA capture panel data;

W.W., C.H. provided samples and clinical and diagnostic data

A.M.H., L.B., C.D.B. wrote the manuscript; and all authors revised and approved the final

version of the manuscript

<u>Conflict of interest:</u>

Lenk: *OSE Immunotherapeutics:* Research Funding. **Burmeister:** *Pfizer Inc.:* Honoraria.

Gökbuget: *Amgen, Astra Zeneca, Autolus, Clinigen, Gilead, Incyte, Jazz Pharmaceuticals,*

Novartis, Pfizer, Sanofi, Servier: Consultancy, Honoraria, Other: Advisory board; *Amgen,*

Clinigen, Incyte, Jazz Pharmaceuticals, Novartis, Pfizer, Servier: Research Funding.

Brüggemann: *Amgen Becton Dickinson AstraZeneca Jazz,Pfizer:* Consultancy, Honoraria,

Research Funding, Speakers Bureau. **Schrappe:** *JazzPharma, Servier, Amgen:* Honoraria,

Research Funding, Speakers Bureau. **Cario:** *Jazz Pharmaceuticals:* Other: travel support.

Baldus: *Janssen, Astellas, Pfizer, Astrazeneca, Servier, BMS:* Consultancy, Honoraria.

References

1. Pieters R, Schrappe M, De Lorenzo P, Hann I, De Rossi G, Felice M, et al. A treatment
protocol for infants younger than 1 year with acute lymphoblastic leukaemia (Interfant-
99): an observational study and a multicentre randomised trial. The Lancet. 2007
July;370(9583):240–50.

2. Szczepański T, Harrison CJ, Van Dongen JJ. Genetic aberrations in paediatric acute
leukaemias and implications for management of patients. Lancet Oncol. 2010
Sept;11(9):880–9.

3. Brady SW, Roberts KG, Gu Z, Shi L, Pounds S, Pei D, et al. The genomic landscape of
pediatric acute lymphoblastic leukemia. Nat Genet. 2022 Sept;54(9):1376–89.

4.   Roberts KG. Genetics and prognosis of ALL in children vs adults. Hematology. 2018 Nov
30;2018(1):137–45.

5.   Meyer C, Larghero P, Almeida Lopes B, Burmeister T, Gröger D, Sutton R, et al. The
KMT2A recombinome of acute leukemias in 2023. Leukemia. 2023 May;37(5):988–
1005.

6.   Pui CH, Gaynon PS, Boyett JM, Chessells JM, Baruchel A, Kamps W, et al. Outcome of
treatment in childhood acute lymphoblastic leukaemia with rearrangements of the 11q23
chromosomal region. Lancet Lond Engl. 2002 June 1;359(9321):1909–15.

7.   Isobe T, Takagi M, Sato-Otsubo A, Nishimura A, Nagae G, Yamagishi C, et al. Multi-
omics analysis defines highly refractory RAS burdened immature subgroup of infant
acute lymphoblastic leukemia. Nat Commun. 2022 Aug 30;13(1):4501.

8.   Kotrova M, Proske C, Darzentas N, Laqua A, Kehden B, Kässens JC, et al. NGS-based
IG/TR rearrangement profiling in acute lymphoblastic leukemia: age dependence of
immunogenetic maturation. Blood J. 2025 Mar 25;blood.2024027175.

9.   Darzentas F, Szczepanowski M, Kotrová M, Kelm M, Hartmann A, Beder T, et al. IGH
Rearrangement Evolution in Adult KMT2A-rearranged B-cell Precursor ALL: Implications
for Cell-of-origin and MRD Monitoring. HemaSphere. 2023 Jan;7(1):e820.

10.   Iacobucci I, Zeng AGX, Gao Q, Garcia-Prat L, Baviskar P, Shah S, et al. Multipotent
lineage potential in B cell acute lymphoblastic leukemia is associated with distinct
cellular origins and clinical features. Nat Cancer [Internet]. 2025 June 27 [cited 2025 July
25]; Available from: https://www.nature.com/articles/s43018-025-00987-2

11.   Khabirova E, Jardine L, Coorens THH, Webb S, Treger TD, Engelbert J, et al. Single-cell
transcriptomics reveals a distinct developmental state of KMT2A-rearranged infant B-cell
acute lymphoblastic leukemia. Nat Med. 2022 Apr;28(4):743–51.

12.   Chen C, Yu W, Alikarami F, Qiu Q, Chen C hui, Flournoy J, et al. Single-cell multiomics
reveals increased plasticity, resistant populations, and stem-cell–like blasts in *KMT2A* -
rearranged leukemia. Blood. 2022 Apr 7;139(14):2198–211.

13.   Yoshimura S, Li Z, Gocho Y, Yang W, Crews KR, Lee SHR, et al. Impact of Age on
Pharmacogenomics and Treatment Outcomes of B-Cell Acute Lymphoblastic Leukemia.
J Clin Oncol. 2024 Oct 10;42(29):3478–90.

14.   Huang X, Li Y, Zhang J, Yan L, Zhao H, Ding L, et al. Single-cell systems pharmacology
identifies development-driven drug response and combination therapy in B cell acute
lymphoblastic leukemia. Cancer Cell. 2024 Apr;42(4):552-567.e6.

15.   Bartram J, Ancliff P, Vora A. How I treat infant acute lymphoblastic leukemia. Blood.
2025 Jan 2;145(1):35–42.

16.   Van Der Sluis IM, De Lorenzo P, Kotecha RS, Attarbaschi A, Escherich G, Nysom K, et
al. Blinatumomab Added to Chemotherapy in Infant Lymphoblastic Leukemia. N Engl J
Med. 2023 Apr 27;388(17):1572–81.

17.   Stewart JP, Gazdova J, Darzentas N, Wren D, Proszek P, Fazio G, et al. Validation of
the EuroClonality-NGS DNA capture panel as an integrated genomic tool for
lymphoproliferative disorders. Blood Adv. 2021 Aug 24;5(16):3188–98.

18. Mihara K, Imai C, Coustan-Smith E, Dome JS, Dominici M, Vanin E, et al. Development
and functional characterization of human bone marrow mesenchymal cells immortalized
by enforced expression of telomerase. Br J Haematol. 2003 Mar;120(5):846–9.

19. Frismantas V, Dobay MP, Rinaldi A, Tchinda J, Dunn SH, Kunz J, et al. Ex vivo drug
response profiling detects recurrent sensitivity patterns in drug-resistant acute
lymphoblastic leukemia. Blood. 2017 Mar 16;129(11):e26–37.

20. Gu Z, Churchman ML, Roberts KG, Moore I, Zhou X, Nakitandwe J, et al. PAX5-driven
subtypes of B-progenitor acute lymphoblastic leukemia. Nat Genet. 2019 Feb;51(2):296–
307.

21. Walter W, Shahswar R, Stengel A, Meggendorfer M, Kern W, Haferlach T, et al. Clinical
application of whole transcriptome sequencing for the classification of patients with acute
lymphoblastic leukemia. BMC Cancer. 2021 Aug 2;21(1):886.

22. Saorin A, Dehler A, Galvan B, Steffen FD, Ray M, Lu D, et al. Transcriptional remodeling
shapes therapeutic vulnerability to necroptosis in acute lymphoblastic leukemia. Blood J.
2025 May 13;blood.2025028938.

23. Beder T, Hansen BT, Hartmann AM, Zimmermann J, Amelunxen E, Wolgast N, et al.
The Gene Expression Classifier ALLCatchR Identifies B-cell Precursor ALL Subtypes
and Underlying Developmental Trajectories Across Age. HemaSphere. 2023
Sept;7(9):e939.

24. Zeng AGX, Iacobucci I, Shah S, Mitchell A, Wong G, Bansal S, et al. Single-cell
Transcriptional Atlas of Human Hematopoiesis Reveals Genetic and Hierarchy-Based
Determinants of Aberrant AML Differentiation. Blood Cancer Discov. 2025 Apr 28;OF1–
18.

25. Burmeister T, Ströh AS, Kehden B, Trautmann H, Meyer C, Marschalek R, et al.
Measurable residual disease quantification in adult patients with KMT2A-rearranged
acute lymphoblastic leukemia. Leukemia. 2024 July;38(7):1600–3.

26. Kaspers GJL, Veerman AJP, Pieters R, Van Zantwijk CH, Smets LA, Van Wering ER, et
al. In Vitro Cellular Drug Resistance and Prognosis in Newly Diagnosed Childhood Acute
Lymphoblastic Leukemia. Blood. 1997 Oct 1;90(7):2723–9.

27. Enblad AP, Krali O, Gezelius H, Lundmark A, Blom K, Andersson C, et al. Ex Vivo Drug
Responses and Molecular Profiles of 597 Pediatric Acute Lymphoblastic Leukemia
Patients [Internet]. Oncology; 2024 [cited 2025 Mar 27]. Available from:
http://medrxiv.org/lookup/doi/10.1101/2024.12.17.24319138

28. Escherich G, Troger A, Gobel U, Graubner U, Pekrun A, Jorch N, et al. The long-term
impact of in vitro drug sensitivity on risk stratification and treatment outcome in acute
lymphoblastic leukemia of childhood (CoALL 06-97). Haematologica. 2011 June
1;96(6):854–62.

Figure Legends

Fig. 1 Age-overriding *KMT2A*r B-ALL patient cohort. A: Included patients (n=465) of different age groups (>=55 n=79, 18-54 n=207, 1.17 n=96, <1 n=63) from three cohorts (UKSH n=288, St. Jude(20) n=136, MLL(21) n=41). **B:** Distribution of common KMT2A fusions across age groups (n=422, 96.6%). **C**: Distribution of rare KMT2A fusions (n=15, 3.4%).

Fig. 2 Gene regulatory landscape of *KMT2A*r fusion partners. **A**: Heatmap of hierarchical clustering of fusion specific genes detected in supervised analysis using multi-comparison ANOVA and subsequent LASSO feature selection for genes separating *AFF1*r, *MLLT1*r and *MLLT3*r (n=148 samples, n=236 genes). Hierarchical clustering splits *AFF1*r and non-*AFF1*r patients, the second split subdivided non-*AFF1*r cases **B**. Mean gene Expression (norm. log2CPMs) per gene of *AFF1*r (y-axis) vs. non-*AFF1*r (x-axis) cases. Highlighted Genes represent genes that were detected by ANOVA and LASSO feature selection (panel A). Genes significantly higher in *AFF1*r cases at log2fc >1.5 are highlighted in orange (above diagonal), genes significantly higher expressed in non-*AFF1*r cases at log2fc >2.5 are highlighted in blue (below diagonal).

Fig. 3 Maturity in KMT2Ar ALL correlates with age and fusion partner. A: Gene set enrichment analysis of n=325 *KMT2A*r ALL cases. Enrichment score represents proximity to five healthy B-cell precursor developmental stages (pro-B, pre-B-I, pre-B-II-Large, pre-B-II-Small, Immature-B), calculated based on the RNAseq data using ALLCatchR tool(23). **B**: Developmental trajectories of *KMT2A*r ALL were condensed into a single maturity score ranging from -1 to 1 by fitting linear regression to the enrichments scores in A and calculating the negative slope. Low maturity scores represent immature transcriptional developmental state and proximity to Pro-B cells, high maturity scores represent more mature (pre-B-II, Immature-B) transcriptional developmental states. **C**. *KMT2A*r samples

were mapped to the expression matrix of the human single-cell B-cell developmental atlas(24) and colored by maturity score (right). Samples with low maturity score map to early cell stages (pro-B-V(D)J / pre-pro-B cycling), samples with higher maturity score map to mature B-cell stages (small Pre-B / Immature-B). **D**: Maturity score distribution between age groups within driver fusion groups (*AFF1*r n=221, *MLLT1*r n=43, *MLLT10*r n=14, *MLLT3*r n=28). Multi-comparison ANOVA between age groups within each fusion group reveals significantly different maturity scores between age groups in *AFF1*r (p=2.6e-05) and *MLLT1*r (p=0.0035) cases.

Fig. 4: MRD response is modulated by age, gene fusion and maturity score. **A**: Distribution of age within MRD clearance categories (MRD fast clearance: n=39, median age: 9.4 years; MRD intermediate clearance: n=53, median age: 32 years; MRD slow clearance: n=122, median age: 37 years). Age differs significantly higher in MRD slow clearance patients (multi-comparison anova, p=0.00021). **B**: Distribution of maturity scores within MRD clearance categories (MRD fast clearance: n=30; MRD intermediate clearance: n=29; MRD slow clearance: n=398). Maturity scores are significantly higher in MRD fast clearance patients (multi-comparison ANOVA, p=0.0013). **C**: Machine-learning trained decision tree, trained to predict MRD clearance categories based on age, fusion and maturity score. The highest-level split predicts patients with maturity score >-0.04 to have fast clearance, the second split predicts patients with maturity score <=-0.04 and *MLLT1*r to have fast MRD clearance, the third split predicts patients with maturity scores between >-0.28 and -0.04 into intermediate MRD clearance and patients with maturity scores <=-0.28 and no *MLLT1*r to have slow MRD clearance. **D** Univariate analysis of factors predicting MRD clearance (n=97). Categorical factors with multiple levels (age group infant, pediatric, adult/elderly and maturity score group <=-0.28 (immature), <=-0.04 and >-0.28, >-0.04 (mature) were tested per level against all others.

Fig. 5: MRD-driven gene expression signatures. A: Gene expression signature detected

from proportional odds linear regression model grouping patients by MRD response. Cluster

1 is enriched for patients with fast MRD clearance [(Cluster 1: n=12/18 (67%) fast, n=5/18

(28%) intermediate, n=1/18 (6%) slow; Cluster 3: n=1/28(4%) fast, n=7/28 (25%)

intermediate, n=20/28 (71%) slow. Chi-squared P<0.0001]). **B:** Resistance score (total score

from single sample gene set enrichment analysis using genes upregulated in Cluster 3 as

up-set and genes upregulated in Cluster 1 as down-set). **C**: Gene ontology annotation for

genes upregulated in MRD Cluster 3. Gene set was annotated with enriched GO terms

(GO:Biological Process). Next, enriched GO-terms were filtered for terms size <1000, terms

including at least five genes from the gene list and genes were filtered for genes present in

at least ten GO-terms. The resulting genes and terms were grouped by functional similarity

and five groups were summarized based on included GO-terms.

Fig 6. *Ex-vivo* drug response to induction therapy drug mirrors slow MRD clearance.

Drug response of *ex-vivo KMT2A*r samples (n=61). **A**: normalized log2 area under the curve

(norm.logAUC) for 27 compounds with at least n=40 cases measured(18,19). Top 6 rows

represent induction therapy drugs. Patients were grouped according to MRD clearance

(n=49) and annotated with resistance score from Figure 5B. **B**: Response (norm.logAUC) to

induction therapy drugs was poorer in patients with slow MRD clearance (n=17; red box)

compared to patients with fast MRD clearance (n=16; green box; mean norm. log AUC

across drugs: 0.61 (slow MRD) vs 0.52 (fast MRD), Wilcox-test per drug: Asparaginase

P=0.02, Dexamethasone P=0.0027, Doxorubicin P=0.041, Vincristine P=0.012, Cytarabine

and Daunorubicin not significant). **C**: Circos plot of patients (n=59) ordered and separated

into quartiles by their mean response (norm.logAUC) to induction phase drugs, and

annotated with maturity score, fusion partner, age group and MRD clearance. Quartiles 1

and 2 (Q1/Q2, n=15 respectively; most sensitive) were enriched for fast MRD clearance,

high maturity score, non-*AFF1*r and pediatric age. Quartiles 3 and 4 (Q3/Q4, n=15

respectively; most resistant) were enriched for slow MRD clearance, low maturity score,

*AFF1*r and older age (Chi-squared tests: MRD clearance P=0.003, maturity score group

P=0.064, fusion P=0.153, age group P=0.025, Supplementary Figure S7). **D**: Response to

induction phase drugs (left) and Venetoclax (right). Patients grouped by maturity score group

(cutoffs -0.04 and -0.28 based on cutoffs from trained decision tree (Figure 4C). Patients in

the lowest maturity score group (<=-0.28) are significantly more resistant to patients in the

highest maturity Score group (>-0.04) (Wilcox-test, p=0.0018). The effect is inverse for

Venetoclax (Wilcox-test p=0.18).

Figure 1

Figure 2

A

B

Figure 3

Figure 4

Figure 5

Figure 6

A

B

C

D

Supplementary

Figure S1

Figure S2

A

B

Figure S3

A

Fusion partner gene signature St Jude

B

Fusion partner gene signature MLL

Fusion
- KMT2A::AFF1
- KMT2A::MLLT1
- KMT2A::MLLT3
- KMT2A::MLLT10
- other

Age Group
- infant
- pediatric
- adult
- elderly

Figure S4

A

B

Figure S5

Figure S6

A

B

C

Figure S7

Figure S8

A

MI-503 Anova, p = 0.02
SNDX-5613 Anova, p = 0.47
VTP-50469 Anova, p = 0.19

logAUC

Fusion (AFF1, MLLT1, MLLT3)

B

MI-503 $R^2 = 0.12$, $P = 0.119$
SNDX-5613 $R^2 = 0.01$, $P = 0.634$
VTP-50469 $R^2 = 0.04$, $P = 0.403$

logAUC

Maturity Score

C

MI-503 Anova, p = 0.28
SNDX-5613 Anova, p = 0.12
VTP-50469 Anova, p = 0.58

logAUC

Age group (Infant, Pediatric, Adult)

D

MI-503 Anova, p = 0.79
SNDX-5613 Anova, p = 0.15
VTP-50469 Anova, p = 0.47

logAUC

MRD clearance (fast, slow)

Fusion
- KMT2A::AFF1
- KMT2A::MLLT1
- KMT2A::MLLT3

Maturity score
1.0 — -1.0

Age group
- infant
- pediatric
- adult
- elderly

MRD clearance
- slow
- intermediate
- fast