Buzzed but not elated? Effect of ethanol on cognitive judgement bias in honeybees

Names of authors

Marta Golańska, Faculty of Psychology, University of Warsaw, Warsaw, Poland 0000-0003-3305-4470

Weronika Antoł, Institute of Systematics and Evolution of Animals, Polish Academy of Sciences, Kraków, Poland 0000-0003-2047-330X

Magdalena Witek, Museum and Institute of Zoology, Polish Academy of Sciences, Warsaw, Poland 0000-0002-6172-8981

Krzysztof Miler, Institute of Systematics and Evolution of Animals, Polish Academy of Sciences, Kraków, Poland 0000-0001-7684-0629

Corresponding author

Marta Golańska

m.golanska@uw.edu.pl

Abstract

Ethanol is a neuroactive compound known to alter cognition, behavior, and physiology in a diverse range of vertebrate and invertebrate animals. It is also naturally present in many environments and might exert ecology-shaping influence on various organisms, such as honeybees feeding on fermenting nectar. In this study, we tested whether exposure of honeybees to low ethanol concentrations (1%) induces 'pessimistic' or 'optimistic' cognitive judgment bias (CJB), an information processing pattern thought to be related to affective processes. Bees were trained in a classical olfactory conditioning paradigm and then presented with conditioned stimuli and ambiguous odor mixtures to assess changes in decision-making under uncertainty. We additionally measured locomotor performance and ethanol absorption via spectrophotometric analysis of hemolymph. Ethanol levels peaked rapidly and remained elevated throughout the testing period. Despite this, ethanol had no detectable effect on choices in the CJB assay or locomotor behavior. These findings suggest that acute exposure to low-dose ethanol does not induce affect-like changes in honeybees. Our study represents the first test of pharmacologically induced judgment bias in an invertebrate and contributes to understanding the resilience of insect cognition to natural neuroactive compounds. The results highlight the limits and potential of CJB paradigms for probing affect-like states across taxa and support the honeybee as a valuable model in comparative cognition and the behavioral ecology of neuroactive substance exposure.

Keywords: ethanol, cognitive bias, affective state, decision-making, learning, *Apis mellifera*

1. Introduction

Ethanol is a neuroactive substance occurring naturally in many environments, and some species have evolved mechanisms to tolerate or utilize it, despite its toxicity (Bowland et al. 2025; Miler 2025). Insects, in particular, frequently encounter ethanol in fermenting plant material. For example, fruit flies (*Drosophila melanogaster*) are attracted to ethanol-rich substrates, using it as a food source and to maintain a competitive advantage over less tolerant species (Merçot et al. 1994). The oriental hornet (*Vespa orientalis*) shows exceptional ethanol tolerance, possibly a result of coevolution with yeasts (Bouchebti et al. 2024). Ambrosia beetles, known for their advanced sociality and symbiosis with fungi, are attracted by ethanol emissions and select the colony sites based on ethanol concentration in the sap of candidate trees (Cavaletto et al. 2021). In addition, the compound has been found to promote the growth of the beetle's nutritional symbiont (Lehenberger et al. 2021). These examples suggest that ethanol exposure can exert strong selective pressures and may have shaped behavioral and physiological adaptations across diverse insect taxa. Honeybees (*Apis mellifera*) also encounter ethanol in their environment, and it plays a significant role in their ecology (Abramson et al. 2025). Yeasts contribute to the fermentation processes involved in the bee bread production, a vital larval food source (Gilliam 1979; Detry et al. 2020 but see Anderson et al. 2014). Ethanol can also be present in flower nectar collected by foragers (e.g. Jakubska et al., 2005; Kevan et al., 1988; Schaeffer et al., 2017). Ethyl oleate, a pheromone produced from ethanol and oleic acid, regulates the division of labor in honeybee colonies (Castillo et al. 2012). Ethanol tolerance varies across worker castes, with higher resistance and alcohol dehydrogenase levels in foragers compared to workers staying inside the nest (Miler et al. 2021).

Much of what we know about the effect of ethanol on bee physiology, behavior, and cognition comes from studies that focus on honeybees as a potential model informing human alcohol addiction research. Honeybees have been found to prefer low ethanol concentrations in sucrose solutions (Abramson et al. 2000; Mustard et al. 2019) and show no conditioned taste aversion to ethanol

(Varnon et al. 2018). Chronic ethanol consumption produces tolerance (Miler et al., 2018; but see

Stephenson et al., 2021), and its discontinuation results in withdrawal-like symptoms in honeybees

(Ostap-Chec et al., 2021; Yang, Z et al., 2025). Ethanol intake in honeybees can be reduced with

disulfiram, a drug used in the treatment of alcohol dependence in humans (Abramson et al. 2003).

Ethanol impairs honeybee motor function (Abramson et al. 2000; Maze et al. 2006; Ahmed et al.

2022) and learning performance (Abramson et al. 2000; Mustard et al. 2008). Social behaviors like

trophallaxis or waggle and tremble dances are also affected (Bozic et al., 2006; Mixson et al., 2010). In

addition, ethanol has been found to increase aggression (Wright et al. 2012; Giannoni-Guzmán et al.

2014), although not all studies support this finding (Abramson et al. 2000). Importantly, most studies

use high ethanol concentrations (typically 2.5-10%), limiting their ecological relevance, as ethanol

levels naturally encountered by honeybees are likely much lower (Wiens et al. 2008; Jones and

Agrawal 2022).

Another limitation of the existing body of work is its limited scope in terms of the broadness of

the cognitive traits investigated, focusing mostly on classical conditioning and memory. In recent

years, more and more studies have uncovered the surprising cognitive complexity of honeybees, such

as contextual and social learning, including also abilities that were thought to be unique to

vertebrates, for example, metacognition, numerical cognition, conceptual learning, and attention-like

processes (Collett et al. 1997; Leadbeater and Chittka 2007; Pahl et al. 2013; Avarguès-Weber and

Giurfa 2013; Perry et al. 2017). One particularly interesting phenomenon is cognitive judgment bias

(CJB), which measures decision-making under uncertainty. The CJB paradigm offers a promising tool

for investigating affect-like processes as well as their evolutionary basis and function. Originally

developed for rodents (Harding et al. 2004), CJB tests assess whether animals show 'optimistic' or

'pessimistic' biases when interpreting ambiguous cues. It has been adapted for invertebrates,

including honeybees (Bateson et al. 2011). In these tests, animals are conditioned to associate two

stimuli with a rewarding stimulus, like food, and a punishment, like an unpalatable substance, and are

later presented with the learned stimuli and novel, ambiguous ones. Reactions to these ambiguous

cues might resemble the animal's responses to the rewarded or punished stimuli, suggesting

'optimistic' or 'pessimistic' biases, respectively. In honeybees, 'pessimistic' biases have been induced

by stressors (shaking; Bateson et al. 2011; Schlüns et al. 2017), and 'optimistic' biases have been

investigated in their relatives, bumblebees (Solvi et al. 2016; Strang and Muth 2023; Romero-

González et al. 2025). Changes in CJB are commonly interpreted as indicators of affect or emotion-like

changes, following the original formulation of the test (Harding et al. 2004; Mendl et al. 2009; Neville

et al. 2020; Lagisz et al. 2020, but see alternative interpretation by Baracchi et al. 2017). Whether

insects possess emotional states or their analogues remains highly debated (see Baracchi et al. 2017;

Perry and Baciadonna 2017). However, CJB responses in invertebrates provide a behavioral window

into valence-based processing. Following Mendl and Paul (2020), we adopt a broad definition of

'affect' that includes all valence-related processes ('pleasantness' or 'unpleasantness'), including

emotions, moods, and their possible manifestations (physiological or behavioral), whether or not

they entail conscious experience. This allows for comparisons across taxa without necessarily

implying that conscious experience must be involved.

Some pharmacological substances have been shown to impact CJB across several, mostly

mammalian, species (see Lagisz et al. 2020, for review and meta-analysis). Ethanol is a potential

pharmacological moderator in this context. It is known to introduce affective changes in humans,

acting as euphoriant in lower, anxiogenic in higher doses, and increasing aggression in some

individuals as well (Freed 1978; Chermack and Giancola 1997). Rodent studies largely find similar

patterns, mostly using traditional behavioral assays (Van Erp and Miczek 1997; Valdez et al. 2002).

Ethanol has never been tested as a mood-manipulating agent in a CJB test; however, one recent study

linked optimistic-like responses to ethanol self-administration in rats (Cieslik-Starkiewicz et al. 2024).

Moreover, no study has tested whether ethanol administration can alter CJB in invertebrates,

including those that are exposed to it under natural conditions.

Because ethanol is a neuroactive substance, even species with relatively high ecological

tolerance may still exhibit transient changes in decision-making or motivational processes upon

exposure. Testing this provides insight into the robustness of insect cognition to naturally occurring

neuroactive compounds. In the present study, we investigated whether a low concentration (1%) of

ethanol can induce CJB in honeybees. We hypothesized that ethanol-exposed bees would exhibit

bias, that is, respond differently to at least one ambiguous cue compared to controls, while

maintaining normal responses to learned stimuli. Additionally, we assessed locomotor behavior and

ethanol absorption via hemolymph sampling to rule out non-specific performance effects and

confirm effective dosing.

2.   Methods

2.1. Cognitive judgment bias (CJB) test

124   2.1.1. Study species and collection procedure

Bees were collected from three families from the experimental apiary located in Kraków, Poland, over

16 days. Animals were collected in one or two blocks daily between 07:30 am and 12:25 pm, with

each collection lasting up to 15 minutes. During the procedure, entrance to the hive was partially

blocked by a foam barrier, allowing the experimenter to select and capture congregating bees. Only

pollen-carrying individuals were chosen to ensure their forager status. Up to 30 bees were restrained

per block. Bees were left for one hour to acclimate and strengthen feeding motivation. From each

block, 12 bees were selected for the conditioning phase based on their overall condition, absence of

proboscis or antennal defects, adequate restraint, and prompt proboscis extension upon brief

antennal stimulation with sucrose solution. The remaining bees were released (see Supplementary

Information 1 for a visualization of the exclusion process).

135   2.1.2. Olfactory conditioning

Selected bees underwent olfactory conditioning using the proboscis extension response (PER)

paradigm (Scheiner et al. 2013). This classical conditioning paradigm utilizes a reflexive extension of

the proboscis in response to sucrose contact and is widely used for appetitive conditioning in

honeybees (Giurfa and Sandoz 2012). Although PER is a reflex, it is known to be modulated by many

factors, including motivational ones (Menzel, R 1990), and it has been successfully used in bee CJB

protocols before (Bateson et al. 2011; Schlüns et al. 2017). Here, the standard appetitive conditioning

protocol also included discrimination learning with NaCl as an aversive (de Brito Sanchez et al., 2015).

One odor (2-octanone, >99%, Thermo Scientific) was paired with an appetitive unconditioned

stimulus (0.5 μl drop of 1.5 M aqueous sucrose solution; hereafter US^+), while a second odor (1-

hexanol, >98%, Tokyo Chemical Industry) was paired with an aversive unconditioned stimulus (0.5 μl

drop of 0.5 M aqueous NaCl solution; US^-). Each odor was delivered from a 24 ml syringe containing a

cotton pad soaked with 6 μl of the odorant. The end of the syringe was placed about 0.5 cm from the

bee's head, and air was expelled for 4 seconds. In the final second of odor delivery, US^+ or US^- was

presented, lasting 3 seconds in total (Matsumoto et al. 2012). The antennae were stimulated with the

solution, and if PER was elicited, a drop was placed on the proboscis using a pipette.

Each bee underwent 16 conditioning trials (8 per odor) in a fixed sequence (CS^+-CS^+-CS^--CS^+-CS^--

CS^+-CS^+-CS^--CS^--CS^+-CS^--CS^--CS^+-CS^--CS^+-CS^-; as used by Schlüns and colleagues (2017), where CS^+ and

CS^- stand for the odors associated with the US^+ (sugar solution) and US^- (salt solution), respectively.

Bee responses during the first 3 seconds of odor presentation (before US delivery) were recorded as

1: full PER, 0: no PER, or partial extension of the proboscis. Inter-trial intervals were 6 minutes. Bees

were included in the experiment if they displayed correct responses in at least four of the final six

trials for each odor. Lack of PER to hexanol and full PER to octanone were considered correct

responses.

2.1.3. Group assignment and experimental manipulation

Bees that met the learning criteria were randomly assigned to either the control group (fed 10 μl of

1.5 M aqueous sucrose solution) or the ethanol group (fed 10 μl of 1.5 M aqueous sucrose solution

with 1% ethanol). For each block, group sizes were balanced wherever possible, or when the number

of bees was odd, the more numerous group was chosen randomly. The individuals were then

randomly assigned to the groups. Successfully fed bees were left for one hour before testing to allow

ethanol absorption and improve feeding motivation. Bees not meeting the learning criteria were

marked to prevent reuse and released.

2.1.4. Testing procedure

The successfully fed bees participated in the CJB test comprising five odor presentations (testing

trials). Each odor was presented for four seconds, and PER responses were coded as 1: full PER and 0:

lack of PER or partial response. No US were delivered, but a pipette was held nearby to mimic the

conditioning context. The odors included the two conditioned odors (6:0 octanone:hexanol =

positive; 0:6 = negative) and three mixtures: 4:2 (near-positive), 3:3 (neutral), and 2:4 (near-

negative). Odors were delivered as in the conditioning phase. The experimenter was blind to the odor

and the testing order, which was randomized per block.

Each testing trial was preceded by two reminder trials (US^+, US^-) to limit extinction of the

conditioned responses (Sandoz and Pham-Delègue 2004). The sequence was RT^+-RT^--**T1**-RT^+-RT^--**T2**-

RT^+-RT^--**T3**- RT^+-RT^--**T4**-RT^+-RT^--**T5** (RT^+ and RT^- = reminder trials with appetitive and aversive US paired

with the conditioned odors; **T1-5** = testing trials). The full procedure lasted 44 minutes, with a 3-

minute inter-trial interval. After testing, bees were marked and released, and odor identities were

unblinded.

2.2. Locomotion test

Bees were collected from seven families from the same apiary over nine days (between 07:00 am and

09:30 am, in two blocks), with one bee collected per minute for up to 30 minutes. Only pollen-

carrying individuals were selected, using the same method as described above, and they were

restrained promptly after being caught. Bees were left for three hours to acclimate and approximate

the conditioning phase duration in the CJB test. The first 15 and last 15 bees of each block were

assigned to the ethanol-fed or control group, with group assignment counterbalanced across families.

Bees were fed 10 µl of either 1.5 M aqueous sucrose solution (control) or the same solution

with 1% ethanol (ethanol-fed). Animals were excluded based on visible leg or mouthpart defects,

poor condition, incomplete consumption, or mortality (for the full inclusion/exclusion process for the

individual bees participating in the test, see Supplementary Information 1). Fifty minutes post-

ingestion, bees were released from restraint and placed under a Petri dish (14.5 cm diameter, 1 cm

high) for 10 minutes to acclimate. One hour after feeding, a 20-minute video of their locomotion

inside the dish was recorded. Videos were processed using AnimalTA (Chiara and Kim 2023) to extract

the total distance travelled, average speed while moving, and meander (average of the change in

direction, or turning angle, divided by the distance travelled for each frame).

2.3. Ethanol levels in hemolymph

Bees were collected from four families from the same apiary over six days. The animals were

collected for one hour, starting at 09:00 am, and assigned to one of four ethanol-fed groups, 15

individuals per group. Only pollen-carrying bees were selected (using the method described in

previous sections), then restrained and fed 10 µl of 1.5 M sucrose solution containing 1% ethanol.

Four ethanol-fed groups differed in the time of hemolymph extraction after feeding (30, 60, 120, or

240 minutes). An additional control group (N = 15) received a sucrose-only solution and was sampled

30 minutes post-ingestion.

All bees were restrained for two hours before feeding to acclimate and mimic the CJB test

procedure. The animals that did not ingest the entire solution were not used (again, for the full

inclusion/exclusion process for the individual bees participating in the test, see Supplementary

Information 1). Hemolymph was extracted using the antennal method (Borsuk et al. 2017) and

collected in end-to-end 10 µl microcapillaries, then frozen until analysis.

Samples were processed using the Ethanol Assay Kit, Megazyme, following the manufacturer's

protocol with some modifications. This method relies on spectrophotometric detection of NADH, a

product of two enzymatic reactions involving ethanol as a substrate. For each reaction, we used the

minimal volume of 3 μl of hemolymph from two samples pooled within a group, or, if the volume was

sufficient, from a single individual. The absorbance at 340 nm was measured twice: before (A1) and 5

min after adding the enzyme initiating the reaction (A2). The difference A2-A1 was used to calculate

ethanol concentration in the sample, based on the calibration curve with blank, performed with each

reaction plate.

2.4. Statistical analysis

All analyses in our experiments were performed using *R* (R Core Team, 2014). Model diagnostics were

conducted using the packages performance (Lüdecke et al. 2021) and DHARMA (Hartig, 2024), and

the emmeans package was used to obtain marginal means (Lenth, R and Piaskowski, J 2025).

2.4.1. CJB test

We analyzed PER responses (1 = full PER; 0 = absent/partial PER) using generalized mixed-effects

binomial models. Odor stimuli consisted of discrete mixtures (0, 2, 3, 4, 6 parts octanone), and so we

first compared a model assuming a strictly linear effect of octanone proportion with a model treating

odor as a categorical predictor. This comparison indicated that the linear model fit worse, suggesting

nonlinearity in the relationship between odor mixture and PER probability. We therefore used a

generalized additive mixed model (GAMM; mgcv package; Wood, S. N., 2017) with a smooth term for

odor mixture and a random-effect smooth for individual bee ID to account for repeated measures.

The model included group (ethanol-treated vs. control) as a fixed effect and a common smooth for

odor mixture. To test judgment bias, we compared this model to a model allowing group-specific

smooths using ANOVA. Model-estimated marginal means and 95% confidence intervals at the tested

odor mixtures were obtained from the fitted model for visualization and contrasts. A total of 736

observations from 148 bees were analyzed. Data for one individual was discarded due to

experimenter error. We additionally explored models that included date, hour, odor presentation

order, and family (as fixed effects), but these covariates were all non-significant and did not improve

model fit and were excluded from consideration. As a robustness check, we fitted the GAMM

described above on the data with a variant coding of PER (1 = full or partial PER, 0 = absent PER) in

order to verify whether decisions regarding coding of the outcome variable meaningfully impact the

result.

For analyzing learning trials, we considered responses from all 274 bees participating in the

learning procedure, for a total of 4230 trials. In order to analyze conditioning acquisition, we used a

binomial generalized linear mixed model with a logit link (package lme4; Bates, D et al., 2015). PER (1

– full PER, 0 – absent/partial PER) was modeled as a function of conditioned stimulus type (CS^+ vs. CS^-

), conditioning trial pair (1-8, treated as a categorical factor), and their interaction. Individual bee ID

was included as a random intercept to account for repeated measurements. Model-based estimated

marginal means and 95% confidence intervals were obtained.

2.4.2. Locomotion test

We analyzed the effect of ethanol on three locomotor variables: distance travelled while moving,

average speed while moving, and meander (change in direction per unit distance). Movement was

defined using a threshold of 0.3 cm/s to minimize the influence of tracking imperfections, with

perspective correction and default smoothing applied (see Chiara and Kim 2023 for explanations of

the parameters). Data from four individuals were excluded due to poor physical condition. Average

speed while moving was analyzed using a linear model (package nlme; Pinheiro, B, 2000) with

treatment group (control vs. ethanol-fed) as a predictor. Distance travelled while moving was

analyzed using a linear model fitted with generalized least squares on log-transformed distances with

group-specific residual variance to account for heteroscedasticity. Meander was analysed using a

linear model on log-transformed values. Model estimates were obtained with 95% confidence

intervals. The final sample size was 280 individuals.

2.4.3. Ethanol levels in hemolymph

Ethanol concentrations in hemolymph were analyzed using a linear model fitted to log-transformed

values after applying a $\log(1 + x)$ transformation to accommodate zero values arising from

measurements below the detection limit. Sampling group (control, 30-, 60-, 120-, or 240-minute post-

ingestion) was included as a fixed effect. Family, assay plate, and the number of individuals pooled

per sample were included as covariates to account for noise. Model-based estimated marginal means

were computed for each group, and contrasts were used to compare ethanol-fed groups to the

control as well as to assess differences among ethanol-fed time points. Model-based estimates were

obtained with 95% confidence intervals. In total, 55 hemolymph samples from 105 individuals were

analyzed.

3. Results

Responses in the cognitive judgment bias (CJB) test were strongly influenced by odor mixture

composition (Figure 1). In both control and ethanol-fed bees, the probability of PER increased

nonlinearly with increasing proportion of the rewarded odor (GAMM smooth terms: control: χ^2 =

65.53, p < 0.001, ethanol-fed: χ^2 = 84.95, p < 0.001). Ethanol treatment had no significant main effect

on PER probability (estimate = −0.27 ± 0.42 SE, z = -0.66, p = 0.51). Importantly, ethanol did not alter

the shape or slope of the odor-response function, as indicated by the absence of group-specific

differences in the nonlinear odor smooths (χ^2 = 5.78, p = 0.22). Responses to ambiguous odor

mixtures (near-positive, neutral, and near-negative cues) were indistinguishable between treatment

groups, providing no evidence for ethanol-induced CJB. For the three ambiguous mixtures, the odds-

ratio confidence intervals were 2:4 (OR = 0.90, 95% CI [0.28, 2.95]), 3:3 (OR = 0.98, 95% CI [0.39,

2.42]), and 4:2 (OR = 0.91, 95% CI [0.38, 2.19]), incompatible with meaningful ethanol-induced shifts

in ambiguous-cue responding, although modest effects remained possible. No differences in learned

odors suggested no post-ingestion changes in memory or proboscis mobility (see Supplementary

Information 2 for all contrasts). The model explained 51.6% of the deviance in PER responses. No

major differences in the model parameters with variant outcome variable coding were detected (see

Supplementary Information 2).

Figure 1. **Predicted probability of PER across odor mixtures in the CJB test**. Curves represent

predictions from a binomial generalized additive mixed model, with shaded bands indicating 95%

confidence intervals. The x-axis shows the proportion of octanone (appetitive CS) in the odor

stimulus, including conditioned cues (6:0 and 0:6) and ambiguous mixtures (4:2, 3:3, 2:4). Both

groups show similar nonlinear odor-response functions, with no evidence of ethanol-induced shifts in

responses to ambiguous cues.

In the locomotor activity test, control and ethanol-fed bees showed similar performance across

the measured parameters (Figure 2). Average speed while moving did not differ between groups

(control: 3.30 cm/s [95% CI 3.13, 3.47]; ethanol-fed: 3.19 cm/s [95% CI 3.02, 3.36]; t = -0.88, p = 0.38).

For distance travelled while moving, there was a non-significant tendency for ethanol-fed bees to

travel less than controls (control: 3470 cm [95% CI 3193, 3772]; ethanol-fed: 3018 cm [95% CI 2654,

3432]; t = -1.80, p = 0.07). Meander while moving also did not differ between groups (control: 227

deg/cm [95% CI 214, 240]; ethanol-fed: 229 deg/cm [95% CI 216, 242]; t = 0.20, p = 0.84).

Figure 2. **Locomotor activity in control and ethanol-fed honeybees.** Raw individual values are shown

as semi-transparent points; larger points and error bars indicate model-estimated marginal means ±

95% confidence intervals. Speed while moving was analyzed with a linear model. Distance travelled

while moving was analyzed using generalized least squares on log-transformed distances with group-

specific residual variance. Meander while moving was analyzed with a linear model on log-

transformed values. Ethanol treatment did not affect speed, showed a non-significant tendency

toward reduced distance, and did not affect meander.

Hemolymph ethanol concentrations differed among sampling groups as a function of time

since ingestion ($F_{4,45}$ = 13.59, p < 0.001; Figure 3). Compared to control bees, ethanol-fed ones

showed significantly elevated ethanol concentrations at 30 minutes (estimate = 0.65, SE = 0.09, t =

6.98, p < 0.001), 60 minutes (estimate = 0.49, SE = 0.09, t = 5.24, p < 0.001), and 120 minutes

(estimate = 0.40, SE = 0.11, t = 3.74, p = 0.005) post-ingestion. At 240 minutes after ingestion, ethanol

concentration was only slightly elevated (estimate = 0.30, SE = 0.12, t = 2.59, p = 0.090). No additional

effects of family, assay plate, or pooling of individuals were detected (see Supplementary Information

2 for full results and contrasts). Together, these results indicate that hemolymph ethanol levels

remained elevated throughout the behavioral testing period (30-120 minutes) and declined toward

baseline within four hours of ingestion.

Figure 3. **Hemolymph ethanol concentration following ingestion of ethanol-containing sucrose**

solution. Bees were sampled at 30, 60, 120, and 240 minutes after feeding on a 1% ethanol solution;

control bees received sucrose without ethanol and were sampled 30 minutes after ingestion. Semi-

transparent points represent individual hemolymph samples. Larger points and error bars indicate

model-estimated marginal means ± 95% confidence intervals from a linear model fitted to log-

transformed ethanol concentrations, adjusted for family, assay plate, and pooling of individuals.

Different letters indicate significant differences among sampling groups based on Tukey-adjusted

pairwise comparisons. Ethanol concentrations were elevated at 30-120 minutes post-ingestion and

approached control levels by 240 minutes.

Analysis of responses during the conditioning phase confirmed effective acquisition. There was a

significant main effect of conditioned stimulus type (CS$^+$ vs. CS$^-$; χ^2 = 7.94, p = 0.0048), a strong effect

of conditioning trial pair (χ^2 = 274.56, p < 0.001), and a pronounced CS type × trial interaction (χ^2 =

150.95, p < 0.001). PER probability increased sharply across successive trial pairs for the rewarded

odor (CS$^+$), while responses to the punished odor (CS$^-$) remained low and diverged progressively from

CS$^+$ across training. This pattern indicates robust associative learning and reliable discrimination

between conditioned stimuli across the conditioning sequence (see Supplementary Information 2 for

predicted probabilities of PER across conditioning and Figure S1).

4. Discussion

Our study investigated whether ethanol exposure influences cognitive judgment bias (CJB) in

honeybees. We found no evidence that ethanol affected bees' responses to ambiguous cues,

suggesting an absence of judgment bias. Additionally, we observed no detectable influence on bees'

responses to the conditioned stimuli themselves, and no clear impairment of general motor function.

Ethanol ingestion and absorption were confirmed by spectrophotometric analysis of hemolymph

samples, which showed elevated levels during the behavioral testing window. Together, these

findings suggest that acute exposure to low doses of ethanol does not produce detectable changes in

judgment bias within a PER-based CJB paradigm in honeybee.

Our findings contrast with earlier investigations using similar paradigms, which have

demonstrated that invertebrates can show affect-like states. For instance, Schlüns et al (2017) found

that honeybees exposed to a physical stressor (shaking) exhibit pessimistic biases (building on the

pioneering, but less methodologically clear report by Bateson et al., 2011). CJB has also been

described in bumblebees (Solvi et al. 2016; Strang and Muth 2023; Procenko et al. 2024; Romero-

González et al. 2025) and fruit flies (Deakin et al. 2018). To our knowledge, however, this is the first

study to investigate whether pharmacological intervention may induce CJB in an invertebrate. The

absence of an effect is compatible with two major non-exclusive interpretations: that ethanol may

not reliably modulate affect-like processing in bees at this dose and time scale, or that the PER-based

CJB paradigm may be relatively insensitive to ethanol-induced state changes. We considered whether

the absence of CJB could reflect insufficient ethanol absorption, but this is unlikely. As in earlier

studies (Bozic et al. 2006; Maze et al. 2006), ethanol was absorbed rapidly and persisted in

hemolymph during the behavioral testing period. Conditioning dynamics also indicated robust

acquisition and discrimination, supporting the effectiveness of the learning procedure.

Dose and exposure regime are plausible explanations. Ethanol's influence may not manifest at

low exposure, even though 1% ethanol has been reported to affect honeybee behavior in some

contexts (e.g. Bozic et al. 2006; Ahmed et al. 2022). Our own data suggests a trend for a change in

gross locomotor behavior after 1% ethanol consumption. Higher concentrations might induce

measurable bias, and further investigation is warranted, but such studies should also account for the

known confounding effects of ethanol on learning and motor performance. From an ecological and

apicultural perspective, exploring a wider ethanol range is relevant because although nectar in

temperate climates is unlikely to reach high ethanol concentrations, bees may encounter high doses

in particular environments, especially human-modified ones such as orchards where fermenting fruits

are available (Ostap-Chec et al., 2025). It remains unclear how often and under what circumstances

bees supplement their diet with fallen fruits (Abramson et al. 2025) and whether this exposure

meaningfully affects their decision-making or welfare. Changes to affect and decision-making might

also only be apparent after chronic exposure, instead of a single acute dose, as in our study. Another

major possible explanation is that the binary measure of PER might not be sensitive enough to detect

the change in behavior, and newer alternatives, such as quantifying the strength of the proboscis

response (see Giurfa & Sandoz, 2012 for discussion), might yield different results. PER is a reflex, and

its suitability for detecting subtle affect-related modulation may be generally questioned.

Additionally, inter-individual variability may have obscured a small existing effect (Van Der Goot et al.

2021).

Our findings are consistent with the broader idea that honeybees may be relatively robust to

low ethanol doses present in their environment. Natural ethanol production dates back to the late

Cretaceous, when yeast gained the ability to ferment sugars in fleshy fruits of early angiosperms

(Benner et al. 2002). Selection could plausibly favor physiological mechanisms that mitigate ethanol's

deleterious neural effects, particularly in foragers that are most exposed to the substance. In fact,

they are known to be the most ethanol-resistant caste and possess elevated alcohol dehydrogenase

levels (Miler et al. 2021). If bees have evolved physiological mechanisms that buffer ethanol's neural

impact, then affect-related cognitive processes might be comparatively resistant to low doses, which

could help explain the absence of detectable bias in our paradigm despite sustained hemolymph

ethanol levels. The persistence of ethanol in hemolymph is likely due to the gradual passage of food

from its storage in the crop to the stomach, where it is digested (Crailsheim 1988). This makes low-

dose robustness potentially consequential: even a single ethanol-containing meal could influence

bees throughout a foraging bout, a vital but risky task that poses challenges to successful navigation,

learning, and predator avoidance. However, despite ethanol's potent, cross-species impact on

neurotransmission (Phillips and Shent 1996), our results suggest that many low-dose behavioral

consequences, if present, are hardly detectable.

Our findings help reconcile mixed reports in the literature. Some previous studies align with

our findings, showing no effect of ethanol on bee learning (Abramson et al. 2000) or flight

performance (Ostap-Chec et al. 2025). However, other works report behavioral changes even at 1%

ethanol, including altered flight kinematics (Ahmed et al. 2022), delayed return to the hive, reduced

trophallaxis and self-grooming, as well as changes in social communication (Bozic et al. 2006). One

possibility is that low concentrations of ethanol impact aspects of body movement that bees can

compensate for, making the effects undetectable at coarser scales (e.g. speed). Another possibility is

that some consequences of ingesting low ethanol concentrations are only apparent on the group

level and might reflect collective decision-making, for example, to reduce foraging when many

individuals carry ethanol-containing food at the same time. For instance, Bozic and colleagues (2006)

reported effects of colony-level inebriation when multiple foragers consumed ethanol

simultaneously. Future studies should explore how low-dose ethanol concentrations influence social

dynamics and collective behavior in honeybees.

5. Conclusion

This study investigated the impact of ethanol on cognitive judgment bias in honeybees. We show that

acute exposure to 1% ethanol produces no detectable judgment bias in honeybees when CJB is

assessed through binary PER responses to ambiguous odor mixtures and does not produce any major

changes in gross locomotor performance. Ethanol presence was confirmed in the hemolymph for

prolonged periods post-ingestion. We propose that honeybees may be behaviorally and

physiologically adapted to tolerate low ethanol levels. Our findings highlight the need to consider

both evolutionary adaptations and methodological sensitivity when evaluating affect-like states in

insects. More sensitive continuous PER measures, explicit assays of sucrose responsiveness and

motivation, and complementary neural or physiological markers will be essential to determine

whether low-dose ethanol modulates arousal or affect-like processing in bees in other ways than

tested here.

6. Funding information

This work was supported by the Faculty of Psychology, University of Warsaw, from the funds awarded

by the Ministry of Science and Higher Education in the form of a subsidy for the maintenance and

development of research potential in 2024 (501-D125-01-1250000 zlec. 501 1000 191). The research

was also supported by the National Science Centre, Poland [grant number Sonata UMO-

2021/43/D/NZ8/01044].

7. Compliance with Ethical Standards

No approval of research ethics committees was required to accomplish the goals of this study,

because experimental work was conducted with an unregulated invertebrate species.

8. Supplementary Information

Supplementary Information 1. Diagram for exclusion/inclusion rationale.

Supplementary Information 2. Tables with additional results.

9. References

Abramson CI, d'Isa R, Wells H (2025) Physiological and behavioral pharmacology of ethanol in honey bees. J Comp Physiol A. https://doi.org/10.1007/s00359-025-01743-8

Abramson CI, Fellows GW, Browne BL, et al (2003) Development of an Ethanol Model Using Social Insects: II. Effect of Antabuse® on Consumatory Responses and Learned Behavior of the Honey Bee (*Apis Mellifera* L.). Psychol Rep 92:365–378. https://doi.org/10.2466/pr0.2003.92.2.365

Abramson CI, Stone SM, Ortez RA, et al (2000) The Development of an Ethanol Model Using Social Insects I: Behavior Studies of the Honey Bee (*Apis mellifera* L.). Alcohol Clin Exp Res 24:1153–1166. https://doi.org/10.1111/j.1530-0277.2000.tb02078.x

Ahmed I, Abramson CI, Faruque IA (2022) Honey bee flights near hover under ethanol-exposure show changes in body and wing kinematics. PLOS ONE 17:e0278916. https://doi.org/10.1371/journal.pone.0278916

Anderson KE, Carroll MJ, Sheehan T, et al (2014) Hive-stored pollen of honey bees: many lines of evidence are consistent with pollen preservation, not nutrient conversion. Mol Ecol 23:5904–5917. https://doi.org/10.1111/mec.12966

Avarguès-Weber A, Giurfa M (2013) Conceptual learning by miniature brains. Proc R Soc B Biol Sci 280:20131907. https://doi.org/10.1098/rspb.2013.1907

Baracchi D, Lihoreau M, Giurfa M (2017) Do Insects Have Emotions? Some Insights from Bumble Bees. Front Behav Neurosci 11:157. https://doi.org/10.3389/fnbeh.2017.00157

Bates, D, Maechler, M, Bolker, B, Walker, S (2015) Fitting Linear Mixed-Effects Models Using lme4. J Stat Softw 67:1–48. https://doi.org/doi:10.18637/jss.v067.i01

Bateson M, Desire S, Gartside SE, Wright GA (2011) Agitated Honeybees Exhibit Pessimistic Cognitive Biases. Curr Biol 21:1070–1073. https://doi.org/10.1016/j.cub.2011.05.017

Benner SA, Caraco MD, Thomson JM, Gaucher EA (2002) Planetary Biology--Paleontological, Geological, and Molecular Histories of Life. Science 296:864–868. https://doi.org/10.1126/science.1069863

Borsuk G, Ptaszyńska AA, Olszewski K, et al (2017) A New Method for Quick and Easy Hemolymph Collection from Apidae Adults. PLOS ONE 12:e0170487. https://doi.org/10.1371/journal.pone.0170487

Bouchebti S, Gershon Y, Gordin A, et al (2024) Tolerance and efficient metabolization of extremely high ethanol concentrations by a social wasp. Proc Natl Acad Sci 121:e2410874121. https://doi.org/10.1073/pnas.2410874121

Bowland AC, Melin AD, Hosken DJ, et al (2025) The evolutionary ecology of ethanol. Trends Ecol Evol
S0169534724002404. https://doi.org/10.1016/j.tree.2024.09.005

Bozic J, Abramson CI, Bedencic M (2006) Reduced ability of ethanol drinkers for social communication
in honeybees (Apis mellifera carnica Poll.). Alcohol 38:179–183.
https://doi.org/10.1016/j.alcohol.2006.01.005

Castillo C, Chen H, Graves C, et al (2012) Biosynthesis of ethyl oleate, a primer pheromone, in the
honey bee (Apis mellifera L.). Insect Biochem Mol Biol 42:404–416.
https://doi.org/10.1016/j.ibmb.2012.02.002

Cavaletto G, Faccoli M, Ranger CM, Rassati D (2021) Ambrosia beetle response to ethanol
concentration and host tree species. J Appl Entomol 145:800–809.
https://doi.org/10.1111/jen.12895

Chermack ST, Giancola PR (1997) The relation between alcohol and aggression: An integrated
biopsychosocial conceptualization. Clin Psychol Rev 17:621–649.
https://doi.org/10.1016/S0272-7358(97)00038-X

Chiara V, Kim S (2023) ANIMALTA: A highly flexible and easy-to-use program for tracking and analysing
animal movement in different environments. Methods Ecol Evol 14:1699–1707.
https://doi.org/10.1111/2041-210X.14115

Cieslik-Starkiewicz A, Piksa M, Noworyta K, et al (2024) Unveiling the power of optimism: Exploring
behavioral and neuromolecular correlates of alcohol seeking and drinking in rats with biased
judgement. Prog Neuropsychopharmacol Biol Psychiatry 135:111124.
https://doi.org/10.1016/j.pnpbp.2024.111124

Collett TS, Fauria K, Dale K, Baron J (1997) Places and patterns -- a study of context learning in
honeybees. J Comp Physiol [A] 181:343–353. https://doi.org/10.1007/s003590050120

Crailsheim K (1988) Regulation of food passage in the intestine of the honeybee (Apis mellifera L.). J
Insect Physiol 34:85–90. https://doi.org/10.1016/0022-1910(88)90158-8

Deakin A, Mendl M, Browne WJ, et al (2018) State-dependent judgement bias in *Drosophila*:
evidence for evolutionarily primitive affective processes. Biol Lett 14:20170779.
https://doi.org/10.1098/rsbl.2017.0779

Detry R, Simon-Delso N, Bruneau E, Daniel H-M (2020) Specialisation of Yeast Genera in Different
Phases of Bee Bread Maturation. Microorganisms 8:1789.
https://doi.org/10.3390/microorganisms8111789

Freed EX (1978) Alcohol and Mood: An Updated Review. Int J Addict 13:173–200.
https://doi.org/10.3109/10826087809039273

Giannoni-Guzmán MA, Giray T, Agosto-Rivera JL, et al (2014) Ethanol-Induced Effects on Sting
Extension Response and Punishment Learning in the Western Honey Bee (Apis mellifera).
PLoS ONE 9:e100894. https://doi.org/10.1371/journal.pone.0100894

Gilliam M (1979) Microbiology of pollen and bee bread: the yeasts. Apidologie 10:43–53.
https://doi.org/10.1051/apido:19790106

Giurfa M, Sandoz J-C (2012) Invertebrate learning and memory: Fifty years of olfactory conditioning
of the proboscis extension response in honeybees. Learn Mem 19:54–66.
https://doi.org/10.1101/lm.024711.111

Harding EJ, Paul ES, Mendl M (2004) Cognitive bias and affective state. Nature 427:312–312.
https://doi.org/10.1038/427312a

Hartig, F (2024) DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression
Models. R Package Version 047. https://github.com/florianhartig/dharma

Jakubska A, Prządo, D, Steininger, M, et al (2005) Why do Pollinators Become "sluggish"? Nectar
Chemical C Onstituents from Epipactis Helleborine (L.) Crantz (orchidaceae). Appl Ecol
Environ Res 3:29–38. https://doi.org/10.15666/aeer/0302_029038

Jones P, Agrawal AA (2022) Caffeine and ethanol in nectar interact with flower color impacting
bumblebee behavior. Behav Ecol Sociobiol 76:103. https://doi.org/10.1007/s00265-022-
03208-3

Kevan PG, Eisikowitch D, Fowle S, Thomas K (1988) Yeast-Contaminated Nectar and its Effects on Bee
Foraging. J Apic Res 27:26–29. https://doi.org/10.1080/00218839.1988.11100777

Lagisz M, Zidar J, Nakagawa S, et al (2020) Optimism, pessimism and judgement bias in animals: A
systematic review and meta-analysis. Neurosci Biobehav Rev 118:3–17.
https://doi.org/10.1016/j.neubiorev.2020.07.012

Leadbeater E, Chittka L (2007) Social Learning in Insects — From Miniature Brains to Consensus
Building. Curr Biol 17:R703–R713. https://doi.org/10.1016/j.cub.2007.06.012

Lehenberger M, Benkert M, Biedermann PHW (2021) Ethanol-Enriched Substrate Facilitates Ambrosia
Beetle Fungi, but Inhibits Their Pathogens and Fungal Symbionts of Bark Beetles. Front
Microbiol 11:590111. https://doi.org/10.3389/fmicb.2020.590111

Lenth, R, Piaskowski, J (2025) emmeans: Estimated Marginal Means, aka Least-Squares Means

Lüdecke D, Ben-Shachar M, Patil I, et al (2021) performance: An R Package for Assessment,
Comparison and Testing of Statistical Models. J Open Source Softw 6:3139.
https://doi.org/10.21105/joss.03139

Matsumoto Y, Menzel R, Sandoz J-C, Giurfa M (2012) Revisiting olfactory classical conditioning of the
proboscis extension response in honey bees: A step toward standardized procedures. J
Neurosci Methods 211:159–167. https://doi.org/10.1016/j.jneumeth.2012.08.018

Maze IS, Wright GA, Mustard JA (2006) Acute ethanol ingestion produces dose-dependent effects on
motor behavior in the honey bee (Apis mellifera). J Insect Physiol 52:1243–1253.
https://doi.org/10.1016/j.jinsphys.2006.09.006

Mendl M, Burman OHP, Parker RMA, Paul ES (2009) Cognitive bias as an indicator of animal emotion
and welfare: Emerging evidence and underlying mechanisms. Appl Anim Behav Sci 118:161–
181. https://doi.org/10.1016/j.applanim.2009.02.023

Mendl M, Paul ES (2020) Animal affect and decision-making. Neurosci Biobehav Rev 112:144–163.
https://doi.org/10.1016/j.neubiorev.2020.01.025

Menzel, R (1990) Learning, Memory, and "Cognition" in Honey Bees. In: Neurobiology of Comparative
Cognition. Psychology Press, pp 237–292

Merçot H, Defaye D, Capy P, et al (1994) Alcohol Tolerance, ADH Activity, and Ecological Niche of
Drosophila Species. Evolution 48:746–757. https://doi.org/10.1111/j.1558-
5646.1994.tb01358.x

Miler K (2025) Ethanol and pollinators: expanding Bowland et al.'s framework. Trends Ecol Evol
40:115–116. https://doi.org/10.1016/j.tree.2024.11.019

Miler K, Kuszewska K, Privalova V, Woyciechowski M (2018) Honeybees show adaptive reactions to
ethanol exposure. Sci Rep 8:8707. https://doi.org/10.1038/s41598-018-27117-6

Miler K, Stec D, Kamińska A, et al (2021) Alcohol intoxication resistance and alcohol dehydrogenase
levels differ between the honeybee castes. Apidologie 52:230–241.
https://doi.org/10.1007/s13592-020-00812-y

Mixson TA, Abramson CI, Božič J (2010) The Behavior and Social Communication of Honey Bees (APIS
Mellifera Carnica Poll.) under the Influence of Alcohol. Psychol Rep 106:701–717.
https://doi.org/10.2466/pr0.106.3.701-717

Mustard JA, Edgar EA, Mazade RE, et al (2008) Acute ethanol ingestion impairs appetitive olfactory
learning and odor discrimination in the honey bee. Neurobiol Learn Mem 90:633–643.
https://doi.org/10.1016/j.nlm.2008.07.017

Mustard JA, Oquita R, Garza P, Stoker A (2019) Honey Bees (Apis mellifera) Show a Preference for the
Consumption of Ethanol. Alcohol Clin Exp Res 43:26–35. https://doi.org/10.1111/acer.13908

Neville V, Nakagawa S, Zidar J, et al (2020) Pharmacological manipulations of judgement bias: A
systematic review and meta-analysis. Neurosci Biobehav Rev 108:269–286.
https://doi.org/10.1016/j.neubiorev.2019.11.008

Ostap-Chec, M, Bajorek, D, Antoł, W, et al (2025) Honey bees are resilient to the long-term presence
of alcohol in their diet. Exotoxicology 1158–1168. https://doi.org/10.1007/s10646-025-
02903-x

Ostap-Chec M, Opalek M, Stec D, Miler K (2021) Discontinued alcohol consumption elicits withdrawal
symptoms in honeybees. Biol Lett 17:20210182. https://doi.org/10.1098/rsbl.2021.0182

Pahl M, Si A, Zhang S (2013) Numerical Cognition in Bees and Other Insects. Front Psychol 4:.
https://doi.org/10.3389/fpsyg.2013.00162

Perry CJ, Baciadonna L (2017) Studying emotion in invertebrates: what has been done, what can be
measured and what they can provide. J Exp Biol 220:3856–3868.
https://doi.org/10.1242/jeb.151308

Perry CJ, Barron AB, Chittka L (2017) The frontiers of insect cognition. Curr Opin Behav Sci 16:111–
118. https://doi.org/10.1016/j.cobeha.2017.05.011

Phillips TJ, Shent EH (1996) Neurochemical Bases of Locomotion and Ethanol Stimulant Effects. In:
International Review of Neurobiology. Elsevier, pp 243–282

Pinheiro, B (2000) Mixed-Effects Models in S and S-PLUS. Springer, New York

Procenko O, Read JCA, Nityananda V (2024) Physically stressed bees expect less reward in an active
choice judgement bias test. Proc R Soc B Biol Sci 291:20240512.
https://doi.org/10.1098/rspb.2024.0512

Romero-González JE, Zhuo Z, Chen L, et al (2025) Positive affective contagion in bumble bees. Science
390:377–380

Sandoz J-C, Pham-Delègue M-H (2004) Spontaneous Recovery After Extinction of the Conditioned
Proboscis Extension Response in the Honeybee. Learn Mem 11:586–597.
https://doi.org/10.1101/lm.81504

Schaeffer RN, Mei YZ, Andicoechea J, et al (2017) Consequences of a nectar yeast for pollinator
preference and performance. Funct Ecol 31:613–621. https://doi.org/10.1111/1365-
2435.12762

Scheiner R, Abramson CI, Brodschneider R, et al (2013) Standard methods for behavioural studies of
Apis mellifera. J Apic Res 52:1–58. https://doi.org/10.3896/IBRA.1.52.4.04

Schlüns H, Welling H, Federici JR, Lewejohann L (2017) The glass is not yet half empty: agitation but
not Varroa treatment causes cognitive bias in honey bees. Anim Cogn 20:233–241.
https://doi.org/10.1007/s10071-016-1042-x

Solvi C, Baciadonna L, Chittka L (2016) Unexpected rewards induce dopamine-dependent positive
emotion–like state changes in bumblebees. Science 353:1529–1531.
https://doi.org/10.1126/science.aaf4454

Stephenson L, Chicas-Mosier A, Black T, et al (2021) Inducing Ethanol Tolerance in Free-Flying Honey
Bees (Apis mellifera L.). Int J Comp Psychol 34:. https://doi.org/10.46867/IJCP.2021.34.00.03

Strang C, Muth F (2023) Judgement bias may be explained by shifts in stimulus response curves. R Soc
Open Sci 10:221322. https://doi.org/10.1098/rsos.221322

Valdez GR, Roberts AJ, Chan K, et al (2002) Increased Ethanol Self-Administration and Anxiety-Like
Behavior During Acute Ethanol Withdrawal and Protracted Abstinence: Regulation by
Corticotropin-Releasing Factor. Alcohol Clin Exp Res 26:1494–1501.
https://doi.org/10.1111/j.1530-0277.2002.tb02448.x

Van Der Goot MH, Kooij M, Stolte S, et al (2021) Incorporating inter-individual variability in
experimental design improves the quality of results of animal experiments. PLOS ONE
16:e0255521. https://doi.org/10.1371/journal.pone.0255521

Van Erp AMM, Miczek KA (1997) Increased aggression after ethanol self-administration in male
resident rats. Psychopharmacology (Berl) 131:287–295.
https://doi.org/10.1007/s002130050295

Varnon CA, Dinges CW, Black TE, et al (2018) Failure to Find Ethanol-Induced Conditioned Taste
Aversion in Honey Bees (*Apis mellifera* L.). Alcohol Clin Exp Res 42:1260–1270.
https://doi.org/10.1111/acer.13761

Wiens F, Zitzmann A, Lachance M-A, et al (2008) Chronic intake of fermented floral nectar by wild
treeshrews. Proc Natl Acad Sci 105:10426–10431. https://doi.org/10.1073/pnas.0801628105

Wood, S. N. (2017) Generalized additive models: an introduction with R (2nd edition). Chapman and
620   Hall/CRC

621   Wright GA, Lillvis JL, Bray HJ, Mustard JA (2012) Physiological State Influences the Social Interactions
622   of Two Honeybee Nest Mates. PLoS ONE 7:e32677.
623   https://doi.org/10.1371/journal.pone.0032677

624   Yang, Z, Xiong, T, Xie, D, et al (2025) Effects of ethanol exposure and withdrawal on olfactory
625   sensitivity and learning and memory in honeybees(Apis mellifera). J Asia-Pac Entomol
626   28:102476

627