Title: Single-cell and cell-type-level insights into stress-driven immune regulation relevant to disease

Authors: Sofia Benavides[1,2], Oksana Kosyk[1], Lillian Carpenter[1], Christina Georgiou[1], Brian C. Miller[3], Natalie
Stanley[4,5], Anthony S. Zannas[1,2,5,6]*

Affiliations:

[1] Department of Psychiatry, University of North Carolina, Chapel Hill, NC 27599, USA

[2] UNC Neuroscience Center, The University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

[3] Department of Medicine, University of North Carolina, Chapel Hill, NC 27599, USA

[4] Department of Computer Science and Computational Medicine Program, The University of North Carolina at
Chapel Hill

[5] Department of Genetics, University of North Carolina, Chapel Hill, NC 27514, USA

[6] Carolina Stress Initiative, University of North Carolina School of Medicine, Chapel Hill, NC 27514, USA

*Corresponding author. Email: anthony_zannas@med.unc.edu

Abstract

Environmental stress contributes to several disease conditions – including cardiovascular disease, mental illness, and
autoimmune disorders – through immune dysregulation. Stress operates through the hypothalamic-pituitary-adrenal
and sympathoadrenal axes, leading to peripheral secretion of cortisol and norepinephrine, respectively. While stress
profoundly influences immune function, the underlying single-cell and cell-type-specific mechanisms remain poorly
understood. To address this knowledge gap, we employed an ex vivo model investigating the transcriptomic
responses of primary peripheral blood mononuclear cells and neutrophils to physiological stress levels (i.e., levels
reached during in vivo stress) of cortisol and norepinephrine at single-cell resolution. We identified novel stress-
hormone-dependent and cell-type-specific transcriptomic changes, including uniquely regulated genes and
inflammation-related signatures within distinct innate (e.g., $CD163^+$ monocytes/macrophages, neutrophils) and other
immune cell types. The single-cell-derived signatures, applied to two independent human cohorts, effectively
distinguished stress-related diseases. Our findings provide cell-type-level insights into how stress hormones
modulate immune function relevant to stress-related conditions.

INTRODUCTION

Environmental stress, an unavoidable characteristic of every society (*1*), has been linked to an increased risk for
various diseases (*2-19*). For example, studies have established an association between excessive psychosocial stress
and an increased risk of cardiovascular disease (*2-7*), mental illness (*20, 21*), and autoimmune disorders (*13-16*).
While the underlying mechanisms are poorly understood, immune dysregulation has been repeatedly highlighted as
a key process shared across stress-related conditions (*22-28*).

Stress operates through two distinct biological pathways: the Hypothalamic-Pituitary-Adrenal (HPA) axis (*29*) and
the sympathoadrenal axis (*30*). These pathways culminate in the secretion of stress hormones – cortisol and
norepinephrine – into the periphery, where they interact with immune cells (*31*). Controversy exists regarding the
effects of stress hormones on immune cells (*32*); some literature suggests an anti-inflammatory response (*33, 34*),
while other studies demonstrate pro-inflammatory effects (*22, 35*). However, to our knowledge, all previous research
on stress hormones and immune cells has focused on analyzing a mixture of cells (i.e., bulk-level analysis)(*36-39*),
thereby not accounting for cell type-specific effects that could play a critical role in mediating stress pathology.
Additionally, most studies investigating stress-induced immune dysfunction have used in-vivo models. While these
models are critical for studying complex effects on body systems, they are confounded by natural shifts in cell
populations and changes in cell-type distribution that constantly occur in vivo, thus providing limited single-cell and
cell-type-specific insights into how stress regulates immune function.

To address these knowledge gaps, here we employ a cell model (ex vivo) approach and single-cell RNA sequencing
(scRNA-seq) to directly assess the immune effects of physiological stress hormone exposure at the single-cell level.
To model physiological stress in culture, primary peripheral blood mononuclear cells (PBMCs) and neutrophils
isolated from human donors were exposed to the endogenous human stress hormones cortisol and norepinephrine at
concentrations known to be reached in human tissues during in vivo stress (*40-47*). This approach enabled us to
model stress ex vivo and conduct a comprehensive analysis of the unique immune transcriptomic signatures of
physiological stress hormone exposure at the single-cell and cell-type levels. The single-cell-derived signatures,
applied to two independent human cohorts, effectively distinguished stress-related diseases. Our findings provide
new insights into how stress hormones influence immune cell function and the immune cell-type-specific
mechanisms underlying stress-related conditions.

RESULTS
Immune single-cell transcriptional profiling after ex vivo exposure to physiological stress hormone levels

To evaluate the effects of stress hormones on immune cell function, we obtained whole blood samples from four
male and four female healthy donors and isolated PBMCs and neutrophils. Cells from each donor were treated with
vehicle control (0.0001% DMSO), 100 nM cortisol (CORT), or 10 nM norepinephrine (NE) (Fig. 1A). These
concentrations were selected based on previous literature indicating that glucocorticoids can reach tissue
concentrations of 100 nM (*40-44*), and norepinephrine reaches 10 nM circulating concentration (*45-47*) during in
vivo stress exposure. This robust experimental design enables within-donor comparisons, thereby accounting for
potential inter-individual variability in transcriptional responses.

After excluding low-quality cells and potential doublets, we retained 62,975 high-quality cells: 23,000 from the
vehicle, 20,707 from the CORT, and 19,268 from the NE group. Quality control metrics for the retained cells are
shown in Fig. S1B–F. Cluster analysis identified 20 distinct cell clusters, visualized using a standard uniform
manifold approximation and projection (UMAP) approach (*48*) (Fig. 1B). Although the treatment groups showed
similar representation across all 20 clusters, initial UMAP visualization by treatment revealed subtle shifts in cell
distribution attributable to the treatments (Figs. 1C, D).

The clusters encompassed all major cell types typically found in peripheral blood, including lymphocytes,
monocytes, and neutrophils. The distribution of cells by individual sample is shown in Fig. S1A. All clusters were
successfully annotated using the top differentially expressed genes (DEGs), canonical markers, and reference data
from previously published scRNA-seq datasets (*49, 50*) (Fig. 1E).

Exposure to physiological stress hormone levels alters immune cell type composition

While stress-induced changes in immune cell type composition have been well-documented in vivo, the functional
impact of CORT and NE on distinct immune cell types remains unclear. Addressing this question in culture allows
for evaluating the direct effects of stress hormones on immune cell composition, free from the confounding effects
of cell mobilization and redistribution observed in vivo. Using density plots, we qualitatively observed treatment-
dependent changes in immune composition. CORT treatment resulted in a general decrease in lymphocyte
proportions and altered the distribution of monocyte/macrophage (MonoMac) populations (Fig. 2A). In contrast, NE
treatment produced a distinct pattern of T cell distributions and a unique MonoMac distribution that differed from
the control and CORT-treated groups (Fig. 2A).

We further quantified these changes, leveraging the strength of our experimental design that enables within-donor
comparisons between control and treated cells. These comparisons revealed significant stress hormone-induced
changes in cell type proportions. Specifically, CORT treatment led to nominally significant reductions in memory
and naïve CD8 T cells, basophils, and plasmacytoid dendritic cells. However, after correcting for multiple testing,
only the decreases in naïve CD8 T cells and plasmacytoid dendritic cells remained statistically significant (Fig. 2B).
Similarly, NE treatment resulted in nominally significant decreases in plasmacytoid dendritic cells and memory B
cells, as well as increases in naïve CD4 T cells and *CD163*+ MonoMac populations (Fig. 2C). After multiple testing
corrections, only the reductions in memory B cells and plasmacytoid dendritic cells remained statistically

significant. Additional patterns of change in cell type proportions were observed, but did not reach statistical
significance (fig. S2A-B).

To validate these findings on cell type composition, we performed RNAScope analysis in four independent donors
(two males and two females). We chose to focus on validating the CORT-driven decrease in naïve CD8+ T cells due
to the greater abundance of this cell type among the statistically significant cell populations (naïve CD8 T ~6.26%,
memory B ~1.5%, plasmacytoid dendritic cells ~0.29%), which allowed for greater statistical power. We observed a
pattern of decreased naïve CD8+ T cells across donors (fig. S3A-C), but the overall result did not reach statistical
significance. This may reflect limitations in power and the lower sensitivity of the RNAScope assay compared to
scRNA-seq.

To further evaluate the potential clinical relevance of stress hormone-induced immune remodeling, we calculated the
monocyte-to-lymphocyte ratio (MLR), an established prognostic marker of conditions such as kidney disease (*51*),
cardiovascular mortality (*52*), and cancer (*53, 54*). Our analysis revealed a significant increase in MLR following
CORT, but not NE, treatment (Fig. 2D). These findings indicate that, even in the absence of cell mobilization and
redistribution, exposure to physiological stress hormone levels can selectively remodel the immune cell
composition.

Exposure to physiological stress hormone levels induces cell type-specific gene expression changes

Next, we investigated the extent to which exposure to CORT and NE concentrations corresponding to physiological
levels of stress alters immune cell-type-specific gene expression.

CORT increased the expression of well-characterized glucocorticoid-responsive genes, such as *FKBP5* and *ZBTB16*
(Fig. 3A), at the bulk level (i.e., when pooling all cell types), confirming the effectiveness of our ex vivo treatment.
In total, we identified 450 bulk-level differentially expressed genes (DEGs) with an absolute log fold change (LFC)
> 0.25 and a false discovery rate (FDR)-corrected p-value (q) < 0.05. Of these, 251 genes were downregulated and
199 were upregulated (table S1). We validated several top DEGs (*VSIG4*, *FKBP5*, *ZBTB16*) using qPCR (Fig. 3C).
We further examined the expression patterns of the top DEGs due to CORT treatment across different immune cell
types. Notably, CORT induced both shared and distinct transcriptional changes across cell types (Fig. 3B). For
instance, *VSIG4* was markedly upregulated in progenitor cells (LFC = 24.4), while it was strongly downregulated in
plasmacytoid dendritic cells (LFC = -23.4). Among all DEGs, *FKBP5* was the only gene consistently upregulated
across all cell types due to CORT treatment (with an absolute LFC > 0.25). Although not part of the top 10 DEGs,
RNU4-2 showed consistent upregulation across most cell types except plasmacytoid dendritic cells, while *CCND3*
exhibited consistent upregulation across all cell types except plasmacytoid dendritic and progenitor cells.

NE exposure led to a total of 472 DEGs with an absolute LFC greater than 0.25 at the bulk level. Strikingly, 451 of
these genes were downregulated, while only 21 were upregulated (Fig. 3D, table S2). We validated several top
DEGs (*RUN4-2*, *CXCL1*, *IL1B*) via qPCR, and although these results did not reach statistical significance, they
followed a similar expression pattern as our scRNA-seq findings (Fig. 3F). We then analyzed the cell type-specific
effects and found that, similar to CORT, NE treatment also induced cell type-specific gene expression patterns. For
example, *CXCL3* was strongly upregulated in neutrophils (LFC = 23.59) and plasmacytoid dendritic cells (LFC =
26.51), whereas it was markedly downregulated in Memory CD8 T cells (LFC = -19.92) and Plasma B cells (LFC =
-25.61) (Fig. 3E). Among all the DEGs, we identified *RNU4-2* as the single gene consistently upregulated in the
same direction across all immune cell types due to NE treatment (with an absolute LFC > 0.25). Given the lack of a
benchmark gene marker for assessing NE signaling, our analyses for the first time identify *RNU4-2* as a strong
candidate for further investigation in immune cells.

Interestingly, CORT and NE treatments shared 100 common DEGs at the bulk level, of which 62 were differentially
regulated in the same direction and 38 were regulated in opposite directions.

To our knowledge, this is the first study showing that stress hormones induce distinct and drastically differential
effects on gene expression that are dependent on immune cell type. These results may help explain the
inconsistencies observed in the literature regarding the effects of stress hormones on immune function (*22, 32, 33,*
55-57), as prior studies were conducted at the bulk level.

Innate immune cell types display distinct inflammatory responses when exposed to CORT and NE

Given our interest in the role of stress in disease-related inflammation, we analyzed how CORT and NE treatments influence differential inflammatory gene expression across immune cell types. To achieve this, we utilized AUCell, a tool designed to assess whether a specific input gene set is enriched within the expression profile of each cell (*58*). For our analysis, we utilized the hallmark inflammatory response gene set, consisting of 200 genes originally developed for gene set enrichment analysis (GSEA)(*59*). The distribution of cells by AUCell score is shown in fig. S4A and S4B. Our findings revealed that cells exposed to stress hormones exhibited cell-type-specific differential expression of inflammation-related genes, with MonoMac cell clusters demonstrating the most pronounced changes (Fig. 4A). To quantify these differences, we compared the AUCell scores for each cell type between the control and treatment groups. Following CORT exposure, neutrophils, *IL7R*[+] MonoMacs, and *CD163*[+] MonoMacs showed significant enrichment of inflammation-related gene expression, with *CD163*[+] MonoMacs exhibiting the most pronounced change (Fig.4B). Similarly, NE treatment resulted in inflammation-related gene enrichment in *IL7R*[+] MonoMacs and *CD163*[+] MonoMacs, with *CD163*[+] MonoMacs again showing the greatest change (Fig. 4C). Interestingly, in both treatment conditions, we observed a significant downregulation of inflammation-related genes in *VCAN*[+] MonoMacs (Fig. 4A and B). This highlights how the effects of different stress hormones can be contrasting even between closely related cell types, such as those in the MonoMac compartment, emphasizing the importance of cell-type-specific analysis.

Characterization and validation of CD163+ MonoMacs as a novel stress-activated innate immune cell subtype

Given the observed robust inflammatory signature enrichment in *CD163*[+] MonoMacs (F-statistic: 75.45, p-value: 4.02e-33) (fig. S4C), we further characterized the stress-driven DEGs specifically in this cell type. CORT exposure resulted in a total of 477 DEGs with an absolute LFC greater than 0.25, of which 218 were downregulated and 259 were upregulated following (table S3). Among the upregulated genes, we identified several well-known pro-inflammatory genes, such as *IL6* (LFC = 27.06), *IL1A* (LFC = 4.02), and *CCL4* (LFC = 3.27) (Fig. 4D). NE treatment resulted in 319 DEGs, with 209 downregulated and 110 upregulated (table S4). Similar to CORT exposure, we observed the upregulation of well-established inflammatory cytokines and chemokines, including *CXCL1* (LFC = 2.46), *IL1B* (LFC = 3.21), and *CXCL8* (LFC = 2.41) (Fig. 4E). These genes—particularly *IL6*, *IL1B*, and the chemokines *CXCL1* and *CXCL8*—are central mediators of inflammation, and their dysregulation has been implicated in the pathogenesis of various inflammatory and autoimmune diseases, highlighting the potential impact of stress-induced immune modulation.

To validate the enrichment of inflammatory gene expression in *CD163*[+] cells following cortisol exposure, we performed RNAScope analysis looking specifically for cells expressing *CD163* that also express *IL6* and *IL1A*. We observed an increase in the number of *CD163*[+] cells expressing *IL6*, as well as a marked increase in *IL6* expression within *IL6*[+] *CD163*[+] cells, as evidenced by a higher number of *IL6* puncta (Fig. 4F & G). Additionally, while the proportion of *CD163*[+] cells expressing *IL1A* did not change, there was a trend (p = 0.08) toward increased *IL1A* expression within these cells, as indicated by an increase in IL1A puncta (Fig. 4F & G). Taken together, these analyses identify *CD163+* MonoMacs as a novel stress-activated innate immune cell subtype and further characterize distinct stress-responsive, disease-relevant gene signatures within this cell subtype.

CD163+ **MonoMacs-specific stress-inflammation transcriptional risk score is associated with disease**

To explore the potential clinical and translational relevance of our findings, we sought to develop a transcriptional risk score (TRS) informed by our previous findings. By integrating the cumulative effects of multiple genomic sites, these composite markers are thought to better capture risk for complex diseases as compared to single risk genes (*60, 61*).

Building on this framework, we developed a Cell-type-specific (*CD163+*) Stress-Inflammation Score (hereafter denoted CytSIS). This parsimonious score was generated by identifying the CORT and NE DEGs in *CD163+* cells that overlapped with the hallmark inflammatory response gene set (n = 10 genes), weighting each gene by its treatment effect size (summed regression coefficient across CORT and NE treatments), and summing the resultant weighted gene expression products (Fig. 5A-B; table S5). Application of CytSIS to our scRNA-seq dataset

demonstrated that stress hormone exposure robustly increased CytSIS in *CD163*+ cells (F-value = 248.25 p-value =
1.18e-104) (Fig. 5C), as expected.

To assess the applicability of CytSIS to cardiovascular disease, we next applied it in a cohort of PBMCs from HIV-
infected individuals, classified into three groups: those with no cardiovascular disease (CVD), those with untreated
CVD, and those with CVD treated with statins (*62*). The analysis focused on *CD163*+ cells (Fig. S5A). CytSIS was
increased in the *CD163*+ cells of the CVD as compared to the non-CVD and statin-treated CVD groups (F-value =
17.5440, p-value = 3.2606e-08; Fig. 5D). To confirm that this difference is driven specifically by the transcriptional
effects of stress and is not due to non-specific differences in inflammation-related gene expression across groups, we
conducted 10,000 permutations whereby we generated TRS by randomly drawing genes from the hallmark
inflammatory gene list expressed in this cohort (n = 4), and weighing them by the scRNA-seq-trained effect sizes.
For each permutation, the TRS was then compared between the CVD and non-CVD groups. CytSIS significantly
outperformed the random permutations (p perm = 0.0003) (Fig. 5E). We observed similar findings when drawing
from genes in the whole transcriptome (Fig. S5B).

We also applied CytSIS to a cohort of individuals with rheumatoid arthritis, who were grouped based on their
symptoms into remitted or symptomatic arthritis (*63*). In *CD163*+ expressing cells (fig. S5C), CytSIS was
significantly elevated in the symptomatic compared to remitted group (T-statistic = 5.310, p-value = 1.356e-06) (Fig.
5F). We again performed 10,000 permutations of randomly generated weighted inflammatory TRS (n genes = 10)
that were compared between remitted vs. symptomatic groups. CytSIS significantly outperformed the random
permutations (pperm = 0.00) (Fig. 5G). We observed similar findings when drawing from genes in the whole
transcriptome (Fig. S5D).

Together, these findings indicate that cell-type-specific measures of stress-related inflammation show promise as
novel markers of disease risk.

DISCUSSION

Prior work underscores immune dysregulation as a potential key process linking stress with disease, but how stress
hormones affect distinct immune cells and cell types has remained unknown. This study leveraged single-cell
transcriptomic signatures to determine how the stress hormones CORT and NE influence immune function and the
relevance of these transcriptomic signatures to stress-related conditions.

Prior research has highlighted a complex relationship between stress exposure, stress hormones, and immune
dysregulation in vivo (*64-67*). Both human and rodent studies have shown that stress exposure induces widespread
shifts in immune cell types (*66, 68-70*). However, in vivo studies cannot disentangle whether these shifts reflect
mere mobilization of particular cell types or reprogramming towards different immune cell types. Using our ex
vivo that is devoid of the confounding effects of immune cell mobilization, we found that CORT exposure reduced
plasmacytoid dendritic cells and CD8 naïve T cells (Fig. 2B). The decrease in CD8 T cells aligns with in
vivo findings, where corticosterone administration in mice significantly reduced circulating CD8 T cells, suggesting
cell death or identity changes (*71*). While corticosterone also changes the proportions of monocytes, neutrophils, and
B cells in vivo (*71*), the lack of significant changes in these populations in our model indicates these shifts are likely
due to cell mobilization only. Similarly, NE exposure decreased plasmacytoid dendritic cells and memory B cells
while increasing CD4 T cells (nominally significant) (Fig. 2C). This aligns with in vivo findings showing a trend
toward increased CD4 T cells after norepinephrine administration. This same in vivo study also demonstrated a
dramatic increase in neutrophils following norepinephrine exposure, attributable to cell mobilization (*71*).

Our compositional analysis also revealed that the monocyte-to-lymphocyte ratio (MLR) increases following cortisol
exposure. Elevated MLR has been reported in individuals experiencing high stress compared to those with low stress
exposure (*64*), suggesting that cortisol may be a key driver of this pathology. Additionally, MLR has been studied as
a marker of disease severity (*51-54, 72-74*), supporting a plausible link between stress and disease.

In this study, we demonstrate that stress hormones induce cell-type-specific transcriptomic changes, shedding new
light that may help reconcile prior research supporting a range of pro- and anti-inflammatory effects of stress
hormone exposure. For instance, some studies report that glucocorticoids reduce pro-inflammatory cytokines such as

TNF-α and IL-1β, while increasing anti-inflammatory cytokines like IL-10 (75). Conversely, other studies suggest
that glucocorticoids can promote the secretion of pro-inflammatory cytokines like IL-1β and IL-18 (22). Similarly,
conflicting evidence exists regarding the inflammatory effects of NE (76, 77). Notably, our findings show that stress
hormones can have opposite effects even within closely related cell subtypes (e.g., *VCAN+* MonoMacs vs. *CD163+*
MonoMacs), highlighting the importance of studying distinct immune cell subpopulations in stress research.
Nevertheless, we also identify a single gene differentially expressed consistently across all cell types for CORT
(*FKBP5*) and another for NE (*RNU4-2*), suggesting that these genes could serve as immune markers of stress
exposure.

Our inflammation analysis revealed that the effects of stress hormones are cell-type-specific, with the potential to
either increase or decrease inflammatory gene expression depending on the cellular context. Notably, *CD163+*
MonoMacs exhibited a particularly significant increase in inflammation-related gene expression following exposure
to both CORT and NE, highlighting their broad importance in stress-driven inflammatory states. This finding is
intriguing given that *CD163+* macrophages are widely described in the literature as having anti-inflammatory
phenotypes, often referred to as alternatively activated macrophages (78, 79). However, it is important to note that
CD163+ macrophages are also frequently observed in inflammatory and tumorigenic microenvironments, where
they can contribute to disease-related inflammation through mechanisms distinct from those of classically activated
macrophages (80-83). These cells are further studied for their pro-inflammatory and pro-atherogenic
roles in atherosclerosis, where they may promote plaque progression and instability (81-83). Thus, our finding
that CORT and NE induce inflammation-related gene expression in *CD163+* MonoMacs could provide a
mechanistic link between environmental stress and highly prevalent conditions such as cardiovascular disease. This
hypothesis is further supported by applying our transcriptional risk score, CytSIS, in cohorts of stress-associated
diseases, highlighting the potential role of stress-induced inflammation in disease pathogenesis.

Our dataset provides a valuable resource for developing transcriptional risk scores (TRS), which are increasingly
being used as biomarkers for disease risk and progression (60, 61). By leveraging cell-type-specific transcriptional
responses to stress, we can create more precise tools for identifying individuals at risk of stress-associated diseases.
For example, we demonstrated that $CD163^+$ cell-specific transcriptional signatures effectively distinguish between
healthy and diseased individuals. This approach provides a potential framework for leveraging single-cell datasets to
promote early risk stratification and personalized intervention strategies. These findings may not only advance the
development of disease biomarkers but also highlight potential therapeutic targets for mitigating the impact of stress
and other environmental exposures on health.

MATERIALS AND MEHTODS

Ethics approval and participant consent

262 Human peripheral blood samples were obtained from healthy donors, aged 19 -28, at the UNC Blood Donation
263 Center with informed consent and IRB approval. Eight healthy volunteers participated in the scRNA-sequencing
264 study and q-PCR experiments, and four independent donors in the RNAscope experiments.

Isolation of PBMCs and Neutrophils

PBMCs and polymorphonuclear neutrophils (PMNs) were isolated from human peripheral blood samples as
previously described (84).

12 mL of fresh peripheral blood was collected using vacutainer EDTA tubes, diluted with 12 mL of DPBS
supplemented with EDTA (2mM) and FBS (1%), and then gently layered over Lymphoprep (STEMCELL
Technologies, USA) in a 50 mL conical tube. Samples were centrifuged at 800g for 15 min at room temperature with
breaks off. After centrifugation, PBMCs in the white foamy layer were carefully transferred to a new tube. After
being washed with DPBS twice, samples were then centrifuged at 250g for 10 min, and the pellets were resuspended
with cell culture media (RPMI supplemented with 5% of human serum). Cell counts were performed on the
automated cell counter and 250.000 cells/well were seeded into 24-well plates, incubated at 37C in a 5% CO2
incubator for 1 hour, and then treated with 100 nM Cortisol, 10 nM Norepinephrine, or 0.0001% DMSO (vehicle
control) overnight.

Given their shorter lifespan in cell culture (*85*), PMNs were recovered from the red blood cells layer following 1 x
RBS lysis buffer digestion procedure and washed twice with 1X PBS and resuspended in cell culture media. Cells
were then seeded into 24-well plates at a seeding density of 250,000 cells/well, incubated at 37C in a 5% CO2
incubator for 1 hour, and then treated with 100 nM Cortisol, 10 nM Norepinephrine, or 0.0001% DMSO for 4 hours.

Library preparation and sequencing

Cells were removed from the incubator, and the plates were centrifuged at 500g for 5 minutes for pelleting. Cells
were washed with 1X PBS and carefully detached from each well by gentle pipetting, then resuspended in 1X PBS
to create a single cell suspension. Cells were fixed immediately using a Fixation Kit (Parse Biosciences), counted,
and stored at -80 °C as per the manufacturer's protocol.

The single-cell libraries were constructed according to the Parse WT kit manufacturer's protocol. 4166 cells per
sample (for a total of 100,000 cells from 24 samples) were barcoded using the reverse transcription barcoding
approach. Barcoded cDNA was amplified and verified via TapeStation for size distribution following fragmentation,
end repair, A-tailing, and adaptor ligation to complete library construction. All libraries were sequenced as paired-
end 150-bp reads using the Illumina NovaSeq X Plus 300 platform with a reading depth of 83-104M reads per
sample.

Preprocessing

Preprocessing was performed using Scanpy (v1.9.3) following a previously established workflow (*86*). Cells
expressing fewer than 300 genes were excluded, as were genes expressed in fewer than 5 cells. To further refine the
dataset, cells expressing more than 8,000 genes or with total gene counts exceeding 20,000—likely representing
doublets—were removed. Additionally, cells with mitochondrial gene expression exceeding 15% of the total counts
were filtered out to exclude unhealthy or dying cells. Clusters were annotated using the top DEGs, canonical
markers, and reference data from previously published single-cell RNA sequencing datasets (*49, 50*). No size cutoff
was used to remove small clusters. Highly expressed genes within each subset were identified using Wilcoxon rank
testing implemented in Scanpy.

Compositional Analysis

Cell densities in each subset were calculated and plotted for treatment samples versus controls using Scanpy
embedding density functions (*86*). For each sample, the cell type composition was calculated as the proportion of
cells from a specific cell type relative to the total cells in that sample. To minimize noise due to inter-individual
differences and thus maximize statistical power, differences in cell type composition were assessed using paired t-
tests, pairing each donor's treatment sample with their corresponding control sample. Monocyte to Lymphocyte
Ratio (MLR) was calculated by adding all cell types that could be classified as monocytes (CD163+ MonoMac,
CD74+ MonoMac, IL7R+ MonoMac, VCAN+ MonoMac, CD14- MonoMac) and divided by the sum of all cells
from cell types classified as lymphocytes (Nai CD4 T, Mem CD4 T, Nai CD8 T, Mem CD8 T, Th17, Nai B, Mem B,
Plasma B). Statistical differences between control and treatment groups were calculated with paired t-tests. Two-
sided $p < 0.05$ was the threshold for statistical significance.

RNAScope

PBMCs were isolated from four healthy donors (two male, two female) and treated as previously described.
Following overnight incubation, cells were collected into centrifuge tubes and centrifuged at 300 × g for 5 minutes.
The resulting cell pellets were resuspended in 10% neutral buffered formalin (NBF) and fixed for 15 minutes at
room temperature. After fixation, cells were centrifuged again at 300 × g for 5 minutes. The supernatant was
carefully removed, and the pellets were resuspended in 20–30 μL of pre-warmed HistoGel (Thermo Scientific, HG-
4000-012). While still in liquid form, the cell-HistoGel mixture was pipetted onto a parafilm to form a small gel
droplet encapsulating the PBMCs. The droplet was allowed to solidify and then immersed in 10% NBF for an
additional hour to ensure thorough fixation. Subsequently, the gel droplet was transferred to 70% ethanol and stored
until further processing.

The gel droplet containing PBMCs was processed as formalin-fixed paraffin-embedded (FFPE) tissue and sectioned
onto slides for RNAscope staining, following the protocol provided by Advanced Cell Diagnostics (ACD).
Depending on the experimental conditions, cells were stained for either CD163, IL1A, and IL6, or CD3E, NELL2,
and IL8A, along with DAPI nuclear counterstaining. Imaging was performed using a Zeiss LSM 980 confocal
microscope equipped with a 64× oil immersion objective. Quantification of RNA signals was conducted using the
ImageJ Cell Counter plugin, enabling the enumeration of individual transcript puncta per cell.

Differential Gene Expression Analysis

Differentially expressed genes (DEGs) were identified using the rank_genes_groups function from Scanpy (*86*).
The analysis compared the CORT and NE treatment groups to the control group using the Wilcoxon rank-sum test.
Gene expression data from the raw count matrix was utilized for this analysis to ensure accurate assessment of gene
expression differences. False discovery rate (FDR) correction was applied using the Benjamini-Hochberg method to
account for multiple hypothesis testing. Adjusted p-values and log-fold change were used to rank significant DEGs.

To identify cell-type-specific gene expression changes, subsets of data from each cluster (cell type) were created,
and the rank_genes_groups function was applied again to detect DEGs. The top 10 DEGs from the bulk analysis
were then visualized based on log fold change to assess heterogeneity in differential expression across cell types
(Fig. 3B and D).

qPCR

Total RNA was extracted with an RNeasy kit (Qiagen, cat. No 74106) from cells exposed to DMSO, cortisol, and
norepinephrine. The Super Script IV VILO cDNA kit (Invitrogen, cat.no 11756500) was used to synthesize cDNA.
mRNA levels of the following geneswere assessed by q-PCR: VSIG4, FKBP5, ZBTB16, RNU4-2, CXCL1, IL1B.

Detailed information on Q-PRC primer sequences is provided later. Reactions were performed with two replicates in
a 96-well plate using the Quant Studio 6 (Applied Biosystems). The housekeeping gene *GAPDH* was used as a
reference control.

AUCell

We employed the AUCell algorithm to score the activity of our gene subset of interest across individual cells,
identifying those with significantly higher transcriptomic activity (*58*). This approach involves ranking all genes
within each cell based on their expression levels, effectively creating a prioritized list of genes per cell. AUCell then
calculates the Area Under the Curve (AUC) for each gene set within these rankings, providing a score that reflects
the enrichment of the gene set among the top-expressed genes in each cell. This method allows for the identification
of cells where specific gene sets are actively expressed, independent of the gene expression units and normalization
procedures used. Given our focus on inflammatory gene expression, we selected the Hallmark Inflammatory
Response gene set from Gene Set Enrichment Analysis (GSEA), consisting of 200 genes, as our gene set of interest
(*59*).

After assigning each cell a single-gene enrichment score, we calculated the difference (delta) between the control
and treatment groups within each cell type's AUCell score. Statistical differences between control and treatment
groups were assessed using Student's T-test, with multiple testing corrected using the Bonferroni method.

CytSIS

Development of the Cell-type-specific Stress-Inflammation Score (CytSIS)

Informed by our findings, we developed a transcriptional risk score (TRS) termed CytSIS. By integrating the
cumulative effects of multiple genomic sites, this composite marker approach is thought to better capture risk for
complex diseases as compared to single risk genes (*60, 61*) . CytSIS was generated by identifying the differentially
expressed genes (DEGs) resulting from CORT and NE treatments in *CD163+* MonoMacs that overlapped with the
Hallmark Inflammatory Response gene set (n = 10 genes) (Table S5). For each DEG, the treatment effect size was

calculated by summing the regression coefficients from CORT and NE treatments compared to the control, as
determined through the Wilcoxon rank-sum test. Each gene was thus weighted by its treatment effect size.

Application of CytSIS

To apply the CytSIS score, we first created a subset with cells in the cohorts that expressed the *CD163* gene. We
then calculated the product of each individual gene's weight (trained using our scRNA-seq treatment dataset) and its
gene expression value for each cell (derived from the human cohort dataset). This calculation was performed only
for the genes that were available in each dataset (4 genes for the CVD cohort and 10 genes for the arthritis cohort).
The final CytSIS score for each cell was obtained by summing these products across all genes in the set, yielding a
single score that reflects the relative stress-related inflammatory activity of the cell.

Permutation Testing

To validate the specificity of the CytSIS score, permutation testing was performed to rule out the possibility of the
score being driven by random gene expression patterns. For each cohort (CVD and arthritis), 10,000 permutations
were conducted where a random selection of genes (matched to the number of genes available in each dataset: 4 for
CVD and 10 for arthritis) was drawn from the hallmark inflammatory response gene set (or the whole genome) and
weighted by the same treatment effect sizes. The resulting permuted scores were then compared to the CytSIS scores
to assess whether CytSIS outperformed the random permutations. Statistical significance was determined by
comparing the observed CytSIS score to the distribution of the permuted scores.

Cohorts

Human cohorts used for CytSIS testing were obtained from publicly available data repositories.

Cardiovascular Disease (CVD) (GEO: GSE205320)

All participants in this study were part of the Women's Interagency HIV Study (WIHS) (*62*). From the initial 1,865
participants in the WIHS vascular sub-study, 32 participants were selected for scRNA-seq analysis of PBMCs. The
presence of carotid artery focal plaque was used to classify participants into four groups of eight participants: HIV−,
HIV+CVD−, HIV+CVD+, and HIV+CVD+ on cholesterol-reducing treatment (CRT). For our analysis, we focused
on the role of stress in CVD and, to rule out confounding by other disease contributions, we excluded the HIV−
group.

Rheumatoid Arthritis (RA)

Participants from this study consist of patients meeting the American College of Rheumatology (ACR) classification
criteria and were recruited from UCSF rheumatology between 2016 and 2020 (*63*). PBMCs were collected for
scRNA-seqThe disease Activity in 28 joints using CRP (DAS28-CRP) was used to stratify the patients in remitted (n
= 2), or symptomatic (n = 14) groups with low or moderate disease activity according to the 2019 updated ACR
recommendation on disease activity measures (*87*).

Supplementary Materials

Figs S1 to S5

Tables S1 to S6

REFERENCES

1.   American Psychological Association. (2023). Stress in America 2023: A nation recovering from collective trauma. https://www.apa.org/news/press/releases/stress/2023/collective-trauma-recovery.

2.   M. Kumari *et al.*, Chronic stress accelerates atherosclerosis in the apolipoprotein E deficient mouse. *Stress* **6**, 297-299 (2003).

3.   L. Roth *et al.*, Chronic intermittent mental stress promotes atherosclerotic plaque vulnerability, myocardial infarction and sudden death in mice. *Atherosclerosis* **242**, 288-294 (2015).

4.   K. N. Kershaw *et al.*, Associations of Stressful Life Events and Social Strain With Incident Cardiovascular Disease in the Women's Health Initiative. *Journal of the American Heart Association* **3**, e000687.

5.   I. M. Carey *et al.*, Increased Risk of Acute Cardiovascular Events After Partner Bereavement: A Matched Cohort Study. *JAMA Internal Medicine* **174**, 598-605 (2014).

6.   W. Ruberman, E. Weinblatt, J. D. Goldberg, B. S. Chaudhary, Psychosocial influences on mortality after myocardial infarction. *N Engl J Med* **311**, 552-559 (1984).

7.   V. Vaccarino *et al.*, Post-traumatic stress disorder and incidence of coronary heart disease: a twin study. *J Am Coll Cardiol* **62**, 970-978 (2013).

8.   L. Leshan, Psychological states as factors in the development of malignant disease: a critical review. *J Natl Cancer Inst* **22**, 1-18 (1959).

9.   B. Mravec, M. Tibensky, L. Horvathova, Stress and cancer. Part I: Mechanisms mediating the effect of stressors on cancer. *J Neuroimmunol* **346**, 577311 (2020).

10. J. W. Eng *et al.*, A nervous tumor microenvironment: the impact of adrenergic stress on cancer cells, immunosuppression, and immunotherapeutic response. *Cancer Immunol Immunother* **63**, 1115-1128 (2014).

11. C. B. Colon-Echevarria, R. Lamboy-Caraballo, A. N. Aquino-Acevedo, G. N. Armaiz-Pena, Neuroendocrine Regulation of Tumor-Associated Immune Cells. *Front Oncol* **9**, 1077 (2019).

12. M. H. Antoni, F. S. Dhabhar, The impact of psychosocial stress and stress management on immune responses in patients with cancer. *Cancer* **125**, 1417-1431 (2019).

13. H. G. Koenig, H. J. Cohen, in *The link between religion and health: Psychoneuroimmunology and the faith factor*. (Oxford University Press, New York, NY, US, 2002), pp. 174-196.

14. L. Stojanovich, D. Marisavljevich, Stress as a trigger of autoimmune disease. *Autoimmun Rev* **7**, 209-213 (2008).

15. B. Hägglöf, L. Blom, G. Dahlquist, G. Lönnberg, B. Sahlin, The Swedish childhood diabetes study: indications of severe psychological stress as a risk factor for type 1 (insulin-dependent) diabetes mellitus in childhood. *Diabetologia* **34**, 579-583 (1991).

16. A. L. Roberts *et al.*, Association of Trauma and Posttraumatic Stress Disorder With Incident Systemic Lupus Erythematosus in a Longitudinal Cohort of Women. *Arthritis Rheumatol* **69**, 2162-2169 (2017).

17. J. Landeo-Gutierrez, E. Forno, G. E. Miller, J. C. Celedón, Exposure to Violence, Psychosocial Stress, and Asthma. *Am J Respir Crit Care Med* **201**, 917-922 (2020).

18. M. S. Rahi *et al.*, The Impact of Anxiety and Depression in Chronic Obstructive Pulmonary Disease. *Adv Respir Med* **91**, 123-134 (2023).

19. A. Pedersen, R. Zachariae, D. H. Bovbjerg, Influence of Psychological Stress on Upper Respiratory Infection—A Meta-Analysis of Prospective Studies. *Psychosomatic Medicine* **72**, (2010).

452   20.   L. Yang *et al.*, The Effects of Psychological Stress on Depression. *Curr Neuropharmacol* **13**,
453   494-504 (2015).

454   21.   G. Richter-Levin, L. Xu, How could stress lead to major depressive disorder? *IBRO Rep* **4**,
455   38-43 (2018).

456   22.   Y. Z. Liu, Y. X. Wang, C. L. Jiang, Inflammation: The Common Pathway of Stress-Related
457   Diseases. *Front Hum Neurosci* **11**, 316 (2017).

458   23.   R. L. Acabchuk, J. Kamath, J. D. Salamone, B. T. Johnson, Stress and chronic illness: The
459   inflammatory pathway. *Soc Sci Med* **185**, 166-170 (2017).

460   24.   Y. Xiang, M. Zhang, D. Jiang, Q. Su, J. Shi, The role of inflammation in autoimmune disease: a
461   therapeutic target. *Front Immunol* **14**, 1267091 (2023).

462   25.   A. Alfaddagh *et al.*, Inflammation and cardiovascular disease: From mechanisms to
463   therapeutics. *Am J Prev Cardiol* **4**, 100130 (2020).

464   26.   P. T. King, Inflammation in chronic obstructive pulmonary disease and its role in
465   cardiovascular disease and lung cancer. *Clin Transl Med* **4**, 68 (2015).

466   27.   A. Danese, C. M. Pariante, A. Caspi, A. Taylor, R. Poulton, Childhood maltreatment predicts
467   adult inflammation in a life-course study. *Proc Natl Acad Sci U S A* **104**, 1319-1324 (2007).

468   28.   T. Heidt *et al.*, Chronic variable stress activates hematopoietic stem cells. *Nat Med* **20**, 754-
469   758 (2014).

470   29.   C. Leistner, A. Menke, Hypothalamic-pituitary-adrenal axis and stress. *Handb Clin Neurol*
471   **175**, 55-64 (2020).

472   30.   J. R. Carter, D. S. Goldstein, Sympathoneural and adrenomedullary responses to mental
473   stress. *Comprehensive Physiology* **5**, 119 (2015).

474   31.   A. Agorastos, G. P. Chrousos, The neuroendocrinology of stress: the stress-related
475   continuum of chronic disease development. *Molecular Psychiatry* **27**, 502-513 (2022).

476   32.   F. S. Dhabhar, Enhancing versus suppressive effects of stress on immune function:
477   implications for immunoprotection and immunopathology. *Neuroimmunomodulation* **16**,
478   300-317 (2009).

479   33.   K. De Bosscher, W. Vanden Berghe, G. Haegeman, The interplay between the glucocorticoid
480   receptor and nuclear factor-kappaB or activator protein-1: molecular mechanisms for gene
481   repression. *Endocr Rev* **24**, 488-522 (2003).

482   34.   A. E. Coutinho, K. E. Chapman, The anti-inflammatory and immunosuppressive effects of
483   glucocorticoids, recent developments and mechanistic insights. *Mol Cell Endocrinol* **335**,
484   2-13 (2011).

485   35.   E. A. Lannan, A. J. Galliher-Beckley, A. B. Scoltock, J. A. Cidlowski, Proinflammatory actions
486   of glucocorticoids: glucocorticoids and TNFα coregulate gene expression in vitro and in
487   vivo. *Endocrinology* **153**, 3701-3712 (2012).

488   36.   K. C. Torres *et al.*, Norepinephrine, dopamine and dexamethasone modulate discrete
489   leukocyte subpopulations and cytokine profiles from human PBMC. *J Neuroimmunol* **166**,
490   144-157 (2005).

491   37.   M. Wahle *et al.*, Beta2-adrenergic receptors mediate the differential effects of
492   catecholamines on cytokine production of PBMC. *J Interferon Cytokine Res* **25**, 384-394
493   (2005).

494   38.   M. G. Ciliberti *et al.*, Peripheral blood mononuclear cell proliferation and cytokine
495   production in sheep as affected by cortisol level and duration of stress. *J Dairy Sci* **100**, 750-
496   756 (2017).

497   39.   T. W. W. Pace *et al.*, Increased peripheral NF-κB pathway activity in women with childhood
498   abuse-related posttraumatic stress disorder. *Brain, Behavior, and Immunity* **26**, 13-17
499   (2012).

40. S. Wüst, I. S. Federenko, E. F. van Rossum, J. W. Koper, D. H. Hellhammer, Habituation of
cortisol responses to repeated psychosocial stress-further characterization and impact of
genetic factors. *Psychoneuroendocrinology* **30**, 199-211 (2005).

41. R. C. Bhake, J. A. Leendertz, A. C. Linthorst, S. L. Lightman, Automated 24-hours sampling
of subcutaneous tissue free cortisol in humans. *J Med Eng Technol* **37**, 180-184 (2013).

42. Z. Tarjányi *et al.*, Free and total cortisol levels are useful prognostic markers in critically ill
patients: a prospective observational study. *Eur J Endocrinol* **171**, 751-759 (2014).

43. A. H. Hamrahian, T. S. Oseni, B. M. Arafah, Measurements of serum free cortisol in critically
ill patients. *N Engl J Med* **350**, 1629-1638 (2004).

44. M. Christ-Crain *et al.*, Measurement of serum free cortisol shows discordant responsivity to
stress and dynamic evaluation. *J Clin Endocrinol Metab* **92**, 1729-1735 (2007).

45. S. K. Pehrsson, P. Hjemdahl, R. Nordlander, H. Aström, A comparison of sympathoadrenal
activity and cardiac performance at rest and during exercise in patients with ventricular
demand or atrial synchronous pacing. *Br Heart J* **60**, 212-220 (1988).

46. H. Ikarugi *et al.*, Norepinephrine, but not epinephrine, enhances platelet reactivity and
coagulation after exercise in humans. *J Appl Physiol (1985)* **86**, 133-138 (1999).

47. G. Granados, M. E. Garay-Sevilla, J. M. Malacara, K. Wróbel-Zasada, A. Rivera-Cisneros,
Plasma epinephrine and norepinephrine response to stimuli in autonomic neuropathy of
type 2 diabetes mellitus. *Acta Diabetol* **37**, 55-60 (2000).

48. J. Healy, L. McInnes, Uniform manifold approximation and projection. *Nature Reviews
Methods Primers* **4**, 82 (2024).

49. N. S. Hackert *et al.*, Human and mouse neutrophils share core transcriptional programs in
both homeostatic and inflamed contexts. *Nature Communications* **14**, 8133 (2023).

50. G. Monaco *et al.*, RNA-Seq Signatures Normalized by mRNA Abundance Allow Absolute
Deconvolution of Human Immune Cell Types. *Cell Reports* **26**, 1627-1640.e1627 (2019).

51. A. V. Mureşan *et al.*, The Predictive Value of NLR, MLR, and PLR in the Outcome of End-
Stage Kidney Disease Patients. *Biomedicines* **10**, (2022).

52. Y. Hua, J.-Y. Sun, Y.-X. Lou, W. Sun, X.-Q. Kong, Monocyte-to-lymphocyte ratio predicts
mortality and cardiovascular mortality in the general population. *International Journal of
Cardiology* **379**, 118-126 (2023).

53. M. Li *et al.*, The pretreatment lymphocyte to monocyte ratio predicts clinical outcome for
patients with urological cancers: A meta-analysis. *Pathol Res Pract* **215**, 5-11 (2019).

54. D. Tan, Y. Fu, W. Tong, F. Li, Prognostic significance of lymphocyte to monocyte ratio in
colorectal cancer: A meta-analysis. *Int J Surg* **55**, 128-138 (2018).

55. M. A. Horowitz *et al.*, Glucocorticoids prime the inflammatory response of human
hippocampal cells through up-regulation of inflammatory pathways. *Brain Behav Immun*
87, 777-794 (2020).

56. H. Y. Lim, N. Müller, M. J. Herold, J. van den Brandt, H. M. Reichardt, Glucocorticoids exert
opposing effects on macrophage function dependent on their concentration. *Immunology*
122, 47-53 (2007).

57. M. P. Yeager, P. A. Pioli, P. M. Guyre, Cortisol exerts bi-phasic regulation of inflammation in
humans. *Dose Response* **9**, 332-347 (2011).

58. S. Aibar *et al.*, SCENIC: single-cell regulatory network inference and clustering. *Nature
Methods* **14**, 1083-1086 (2017).

59. A. Liberzon *et al.*, The Molecular Signatures Database (MSigDB) hallmark gene set
collection. *Cell Syst* **1**, 417-425 (2015).

60. U. M. Marigorta *et al.*, Transcriptional risk scores link GWAS to eQTLs and predict
complications in Crohn's disease. *Nature Genetics* **49**, 1517-1521 (2017).

61. N. Shan *et al.*, A novel transcriptional risk score for risk prediction of complex human
diseases. *Genet Epidemiol* **45**, 811-820 (2021).
62. J. Vallejo *et al.*, Combined protein and transcript single-cell RNA sequencing in human
peripheral blood mononuclear cells. *BMC Biology* **20**, 193 (2022).
63. M. Binvignat *et al.*, Single-cell RNA-Seq analysis reveals cell subsets and gene signatures
associated with rheumatoid arthritis disease activity. *JCI Insight* **9**, (2024).
64. T. J. Barrett *et al.*, Chronic stress primes innate immune responses in mice and humans.
Cell Reports **36**, (2021).
65. C. C. Li *et al.*, Restraint Stress-Induced Immunosuppression Is Associated with Concurrent
Macrophage Pyroptosis Cell Death in Mice. *Int J Mol Sci* **24**, (2023).
66. L. Tang *et al.*, Acute stress induces an inflammation dominated by innate immunity
represented by neutrophils in mice. *Front Immunol* **13**, 1014296 (2022).
67. M. Bodemeier Loayza Careaga, T. J. Wu, Chronically stressed male and female mice show a
similar peripheral and central pro-inflammatory profile after an immune challenge. *PLoS
One* **19**, e0297776 (2024).
68. Y. Ishikawa *et al.*, Repeated social defeat stress induces neutrophil mobilization in mice:
maintenance after cessation of stress and strain-dependent difference in response. *British
Journal of Pharmacology* **178**, 827-844 (2021).
69. C. Xu, S. K. Lee, D. Zhang, P. S. Frenette, The Gut Microbiome Regulates Psychological-
Stress-Induced Inflammation. *Immunity* **53**, 417-428.e414 (2020).
70. A. Niraula, Y. Wang, J. P. Godbout, J. F. Sheridan, Corticosterone Production during
Repeated Social Defeat Causes Monocyte Mobilization from the Bone Marrow,
Glucocorticoid Resistance, and Neurovascular Adhesion Molecule Expression. *The Journal
of Neuroscience* **38**, 2328-2340 (2018).
71. F. S. Dhabhar, W. B. Malarkey, E. Neri, B. S. McEwen, Stress-induced redistribution of
immune cells—From barracks to boulevards to battlefields: A tale of three hormones – Curt
Richter Award Winner. *Psychoneuroendocrinology* **37**, 1345-1368 (2012).
72. J. Wang *et al.*, The MLR, NLR, PLR and D-dimer are associated with clinical outcome in lung
cancer patients treated with surgery. *BMC Pulm Med* **22**, 104 (2022).
73. Y. Wen *et al.*, Monocyte/Lymphocyte Ratio and Cardiovascular Disease Mortality in
Peritoneal Dialysis Patients. *Mediators Inflamm* **2020**, 9852507 (2020).
74. R. Muto *et al.*, Increased Monocyte/Lymphocyte Ratio as Risk Marker for Cardiovascular
Events and Infectious Disease Hospitalization in Dialysis Patients. *Blood Purif* **51**, 747-755
(2022).
75. K. De Bosscher, W. Vanden Berghe, G. Haegeman, The interplay between the glucocorticoid
receptor and nuclear factor-kappaB or activator protein-1: molecular mechanisms for gene
repression. *Endocr Rev* **24**, 488-522 (2003).
76. G. Pongratz, R. H. Straub, The sympathetic nervous response in inflammation. *Arthritis Res
Ther* **16**, 504 (2014).
77. D. Sharma, J. D. Farrar, Adrenergic regulation of immune cell function and inflammation.
Semin Immunopathol **42**, 709-717 (2020).
78. L. Fischer-Riepe *et al.*, CD163 expression defines specific, IRF8-dependent, immune-
modulatory macrophages in the bone marrow. *Journal of Allergy and Clinical Immunology*
146, 1137-1151 (2020).
79. S. Gordon, Alternative activation of macrophages. *Nat Rev Immunol* **3**, 23-35 (2003).
80. M. K. Skytthe, J. H. Graversen, S. K. Moestrup, Targeting of CD163(+) Macrophages in
Inflammatory and Malignant Diseases. *Int J Mol Sci* **21**, (2020).

81. L. Guo *et al.*, CD163+ macrophages promote angiogenesis and vascular permeability
accompanied by inflammation in atherosclerosis. *J Clin Invest* **128**, 1106-1124 (2018).
82. B. Pourcet, B. Staels, Alternative macrophages in atherosclerosis: not always protective! *J*
Clin Invest **128**, 910-912 (2018).
83. M. Mori *et al.*, CD163+ Macrophages Induce Endothelial-to-Mesenchymal Transition in
Atheroma. *Circulation Research* **135**, e4-e23 (2024).
84. C. Cui, K. Q. Schoenfelt, K. M. Becker, L. Becker, Isolation of polymorphonuclear neutrophils
and monocytes from a single sample of human peripheral blood. *STAR Protocols* **2**, 100845
(2021).
85. M. C. Bonilla *et al.*, How Long Does a Neutrophil Live?-The Effect of 24 h Whole Blood
Storage on Neutrophil Functions in Pigs. *Biomedicines* **8**, (2020).
86. F. A. Wolf, P. Angerer, F. J. Theis, SCANPY: large-scale single-cell gene expression data
analysis. *Genome Biol* **19**, 15 (2018).
87. B. R. England *et al.*, 2019 Update of the American College of Rheumatology Recommended
Rheumatoid Arthritis Disease Activity Measures. *Arthritis Care Res (Hoboken)* **71**, 1540-
1555 (2019).

Acknowledgments: The authors acknowledge Laura Huff for her valuable feedback on this manuscript. We also
thank Dr. Tessa-Jonne Ropp of the UNC Neuroscience Microscopy Core for her assistance with image acquisition,
and Ashley Ezzell for her guidance in developing the methodology for PBMC RNAscope experiments.

Funding: Research reported here was supported by the National Heart, Lung, and Blood Institute of the National
Institutes of Health under Award Number R01HL163031.

Author contributions:

Conceptualization: SB, OK, AZ

Methodology: SB, OK, LC, NS, BM

Investigation: SB

Visualization: SB, LC

Funding acquisition: AZ

Project administration: SB, AZ

Supervision: AZ

Writing – original draft: SB, OK

Writing – review & editing: SB, AZ

Competing interests:

Authors declare that they have no competing interests.

Figures:

Fig. 1 Single-cell RNA sequencing analysis of immune cells after ex vivo exposure to stress hormones.

(A) Graphical representation of experimental model. PBMCs and neutrophils from healthy human donors were
isolated and exposed to control (DMSO) and treatment (cortisol (CORT), norepinephrine (NE)) groups.
After 4 hours of incubation for neutrophils (due to their shorter lifespan (*85*)) and overnight incubation for
PBMCs, cells were prepared for single-cell RNA sequencing.

(B) UMAP plot showing 62,975 cells from all conditions, colored by cell type.

(C) UMAP plot of all cells colored by treatment group (DMSO, CORT, NE).

(D) UMAP plots separated by treatment group, highlighting that all cell type clusters are represented in each
condition.

(E) Heatmap displaying differentially expressed genes (DEGs), canonical marker genes, and genes identified in
previous PBMC studies used for cell type annotation.

A.

DMSO

CORT

NE

B. CORT vs DMSO

Basophils

Memory CD8 T

Naive CD8 T

Plasmacytoid Dendritic

C. NE vs DMSO

CD163+ MonoMacs

Naive CD4 T

Memory B

Plasmacytoid Dendritic

D.

CORT - Monocyte to Lymphocyte Ratio

NE - Monocyte to Lymphocyte Ratio

Fig. 2. Stress hormone treatment induces treatment-specific immune remodeling.

    (A) Density plots showing cell type distributions in control and treatment groups. Blue represents lower cell density, while red indicates higher density.

    (B) Changes in the percentages of plasmacytoid dendritic cells, naïve CD8 T cells, memory CD8 T cells, and basophils following CORT exposure. While all cell types shown were nominally significant, only naïve CD8 T cells and plasmacytoid dendritic cells remained significant after multiple testing correction using the Bonferroni method. Graphs depict the deltas between CORT and DMSO. Paired t-tests were used to test for significance (*$p < 0.0025$).

    (C) Changes in the percentages of plasmacytoid dendritic cells, naïve CD4 T cells, memory B cells, and CD163$^+$ MonoMacs following NE exposure. While all cell types shown were nominally significant, only memory B cells and plasmacytoid dendritic cells remained significant after multiple testing correction using the Bonferroni method. Graphs show the deltas between NE and DMSO. Paired t-tests were used to test for significance (*$p < 0.0025$).

    (D) Differences in the MLR between treatment and control groups. Graphs represent the deltas for CORT (green) and NE (purple) compared to control. Paired t-tests were used to test for significance (*$p < 0.05$).

659   **Fig. 3. CORT and NE treatment lead to cell-type-specific gene expression changes.**

660   (A) Volcano plot showing bulk gene expression changes due to CORT treatment. Green dots represent
661   significant genes with an absolute log fold change (LFC) greater than 0.25 and an FDR-corrected p-value
662   (q) of less than 0.05. The top 10 genes, based on LFC, are labeled.
663   (B) Heatmap of the top 10 differentially expressed genes (DEGs) based on LFC, categorized by cell type.
664   (C) qPCR validation of gene expression changes due to CORT treatment. Statistical significance was
665   determined by Student's t-test. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.
666   (D) Volcano plot showing bulk gene expression changes due to NE treatment. Purple dots indicate significant
667   genes with an absolute LFC greater than 0.25 and q less than 0.05. The top 10 genes according to LFC are
668   labeled.
669   (E) Heatmap of the top 10 DEGs based on LFC, categorized by cell type
670   (F) qPCR validation of gene expression changes due to NE treatment. Statistical significance was determined
671   by Student's t-test.

Fig. 4. *CD163*⁺ MonoMacs show a distinct inflammatory response to stress hormone treatment.

(A) Heatmaps of UMAPs by treatment. Heatmap shows individual cell's AUCell score depicting inflammatory
gene expression enrichment.
(B) Quantification of the difference in AUCell enrichment score between CORT and control. Bonferroni–
corrected p values: $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.
(C) Quantification of the difference in AUCell enrichment score between NE and control. Bonferroni–corrected
p values: $***p < 0.001$.
(D) Volcano plot showing gene expression changes due to CORT treatment in *CD163*⁺ MonoMacs. Green dots
represent significant genes with an absolute log fold change (LFC) greater than 0.25. The top 10 genes,
based on LFC, are labeled.
(E) Volcano plot showing gene expression changes due to NE treatment in *CD163*⁺ MonoMacs. Green dots
represent significant genes with an absolute log fold change (LFC) greater than 0.25. The top 10 genes,
based on LFC, are labeled.
(F) Representative images of RNAScope analysis. Cells were selected for *CD163+* expression (green) and then
IL6 (red) and *IL1A* (magenta) puncta were analyzed.
(G) Image quantifications of RNAScope analysis. Significance was established using student's t- test. : $*p <$
0.05, $**p < 0.01$, $***p < 0.001$.

A.

B.

C.

D.

E.

F.

G.

Fig. 5. Cell-type-specific (*CD163*+) Stress-Inflammation Score (CytSIS) and its application in stress-associated diseases

(A) Overview of the methodology for generating the transcriptional risk score (TRS) CytSIS. The score was developed using *CD163*+ MonoMac inflammatory gene expression, and this score was applied across different single-cell RNA sequencing datasets, including those investigating stress-associated diseases.

(B) Identification of 29 overlapping genes present in both the CORT and NE DEGs and the hallmark inflammatory gene set. These genes were used to generate CytSIS (Supp Table 5).

(C) Application of CytSIS to our dataset shows significantly higher CytSIS scores in stress hormone-treated groups compared to control CD163+ cells, as expected. Statistical significance was determined by a One-way ANOVA and Tukey Post-hoc test. ****$p < 0.0001$.

(D) Application of CytSIS to a cohort of HIV-infected individuals, categorized by cardiovascular disease (CVD) status. CytSIS scores are significantly higher in individuals with CVD compared to those without CVD, with scores decreasing in individuals with CVD who were treated with statin. Statistical significance was determined by a One-way ANOVA and Tukey Post-hoc test. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

(E) 10,000 permutations were conducted, generating TRS by randomly drawing the same number of genes (n = 3) from the hallmark inflammatory genes and weighing them by the same treatment effect sizes. For each permutation, the TRS was then compared between the *CD163*+ cells in the CVD and non-CVD groups. CytSIS significantly outperformed the random permutations.

(F) Application of CytSIS to a cohort of individuals with rheumatoid arthritis, grouped based on symptom severity (remission, low, and moderate arthritis). CytSIS was significantly elevated in the low and moderate groups compared to the remission group. Statistical significance was determined by a One-way ANOVA and Tukey Post-hoc test. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

(G) 10,000 permutations were conducted, generating TRS by randomly drawing the same number of genes (n = 10) from the hallmark inflammatory genes and weighing them by the same treatment effect sizes. For each permutation, the TRS was then compared between the remission and low arthritis groups. CytSIS significantly outperformed the random permutations.