Efficacy of natural marine sponges as a passive environmental DNA sampler for freshwater

fish diversity monitoring

Author's Name: **Ryohei Nakao[1], Manami Inaba[1], Seiji Miyazono[1], Minoru Saito[1,2], Keita**

Maruyama[1,4], Fumiko Imamura[3], Yoshihisa Akamatsu[1]

Institution:

[1] Graduate School of Science and Technology for Innovation, Yamaguchi University, Yamaguchi

University, 2-16-1, Tokiwadai, Ube, Yamaguchi, Japan

[2] Japan International Research Center for Agricultural Sciences, 1-1 Ohwashi, Tsukuba, Ibaraki,

Japan

[3] Research and Development Center, Nippon Koei Co., Ltd, 2304, Inarihara, Tsukuba, Ibaraki,

Japan

[4] Department of Ocean Sciences, Tokyo University of Marine Science and Technology, 4-5-7

Konan, Minato-ku, Tokyo 108-8477, Japan

Correspond Author: Ryohei Nakao, Graduate School of Science and Technology for Innovation,

Yamaguchi University, 2-16-1, Tokiwadai, Ube, Yamaguchi, 755-8611, Japan.

E-mail: lineck92@yahoo.co.jp

Abstract

Environmental DNA (eDNA) analysis is a cost-effective and non-invasive tool for species and

biodiversity monitoring in aquatic environments. Passive eDNA sampling method is focused as a

novel and alternative to conventional sampling methods, such as water filtration. In this study, we

examined the efficacy of the sponge skeleton as an eDNA passive sampler for the freshwater fish

monitoring tool. We compared the performance of passive sampling method with standard water

filtration in the river environment. A total of five different DNA extraction methods were also

conducted in three experiments, and a suitable DNA extraction method from sponge skeleton was

estimated. Quantitative fish metabarcoding using MiFish primers revealed no significant difference

in species richness between water and passive sampling methods. While both sampling methods

showed comparable trends in fish community structures, different clusters were identified for water

sampling and passive samplers by the differences in DNA concentration of each fish species. Fish

diversity in passive samples was comparable among the four DNA extraction methods using the

Direct Capture method, except for the filtration method. Our results demonstrate the efficacy of

passive eDNA sampling for monitoring freshwater fish diversity and potential use of sponge

skeletons for the absorption material of the eDNA passive sampler.

Keywords: environmental DNA, metabarcoding, passive sampler, river ecosystem, freshwater fish

Introduction

Environmental DNA (eDNA) analysis is increasingly used to monitor the biodiversity in various ecosystems. This method can detect diverse organisms by capturing, extracting DNA from excrements and extracellular DNA from different environmental samples, such as water, soil, and air (Turner et al. 2015, Tsuji et al. 2019, Lynggaard et al. 2022). The eDNA analysis is also a sensitive, non-invasive and broadly applicable tool for species detection. One of eDNA approach, eDNA metabarcoding, which can detect multiple species simultaneously (Miya et al., 2015; Deiner et al., 2017), is useful for performing rapid and large-scale multi-species surveys (Olds et al., 2016; Yamamoto et al., 2017). In particular, eDNA metabarcoding has been used in fish diversity monitoring in a variety of water bodies, ranging in size and type of aquatic ecosystems (Hänfling et al. 2016, Nakagawa et al. 2018, Oka et al. 2021). Many studies generally show that eDNA metabarcoding is a more effective and sensitive method for fish diversity assessment compared with the conventional approaches such as visual census, fyke net capturing and electrofishing (Yamamoto et a. 2017, Shaw et al. 2016).

Water eDNA, an environmental DNA collected from water samples, is most commonly used for fish diversity monitoring in the eDNA approach. Water eDNA of fishes is generally collected and concentrated from water samples by the filtration using artificial membranes (Yamamoto et al. 2017, Ushio 2019, Jeunen et al. 2019). However, fish diversity in water eDNA is often a snapshot information, and insufficient sampling effort could underestimate the fish diversity

(Zinger et al. 2019). For example, if the concentration of target DNA is low, such as rare species,

small sampling volumes and few replicates can lead to false negatives of these species.

Furthermore, since eDNA concentrations reflect the activity levels of target species (e.g., diurnal or

nocturnal), the mismatch of timing between the water sampling and fish ecology may increase false

negative detections. Large volume of water filtration and sampling replicate should be needed to

remove these false negatives and to provide reproducibility in fish diversity. However, the

conventional sampling method with water filtration requires a lot of time and cost to achieve such

improvements.

Passive eDNA collection methods, which directly submerge the sampling material in the

water body, facilitate the increase of reproducible biomonitoring as a simpler and more cost-

effective sampling method than traditional active filtration methods (Bessey et al., 2021; Kirtane et

al., 2020; Zhang et al. 2024). Absorption materials of passive eDNA sampling used in previous

studies generally have porous substances, adsorption capabilities enough to detect the target eDNA,

and simple or user-friendly form. For example, submersion materials in the passive eDNA sampling

method include activated carbon, montmorillonite (Kirtane et al., 2020), living sponge (Mariani et

al., 2019; Cai et al., 2022), and biofilm (Rivera et al., 2021). In the case of artificial materials,

Verdier et al. (2022) created the passive, 3D-printed and easy-to-use eDNA sampler using

hydroxyapatite, a natural mineral with a high DNA adsorption capacity. Yan et al. (2024) deployed

the passive sampling using GF filters by angling (short time immersion at site) and trolling (towing

in the water) method in large lake environments. These studies showed that passive collection

methods detected fish as effectively as conventional active filtration methods and provided similar

estimates of total fish biodiversity. In addition, because passive eDNA samplers require minimal or

no supporting equipment (e.g., no vacuum or peristaltic pumps), it is suitable deployment to remote

environments, and for non-experts.

In particular, natural and artificial sponges are expected to be superior eDNA passive

samplers due to their ubiquitous distribution, high regeneration ability, and ease of sampling. For

example, filter-feeding organisms have been shown to naturally accumulate eDNA in their tissue

matrices (Jeunen et al., 2021). Mariani et al. (2019) showed that eDNA extracted from alive marine

sponges can detect variety of marine fishes and mammals in the Mediterranean Sea and Antarctic

Ocean. In addition, Turon et al. (2020) highlighted the suitability and cost-effectiveness of alive

marine sponges as an eDNA passive sampler to describe tropical fish communities from coral reefs

in Southeast Asia. Jeunen et al. (2022) suggest that more porous passive filter materials, such as

artificial sponges, potentially achieve higher detection rates through an adsorption and retaining

effect of less-abundant DNA and DNA bound to particulate matter. Therefore, natural sponge

skeletons potentially achieve the effective material of passive eDNA sampler. However, knowledge

on the application of sponge skeletons to eDNA studies is insufficient, not only in marine but also

in freshwater environments, especially in rivers.

In this study, we examined the efficacy of the sponge skeleton as an eDNA passive sampler

(sponge sampler) in freshwater river environment. We submerged sponge samplers in the river and

estimated the freshwater fish diversity from the sponge eDNA samples using eDNA metabarcoding.

In addition, to examine the performance of eDNA accumulation with long-term immersion of

sponges, we also collected river water eDNA samples, the major material in eDNA surveys, in

parallel with sponge eDNA samples, and compared differences in the number of fish species and

fish species composition between passive and water samples. Furthermore, several sampling and

DNA purification methods in passive samplers were examined to estimate the optimal passive

sampling method using the sponge skeletons.

Materials and Methods

Study field and environmental DNA samplings

In this study, field surveys were conducted in the middle reaches of the Saba River Basin,

Yamaguchi Prefecture, Japan in July 30–31, and October 3–4 2021, and May 9–10 2022 (Fig.1A,

B). Fish activity times in rivers are different for species (e.g., diurnal or nocturnal), so the species

composition detected may not be identical depending on the timing of the water sampling. Sponge

passive samplers could potentially overcome these variabilities of species detection through long-

term eDNA adsorption on the sponge skeleton in the water. In order to accurately assess the

function of eDNA adsorption on the passive sampler, we carried out multi-time water sampling

during the immersion of the passive sampler. In water sampling, 3 × 1 L of surface water samples

were directly collected in the center of riffle using bleached plastic bottles every 2 hours (0, 2, 4, 6,

8, 10, 12, 14, 16, 18, 20, 22, 24 h in 2021) or 3 hours (0, 3, 6, 9, 12, 15, 18, 21, 24 h in 2022) after

submerging the equipment (Fig.1C). In this study, we started all experiments at 11.00. Therefore, in

Experiments 1 and 2 in this study, we defined 0-6 h (11:00-17:00) and 20-24 h (7:00-11:00) as

daytime and 8-18 h (19:00-5:00) as nighttime. In Experiment 3, we also defined 0-6 hours (11:00-

17:00) and 21-24 hours (8:00-11:00) as daytime and 9-18 hours (20:00-5:00) as nighttime. Water

samplings were also carried out the downstream of the equipment to prevent the contamination due

to river-bed disturbance caused by human movement. Water samples were immediately filtered on

site using a portable pump (Sentino microbiology pump; Cytiva, Tokyo, Japan) and GF/F glass

fiber filter (pore size 0.7μm; Cytiva). After filtration, used equipment was bleached to prevent DNA

cross-contamination between water samples. Filters were stored at -16 °C in the ice box and taken

back to the laboratory at 12 and 24 hours, respectively. Then, filters were stored at -20°C until DNA

extraction.

We used a specific passive sampling equipment for submerging the sponge samplers (Fig.

1B, C). This equipment consists of the weight and float connected by ropes. A sponge skeleton of

bath sponge *Spongia* spp. species was used as the adsorption material for the sponge sampler. To let

the sponges float in the middle layer of the water, sponge samplers were attached to the midpoint of

the rope tie-wraps before submerging the equipment. The equipment was placed in run near the

thalweg of the river for 24 hours for passive eDNA sampling. After 24 hours, the sponge samplers

were collected from the equipment and stored according to the specific procedures in each

experiment. After collection of the sponge samplers, the equipment was removed from the river. In

this study, two different sizes of sponges were used as the passive sampler: medium-sized sponges

(M-sponge; ca. 100 mm in diameter) and small-sized sponges (S-sponge; ca. 3–40 mm in diameter).

In this study, we compared various methods in each experiment to examine the optimal

method to recover eDNA from sponge samplers. In Experiment 1 (July 30–31, 2021), six M-sponge

samplers were used and 100 mL of pressed solutions were extracted from each samplers using a

handy juicer (Fig. 1C). Then, 100 mL of distilled water was added to the M-sponge to obtain an

additional 100 mL of pressed solution. Addition of distilled water and re-extraction were carried out

twice and a total of 300 mL of the pressings were obtained. Three of six pressed solutions were

additionally separated into 3 × 100 mL using 100 mL polyethylene bottles (M-solution). M-

solutions were stored at 4°C in a cooler box and taken back to the laboratory, and then stored at -

20°C until DNA extraction. The 300 mL of other three M-sponge solutions in Experiment 1 were

filtered with GF/F and filters were stored at -16 °C in the ice box and taken back to the laboratory,

and then stored at -20°C until DNA extraction (M-filter). In Experiment 2 (October 3–4, 2021), six

M-sponge samplers were used and all pressed solutions were handled as M-solution samples and

stored in the same methods as in Experiment 1 (Fig. 1C). In Experiment 3 (May 21–22, 2022), four

different passive sampling methods were examined. Three M-sponge samplers and six S-sponge

samplers were submerged in the river and collected after 24 hours. In Experiment 3, 100 mL of first

pressed solutions were only extracted from each M-sponge sampler and used as M-solution2

samples (Fig. 1C). Six S-sponge samplers were collected from the equipment and stored in separate

syringes. Three of the six S-sponge samplers were pressed in syringes and separated into syringe

samples (S-syringe) and pressed solutions (S-solution). Approximate 50 mL S-solution sample was

extracted from each S-sponge sampler and collected using 100 mL polyethylene bottles. The other

three S-sponge samplers were stored in syringes without pressing (S-syringe-NP). In Experiment 3,

all passive samples were stored at 4°C in a cooler box and taken back to the laboratory, and then

stored at -20°C until DNA extraction.

DNA extraction from water and sponge samples

Total eDNA of filter sample was extracted and purified following by Tsuji et al. (2022).

Briefly, each filter was placed in the upper part of a salivette tube (Sarstedt, Nümbrecht, Germany)

and pre-centrifuged at 5000 ×g for 1 min to remove the excess water. Each filter was soaked with

440 μL of a mixture composed of 200 μL Buffer AL, 200μL of H_2O and 20 μL of proteinase K, and

were incubated at 56°C for 45 min. Afterward, the cartridges were centrifuged at 5000 ×g for 3 min,

and the extract solutions of each tube were collected. Total eDNA was purified from extract

solution using DNeasy Blood & Tissue Kit (QIAGEN, Hilden, Germany) and was eluted in 100 μL

buffer AE from spin column according to the manufacturer's instructions.

Total eDNA of solution samples in passive sampling methods (M-solution and S-solution)

was extracted and purified using Maxwell® RSC Enviro Total Nucleic Acid Kit (Promega,

Madison, WI, USA) according to the manufacturer's instructions (hereafter, Direct Capture (DC)

method). Briefly, 12.5 µL of proteinase solution from the DC kit was added per 1 mL of pressed

solution, and samples were incubated for 30 min at ambient temperature. The samples were then

centrifuged at 3000 xg for 10 min at room temperature to remove the precipitation. The supernatant

was divided equally into two 50 mL tube and 6.0 mL of Binding Buffer 1 and 0.5 mL of Binding

Buffer 2 were added to the supernatant, respectively. The 24 mL of isopropanol was added to the

mixture, gently mixed and passed through a PureYield Midi Binding Column using a VacMan

Vacuum Manifold. Nucleic acid captured on the PureYield Midi Binding Column was washed with

5 mL of Column Wash 1 followed by 20 mL of Column Wash 2. Nucleic acid was eluted with 750

µL of pre-warmed (60°C) nuclease-free water. Then, the eluate was transferred to a Maxwell® RSC

Cartridge and the cartridge was held in a Maxwell® deck tray. DNA purification and elution of

nuclease-free water were performed on a Maxwell® instrument according to the manufacturer's

instructions. After completed procedures, the eluate was collected from the deck tray and 100 µL of

eDNA solution was obtained. UV light treatment was then performed on the instrument to avoid the

cross-contamination among samples.

Total eDNA of syringe samples in passive sampling methods (S-syringe and S-syringe-NP)

was extracted and purified using the extraction kit same as sponge solution samples with the slight

modification in the incubation step). Briefly, proteinase K powder (approximate 30 units/mg;

Nacalai tesque, Kyoto, Japan) was dissolved in TE buffer (pH 8.0) to prepare the proteinase K stock

solution (10 units/mL). Then, DNA elution mixture was prepared by adding 2.0 mL of Proteinase K

stock solution to 18 mL of TE buffer (pH 8.0). S-syringe sample was submerged in this solution in

each syringe, and incubated for 45 min at 56°C. The solution sample was discharged to a 50 mL

tube from each syringe and 6.0 mL of Binding Buffer 1 and 500 μL of Binding Buffer 2 were added

to the sample, respectively. The samples were then centrifuged at 3000 xg for 10 min at room

temperature to remove the precipitation. Following protocols (DNA purification, elution and

collection) were same as that of sponge solution samples. All eDNA samples were stored at -20°C

until 1st polymerase chain reaction (PCR) step in the eDNA metabarcoding.

Quantitative fish environmental DNA metabarcoding

In this study, we used the quantitative eDNA metabarcoding (qMiSeq, Ushio et al. 2019)

method to identify fish species composition and to quantify fish eDNA concentration,

simultaneously. In general, sequence reads obtained from high-throughput sequencing does not

necessarily correspond to the eDNA concentration because of the differences in PCR efficiency

(primer bias, PCR inhibition, etc.). In qMiSeq method, sequence reads can be converted to DNA

concentration (DNA copies/template) using sample-specific standard regression lines.

Amplicon libraries were prepared according to the following protocols (detailed

information in Supplementary Information S1). Fish eDNA of each sample was amplified using

MiFish primers (MiFish-U, -O, and -L; Miya et al. 2020), and internal standard DNA mix (5, 25, 50

copies /template) was added in each PCR reaction mix. The first PCR was repeated eight times for

each sample, and the replicated samples were pooled as a single first PCR product for use in the

subsequent step. In the first PCR step, the PCR negative controls (eight replicates) were included in

each experiment. In second PCR, the 8-bp dual indices were added to the purified first PCR

amplicons. The second PCR amplicons were pooled and the target library size (ca. 370 bp) of

second PCR was collected using E-Gel SizeSelect agarose gels (Thermo Fisher Scientific). Purified

library was sequenced using iSeq i1 Reagent and iSeq 100 platform with 30% PhiX control

(Illumina, San Diego, CA, USA). Sequencing dataset output from iSeq platform was subjected to

bioinformatic analysis. After bioinformatic analysis, sequence reads of each fish species and

standard DNA mix in each sample were obtained and subjected to qMiSeq analysis. To obtain

sample-specific regression lines, the regression analysis was performed using sequence reads and

DNA concentration of internal standard DNA mix. Slopes and R^2 values of each regression line

were shown in Supplementary Table S1–3. The R^2 values of all regression lines were higher than

0.98. In this study, no obvious PCR inhibition was identified in all water and passive DNA samples.

Statistical analysis

In this study, differences in the number of detected species among the methods (water and

each passive sampling) were assessed in each experiment. In each experiment, to examine the

difference in the diurnal conditions, the number of species in water samples was pooled together

into three categories: water (all), water (day) and water (night). The water (day) contains daytime

samples (0–6h and 20–24h), water (night) contains nighttime samples (8h–18h), and water (all)

contains all water samples (day and night). Kruskal-Wallis test was conducted to examine

differences in the number of species among all survey protocols (three conditions of water samples

and passive sampling methods) in each experiment. Then, Steel-Dwass multiple comparison was

performed in the experiment with significant differences.

Using fish eDNA concentration dataset, non-metric Multidimensional Scaling (nMDS) in

1000 separate runs was performed to visualize differences in fish community structure among

sampling methods. Stress value was used to measure the indicator of nMDS fitting. Threshold

stress values in the nMDS were less than 0.2, and the lowest of result was chosen. In addition,

permutational multivariate analysis of variance (PERMANOVA) were used to examine the

statistical differences in fish community structure among the sampling methods, and permutational

analysis of multivariate dispersions (PERMDISP) were used to assess the variation in species

composition among the sampling methods. Bray-Curtis dissimilarity of fish community structure

between samples in each experiment was calculated and β-diversity plots of each samples were

visualized in each experiment. In each experiment, fish species that affected differences in

community structure between sampling methods were estimated using "envfit" function. In this

study, the threshold of p-value was <0.01 in order to choose species which have more obvious

effects on community structure. All statistical analyses and plotting were performed using R ver.

4.02. For nMDS, PERMANOVA and PERMDISP, the "metaMDS", "adonis" and "betadisper"

functions of the vegan package were used.

Results

In this study, 8,296,115 reads were obtained from 105 water samples and 41 passive

samples, and 6,903,093 reads were remained after pre-processing (Supplementary Table S4). The

number of species detected in Experiment 1 (July, 2021) ranged from 15–22 for water samples (n =

39) and 7–14 (M-filter, n = 3) and 15–22 (M-solution, n = 8) for passive samples, respectively (Fig.

2). Due to the loss of one of the M-solution samples from Experiment 1 (M-solution 3-3) from the

equipment, the total number of M-solution samples in Experiment 1 was eight (Supplementary

Table S1). The number of species detected in Experiment 2 (October, 2021) ranged from 15–23 for

water samples (n = 39) and 19–25 for M-solution samples (n = 18). In Experiment 3 (May, 2022),

the number of species ranged from 18–24 for water samples (n = 27) and 21–23 (M-solution, n = 3),

21–23 (S-solution, n = 3), 16–19 (S-syringe, n = 3) and 19 (S-syringe-NP, n = 3) for passive

samples, respectively (Fig. 2). Steel-Dwass multiple comparison was performed, as the significant

differences were confirmed in results of Kruskal-Wallis test in all experiments (Supplementary

Table S5). However, significant differences were only confirmed in Experiment 1 in the results of

the Steel-Dwass test (Fig. 2). In Experiment 1, significant differences were observed between the

number of species detected from M-filter and that of water (all), water (day), water (night) and

between water (day) and water (night), respectively ($p < 0.05$). In Experiment 2, the number of

species detected from M-solution sample tended to be higher than water (all) and water (day),

although the differences were not statistically significant ($p = 0.055$ and 0.072). In Experiment 3,

there were no statistically significant differences in the number of species among all sample types

(Supplementary Table S5).

Cumulative fish diversity was highest for water sampling accumulated between 0 and 24

hours (3 samples × 13 timings) than passive sampling methods in all Experiments (Fig. 3).

Specifically, cumulative number of fish species in Experiment 1 was 25, 15, 23, 24 and 19 in water,

M-filter, M-solution 1, 2, and 3 sample, respectively. Cumulative number of fish species in

Experiment 2 was 28 and 31 in water and M-solution sample. In Experiment 3, cumulative number

of fish species was 26, 23, 24, 21 and 21 in water, M-solution, S-solution, S-syringe and S-syringe-

NP sample, respectively. Passive sampling methods also showed high cumulative fish diversity,

similar to water, with the M-solution in Experiment 2 in particular showing the highest diversity

(Supplementary Table S1–3).

Fish species composition was substantially similar in all experiments (Fig. 4). The number

of common species between water (day) or water (night) and passive samples (M-solution, S-

284 solution, S-syringe or S-syringe-NP) was much higher than other combinations in all experiments.

285 However, common fish species in the two water samples (9 species) were not detected in M-filter

286 samples in Experiment 1 (Fig. 4). On the other hand, the comparison with M-solution showed a

287 decrease in the number of species detected only in the two water samples and an increase in the

288 number of species common to the three methods, compared to the M-filter. Cumulative fish species

289 compositions detected by each sampling method in Experiments 1-3 are listed in Supplementary

290 Table S6. In Experiment 1, *Oncorhynchus masou* subsp. was specifically detected from water (day)

291 and *Pseudogobio agathonectris* in passive samples (M-filter and M-solution). In Experiment 2,

292 three fish species (*Acheilognathus rhombeus*, *Hemibarbus* sp. and *Salvelinus leucomaenis* subsp.)

293 were only detected in M-solution samples and the species detected only in two types of water

294 samples were not observed. In Experiment 3, *Tanakia limbata* was only detected in water (day)

295 samples and *Hemibarbus* sp. in S-syringe samples. In Experiment 3, the number of common species

296 between the two types of water sample and four passive sampling methods was much similar. Fish

297 species detected by eDNA metabarcoding were generally known to inhabit the study site in the

298 previous years. (River Environmental Database system, 2018). However, a tropical garfish

299 *Atractosteus tropicus*, listed as alien species in Japan, was newly detected with high DNA

300 concentration from water (night), M-solution, S-solution samples.

301   In this study, nMDS plot was used to assess the differences in community structure

302 considering the eDNA concentration of each fish species (Fig. 5). Fish community structure in

Experiment 1 was much different among sample types (PERMANOVA: $p < 0.01$, PERMDISP: $p =$

0.441). In particular, M-filter samples showed greater variation among samples than the other

sample types. While the number of species in the M-solution samples was similar to the two types

of water samples, a few differences were observed in the community structure. Number of fish

species affecting the differences in community structure was 8 species, and 2 of these species

(*Hemibarbus longirostris* and *Tachysurus nudiceps*) were not detected in M-filter samples. Other

species were detected in all sample types, but DNA concentrations of these species were much

higher in water samples than in passive samples (Supplementary Table S1). In Experiment 2, the

fish community structure of water (night) was included in that of water (day) and the M-solution

fish community was plotted outside the two types of water samples. However, the differences in

community structure among the three types could not be precisely assessed (PERMANOVA: $p <$

0.01, PERMDISP: $p < 0.05$). There could not be also assessed the differences in community

structure between water (all) and M-solution samples (PERMANOVA: $p < 0.01$, PERMDISP: $p <$

0.05). Number of fish species affecting the differences in community structure was 14 species in

Experiment 2, and *Hemibarbus* sp. was only detected in M-solution samples. Other species were

detected in all sample types, and trend in DNA concentration of these species was similar to

Experiment 1 (Supplementary Stable S2). In Experiment 3, there were slight differences in fish

community structure between water (day) and water (night) samples. Also in the passive samples,

the community structure was different between the solution types (M-solution and S-solution) and

the syringe types (S-syringe and S-syringe-NP). In Experiment 3, Fish community structure was

much different among sample types (PERMANOVA: $p < 0.01$, PERMDISP: $p = 0.11$). Number of

fish species affecting the differences in community structure was 13 species in Experiment 3. All

species were detected in all sample types, except for Misgurnus anguillicaudatus in S-syringe

samples. Trend in DNA concentration of these species was also similar to Experiment 1 and 2

(Supplementary Table S3).

Discussion

In this study, we assessed the effectiveness of eDNA passive sampling methods compared

to water sampling in riverine environment. In addition, in the passive sampling method, some DNA

extraction methods were examined to estimate the optimal eDNA recovery method from sponges.

Our results showed that the passive sampling method using sponge skeletons had similar or higher

number of fish species than water samples. Our results in the present study would support previous

studies showing similar trends (Jaunen et al. 2024, Zhan et al. 2024, Mlinarec et al. 2025).

However, in Experiment 1, the number of species of M-filter sample was significantly lower than

other methods (water and other passive samples) and the variation of fish community composition

was higher than other methods. However, as no PCR inhibition was observed in all DNA samples,

other factors may have affected the reduction in the number of species in the M-filter samples.

These suggest that DNA recovery rate of M-filter was less efficient than other methods. Water

samples pressed from sponges contain large amount of material besides eDNA (e.g., sand, mud,

biofilm, etc.) and DNA in pressed solutions could be captured to these materials. This can easily be

assumed from the work of previous studies using soil particles and biofilms as the capturing

materials of passive sampling (Kirtane et al., 2020, Rivera et al., 2021). DNA extraction method

from GF/F filter used in the M-filter might not have efficiently recovered the eDNA captured on

these materials. On the other hand, DC method applied to passive samplers in this study suggests

that efficiency of DNA recovery was improved by direct DNA extraction from pressed solutions or

by direct immersion of sponge skeletons in the elution mixture. In fact, M-solution in the passive

sampling method showed similar fish diversity in all experiments, both in the number of species

and the cumulative number of species, as in the water samples. Furthermore, in Experiment 2, fish

species detected only in M-solution were observed. The cumulative number of samples in passive

sampling methods was larger than for water samples (Fig.4). These results show that passive

samples (sponge skeletons) of 24-hour immersion have the same detection performance as 24-hour

water samples taken every two hours. Slightly lower number of species were observed in other

passive sampling methods (S-solution, S-syringe and S-syringe-NP) compared to water samples,

but the statistically significant differences were not found. The high cumulative fish diversity for S-

solution and the presence of specific species for S-syringe, respectively, may also indicate the

effectiveness of each method (Fig. 3, Fig.4). However, the passive sampling method in Experiment

3 could have been difficult to find the difference from other methods because of the lack of

replicate samples (triplicate in each sample). Therefore, the effectiveness of these methods may

need to be carefully evaluated with further work.

In Experiments 1 and 3, the number of detected species at night-time tended to be higher

than that of daytime in the water samples (Fig.3). Cumulative number of species in the water

samples in Experiments 1 and 2 varied considerably with collection timing (Fig.4). On the other

hand, the number of detected and cumulative species in passive samples, except for the M-filter

sample, tended to be more homogeneous than in water samples in all experiments. These

differences could be due to dissimilarities in the DNA information of freshwater fishes in the water

and passive samples. In general, eDNA information on water samples contains the snapshot

information reflecting the collection timing (Spear et al. 2015, Bowers et al., 2021; Jensen et al.,

2022). Therefore, eDNA shed from rare species or species with coincidental appearance could be

rapidly diluted and dispersed, and elude detection via instantaneous water sampling (Kirtane et al.,

2020). On the other hand, passive sampling methods could have accumulated more eDNA by

filtering the large water volume through long-term immersion in the water. Jeunen et al. (2022)

indicated that more porous passive filter materials potentially achieve higher detection rates through

an adsorption and retaining effect of low-abundant DNA and DNA bound to particulate matter.

Differences in temporal resolution between sampling methods could be reflected in such variations

and differences in fish species diversity.

Experiments 1 and 3 showed differences in fish community structure between methods in

the nMDS plot and statistical analyses. In all experiments, some fish species also affected

differences in community structure among sample types. In Experiment 1, this is reasonable as fish

diversity and DNA concentrations of each fish species in the M-filter samples were much lower

than other methods. However, in other experiments, the fish diversity and species composition are

few different among sampling methods (Fig.3, Fig. 4). Therefore, these variations in community

structure among methods could be attributed to differences in the eDNA concentration in each fish

species. In fact, there were differences in fish DNA concentrations among sampling methods, and

that of common fish species (*Zacco platypus*, *Nipponocypris temminckii* and *Pungtungia herzi*)

observed in the survey site showed large differences between water and passive samples in all

experiments (Supplementary Table 1–3). As fish DNA concentration of each sample was

standardized by adding the standard DNA mix, these differences indicate that eDNA concentration

of fishes in passive samples is lower than in water samples. Passive sampling methods store the

aquatic eDNA by capturing and accumulating the collection materials. Cananzi et al. (2025) showed

that passive samplers contain eDNA of diverse taxa and particle sizes and are more efficient for

detecting micro-invertebrates and amphibians compared to water samples by using multi-marker

approach. Similarly in our study, large amounts of eDNA from non-fish organisms (e.g., aquatic

plants, insects and micro-organisms) may have been captured by sponge skeletons during long-term

immersion. As a result, fish eDNA could be relatively reduced in passive samples, causing the

differences in proportion of DNA concentration between water samples and passive samples.

Differences in eDNA concentrations may also reflect differences in DNA extraction efficiency

between sponge skeletons and GF/F filters. While optimization of DNA extraction procedures from

GF/F filters has been examined by previous studies (Minamoto et al. 2021), that of sponge

skeletons may still be insufficient. Previous studies have also shown that the efficiency of DNA

recovery in passive samples varied greatly depending on the sampler material and immersion

method (Verdier et al. 2022, Saltonstall et al. 2024, Stevens et al. 2024). Therefore, future

improvements in methodology will enhance the efficacy of the passive sampling method using

sponge skeletons.

In conclusion, we showed that the eDNA passive sampling method using sponge skeletons

is one of the most efficient methods for monitoring freshwater fishes in riverine environments.

Passive sampling methods enable more cost-effective and efficient eDNA monitoring than

conventional filtering methods. Passive sampling methods also requires no need for time-

consuming filtration or the preservation of large volumes of water until filtering. Furthermore,

passive sampling methods minimize sampling and processing time in the field and accumulate

eDNA with long-term immersion, and enable time-efficient biomonitoring. Indeed, our passive

sampling method could detect similar or higher fish species diversity than that of water sampling

due to long-term and cumulative eDNA adsorption on sponge skeletons. However, there are some

challenges in the passive sampling method. One of the challenges of passive sampling method is

that the actual amount of water passing through the passive collection device during submersion is

difficult to quantify (Zhang et al. 2024). Currently, this method may be unsuitable for estimating

eDNA concentration of target species and quantitative assessments of species abundance and

biomass. In this study, eDNA concentration of each fish species was estimated by qMiSeq method,

but the accuracy of DNA quantification could not be evaluated. This challenge could be solved by

measuring flow rates and by estimating the water volume through the sponge skeleton.

Furthermore, passive sampling methods could concentrate PCR inhibitors as well as eDNA with

long-term immersion. Organic compounds (e.g., humic and fulvic acid) produced by biological

processes of phytoplankton and from terrestrial detritus inputs were contained in aquatic systems,

and can be the major cause of PCR inhibition (Hellings et al. 1999, Albers et al., 2013, Uchii et al.

2019). These substances may be adsorbed and concentrated on the sponge skeleton as well as

eDNA, increasing the probability of PCR inhibition. Further improvements of methodology, such as

the effect of PCR inhibition, quantitative assessment, and immersion period, should be required so

that passive sampling method using sponge skeletons will be an effective tool for the eDNA survey.

Acknowledgements

We thank students of Akamatsu laboratory in Yamaguchi University for supporting the field

experiments. We also thank Saba River Fisheries Cooperative and Yamaguchi River and National

Highway Office, Chugoku Regional Development Bureau, Ministry of Land, Infrastructure,

Transport and Tourism for the permission of field surveys. This work was supported by Grant-in-

Aid for Challenging Research (Exploratory) of Japan Society for the Promotion of Science (Grant

ID: 22K18298).

Conflicts of interest

We have read the journal's policy and the authors of this manuscript have the following competing

interests: F.I. is currently a paid employee of Nippon Koei Co., Ltd. We further declare that the

methodology and device of the passive sampling method using sponge skeletons has a patent

pending under the application number PCT/JP2023/036721.

Data accessibility

All sequence data will be published in DRA (Accession number:) after accept of the manuscript and

detailed information of fish community structure and methodological descriptions of library

preparation are available in the Supplementary materials.

References

Albers, C.N., Jensen, A., Bælum, J., Jacobsen, C.S. (2013) Inhibition of DNA polymerases used in

Q-PCR by structurally different soil-derived humic substances. Geomicrobiol. J. 30, 675–681.

https://doi.org/10.1080/01490451.2012.758193

Bessey, C., Jarman, S. N., Simpson, T., Miller, H., Stewart, T., Keesing, J. K., & Berry, O. (2021).

Passive eDNA collection enhances aquatic biodiversity analysis. Communications Biology, 4,

236. https://doi. org/10.1038/s42003-021-01760-8

Bowers, H. A., Pochon, X., von Ammon, U., Gemmell, N., Stanton, J. A. L., Jeunen, G. J.,

Sherman, C. D. H., Zaiko, A. (2021) Towards the optimization of eDNA/eRNA sampling

Technologies for Marine Biosecurity Surveillance. Water, 13(8), 13081113.

https://doi.org/10.3390/w13081113

Cai, W., Harper, L. R., Neave, E. F., Shum, P., Craggs, J., Arias, M. B., Riesgo, A., & Mariani, S.

(2022). Environmental DNA persistence and fish detection in captive sponges. Molecular

Ecology Resources, 22, 2956–2966. https://doi.org/10.1111/1755-0998.13677

Cananzi, G., Tatini, I., Li, T., Montagna, M., Serra, V., Petroni, G. (2025) Active or passive? A

multi-marker approach to compare active and passive eDNA sampling in riverine environments.

Science of The Total Environment, 974, 179247. https://doi.org/10.1016/j.scitotenv.2025.179247

Hänfling, B., Handley,L.L., Read, D.S., Hahn, C., Li, J., Nichols, P., Blackman, R.C., Oliver, A.,

Winfield, I.J. (2016) Environmental DNA metabarcoding of lake fish communities reflects long-

term data from established survey methods. Molecular Ecology, 25, 3101–3119.

https://doi.org/10.1111/mec.13660

Hellings, L., Dehairs, F., Tackx, M., Keppens, E., Baeyens, W. (1999) Origin and fate of organic

carbon in the freshwater part of the Scheldt Estuary as traced by stable carbon isotope

composition. Biogeochemistry 47, 167–186. https://doi.org/ 10.1023/A:1006143827118

Jensen, M. R., Sigsgaard, E. E., Ávila, M. de P., Agersnap, S., Brenner-Larsen, W., Sengupta, M.

E., Xing, Y., Krag, M. A., Knudsen, S. W., Carl, H., Møller, P. R., Thomsen, P. F. (2022) Short-

term temporal variation of coastal marine eDNA. Environmental DNA, 4, 747–762.

https://doi.org/10.1002/edn3.285

Jeunen, G.J. Knapp, M., Spencer, H.G., Taylor, H.R., Lamare, M.D., Stat, M., Bunce, M.,

Gemmell, N.J. (2019) Species-level biodiversity assessment using marine environmental DNA

metabarcoding requires protocol optimization and standardization. Ecology and Evolution, 9,

1323–1335. https://doi.org/10.1002/ece3.4843

Jeunen, G.J., Mills, S., Mariani, S., Treece, J., Ferreira, S., Stanton J.A.L., Dur´an-Vinet, B.,

Duffya, G.A., Gemmell, N.J., Lamare, M. (2024) Streamlining large-scale oceanic biomonitoring

using passive eDNA samplers integrated into vessel's continuous pump underway seawater

systems. Sci. Total Environ., 946, 174354. https://doi.org/10.1016/j.scitotenv.2024.174354

Jeunen, G.J., von Ammon, U., Cross, H., Ferreira, S., Lamare, M., Day, R., Treece, J.,Pochon, X.,

Zaiko, A., Gemmell, N. J., Stanton, J.A. (2022) Moving environmental DNA (eDNA)

technologies frombenchtop to the field using passive sampling and PDQeXextraction.

Environmental DNA, 4, 1420–1433. https://doi.org/10.1002/edn3.356

Kirtane, A., Atkinson, J. D., & Sassoubre, L. (2020). Design and validation of passive

environmental DNA samplers using granular activated carbon and montmorillonite clay.

Environmental Science and Technology. Environmental Science & Technology, 54, 11961–

11970. https://doi.org/10.1021/acs.est.0c01863

Lynggaard, C., Bertelsen, M.F., Jensen, C.V., Johnson, M.S., Frøslev, T.G., Olsen, M.T., Bohmann,

K. (2022) Airborne environmental DNA for terrestrial vertebrate community monitoring. Current

Biology, 32, 701–707. https://doi.org/10.1016/j.cub.2021.12.014

Mariani, S., Baillie, C., Colosimo, G., & Riesgo, A. (2019) Sponges as natural environmental DNA

samplers. Current Biology, 29(11), R401–R402. https://doi.org/10.1016/j.cub.2019.04.031

Minamoto, T., Miya, M., Sado, T., Seino, S., Doi, H., Kondoh, M., Nakamura, K., Takahara, T.,

Yamamoto, S., Yamanaka, H., Araki, H., Iwasaki, W., Kasai, A., Masuda, R., Uchii, K. (2021)

An illustrated manual for environmental DNA research: Water sampling guidelines and

experimental protocols. Environmental DNA 3, 8-13 https://doi.org/10.1002/edn3.121

Ministroy of Land, Infrastructure, Transport and Tourism, Japan. (2018) Chugoku District in River

Environment Database system. https://www.nilim.go.jp/lab/fbg/ksnkankyo/dl_87_index.html

(referred in October 10, 2022)

Miya, M., Gotoh, R.O., Sado, T. (2020) MiFish metabarcoding: a high-throughput approach for

simultaneous detection of multiple fish species from environmental DNA and other samples.

Fisheries Science, 86, 939–970. https://doi.org/10.1007/s12562-020-01461-x

Mlinarec, J., Svetličić, I., Kresonja, M., Rubinić, M., Megyery, T., Kaliger, I., Mihaljević, D. (2025)

Temporal and Spatial Dynamics of the Lotic Fish Communities: A Comparison of Coffee Filter-

Based Passive eDNA Collection Versus Active eDNA Filtering. Environmental DNA, 7.

https://doi.org/10.1002/edn3.70065o

Nakagawa, H., Yamamoto, S., Sato, Y., Sado, T., Minamoto, T., Miya M. (2018) Comparing local-

and regional-scale estimations of the diversity of stream fish using eDNA metabarcoding and

conventional observation methods. Freshwater Biology, 63, 569–580.

https://doi.org/10.1111/fwb.13094

Oka S.I., Doi, H., Miyamoto, K., Hanahara, N., Sado ,T., Miya, M. (2021) Environmental DNA

metabarcoding for biodiversity monitoring of a highly diverse tropical fish communityin a coral

reef lagoon: Estimation of species richness and detection of habitat segregation. Environmental

DNA, 3, 55–69. https://doi.org/10.1002/edn3.132

Rivera, S. F., Rimet, F., Vasselon, V., Vautier, M., Domaizon, I., & Bouchez, A. (2021). Fish

eDNA metabarcoding from aquatic biofilm samples: Methodological aspects. Molecular Ecology

Resources, 22, 1440–1453. https://doi. org/10.1111/1755-0998.13568

Saltonstall, K., Delgado, J., Vargas, M., Collin, R. (2024) Are passive collectors effective samplers

of microbes in natural aquatic systems? Front. Freshw. Sci., 2, 1460713.

https://doi.org/10.3389/ffwsc.2024.1460713

Shaw, J.L., Clarke, L.J., Wedderburn, S.D., Barnes, T.C., Weyrich, L.S., Cooper, A. (2016)

Comparison of environmental DNA metabarcoding and conventional fish survey methods in a

river system. Biological Conservation, 197, 131–138.

https://doi.org/10.1016/j.biocon.2016.03.010

Stevens, E.R., Hyde, J., Beesley, L.S., Gwinn, D.C., Thompson, S., Morris, L., Wilson, P.R.,

Gleeson, D.B. (2024) Fishy Business—Assessing the Efficacy of Active and Passive eDNA to

Describe the Fish Assemblage of a River in Southwestern Western Australia to Support Effective

Monitoring. Environmental DNA, 6, 6. https://doi.org/10.1002/edn3.70040

Tsuji, S., Inui, R., Nakao, R., Miyazono, S., Saito, M., Kono, T., Akamatsu, Y. (2022) Quantitative

environmental DNA metabarcoding shows high potential as a novel approach to quantitatively

assess fish community. Scientific Reports, 12, 21524. https://doi.org/10.1038/s41598-022-25274-

3

Tsuji, S., Takahara, T., Doi, H., Shibata, N., Yamanaka, H. (2019) The detection of aquatic

macroorganisms using environmental DNA analysis—A review of methods for collection,

extraction, and detection. Environmental DNA, 1, 99–108. https ://doi.org/10.1002/edn3.21

Turner, C.R., Uy, K.L., Everhart, R.C. (2015) Fish environmental DNA is more concentrated in

aquatic sediments than surface water. Biological Conservation, 183, 93–102.

https://doi.org/10.1016/j.biocon.2014.11.017

Turon, M., Angulo-Preckler, C., Antich, A., Præbel, K., & Wangensteen, O. S. (2020). More than

expected from old sponge samples: A natural sampler DNA metabarcoding assessment of marine

fish diversity in Nha Trang Bay (Vietnam). Frontiers in Marine Science, 7, 1–14.

https://doi.org/10.3389/fmars.2020.605148

Uchii, K., Doi, H., Okahashi, T., Katano, I., Yamanaka, H., Sakata, M.K., Minamoto, T. (2019)

Comparison of inhibition resistance among PCR reagents fordetection and quantification of

environmental DNA.Environmental DNA. 1:359–367. https://doi.org/10.1002/edn3.37

Ushio, M. (2019) Use of a filter cartridge combined with intra-cartridge bead-beating improves

detection of microbial DNA from water samples. Methods in Ecology and Evolution, 10, 1142–

1156. https://doi.org/10.1111/2041-210X.13204

Ushio, M., Murakami, H., Masuda, R., Sado, T., Miya, M., Sakurai, S., Yamanaka, H., Minamoto,

T., Kondoh, M. (2018) Quantitative monitoring of multispecies fish environmental DNA using

high-throughput sequencing. Metabarcoding and Metagenomics, 2, e23297.

https://doi.org/10.3897/mbmg.2.23297

Verdier, H., Konecny-Dupre, L., Marquette, C., Reveron, H., Tadier, S., Grémillard, L., Barthès, A.,

Datry, T., Bouchez, A., & Lefébure, T. (2022). Passive sampling of environmental DNA in

aquatic environments using 3D-printed hydroxyapatite samplers. Molecular Ecology Resources, 22, 2158–2170. https://doi. org/10.1111/1755-0998.13604

Yamamoto, S., Masuda, M., Sato, Y., Sado, T., Araki, H., Kondoh, M., Minamoto, T., Miya, M. (2017) Environmental DNA metabarcoding reveals local fish communities in a species-rich coastal sea. Sci Rep, 7, 40368.

Yan, Z., Luo, Y., Chen, X., Yang, L., Yao, M. (2024) Angling and trolling for eDNA: A novel and effective approach for passive eDNA capture in natural waters. Environmental International, 194, 109175. https://doi.org/10.1016/j.envint.2024.109175

Zhang, L., Zhou, W., Jiao, M., Xie, T., Xie, M., Li, H., Suo, A., Yue, W., Ding, D., He, W. (2024) Use of passive sampling in environmental DNA metabarcoding technology: Monitoring of fish diversity in the Jiangmen coastal waters. Sci. Total Environ., 908, 168298. https://doi.org/10.1016/j.scitotenv.2023.168298

Zinger, L., Bonin, A., Alsos, I. G., Balint, M., Bik, H., Boyer, F., Chariton, A. A., Creer, S., Coissac, E., Deagle, B. E., De Barba, M., Dickie, I. A., Dumbrell, A. J., Ficetola, G. F., Fierer, N., Fumagalli, L., Gilbert, M. T. P., Jarman, S., Jumpponen, A., ... Taberlet, P. (2019). DNA me-tabarcoding – Need for robust experimental designs to draw sound ecological conclusions. Molecular Ecology, 28, 1857–1862.

Fig.1

582 **Figure 1**

583 Study site information and descriptions of experimental design in this study. (A) descriptions of

584 survey site, (B) detailed information of survey site and passive sampler, (C) Methodological

585 procedures of water and passive sampling in each Experiment.

Figure 2

Boxplots in the number of detected species in each experiment. Asterisks indicate the significant

differences among sample categories (**: $p < 0.01$, *: $p < 0.05$).

Fig.3 Cumulative number of species detected in each collection timing and DNA extraction method.

Figure 4

Venn-diagrams showing the number of fish species belonging to each sampling method that were

detected by eDNA metabarcoding in each Experiment. Detailed information of fish species

compositions in each Experiment was described in Supplementary Table S6.

Figure 5 The nMDS plots in each experiment. Green arrows represent the fish responsible for

causing the significant difference among sampling methods.