Preliminary result

AsCas12a Exhibits Intrinsic, DNA-Independent ATPase Activity

Supreet Bhattacharya[1]

1-Department of Biochemistry, Indian Institute of Science, Bangalore, India

Correspondence to Supreet Bhattacharya, ORCID ID- 0000-0002-9880-5127

Abstract

Cas12a (Cpf1) is a class 2 CRISPR-Cas effector protein with RNA-guided DNA endonuclease activity widely used for genome editing. While its DNA cleavage and target recognition mechanisms have been studied extensively, the possibility of auxiliary enzymatic functions remains underexplored. Here, I report that *Acidaminococcus* sp. Cas12a (AsCas12a) possesses intrinsic ATPase activity, despite lacking canonical nucleotide-binding or hydrolysis motifs. Using a radiometric thin-layer chromatography (TLC) assay, I demonstrate that AsCas12a hydrolyzes ATP in a concentration and time-dependent manner. Importantly, this activity occurs independently of DNA cofactors, as neither single-stranded nor double-stranded DNA influenced the rate or extent of ATP hydrolysis. Bioinformatic analyses using NsitePred and SwissDock identified potential ATP-binding residues with predicted favorable binding energies. This preliminary finding uncovers a previously unrecognized biochemical property of AsCas12a and raises questions regarding the physiological role of this ATPase activity in CRISPR function.

 Keywords: Cas12a, CRISPR-Cas, ATPase, DNA-independent activity, enzyme mechanism, AsCas12a

1. Introduction

Cas12a (formerly Cpf1) is a type V-A CRISPR-associated endo- and exonuclease that has garnered significant attention for its programmable DNA cleavage ability in genome editing and diagnostics [1–3]. Like Cas9, Cas12a is guided by a CRISPR RNA (crRNA) to introduce double-stranded breaks at target sites. While its structural domains and DNA targeting functions have been well characterized [4–6], auxiliary enzymatic activities associated with Cas12a remain relatively unexplored. ATP hydrolysis plays critical roles in the function of many DNA-processing enzymes, including helicases, translocases, and remodeling factors [7–9]. However, Cas12a lacks conserved nucleotide-binding motifs such as Walker A or B, raising the question of whether it could still exhibit ATPase activity through non-canonical mechanisms [2]. Previous studies on CRISPR-associated proteins have uncovered unexpected enzymatic functions, prompting us to investigate whether AsCas12a could similarly possess cryptic enzymatic activities. In this study, I used a radiolabeled ATP hydrolysis assay to test

AsCas12a's ability to catalyze ATP turnover in the presence and absence of DNA cofactors. The ATPase activity of the AsCas12a protein was discovered serendipitously-when in one of the assay to test for *E. coli.* RecA activity, it was kept as a control (10). Further, bioinformatics analysis identified a total of six ATP-binding residues in AsCas12a, which needs to be validated further through other experiments. Overall, the findings reveal a DNA-independent ATPase activity intrinsic to AsCas12a, suggesting potential auxiliary roles beyond its endonuclease function.

2 Materials and Methods

2.1 Purification of AsCas12a

AsCas12a was purified as previously described (1).

2.2 Preparation of DNA substrates

The sequences of oligonucleotides (ODNs) used in this study are listed in Table 1. ODNs were 5′ end-labelled with [γ-^{32}P] ATP using T4 polynucleotide kinase, and DNA substrates were made as described previously (Table 2, ref. 11).

Oligonucleotide sequences for making DNA substrates-

Table 1. The sequences of oligonucleotides used to make various DNA substrates.

ODN	Sequence (5'-3')	Length
HJds	AACGGCATAAGCTTGACGATTACATTGCTAGGACATCT TTGCCCACCTGCAGGTTCACCCA	61
HJ02	TGGGTGAACCTGCAGGTGGGCAAAGATGTCCTAGCAA T GTAATCGTCAAGCTTATGCCGTT	61

Table 2. Construction of synthetic DNA substrates by annealing respective oligos (see Methods in main text). Asterisks (*) indicate the ^{32}P labeled strand.

Substrate	Oligonucleotides used
Single-stranded DNA	HJ02*
Double-stranded DNA	HJ02* + HJds

2.3 ATPase assay

The assay was performed as previously described (11). Reaction mixes (20 µl), which contained 20 mM Tris-HCl buffer (pH 7.5), 12 mM $MgCl_2$, 2 mM DTT, 15% glycerol, 100 µg/ml BSA, 200 µM cold ATP, 200 pM [$\square$-^{32}P] ATP and indicated concentrations of AsCas12a, were incubated at 37 ºC for 1 h. The reaction was stopped by adding EDTA to a concentration of 25 mM. After incubation at 37 ºC for 10 min, aliquots (2 µl) of the reaction volume were spotted on a polyethyleneimine (PEI)-cellulose thin-layer plate (Whatman), and the reaction products were separated using a solution of 1 M formic acid and 0.5 M LiCl. The plate was air-dried, exposed to a phosphorimager screen, and visualised using Fuji FLA-9000 laser scanner. The band intensities were analysed and quantified using UVI- BandMap (version 97.04). The data was plotted using Graph-Pad Prism software (version 5.0).

2.4 Bioinformatics analysis

The amino acid sequence of AsCas12a was retrieved from the Protein Data Bank using the PDB ID, 5B43. This sequence, formatted in FASTA, was then input into the NsitePred tool to identify potential nucleotide-binding residues. Further, the most probable binding site was subject to molecular docking using SwissDock.

3 Results

3.1 AsCas12a hydrolyzes ATP in a concentration and time-dependent manner

Despite lacking canonical ATP-binding motifs or domains typically required for ATP hydrolysis, AsCas12a unexpectedly exhibited ATPase activity when included as a control in assays for RecA (2, 7). To systematically test for intrinsic ATPase activity, I incubated purified wild-type AsCas12a with [γ-^{32}P]ATP and monitored hydrolysis via TLC. Separation of substrate and product allowed visualization of free ^{32}Pi release. AsCas12a catalyzed ATP hydrolysis in a concentration-dependent manner, with increasing amounts of ^{32}Pi correlating with enzyme concentration up to 480 nM (Fig. 1A, B). These data indicate that AsCas12a is capable of catalyzing ATP turnover in the absence of DNA or RNA cofactors.

Figure 1. AsCas12a catalyzes hydrolysis of ATP to yield ADP and inorganic phosphate. (A), Representative image showing the separation of [□-^{32}P] ATP (substrate) and ^{32}Pi (product) by thin-layer chromatography (TLC) after incubation with indicated concentration of AsCas12a (lanes 2-12). Control lane 1 lacked AsCas12a. The assay was performed as described in Material and methods section. Varying concentrations of AsCas12a is depicted by an empty triangle above the gel image. (B), Graph shows the amount (%) of ATP hydrolyzed versus increasing concentration of AsCas12a. (C), (D), Kinetics of ATP hydrolysis reaction mediated by AsCas12a and its quantification. The data shows mean ± S.D.from three independent experiments. Note that in the first lane of both panels (A) and (C), [□ -^{32}P]ATP undergoes spontaneous degradation releasing Pi, and the products of ATP hydrolysis by AsCas12a migrate on TLC at same level.

3.2 ATPase activity of AsCas12a is independent of DNA cofactors

For many bacterial proteins, the property of ATPase activity is tightly linked to DNA metabolism, while in some others these are independent (12, 13). To determine whether DNA enhances or regulates ATP hydrolysis, I repeated the assay in the presence of either single-stranded DNA (ssDNA) or double-stranded DNA (dsDNA). The protocol involved pre-incubation of DNA with AsCas12a at 37 ºC for 10 minutes, followed by carrying out the ATPase assay. No significant difference was observed in

the rate or total extent of ATP hydrolysis with either cofactor (single-stranded or double-stranded DNA) compared to the DNA-free condition (Fig. 2A–D). This indicates that AsCas12a's ATPase activity is DNA-independent and likely intrinsic to the protein.

Figure 2. AsCas12a possesses DNA-independent ATPase activity. TLC-based ATPase assays were performed in the absence (panel A, lanes 2-12) or presence of ssDNA (panel B, lanes 2-12) or dsDNA (panel C, lanes 2-12) and varying amounts of AsCas12a as described in Materials and methods section. (A-C), control lane 1 lacked AsCas12a. The empty triangles above the gel images indicate increasing concentrations of AsCas12a. (D) Graph shows quantitative comparison of ATPase activity (shown in panels A-C) in the presence of different DNA cofactors. The amount of ATP hydrolyzed was quantified, and the data were plotted as the percentage of ATP hydrolyzed versus increasing concentrations of AsCas12a. Each bar is the mean ± standard error of three independent experiments.

3.3 Predicting AsCas12a's potential nucleotide-binding residues

Given the robust ATP hydrolysis activity exhibited by AsCas12a, it was pertinent to investigate its potential nucleotide-binding residues. The amino acid sequence of AsCas12a was obtained from the Protein Data Bank (PDB ID: 5B43) and analyzed using NsitePred, a freely accessible web-based tool that integrates sequence-derived

structural features, evolutionary profiles, and alignment data to predict nucleotide-binding residues with high accuracy (14). NsitePred identified four ATP-binding residues, along with additional sites potentially involved in binding other nucleotides, including ADP, AMP, GTP, and GDP (Table 3). Notably, residues Pro1069, Ala1070, and Tyr1072 were predicted with high confidence to constitute an ATP-binding site, which may function as a catalytic triad. These residues, all located within the NUC domain, were designated as the region of interest for molecular docking. Using SwissDock, ATP was docked as a ligand within this region (15) (Figure 3). The observed negative AC and SwissParam scores indicated favorable binding energy and strong predicted affinity (Table 4). Nonetheless, further validation through molecular dynamics simulations and site-directed mutagenesis will be required to confirm the functional relevance of the predicted ATP-binding site.

Table 3. Possible nucleotide binding site of AsCas12a.

Ligand	Amino acid residues
ATP	Ser925, Pro1069, Ala1070 and Tyr1072
ADP	Asn522, Asp924 and Thr1073
AMP	Tyr566 and Ile910
GTP	Arg915, Asp924, Ser925, Thr926, Asn997 and Thr1073
GDP	Tyr882, Gln883 and Ala1070

Table 4. SwissDock docking scores for ATP binding to Pro1069, Ala1070 and Tyr1072 residues of AsCas12a.

Cluster number	Cluster member	AC Score	SwissParam Score		Cluster number	Cluster member	AC Score	SwissParam Score
0	1	-432.544854	-7.7608					
1	1	-432.227006	-7.7293		Cluster number	Cluster member	AC Score	SwissParam Score
2	1	-432.123434	-7.7854		28	1	-420.697526	-7.3170
3	1	-429.881691	-7.7600		29	1	-420.624931	-6.9779
4	1	-429.714240	-7.8170		30	1	-420.388982	-6.9535
5	1	-429.089248	-7.0285		31	1	-419.902469	-7.3634
6	1	-428.579477	-7.5557		32	1	-419.866921	-6.8731
7	1	-427.612326	-7.1577		33	1	-419.480804	-7.1141
8	1	-427.214503	-7.0898		34	1	-419.432935	-6.8469
9	1	-427.208456	-7.1664		35	1	-419.237178	-7.4396
10	1	-427.031676	-8.0290		36	1	-419.167773	-6.6096
11	1	-426.958608	-7.7012		37	1	-418.901965	-7.0800
12	1	-426.795874	-7.0641		38	1	-418.044128	-6.5219
13	1	-425.753930	-7.6909		39	1	-417.611404	-7.4589
14	1	-424.746042	-7.4473		40	1	-417.270540	-7.5150
15	1	-424.576516	-7.4783		41	1	-416.923222	-7.0968
16	1	-424.564647	-7.6569		42	1	-416.800615	-6.7108
17	1	-424.486993	-6.8633		43	1	-416.727083	-6.8288
18	1	-424.100708	-7.3281		44	1	-416.623138	-6.2680
19	1	-424.081004	-7.4876		45	1	-416.604094	-6.9205
20	1	-423.853729	-7.3595		46	1	-415.999844	-6.7247
21	1	-423.220309	-7.3480		47	1	-415.590540	-6.7422
22	1	-422.812718	-7.7829		48	1	-415.561912	-6.9222
23	1	-422.479710	-7.1725		49	1	-415.531563	-6.4181
24	1	-422.474212	-7.0627					
25	1	-422.375010	-6.5444					
26	1	-421.676616	-7.4628					
27	1	-420.725179	-6.9334					

Figure 3. *In silico* prediction of ATP-binding residues in AsCas12a.(A) Schematic representation of the domain architecture of AsCas12a, with predicted ATP-binding residues indicated as red bars within the NUC (nuclease) domain.(B) Surface rendering of the AsCas12a structure generated using SwissDock, illustrating the docked ATP molecule positioned deep within a defined region of the protein, marked by an oval. (C) Enlarged schematic view of the predicted ATP-binding pocket in AsCas12a, highlighting the spatial arrangement of the binding site within the selected region.

4. Discussion

I report the unexpected discovery that AsCas12a, a well-characterized CRISPR-Cas effector, possesses intrinsic ATPase activity. Despite the absence of identifiable ATP-binding motifs, AsCas12a hydrolyzes ATP efficiently and independently of DNA cofactors. This distinguishes it from classical DNA-dependent ATPases and raises intriguing mechanistic and functional questions. The biological relevance of this activity remains to be determined. It is possible that ATP hydrolysis could modulate conformational changes in Cas12a during its activation cycle or interactions with other cellular factors (16). Alternatively, the ATPase activity may reflect a vestigial or context-specific enzymatic function with regulatory implications. Notably, ATPase activity is not

a known requirement for Cas12a's DNA cleavage function, which relies on a metal-dependent RuvC domain ([17](#)). However, auxiliary ATPase and ATP deamination activities in other CRISPR systems have been documented, particularly in Cas3 and CRISPR-associated Cad1 complexes, respectively, supporting the idea that CRISPR proteins may carry multifunctional capabilities ([18](#), [19](#)). Further biochemical, structural and *in vivo* studies will be necessary to elucidate the mechanistic basis and potential physiological role of this novel ATPase activity.

Authorship contribution

SB: Investigation, Visualization, Methodology; Writing- final version.

Declaration of competing interest

The author declares no competing interests are involved.

Acknowledgements

I thank Osamu Nureki (University of Tokyo) for sharing the pE-AsCas12a expression plasmid. This work was funded the Indian Institute of Science fellowship to SB. I also thank the Department of Biochemistry, Indian Institute of Science for providing facilities to conduct protein purifications and biochemical assays.

Abbreviations: ATP, Adenosine triphosphate; CRISPR-Cas, Clustered regularly interspaced short palindromic repeats-CRISPR associated; dsDNA, Double-stranded DNA; IPTG, Isopropyl-β-D-thiogalactopyranoside; ODN, oligonucleotide; ssDNA, Single-stranded DNA

Disclaimer:

This study presents preliminary findings that have not yet been peer-reviewed. The data and interpretations should be considered with caution, and further validation, including molecular dynamics simulations and site-directed mutagenesis, is required to confirm the proposed finding. Feedback from the scientific community is welcome to refine these conclusions through continued investigation.

References

[1] Bhattacharya S, Agarwal A, Muniyappa K. Deciphering the Substrate Specificity Reveals that CRISPR-Cas12a Is a Bifunctional Enzyme with Both Endo- and Exonuclease Activities. J Mol Biol. 2024; 436(10):168550. doi:10.1016/j.jmb.2024.168550.

[2] Zetsche B, Gootenberg JS, Abudayyeh OO, Slaymaker IM, Makarova KS, Essletzbichler P, Volz SE, Joung J, van der Oost J, Regev A, Koonin EV, Zhang F. Cpf1

is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system. Cell. 2015 ;163(3):759-71. doi: 10.1016/j.cell.2015.09.038.

[3] Ma E, Chen K, Shi H, Stahl EC, Adler B, Trinidad M, Liu J, Zhou K, Ye J, Doudna JA. Improved genome editing by an engineered CRISPR-Cas12a. Nucleic Acids Res. 2022; 50(22):12689-12701. doi: 10.1093/nar/gkac1192.

[4] Jianwei L, Jobichen C, Machida S, Meng S, Read RJ, Hongying C, Jian S, Yuan YA, Sivaraman J. Structures of apo Cas12a and its complex with crRNA and DNA reveal the dynamics of ternary complex formation and target DNA cleavage. PLoS Biol. 2023; 21(3):e3002023. doi: 10.1371/journal.pbio.3002023.

[5] Swarts DC, van der Oost J, Jinek M. Structural Basis for Guide RNA Processing and Seed-Dependent DNA Targeting by CRISPR-Cas12a. Mol Cell. 2017; 66(2):221-233.e4. doi: 10.1016/j.molcel.2017.03.016.

[6] Swarts DC, Jinek M. Mechanistic Insights into the cis- and trans-Acting DNase Activities of Cas12a. Mol Cell. 2019; 73(3):589-600.e4. doi: 10.1016/j.molcel.2018.11.021.

[7] Toseland CP, Webb MR. ATPase mechanism of the 5'-3' DNA helicase, RecD2: evidence for a pre-hydrolysis conformation change. J Biol Chem. 2013; 288(35):25183-25193. doi: 10.1074/jbc.M113.484667.

[8] Chand MK, Carle V, Anuvind KG, Saikrishnan K. DNA-mediated coupling of ATPase, translocase and nuclease activities of a Type ISP restriction-modification enzyme. Nucleic Acids Res. 2020; 48(5):2594-2603. doi: 10.1093/nar/gkaa023.

[9] Johnson SL, Narlikar GJ. ATP Hydrolysis Coordinates the Activities of Two Motors in a Dimeric Chromatin Remodeling Enzyme. J Mol Biol. 2022; 434(14):167653. doi: 10.1016/j.jmb.2022.167653.

[10] Fu H, Le S, Chen H, Muniyappa K, Yan J. Force and ATP hydrolysis dependent regulation of RecA nucleoprotein filament by single-stranded DNA binding protein. Nucleic Acids Res. 2013; 41(2):924-32. doi: 10.1093/nar/gks1162.

[11] Thakur M, Kumar MB, Muniyappa K. Mycobacterium tuberculosis UvrB Is a Robust DNA-Stimulated ATPase That Also Possesses Structure-Specific ATP-Dependent DNA Helicase Activity. Biochemistry. 2016; 55(41):5865-5883. doi: 10.1021/acs.biochem.6b00558.

[12] Zylicz M, LeBowitz JH, McMacken R, Georgopoulos C. The dnaK protein of Escherichia coli possesses an ATPase and autophosphorylating activity and is essential in an in vitro DNA replication system. Proc Natl Acad Sci U S A. 1983; 80(21):6431-5. doi: 10.1073/pnas.80.21.6431.

[13] Biswas T, Pero JM, Joseph CG, Tsodikov OV. DNA-dependent ATPase activity of bacterial XPB helicases. Biochemistry. 2009; 48(12):2839-48. doi: 10.1021/bi8022416.

[14] Chen K, Mizianty MJ, Kurgan L. Prediction and analysis of nucleotide-binding residues using sequence and sequence-derived structural descriptors. Bioinformatics. 2012;28(3):331-41. doi: 10.1093/bioinformatics/btr657.

[15] Bugnon M, Röhrig UF, Goullieux M, Perez MAS, Daina A, Michielin O, Zoete V. SwissDock 2024: major enhancements for small-molecule docking with Attracting Cavities and AutoDock Vina. Nucleic Acids Res. 2024;52(W1):W324-W332. doi: 10.1093/nar/gkae300.

[16] Stella S, Mesa P, Thomsen J, Paul B, Alcón P, Jensen SB, Saligram B, Moses ME, Hatzakis NS, Montoya G. Conformational Activation Promotes CRISPR-Cas12a Catalysis and Resetting of the Endonuclease Activity. Cell. 2018; 175(7):1856-1871.e21. doi: 10.1016/j.cell.2018.10.045.

[17] Strohkendl I, Saha A, Moy C, Nguyen AH, Ahsan M, Russell R, Palermo G, Taylor DW. Cas12a domain flexibility guides R-loop formation and forces RuvC resetting. Mol Cell. 2024 Jul 25;84(14):2717-2731.e6. doi: 10.1016/j.molcel.2024.06.007.

[18] Sinkunas T, Gasiunas G, Fremaux C, Barrangou R, Horvath P, Siksnys V. Cas3 is a single-stranded DNA nuclease and ATP-dependent helicase in the CRISPR/Cas immune system. EMBO J. 2011; 30(7):1335-42. doi: 10.1038/emboj.2011.41.

[19] Baca CF, Majumder P, Hickling JH, Ye L, Teplova M, Brady SF, Patel DJ, Marraffini LA. The CRISPR-associated adenosine deaminase Cad1 converts ATP to ITP to provide antiviral immunity. Cell. 2024; 187(25):7183-7195.e24. doi: 10.1016/j.cell.2024.10.002.